

**First records of molluscs naturally infected with  
*Angiostrongylus cantonensis* (Nematoda: Metastrongyloidea)  
in Northeastern Brazil, including new global records of natural  
intermediate hosts**

**Jucicleide Ramos-de-Souza<sup>1</sup>, Silvana Carvalho Thiengo<sup>2</sup>, Monica Ammon Fernandez<sup>2</sup>, Suzete Rodrigues Gomes<sup>2</sup>, Jéssica Corrêa Antônio<sup>2</sup>, Marianna de Carvalho Clímaco<sup>1</sup>, Juberlan Silva Garcia<sup>3</sup>, Arnaldo Maldonado-Junior<sup>3</sup>, Luciene Barbosa<sup>1</sup>, Silvio Santana Dolabella<sup>1</sup>**

**ABSTRACT**

Human neural angiostrongyliasis is an emerging infectious disease caused by nematode *Angiostrongylus cantonensis*. The present study investigated the presence of *Angiostrongylus* spp. in terrestrial molluscs collected from the following areas in the Metropolitan Region of Aracaju, Sergipe State, Brazil: Barra dos Coqueiros, Nossa Senhora do Socorro, Sao Cristovao and Aracaju. In total, 703 specimens representing 13 mollusc species were screened for *Angiostrongylus* spp. Larvae of *Angiostrongylus* spp. were found in three species. Larvae recovered from *Achatina fulica* were used for experimental infection in Wistar rats (*Rattus norvegicus*). For specific identification of nematodes, the mitochondrial cytochrome c oxidase subunit I (COI) was sequenced from both larvae and adults recovered from molluscs and rats, respectively. Infection with *A. cantonensis* was detected in all municipalities and in the following three host species: *Bulimulus tenuissimus*, *Cyclodontina fasciata* (Barra dos Coqueiros), and *A. fulica* (Aracaju, Nossa Senhora do Socorro and Sao Cristovao). Coinfections were also found with *Caenorhabditis* sp. and *Strongyluris* sp. larvae. This is the first study of the helminth fauna associated with the terrestrial malacofauna in Sergipe State, and confirms that these three snail species are involved in the transmission of *A. cantonensis* in the state. In addition, *B. tenuissimus* and *C. fasciata* are newly reported natural hosts of the parasite.

**KEYWORDS:** Eosinophilic meningitis. *Angiostrongylus*. *Achatina fulica*. *Bulimulus tenuissimus*. *Cyclodontina fasciata*. Land snails.

**INTRODUCTION**

*Achatina fulica* Bowdich, 1822 is listed among 100 of the worst invasive species in the world and is present in almost all Brazilian states<sup>1,2</sup>. This species' rapid proliferation is the main reason it is considered an agricultural pest. It can also act as an intermediate host of *Angiostrongylus cantonensis* (Chen, 1935) and *Angiostrongylus costaricensis* Morera and Céspedes, 1971. These two species are the etiological agents of human neural angiostrongyliasis and human abdominal angiostrongyliasis, respectively<sup>2-4</sup>.

Eosinophilic meningitis is the principal clinical manifestation of human neural angiostrongyliasis, an emerging infectious disease that causes severe impairment of the central nervous system and can lead to death<sup>5-7</sup>. This nematode has low specificity

<sup>1</sup>Universidade Federal de Sergipe, Departamento de Morfologia, Laboratório de Entomologia e Parasitologia Tropical, São Cristóvão, Sergipe, Brazil

<sup>2</sup>Fundação Oswaldo Cruz, Instituto Oswaldo Cruz, Laboratório de Referência Nacional em Esquistossomose – Malacologia, Rio de Janeiro, Rio de Janeiro, Brazil

<sup>3</sup>Fundação Oswaldo Cruz, Instituto Oswaldo Cruz, Laboratório de Biologia e Parasitologia de Mamíferos Silvestres Reservatórios, Rio de Janeiro, Rio de Janeiro, Brazil

**Correspondence to:** Silvio Santana Dolabella

Universidade Federal de Sergipe, Departamento de Morfologia, Laboratório de Entomologia e Parasitologia Tropical, Av. Marechal Rondon, s/n, Jd. Rosa Elze, CEP 49100-000, São Cristóvão, SE, Brazil

**E-mail:** [dolabella@ufs.br](mailto:dolabella@ufs.br)

**Received:** 16 April 2018

**Accepted:** 16 August 2018

for its intermediate hosts, which means that many mollusc species have already been found to be naturally infected globally<sup>8,9</sup>. However, *A. fulica* is an important intermediate host for *A. cantonensis*<sup>4,8,10-12</sup>.

Human neural angiostrongyliasis occurs primarily by ingestion of snails that are infected with L<sub>3</sub> larvae<sup>13</sup>. These larvae reach the brain where they mature into L<sub>4</sub> and L<sub>5</sub> but are unable to migrate from the brain to the heart (as they do in the definitive rat hosts) but move around in the brain causing neurological damage and, especially when they die, cause intense inflammatory reactions that result in additional neurological damage and symptoms<sup>5,7,14</sup>.

More than 2,800 human cases of eosinophilic meningitis caused by *A. cantonensis* have been reported in more than 30 countries<sup>11</sup> since it was first reported in Taiwan in 1945<sup>15</sup>. In Brazil, out of 84 suspected cases, there have been 34 confirmed cases of *A. cantonensis* infections in humans as well as the confirmation of definitive hosts and infected intermediate hosts in different regions of the country<sup>10</sup>.

In Sergipe State, 19 municipalities have reported infestation of *A. fulica* (Comissao de Combate ao Caramujo Africano, personal communication) in addition to Barra dos Coqueiros municipality<sup>8</sup>. The objective of this study was to verify the infection rate by nematode larvae in *A. fulica* and other terrestrial molluscs collected in the Aracaju Metropolitan Region, Sergipe State, Northeastern Brazil.

## MATERIALS AND METHODS


This study was undertaken in urban environments of four municipalities that constitute the Metropolitan Region of Aracaju, Sergipe State (Figure 1): Aracaju, Barra dos Coqueiros, Nossa Senhora do Socorro and Sao Cristovao (Table 1).

Each point of collection was georeferenced, followed by manual random sampling of terrestrial molluscs using forceps

and gloves. The collections were carried out in the following three seasons: April (autumn), July (winter) and October (spring) of 2016, at three locations in each municipality. The samples were sent to the National Reference Laboratory for Schistosomiasis - Malacology (LRNEM) of the Instituto Oswaldo Cruz, Fundação Oswaldo Cruz (IOC/Fiocruz, Rio de Janeiro), where they were analyzed. The identification of the molluscs species was done based on conchological characteristics compared to photos and descriptions from catalogues<sup>16-19</sup>. We also compared our material with material deposited in the Instituto Oswaldo Cruz. Samples of each species and from each collection locality were deposited in the Mollusca Collection of the Instituto Oswaldo Cruz (CMIOC 10103 - 10134, 11206).

To collect the nematode larvae, the molluscs were artificially digested by a modified Wallace and Rosen technique<sup>20</sup>. The recovered material was observed under a stereoscopic microscope following the LRNEM Identification Guide and based on the literature<sup>4,21</sup>. The *Angiostrongylus* sp. (Nematoda: Metastrongylidae) larvae were identified under a compound microscope according to Thiengo *et al.*<sup>4</sup>. Ten *Angiostrongylus* sp. larvae from each sample were collected and cryopreserved at -20 °C until DNA extraction for subsequent molecular analysis. Some rhabditiform larvae were observed and prepared for DNA extraction. In addition, *Strongyluris* sp. was morphologically identified, based on Thiengo<sup>21</sup>.

To assess the viability of the parasites, *Angiostrongylus* sp. larvae recovered from *A. fulica* from the municipality of Aracaju were used to infect two specimens of *Rattus norvegicus* through an orogastric tube at a concentration of 50 L<sub>3</sub> larvae per rodent. Fifty days after infection, the animals were killed and necropsied to confirm the presence of adult *A. cantonensis*<sup>22</sup>. This procedure was carried out following the Ethics Commission on Animal Use of the Oswaldo Cruz Foundation (LW-47/14).



**Figure 1** - Study area: Aracaju Metropolitan Region, including the capital of Sergipe State, the municipalities of Barra dos Coqueiros, Nossa Senhora do Socorro and Sao Cristovao. The black spots represent the collection localities

**Table 1** - Geographical coordinates by location of collection of terrestrial molluscs

Municipalities	Description of locality	Geographical coordinates
Barra dos Coqueiros	Square / open area	10°49'30.1"S 36°56'46.7"W
	Outer clothing area	10°49'42.6"S 36°56'08.2"W
	School backyard / Rubble	10°49'12.2"S 36°56'58.9"W
Sao Cristovao	Street / close to houses	10°55'57.8"S 37°06'26.6"W
	Street / close to houses	10°55'27.7"S 37°06'43.2"W
	Street and house backyard	10°55'43.1"S 37°07'16.2"W
Aracaju	Street / close to houses	10°58'29.7"S 37°06'16.7"W
	Ground / adjacent to a commercial establishment	11°00'11.9"S 37°05'03.2"W
	Close to sewer-line	10°54'24.1"S 37°05'23.3"W
Nossa Senhora do Socorro	Garden and houses backyard	10°50'19.0"S 37°03'15.0"W
	Houses backyard	10°51'02.2"S 37°05'38.1"W
	Street and wasteland	10°53'23.0"S 37°08'50.9"W

### Molecular diagnosis of nematode larvae

DNA was extracted from 10 *Angiostrongylus* sp. larvae obtained from each mollusc in which larvae had been found and resuspended in 30 µL of PCR Buffer solution (Thermo Fisher Scientific, Massachusetts, USA). The polymerase chain reaction (PCR) mixtures were prepared in a volume of 20 µL containing 8 µL of ultrapure water, 5 µL of 10% trehalose, 2.5 µL of 10x PCR Reaction Buffer, 2 µL of 2.5 mM dNTPs, 1.25 µL 50 mM MgCl<sub>2</sub>, 0.5 µL each of 5 µM forward and reverse primer (Nem 3 from Prosser *et al.*<sup>23</sup>), and 0.25 µL of recombinant Taq DNA polymerase (Thermo Fisher Scientific). A total of 5 µL of the DNA sample was added to the mixture, producing a final volume of 25 µL for each reaction. For all the reactions, ultrapure water was used as a negative control template, and the positive control was performed with genomic DNA of *A. cantonensis* as template. The 700 bp fragment of the *Angiostrongylus* sp. larvae mitochondrial cytochrome *c* oxidase subunit I (COI) gene was amplified using the following PCR conditions: initial denaturation at 94 °C for 1 min, five cycles at 94 °C for 40 s, 45 °C for 40 s, and 72 °C for 1 min, followed by 30 cycles at 94 °C for 40 s, 51 °C for 40 s, and 72 °C for 1 min and a final extension at 72 °C for 1 min.

For rhabditiform larvae, individual DNA from some isolates was used to amplify the 480 bp ribosomal 18S region<sup>24</sup>.

The PCR products were purified using the Illustra GFX PCR DNA and Gel Band Purification Kit (GE Healthcare, Little Chalfont, UK) following the manufacturer's protocol. Purified products after amplification were bidirectionally sequenced using BigDye Terminator v3.1 Cycle Sequencing kit (Applied Biosystems, California, USA) according

to the manufacturer's instructions. Chromatograms of the sequences obtained were analyzed and edited using Geneious version R9 software (<http://www.geneious.com>), resulting in a consensus sequence (contig). Then, a comparative similarity search was performed on GenBank ([www.ncbi.nlm.nih.gov/genbank/](http://www.ncbi.nlm.nih.gov/genbank/))<sup>25</sup> using BLAST (Basic Local Alignment Search Tool) to identify the closest match. The obtained sequences are available in GenBank under the access N° MH511539 - MH511542, and MH547424.

### RESULTS

In total, 703 terrestrial gastropods were analyzed (190 in Aracaju, 250 in Barra dos Coqueiros, 159 in Nossa Senhora do Socorro, and 104 in São Cristovão) and identified as the following 13 species: *Achatina fulica* (Bowdich, 1822), *Allopeas gracile* (Hutton, 1834), *Subulina octona* (Bruguière, 1789), *Leptinaria unilamellata* (d'Orbigny, 1835), *Bulimulus tenuissimus* (d'Orbigny, 1835), *Cyclodontina fasciata* (Potiez & Michaud, 1838), *Latipes erinaceus* (Colosi, 1921), *Sarasinula linguaeformis* (Semper, 1885), *Streptartemon cookeanus* (Baker, 1914), *Streptartemon quixadensis* (Baker, 1914), *Tamayoa banghaasi* (Thiele, 1927), *Helicina* sp. and *Omalonyx* sp.

*Angiostrongylus* sp. larvae were found in specimens of *A. fulica*, *B. tenuissimus* and *C. fasciata* and were later confirmed by molecular analysis as *A. cantonensis* (Table 2).

*Achatina fulica* was found in all four municipalities, with snails harboring infection by *A. cantonensis* in three of them (Aracaju, Sao Cristovao and Nossa Senhora do Socorro). *Bulimulus tenuissimus* occurred in all municipalities, while specimens of *C. fasciata* were found only in Aracaju and Barra dos Coqueiros. Both *B. tenuissimus* and *C. fasciata*

**Table 2** - Prevalence of nematodes in terrestrial molluscs from the Metropolitan Region of Aracaju, Sergipe State, Northeast Brazil, from April to October 2016

Locality	Species	N° specimens (n)	Positive molluscs (n)		
			<i>A. cantonensis</i>	<i>Caenorhabditis</i> sp.	<i>Strongyluris</i> sp.
Barra dos Coqueiros	<i>A. fulica</i>	110	-	48	22
	<i>B. tenuissimus</i>	23	1	3	-
	<i>C. fasciata</i>	112	2	48	23
	<i>Helicina</i> sp.	-	-	-	-
	<i>S. cookeanus</i>	1	-	-	-
	<i>S. linguaeformis</i>	2	-	-	1
	<i>S. octona</i>	2	-	-	-
	<i>S. quixadensis</i>	-	-	-	-
Sao Cristovao	<i>T. banghaasi</i>	-	-	-	-
	<i>A. fulica</i>	63	22	25	1
	<i>B. tenuissimus</i>	17	-	1	-
	<i>S. linguaeformis</i>	10	-	1	-
Aracaju	<i>S. octona</i>	14	-	1	-
	<i>A. fulica</i>	97	36	46	12
	<i>A. gracille</i>	-	-	-	-
	<i>B. tenuissimus</i>	19	-	2	-
	<i>C. fasciata</i>	43	-	4	1
	<i>S. cookeanus</i>	2	-	-	-
	<i>S. linguaeformis</i>	5	-	-	-
Nossa Senhora do Socorro	<i>S. octona</i>	24	-	2	-
	<i>T. banghaasi</i>	-	-	-	-
	<i>A. fulica</i>	85	5	34	-
	<i>B. tenuissimus</i>	13	-	3	-
	<i>L. erinaceus</i>	-	-	-	-
	<i>L. unilamellata</i>	47	-	5	-
	<i>Omalonyx</i> sp.	2	-	1	-
<i>S. linguaeformis</i>	2	-	-	-	
<i>S. octona</i>	10	-	4	-	
<i>T. banghaasi</i>	-	-	-	-	

were infected by *A. cantonensis* in the municipality of Barra dos Coqueiros, near the port area of Sergipe State.

Co-infections were observed between *A. cantonensis* and other nematodes, such as *Strongyluris* sp. and rhabditiform larvae (Table 3). The rhabditiform larvae were analyzed molecularly and were found to be 99% similar to a sequence from GenBank identified as *Caenorhabditis* sp.

Infections of *Caenorhabditis* sp. were observed in *A. fulica*, *B. tenuissimus*, *C. fasciata*, *S. octona*, *L. unilamellata*, *S. linguaeformis* and *Omalonyx* sp. Infections by *Strongyluris* sp. were observed in *A. fulica*, *C. fasciata* and *S. linguaeformis*.

Two rats that were experimentally infected with *Angiostrongylus* sp. isolates displayed symptoms 50 days after infection. Forty *A. cantonensis* adult worms were recovered in the first rat (37.5% males and 62.5% females), and 42 adult worms were recovered in the second (40.4% males and 59.5% females).

## DISCUSSION

The snail *A. fulica* has already been reported as being parasitized with *A. cantonensis* by several authors<sup>2,4,8,10,19,26</sup>. In Brazil, the species is considered one of the main


**Table 3** - Prevalence of co-infection between *Angiostrongylus cantonensis*, *Caenorhabditis* sp. and *Strongyluris* sp. in land molluscs from the Aracaju Metropolitan Region, Sergipe State, Northeast Brazil, from April to October 2016

Locality	Species	Specimens analyzed (n)	Positive specimens for <i>A. cantonensis</i>		Co-infection (n)		
			(n)	(%)	<i>A. cantonensis</i> + <i>Strongyluris</i> sp.	<i>A. cantonensis</i> + <i>Caenorhabditis</i> sp.	<i>A. cantonensis</i> + <i>Caenorhabditis</i> sp. + <i>Strongyluris</i> sp.
Barra dos Coqueiros	<i>B. tenuissimus</i>	23	1	4.3	-	1	-
	<i>C. fasciata</i>	109	2	1.8	-	-	-
São Cristovao	<i>A. fulica</i>	63	22	34.9	-	6	-
Aracaju	<i>A. fulica</i>	99	36	36.4	3	10	7
Nossa Senhora do Socorro	<i>A. fulica</i>	85	5	5.9	-	-	-

potential transmitters of human neural angiostrongyliasis, considering their widespread distribution, high population densities and proximity to humans<sup>10,26</sup>.

Despite knowledge of the presence of the giant African snail in Sergipe State since 2006<sup>27,28</sup>, only one previous investigative study has been carried out in this region<sup>8</sup>. Ten years after the first detection of *A. cantonensis* in Brazil, this study records its presence in Sergipe State for the first time; it is also the first report of *C. fasciata* and *B. tenuissimus* as natural intermediate hosts of the parasite. In addition, we observed co-infection with other nematodes, which is consistent with other studies<sup>12,26,28</sup>.

Since 2007, when the first Brazilian case of eosinophilic meningitis caused by *A. cantonensis* was recorded, in Espírito Santo State<sup>29</sup>, reports of the disease have increased. Currently, the parasite has been found in 12 states (Figure 2)<sup>2,7,10,29-31</sup>.



**Figure 2** - Distribution in Brazil of *Angiostrongylus cantonensis* in natural vertebrate and invertebrate hosts, and in humans. Adapted from: Thiengo *et al.*<sup>5</sup>, Carvalho *et al.*<sup>8</sup>, Morassutti *et al.*<sup>10</sup>, Espírito Santo *et al.*<sup>30</sup>, Moreira *et al.*<sup>31</sup>. AM: Amazonas; PA: Para; CE: Ceara; PE: Pernambuco; SE: Sergipe; BA: Bahia; ES: Espírito Santo; RJ: Rio de Janeiro; SP: Sao Paulo; PR: Parana; SC: Santa Catarina; RS: Rio Grande do Sul

The first known occurrence of *A. fulica* naturally infected by *A. cantonensis* in the northeast region of Brazil was reported in 2008, with human cases of eosinophilic meningitis and naturally infected intermediate hosts in the municipalities of Escada and Olinda, Pernambuco State<sup>4</sup>. A study carried out in Brazilian port areas, including the port of Sergipe State, reported the presence of terrestrial and aquatic molluscs that, although previously negative for *A. cantonensis*, tested positive in the bordering Bahia State<sup>8</sup> and in Ceara State<sup>10</sup>.

Several other terrestrial molluscs have also been found to be naturally infected with this nematode in Brazil<sup>2,3,8,19</sup> and in other countries<sup>9,13,14</sup>. Among these species are *S. linguaeformis* and *S. octona*<sup>2,3,8,9</sup>, which, however, were not found to be infected in the present study. A possible reason for this is the lower abundance of these species in the sampled localities. The three species found infected by *A. cantonensis* were also those that were more abundant in the environment.

The present study identifies, for the first time, the following two species as natural hosts of the parasite: *C. fasciata*, which had not yet been described in the literature as an intermediate host for *A. cantonensis* or for other nematodes of human and veterinary interest, and *B. tenuissimus*, described as a host of metacercariae of the genus *Postharmostomum*<sup>32</sup>. This interaction between exotic and native species has become a new concern in the ecology of the disease and may result in increased parasitic spread as well as pose a threat to native species<sup>9</sup>.

Studies indicate the need to implement control measures against molluscs associated with public health, particularly the exotic species *A. fulica*. However, passive geographic dispersal is the main form of spread of the giant African snail, which hinders its control<sup>33</sup>.

The present study reinforces the need to investigate the current situation of *A. cantonensis* dispersal in the Northeastern region of Brazil, since there are reports of

its presence in four of its nine states, i.e., Bahia, Ceara, Pernambuco and Sergipe<sup>7,8</sup>, as well as the evident dispersal of *A. fulica* in several municipalities in the Sergipe State.

## ACKNOWLEDGMENTS

The authors are grateful to the Genomic Platform DNA Sequencing - RPT01A (Rede de Plataformas Tecnológicas FIOCRUZ) and for the technical support of the LRNEM/IOC/Fiocruz team, Eduardo Cinilha for help with map preparation, and the CAPES for the scholarship to Jucicleide R. Souza.

## CONFLICT OF INTERESTS

The authors declare that there are no conflicts of interest.

## REFERENCES

1. Lowe S, Browne M, Boudjelas S, De Poorter M. 100 of the world's worst invasive alien species: a selection from the Global Invasive Species Database. Auckland: Invasive Species Specialist Group; 2000.
2. Thiengo SC, Fernandez, MA. Gastrópodes neotropicais continentais de importância médica. In: Coura JR, editor. Dinâmica das doenças infecciosas e parasitárias. 2ª ed. Rio de Janeiro: Guanabara Koogan; 2018. p.131-53.
3. Laitano AC, Genro JP, Fontoura R, Branco SS, Maurer RL, Graeff-Teixeira C, et al. Report on the occurrence of *Angiostrongylus costaricensis* in southern Brazil, in a new intermediate host from the genus *Sarasinula* (Veronicellidae, Gastropoda). *Rev Soc Bras Med Trop.* 2001;34:95-7.
4. Thiengo SC, Maldonado Júnior A, Mota EM, Torres EJ, Caldeira R, Carvalho OS, et al. The giant African snail *Achatina fulica* as natural intermediate host of *Angiostrongylus cantonensis* in Pernambuco, northeast Brazil. *Acta Trop.* 2010;115:194-9.
5. Thiengo SC, Simões RO, Fernandez M, Maldonado Júnior A. *Angiostrongylus cantonensis* and rat lung worm disease in Brazil. *Hawaii J Med Public Health.* 2013;72 Suppl 2:18-22.
6. Graeff-Teixeira C, Silva AC, Yoshimura K. Update on eosinophilic meningoencephalitis and its clinical relevance. *Clin Microbiol Rev.* 2009;22:322-48.
7. Cowie RH. Biology, systematics, life cycle, and distribution of *angiostrongylus cantonensis*, the cause of rat lungworm disease. *Hawaii J Med Public Health.* 2013;72 Suppl 2:6-9.
8. Carvalho OS, Scholte RG, Mendonça CL, Passos LK, Caldeira RL. *Angiostrongylus cantonensis* (Nematode: Metastrongyloidea) in molluscs from harbour areas in Brazil. *Mem Inst Oswaldo Cruz.* 2012;107:740-6.
9. Kim JR, Hayes KA, Yeung NW, Cowie RH. Diverse gastropod hosts of *angiostrongylus cantonensis*, the rat lungworm, globally and with a focus on the Hawaiian Islands. *PloS One.* 2014;9:e94969.
10. Morassutti AL, Thiengo SC, Fernandez M, Sawanyawisuth K, Graeff-Teixeira C. Eosinophilic meningitis caused by *Angiostrongylus cantonensis*: an emergent disease in Brazil. *Mem Inst Oswaldo Cruz.* 2014;109:399-407.
11. Wang QP, Lai DH, Zhu XQ, Chen XG, Lun ZR. Human *angiostrongyliasis*. *Lancet Infect Dis.* 2008;8:621-30.
12. Maldonado Júnior A, Simões RO, Oliveira AP, Motta EM, Fernandez MA, Pereira ZM, et al. First report of *Angiostrongylus cantonensis* (Nematoda: Metastrongylidae) in *Achatina fulica* (Mollusca: Gastropoda) from Southeast and South Brazil. *Mem Inst Oswaldo Cruz.* 2010;105:938-41.
13. Vitta A, Polsut W, Fukruksa C, Yimthin T, Thanwisai A, Dekumyoy P. Levels of infection with the lungworm *Angiostrongylus cantonensis* in terrestrial snails from Thailand, with *Cryptozonia siamensis* as a new intermediate host. *J Helminthol.* 2016;90:737-41.
14. Maldonado Júnior A, Simões R, Thiengo SC. *Angiostrongyliasis* in the Americas. In: Morales-Lorenzo J, editor. *Zoonosis.* London: IntechOpen; 2012. p.303-20.
15. Beaver PC, Rosen L. Memorandum on the first report of *Angiostrongylus* in Man, by Nomura and Lin, 1945. *Am J Trop Med Hyg.* 1964;13:589-90.
16. Thomé JW. Estado atual da sistemática dos Veronicellidae (Mollusca, Gastropoda) americanos, com comentários sobre sua importância econômica, ambiental e na saúde. *Biociências.* 1993;1:61-75.
17. Salgado NC, Coelho AC. Moluscos terrestres do Brasil (Gastrópodes operculados ou não, exclusive Veronicellidae, Milacidae e Limacidae). *Rev Biol Trop.* 2003;5 Suppl 3:149-89.
18. Simone LR. Land and freshwater molluscs of Brazil: an illustrated inventory on the Brazilian malacofauna, including neighbor regions of the South America, respect to the terrestrial and freshwater ecosystems. São Paulo: EGB; 2006.
19. Ohlweiler FP, Takahashi FY, Guimarães MC, Gomes SR, Kawano T. Manual de gastrópodes límnicos e terrestres do Estado de São Paulo associados às helmintoses. Porto Alegre: Redes; 2010.
20. Wallace GD, Rosen L. Techniques for recovering and identifying larvae of *Angiostrongylus cantonensis* from molluscs. *Malacologia.* 1969;7:427-38.
21. Thiengo SC. Presence of Strongylurid-like larvae (Nematoda) in some terrestrial molluscs in Brazil. *Mem Inst Oswaldo Cruz.* 1995;90:619-20.
22. Garcia JS, Lúcio CS, Bonfim TC, Maldonado Júnior A, Tunholi VM, Tunholi-Alves VM, et al. Metabolic and histopathological profile of *Rattus norvegicus* (Wistar) experimentally infected by *Angiostrongylus cantonensis* (Chen, 1935). *Exp Parasitol.* 2014;137:35-40.

23. Prosser SW, Velarde-Aguilar MG, León-Règagnon V, Hebert PDN. Advancing nematode barcoding: a primer cocktail for the cytochrome c oxidase subunit I gene from vertebrate parasitic nematodes. *Mol Ecol Resour.* 2013;13:1108-15.
24. Blaxter ML, De Ley P, Garey JR, Liu LX, Scheldeman P, Vierstraete A, et al. A molecular evolutionary framework for the phylum Nematoda. *Nature.* 1998;392:71-5.
25. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics.* 2012;28:1647-9.
26. Oliveira AP, Gentile R, Maldonado Júnior A, Torres EJ, Thiengo SC. *Angiostrongylus cantonensis* infection in molluscs in the municipality of São Gonçalo, a metropolitan area of Rio de Janeiro, Brazil: role of the invasive species *Achatina fulica* in parasite transmission dynamics. *Mem Inst Oswaldo Cruz.* 2015;110:739-44.
27. Ramos-de-Souza J, Thiengo SC, Gomes SR, Fernandez MA, Clímaco MC, Jain S, et al. Native and alien terrestrial snails in urban areas of Sergipe, Northeast Brasil. *Tentacle.* 2017;25:9-13.
28. Thiengo SC. Helminthoses de interesse médico-veterinário transmitidas por moluscos no Brasil. In: Santos SB, Pimenta AD, Thiengo SC, Fernandez MA, Absalão RS, organizadores. Tópicos em malacologia: ecos do XVIII EBRAM. Rio de Janeiro: Sociedade Brasileira de Malacologia; 2007. p.287-94.
29. Caldeira RL, Mendonça CL, Goveia CO, Lenzi HL, Graeff-Teixeira C, Lima WS, et al. First record of molluscs naturally infected with *Angiostrongylus cantonensis* (Chen, 1935) (Nematoda: Metastrongylidae) in Brazil. *Mem Inst Oswaldo Cruz.* 2007;102:887-9.
30. Espírito-Santo MC, Pinto PL, Mota DJ, Gryscek RC. The first case of *Angiostrongylus cantonensis* eosinophilic meningitis diagnosed in the city of São Paulo, Brazil. *Rev Inst Med Trop Sao Paulo.* 2013;55:129-32.
31. Moreira VL, Giese EG, Melo FT, Simões RO, Thiengo SC, Maldonado Júnior A, et al. Endemic angiostrongyliasis in the Brazilian Amazon: natural parasitism of *Angiostrongylus cantonensis* in *Rattus rattus* and *R. norvegicus*, and sympatric giant African land snails, *Achatina fulica*. *Acta Trop.* 2013;125:90-7.
32. Thiengo SC, Amato SB. *Phyllocaulis variegatus* (Mollusca: Veronicellidae), a new intermediate host for *Brachylaima* sp. (Digenea: Brachylaimatidae). *Mem Inst Oswaldo Cruz.* 1995;90:621-2.
33. Colley E, Fisher ML. Avaliação dos problemas enfrentados no manejo do caramujo gigante africano *Achatina fulica* (Gastropoda: Pulmonata) no Brasil. *Zoologia.* 2009;26:674-83.