

Morphology and Molecular Characterization of *Alternaria argyranthemi* on *Chrysanthemum coronarium* in China

Huan Luo^a, Zhen Zhou Xia^a, Yun Yun Chen^b, Yi Zhou^a and Jian Xin Deng^a

^aDepartment of Plant Protection, College of Agriculture, Yangtze University, Jingzhou, China; ^bPlant Protection, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China

ABSTRACT

Chrysanthemum coronarium is an economically important plant in Asia, and used medicinally, ornamentally and as a vegetable. In April 2017, leaf spot disease on *C. coronarium* was observed in Shiyan, Hubei, China. A single-spore isolate was obtained and identified based on morphology and sequence analysis using four regions (rDNA ITS, GAPDH, EF-1 α , and RPB2). The results indicated that the fungus is *Alternaria argyranthemi*. The pathogenicity tests revealed that the species could cause severe leaf spot and blight disease on the host. This is the first report of leaf spot disease on *C. coronarium* caused by *A. argyranthemi* in the world, which is also a new record of *Alternaria* species in China.

ARTICLE HISTORY

Received 4 April 2018
Revised 14 June 2018
Accepted 21 June 2018

KEYWORDS

Asteraceae; leaf spot disease; morphology; multigene phylogenetic analysis

1. Introduction

Chrysanthemum coronarium L. (= *Glebionis coronaria*), commonly called garland chrysanthemum, is annual herbaceous plant belonging to the family Asteraceae and widely distributed in the Mediterranean region and several Asian countries [1]. The green leaves and shoots are appreciated in China, Korea, and Japan as a popular vegetable due to their fragrance, flavor, and nutritional quality [2]. The species is also used as an ornamental and in folk medicine. Furthermore, the antioxidative, antimicrobial, anticancerous, nematicidal, phytotoxic and insecticidal biological properties have been found in the numerous components extracted from *C. coronarium* flowerheads [3–5].

The plant can be infested by various fungi, and there have been reports of *Cercospora* spp., *Entyloma* spp., *Peronospora* spp., *Puccinia* spp. and so on (<https://fungaldatabases/>). Uematsu et al. [6] found that the *Colletotrichum carthami* could induce severe anthracnose on *C. coronarium* var. *spatiosum*. A new species, *Gibellulopsis chrysanthemi* was isolated from rotted leaves of *C. coronarium* and demonstrated to reproduce the natural symptoms in pathogenicity test [7, 8]. Though *Alternaria* leaf spots and blight diseases are commonly found on asteraceous plants, and over 30 *Alternaria* species have been reported [9], there is no report about *Alternaria* spp. causing diseases on *C. coronarium*.

During a survey of large-spored *Alternaria* species in China, a unique species of the genus

Alternaria with solitary conidia was observed on leaf spots of *C. coronarium* from Shiyan, Hubei, China, in April 2017. The study aimed to identify the *Alternaria* species associated with a leaf spot disease of *C. coronarium* based on morphological and molecular characteristics and to reveal their pathogenicity on *C. coronarium*.

2. Symptoms and isolation

The leaf spot or blight symptoms were most severe on the lower part of the plants (Figure 1(A)). The initial symptoms appeared as irregular small brown spots with faint yellowish halos (Figure 1(B)). The typical character was round or irregular necrotic spots surrounded by yellow halo. There was a tiny white spot in the center of each larger necrotic spot. Leaf spots enlarged and their colors darkened as infection progressed, and developed into full blight symptoms (Figure 1(C)). Severely infected leaves fell off.

Specimens with leaf spot symptoms were randomly sampled. Infected tissues were cut into small pieces and incubated on sterile wet paper at 25 °C in the dark. After 1 d, a type of *Alternaria* conidia was observed. A single conidium was picked up under a stereomicroscope and placed on the surface of potato dextrose agar (PDA; Difco, Montreal, Canada). The plates were transferred into an incubator and kept at 25 °C without light. Six pure cultures were placed on PDA slants and in glycerol




Figure 1. Symptoms on *Chrysanthemum coronarium* caused by *Alternaria argyranthemi* in field. (A) Leaf spots in field. (B, C) Different stages of natural leaf spots. (D) Necrotic spots after 3-day-inoculation of spore suspension for the pathogenicity tests. (E) Leaf blight symptoms observed after 6-day-inoculation.

stock solutions and deposited in the Culture Collection of Yangtze University (YZU). One of deposited strains, YZU 171067 was selected for morphological characterization and DNA sequencing.

3. Morphological observation

Isolates were grown on PDA and PCA (containing 20 g potato, 20 g carrot, 20 g agar per liter) at 25 °C for 3–5 d. Mycelial plugs (6 mm diameter) were collected from the edge of colonies and placed onto the center of PDA and PCA plates (90 mm diameter). The cultures were maintained at 25 °C for 7 d in darkness to determine the cultural characteristics. To assess conidial morphology, mycelia of the isolate YZU 171067 were transferred to PCA and incubated at 22 °C for 7 d [9]. However, conidia were rarely found on PCA, but chlamydo-spores (microsclerotia) were abundant. Then, the mycelia were placed on healthy leaves of *C. coronarium* to induce disease symptoms. Diseased leaves were cut into 1 cm length, and placed onto wet filter paper in plates and incubated at 22 °C with light/dark periods of 8 h/16 h for conidial production. After 1 d, the conidiophores were examined using a Leica M205A stereomicroscope (Germany, L 161). Conidia were collected and mounted in lactophenol picric acid solution for further measurement. Conidia ($n = 50$) were randomly chosen and measured under an Olympus BX50 light microscope (Japan, L 165).

Colonies of the isolate YZU 171067 on PDA and PCA medium were strikingly different from each other after 7 d. Colonies on PDA (Figure 2(A)) were 23–30 mm across, irregular effuse, white to dark olive brown, and velvety with thick hyphae covering the center and dark olive brown on the

reverse side. Colonies on PCA (Figure 2(B)) were 33–38 mm across, densely radial, pale gray to dark olive brown with a similar color on the reverse side. Chlamydo-spores were golden brown to dark brown with punctate ornamentation, toruloid or oval, 30–100 × 18–30 μm in the middle of hyaline hyphae (Figure 2(C–G)). However, there were few conidia found on PCA colonies. To measure the conidia, the strain YZU 171067 was stimulated to produce conidia by inoculation onto the host *C. coronarium* and incubation at 22 °C (Figure 2(H)). Conidia produced after 1 d were predominantly solitary, medium yellowish tan to brown, smooth, narrowly ellipsoid to obclavate, 50–120 × 19–30 μm, with 4–9 transversal septa and 0–3 longitudinal septa in some of the transverse divisions, normally with blunt tapered beaks, 10–25 × 4–8 μm (Figure 2(I)). The conidial morphology is in agreement with *Alternaria argyranthemi* described by Simmons [9]. The species produces enormous numbers of chlamydo-spores (microsclerotia) on PCA and V8 media [9], which was also observed during this study. However, few conidia were observed on PCA, which is in contrast to the results of Simmons [9]. Chlamydo-spores are readily identifiable characteristics among the morphologically similar species (*A. sonchi*, *A. brassicae*, and *A. panax*) [9]. This is a particularly key character to identify this species.

4. Molecular analyses

Genomic DNA of the fungus was extracted using the method described by Cenis [10]. The DNA pellet was dissolved in 40 μL of ddH₂O and stored at –20 °C until required for further use. Four target regions were selected for PCR amplification: rDNA ITS (ITS4/ITS5)




Figure 2. Morphology of *Alternaria argyranthemii* from *Chrysanthemum coronarium*. (A) Colony on PDA. (B) Colony on PCA. (C–G) Chlamydospores (microsclerotia) formed on PCA at 22 °C for 7 days (bar = 20 μm). (H) Sporulation patterns on the host plant (bar = 50 μm). (I) Conidia produced on diseased leaves inoculated at 22 °C (bar = 20 μm).

[11], GAPDH (gpd1/gpd2) [12], EF-1 α (EF1-728F/EF1-986R) [13] and RPB2 (RPB2-6F/RPB2-7cR) [14]. PCR amplification was performed in 20 μ L (10 μ L 2 \times Taq PCR StarMix, 7 μ L ddH₂O, 2 μ L DNA, 0.5 μ L of each primer) reactions using an ABI Veriti 96-well Thermal Cycler. Successfully amplified DNA was purified and sequenced with the same primers by BGI (Beijing Genomics Institute). The sequences were compared to GenBank (<https://www.ncbi.nlm.nih.gov/>) using BLASTn, and most highly matching sequences were downloaded. The sequences were adjusted and compared using PHYDIT 3.2 software (Korea, L 281). Four regions were combined, aligned and constructed a neighbor-joining tree by MEGA 7.0 [15], the evolutionary distances were computed using Kimura 2-parameter method and with 1000 bootstrap replicates.

The resulting sequences of the rDNA ITS, GAPDH, EF-1 α and RPB2 region have been deposited in GenBank (Table 1). Sequences of the rDNA ITS, GAPDH and RPB2 showed 100% identity to the type strain of *A. argyranthemii* CBS 116530, with two nucleotides different in the EF-1 α sequence. Their phylogenetic relationship was exhibited in the neighbor-joining phylogram generated from the four gene sequences. The isolates YZU 171067 and CBS 116530 were clustered together supported by 100% bootstrap value (Figure 3). In a taxonomic analysis of the genus *Alternaria*, *A. argyranthemii* fell into a monotypic lineage (containing one species), and was not assigned to any of the 24 sections [16]. The gene sequences of Alt a1 (Alt-a1-for/Alt-a1-rev) [17] and ATPase (ATPDF1/ATPDR1) [18] were also amplified, sequenced and submitted to GenBank

Table 1. Isolates used in this study for the phylogenetic analysis.

Species	Strain	Host	Location	GenBank accession No.			
				rDNA ITS	GAPDH	EF-1 α	RPB2
<i>A. argyranthemii</i>	CBS 116530	<i>Argyranthemum frutescens</i>	New Zealand	KC584181	KC584098	KC584637	KC584378
	YZU 171067	<i>Chrysanthemum coronarium</i>	China	MG647618	MG674139	MG647619	MG647617
<i>A. brassicae</i>	CBS 116528	<i>Brassica oleracea</i>	USA	KC584185	KC584102	KC584641	KC584382
<i>A. botrytis</i>	CBS 197.67	Contaminant	USA	KC584243	KC584168	KC584736	KC584476
<i>A. capsici-annui</i>	CBS 504.74	<i>Capsicum annuum</i>	Unkonw	KC584187	KC584105	KC584644	KC584385
<i>A. panax</i>	CBS 482.81	<i>Aralia racemosa</i>	USA	KC584209	KC584128	KC584675	KC584417
<i>A. penicillata</i>	CBS 116607	<i>Papaver rhoeas</i>	Austria	KC584229	KC584153	KC584706	KC584447
<i>A. sonchi</i>	CBS 119675	<i>Sonchus asper</i>	Canada	KC584220	KC584142	KC584691	KC584433




Figure 3. Neighbor-joining tree generated from *Alternaria argyranthemii* YZU 171067 from *Chrysanthemum coronarium* and its related species based on a combined datasets of rDNA ITS, GAPDH, EF-1 α and RPB2 gene sequences. Bootstrap values were determined using 1000 replicates.

with accession No. MG647616 and MG647615, respectively. The results also supported the identification of YZU 171067 as *A. argyranthemii*.

5. Pathogenicity tests

The chlamydospore of YZU 171067 grown on PCA after 7 d was flushed by ddH₂O, using two layers of sterile gauze to filter out hyphae to obtain a chlamydospore suspension, the suspension was adjusted to 1×10^5 spores/mL. Intact potted *C. coronarium* plants were sprayed with the spore suspension until the leaves were uniformly wet. As a control, plants were inoculated with sterile distilled water. All the plants were incubated at 80–90% relative humidity in separate containers around 25 °C. Disease progress was monitored daily for 7 d. The fungus was re-isolated from symptomatic tissues to address Koch's postulates. The tests were repeated three times.

After inoculation with the chlamydospores suspension, the symptoms on leaves appeared as little brown to black, water-soaked spots after 24 h. It enlarged on leaves with large irregular spots after 3 d. Furthermore, stems were also infected with small long fusiform necrotic symptoms, and the top shoots were severely infected and became blighted (Figure 1(D)). Later, foliar lesions enlarged and coalesced to form large areas of necrosis on leaves and stem, finally resulting in defoliation (Figure 1(E)).

No symptoms were observed in the control leaves inoculated with sterile distilled water. Also, the pathogenicity tests revealed that the fungus is the causal agent causing leaf spot and blight disease on the host. Nishikawa and Nakashima [19] had been tested the pathogenicity of *A. cinerariae* on *C. coronarium* (*Glebionis coronaria*) and found it to be non-pathogenic on this plant.

6. Conclusions

In this study, the fungus isolated from *C. coronarium* was identified as *Alternaria argyranthemii*. E.G. Simmons & C.F. Hill based on the morphology and molecular analysis. To our knowledge, this is the first report of leaf spot disease on *C. coronarium* caused by *A. argyranthemii* in the world, which is also a new record of *Alternaria* species in China.

Acknowledgements

We sincerely thank T. Hsiang for his assistance during the preparation of this manuscript.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

The work was supported by the National Natural Science Foundation of China [31400014] and under the Young Scientist Foundation of Yangtze University [2016cqr08].

References

- [1] Basta A, Pavlovi M, Couladis M, et al. Essential oil composition of the flower heads of *Chrysanthemum coronarium* L from Greece. *Flavour Fragr J.* 2007;22:197–200.
- [2] Chung K, Park Y. Development of an agrobacterium-mediated transformation system for regenerating garland chrysanthemum (*Chrysanthemum coronarium* L.). *J Plant Biol.* 2005;48:136–141.
- [3] Bar-Eyal M, Sharon E, Spiegel Y. Nematicidal activity of *Chrysanthemum coronarium*. *Eur J Plant Pathol.* 2006;114:427–433.

- [4] Haouas D, Cioni PL, Halima-Kamel MB, et al. Chemical composition and bioactivities of three *Chrysanthemum* essential oils against *Tribolium confusum*, (du Val) (Coleoptera: Tenebrionidae). *J Pest Sci.* 2012;85:367–379.
- [5] Hosni K, Hassen I, Sebei H, et al. Secondary metabolites from *Chrysanthemum coronarium* (Garland) flowerheads: chemical composition and biological activities. *Ind Crop Prod.* 2013;44:263–271.
- [6] Uematsu S, Kageyama K, Moriwaki J, et al. *Colletotrichum carthami* comb. nov., an anthracnose pathogen of safflower, garland chrysanthemum and pot marigold, revived by molecular phylogeny with authentic herbarium specimens. *J Gen Plant Pathol.* 2012;78:316–330.
- [7] Kawaradani M, Taguchi K, Okada K, et al. Seedling rot of garland chrysanthemum caused by *Gibellulopsis chrysanthemi* and ecological characters of the causal fungus. *J Gen Plant Pathol.* 2013;79:346–349.
- [8] Hirooka Y, Kawaradani M, Sato T. Description of *Gibellulopsis chrysanthemi* sp. nov. from leaves of garland chrysanthemum. *Mycol Progress.* 2014;13:13–19.
- [9] Simmons EG. *Alternaria: an identification manual.* CBS Biodiversity Series 6. Utrecht: CBS Fungal Biodiversity Centre; 2007.
- [10] Cenis JL. Rapid extraction of fungal DNA for PCR amplification. *Nucleic Acids Res.* 1992;20:2380
- [11] White TJ, Bruns TD, Lee SB, et al. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds.). *PCR protocols: a guide to methods and applications.* San Diego: Academic Press. 1990. p. 315–322.
- [12] Berbee ML, Pirseyedi M, Hubbard S. *Cochliobolus* phylogenetics and the origin of known, highly virulent pathogens, inferred from its and glyceraldehyde-3-phosphate dehydrogenase gene sequences. *Mycologia.* 1999;91:964–977.
- [13] Carbone I, Kohn LM. A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia.* 1999;91:553–556.
- [14] Liu YJ, Whelen S, Hall BD. Phylogenetic relationships among ascomycetes: evidence from an RNA Polymerase II Subunit. *Mol Biol Evol.* 1999;16:1799–1808.
- [15] Kumar S, Stecher G, Tamura K. Mega7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol.* 2016;33:1870.
- [16] Woudenberg JHC, Groenewald JZ, Binder M, et al. *Alternaria* redefined. *Stud Mycol.* 2013;75:171–212.
- [17] Hong SG, Cramer RA, Lawrence CB, et al. Alt a 1 allergen homologs from *Alternaria*, and related taxa: analysis of phylogenetic content and secondary structure. *Fungal Genet Biol.* 2005;42:119–129.
- [18] Lawrence DP, Gannibal PB, Dugan FM, et al. Characterization of *Alternaria* isolates from the *infectoria* species-group and a new taxon from *Arrhenatherum*, *Pseudoalternaria arrhenatheria* sp. nov. *Mycol Progress.* 2014;13:257–276.
- [19] Nishikawa J, Nakashima C. Morphological variation and experimental host range of *Alternaria cinerariae*. *Mycoscience.* 2015;56:141–149.