


RESEARCH ARTICLE

Expected changes in obesity after reformulation to reduce added sugars in beverages: A modeling study

Ana Basto-Abreu¹, Ariela Braverman-Bronstein¹, Dalia Camacho-García-Formentí¹, Rodrigo Zepeda-Tello¹, Barry M. Popkin², Juan Rivera-Dommarco³, Mauricio Hernández-Ávila⁴, Tonatiuh Barrientos-Gutiérrez^{1*}

1 National Institute of Public Health, Center for Population Health Research, Cuernavaca, Mexico, **2** Department of Nutrition, University of North Carolina Gillings School of Global Public Health, Carolina Population Center, Chapel Hill, North Carolina, United States of America, **3** National Institute of Public Health, Cuernavaca, Mexico, **4** University Center of Los Altos, Tapatitlan de Morelos, University of Guadalajara, Guadalajara, Jalisco, Mexico

* tbarrientos@insp.mx



OPEN ACCESS

Citation: Basto-Abreu A, Braverman-Bronstein A, Camacho-García-Formentí D, Zepeda-Tello R, Popkin BM, Rivera-Dommarco J, et al. (2018) Expected changes in obesity after reformulation to reduce added sugars in beverages: A modeling study. *PLoS Med* 15(10): e1002664. <https://doi.org/10.1371/journal.pmed.1002664>

Academic Editor: Ed Gregg, Centers for Disease Control and Prevention, UNITED STATES

Received: April 19, 2018

Accepted: August 31, 2018

Published: October 5, 2018

Copyright: © 2018 Basto-Abreu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Baseline added sugar consumption from SSBs and weight were obtained from the 2012 Mexico National Health and Nutrition Survey (<http://ensanut.insp.mx>). The final dataset after running the Chow and Hall's model is available at the open science framework, available at <https://osf.io/vfcm8/> (DOI: [10.17605/OSF.IO/VFCM8](https://doi.org/10.17605/OSF.IO/VFCM8)). The mathematical model used to estimate the impact on body weight is freely available within an R package version 1.0.0 named

Abstract

Background

Several strategies have been proposed to reduce the intake of added sugars in the population. In Mexico, a 10% sugar-sweetened beverages (SSBs) tax was implemented in 2014, and the implementation of other nutritional policies, such as product reformulation to reduce added sugars, is under discussion. WHO recommends that all individuals consume less than 10% of their total energy intake (TEI) from added sugars. We propose gradually reducing added sugars in SSBs to achieve an average 10% consumption of added sugars in the Mexican population over 10 years and to estimate the expected impact of reformulation in adult body weight and obesity.

Methods and findings

Baseline consumption for added sugars and SSBs, sex, age, socioeconomic status (SES), height, and weight for Mexican adults were obtained from the 2012 Mexico National Health and Nutrition Survey (ENSANUT). On average, 12.6% of the TEI was contributed by added sugars; we defined a 50% reduction in added sugars in SSBs over 10 years as a reformulation target. Using a dynamic weight change model, sugar reductions were translated into individual expected changes in body weight assuming a 43% caloric compensation and a 2-year lag for the full effect of reformulation to occur. Results were stratified by sex, age, and SES. Twelve years after reformulation, the TEI from added sugars is expected to decrease to 10%, assuming no compensation from added sugars; 44% of the population would still be above WHO recommendations, requiring further sugar reductions to food. Body weight could be reduced by 1.3 kg (95% CI -1.4 to -1.2) in the adult population, and obesity could decrease 3.9 percentage points (pp; -12.5% relative to baseline). Our sensitivity analyses suggest that the impact of the intervention could vary from 0.12 kg after 6 months to 1.52 kg in the long term.

'bw' in <https://cran.r-project.org/web/packages/bw/index.html>.

Funding: This project was funded by Bloomberg Philanthropies (<https://www.bloomberg.org/>; JRD received the grant). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: I have read the journal's policy and the authors of this manuscript have the following competing interests: BMP is a member of the Editorial Board of PLOS Medicine.

Abbreviations: BMI, body mass index; CONAPO, Mexico National Population Council; ENSANUT, Mexico National Health and Nutrition Survey; NNS, noncaloric sweetener; pp, percentage points; RCT, randomized controlled trial; SES, socioeconomic status; SSB, sugar-sweetened beverage; TEI, total energy intake.

Conclusions

Reformulation to reduce added sugars in SSBs could produce large reductions in sugar consumption and obesity in the Mexican adult population. This study is limited by the use of a single dietary recall and by data collected in all seasons except summer; still, these limitations should lead to conservative estimates of the reformulation effect. Reformulation success could depend on government enforcement and industry and consumer response, for which further research and evidence are needed.

Author summary

Why was this study done?

- High sugar intake is associated with obesity.
- Adults in Mexico consume 12.6% of their everyday energy intake from added sugars, exceeding WHO recommendations (<10% of total energy intake [TEI]).
- The Mexican government imposed a 10% sugar-sweetened beverages (SSBs) tax to reduce sugar consumption; other interventions such as reformulation and gradual elimination of sugar are being discussed.

What did the researchers do and find?

- We estimated the proportion of added sugars that should be reduced in SSBs to reduce sugar intake to less than 10% of everyday energy intake. We found that we could achieve this goal by halving the sugar content in SSBs.
- Using mathematical models, we translated the reductions in sugar consumption into expected changes in body weight and the prevalence of obesity in adults.
- Allowing for 43% caloric compensation, we estimated that 12 years after starting the sugar reduction in SSBs, the Mexican population would experience an average reduction of 1.3 kg and 12.5% less obesity.
- Larger effects of the sugar regulation will be expected among males, young adults, and the middle socioeconomic status (SES) group.

What do these findings mean?

- Our study shows that a national regulation to reduce added sugars in SSBs could be an effective intervention to reduce sugar intake and curb the obesity epidemic in Mexico.
- The sugar regulation should be considered as a part of an integral strategy by the Mexican government to reduce obesity and avert obesity-related diseases.

Introduction

High sugar intake is a primary risk factor for obesity. Sugar intake has been associated with weight gain [1,2], diabetes, and other diseases [3–5]. Recent calls have been made to reduce sugar intake, in particular targeting sugar-sweetened beverages (SSBs) [6,7]. SSBs provide 39% of daily sugar intake in the United Kingdom [6] and 60% in the United States [8]. In 2011, Mexico was the largest SSB consumer worldwide, with an average of 163 liters per capita [9]; SSBs provided 69% of total added sugar in the population's diet [10]. Adults in Mexico consume 12.6% of their total energy intake (TEI) from added sugars, exceeding the intake recommended by US dietary guidelines (<10% TEI from added sugars) and WHO (<10% TEI from free sugars) [7,11].

A first step towards regulating SSB consumption was taken in January 2014 by the Mexican Government when it imposed a 10% tax on all industrialized SSBs, which produced a 9.7% purchase reduction by 2015 [12]. Still, more interventions are needed to further decrease sugar consumption and produce larger health benefits [13]. In recent years, a movement towards defining maximum concentrations of specific substances (with sodium intake being most prominent) in the food industry has gained momentum [14–16]. Countries have also used more complex grams of sugar and tiered taxation schemes to foster sugar reformulation [17]. For instance, in 2016, the UK government challenged the industry to reduce added sugars in products consumed by children, leading to an 11% reduction in the sugar concentration of SSBs [18]. In this context, setting a maximum limit to added sugars in SSBs is a reasonable intervention to decrease sugar intake and obesity.

Mexico is an ideal site to analyze the potential implementation of a national regulation to decrease added sugar content in beverages. The health secretary has declared obesity and diabetes public health emergencies, calling for decisive actions to curb these epidemics [19], and the National Strategy to Control and Prevent Overweight, Obesity and Diabetes considers the reformulation of beverages and food as well as encourages the gradual elimination of sugar, sodium, and saturated fats [20]. We aimed to estimate the proportion of sugar that should be reduced in SSBs to reduce added sugar intake in the population to an average 10% TEI, assuming no changes in consumption. Using a representative sample of adults, we estimated the expected impact of a sugar regulation of SSBs on weight and prevalence in obesity and overweight over 12 years.

Methods

Data collection

We obtained baseline SSB consumption, weight, and prevalence of overweight and obesity among adults in Mexico using the 2012 National Health and Nutrition Survey (ENSANUT). The ENSANUT 2012 is a cross-sectional, multistage, probabilistic survey representative of the Mexican population that measures the health and nutrition status in the Mexican population. The ENSANUT 2012 surveys 50,528 households, with a total of 96,031 individuals, including children, adolescents, and adults. This survey includes demographic, socioeconomic, nutritional, and health-related data. Nutritional information was collected on a representative subsample of 10,886 individuals (approximately 11%). The National Institute of Public Health estimated the individual sampling weights in ENSANUT needed to represent the age and sex distribution of the Mexican population and to account for survey nonresponse [21].

To collect baseline information from dietary intake and body mass index (BMI), we used the food 24-hour recall and anthropometry databases, including adults 20 years of age and older. Extreme values of energy intake (more than ± 3 SDs $\ln\left(\frac{\text{total energy intake}}{\text{total energy requirements}}\right)$, $n = 125$) [22], extreme values of SSB consumption ($>3,000$ kcal; $n = 1$), children <1 year, breastfeeding

children, and pregnant and lactating women were excluded ($n = 665$) because energy intake varies during these periods. Participants younger than 20 years ($n = 6,926$) and those with missing data for weight and height ($n = 164$) were excluded from the analysis.

Dietary intake

We collected added sugar consumption (from SSBs and other products) using the 24-hour diet recall from ENSANUT for the adult population. SSBs included all industrialized nonalcoholic beverages taxed in 2014 in Mexico (a list of beverages is available in section 1.1 of Supporting Information [S1 Methods](#)). The food composition table compiled to analyze ENSANUT's 24-hour diet recall does not include added sugar. This nutrient was previously estimated by Sánchez-Pimienta and colleagues [10]. Briefly, added sugar was set to zero for all foods without sugar or foods for which all sugar is intrinsic (such as fruits, unprocessed cereals, or legumes); added sugar was set to the total sugar content for foods for which all sugar is added (such as sodas, confectionary, and sweeteners). Added sugar was estimated as a share of the total sugar content for all other foods having a mix of intrinsic and added sugar. This share was estimated based on either lactose content or similar foods that had no added sugars, for instance, comparing the sugar content from 100% juice versus industrialized juice (understanding industrialized juices as beverages that contain some proportion of fruit juices but with added ingredients such as sugar, additives, or conservatives).

BMI

Weight and height were directly measured by trained interviewers during home visits using standardized examination procedures [21]. BMI was calculated using weight (kg) divided by squared height (m^2). We classified each adult following WHO guidelines, but given the low prevalence of underweight (0.8% with $<18.5 \text{ kg}/m^2$), we collapsed this category with the normal range ($BMI \geq 18.5$ and $< 25 \text{ kg}/m^2$); overweight ($BMI \geq 25$ and $< 30 \text{ kg}/m^2$) and obesity ($BMI \geq 30 \text{ kg}/m^2$) were reported as usual.

Stratification variables

SSB consumption varies according to sex, age, and socioeconomic status (SES) [23]. Continuous age was introduced in the weight change model, yet we categorized it into young adults 20 to 40 years old, adults 40 to 60 years old, and elderly (age 60 and older) to facilitate the presentation of results. SES was constructed for the ENSANUT 2012 using demographic and socioeconomic characteristics of the household head (years of education, sex, and employment status), sociodemographic structure (individuals living in the household, individuals employed, adults employed, children working, and index of economic dependence), household characteristics (number of rooms, exclusive kitchen, bathroom, and type of fuel, among others), household assets (TV, microwave, and computer, among others), family consumption patterns (housing rent and logarithm of household expenditures), and characteristics of the place of residence using the 2010 marginalization index from the National Council on Population, which considers education, housing, and income at the local level [24]. For this analysis, we used three categories of SES (low, middle, and high), which represent SES tertiles of the adult population in the anthropometric ENSANUT database.

Reduction of the amount of added sugar

To estimate the amount of added sugar that should be reduced in SSBs to reduce the population intake to less than 10% TEI, we calculated the total added sugar consumption from all

dietary sources for each ENSANUT participant. At baseline, adults in Mexico consumed on average 12.6% of TEI from added sugar, exceeding WHO and US guideline recommendations [7,11]. While the WHO recommendation is for all individuals, we estimated the sugar reduction required to achieve an average sugar consumption of 10% at the population level, which implies that 44% of individuals would still be above the recommended 10% TEI (assuming no compensation for added sugars).

To calculate the average reduction in sugar from SSBs required to reach the average 10% TEI goal, we first estimated the individual maximum added sugar consumption from SSBs, as follows (Eq 1):

$$10\% = \frac{SSB_{max}^i + others^i}{TEI_{init}^i + (SSB_{max}^i - SSB_{init}^i)}, \tag{1}$$

where “others” is the consumption of added sugars from sources other than SSBs, TEI_{init} is the TEI at baseline, and SSB_{init} is the added sugars from SSBs. If current consumption from SSBs was lower than the maximum calculated ($SSB_{init} < SSB_{max}$), no changes were made $\Delta_{change} = 0$; if $SSB_{init} > SSB_{max}$, the change in added sugars was $\Delta_{change} = SSB_{init} - SSB_{max}$. If added sugar consumption from other sources was above 10% TEI, then we reduced all sugar from SSBs so that $\Delta_{change} = -SSB_{init}$. The average proportion of added sugars in SSBs was calculated based on the individual change, $\sum_{i=1}^N \omega_i \cdot prop_i = \sum_{i=1}^N \omega_i \cdot \frac{\Delta_{change}^i}{SSB_{init}^i} = 52.2\%$, which we rounded down to 50%. A detailed description of how we estimated the target reduction in added sugars can be found in Supporting Information [S1 Methods](#), section 2.1 and 2.2.

To halve the sugar content, we proposed a national regulation to gradually reduce added sugars in SSBs to reach the goal over a 10-year period. We set a long implementation period with small yearly sugar reductions to allow for a gradual adaption to the new level of sweetness. The first-year reduction was estimated as follows (Eq 2):

$$Reduction(y_1) = 1 - (1 - Reduction(y_k))^{1/k} \tag{2}$$

where k denotes the number of years the regulation takes place. Considering $Reduction(y_{10}) = 0.5 = 50\%$. We obtained a first-year reduction of $Reduction(y_1) 0.067 = 6.7\%$. The accumulated reductions in the following years were estimated as follows (Eq 3):

$$Reduction(y_k) = 1 - (1 - Reduction(y_1))^k, \tag{3}$$

where $Reduction(y_k)$ is the estimated reduction for each year $k = 1, 2, \dots, 10$. Hence, the reduction at y_2 would be 12.9%, at y_3 would be 18.8%, and so on to achieve a 50% reduction by y_{10} . A detailed description of the yearly reduction estimation can be found in Supporting Information [S1 Methods](#) (section 2.3).

Change in energy intake

To estimate the daily change in caloric intake, $\Delta Caloric\ intake$ attributable to the sugar reduction in SSBs, we multiplied the individual added sugar intake from SSBs by the cumulative percent reduction in sugar content for that year considering compensation, as follows (Eq 4):

$$\Delta Caloric\ intake(y_k) = Added\ sugar\ SSBs \times Reduction(y_k) \times \left(1 - \frac{Compensation}{100}\right). \tag{4}$$

Compensation is understood as the adjustments in energy intake that follow to the reduction induced by the intervention (in this case, the reduction in added sugars in SSBs). We assumed a 43% compensation based on a recent meta-analysis that analyzed the substitution

of added sugars by low-caloric artificial sweeteners [25]. Under 43% compensation, an individual reducing 50 kcal/day would compensate 21.5 kcal through sources other than added sugars, resulting in a net reduction of 28.5 kcal/day. The average change in energy intake was stratified by sex, age group, and SES. Aggregated changes in energy intake were estimated using Stata/MP version 14.1 [26] taking into account the complex survey design (see Supporting Information [S1 Methods](#), section 1.2).

Change in body weight and obesity prevalence

The change in body weight was estimated using the dynamic weight change model proposed by Chow and Hall [27], implemented individual by individual for all participants in the ENSANUT. This model accounts for the dynamic physiological adaptation that occurs following weight loss, and it has been validated with experimental data [27–29]. Briefly, the model uses a system of differential equations to predict the expected body weight change of an individual. It is estimated as the sum of extracellular fluid, glycogen, fat mass, and lean tissue, dependent on sodium and energy intake, energy expenditure, and other characteristics such as sex, age, height, initial body weight, and physical activity level [27]. Our model assumes steady-state for body weight and no changes in physical activity nor in energy intake (apart from the sugar regulation) during the modeling period. A step-by-step explanation of the implementation of the model is provided in [S1 Methods](#), section 3.

Using the expected reduction in energy intake, we implemented Chow and Hall’s model to each adult in ENSANUT 2012 to obtain the expected reduction in weight after 12 years (considering a 2-year time lag for caloric changes to influence weight). Final weights were transformed to BMI and BMI categories (normal, overweight, obese) to estimate prevalence changes. Assuming that the sugar regulation would be implemented in 2020, we estimated the number of prevented overweight and obesity cases by 2032 by multiplying the change in prevalence in percentage points (pp) by the expected adult population in 2032, using population projections published by the National Population Council (CONAPO) [30]. The dynamic body weight model was programmed using R statistical software version 3.4.1 (2017-06-30) [31] using the *bw* package [31] and the *Rcpp* package [32–35]. Aggregated changes in body weight and BMI status by sex, age group, and SES were estimated using Stata/MP 14.1 [26], taking into account the complex survey design (see Supporting Information [S1 Methods](#), section 1.2). The final dataset used is available at <https://osf.io/vfcm8/> (doi: [10.17605/OSF.IO/VFCM8](https://doi.org/10.17605/OSF.IO/VFCM8)), and the variables are explained in [S1 Methods](#) (section 1).

Sensitivity analyses

In the main scenario, we assumed 43% energy compensation based on a recent meta-analysis [25]. This was a request from the peer review process. In our first analysis, we had assumed no energy compensation given the discordant heterogeneity of results included in the meta-analysis and other sources suggesting minimal or zero dietary compensation for liquids [36–38]. Results with 0% compensation were included as a sensitivity analysis (Table A in [S1 Results](#)). We also conducted a sensitivity analysis to estimate all possible compensation scenarios (from 0% to 100%) and different added sugar reduction targets (from 0% to 100%) in a 10-year period following the “decreasing yearly reduction” (Eq 3). The overall reduction for each added sugar target and each compensation level was estimated as follows (Eq 5):

$$Reduction(y_1) = Target \times \left(1 - \frac{Compensation}{100} \right). \quad (5)$$

where *Target* represents the maximum reduction achieved with the regulation. The change in

energy intake was estimated using Eq 4 and applied it to the individual weight change model. For additional information, including a matrix summarizing weight reductions, see Supporting Information S1 Methods (section 4).

For the main analysis, the regulation set a 6.7% added sugar reduction in the first year, followed by lower percent reductions for the subsequent years. We called this scenario “decreasing yearly reduction.” Two other implementation scenarios were estimated with a 50% reduction of added sugar gradually achieved over 10 years, yet we decided to present results after 12 years because, at that point, the three scenarios converged. The “constant yearly reduction” scenario reduces added sugar by 5% yearly from the baseline content. The “increasing yearly reduction” scenario considers the same percent changes as the “decreasing yearly reduction” but in reverse order: starting with a small decrease ($y_1 = 3.6\%$) and augmenting the percent reduction ($y_{10} = 6.7\%$) to reach 50% by year 10. This last scenario is given by Eq 6:

$$Reduction(y_k) = (1 - Reduction(y_{10}))^{\frac{10-k}{k}} \times Reduction(y_{10}) \tag{6}$$

See Supporting Information S1 Methods section 2.3 for more details.

Results

Table 1 shows baseline characteristics of the Mexican adults. The final sample included 3,005 individuals, which expanded to 64,885,715 adults. The average added sugar consumption from all sources was 244.9 kcal/day (12.6%) and from SSBs was 101.1 kcal/day (5.1% TEI). Men consumed more added sugar in total and from SSBs. Young adults, 20 to 39 years old, were the largest consumers of added sugar in total and from SSBs in comparison with other age groups. Smaller differences in added sugar consumption were observed across SES groups; in general, the low SES group had the lowest consumption of total and SSB added sugars.

Table 2 shows the estimated results of halving the sugar content in SSBs. Twelve years after implementation, assuming 43% compensation, the energy intake could be reduced by 28.8 kcal/day (95% CI 26.6–31.1), which translates into a 1.3 kg (95% CI 1.2–1.4) reduction in weight and a 0.5 kg/m² (95% CI 0.5–0.6) decrease in BMI. Stratifying by sex, the sugar

Table 1. Sample characteristics, added-sugar intake (total and from SSBs) in and their contribution to the %TEI for the adult population in Mexico before regulation.

	Proportion of individuals in the sample (%)	Total added-sugar intake (kcal/day) (95% CI)	%TEI	Added-sugar intake from SSBs (kcal/day) (95% CI)	%TEI
Overall		244.9 (232.9–256.9)	12.6	101.1 (93.3–109.0)	5.1
Sex					
Male	47.7	278.9 (259.9–297.9)	12.9	128.0 (115.4–140.6)	6.1
Female	52.3	213.9 (199.6–228.2)	12.2	76.7 (67.4–86.0)	4.2
Age group					
20–39	44.6	285.7 (264.6–306.8)	13.8	130.8 (117.5–144.2)	6.4
40–59	36.8	235.7 (217.4–254.1)	12.2	90.3 (78.1–102.4)	4.5
60+	18.6	165.2 (148.2–182.2)	10.3	51.4 (39.3–63.5)	3.2
SES					
Low	29.7	213.1 (196.0–230.2)	11.3	84.5 (71.8–97.2)	4.5
Middle	30.1	255.0 (233.1–277.0)	12.7	118.0 (102.6–133.3)	5.7
High	40.2	260.7 (239.3–282.2)	13.4	100.8 (87.6–114.0)	5.2

Abbreviations: %TEI, percent of total energy intake; SES, socioeconomic status; SSB, sugar-sweetened beverage.

<https://doi.org/10.1371/journal.pmed.1002664.t001>

Table 2. Predicted reduction in added sugars, energy intake, body weight, and BMI from a 50% gradual reduction of added sugars to SSBs, considering 43% compensation after 12 years.

	Added sugars (kcal/day) (95% CI)	Energy intake* (kcal/day) (95% CI)	Weight (kg) (95% CI)	BMI (kg/m ²) (95% CI)
Average	-50.6 (-54.5 to -46.6)	-28.8 (-31.1 to -26.6)	-1.3 (-1.4 to -1.2)	-0.5 (-0.6 to -0.5)
Sex				
Male	-64.0 (-70.3 to -57.7)	-36.5 (-40.1 to -32.9)	-1.6 (-1.7 to -1.4)	-0.6 (-0.6 to -0.5)
Female	-38.3 (-43.0 to -33.7)	-21.8 (-24.5 to -19.2)	-1.1 (-1.2 to -0.9)	-0.5 (-0.5 to -0.4)
Age group				
20–39	-65.4 (-72.1 to -58.7)	-37.3 (-41.1 to -33.5)	-1.6 (-1.8 to -1.5)	-0.6 (-0.7 to -0.6)
40–59	-45.1 (-51.2 to -39.1)	-25.7 (-29.2 to -22.3)	-1.2 (-1.4 to -1.1)	-0.5 (-0.6 to -0.4)
60+	-25.7 (-31.8 to -19.7)	-14.7 (-18.1 to -11.2)	-0.7 (-0.9 to -0.5)	-0.3 (-0.3 to -0.2)
SES				
Low	-42.2 (-48.6 to -35.9)	-24.1 (-27.7 to -20.5)	-1.1 (-1.2 to -0.9)	-0.4 (-0.5 to -0.4)
Middle	-59.0 (-66.7 to -51.3)	-33.6 (-38.0 to -29.2)	-1.5 (-1.7 to -1.3)	-0.6 (-0.7 to -0.5)
High	-50.4 (-57.0 to -43.8)	-28.7 (-32.5 to -25.0)	-1.3 (-1.5 to -1.2)	-0.5 (-0.6 to -0.4)

*Net energetic change assuming 43% compensation for sources other than added sugars.

Abbreviations: BMI, body mass index; SES, socioeconomic status; SSB, sugar-sweetened beverage.

<https://doi.org/10.1371/journal.pmed.1002664.t002>

regulation is estimated to have larger weight decreases in males (1.6 kg; 95% CI 1.4–1.7) than in females (1.1 kg; 95% CI 0.9–1.2). Younger adults, 20 to 39 years old, experience larger decreases in weight (1.6 kg; 95% CI 1.5–1.8) compared to older adults (0.7 kg; 95% CI 0.5–0.9). Stratifying by SES, people in the middle SES group are expected to experience larger weight decreases (1.5 kg; 95% CI 1.3–1.7) compared to low SES (1.1 kg; 95% CI 0.9–1.2) and high SES (1.3 kg; 95% CI 1.2–1.5).

Table 3 shows the expected change in the prevalence and prevented cases of overweight and obesity 12 years after the implementation of the sugar regulation. The regulation could increase the prevalence of people with normal BMI by 3.8 pp (95% CI 2.8–4.9), which translates into 3.5 million individuals more with normal weight by the year 2032, along with a reduction in obesity of 3.9 pp (95% CI 2.6–5.1), which translates into 3.5 million less individuals in the obesity category by 2032. Larger reductions in obesity are expected for adults 20 to 39 years old (4.7 pp; 95% CI 2.8–6.6) than for those 40 to 59 years old (4.1 pp; 95% CI 2.0–6.3) or older adults (1.3 pp; 95% CI 0.4–2.3). People in the low SES group are expected to experience the highest decrease in obesity (4.8 pp; 95% CI 2.1–7.6) followed by the high SES (3.7 pp; 95% CI 2.0–5.5) and middle SES (3.1 pp; 95% CI 1.4–4.8) groups. However, the middle SES group could also experience the largest increases in the prevalence of normal weight (4.3 pp; 95% CI 2.2–6.5) followed by high SES (3.8 pp; 95% CI 2.3–5.4) and low SES (3.4 pp; 95% CI 1.8–5.0).

Fig 1 shows a sensitivity analysis measuring the impact of several sugar reduction targets, in combination with different compensation rates. Our main analysis considers a 50% reduction in SSBs with 43% compensation, which results in a 1.3 kg reduction after 12 years; this effect could be 0 kg with 100% compensation or an increase of up to 2.3 kg assuming no compensation. Considering no compensation, or compensation with noncaloric beverages, reducing sugar content in SSBs from 0% to 100% could reduce body weight by between 0 and 4.5 kg, respectively.

Discussion

We estimated the impact of reformulating SSBs to reduce added sugars and achieve an average 10% of the TEI in the Mexican adult population. This goal would be achieved if added sugars in SSBs were reduced by 50%. Assuming that halving the sugar content takes up to 10 years

Table 3. Predicted change in absolute and relative prevalence and absolute number of individuals for normal, overweight, and obesity after 12 years of the implementation of a 50% gradual reduction of added sugar in SSBs, considering a 43% compensation (data projected to year 2032).

		Baseline prevalence (%; 95% CI)	Change in prevalence (absolute pp; 95% CI)	Percent change in prevalence (% baseline; 95% CI)	Change in number of individuals (in millions; 95% CI)
Average	Normal	30.4 (27.9 to 33.1)	3.8 (2.8 to 4.9)	12.6 (9.0 to 16.3)	3.5 (2.6 to 4.5)
	Overweight	38.7 (36.1 to 41.4)	0.0 (-1.6 to 1.7)	0.0 (-4.2 to 4.3)	0.0 (-1.5 to 1.5)
	Obesity	30.9 (28.5 to 33.3)	-3.9 (-5.1 to -2.6)	-12.5 (-16.2 to -8.8)	-3.5 (-4.7 to -2.4)
Sex					
Male	Normal	36.4 (32.3 to 40.7)	5.0 (3.5 to 6.5)	13.7 (9.0 to 18.3)	2.2 (1.5 to 2.8)
	Overweight	39.5 (35.4 to 43.7)	-1.0 (-3.5 to 1.4)	-2.6 (-8.7 to 3.6)	-0.4 (-1.5 to 0.6)
	Obesity	24.1 (20.9 to 27.7)	-4.0 (-5.8 to -2.1)	-16.4 (-23.4 to -9.4)	-1.7 (-2.5 to -0.9)
Female	Normal	25.0 (22.2 to 28.0)	2.8 (1.5 to 4.2)	11.3 (5.6 to 17.0)	1.4 (0.7 to 2.0)
	Overweight	38.0 (34.4 to 41.7)	1.0 (-1.2 to 3.1)	2.5 (-3.3 to 8.3)	0.5 (-0.6 to 1.5)
	Obesity	37.0 (33.7 to 40.4)	-3.8 (-5.4 to -2.2)	-10.2 (-14.3 to -6.1)	-1.8 (-2.6 to -1.1)
Age group					
20–39	Normal	39.4 (35.4 to 43.6)	4.9 (3.0 to 6.9)	12.5 (7.2 to 17.9)	2.0 (1.2 to 2.8)
	Overweight	35.2 (31.2 to 39.5)	-0.3 (-3.1 to 2.6)	-0.7 (-8.7 to 7.3)	-0.1 (-1.3 to 1.1)
	Obesity	25.3 (22.2 to 28.8)	-4.7 (-6.6 to -2.8)	-18.5 (-25.5 to -11.5)	-1.9 (-2.7 to -1.1)
40–59	Normal	19.1 (15.9 to 22.9)	3.6 (2.1 to 5.1)	18.8 (9.9 to 27.6)	1.2 (0.7 to 1.7)
	Overweight	41.2 (36.8 to 45.7)	0.5 (-2.2 to 3.3)	1.3 (-5.3 to 7.9)	0.2 (-0.7 to 1.1)
	Obesity	39.7 (35.4 to 44.1)	-4.1 (-6.3 to -2.0)	-10.4 (-15.7 to -5.2)	-1.4 (-2.1 to -0.7)
60+	Normal	31.3 (26.5 to 36.4)	1.7 (0.7 to 2.7)	5.5 (2.0 to 8.9)	0.3 (0.1 to 0.5)
	Overweight	42.1 (37.0 to 47.4)	-0.4 (-1.8 to 1.0)	-0.9 (-4.3 to 2.5)	-0.1 (-0.3 to 0.2)
	Obesity	26.6 (22.5 to 31.3)	-1.3 (-2.3 to -0.4)	-5.0 (-8.6 to -1.3)	-0.2 (-0.4 to -0.1)
SES					
Low	Normal	37.9 (33.4 to 42.6)	3.4 (1.8 to 5.0)	8.9 (4.4 to 13.4)	0.9 (0.5 to 1.3)
	Overweight	35.3 (31.1 to 39.7)	1.4 (-1.8 to 4.7)	4.1 (-5.3 to 13.4)	0.4 (-0.5 to 1.3)
	Obesity	26.8 (22.9 to 31.1)	-4.8 (-7.6 to -2.1)	-18.0 (-26.9 to -9.0)	-1.3 (-2.1 to -0.6)
Middle	Normal	30.6 (26.4 to 35.1)	4.3 (2.2 to 6.5)	14.2 (6.4 to 22.0)	1.2 (0.6 to 1.8)
	Overweight	37.1 (33.0 to 41.4)	-1.2 (-4.1 to 1.6)	-3.3 (-10.9 to 4.2)	-0.3 (-1.1 to 0.4)
	Obesity	32.3 (28.4 to 36.5)	-3.1 (-4.8 to -1.4)	-9.6 (-14.7 to -4.5)	-0.9 (-1.3 to -0.4)
High	Normal	24.8 (20.6 to 29.6)	3.8 (2.3 to 5.4)	15.4 (8.2 to 22.5)	1.4 (0.8 to 2.0)
	Overweight	42.4 (37.7 to 47.3)	-0.1 (-2.5 to 2.3)	-0.2 (-5.9 to 5.5)	0.0 (-0.9 to 0.9)
	Obesity	32.7 (28.5 to 37.2)	-3.7 (-5.5 to -2.0)	-11.4 (-16.5 to -6.3)	-1.4 (-2.0 to -0.7)

pp to reflect the absolute difference.

“Normal” includes underweight <18.5 kg/m².

Abbreviations: pp, percentage points; SES, socioeconomic status; SSB, sugar-sweetened beverage.

<https://doi.org/10.1371/journal.pmed.1002664.t003>

and a 43% caloric compensation, this regulation would produce a reduction of 1.3 kg of body weight and 0.5 kg/m² of BMI, reducing by 3.9 pp the prevalence of obesity (12.6% relative to the baseline). Larger decreases will be expected among males, young adults, and the middle SES group.

Regulating added sugars through national reformulation has been described as cost-effective, in some cases having a larger impact than reducing portion sizes [17,39,40]. Several countries have forecasted the long-term impact of reformulating SSBs to reduce sugar content. In Australia, a mandatory reduction of added sugar by 30% in SSBs (assuming no compensation) was estimated to produce a 1.39 kg reduction in weight (0.56 kg/m² of BMI) over a lifetime [39]. In the UK, Ma and colleagues estimated that reducing 40% of added sugar in SSBs

been validated against experimental data and can be implemented using individual-level or aggregated data; however, other models to estimate weight change are available. Christiansen and Garby proposed an individual-based model that expresses body weight as a function of energy intake and physical activity level and can also be implemented at an individual level [42]. Briggs and colleagues used experimental data from RCTs to estimate the potential impact of sugar reformulation in the UK using the comparative assessment framework [17]. To reflect the heterogeneity of models, we conducted a sensitivity analysis following Briggs methodology, a modified version of Briggs using effect estimates of SSB consumption on body weight in Mexican women [43] and Christiansen and Garby's model [42]. The estimates obtained using comparative risk assessment are smaller than our results (see Table C in [S1 Results](#)): under Briggs approach, the expected body weight reduction would reach 0.12 kg after 6 months, increasing to 0.44 kg at 2 years if the association of SSB consumption and weight in Mexican women is used as the effect estimate; in contrast, Christiansen and Garby's model with 43% compensation produces a higher estimate than ours (1.52 kg compared to 1.31 kg). The comparison across models is difficult because they consider different time frames; still, we could consider Briggs to be the lowest boundary and Christiansen the highest.

Implementation process

Sugar reductions to SSBs have been previously proposed, although the mechanism for implementation and enforcement has not been clearly laid out other than the UK study on reformulation by use of a tiered sugar tax [17]. Sugar reduction could be part of a self-regulatory plan or a national regulation. Industry self-regulation has proven to be ineffective because its main focus is usually restricted to improve public perception: an image of concern and care is created, while at the same time, standards and enforcement are relaxed [44]. In contrast, a regulation enforced by the government has proven to be effective to limit food additives. For instance, trans fat bans in New York restaurants led to a reduction in trans fat consumption [45]. Also, government programs to reduce salt content in industrialized products produced significant decreases in salt consumption in China, Finland, France, Ireland, South Africa, and the UK [46,47]. The UK, Ireland, and South Africa are three countries that are using tiered and/or tiered plus grams of sugar tax structures to promote reformulation of SSBs [17,48]. Consequently, we consider that a national regulation to halve the sugar content over 10 years is feasible and necessary to reduce obesity in Mexico and should be promoted along with other simultaneous interventions, such as food taxes, information campaigns, improvement of food systems, and strengthening of product labeling [13,49]. In our opinion, sugar reduction in SSBs should be implemented as a national regulation to avoid pressure from industry, provide compliance, provide monitoring, and establish independent evaluations of effectiveness.

To be successful a state-enforced sugar reduction requires a careful analysis of the implementation process, taking into the account the consumer's preference and the impact on flavor. Recent evidence suggests that small reductions in added sugar content in SSBs have no effect on acceptance or pleasantness and could further increase the preference for less sweetened food or beverages [50,51]. Furthermore, experimental evidence has shown low ability of consumers to discriminate across SSB brands or type of sweetener [52,53]. In our models, the decreasing reduction scenario, which sets larger reductions on sugar content at the beginning, had the highest impact on weight at year 10; however, this scenario could imply a stronger change in flavor. The other two scenarios (Fig A in [S1 Results](#)) imply smaller sugar reductions at the beginning and could be more easily accepted by consumers, achieving the same weight reductions than the decreasing reduction scenario by year 12. Further research is needed to

estimate the effect of lowering added sugars on product preference, acceptance, and consumption to inform the best implementation scenario and improve the odds of success.

Finally, a state-enforced sugar regulation must carefully weigh the use of noncaloric sweeteners (NNS). Advocates of NNSs propose their use to maintain the same level of sweetness while decreasing caloric content. In this study, we assumed no substitution by NNS given reported unpleasant taste or a slower-onset sweetness in comparison with sugar [51]. Also, in recent complex research, cohort studies suggest that NNS could increase caloric intake and body weight, as well as hypertension, metabolic syndrome, type 2 diabetes, and cardiovascular events [54,55]. Studies that control for diet modification and RCTs suggest no effect, leading dietary guideline committees in the US and UK and other organizations to continue to recommend the intake of NNSs [11,55–58]. Other concerns relate to NNS addictiveness as a result of an increase in sugar tolerance and the overstimulation of the sugar receptors, yet RCT results suggest no effect [59–62]. NNS addictiveness results are often taken from mice models, which might be an inappropriate comparison because NNS concentration by mice per body weight have about 100 times the sweetness preference of humans. Although the negative evidence of NNS is still insufficient, ideally, we would want the sweetness threshold to be reduced and—where potable water is available—have people shift to less sweetened beverages and nonsweetened beverages.

Strengths and limitations

Some limitations for input data must be acknowledged. First, both BMI and sugar intake from SSBs come from nationally representative data. We used a single 24-hour dietary recall to estimate sugar intake at the population level and subgroups; the 24-hour recall is limited to representing usual individual intake as it fails to capture day-to-day variation, yet it is adequate to estimate energy means at an aggregated level [63]. Also, the 24-hour recall could underestimate sugar intake (a source of variability that is not included in the model), particularly in individuals with higher BMI; however, if anything, this error would lead to conservative estimates of the regulation effect [64]. ENSANUT 2012 obtained dietary recall from all seasons, except for summer (June to September), which ignores the variations in consumption observed in those months. In summer, SSB consumption is higher, which could lead to the underestimation of the total effect of the sugar regulation [65]. Finally, we excluded 164 observations from the ENSANUT sample due to missing anthropometric data. Excluded observations were older and more men but had comparable total and SSB-related caloric intakes to the analytical sample; given the small proportion of missing observations and the similar caloric intake, we assume that these exclusions will not fundamentally change our estimates.

Some limitations and strengths for the modeling process must be acknowledged. In our main scenario, we assume steady-state conditions along the 12-year simulation and a 43% compensation, but under the assumption that the industry would not interfere with the caloric reduction derived from sugar reductions. However, the industry could find ways to circumvent the regulation, such as increasing the black market, in which case our primary scenario would overestimate the expected weight decreases. A possible downside of the SSB regulation would be the increase of the prevalence of underweight (BMI < 18.5 kg/m²). We included underweight in the normal BMI group due to its small percentage and to facilitate the presentation of results; in Mexico, a very small percentage of individuals are underweight (0.8%), but following reformulation, it could increase to 1.7%. Low body weight in adults can be produced by undernutrition but also by medical conditions such as cancer, diabetes, or anorexia; the inclusion of these conditions is beyond the scope of our paper, although the potential increase in underweight following reformulation should be considered to provide adequate detection

and medical care to underweight individuals. Modeling the impact of reformulation in children and adolescents is important, given that SSBs contribute to a large proportion of their energy consumption [10]. However, weight modeling in these age groups is challenging because it requires additional energy functions to take into account growth [66]; unfortunately, no such functions are available for Mexican children and adolescents. Finally, Hall's model does not provide CIs to the estimated weight change, therefore our CIs only consider the sources of error captured by the survey data. However, this model has been previously validated and used to estimate the impact of SSB tax in Mexico [67]; also, it has the advantage of using individual-level data, which allows for result stratification.

Conclusions

Increasing awareness about excessive sugar contents and implementing effective policies to decrease its consumption is paramount to prevent obesity. Considering the variables involved, providing a precise prediction of the impact of sugar reformulation in beverages is difficult; our study suggests that reformulation could reduce obesity by 12.5%, yet this result could vary depending on implementation conditions, caloric substitution, and industry response. National regulation of sugar content is an effective and feasible action that could be implemented in Mexico and other countries, to gradually adapt the population taste to lower sugar intake and promote healthier diets.

Supporting information

S1 Methods. Supporting information for the input data, estimation of the sugar reduction target, implementation of the weight change model, and sensitivity analysis.
(PDF)

S1 Results. Supporting information for the impact of sugar reduction program using different assumptions.
(DOCX)

Acknowledgments

We are grateful to Carolina Batis from the National Institute of Public Health, Mexico for her nutritional insight and support in managing the 24-hour dietary recall data from ENSANUT 2012. We are also thankful to D  sir  e Vida  a from the National Institute of Public Health, Mexico for her valuable comments on and feedback of the study.

Author Contributions

Conceptualization: Ana Basto-Abreu, Mauricio Hern  ndez-  vila, Tonatiuh Barrientos-Guti  rrez.

Data curation: Ana Basto-Abreu, Tonatiuh Barrientos-Guti  rrez.

Formal analysis: Ana Basto-Abreu, Ariela Braverman-Bronstein, Rodrigo Zepeda-Tello, Tonatiuh Barrientos-Guti  rrez.

Funding acquisition: Barry M. Popkin, Juan Rivera-Dommarco, Mauricio Hern  ndez-  vila, Tonatiuh Barrientos-Guti  rrez.

Investigation: Ana Basto-Abreu, Tonatiuh Barrientos-Guti  rrez.

Methodology: Ana Basto-Abreu, Ariela Braverman-Bronstein, Dalia Camacho-García-Formentí, Rodrigo Zepeda-Tello, Barry M. Popkin, Juan Rivera-Dommarco, Tonatiuh Barrientos-Gutiérrez.

Project administration: Tonatiuh Barrientos-Gutiérrez.

Resources: Juan Rivera-Dommarco, Tonatiuh Barrientos-Gutiérrez.

Software: Ana Basto-Abreu, Dalia Camacho-García-Formentí, Rodrigo Zepeda-Tello, Tonatiuh Barrientos-Gutiérrez.

Supervision: Ana Basto-Abreu, Rodrigo Zepeda-Tello, Barry M. Popkin, Juan Rivera-Dommarco, Mauricio Hernández-Ávila, Tonatiuh Barrientos-Gutiérrez.

Validation: Ana Basto-Abreu, Dalia Camacho-García-Formentí, Rodrigo Zepeda-Tello, Tonatiuh Barrientos-Gutiérrez.

Visualization: Ana Basto-Abreu, Tonatiuh Barrientos-Gutiérrez.

Writing – original draft: Ana Basto-Abreu, Ariela Braverman-Bronstein, Tonatiuh Barrientos-Gutiérrez.

Writing – review & editing: Ana Basto-Abreu, Ariela Braverman-Bronstein, Dalia Camacho-García-Formentí, Rodrigo Zepeda-Tello, Barry M. Popkin, Juan Rivera-Dommarco, Mauricio Hernández-Ávila, Tonatiuh Barrientos-Gutiérrez.

References

1. Te Morenga L, Mallard S, Mann J. Dietary sugars and body weight: systematic review and meta-analyses of randomised controlled trials and cohort studies. *Bmj*. 2012; 346: e7492. <https://doi.org/10.1136/bmj.e7492> PMID: 23321486
2. Malik VS, Pan A, Willett WC, Hu FB. Sugar-sweetened beverages and weight gain in children and adults: a systematic review and meta-analysis. *Am J Clin Nutr*. 2013; <https://doi.org/10.3945/ajcn.113.058362> PMID: 23966427
3. Te Morenga LA, Howatson AJ, Jones RM, Mann J. Dietary sugars and cardiometabolic risk: systematic review and meta-analyses of randomized controlled trials of the effects on blood pressure and lipids. *Am J Clin Nutr*. 2014; 100: 65–79. <https://doi.org/10.3945/ajcn.113.081521> PMID: 24808490
4. Malik VS, Popkin BM, Bray GA, Després J-P, Hu FB. Sugar-sweetened beverages, obesity, type 2 diabetes mellitus, and cardiovascular disease risk. *Circulation*. 2010; 121: 1356–1364. <https://doi.org/10.1161/CIRCULATIONAHA.109.876185> PMID: 20308626
5. Weeratunga P, Jayasinghe S, Perera Y, Jayasena G, Jayasinghe S. Per capita sugar consumption and prevalence of diabetes mellitus—global and regional associations. *BMC Public Health*. 2014; 14: 186. <https://doi.org/10.1186/1471-2458-14-186> PMID: 24555673
6. Tedstone A, Anderson S, Allen R, Targett V, Allen R. Sugar Reduction: The evidence for action [Internet]. Public Health England. London; 2015. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/470179/Sugar_reduction_The_evidence_for_action.pdf
7. World Health Organization. WHO Guideline: Sugars intake for adults and children. Geneva; 2015.
8. Johnson RK, Appel LJ, Brands M, Howard B V, Lefevre M, Lustig RH, et al. Dietary sugars intake and cardiovascular health a scientific statement from the american heart association. *Circulation*. 2009. pp. 1011–1020. <https://doi.org/10.1161/CIRCULATIONAHA.109.192627> PMID: 19704096
9. Euromonitor. Euromonitor International's Passport Global Market. [Internet]. 2011. <http://www.euromonitor.com/soft-drinks-in-mexico/report>. [cited 2018 September 18].
10. Sánchez-Pimienta TG, Batis C, Lutter CK, Rivera JA. Sugar-Sweetened Beverages Are the Main Sources of Added Sugar Intake in the Mexican Population. *J Nutr*. 2016; 146: 1888S–1896S. <https://doi.org/10.3945/jn.115.220301> PMID: 27511931
11. U.S. Department of Health and Human Services, U.S. Department of Agriculture. 2015–2020 Dietary Guidelines for Americans. 8th edition [Internet]. 2015.
12. Arantxa Colchero M, Rivera-Dommarco J, Popkin BM, Ng SW, Arantxa Cochero M, Rivera-Dommarco J, et al. In Mexico, evidence of sustained consumer response two years after implementing a sugar-

- sweetened beverage tax. *Health Aff.* 2017; 36: 564–571. <https://doi.org/10.1377/hlthaff.2016.1231> PMID: 28228484
13. Popkin BM, Hawkes C. Sweetening of the global diet, particularly beverages: patterns, trends, and policy responses. *Lancet Diabetes Endocrinol.* 2015; 4: 174–186. [https://doi.org/10.1016/S2213-8587\(15\)00419-2](https://doi.org/10.1016/S2213-8587(15)00419-2) PMID: 26654575
 14. Trieu K, Neal B, Hawkes C, Dunford E, Campbell N, Rodriguez-Fernandez R, et al. Salt Reduction Initiatives around the World—A Systematic Review of Progress towards the Global Target. DeAngelis MM, editor. *PLoS ONE.* San Francisco, CA USA: Public Library of Science; 2015; 10: e0130247. <https://doi.org/10.1371/journal.pone.0130247> PMID: 26201031
 15. Tedstone A, Targett V, Owtram G, Pyne V, Allen R, Bathrellou K, et al. Sugar Reduction: Achieving the 20%. A technical report outlining progress to date, guidelines for industry, 2015 baseline levels in key foods and next steps [Internet]. London; 2017. <https://www.gov.uk/government/publications/sugar-reduction-achieving-the-20>. [cited 2018 September 18].
 16. World Health Organization. Eliminating trans fats in europe. A policy brief [Internet]. Copenhagen; 2015. http://www.euro.who.int/_data/assets/pdf_file/0010/288442/Eliminating-trans-fats-in-Europe-A-policy-brief.pdf. [cited 2018 September 18].
 17. Briggs ADM, Mytton OT, Kehlbacher A, Tiffin R, Elhoussein A, Rayner M, et al. Health impact assessment of the UK soft drinks industry levy: a comparative risk assessment modelling study. *Lancet Public Heal.* 2017; 2: e15–e22. [https://doi.org/10.1016/S2468-2667\(16\)30037-8](https://doi.org/10.1016/S2468-2667(16)30037-8) PMID: 28804786
 18. Public Health England. Sugar reduction and wider reformulation programme: Report on progress towards the first 5% reduction and next steps [Internet]. London; 2018. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/709008/Sugar_reduction_progress_report.pdf
 19. Secretaría de la Salud. Emite la Secretaría de Salud emergencia epidemiológica por diabetes mellitus y obesidad [Internet]. Mexico City; 2016. <https://www.gob.mx/salud/prensa/emite-la-secretaria-de-salud-emergencia-epidemiologica-por-diabetes-mellitus-y-obesidad>. [cited 2018 September 18].
 20. D.R. Secretaría de Salud. Estrategia nacional para la prevención y el control del sobrepeso, la obesidad y la diabetes. [Internet]. IEPSA, Entidad paraestatal del Gobierno Federal. Ciudad de México; 2013. https://www.gob.mx/cms/uploads/attachment/file/276108/estrategia_sobrepeso_diabetes_obesidad.pdf
 21. Romero-Martínez M, Shamah-Levy T, Franco-Núñez A, Villalpando S, Cuevas-Nasu L, Gutiérrez JP, et al. Encuesta Nacional de Salud y Nutrición 2012: diseño y cobertura. *Salud Publica Mex.* 2013; 55: S332–S340.
 22. Goldberg GR, Black AE, Jebb SA, Colte T, Murgatroyd PR, Coward WA, et al. Critical evaluation of energy intake data using fundamental principles of energy physiology: 1. Derivation of cut-off limits to identify under-recording. *Eur J Clin Nutr.* 1991; 45: 569–81. PMID: 1810719
 23. Han E, Powell LM. Consumption Patterns of Sugar-Sweetened Beverages in the United States. *J Acad Nutr Diet.* 2013; 113: 43–53. <https://doi.org/10.1016/j.jand.2012.09.016> PMID: 23260723
 24. Gutiérrez JP. [Household socioeconomic classification in the National Health and Nutrition Survey 2012]. *Salud Pública de México.* 2013. pp. S341–S346.
 25. Rogers PJ, Hogenkamp PS, De Graaf C, Higgs S, Lluch A, Ness AR, et al. Does low-energy sweetener consumption affect energy intake and body weight? A systematic review, including meta-analyses, of the evidence from human and animal studies. *Int J Obes.* 2016; 40: 381–394. <https://doi.org/10.1038/ijo.2015.177> PMID: 26365102
 26. StataCorp. Stata Statistical Software: Release 14. 2015. 2015.
 27. Chow CC, Hall KD. The dynamics of human body weight change. *PLoS Comput Biol.* 2008; 4: e1000045. <https://doi.org/10.1371/journal.pcbi.1000045> PMID: 18369435
 28. Hall KD. Predicting metabolic adaptation, body weight change, and energy intake in humans. *Am J Physiol Metab. Am Physiological Soc;* 2010; 298: E449–E466. <https://doi.org/10.1152/ajpendo.00559.2009> PMID: 19934407
 29. Hall KD, Jordan PN. Modeling weight-loss maintenance to help prevent body weight regain. *Am J Clin Nutr. Am Soc Nutrition;* 2008; 88: 1495–1503. <https://doi.org/10.3945/ajcn.2008.26333> PMID: 19064508
 30. CONAPO. Proyecciones de los hogares en México y las Entidades Federativas, 2010–2030. In: México en cifras [Internet]. 2014 p. 1. <http://www.snieg.mx/cni/sreferencia.aspx>. [cited 2018 September 18].
 31. Camacho-García-Formentí D, Zepeda-Tello R. bw: Dynamic Body Weight Models for Children and Adults. [Internet]. Mexico City: R package version 1.0.0; 2018. <https://cran.r-project.org/web/packages/bw/index.html>

32. Lumley T. Analysis of complex survey samples. *J Stat Softw.* 2004; 9: 1–19. <https://doi.org/10.18637/jss.v009.i08>
33. Lumley T. survey: analysis of complex survey samples [Internet]. R package version 3.32; 2017. <https://cran.r-project.org/web/packages/survey/index.html>
34. Eddelbuettel D. Seamless R and C++ integration with Rcpp. *Seamless R and C++ Integration with Rcpp.* 2013. <https://doi.org/10.1007/978-1-4614-6868-4>
35. Eddelbuettel D, Francois R. Rcpp: Seamless R and C++ Integration. *J Stat Software*; Vol 1, Issue 8. 2011; Available from: <https://www.jstatsoft.org/v040/i08>.
36. Mattes RD. Dietary compensation by humans for supplemental energy provided as ethanol or carbohydrate in fluids. *Physiol Behav.* 1996; 59: 179–187. [https://doi.org/10.1016/0031-9384\(95\)02007-1](https://doi.org/10.1016/0031-9384(95)02007-1) PMID: 8848479
37. Mourao DM, Bressan J, Campbell WW, Mattes RD. Effects of food form on appetite and energy intake in lean and obese young adults. *Int J Obes.* 2007; 31: 1688–1695. <https://doi.org/10.1038/sj.ijo.0803667> PMID: 17579632
38. DiMeglio DP, Mattes RD. Liquid versus solid carbohydrate: effects on food intake and body weight. *Int J Obes Relat Metab Disord.* 2000; 24: 794–800. <https://doi.org/10.1038/sj.ijo.0801229> PMID: 10878689
39. Crino M, Herrera AMM, Ananthapavan J, Wu JHY, Neal B, Lee YY, et al. Modelled cost-effectiveness of a package size cap and a kilojoule reduction intervention to reduce energy intake from sugar-sweetened beverages in Australia. *Nutrients.* 2017; 9: 1–17. <https://doi.org/10.3390/nu9090983> PMID: 28878175
40. Dobbs R, Sawers C, Thompson F, Manyika J, Woetzel J, Child P, et al. Overcoming obesity: An initial economic analysis [Internet]. McKinsey Global Institute. 2014. https://www.mckinsey.com/~/media/McKinsey/Business%20Functions/Economic%20Studies%20TEMP/Our%20Insights/How%20the%20world%20could%20better%20fight%20obesity/MGI_Overcoming_obesity_Full_report.ashx. [cited 2018 September 18].
41. Ma Y, He FJ, Yin Y, Hashem KM, MacGregor GA. Gradual reduction of sugar in soft drinks without substitution as a strategy to reduce overweight, obesity, and type 2 diabetes: a modelling study. *Lancet Diabetes Endocrinol.* 2016; 4: 105–114. [https://doi.org/10.1016/S2213-8587\(15\)00477-5](https://doi.org/10.1016/S2213-8587(15)00477-5) PMID: 26777597
42. Christiansen E, Garby L. Prediction of body weight changes caused by changes in energy balance. *Eur J Clin Invest.* 2002; 32: 826–830. <https://doi.org/10.1046/j.1365-2362.2002.01036.x> PMID: 12423323
43. Stern D, Middaugh N, Rice MS, Laden F, López-Ridaura R, Rosner B, et al. Changes in sugar-sweetened soda consumption, weight, and waist circumference: 2-year cohort of Mexican women. *Am J Public Health.* 2017; <https://doi.org/10.2105/AJPH.2017.304008> PMID: 28933937
44. Sharma LL, Teret SP, Brownell KD. The food industry and self-regulation: Standards to promote success and to avoid public health failures. *Am J Public Health.* 2010; 100: 240–246. <https://doi.org/10.2105/AJPH.2009.160960> PMID: 20019306
45. Restrepo BJ, Rieger M. Trans fat and cardiovascular disease mortality: Evidence from bans in restaurants in New York. *J Health Econ.* 2016; 45: 176–196. <https://doi.org/10.1016/j.jhealeco.2015.09.005> PMID: 26620830
46. Barberio AM, Sumar N, Trieu K, Lorenzetti DL, Tarasuk V, Webster J, et al. Population-level interventions in government jurisdictions for dietary sodium reduction: a Cochrane Review. *Int J Epidemiol.* 2017; 46: 1405–1551. Available: <http://dx.doi.org/10.1093/ije/dyw361>
47. Charlton K, Webster J, Kowal P. To legislate or not to legislate? A comparison of the UK and South African approaches to the development and implementation of salt reduction programs. *Nutrients.* 2014; 6: 3672–3695. <https://doi.org/10.3390/nu6093672> PMID: 25230210
48. Backholer K, Blake M, Vandevijvere S. Sugar-sweetened beverage taxation: an update on the year that was 2017. *Public Health Nutr.* 2017; 20: 3219–3224. <https://doi.org/10.1017/S1368980017003329> PMID: 29160766
49. Grupo Multidisciplinario sobre obesidad de la Academia Nacional de Medicina. *Obesidad en México: Recomendaciones para una política de Estado.* Rivera Dommarco JA, Hernández Ávila M, Aguilar Salinas, CA, Vadillo Ortega F, Murayama Rendón C, editors. Mexico City: UNAM; 2015.
50. Pineli L L O, Aguiar LA, Fiusa A, Botelho RB A, Zandonadi RP, Melo L. Sensory impact of lowering sugar content in orange nectars to design healthier, low-sugar industrialized beverages. *Appetite.* 2016; 96: 239–244. <https://doi.org/10.1016/j.appet.2015.09.028> PMID: 26428862
51. Wise PM, Nattress L, Flammer LJ, Beauchamp GK. Reduced dietary intake of simple sugars alters perceived sweet taste intensity but not perceived pleasantness. *Am J Clin Nutr.* 2016; 103: 50–60. <https://doi.org/10.3945/ajcn.115.112300> PMID: 26607941
52. Raghubir P, Tyebjee TT, Lin Y-C. The Sense and Nonsense of Consumer Product Testing: How to Identify Whether Consumers Are Blindly Loyal? *Found Trends® Mark.* Now Publishers; 2009; 3: 127–176. <https://doi.org/10.1561/1700000009>

53. Delogu F, Huddas C, Steven K, Hachem S, Lodhia L, Fernandez R, et al. A Dissociation Between Recognition and Hedonic Value in Caloric and Non-caloric Carbonated Soft Drinks. *Front Psychol*. 2016; 7: 36. <https://doi.org/10.3389/fpsyg.2016.00036> PMID: 26858681
54. Imamura F, O'Connor L, Ye Z, Mursu J, Hayashino Y, Bhupathiraju SN, et al. Consumption of sugar sweetened beverages, artificially sweetened beverages, and fruit juice and incidence of type 2 diabetes: systematic review, meta-analysis, and estimation of population attributable fraction. *BMJ*. 2015; 351: h3576. <https://doi.org/10.1136/bmj.h3576> PMID: 26199070
55. Azad MB, Abou-Setta AM, Chauhan BF, Rabbani R, Lys J, Copstein L, et al. Nonnutritive sweeteners and cardiometabolic health: a systematic review and meta-analysis of randomized controlled trials and prospective cohort studies. *Can Med Assoc J*. 2017; 189: E929–E939. <https://doi.org/10.1503/cmaj.161390> PMID: 28716847
56. Duffey KJ, Steffen LM, Van Horn L, Jacobs Jr, DR, Popkin BM. Dietary patterns matter: diet beverages and cardiometabolic risks in the longitudinal Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Am J Clin Nutr*. 2012; 95: 909–915. <https://doi.org/10.3945/ajcn.111.026682> PMID: 22378729
57. de Koning L, Malik VS, Kellogg MD, Rimm EB, Willett WC, Hu FB. Sweetened Beverage Consumption, Incident Coronary Heart Disease and Biomarkers of Risk in Men. *Circulation*. 2012; 1735–1741. <https://doi.org/10.1161/CIRCULATIONAHA.111.067017> PMID: 22412070
58. Raben A, Vasilaras TH, Moller AC, Astrup A. Sucrose compared with artificial sweeteners: different effects on ad libitum food intake and body weight after 10 wk of supplementation in overweight subjects. *Am J Clin Nutr*. 2002; 76: 721–729. <https://doi.org/10.1093/ajcn/76.4.721> PMID: 12324283
59. Lenoir M, Serre F, Cantin L, Ahmed SH. Intense Sweetness Surpasses Cocaine Reward. *PLoS ONE*. Public Library of Science; 2007; 2: e698. Available: <https://doi.org/10.1371/journal.pone.0000698> PMID: 17668074
60. Bloomgarden ZT. Nonnutritive Sweeteners, Fructose, and Other Aspects of Diet. *Diabetes Care*. American Diabetes Association; 2011; 34: e46–e51. <https://doi.org/10.2337/dc11-0448> PMID: 21525491
61. Piernas C, Tate DF, Wang X, Popkin BM. Does diet-beverage intake affect dietary consumption patterns? Results from the Choose Healthy Options Consciously Everyday (CHOICE) randomized clinical trial. *Am J Clin Nutr*. 2013; 97: 604–611. <https://doi.org/10.3945/ajcn.112.048405> PMID: 23364015
62. Tate DF, Turner-McGrievy G, Lyons E, Stevens J, Erickson K, Polzien K, et al. Replacing caloric beverages with water or diet beverages for weight loss in adults: main results of the Choose Healthy Options Consciously Everyday (CHOICE) randomized clinical trial. *Am J Clin Nutr*. 2012; 95: 555–63. <https://doi.org/10.3945/ajcn.111.026278> PMID: 22301929
63. Baranowski T. 24-Hour Recall and Diet Record Methods. *Nutritional Epidemiology*. Third Edit. Oxford University Press; 2013. pp. 49–69. <https://doi.org/10.1093/acprof:oso/9780199754038.003.0004>
64. Macdiarmid J, Blundell J. Assessing dietary intake: Who, what and why of under-reporting. *Nutr Res Rev*. 1998; 11: 231. <https://doi.org/10.1079/NRR19980017> PMID: 19094249
65. Malisova O, Bountziouka V, Zampelas A, Kapsokefalou M. Evaluation of Drinks Contribution to Energy Intake in Summer and Winter. *Nutrients*. MDPI; 2015; 7: 3724–3738. <https://doi.org/10.3390/nu7053724> PMID: 25988765
66. Hall KD, Butte NF, Swinburn BA, Chow CC. Dynamics of childhood growth and obesity: development and validation of a quantitative mathematical model. *Lancet Diabetes Endocrinol*. 2013; 1: 97–105. PMID: 24349967
67. Barrientos-Gutierrez T, Zepeda-Tello R, Rodrigues ER, Colchero-Aragónés A, Rojas-Martínez R, Lazcano-Ponce E, et al. Expected population weight and diabetes impact of the 1-peso-per-litre tax to sugar sweetened beverages in Mexico. *PLoS ONE*. 2017; 12: e0176336. <https://doi.org/10.1371/journal.pone.0176336> PMID: 28520716