

VOLUME 290 (2015) PAGES 27090–27100

DOI 10.1074/jbc.W118.005758

Withdrawal: Antioncogenic and oncogenic properties of Nrf2 in arsenic-induced carcinogenesis.

Young-Ok Son, Poyil Pratheeshkumar, Ram Vinod Roy, John Andrew Hitron, Lei Wang, Sasidharan Padmaja Divya, Mei Xu, Jia Luo, Gang Chen, Zhuo Zhang, and Xianglin Shi

This article has been withdrawn by the authors. The corresponding author identified some issues and brought them to the attention of the Journal. The following issues were discovered in the article. The GAPDH immunoblot in Fig. 5A was reused in Figs. 1H, 2F, 2I, 3B, 3G, and 5B. The ESR spectrum from BEAS-2B cells in Fig. 2A was previously published in Son *et al.* ((2014) *J. Biol. Chem.* **289**, 28660–28675) without attribution. The BEAS-2B image in Fig. 2C was reused in Fig. 6A as control siRNA. The GAPDH immunoblot from Fig. 3A was previously published in Fig. 4A of Son *et al.* ((2014) *J. Biol. Chem.* **289**, 28660–28675), representing different experimental conditions. The GAPDH immunoblot from Fig. 3D was reused in Fig. 3I. A portion of the same immunoblot was reused in Fig. 4A as actin. Lanes 9–12 of the β -actin gel in Fig. 4B was reused in lanes 9–10 of the Bcl-xL ARE F1 gel in Fig. 4E. Fig. 5B contained several undeclared splices. In Fig. 5C, the control images for BEAS-2B and AsT cells were reused.