

Readmissions in adult patients following hospitalization for influenza: a nationwide cohort study

Srikanth Yandrapalli, Wilbert S. Aronow, William H. Frishman

Department of Medicine, New York Medical College at Westchester Medical Center, Valhalla, NY, USA

Contributions: (I) Conception and design: All authors; (II) Administrative support: None; (III) Provision of study materials or patients: None; (IV) Collection and assembly of data: None; (V) Data analysis and interpretation: All authors; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Srikanth Yandrapalli, MD. Division of Internal Medicine, Westchester Medical Center, 100 Woods Road, Taylor Pavilion, Valhalla, NY 10595, USA. Email: srikanth.yn@gmail.com.

Background: Influenza epidemics are a major health care concern in the US. Influenza related complications can increase in-hospital complications, and readmissions following a hospitalization for influenza. We sought to determine the 30-day readmission rate, etiologies, outcomes, and healthcare burden of 30-day readmissions in adults hospitalized for influenza.

Methods: The 2014 US National Readmissions Database (NRD) was retrospectively analyzed to identify patients ≥ 18 years of age hospitalized for influenza and discharged between January and November 2014. We used this time frame as this was the most recent data available for analysis and included patients who had 30-day follow-up. Survey design based multivariable logistic regression models were used to identify factors associated with a 30-day readmission.

Results: Of the 46,117 patients who were hospitalized for influenza and survived to discharge, 4,721 (10.2%) patients had 5,275 30-day readmissions, estimated to 11.4 readmissions per 100 patients. Non-influenza pneumonia was the most common etiology of 30-day readmissions (10.4%) followed by sepsis (9.8%). The median costs of readmissions were \$8,538 (IQR, \$5,053–15,262), which were significantly higher than the median costs of their index hospitalizations [\$7,863 (IQR, \$4,875–13,212); $P < 0.001$]. Around 6.5% of the patients died during a readmission.

Conclusions: Adult patients hospitalized for influenza had 11.4 30-day readmissions per 100 patients, most commonly for non-influenza pneumonia. Thirty-day readmissions were associated with higher costs of care and considerable mortality.

Keywords: Influenza; acute myocardial infarction (AMI); stroke; readmissions; pneumonia

Submitted Apr 20, 2018. Accepted for publication Jul 19, 2018.

doi: 10.21037/atm.2018.07.18

View this article at: <http://dx.doi.org/10.21037/atm.2018.07.18>

Introduction

Influenza is a major public health problem in the US with significant variations in its annual incidence and public health burden (1). While the majority of influenza patients are managed in the ambulatory setting, influenza and related complications can result in hospitalizations (1,2). The Center for Disease Control (CDC) reported an estimated minimum 140,000 and maximum 710,000

hospitalizations for influenza and related complications between years 2010 and 2016 in the US (1,2). In the US in year 2003, influenza-related hospitalizations amounted to \$6.0 billion in costs, which were only 7% of the total economic burden of influenza during that year (3). Among adult patients hospitalized for influenza, more than two-thirds have underlying medical conditions, most commonly cardiovascular (CV) disease, metabolic disorder, obesity, and chronic lung disease (4). Further, available data

suggest an increased risk of adverse CV events like acute myocardial infarction (AMI) and ischemic cerebrovascular events (CVEs) in the acute phase following an influenza infection (5-14).

While such risk of increased adverse events can increase in-hospital complications, they can also cause readmissions in these patients in the acute phase following influenza infection. Readmissions are usually measured within 30 days because readmissions and deaths after a longer time period may be less influenced by the admitting condition and care gotten in the hospital and more to do with other complicating illnesses, patients' behavior, or care provided to patients after hospital discharge (15). Data are scarce regarding the etiology and burden of 30-day readmissions following a hospitalization for influenza. We sought to determine the 30-day readmission rates and associated factors, reasons for readmission and their healthcare burden, after hospitalization for either influenza-pneumonia or influenza-related organ system manifestations in adult patients.

Methods

Data source

In this retrospective observational study, data were collected from the publicly available 2014 US National Readmissions Database (NRD). The NRD is part of the Healthcare Cost and Utilization Project (HCUP) that is sponsored by the Agency for Healthcare Research and Quality (AHRQ) and is a database of all-payer hospital inpatient stays that can be used to generate national estimates of readmissions (16). The NRD is drawn from HCUP State Inpatient Databases which contain verified patient linkage numbers that can be used to track a person across readmissions while maintaining privacy. The NRD includes community hospitals and excludes rehabilitation or long-term acute care hospitals. Unweighted, the 2014 NRD contains data from approximately 15 million discharges from 22 partner states: Arkansas, California, Florida, Georgia, Hawaii, Iowa, Louisiana, Maryland, Massachusetts, Missouri, Nebraska, New Mexico, Nevada, New York, South Carolina, South Dakota, Tennessee, Utah, Virginia, Vermont, Washington, and Wisconsin. These states are geographically dispersed and account for 51.2% of the total US resident population and 49.3% of all hospitalizations. Weighted, it estimates roughly 35 million discharges (16).

The NRD was post-stratified by hospital characteristics

(census region, urban/rural location, hospital teaching status, size of the hospital defined by the number of beds, and hospital control) and patient characteristics [sex and five age groups (0, 1–17, 18–44, 45–64, and 65 and older)] (16). Discharge weights (variable DISCWT) were provided to obtain national estimates.

Patient population and baseline characteristics

The study sample included all adult patients (≥ 18 years) who were discharged alive between January and November 2014 with a primary discharge diagnosis of influenza pneumonia or other influenza-related manifestations. These cases formed our index cohort. Patients discharged in December were excluded because the NRD design will not allow for 30-day follow-up of patients discharged in this month. International Classification of Diseases, Ninth Edition, Clinical Modification (ICD-9-CM) diagnosis codes 487 and 488 were used to identify influenza diagnoses. We chose data from 2014 for our study as these were the most recently available data which used ICD-9 codes throughout for coding diagnosis. If a patient had multiple hospitalizations for influenza during the study period, only the first hospitalization for influenza was included in the index cohort. We did not include hospitalizations with a secondary discharge diagnosis of influenza because such hospitalizations could include patients with previously diagnosed influenza, co-incident influenza diagnosis, and also hospital-acquired influenza when admitted for other reasons.

Baseline patient characteristics identified in the index cohort were age, sex, primary expected payer, median household income, comorbidities (anemia, atrial fibrillation, cancer, chronic lung disease, coagulopathy, depression, drug abuse, dementia, diabetes mellitus, hypertension, hypothyroidism, fluid-electrolyte imbalances, liver disease, obesity, other neurological disorders, paralysis, psychoses, peripheral vascular disease, pulmonary circulation disorders, chronic renal failure, and heart failure), in-hospital complications [septicemia and shock, AMI, acute kidney injury (AKI), ischemic stroke and transient cerebral ischemic events, gastrointestinal hemorrhage, and respiratory failure), type and day of admission, and discharge disposition. Hospital characters (location, bed-size, and teaching status) were also identified. Elixhauser comorbidity data were extracted from the NRD. ICD-9-CM and the HCUP Clinical Classification Software

(CCS) codes used to identify comorbidities and in-hospital complications are available in *Table S1*.

Study outcomes

The primary outcomes of this study were the 30-day readmission rate from the day of discharge, etiology for readmission, and the factors associated with 30-day readmissions. Secondary outcomes were comparison of length of stay (LOS) and costs of care between index hospitalization and readmission, and all-cause in-hospital mortality during 30-day readmissions only because all patients in our index cohort survived to discharge. LOS and costs of hospitalization were identified in both index cases and readmission, whereas, all-cause inpatient mortality was identified in 30-day readmissions. Methodology described by the HCUP to identify readmissions was used to identify 30-day readmissions (16). Total number of 30-day readmissions and the number of index-patients readmitted within 30-day were identified and expressed as proportion of the index hospitalizations. Time to readmission was calculated as the duration between index-hospital discharge and the first day of hospital readmission.

Etiologies of readmissions were recognized by identifying the ICD-9-CM diagnosis codes and the HCUP CCS codes of the primary discharge diagnosis of each individual readmission (available in *Table S2*). Causes of readmissions were classified on the basis of organ system involvement into respiratory, CV, infection-related, neurovascular, peripheral vascular, hematologic and cancer related, gastrointestinal, renal and fluid-electrolyte disorders, endocrine and metabolic, neuropsychiatric and substance abuse-related, musculoskeletal and injury related, symptom related (fever of unknown origin, malaise, abdominal pain, nausea/vomiting), complication of device-graft-implant, or others. Respiratory causes included non-influenza pneumonia and empyema, influenza, obstructive lung disease, pleural disease, respiratory failure, aspiration pneumonitis, and others. CV causes included heart failure, arrhythmias and conduction disorders, AMI, hypertension, coronary artery disease, pulmonary embolism, syncope, and chest pain. Infection-related etiologies included sepsis, intestinal infections, skin infections, urinary tract infections (UTIs), and others.

To identify costs of hospitalization, hospital reported charges (variable TOTCHG) were multiplied with the cost-to-charge ratios developed and reported by the AHRQ.

Statistical analysis

All statistical analyses were performed using IBM SPSS Statistics 25.0 (IBM Corp., Armonk, NY, USA). Weighted data were used for all statistical analyses. Categorical variables are expressed as percentages, and continuous variables as mean \pm standard deviation (SD). Because the distributions of LOS and total hospital costs were skewed to the right, median [interquartile range (IQR)] were also reported. Baseline patient characters, comorbidities and complications, and hospital characteristics were compared between patients with and without 30-day readmission using Pearson χ^2 test for categorical variables and student *t*-test for continuous variables. Nonparametric Mann-Whitney U test was used to compare medians between patient groups. Since the NRD has a complex semi-random multistage sampling survey design, survey design based multivariable logistic regression model [designed to test for 95% confidence intervals (CIs) and a two-sided P value threshold of <0.05] was designed to identify factors associated with 30-day readmissions. The regression model included all the variables which achieved a P value <0.2 in unadjusted analysis. In the NRD, patients' aged ≥ 90 years were reported as 90-year-old. This can affect the results when age (in years) is used as a covariate in statistical analysis. Hence, patients were sub grouped into three age categories (18–44, 45–64, and ≥ 65 years). LOS of index hospitalizations was included in the regression model as a categorical variable divided into 2 groups based on median LOS cut-off. Results of the logistic regression analyses were reported as odds ratios (ORs) and 95% CIs.

Around $\leq 1\%$ data were missing for the variables primary payer status, median household income, discharge disposition, and type of admission (elective versus non-elective). Missing values were replaced with the dominant categories. Total hospital charge data were missing for around 1% index-hospitalizations and 0.9% of the 30-day readmission. Reported cost data were for the remainder of the cases.

Results

Baseline characteristics of the study cohort and in-hospital complications

The index-study cohort included 46,117 adult patients who had influenza hospitalization meeting our index-case criteria and survived to discharge. These patients had a

mean age \pm SD of 61.1 \pm 17.9 years and 53.6% were female. Of these, 22,457 (48.7%) were primarily hospitalized for influenza-pneumonia, 22,375 (48.5%) for influenza-related respiratory symptoms, and 1,285 (2.8%) for other organ-system manifestations of influenza. Overall, around 44.5% patients were elderly with age \geq 65 years. Baseline patient demographics, comorbidities, in-hospital complications, and hospital characteristics for the overall index-study cohort are presented in *Table 1*. More patients (31.1%) were in the lowest median income quartile (\$1–39,999) group and around two-thirds had either Medicare or Medicaid health insurance. Hypertension, chronic pulmonary disease, diabetes mellitus, fluid-electrolyte disturbances, and anemia were the most prevalent comorbidities. While respiratory failure occurred in 22.1% of index hospitalizations, AKI complicated 12.8%, an AMI in 1.2%, and ischemic CVE in 0.3%. More patients in the index-cohort were hospitalized in large-sized urban-based teaching hospitals. Median LOS was relatively short at 3 days [IQR, 2–6 days] and mean LOS \pm SD was 5.1 \pm 6.9 days. The total costs of index hospitalizations were \$489.9 million, mean \$10,729 \pm 22,255 and median \$6,082 (IQR, \$3,921–10,220).

Thirty-day readmissions and characteristics of index hospitalizations by readmission status

Of the 46,117 index-group patients, 4,721 (10.2%) patients had 5,275 30-day readmissions. This estimates to 11.4 readmissions per 100 patients within a 30-day period from discharge. Baseline patient demographics, comorbidities, in-hospital complications, and hospital characters for the index cases with and without a readmission are presented in *Table 1*.

Patients who had a 30-day readmission were generally older (63.9 \pm 17.5 years) compared to those without a 30-day readmission (60.7 \pm 17.9 years; $P=0.02$), more likely to be insured by Medicare/Medicaid (74.9% *vs.* 64.6%; $P<0.001$), and more likely to have a higher burden of comorbidities and in-hospital complications. Ischemic CVEs (0.6% *vs.* 0.3%; $P=0.005$), AMI (2.1% *vs.* 1.2%; $P<0.001$), and gastrointestinal bleeding (1.4% *vs.* 0.7%; $P<0.001$) occurred almost two-times the rate in patients who had a 30-day readmission compared to those who did not. Patients who had a 30-day readmission had a longer median initial LOS [4 days (IQR, 3–7 days) *vs.* 3 days (IQR, 2–5 days); $P<0.001$], more likely to be discharged to a nursing home or other medical facility (20.4% *vs.* 12.3%; $P<0.001$) and had higher median index hospitalization costs

[\$7,863 (IQR \$4,875–13,212) *vs.* \$5,946 (\$3,830–9,804); $P<0.001$].

Factors associated with a 30-day readmission

In multivariable logistic regression analysis results presented in *Table 2*, presence of chronic pulmonary disease (OR 1.15; 95% CI, 1.03–1.27), drug abuse (OR 1.53; 95% CI, 1.18–1.99), liver disease (OR 1.47; 95% CI, 1.12–1.92), chronic renal failure (OR 1.42; 95% CI, 1.23–1.64), heart failure (OR 1.45; 95% CI, 1.25–1.69), AKI (OR 1.24; 95% CI, 1.07–1.44), anemia (OR 1.32; 95% CI, 1.17–1.50), cancer (OR 1.93; 95% CI, 1.56–2.39), and sepsis (OR 1.31; 95% CI, 1.01–1.68) were associated with a 30-day readmission. A longer LOS >3 days (OR 1.48; 95% CI, 1.31–1.67), discharge to nursing home/other medical facility (OR 1.23; 95% CI, 1.07–1.44), and admission to a teaching hospital (OR 1.15; 95% CI, 1.03–1.29) also predicted a 30-day readmission.

Timing and causes of 30-day readmissions

Of the 5,275 readmissions, 45.8% were within 10 days from discharge, 29.3% between days 11–20, and 24.8% between days 21–30. Median time to readmission was 12 days (IQR, 5–20 days). Etiologies of readmissions were presented in *Table 3*. When stratified according to organ-system involvement, most common causes of 30-day readmission were respiratory (31.2%), infectious (15.8%), CV (14.9%), gastrointestinal (7.6%), and hematologic-cancer related (4.9%). Diagnosis wise, non-influenza pneumonia was the most common etiology of 30-day readmissions (10.4%), followed by sepsis (9.8%), obstructive lung disease (8.0%), heart failure (6.2%), AKI and UTIs at 2.3% each. While a microbiological diagnosis was not made in most non-influenza pneumonias (78.4%), methicillin-resistant *Staphylococcus aureus* (MRSA) was the most frequently identified organism (4.8%) in the rest. Similarly, around 81% of the sepsis-related admissions did not have a microbiologic diagnosis, but when an organism was identified, it was more often gram-positive (12%; 5% streptococcal and 4% MRSA).

LOS, costs, and mortality related to 30-day readmissions

Overall, 6.5% of the patients readmitted within 30-days from discharge died during a readmission. Mortality was highest when the readmission etiology was infectious,

Table 1 Baseline characteristics, comorbidities, in-hospital complication, and outcomes of patients in the index-cohort with and without a 30-day readmission

Baseline characteristics	Total, % (N=46,117)	30-day readmission		P value
		No, % (N=41,396)	Yes, % (N=4,721)	
Patient variables				
Age (years)				<0.001
18–44	18.7	19.2	14.2	
45–64	36.8	37.1	34.1	
65 and above	44.5	43.7	51.7	
Sex				0.081
Female	53.6	53.8	52.4	
Male	46.4	46.2	47.6	
Median household income				0.394
\$1–39,999	31.1	31.2	30.2	
\$40,000–50,999	26.4	26.2	27.6	
\$51,000–65,999	21.9	22.0	20.7	
\$66,000 or more	20.7	20.6	21.5	
Primary payer				<0.001
Medicare/Medicaid	65.7	64.6	74.9	
Private	25.4	26.1	19.6	
Uninsured	6.6	6.9	3.8	
Others	2.3	2.4	1.8	
Comorbidities				
Anemia	17.5	16.4	26.4	<0.001
Atrial fibrillation	13.1	12.5	18.5	<0.001
Cancer	5.1	4.7	8.9	<0.001
Chronic pulmonary disease	40.3	39.9	43.6	<0.001
Chronic renal failure	15.6	14.4	25.8	<0.001
Coagulopathy	8.0	7.7	9.9	<0.001
Heart failure	15.1	13.9	25.5	<0.001
Dementia	7.8	7.5	10.3	<0.001
Depression	13.1	12.8	15.2	<0.001
Diabetes mellitus	30.0	29.4	35.5	<0.001
Drug abuse	3.3	3.2	4.6	<0.001
Fluid and electrolyte disorders	41.7	41.5	43.3	0.021
Hypertension	58.9	58.2	64.8	<0.001
Hypothyroidism	13.0	12.8	14.3	0.004
Liver disease	2.8	2.7	4.4	<0.001
Obesity	16.6	16.8	15.4	0.019
Other neurological disorders	9.7	9.4	11.6	<0.001

Table 1 (continued)

Table 1 (continued)

Baseline characteristics	Total, % (N=46,117)	30-day readmission		P value
		No, % (N=41,396)	Yes, % (N=4,721)	
Paralysis	2.3	2.2	3.2	<0.001
Peripheral vascular disease	4.4	4.1	6.4	<0.001
Psychoses	5.2	5.1	6.0	0.006
Pulmonary circulation disorders	4.2	3.9	6.8	<0.001
In-hospital complications				
AMI	1.2	1.1	2.1	<0.001
AKI	12.8	12.1	19.7	<0.001
Sepsis and shock	2.8	2.6	4.4	<0.001
Respiratory failure	22.1	21.4	28.2	<0.001
Gastrointestinal hemorrhage	0.8	0.7	1.4	<0.001
Ischemic CVE	0.3	0.3	0.6	0.005
Median LOS	3 days	3 days	4 days	<0.001
≤3 days	51.9	53.5	37.4	
>3 days	48.1	46.5	62.6	
Admission day				0.026
Weekday	74.2	74.0	75.5	
Weekend	25.8	26.0	24.5	
Admission type				0.110
Elective	4.6	4.6	4.1	
Disposition				<0.001
Home	86.9	87.7	79.6	
Short term hospital, nursing home, or other facility	13.1	12.3	20.4	
Hospital characteristics				
Location				0.001
Rural	4.6	4.7	3.6	
Urban	95.4	95.3	96.4	
Teaching status				<0.001
Non-teaching	40.6	41.1	35.9	
Teaching	59.4	58.9	64.1	
Bed size				<0.001
Small	18.7	19.0	16.4	
Medium	27.8	27.8	27.6	
Large	53.5	53.3	56.0	
Median cost of care (US \$)	6,082	5,946	7,863	<0.001

AMI, acute myocardial infarction; AKI, acute kidney injury; LOS, length of stay; CVE, cerebrovascular event.

Table 2 Results of the multivariable regression analysis to identify factors associated with increased 30-day readmissions

Factors	Reference category	Multivariable analysis*, OR (95% CI)
LOS >3 days	LOS ≤3 days	1.48 (1.31–1.67)
Sepsis and septic shock	Absent	1.31 (1.01–1.68)
Cancer	Absent	1.93 (1.56–2.39)
Anemia	Absent	1.32 (1.17–1.50)
AKI	Absent	1.24 (1.07–1.44)
Heart failure	Absent	1.45 (1.25–1.69)
Chronic renal failure	Absent	1.42 (1.23–1.64)
Liver disease	Absent	1.47 (1.12–1.92)
Drug abuse	Absent	1.53 (1.18–1.99)
Chronic pulmonary disease	Absent	1.15 (1.03–1.27)
Discharge to nursing home/other facility	Home discharge	1.24 (1.07–1.44)
Teaching hospital	Non-teaching hospital	1.15 (1.03–1.29)
Private insurance	Medicare/Medicaid	0.85 (0.74–0.97)
Uninsured	Medicare/Medicaid	0.67 (0.52–0.85)
Obesity	Non-obese	0.82 (0.69–0.96)
Age (years)		
45–64	18–44	1.02 (0.85–1.23)
65 and above	18–44	1.01 (0.84–1.21)
Male sex	Female sex	1.02 (0.93–1.13)
AMI	Absent	1.08 (0.75–1.56)
Ischemic CVE	Absent	1.05 (0.45–2.47)
Atrial fibrillation	Absent	1.10 (0.95–1.28)
Coagulopathy	Absent	1.04 (0.87–1.24)
Hypertension	Absent	1.01 (0.90–1.13)
Hypothyroidism	Absent	0.98 (0.83–1.15)
Depression	Absent	1.13 (0.97–1.32)
Dementia	Absent	1.06 (0.87–1.29)
Diabetes mellitus	Absent	1.12 (0.99–1.26)
Gastrointestinal bleed	Absent	1.33 (0.87–2.05)
Fluid and electrolyte disorder	Absent	0.93 (0.84–1.04)
Other neurological disorder	Absent	1.09 (0.92–1.30)
Paralysis	Absent	1.28 (0.95–1.74)
Peripheral vascular disease	Absent	1.09 (0.88–1.34)
Psychoses	Absent	1.14 (0.91–1.44)
Pulmonary circulation disorders	Absent	1.22 (0.99–1.49)

Table 2 (continued)

Table 2 (continued)

Factors	Reference category	Multivariable analysis*, OR (95% CI)
Respiratory failure	Absent	1.12 (0.98–1.28)
Elective admission	Non-elective admission	0.95 (0.72–1.26)
Weekend admission	Weekday admission	0.89 (0.79–1.01)
Rural-based hospital	Urban-based hospital	1.02 (0.74–1.39)
Bed size of hospital		
Large	Small	1.15 (0.99–1.33)
Medium	Small	1.16 (0.99–1.37)

*; survey design based multivariable logistic regression model designed to test for 95% confidence intervals and a two-sided P value of <0.05. OR, odds ratio; CI, confidence intervals; LOS, length of stay; AKI, acute kidney injury; AMI, acute myocardial infarction; CVE, cerebrovascular event.

Table 3 System based and disease specific etiologies of 30-day readmission after hospitalization for influenza

Etiology of readmission	30-day readmissions (N=5,275, %)
Respiratory	1,643 (31.1)
Non-influenza pneumonia and empyema	549 (10.4)
Influenza	190 (3.6)
Chronic obstructive lung disease and asthma	422 (8.0)
Aspiration pneumonitis	127 (2.4)
Pleural disease	32 (0.6)
Respiratory failure	223 (4.2)
Other respiratory tract disease	100 (1.9)
Cardiovascular	784 (14.9)
Heart failure	327 (6.2)
Conduction disease and arrhythmias	127 (2.4)
Acute myocardial infarction	63 (1.2)
Coronary artery disease	37 (0.7)
Hypertension	63 (1.2)
Pulmonary embolism	84 (1.6)
Syncope	32 (0.6)
Nonspecific chest pain	42 (0.8)
Infection-related and septicemia	835 (15.8)
Septicemia and shock	516 (9.8)
Intestinal infection	92 (1.7)
Urinary tract infections	119 (2.3)

Table 3 (continued)

Etiology of readmission	30-day readmissions (N=5,275, %)
Skin and soft-tissue infection	60 (1.1)
Others	42 (1.0)
Neurovascular	104 (2.0)
Ischemic cerebrovascular events	59 (1.1)
Peripheral vascular	91 (1.7)
Endocrine and metabolic	119 (2.3)
Gastrointestinal	402 (7.6)
Gastrointestinal hemorrhage	106 (2.0)
Renal diseases and fluid/electrolyte imbalance	243 (4.6)
Acute kidney injury	119 (2.3)
Fluid and electrolyte disorder	96 (1.8)
Neuropsychiatric and substance-related	242 (4.6)
Musculoskeletal and injury related	229 (4.3)
Symptom-related	109 (2.0)
Fever of unknown origin	38 (0.7)
Nausea/vomiting	22 (≤0.5)
Abdominal pain	28 (≤0.5)
Malaise and fatigue	15 (≤0.5)
Device, implant, graft complications	129 (2.4)
Others	90 (1.7)

Table 3 (continued)

reaching a high of 21.5% in patient admitted for septicemia. Median LOS was 4 days (IQR, 3–8 days) with a mean of 6.5 ± 7.6 days. Total costs associated with 30-day readmissions were around \$77.2 million during the study period. Median readmission costs were \$8,538 (IQR, \$5,053–15,262) and mean costs were $\$14,772 \pm 23,348$. The median LOS of the readmissions (4 days) was significantly higher than the median LOS of all index admissions (3 days; $P < 0.001$), but was similar to the median LOS of the index cases which had a readmission (4 vs. 4 days; $P = 0.77$). The median costs of readmissions (\$8,538) were significantly higher compared to median costs of all index cases (\$6,082; $P < 0.001$) and index cases with a 30-day readmission (\$7,863; $P < 0.001$).

Discussion

In this retrospective observational study of a large group of patients who were representative of the US population and were primarily hospitalized for influenza in 2014, a substantial number of patients were readmitted within 30-days from discharge, with an estimated 11.4% readmission rate.

Respiratory system diseases and infections were the most common etiologies for a 30-day readmission. Influenza infection increases the risk of secondary pneumonias' particularly bacterial (17). In our study, non-influenza pneumonia was the most common cause of 30-day readmissions. A microbiological diagnosis was not made in around four-fifths of the cases in our study, but the most commonly identified organism was MRSA in 5%. During influenza epidemics, both *Streptococcus pneumoniae* and *Staphylococcus aureus* (more recently MRSA) were identified as common causes of bacterial pneumonias (17).

Prior studies have reported an increased risk of AMI and stroke in the acute phase following influenza (5-7,9-11,13,14,18,19). In our study, AMI and ischemic CVEs had incident rates of 1.2% and 0.3% respectively and were infrequent causes of 30-day readmissions (between 1–2%). However, it must be noted that we only included patients who survived to discharge during an index hospitalization in our analysis and the hence did not fully capture incidence rate of AMI or ischemic CVEs in all adults hospitalized for influenza. Influenza infection has not been clearly implicated to cause heart failure and obstructive lung disease exacerbations (20–22). In our study, hospitalizations for heart failure, chronic obstructive pulmonary disease, and asthma were important causes 30-day readmissions.

In our study, increasing age categories and differences

in sex did not influence 30-day readmission risk. Anemia, cancer, heart failure, chronic lung disease, liver disease, and chronic renal failure independently predicted a 30-day readmission. These conditions increase the comorbid disease burden in patients and can result in multiple hospitalizations. AMI and ischemic CVEs, which occurred infrequently in our index group patients, were not associated with a 30-day readmission. Interestingly, LOS more than 3 days was independently associated with a readmission. A longer LOS might suggest a sicker patient but a LOS higher than the median predicted a readmission after accounting for common in-hospital complications and comorbidities. Patients had a higher 30-day readmission risk when discharged to a nursing home or other medical facility. Patients discharged to a nursing home should have more supervision of care to help reduce hospitalizations. However, patients typically discharged to a nursing facility are older with a higher comorbidity burden and severity of illness associated with increased readmissions (23). Early and appropriate discharge planning to reduce LOS and to aim for home-discharge with adequate support might help reduce the readmission rate in this patient group.

Readmissions also represented a significant health-care and economic burden. Among patients hospitalized within 30 days from discharge, 6.5% died. More patients died when the cause of readmission was infectious. Appropriate infection control strategies during index hospitalization and early identification and treatment of septicemia should be emphasized to decrease this high mortality rate. The economic burden of readmissions was higher than the index admissions despite having a similar LOS. These findings suggest higher health-care resource utilization during readmissions.

The strengths of our study are utilization of a large nationally representative administrative database which was collected using validated methods to reduce selection bias. Using validated methodology, we were able to determine the 30-day readmission rates, identify the causes and predictors of readmissions, and determine their economic impact in the real-world patients hospitalized for influenza (16).

Our study has important limitations. Firstly, administrative databases lack clinical and laboratory variables which can more accurately determine duration of infection and patient status. It is not known if influenza was diagnosed based on clinical symptoms or laboratory testing in our study. Secondly, using billing codes to identify causes of hospitalizations and in-hospital complications can result in misclassification

bias. Thirdly, data on anti-influenza medication use and vaccination was not available. We were unable to test the association of immunization and anti-viral treatment on readmission risk. Fourthly, our sample included influenza patients across 2 different seasons (January–November) and excluded patients discharged in December. The design of the NRD, which is a yearly database, limited our ability to perform analysis bases on flu-season. We could not adjust for viral-strain and seasonal based differences in readmission rates. Lastly, the NRD does not report out of hospital and emergency room mortality data, which could influence the readmission rates.

Conclusions

In conclusion, adult patients hospitalized for influenza had 11.4 30-day readmissions per 100 patients, most commonly for non-influenza pneumonia and other infections. Thirty-day readmissions were associated with significant healthcare impact with higher costs of care and considerable mortality.

Acknowledgements

None.

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

References

1. Disease Burden of Influenza. Seasonal Influenza (Flu). Available online: <https://www.cdc.gov/flu/about/disease/burden.htm>
2. Estimated Influenza Illnesses, Medical Visits, Hospitalizations, and Deaths Averted by Vaccination in the United States. Seasonal Influenza (Flu). Available online: <https://www.cdc.gov/flu/about/disease/2015-16.htm>
3. Molinari NA, Ortega-Sanchez IR, Messonnier ML, et al. The annual impact of seasonal influenza in the US: measuring disease burden and costs. *Vaccine* 2007;25:5086-96.
4. Weekly U.S. Influenza Surveillance Report. Seasonal Influenza (Flu). Available online: <https://www.cdc.gov/flu/weekly/index.htm>
5. Barnes M, Heywood AE, Mahimbo A, et al. Acute myocardial infarction and influenza: a meta-analysis of case-control studies. *Heart* 2015;101:1738-47.
6. Estabragh ZR, Mamas MA. The cardiovascular manifestations of influenza: a systematic review. *Int J Cardiol* 2013;167:2397-403.
7. Madjid M, Naghavi M, Litovsky S, et al. Influenza and cardiovascular disease: a new opportunity for prevention and the need for further studies. *Circulation* 2003;108:2730-6.
8. Flu and Heart Disease & Stroke. <https://www.cdc.gov/flu/heartdisease/index.htm>
9. Urbanek C, Palm F, Grau AJ. Influenza and stroke risk: a key target not to be missed? *Infect Disord Drug Targets* 2010;10:122-31.
10. Warren-Gash C, Smeeth L, Hayward AC. Influenza as a trigger for acute myocardial infarction or death from cardiovascular disease: a systematic review. *Lancet Infect Dis* 2009;9:601-10.
11. Toschke AM, Heuschmann PU, Wood O, et al. Temporal relationship between influenza infections and subsequent first-ever stroke incidence. *Age Ageing* 2009;38:100-3.
12. Nguyen JL, Yang W, Ito K, et al. Seasonal Influenza Infections and Cardiovascular Disease Mortality. *JAMA Cardiol* 2016;1:274-81.
13. Muhammad S, Haasbach E, Kotchourko M, et al. Influenza virus infection aggravates stroke outcome. *Stroke* 2011;42:783-91.
14. Kwong JC, Schwartz KL, Campitelli MA, et al. Acute Myocardial Infarction after Laboratory-Confirmed Influenza Infection. *N Engl J Med* 2018;378:345-53.
15. The Official U.S. Government Site for Medicare. Available online: <https://www.medicare.gov/hospitalcompare/Data/30-day-measures.html>
16. The Hcup Nationwide Readmissions Database (NRD), 2014. Available online: https://www.hcup-us.ahrq.gov/db/nation/nrd/NRD_Introduction_2014.jsp
17. Metersky ML, Masterton RG, Lode H, et al. Epidemiology, microbiology, and treatment considerations for bacterial pneumonia complicating influenza. *Int J Infect Dis* 2012;16:e321-31.
18. Clar C, Oseni Z, Flowers N, et al. Influenza vaccines for preventing cardiovascular disease. *Cochrane Database Syst Rev* 2015;5:CD005050.
19. Udell JA, Zawi R, Bhatt DL, et al. Association between influenza vaccination and cardiovascular outcomes in high-risk patients: a meta-analysis. *JAMA* 2013;310:1711-20.
20. Kurai D, Saraya T, Ishii H, et al. Virus-induced exacerbations in asthma and COPD. *Front Microbiol* 2013;4. doi:10.3389/fmicb.2013.00293

21. Mallia P, Johnston SL. Influenza infection and COPD. *Int J Chron Obstruct Pulmon Dis* 2007;2:55-64.
22. Sandoval C, Walter SD, Krueger P, et al. Risk of hospitalization during influenza season among a cohort of patients with congestive heart failure. *Epidemiol Infect* 2007;135:574-582.
23. Kripalani S, Theobald CN, Anctil B, et al. Reducing hospital readmission: current strategies and future directions. *Annu Rev Med* 2014;65:471-85.

Cite this article as: Yandrapalli S, Aronow WS, Frishman WH. Readmissions in adult patients following hospitalization for influenza: a nationwide cohort study. *Ann Transl Med* 2018;6(16):318. doi: 10.21037/atm.2018.07.18

Supplementary

Table S1 ICD-9CM and the CCS codes used to identify primary discharge diagnoses, baseline comorbidities, and in-hospital complications

Diagnosis	ICD-9-CM codes	CCS codes
Influenza		123
Influenza-pneumonia	487.0, 488.01, 488.11, 488.81	
Other respiratory manifestations	487.1, 488.02, 488.12, 488.82	
Other manifestations	487.8, 488.09, 488.19, 488.89	
Comorbidities		
Dementia	290.xx, 294.1x, 294.2x, 331.0, 331.1x, 331.82, 331.89	
Atrial fibrillation	427.31	
In-hospital complications		
Sepsis	788.52	2
Respiratory failure		131
Gastrointestinal hemorrhage		153
Acute myocardial infarction	410.x1, 411.1	
Acute kidney injury	584.x	
Ischemic stroke and transient cerebral ischemia	433.x1	112

x, the ICD-9CM subcode; ICD-9CM, International Classification of Diseases, Ninth Edition, Clinical Modification; CCS, Clinical Classification Software.

Table S2 ICD-9CM and the CCS codes in the primary discharge diagnosis position used to identify causes of 30-day readmissions

Organ system/diagnosis	CCS codes	ICD-9-CM codes
Respiratory	56, 122–134	
Pneumonia and empyema	122	
Influenza	123	
COPD and asthma	127, 128	
Aspiration pneumonitis	129	
Pleural disease		510.9, 511.0, 511.89, 511.9, 512.0, 512.89
Respiratory failure		518.8x
Other respiratory tract disease	125,126,133,134	
Cardiovascular	96–108, 245	
Heart failure	108	402.91, 404.91, 404.93, 788.51
Conduction disease and arrhythmia	105, 106, 107	
AMI	100	
CAD	101	414.04
Hypertension	98,99	
Pulmonary embolism		415.1x
Syncope	245	
Nonspecific chest pain	102	
Neurovascular	109–113	
Ischemic stroke and transient cerebral ischemia	112	434.11, 434.91
Peripheral vascular	114–121	
Hematologic and cancer-related	14, 15, 16, 18, 19, 24, 33, 35, 38, 39, 40, 42, 44, 45, 47, 59–63	
Endocrine and metabolic	50, 51, 58	
Gastrointestinal	138–141, 143–155	
GIB	153	
Renal diseases and fluid/electrolyte imbalance	55, 157,158, 160–163	
AKI		584.x
Fluid and electrolyte disorders	55	
Neuropsychiatric and substance-related	81, 83, 84, 85, 93, 95, 651, 653, 654, 657, 659, 660, 661	
MSK and injury related	54, 203–205, 207, 210–212, 226, 229–234, 238–244	
Symptoms		
Fever of unknown origin	246	
Nausea/vomiting	250	
Abdominal pain	251	
Malaise and fatigue	252	
Device, implant, graft complications	237	
Others	89, 92, 136, 137, 165–167, 171,175,182,187,189,193, 199, 253, 254, 257, 259	

x, the ICD-9CM subcode; ICD-9CM, International Classification of Diseases, Ninth Edition, Clinical Modification; CCS, Clinical Classification Software; COPD, chronic obstructive lung disease; AMI, acute myocardial infarction; CAD, coronary artery disease; UTI, urinary tract infection; AKI, acute kidney injury.