

LETTERS

A patient with polytrauma, hypothermia and cardiac arrest after delayed mountain rescue

With interest, we followed the case report by Ting and Brown¹ of a severely hypothermic patient with trauma treated with extracorporeal life support. We fully agree with the authors that extracorporeal life support should be taken into consideration in distinct cases of multiple trauma and severe hypothermia. We report a case of a hypothermic patient with polytrauma who arrived in cardiac arrest after a fall of 30 m in the mountains.²

Owing to bad weather, the helicopter emergency medical service team had reached the 59-year-old man four hours after the fall occurred. The patient had signs of cardiac instability and a Glasgow Coma Scale of 9/15. During transport to the level I trauma centre, he sustained cardiac arrest. On arrival, he was undergoing cardiopulmonary resuscitation, and had ventricular fibrillation, a core temperature of 25.3°C and a potassium level of 2.7 mmol/L.

The receiving team decided to proceed according to our Bernese Hypothermia Algorithm³ and to rewarm the patient, with limited trauma assessment. The assessment, which was done simultaneously with cannulation of the groin, included a primary survey and extended focused assessment with sonography in trauma to exclude major hemorrhage. As large bleeding was excluded, the team decided to start active invasive rewarming using minimal extracorporeal circulation.⁴ During rewarming, the patient underwent a full trauma assessment using our local polytrauma protocol. The contrast computed tomography scan — among others — showed serial rib frac-

tures with hemopneumothorax, an open book pelvic fracture, a pertrochanteric femur fracture and an active hemorrhage from a lumbar artery. The arterial bleeding was successfully embolized. Subsequently, the patient underwent several orthopedic interventions. Fourteen days after the fall, he was discharged home with full neurologic recovery.

Even in otherwise usually hopeless combinations of cardiac arrest in multiple trauma, a good outcome is possible if the cardiac arrest is caused by accidental hypothermia, as in our case. We hope to motivate colleagues to apply the concept of “no one is dead until he is warm and dead”⁵ even in patients with polytrauma, because rewarming could be the only strategy to improve coagulation and allow survival.⁶

Monika Brodmann Maeder MD MME

Emergency physician, Department of Emergency Medicine, Bern University Hospital and Institute of Mountain Emergency Medicine, EURAC rescue, Bozen, Italy

Volker Lischke MD

Anesthesiologist, Department of Anaesthesiology and Intensive Care Medicine, Hochtaunus-Kliniken gGmbH, Bad Homburg, Germany; German Red Cross Mountain Rescue, Berlin, Germany

Armin Berner MD

Anesthesiologist, Department of Anesthesia and Surgical Intensive Care, Hospital Garmisch-Partenkirchen, Garmisch-Partenkirchen, Germany; Mountain Rescue Bavaria, Bad Tolz, Germany

Oliver Reisten MD PhD

Anesthesiologist, Emergency Medical Service, Zermatt, Switzerland, and International Commission for Mountain Emergency Medicine (ICAR MEDCOM)

Urs Pietsch MD

Anesthesiologist, Department of Anaesthesiology and Intensive Care Medicine, Cantonal Hospital, St. Gallen, Switzerland; Air Zermatt, Zermatt, Switzerland

Mathieu Pasquier MD

Emergency physician, Emergency Service, Lausanne University Hospital Center, CHUV, Lausanne, Switzerland; International Commission for Mountain Emergency Medicine

■ Cite as: *CMAJ* 2018 October 22;190:E1263. doi: 10.1503/cmaj.70338

References

1. Ting DK, Brown DJ. Use of extracorporeal life support for active rewarming in a hypothermic, nonarrested patient with multiple trauma. *CMAJ* 2018;190:E718-21.
2. Lischke V, Berner A, Pietsch U, et al. Cardiac arrest in mountain areas during winter season. *Der Notarzt* 2014;30:58-65.
3. Monika BM, Martin D, Balthasar E, et al. The Bernese Hypothermia Algorithm: a consensus paper on in-hospital decision-making and treatment of patients in hypothermic cardiac arrest at an alpine level 1 trauma centre. *Injury* 2011;42:539-43.
4. Anastasiadis K, Murkin J, Antonitsis P, et al. Use of minimal invasive extracorporeal circulation in cardiac surgery: principles, definitions and potential benefits. A position paper from the Minimal Invasive Extra-Corporeal Technologies International Society (MiECTIS). *Interact Cardiovasc Thorac Surg* 2016;22:647-62.
5. Hilmo J, Naesheim T, Gilbert M. “Nobody is dead until warm and dead”: prolonged resuscitation is warranted in arrested hypothermic victims also in remote areas — a retrospective study from northern Norway. *Resuscitation* 2014;85:1204-11.
6. Paal P, Gordon L, Strapazzon G, et al. Accidental hypothermia—an update: The content of this review is endorsed by the International Commission for Mountain Emergency Medicine (ICAR MEDCOM). *Scand J Trauma Resusc Emerg Med* 2016;24:111.

Competing interests: None declared.

The authors have obtained patient consent.