

# Bacterial infections and cancer

Daphne van Elsland & Jacques Neefjes\* 

## Abstract

Infections are estimated to contribute to 20% of all human tumours. These are mainly caused by viruses, which explains why a direct bacterial contribution to cancer formation has been largely ignored. While epidemiological data link bacterial infections to particular cancers, tumour formation is generally assumed to be solely caused by the ensuing inflammation responses. Yet, many bacteria directly manipulate their host cell in various phases of their infection cycle. Such manipulations can affect host cell integrity and can contribute to cancer formation. We here describe how bacterial surface moieties, bacterial protein toxins and bacterial effector proteins can induce host cell DNA damage, and thereby can interfere with essential host cell signalling pathways involved in cell proliferation, apoptosis, differentiation and immune signalling.

**Keywords** bacteria; cancer; effectors; infection; signalling

**DOI** 10.15252/embr.201846632 | Received 25 June 2018 | Revised 10 August 2018 | Accepted 24 September 2018 | Published online 22 October 2018

**EMBO Reports (2018) 19: e46632**

See the Glossary for abbreviations used in this article.

## Introduction

Cancer development is the result of a series of genetic modifications that alter the normal control of cell growth and survival. These genetic alterations can be induced by a wide variety of external factors [1], including smoking, alcohol [2] and sunlight [3,4]. At least 75% of the head and neck cancers are caused by tobacco and alcohol [5] and 65–86% of the skin cancer risk can be attributed to sun exposure [4]. In addition to these external factors, viral genomes have been retrieved from a variety of tumour samples [6] and this link has been further substantiated by many epidemiological studies (Table 1). For example, viral infections such as human papillomavirus and hepatitis B virus and hepatitis C virus have been associated with ~90% of cervical cancer cases [7] and ~80% of hepatocellular carcinoma cases [8], respectively.

An even more compelling case for the link between viral infections and cancer arose from experiments showing that viruses exploit the host cell niche for their infection cycle and as a result stimulate mammalian growth-inducing genes, leaving the cells in a

cancerous state of uncontrolled cell division. It is now understood how viruses such as hepatitis B virus and human papillomavirus types 5 and 8 cause cellular transformation by inducing genetic instability through viral integration and through the activation of a large number of signalling pathways and cellular genes involved in oncogenesis, proliferation, inflammation and immune responses [9,10].

Viruses do, however, represent only one segment of the microbiome that exploits the mammalian host during its infection cycle. Pathogenic moulds, helminths and bacteria intensively interact with mammalian host cells to ensure their survival. Although these microorganisms usually do not leave a genetically recognizable trait or piggyback on mammalian genes, such as illustrated by viral infections, strong epidemiological links exist between various microbiological infections and cancers (Table 1). Examples include connections between *Schistosoma haematobium* infections and bladder cancer [11], *Helicobacter pylori* (*H. pylori*) infections and gastric cancer [12], chronic *Salmonella* Typhi (*S. Typhi*) infections and gallbladder carcinoma [13], and *Salmonella* Enteritidis (*S. Enteritidis*) infections and colon carcinoma [14]. Moreover, studies in germ-free and antibiotic-treated animals have indicated cancer-promoting effects of microbiota in various experimental systems, varying from gastric [15,16], colon [17,18] and liver [19] cancers.

However, since microbiome–host interactions are extremely diverse, their exact contributions to cancer development are hard to pinpoint. Especially, pathogenic bacteria have been shown to manipulate and exploit the human host cell niche in various ways throughout various stages of their infection cycle. In this review, we will discuss how bacterial surface moieties, bacterial protein toxins and bacterial effector proteins interact with host cells, and how such encounters can result in the modification of essential host cell signalling pathways involved in cancer formation.

## Bacterial cell-surface components and cancer development

The bacterial outer surface directly contacts host cells and consists of complex structures that include various antigenic moieties that activate host innate and adaptive immune responses. As a consequence, pathogenic bacteria have evolved a wide variety of outer-surface modifications that ensure immune escape to afford significant survival opportunities. To abolish immune recognition and clearance, Gram-negative bacteria cover their complex outer-surface macromolecules with a polysaccharide-rich capsule. These capsules

## Glossary

<b>Apc</b>	Adenomatous polyposis coli
<b>BFT</b>	<i>Bacteroides fragilis</i> toxin
<b>CagA</b>	Cytotoxin-associated gene A
<b>CCL5</b>	Chemokine (C-C motif) ligand 5
<b>CDK1</b>	Cyclin-dependent kinase 1
<b>CDT</b>	Cytolethal distending toxin
<b>DDR</b>	DNA damage responses
<b>DSBs</b>	Double-strand DNA breaks
<b>EF-2</b>	Elongation factor 2
<b>EGFR</b>	Epidermal growth factor receptor
<b>ER</b>	Endoplasmic reticulum
<b>ERK</b>	Extracellular signal-regulated kinase
<b>FadA</b>	Fusobacterium adhesion A
<b>FCP</b>	<i>Francisella</i> -containing phagosome
<b>IKK</b>	I $\kappa$ B kinase
<b>IL</b>	Interleukin
<b>JNK</b>	C-Jun N-terminal kinase
<b>LPS</b>	Lipopolysaccharides
<b>MALT</b>	Mucosa-associated lymphoid tissue
<b>MAPK</b>	Mitogen-activated protein kinase
<b>MEK</b>	Mitogen-activated protein kinase kinase
<b>MyD88</b>	Myeloid differentiation primary response 88
<b>NET1</b>	Neuroepithelial cell-transforming gene 1 protein
<b>NF-<math>\kappa</math>B</b>	Nuclear factor- $\kappa$ B
<b>OipA</b>	Outer inflammatory protein A
<b>PAK</b>	p21-activated kinase
<b>Pks</b>	Polyketide synthetase
<b>Raf</b>	Rapidly Accelerated Fibrosarcoma
<b>SCV</b>	<i>Salmonella</i> -containing Vacuole
<b>Tcf</b>	T-cell factor
<b>TLR</b>	Toll-like receptor
<b>VacA</b>	Vacuolating cytotoxin A

limit complement activation by shielding deeper structures on the membranes of pathogenic variants of *Escherichia coli* (*E. coli*), *Streptococcus pneumoniae*, *Haemophilus influenzae* type b, *Neisseria meningitidis* and others, and prevent engulfment by professional phagocytes [20–23]. Unencapsulated mutants of these bacteria rarely cause an invasive infection and are highly attenuated in various infection models due to better opsonophagocytic clearance [22,24,25].

In addition to their shielding capsules, many bacterial pathogens have modified their surface-exposed molecules, including lipopolysaccharides (LPS), flagella and peptidoglycans, to limit immune recognition. For example, *H. pylori* has LPS surface molecules that harbour “underacylated” lipid A molecules that are a poor substrate for host Toll-like receptor (TLR)4 and as such evade innate immune sensing [26,27]. *Helicobacter pylori* also produces modified flagellin molecules that are not recognized by TLR5 to prevent TLR5-mediated interleukin (IL)-8 secretion and subsequent immune signalling [28]. *Salmonella typhimurium* (*S. typhimurium*) expresses lipid A deacetylase PagL and a lipid A palmitoyltransferase PagP to modify lipid A, resulting in a 100-fold decrease in lipid A-mediated TLR4 activation and nuclear factor- $\kappa$ B (NF- $\kappa$ B) activation [29]. These examples illustrate how bacterial pathogens modify their outer surface to escape immune recognition.

Pathogenic bacteria that favour an intracellular lifestyle express surface proteins that promote both host cell attachment and internalization. For example, pathogenic species of the *Neisseria* family

express a variety of surface adhesins that mediate selective interaction with certain cell types, thereby allowing the exploitation of specialized host cell niches [30]. In a similar fashion, fibronectin-binding proteins of *Staphylococcus aureus* and *Borrelia burgdorferi* mediate the interaction between bacterium and host cell through the formation of tandem  $\beta$ -zippers that stimulate bacterial engulfment by non-phagocytic cells [31,32].

In general, these surface-mediated assault strategies are aimed at facilitating bacterial survival within the host through both immune evasion and host invasion. However, to further control the host cell machinery, bacterial surface molecules also manipulate host cell signalling cascades and affect host cell integrity, which can coincidentally induce cellular malignancies. CagL is a type IV pilus adhesin of *H. pylori* that ensures the adherence of *H. pylori* to gastric epithelial cells and then controls a signalling cascade that induces upregulation of gastrin secretion. This results in hypergastrinemia, a major risk factor for the development of gastric adenocarcinoma. CagL binds  $\beta$ 5-integrin thus manipulating integrin-linked kinase complexes and the downstream rapidly accelerated fibrosarcoma (Raf) kinase, the mitogen-activated protein kinase kinase (MEK) and the extracellular signal-regulated kinase (ERK) pathways (Fig 1A) [33]. The outer inflammatory protein A (OipA) of *H. pylori* activates EGFR (epidermal growth factor receptor) and stimulates Akt and  $\beta$ -catenin signalling, a phenotype observed in a number of different cancers, including gastric cancer (Fig 1B) [34,35]. OipA inactivation decreases  $\beta$ -catenin nuclear localization *in vitro* and reduces the incidence of cancer in animal models [36]. In addition, the blood group antigen-binding adhesin BabA of *H. pylori* can bind human Lewis(b) surface epitopes which indirectly increases mRNA levels of proinflammatory cytokines chemokine (C-C motif) ligand 5 (CCL5) and IL-8, and the precancer-related factors CDX2 and MUC2 (Fig 1C) [37]. The fusobacterium adhesion A (FadA) of *Fusobacterium nucleatum* (*F. nucleatum*) can bind the extracellular domain of E-cadherin, thereby inducing phosphorylation and internalization of E-cadherin. This then releases  $\beta$ -catenin to activate  $\beta$ -catenin–T-cell factor (Tcf)/LEF, downstream in the Wnt signalling pathway to control transcription of genes involved in apoptosis, cell proliferation and transformation (Fig 1D) [38]. In patients with colon adenomas or adenocarcinomas, high expression levels of *F. nucleatum* fadA have been associated with upregulated expression of oncogenic and inflammatory genes associated with the Wnt signalling pathway [39,40].

The major surface-exposed component of Gram-negative bacteria, LPS additionally activates signalling cascades that promote cancer development. LPS is present in both pathogenic and commensal bacteria and plays a central role in the activation of TLR4. TLR4-mediated signalling is critical for the downstream activation of numerous signalling pathways that underlie a variety of inflammatory and immune responses, and can promote the development of adenomatous polyposis coli (Apc)-dependent colorectal cancers and inflammation-associated colorectal cancers in mice. The role of TLR signalling in intestinal tumorigenesis has been studied through the crossing of myeloid differentiation primary response 88 (MyD88)-deficient mice that have impaired TLR4 signalling, with *Apc* (*Apc*<sup>Min/+</sup>) mice that mimic sporadic cancer and familial adenomatous polyposis. These MYD88-deficient  $\times$  *Apc*<sup>Min/+</sup> mice showed a reduction in both tumour number and size compared to the

**Table 1. Epidemiological and experimental evidence for a link between microbial infections and cancer.**

Infectious agent	Type of micro-organism	Cancer type
Epstein–Barr virus	Virus	Nasopharyngeal carcinoma, Burkitt lymphoma, immune suppression-related non-Hodgkin lymphoma, Hodgkin lymphoma, extranodal natural killer/T-cell lymphoma (nasal type) [102]
Hepatitis B virus	Virus	Hepatocellular carcinoma [102]
Hepatitis C virus	Virus	Hepatocellular carcinoma, non-Hodgkin lymphoma [102]
Kaposi sarcoma herpesvirus	Virus	Kaposi sarcoma, primary effusion lymphoma [102]
Human immunodeficiency virus 1	Virus	Kaposi sarcoma, non-Hodgkin lymphoma, Hodgkin lymphoma, carcinoma of the cervix, anus, conjunctiva [102]
Human papillomavirus type 16	Virus	Carcinoma of the cervix, vulva, vagina, penis, anus, oral cavity, and oropharynx and tonsil [102]
Human T-cell lymphotropic virus type 1	Virus	Adult T-cell leukaemia and lymphoma [102]
Merkel cell polyomavirus	Virus	Merkel cell carcinoma [103]
<i>Opisthorchis viverrini</i>	Trematode	Cholangiocarcinoma [102]
<i>Clonorchis sinensis</i>	Helminth	Cholangiocarcinoma [102]
<i>Schistosoma haematobium</i>	Trematode	Urinary bladder cancer [102]
<i>Helicobacter pylori</i>	Bacterium	Non-cardia gastric carcinoma, low-grade B-cell MALT gastric lymphoma [102]
Alfatoxin (B1)	Mould ( <i>Aspergillus flavus</i> )	Liver cancer [102]
<i>Salmonella</i> Typhi	Bacterium	Gallbladder carcinoma [13]
<i>Salmonella</i> Enteritidis	Bacterium	Colon carcinoma in the ascending and transverse parts of the colon [14]
<i>Chlamydia trachomatis</i>	Bacterium	Carcinoma of the cervix and ovaries [104,105]


*Apc*<sup>Min/+</sup> control mice, suggesting that TLR4 signalling further promotes tumour growth [41,42]. Tumour tissues of mice lacking MyD88 showed lower expression of the Cox2 gene that is involved in inflammation, indicating a role of this gene in reduced tumour formation [43]. It has furthermore been shown that Cox2 inhibitors, such as aspirin, reduce colorectal cancer risk in people that overexpress the 15-PGDH gene which encodes for an enzyme that disrupts Cox2 activity [44]. Studies with germ-free and wild-type mice showed that TLR4 activation by LPS from the intestinal microbiota pool contributes to the promotion of injury- and inflammation-driven hepatocellular carcinoma by activating proliferative and anti-apoptotic signals [19]. Findings from these animal studies were further corroborated by human studies in which enhanced expression of the TLR4/MyD88 complex was detected in 20% of colorectal patient samples [45].

### Bacterial toxin-mediated host cell transformation

To ensure immune escape, rapid replication and spreading, pathogenic bacteria do not only use immune-evasion strategies to avoid host cell clearance, but are also capable of immune cell elimination. One of the strategies employed by bacteria is the secretion of protein toxins that have cytolytic properties. Bacteria can express protein toxins from their pathogenicity islands and secrete them through specialized secretion systems for transport across bacterial outer membranes [46]. The interaction of proteins toxins with the host generally occurs in an ordered series of events and can be illustrated by the mode of action of the diphtheria toxin that inhibits the synthesis of host cell proteins through the inactivation of the host

elongation factor 2 (EF-2) protein. The diphtheria toxin consists of three subunits and is secreted by *Corynebacterium diphtheriae* as a single polypeptide chain. Diphtheria toxin then binds to the host's heparin-binding epidermal growth factor-like surface receptor that then is internalized in the endosomal system. Here, the transmembrane domain of the toxin is unfolded, which translocates the toxin to the cytosolic side of the endosomal membrane. This is followed by a reduction in the disulphide bond between toxin fragments A and B and release of the C-domain into the cytoplasm. The C-domain is then refolded into an enzymatically active conformation that catalyses NAD<sup>+</sup>-dependent ADP-ribosylation of EF-2. This then inhibits protein synthesis, ultimately resulting in cell death of the targeted cells [47].

Although pathogenic bacteria primarily use their toxin-mediated assault strategies to create a favourable host cell environment, their toxins, likely as a side effect of their mode of action, can also contribute to carcinogenesis. Toxin-mediated carcinogenesis can occur in multiple ways, including the induction of genomic instability, the induction of cell death resistance cell signalling and the induction of proliferative signalling [48]. Genome instability is most readily caused by bacterial protein toxins that induce host cell double-stranded DNA breaks, including the cytolethal distending toxin (CDT), the colibactin, the Shiga toxin and endonucleases. CDT is secreted by various Gram-negative bacteria that belong to the Gamma and Epsilon class of Proteobacteria, including *S. Typhi*, *E. coli*, *Shigella dysenteriae* and *Campylobacter jejuni*. CDT is comprised of three subunits, CdtA, CdtB and CdtC. CdtA and CdtC ensure the uptake and cellular delivery of CdtB, which harbours the catalytic activity of CDT and causes double-strand DNA breaks (DSBs) in host cells. After host cell binding and internalization by




**Figure 1. Bacterial outer-surface components that manipulate host cell signalling cascades involved in cellular malignancy.**

(A) *Helicobacter pylori* CagL binds  $\beta 5$ -integrin and induces downstream signalling of Raf, MEK and ERK pathways that play a central role in *H. pylori*-induced gastrin production and cellular transformation. (B) *H. pylori* OipA activates EGFR and stimulates Akt and  $\beta$ -catenin signalling, causing cell proliferation. (C) *H. pylori* BabA binds human Lewis(b) surface epitopes which increases levels of CCL5, IL-8, CDX2 and MUC2, causing cell proliferation. (D) *Fusobacterium nucleatum* FadA binds to E-cadherin, which releases  $\beta$ -catenin that activates transcription factor Tcf/LEF which controls the transcription of genes involved in apoptosis, cell proliferation and transformation.

subunits CdtA and CdtC, CdtB undergoes retrograde transport via the endosomes and Golgi to the endoplasmic reticulum (ER), where it undergoes ER-associated protein degradation-mediated translocation into the cytosol. The CdtB subunit is then imported in the

nucleus where it induces DSBs [49]. These DSBs result in DNA damage responses (DDR) that cause G1-S cell cycle arrest in endothelial and epithelial cells, and both G1-S and G2-M cell cycle arrest in fibroblasts and apoptosis in haematopoietic cells that are



**Figure 2. Host cell signalling pathways involved in cell growth and transformation manipulated by bacterial toxins.**

(A) The CdtB subunit of the CDT toxin is delivered to the nucleus where it causes DSBs and impairs DNA DDR sensor functionality. At the same time, NET1 and RhoA are activated which ensure upregulation of MAPK and cellular survival. (B) *Bacteroides fragilis* BFT binds to E-cadherin and involved in cellular signalling, proliferation and transformation via activation of the  $\beta$ -catenin/Wnt and NF- $\kappa$ B signalling pathways.

particularly sensitive to these toxins. As a result, this toxin can locally eliminate immune cells, providing an obvious advantage for the bacteria. However, prolonged exposure to sublethal doses of CDT can impair DDR sensor functionality, resulting in impaired detection of DNA damage and the accumulation of mutations. At the same time, mitogen-activated protein kinase (MAPK) activity is upregulated by activation of the neuroepithelial cell-transforming gene 1 protein (NET1) and the GTPase RhoA, which supports survival of the toxin-exposed cells (Fig 2A) [50]. As a consequence, these cells can propagate with DNA mutations and deletions that arise during the repair process, thus inducing genomic errors that underlie cancer formation.

In addition to the CDT toxins, the DNA interacting colibactin toxin has also been associated with the formation of DSBs and the introduction of genomic instability. Colibactin is secreted by *E. coli* strains of the phylogenetic group B2 that harbours the polyketide synthetase (pks) island [51]. Bacteria that harbour the pks genomic island are able to induce DSBs in eukaryotic cells, which results in the activation of the DNA damage checkpoint pathways ATM, CHK1 and CHK2. This then results in CDC25 and cyclin-dependent kinase 1 (CDK1)-mediated G<sub>2</sub>- to M-phase cell cycle arrest and finally in apoptotic cell death. As a side effect of their mode of action, colibactin-producing bacteria also induce incomplete DNA repair, chromosomal instability and anchorage-dependent colony formation, phenotypes that can promote cancer formation [52,53]. This is further substantiated by epidemiological studies showing that colibactin-producing *E. coli* bacteria appear with high prevalence in biopsies of patients with human colorectal tumours [54,55]. Moreover, colitis-susceptible IL-10-deficient mice showed increased formation of invasive carcinoma when colonized with *E. coli*

secreting colibactin, whereas deletion of the pks genotoxic island from these *E. coli* strains decreased tumour multiplicity and invasion [56].

Besides toxins that contribute to carcinogenesis by introducing DSBs and genomic instability, toxins have been reported that promote carcinogenesis by inducing resistance to cell death signalling and by promoting proliferative signalling. These toxins are generally secreted by pathogenic bacteria that favour an intracellular host cell life as part of their infectious cycle and thus directly benefit from host cell survival. An example of such a toxin is the *Bacteroides fragilis* (*B. fragilis*) toxin (BFT) that binds to intestinal epithelial cell receptors and stimulates cell proliferation by cleavage of the tumour suppressor protein E-cadherin [57,58]. E-cadherin is involved in the formation of intercellular adhesion junctions in the intestinal epithelium and is involved in cellular signalling, proliferation and differentiation via activation of the  $\beta$ -catenin/Wnt and NF- $\kappa$ B signalling pathways (Fig 2B) [59–61]. BFT induced acute and chronic colitis in C57BL/six mice, and colon tumours in the multiple intestinal neoplasia (*Apc*<sup>Min/+</sup>) mouse model for human colon carcinoma. This is the same mouse model where *H. pylori* triggers a pro-carcinogenic multi-step inflammatory cascade that requires IL-17R, NF- $\kappa$ B and STAT3 signalling in colonic epithelial cells [62,63]. These mouse experiments are further substantiated by epidemiology, indicating that infections with enterotoxigenic variants of *B. fragilis*, as opposed to non-toxigenic variants, are more prevalent in people with colorectal cancers. More specifically, the enterotoxigenic variant is present in only 10–20% of the healthy population, whereas 40% of CRC patients present enterotoxigenic *B. fragilis* in their faeces [64]. In addition to BFT, multiple biologically plausible mechanisms have been reported that explain how the vacuolating



cytotoxin A (VacA) of *H. pylori* enhances gastric cancer risk. Similar as the *H. pylori* outer membrane protein OipA, VacA activates the EGFR receptor that triggers PI3K–Akt signalling, and inactivates glycogen synthase kinase 3 $\beta$  [34,65]. As a result,  $\beta$ -catenin degradation is abolished, which promotes Tcf/LEF-controlled transcription that promotes cell growth and transformation [34,65,66]. Another *H. pylori* virulence factor, cytotoxin-associated gene A (CagA), which depends on the type IV pilus cell-surface adhesion CagL for its host cell targeting, interacts with the c-Met receptor to activate epithelial proliferation, as shown in human gastric organoids [67]. Phosphorylated and unphosphorylated CagA can also interact with a variety of host proteins involved in the MEK, ERK, NF- $\kappa$ B and  $\beta$ -catenin pathways that are all involved in host cell proliferation and cancer formation [68,69].

### Bacterial effector proteins that mediate host cell transformation

Various intracellular bacterial pathogens have developed molecular mechanisms to ensure a persistent infection within the protective environment of the host cell's interior. This requires host cell control at various steps of the intracellular infection cycle, including host cell internalization through receptor-mediated endocytosis or phagocytosis, intracellular survival and growth, and release from the infected host cell.


After host cell internalization bacterial-cargo generally routes across the endosomal system that usually terminates in a highly degradative organelle, the phagolysosome. To avoid phagolysosomal degradation, intracellular bacterial pathogens have evolved various mechanisms that can be broadly grouped into pathways where pathogenic bacteria either escape the phagosome or enter in the cytosol, and pathways where the phagosome is hijacked and tailored to the preferences of the bacteria. Cytosolic pathogens like *Listeria*, *Shigella flexneri* (*S. flexneri*), *Rickettsia* and *Francisella* are known to rapidly escape the phagosome to enter the host cytosol and thereby avoid lysosomal fusion and degradation [70]. This generally involves secretion of bacterial effector proteins that induce pore formation of the endolysosomal vacuole and ensure its subsequent rupture. It has, for example, been shown that *S. flexneri* secretes the effector protein Invasion plasmid antigen B that forms ion channels in eukaryotic membranes and can mediate potassium influx and subsequent endolysosomal leakage [71]. In addition, *Listeria* can secrete the listeriolysin-O protein that induces small-membrane perforations, which causes Ca<sup>2+</sup> leakage from vacuoles and an increase in the vacuolar pH. Subsequently, vacuolar maturation is prevented [72,73]. *Francisella tularensis* (*F. tularensis*) also escapes into the host cytoplasm. After phagocytic uptake by macrophages, *F. tularensis* resides in the *Francisella*-containing phagosome (FCP) that over time matures from a phagosome with an early endosomal character into a more acidic late endosomal phagosome. Since inhibition of FCP acidification delays the escape of *F. tularensis* into the cytosol, further acidification during phagosome maturation apparently stimulates *F. tularensis* to express unique, as-yet-undefined factors to disrupt the phagosomal membrane [74–76].

In contrast to bacteria that escape the phagosome, pathogenic bacteria have been reported that hijack the phagosome to ensure a favourable replication niche. An example of such a pathogen is

*Legionella pneumophila* that redirects the *Legionella*-containing phagosome to the ER via the secretion of bacterial proteins through the Dot-Icm secretion system. This rearrangement prevents lysosomal degradation and ensures *Legionella* replication within the phagosome [77,78]. Bacterial control of phagosomal maturation has also been reported for the intracellular pathogen *Salmonella*. After its host cell internalization, *Salmonella* ends up in a membrane-bound phagosome-like vacuolar compartment called the *Salmonella*-containing vacuole (SCV). The SCV then matures and acquires characteristics of late endocytic compartments including acidification. It does, however, not become bactericidal. Under control of the *Salmonella* effectors, SifA, SseJ, SseG, SseF, SopD2, and PipB2, cellular host processes are manipulated to turn the SCV into a compartment that facilitates *Salmonella* replication [79]. SifA, which is critical in this process [80], interacts with the host cell effector of the GTPase Arl8b, the SifA and kinesin-interacting protein SKIP. This interaction results in the formation of a tubular membrane network, known as *Salmonella*-induced filaments, that is essential for the supply of nutrients to the SCV and prevents endosomal antimicrobial activities due to constant mixing of antimicrobial agents with late endosomes and lysosomes [81,82].

Intracellular pathogenic bacteria that engage effector proteins during their intracellular life cycle manipulate host cell integrity in a major way. To this end, some of these infections have been epidemiologically linked to particular cancer types. Infections by two food-borne *Salmonella* serovars, *S. Typhi* and *S. Enteritidis*, are linked to gallbladder carcinoma and colon cancer, respectively [13,14]. These bacteria introduce a series of effector proteins in the host cell to take over host cell biology and—depending on host pathway affected—can contribute to cancer formation. A *Salmonella* effector protein that has been linked to colon cancer formation is the acetyltransferase AvrA that alters a variety of host-signalling pathways and modulates immune responses, apoptosis and proliferation [83,84]. AvrA modifies and stabilizes  $\beta$ -catenin, thereby enhancing signalling and promoting epithelial cell proliferation (Fig 3A) [85–87]. AvrA also suppresses the host immune system and its apoptotic defences via the inhibition of the c-Jun N-terminal kinase (JNK) and NF- $\kappa$ B signalling pathways (Fig 3A) [88]. In addition to AvrA, three AvrA orthologues have been reported that similarly interact with essential host cell signalling pathways. However, in contrast to AvrA these orthologues have primarily only inhibitory effects on the host immune system. YopJ is expressed by *Yersinia pestis* and attenuates the ERK, p38, JNK and I $\kappa$ B kinase (IKK) pathways involved in the synthesis of cytokines as well as anti-apoptotic factors [89]. VopA of *Vibrio parahaemolyticus* can similarly inhibit host ERK, p38 and JNK signalling, but not the IKK pathway [90,91], and AopP of *Aeromonas salmonicida* interacts with the IKK pathway [92].

In epithelial cells infected with *S. typhimurium*, the effector proteins SopE, SopE2 and SopB can manipulate host Rho-family GTPases, p21-activated kinase (PAK) and ABL tyrosine kinase to activate STAT3 and alter transcription regulation, which [93] can mediate transformation of cells (Fig 3B). In addition, cellular transformation can occur through *Salmonella* effector SopE, SopE2, SopB and SptP-mediated activation of the MAPK and AKT pathways (Fig 3B) [94]. The activation of these signalling pathways enables the transformation of fibroblasts and gallbladder organoids that harbour a pre-transformed phenotype whereby the tumour suppressor gene p53 is inactivated and the MYC oncogene is overexpressed. These findings



**Figure 3. Examples of bacterial effector proteins involved in cellular transformation.**

(A) *Salmonella* Enteritidis AvrA stabilizes  $\beta$ -catenin, which results in proliferative Wnt signalling. At the same time, AvrA inhibits JNK and NF- $\kappa$ B signalling pathways involved in inflammation and apoptosis. (B) The effector proteins SopB, SopE and SopE2 of *Salmonella typhimurium* activate the small GTPases RhoG, Rac1 and Cdc42 and activate members of the PAK that phosphorylate members of the Abl kinase family, leading to the activation the cytoplasmic transcription factor STAT3, which contributes to cellular transformation. The effector proteins SopB, SopE, SopE2 and SptP of *S. typhimurium* additionally mediate activation of the MAPK and Akt pathways, which transforms premutated fibroblasts and gallbladder organoids.

are supported by pathology on gallbladder carcinoma samples from Indian patients that contain both *S. Typhi* DNA and the pre-transformed modifications also observed in the laboratory experiments, and by an *Apc*<sup>Min/+</sup> mouse model in which oral infection with *S. typhimurium* results in the development colorectal adenocarcinomas in a *Salmonella* effector-dependent manner [13].

## Conclusions

Although bacterially induced host cell manipulation can promote cancer formation, it is unlikely that bacterial pathogens themselves experience any evolutionary benefit from their carcinogenic actions. Bacterially induced cancer formation is more likely an unfortunate

**Box 1: In need of answers**

- (i) Do bacterial infections only decrease the threshold for cellular transformation or can they also initiate tumour formation?
- (ii) How is transformation by activation of host signalling pathways imprinted in host cells?
- (iii) How can correlations from microbiome studies be translated to causalities?
- (iv) Does transformed tissue cause microbial dysbiosis [100]?
- (v) It has been shown that there is a distal oncogenic effect of the gut microbiome [101]. How does the gut microbiome affect tumour formation at a distance?
- (vi) What is the total contribution of bacteria to cancer formation?
- (vii) How to translate the collective knowledge on bacteria and cancer formation into treatment or prevention measures?

consequence of the bacterial infection cycle since cancer usually occurs long after the bacterium and its effectors have left the host [13,14]. Moreover, bacterial host cell manipulations involved in the induction of cancer formation usually account for only one step in the multi-step process required for actual cellular transformation and cancer formation. This can be illustrated by *Salmonella* infections that only in combination with pre-mutations allow cellular transformation in tissue culture fibroblasts and gallbladder organoids and is supported by observations of Indian gallbladder cancer patients who showed the corresponding pre-mutations in the p53 gene, c-MYC amplification in their tumours and had a history of *S. Typhi* infection. [11] In other words, *Salmonella* will only induce cancer when the cell has made already one or multiple pretransforming steps. This would explain why chronic bacterial infections have a higher statistical chance of initiating tumorigenesis as the likelihood of encountering a pre-transformed cell would then be markedly increased. This may also explain the correlations of persistent *Mycobacterium tuberculosis* infections and pulmonary cancers [95] and chronic *Coxiella burnetii* infections and B-cell non-Hodgkin lymphoma [96].

Since many epidemiological studies reveal a link between bacterial infections and cancer incidence, and the number of bacterial mechanisms that can contribute to cellular transformation are most likely considerably larger than reported to date, we expect that the number of examples illustrating the role of bacterial infections in cancer formation will increase the coming years. It is also known that bacterial effectors from different species can act synergistically during host cell manipulation and then act in a symbiotic interspecies manner [55,97]. These combined mechanisms can induce cell transformation and cancer in an even more complex manner and further contribute to the complexity of bacterial contributions to cancer.

While the first examples of bacterial mechanisms contributing to cancer are uncovered, it is likely that bacteria will provide many new and surprising mechanisms for host cell manipulation, some of which may participate in cell transformation. These may include an expansion of mechanisms involved in immune evasion, DNA damage and signalling pathways, but may also include more indirect routes, as, for example, via the formation of carcinogenic metabolites [98]. When the role of defined bacterial mechanisms in cancer formation will become more apparent and accepted (see also Box 1), studies on their prevention or control can help reduce

cancer formation. On this note, antibiotic therapy during cancer treatment, which is already a standard of care in patients with gastric mucosa-associated lymphoid tissue (MALT) lymphoma [99], might become a valuable addition to current tumour-targeting therapies. This, however, may only help when the presence of a bacterial species is required to continuously provide signals to maintain the transformed state. Otherwise, patients diagnosed with a bacterial pathogen known to participate in cancer formation—but not necessarily maintenance—may be incorporated in cancer screening programs.

**Acknowledgements**

This work is supported by an ERC Advanced grant and a grant from the Dutch Cancer Society KWF to JN.

**Conflict of interest**

The authors declare that they have no conflict of interest.

**References**

1. Wu S, Powers S, Zhu W, Hannun YA (2015) Substantial contribution of extrinsic risk factors to cancer development. *Nature* 529: 43
2. Siegel RL, Jacobs EJ, Newton CC, Feskanich D, Freedman ND, Prentice RL, Jemal A (2015) Deaths due to cigarette smoking for 12 smoking-related cancers in the United States. *JAMA Intern Med* 175: 1574–1576
3. Koh HK, Geller AC, Miller DR, Grossbart TA, Lew RA (1996) Prevention and early detection strategies for melanoma and skin cancer. Current status. *Arch Dermatol* 132: 436–443
4. Parkin DM, Mesher D, Sasieni P (2011) 13. Cancers attributable to solar (ultraviolet) radiation exposure in the UK in 2010. *Br J Cancer* 105 (Suppl 2): S66–S69
5. Blot WJ, McLaughlin JK, Winn DM, Austin DF, Greenberg RS, Preston-Martin S, Bernstein L, Schoenberg JB, Stemhagen A, Fraumeni JF Jr (1988) Smoking and drinking in relation to oral and pharyngeal cancer. *Can Res* 48: 3282–3287
6. Khoury JD, Tannir NM, Williams MD, Chen Y, Yao H, Zhang J, Thompson EJ, TCGA Network, Meric-Bernstam F, Medeiros LJ *et al* (2013) Landscape of DNA virus associations across human malignant cancers: analysis of 3,775 cases using RNA-seq. *J Virol* 87: 8916–8926
7. Bosch FX, Manos MM, Munoz N, Sherman M, Jansen AM, Peto J, Schiffman MH, Moreno V, Kurman R, Shah KV (1995) Prevalence of human papillomavirus in cervical cancer: a worldwide perspective. International biological study on cervical cancer (IBSCC) Study Group. *J Natl Cancer Inst* 87: 796–802
8. El-Serag HB (2012) Epidemiology of viral hepatitis and hepatocellular carcinoma. *Gastroenterology* 142: 1264–1273.e1
9. Neuveut C, Wei Y, Buendia MA (2010) Mechanisms of HBV-related hepatocarcinogenesis. *J Hepatol* 52: 594–604
10. Münger K, Baldwin A, Edwards KM, Hayakawa H, Nguyen CL, Owens M, Grace M, Huh K (2004) Mechanisms of human papillomavirus-induced oncogenesis. *J Virol* 78: 11451–11460
11. Mostafa MH, Sheweita SA, O'Connor PJ (1999) Relationship between schistosomiasis and bladder cancer. *Clin Microbiol Rev* 12: 97–111
12. Kikuchi S (2002) Epidemiology of *Helicobacter pylori* and gastric cancer. *Gastric Cancer* 5: 6–15



13. Scanu T, Spaapen RM, Bakker JM, Pratap CB, Wu LE, Hofland I, Broeks A, Shukla VK, Kumar M, Janssen H *et al* (2015) *Salmonella* manipulation of host signaling pathways provokes cellular transformation associated with gallbladder carcinoma. *Cell Host Microbe* 17: 763–774
14. Mughini-Gras L, Schaapveld M, Kramers J, Mooij S, Neeffes-Borst EA, Pelt WV, Neeffes J (2018) Increased colon cancer risk after severe *Salmonella* infection. *PLoS One* 13: e0189721
15. Lofgren JL, Whary MT, Ge Z, Muthupalani S, Taylor NS, Mobley M, Potter A, Varro A, Eibach D, Suerbaum S *et al* (2011) Lack of commensal flora in *Helicobacter pylori*-infected INS-GAS mice reduces gastritis and delays intraepithelial neoplasia. *Gastroenterology* 140: 210–220
16. Lee CW, Rickman B, Rogers AB, Ge Z, Wang TC, Fox JG (2008) *Helicobacter pylori* eradication prevents progression of gastric cancer in hypergastrinemic INS-GAS mice. *Can Res* 68: 3540–3548
17. Vannucci L, Stepankova R, Kozakova H, Fiserova A, Rossmann P, Tlaskalova-Hogenova H (2008) Colorectal carcinogenesis in germ-free and conventionally reared rats: different intestinal environments affect the systemic immunity. *Int J Oncol* 32: 609–617
18. Li Y, Kundu P, Seow SW, de Matos CT, Aronsson L, Chin KC, Karre K, Pettersson S, Greicius G (2012) Gut microbiota accelerate tumor growth via c-jun and STAT3 phosphorylation in APCMin/+ mice. *Carcinogenesis* 33: 1231–1238
19. Dapito Dianne H, Mencin A, Gwak G-Y, Pradere J-P, Jang M-K, Mederacke I, Caviglia Jorge M, Khiabanian H, Adeyemi A, Bataller R *et al* (2012) Promotion of hepatocellular carcinoma by the intestinal microbiota and TLR4. *Cancer Cell* 21: 504–516
20. Pluschke G, Mayden J, Achtman M, Levine RP (1983) Role of the capsule and the O antigen in resistance of O18:K1 *Escherichia coli* to complement-mediated killing. *Infect Immun* 42: 907–913
21. Abeyta M, Hardy GG, Yother J (2003) Genetic alteration of capsule type but not PspA type affects accessibility of surface-bound complement and surface antigens of *Streptococcus pneumoniae*. *Infect Immun* 71: 218–225
22. Brown EJ, Hosea SW, Frank MM (1983) The role of antibody and complement in the reticuloendothelial clearance of pneumococci from the bloodstream. *Rev Infect Dis* 5(Suppl 4): S797–S805
23. Winkelstein JA, Tomasz A (1978) Activation of the alternative complement pathway by pneumococcal cell wall teichoic acid. *J Immunol* 120: 174–178
24. Watson DA, Musher DM (1990) Interruption of capsule production in *Streptococcus pneumoniae* serotype 3 by insertion of transposon Tn916. *Infect Immun* 58: 3135–3138
25. Geno KA, Gilbert GL, Song JY, Skovsted IC, Klugman KP, Jones C, Konradsen HB, Nahm MH (2015) Pneumococcal capsules and their types: past, present, and future. *Clin Microbiol Rev* 28: 871–899
26. Mattsby-Baltzer I, Mielniczuk Z, Larsson L, Lindgren K, Goodwin S (1992) Lipid A in *Helicobacter pylori*. *Infect Immun* 60: 4383–4387
27. Tran AX, Stead CM, Trent MS (2005) Remodeling of *Helicobacter pylori* lipopolysaccharide. *J Endotoxin Res* 11: 161–166
28. Gewirtz AT, Yu Y, Krishna US, Israel DA, Lyons SL, Peek RM Jr (2004) *Helicobacter pylori* flagellin evades toll-like receptor 5-mediated innate immunity. *J Infect Dis* 189: 1914–1920
29. Kawasaki K, Ernst RK, Miller SI (2004) 3-O-deacylation of lipid A by PagL, a PhoP/PhoQ-regulated deacylase of *Salmonella typhimurium*, modulates signaling through toll-like receptor 4. *J Biol Chem* 279: 20044–20048
30. Popp A, Billker O, Rudel T (2001) Signal transduction pathways induced by virulence factors of *Neisseria gonorrhoeae*. *Int J Med Microbiol* 291: 307–314
31. Meenan NA, Visai L, Valtulina V, Schwarz-Linek U, Norris NC, Gurusiddappa S, Hook M, Speziale P, Potts JR (2007) The tandem beta-zipper model defines high affinity fibronectin-binding repeats within *Staphylococcus aureus* FnBPA. *J Biol Chem* 282: 25893–25902
32. Raibaud S, Schwarz-Linek U, Kim JH, Jenkins HT, Baines ER, Gurusiddappa S, Hook M, Potts JR (2005) *Borrelia burgdorferi* binds fibronectin through a tandem beta-zipper, a common mechanism of fibronectin binding in staphylococci, streptococci, and spirochetes. *J Biol Chem* 280: 18803–18809
33. Wiedemann T, Hofbauer S, Tegtmeyer N, Huber S, Sewald N, Wessler S, Backert S, Rieder G (2012) *Helicobacter pylori* CagL dependent induction of gastrin expression via a novel alphavbeta5-integrin-integrin linked kinase signalling complex. *Gut* 61: 986–996
34. Tabassam FH, Graham DY, Yamaoka Y (2009) *Helicobacter pylori* activate epidermal growth factor receptor- and phosphatidylinositol 3-OH kinase-dependent Akt and glycogen synthase kinase 3beta phosphorylation. *Cell Microbiol* 11: 70–82
35. Polk DB, Peek RM Jr (2010) *Helicobacter pylori*: gastric cancer and beyond. *Nat Rev Cancer* 10: 403–414
36. Franco AT, Johnston E, Krishna U, Yamaoka Y, Israel DA, Nagy TA, Wroblewski LE, Piazzuelo MB, Correa P, Peek RM Jr (2008) Regulation of gastric carcinogenesis by *Helicobacter pylori* virulence factors. *Can Res* 68: 379–387
37. Ishijima N, Suzuki M, Ashida H, Ichikawa Y, Kanegae Y, Saito I, Boren T, Haas R, Sasakawa C, Mimuro H (2011) BabA-mediated adherence is a potentiator of the *Helicobacter pylori* type IV secretion system activity. *J Biol Chem* 286: 25256–25264
38. Rubinstein MR, Wang X, Liu W, Hao Y, Cai G, Han YW (2013) *Fusobacterium nucleatum* promotes colorectal carcinogenesis by modulating E-cadherin/beta-catenin signaling via its FadA adhesin. *Cell Host Microbe* 14: 195–206
39. Kostic AD, Gevers D, Pedamallu CS, Michaud M, Duke F, Earl AM, Ojesina AI, Jung J, Bass AJ, Taberero J *et al* (2012) Genomic analysis identifies association of *Fusobacterium* with colorectal carcinoma. *Genome Res* 22: 292–298
40. Castellari M, Warren RL, Freeman JD, Dreolini L, Krzywinski M, Strauss J, Barnes R, Watson P, Allen-Vercoe E, Moore RA *et al* (2012) *Fusobacterium nucleatum* infection is prevalent in human colorectal carcinoma. *Genome Res* 22: 299–306
41. Coleman OI, Haller D (2017) Bacterial signaling at the intestinal epithelial interface in inflammation and cancer. *Front Immunol* 8: 1927
42. Rakoff-Nahoum S, Medzhitov R (2007) Regulation of spontaneous intestinal tumorigenesis through the adaptor protein MyD88. *Science* 317: 124–127
43. Abreu MT (2010) Toll-like receptor signalling in the intestinal epithelium: how bacterial recognition shapes intestinal function. *Nat Rev Immunol* 10: 131–144
44. Fink SP, Yamauchi M, Nishihara R, Jung S, Kuchiba A, Wu K, Cho E, Giovannucci E, Fuchs CS, Ogino S *et al* (2014) Aspirin and the risk of colorectal cancer in relation to the expression of 15-hydroxyprostaglandin dehydrogenase (15-PGDH, HPGD). *Sci Transl Med* 6: 233re2
45. Wang EL, Qian ZR, Nakasono M, Tanahashi T, Yoshimoto K, Bando Y, Kudo E, Shimada M, Sano T (2010) High expression of Toll-like receptor 4/myeloid differentiation factor 88 signals correlates with poor prognosis in colorectal cancer. *Br J Cancer* 102: 908–915
46. Costa TRD, Felisberto-Rodrigues C, Meir A, Prevost MS, Redzej A, Trokter M, Waksman G (2015) Secretion systems in Gram-negative bacteria: structural and mechanistic insights. *Nat Rev Microbiol* 13: 343

47. Murphy JR (2011) Mechanism of diphtheria toxin catalytic domain delivery to the eukaryotic cell cytosol and the cellular factors that directly participate in the process. *Toxins* 3: 294–308
48. Rosadi F, Fiorentini C, Fabbri A (2016) Bacterial protein toxins in human cancers. *Pathog Dis* 74: ftv105
49. Cortes-Bratti X, Chaves-Olarte E, Lagergard T, Thelestam M (2000) Cellular internalization of cytolethal distending toxin from *Haemophilus ducreyi*. *Infect Immun* 68: 6903–6911
50. Guerra L, Carr HS, Richter-Dahlfors A, Masucci MG, Thelestam M, Frost JA, Frisan T (2008) A bacterial cytotoxin identifies the RhoA exchange factor Net1 as a key effector in the response to DNA damage. *PLoS One* 3: e2254
51. Putze J, Hennequin C, Nougayrede JP, Zhang W, Homburg S, Karch H, Bringer MA, Fayolle C, Carniel E, Rabsch W *et al* (2009) Genetic structure and distribution of the colibactin genomic island among members of the family Enterobacteriaceae. *Infect Immun* 77: 4696–4703
52. Nougayrede JP, Homburg S, Taieb F, Boury M, Brzuszkiewicz E, Gottschalk G, Buchrieser C, Hacker J, Dobrindt U, Oswald E (2006) *Escherichia coli* induces DNA double-strand breaks in eukaryotic cells. *Science* 313: 848–851
53. Cuevas-Ramos G, Petit CR, Marcq I, Boury M, Oswald E, Nougayrede JP (2010) *Escherichia coli* induces DNA damage *in vivo* and triggers genomic instability in mammalian cells. *Proc Natl Acad Sci USA* 107: 11537–11542
54. Buc E, Dubois D, Sauvanet P, Raisch J, Delmas J, Darfeuille-Michaud A, Pezet D, Bonnet R (2013) High prevalence of mucosa-associated *E. coli* producing cyclomodulin and genotoxin in colon cancer. *PLoS One* 8: e56964
55. Dejea CM, Fathi P, Craig JM, Boleij A, Taddese R, Geis AL, Wu X, DeStefano Shields CE, Hechenbleikner EM, Huso DL *et al* (2018) Patients with familial adenomatous polyposis harbor colonic biofilms containing tumorigenic bacteria. *Science* 359: 592–597
56. Arthur JC, Perez-Chanona E, Muhlbauer M, Tomkovich S, Uronis JM, Fan TJ, Campbell BJ, Abujamel T, Dogan B, Rogers AB *et al* (2012) Intestinal inflammation targets cancer-inducing activity of the microbiota. *Science* 338: 120–123
57. Wu S, Rhee KJ, Zhang M, Franco A, Sears CL (2007) *Bacteroides fragilis* toxin stimulates intestinal epithelial cell shedding and gamma-secretase-dependent E-cadherin cleavage. *J Cell Sci* 120: 1944–1952
58. Sears CL (2009) Enterotoxigenic *Bacteroides fragilis*: a rogue among symbiotes. *Clin Microbiol Rev* 22: 349–369, Table of Contents
59. Wu S, Lim KC, Huang J, Saidi RF, Sears CL (1998) *Bacteroides fragilis* enterotoxin cleaves the zonula adherens protein, E-cadherin. *Proc Natl Acad Sci USA* 95: 14979–14984
60. Wu S, Powell J, Mathioudakis N, Kane S, Fernandez E, Sears CL (2004) *Bacteroides fragilis* enterotoxin induces intestinal epithelial cell secretion of interleukin-8 through mitogen-activated protein kinases and a tyrosine kinase-regulated nuclear factor-kappaB pathway. *Infect Immun* 72: 5832–5839
61. Nelson WJ, Nusse R (2004) Convergence of Wnt, beta-catenin, and cadherin pathways. *Science* 303: 1483–1487
62. Wu S, Rhee KJ, Albesiano E, Rabizadeh S, Wu X, Yen HR, Huso DL, Brancati FL, Wick E, McAllister F *et al* (2009) A human colonic commensal promotes colon tumorigenesis via activation of T helper type 17 T cell responses. *Nat Med* 15: 1016–1022
63. Rhee KJ, Wu S, Wu X, Huso DL, Karim B, Franco AA, Rabizadeh S, Golub JE, Mathews LE, Shin J *et al* (2009) Induction of persistent colitis by a human commensal, enterotoxigenic *Bacteroides fragilis*, in wild-type C57BL/6 mice. *Infect Immun* 77: 1708–1718
64. Toprak NU, Yagci A, Gulluoglu BM, Akin ML, Demirkalem P, Celenk T, Soyletir G (2006) A possible role of *Bacteroides fragilis* enterotoxin in the aetiology of colorectal cancer. *Clin Microbiol Infect* 12: 782–786
65. Nakayama M, Hisatsune J, Yamasaki E, Isomoto H, Kurazono H, Hatakeyama M, Azuma T, Yamaoka Y, Yahiro K, Moss J *et al* (2009) *Helicobacter pylori* VacA-induced inhibition of GSK3 through the PI3K/Akt signaling pathway. *J Biol Chem* 284: 1612–1619
66. Sokolova O, Bozko PM, Naumann M (2008) *Helicobacter pylori* suppresses glycogen synthase kinase 3beta to promote beta-catenin activity. *J Biol Chem* 283: 29367–29374
67. McCracken KW, Cata EM, Crawford CM, Sinagoga KL, Schumacher M, Rockich BE, Tsai YH, Mayhew CN, Spence JR, Zavros Y *et al* (2014) Modelling human development and disease in pluripotent stem-cell-derived gastric organoids. *Nature* 516: 400–404
68. Xu X, Liu Z, Fang M, Yu H, Liang X, Li X, Liu X, Chen C, Jia J (2012) *Helicobacter pylori* CagA induces ornithine decarboxylase upregulation via Src/MEK/ERK/c-Myc pathway: implication for progression of gastric diseases. *Exp Biol Med (Maywood)* 237: 435–441
69. Wang F, Meng W, Wang B, Qiao L (2014) *Helicobacter pylori*-induced gastric inflammation and gastric cancer. *Cancer Lett* 345: 196–202
70. Fredlund J, Enninga J (2014) Cytoplasmic access by intracellular bacterial pathogens. *Trends Microbiol* 22: 128–137
71. Senerovic L, Tsunoda SP, Goosmann C, Brinkmann V, Zychlinsky A, Meissner F, Kolbe M (2012) Spontaneous formation of IpaB ion channels in host cell membranes reveals how Shigella induces pyroptosis in macrophages. *Cell Death Dis* 3: e384
72. Henry R, Shaughnessy L, Loessner MJ, Alberti-Segui C, Higgins DE, Swanson JA (2006) Cytolysin-dependent delay of vacuole maturation in macrophages infected with *Listeria monocytogenes*. *Cell Microbiol* 8: 107–119
73. Shaughnessy LM, Lipp P, Lee K-D, Swanson JA (2007) Localization of protein kinase C  $\epsilon$  to macrophage vacuoles perforated by *Listeria monocytogenes* cytolysin. *Cell Microbiol* 9: 1695–1704
74. Santic M, Asare R, Skrobbonja I, Jones S, Abu Kwaik Y (2008) Acquisition of the vacuolar ATPase proton pump and phagosome acidification are essential for escape of *Francisella tularensis* into the macrophage cytosol. *Infect Immun* 76: 2671–2677
75. Ozanic M, Marecic V, Abu Kwaik Y, Santic M (2015) The divergent intracellular lifestyle of *Francisella tularensis* in evolutionarily distinct host cells. *PLoS Pathog* 11: e1005208
76. Chong A, Wehrly TD, Nair V, Fischer ER, Barker JR, Klose KE, Celli J (2008) The early phagosomal stage of *Francisella tularensis* determines optimal phagosomal escape and *Francisella* pathogenicity island protein expression. *Infect Immun* 76: 5488–5499
77. Nagai H, Kagan JC, Zhu X, Kahn RA, Roy CR (2002) A bacterial guanine nucleotide exchange factor activates ARF on *Legionella* phagosomes. *Science* 295: 679–682
78. Tilney LG, Harb OS, Connelly PS, Robinson CG, Roy CR (2001) How the parasitic bacterium *Legionella pneumophila* modifies its phagosome and transforms it into rough ER: implications for conversion of plasma membrane to the ER membrane. *J Cell Sci* 114: 4637–4650
79. Rajashekar R, Liebl D, Chikkaballi D, Liss V, Hensel M (2014) Live cell imaging reveals novel functions of *Salmonella enterica* SPI2-T3SS effector proteins in remodeling of the host cell endosomal system. *PLoS One* 9: e115423

80. Beuzon CR, Meresse S, Unsworth KE, Ruiz-Albert J, Garvis S, Waterman SR, Ryder TA, Boucrot E, Holden DW (2000) *Salmonella* maintains the integrity of its intracellular vacuole through the action of SifA. *EMBO J* 19: 3235–3249
81. Sindhvani A, Arya SB, Kaur H, Jagga D, Tuli A, Sharma M (2017) *Salmonella* exploits the host endolysosomal tethering factor HOPS complex to promote its intravacuolar replication. *PLoS Pathog* 13: e1006700
82. Stein MA, Leung KY, Zwick M, Garcia-del Portillo F, Finlay BB (1996) Identification of a *Salmonella* virulence gene required for formation of filamentous structures containing lysosomal membrane glycoproteins within epithelial cells. *Mol Microbiol* 20: 151–164
83. Lu R, Wu S, Zhang YG, Xia Y, Zhou Z, Kato I, Dong H, Bissonnette M, Sun J (2016) *Salmonella* protein AvrA activates the STAT3 signaling pathway in colon cancer. *Neoplasia* 18: 307–316
84. Lu R, Bosland M, Xia Y, Zhang YG, Kato I, Sun J (2017) Presence of *Salmonella* AvrA in colorectal tumor and its precursor lesions in mouse intestine and human specimens. *Oncotarget* 8: 55104–55115
85. Lu R, Wu S, Zhang YG, Xia Y, Liu X, Zheng Y, Chen H, Schaefer KL, Zhou Z, Bissonnette M *et al* (2014) Enteric bacterial protein AvrA promotes colonic tumorigenesis and activates colonic beta-catenin signaling pathway. *Oncogenesis* 3: e105
86. Ye Z, Petrof EO, Boone D, Claud EC, Sun J (2007) *Salmonella* effector AvrA regulation of colonic epithelial cell inflammation by deubiquitination. *Am J Pathol* 171: 882–892
87. Sun J, Hobert ME, Rao AS, Neish AS, Madara JL (2004) Bacterial activation of beta-catenin signaling in human epithelia. *Am J Physiol Gastrointest Liver Physiol* 287: G220–G227
88. Jones RM, Wu H, Wentworth C, Luo L, Collier-Hyams L, Neish AS (2008) *Salmonella* AvrA coordinates suppression of host immune and apoptotic defenses via JNK pathway blockade. *Cell Host Microbe* 3: 233–244
89. Orth K, Palmer LE, Bao ZQ, Stewart S, Rudolph AE, Bliska JB, Dixon JE (1999) Inhibition of the mitogen-activated protein kinase superfamily by a Yersinia effector. *Science* 285: 1920–1923
90. Trosky JE, Mukherjee S, Burdette DL, Roberts M, McCarter L, Siegel RM, Orth K (2004) Inhibition of MAPK signaling pathways by VopA from *Vibrio parahaemolyticus*. *J Biol Chem* 279: 51953–51957
91. Trosky JE, Li Y, Mukherjee S, Keitany G, Ball H, Orth K (2007) VopA inhibits ATP binding by acetylating the catalytic loop of MAPK kinases. *J Biol Chem* 282: 34299–34305
92. Fehr D, Casanova C, Liverman A, Blazkova H, Orth K, Dobbelaere D, Frey J, Burr SE (2006) AopP, a type III effector protein of *Aeromonas salmonicida*, inhibits the NF-kappaB signalling pathway. *Microbiology* 152: 2809–2818
93. Hannemann S, Gao B, Galán JE (2013) *Salmonella* modulation of host cell gene expression promotes its intracellular growth. *PLoS Pathog* 9: e1003668
94. Kuijl C, Savage ND, Marsman M, Tuin AW, Janssen L, Egan DA, Ketema M, van den Nieuwendijk R, van den Eeden SJ, Geluk A *et al* (2007) Intracellular bacterial growth is controlled by a kinase network around PKB/AKT1. *Nature* 450: 725–730
95. Kuo SC, Hu YW, Liu CJ, Lee YT, Chen YT, Chen TL, Chen TJ, Fung CP (2013) Association between tuberculosis infections and non-pulmonary malignancies: a nationwide population-based study. *Br J Cancer* 109: 229
96. Melenotte C, Million M, Audoly G, Gorse A, Dutronc H, Roland G, Dekel M, Moreno A, Cammilleri S, Carrieri MP *et al* (2016) B-cell non-Hodgkin lymphoma linked to *Coxiella burnetii*. *Blood* 127: 113–121
97. Fulbright LE, Ellermann M, Arthur JC (2017) The microbiome and the hallmarks of cancer. *PLoS Pathog* 13: e1006480
98. Louis P, Hold GL, Flint HJ (2014) The gut microbiota, bacterial metabolites and colorectal cancer. *Nat Rev Microbiol* 12: 661
99. Wundisch T, Thiede C, Morgner A, Dempfle A, Gunther A, Liu H, Ye H, Du MQ, Kim TD, Bayerdorffer E *et al* (2005) Long-term follow-up of gastric MALT lymphoma after *Helicobacter pylori* eradication. *J Clin Oncol* 23: 8018–8024
100. Dmitrieva O, Grivennikov SI (2017) Microbiota and cancer: a complex equation with a lot of exciting unknowns. *Semin Immunol* 32: 1–2
101. Erdman SE, Poutahidis T (2015) Gut bacteria and cancer. *Biochem Biophys Acta* 1856: 86–90
102. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans (2012) Biological agents. Volume 100 B. A review of human carcinogens. *IARC Monogr Eval Carcinog Risks Hum* 100: 1–441
103. Liu W, MacDonald M, You J (2016) Merkel cell polyomavirus infection and Merkel cell carcinoma. *Curr Opin Virol* 20: 20–27
104. Zhu H, Shen Z, Luo H, Zhang W, Zhu X (2016) *Chlamydia trachomatis* infection-associated risk of cervical cancer: a meta-analysis. *Medicine (Baltimore)* 95: e3077
105. Trabert B, Waterboer T, Idahl A, Brenner N, Brinton LA, Butt J, Coburn SB, Hartge P, Hufnagel K, Inturrisi F *et al* (2018) Antibodies against *Chlamydia trachomatis* and ovarian cancer risk in two independent populations. *J Natl Cancer Inst* 111: djy084



**License:** This is an open access article under the terms of the Creative Commons Attribution 4.0 License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.