

NEWS AND VIEWS

Open Access

More than two decades trapped

Cheng-Wei Qiu¹ and Lei-Ming Zhou¹ 

Abstract

Optical tweezers, crowned by Nobel Prize the first time in 1990s, have widely impacted the research landscape of atom cooling, particle manipulation/sorting, and biology. After more than two decades of steady development, it received the deserving recognition once again in 2018. Unprecedented advancements across various disciplines are believed to be spurred furthermore by this important tool of optical manipulation.

Since the discovery of optical trapping force in half a century ago, it has been used and known widely as the optical tweezers^{1,2}. In 1997, Steven Chu, Claude Cohen-Tannoudji, and William D. Phillips won the Nobel Prize for atom trapping and cooling by laser^{3,4}. Through trapping and manipulating larger items such as bacteria and cells^{5,6}, it becomes as a powerful paradigm in biological and medical science, which won Arthur Ashkin the Nobel Prize in 2018. In between those two Nobel Prizes in Physics, more than two decades have elapsed, witnessing significant progress in advanced optical micro-/nano-manipulations based on the optical tweezer concept. This “trapped” state, though not short, provides the whole community a steady yet profound opportunity to sit back, searching for groundbreaking application values. And, the optical tweezer concept eventually returns with an epic laureate, owing to its powerful and promising applications in biology.

The first observation of the mechanical effect of light force phenomenon can be traced back to 1619, i.e., more than 400 years ago, when Kepler^{7,8} observed the comet tails pointed away from the sun and captivated that it was caused by the force of light. Then, 250 years later, Maxwell⁹ crafted his theory of classical electrodynamics, in which he showed that the light carried momentum and exerted a pressure on an object if the object reflected the light. The force along with the pressure could push the object forward, in an analog to what happened in a comet

tail. His theory was confirmed experimentally by Nicolas and Hull¹⁰.

The force caused by light is called radiative pressure since then. It was taken for granted that the radiative force can push the particle forward due to the momentum conservation law. Ashkin¹ counter-intuitively demonstrated the gradient of the light field distribution could drag and trap the particle in the liquid with two counter-propagating laser beams. In this work, he also stated the idea of levitating atoms and molecules using resonant light with the atom transition. Ashkin and Chu² further demonstrated the trapping of a dielectric particle with a single strongly focused beam and extended the trapping size range to 10 μm –25 nm, which paves the most fundamental platform of optical tweezers².

Chu et al.⁴ demonstrated the trapping of atoms using laser beam and cooling of the atoms to extremely low temperature. The cooled atoms empower a plethora of applications, especially in high sensitivity metrology including atomic interferometry and atomic clock. In parallel, Ashkin continued to flourish the realm of optical tweezers. He managed to demonstrate the manipulation of single viruses and bacteria⁵, and single cells alive⁶. The 1064 nm-wavelength infrared light has been employed, providing sufficiently large force with greatly reduced damage to the biology cell. It was followed by tremendous investigations and developments in biology science based on optical tweezers. Block et al.¹¹ studied the bead movement by single kinesin molecules with optical tweezers. Yin et al.¹² measured the force produced by a single molecule of RNA polymerase during transcription. Using an optical trapping interferometer with feedback


Correspondence: C-W. Qiu (chengwei.qiu@nus.edu.sg)

¹Department of Electrical and Computer Engineering, National University of Singapore, 4 Engineering Drive 3, Singapore 117583, Singapore

© The Author(s) 2018



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.



control, Wang et al.¹³ measured the force-extension relationships of single DNA molecules. Optical tweezers can trap micrometer-/nanometer-size items with an exerted force from 100 aN to 100 pN, right in the range of the forces within cell and macromolecular systems. Thus, optical tweezers fit perfectly for investigating and even engineering various biological process, e.g., characterization of the forces of kinesin molecules¹¹, probing the viscoelastic properties¹³, and doing intracellular surgery¹⁴. Over the past decades, the territorial boundary of optical tweezers has also been significantly extended to various other areas, including colloid and interface science¹⁵, microfluidic sorting by light¹⁶, and even quantum science and technology based on levitated opto-mechanical system¹⁷. The sophistication and powerfulness of the tweezer have also been greatly boosted, e.g., nanometric optical tweezers¹⁸ and holographic optical tweezers¹⁹. A schematic illustration of its historical development is shown in Fig. 1.

Although the optical tweezers have received the grand recognition of Nobel Prize twice in the past three decades, the novel physics behind the optical force still fascinate the researchers, especially when optical force meets with structured lights or materials. Novel mechanisms of optical force have been revealed. Using vortex beams, researchers reported the complex stiffness and trapping mechanism of beams with orbital angular momentum²⁰. With a birefringent microparticle in vacuum, Arita et al.²¹ demonstrated 5 MHz frequency of rotation using the circularly polarized beam and proposed its application in micro-gyroscope. Using beams with angular momentum, researchers managed to trap and spin the particles^{22,23}. Another unprecedentedly interesting topic is the optical

pulling force—the light could pull the particle toward the light source instead of pushing away. The pulling phenomena is first studied in optical solenoid beams²⁴ and afterwards the full framework for the theory of optical “tractor beam” is established^{25–27}.

Apart from those exciting achievements in aforementioned areas, the optical tweezer technology rapidly expands its “contour” and synergizes with other disciplines^{28–30}. For instance, it spurs the advancement in the ground-state cooling of macro-particles¹⁷, detection of non-Newtonian gravity³¹, detection of gravitational wave³², and Brownian Carnot engine³³. Therefore, it is believed that the optical tweezers will continue to sail and explore the new edges, and return with more in near future.

Acknowledgements

C.-W.Q. acknowledges the support from Ministry of Education, Singapore (Grant No. R-263-000-D11-114).

Conflict of interest

The authors declare that they have no conflict of interest.

Received: 13 October 2018 Accepted: 21 October 2018

Published online: 07 November 2018

References

- Ashkin, A. Acceleration and trapping of particles by radiation pressure. *Phys. Rev. Lett.* **24**, 156–159 (1970).
- Ashkin, A., Dziedzic, J. M., Bjorkholm, J. E. & Chu, S. Observation of a single-beam gradient force optical trap for dielectric particles. *Opt. Lett.* **11**, 288–290 (1986).
- Chu, S., Hollberg, L., Bjorkholm, J. E., Cable, A. & Ashkin, A. Three-dimensional viscous confinement and cooling of atoms by resonance radiation pressure. *Phys. Rev. Lett.* **55**, 48–51 (1985).

4. Chu, S., Bjorkholm, J. E., Ashkin, A. & Cable, A. Experimental observation of optically trapped atoms. *Phys. Rev. Lett.* **57**, 314–317 (1986).
5. Ashkin, A. & Dziedzic, J. M. Optical trapping and manipulation of viruses and bacteria. *Science* **235**, 1517–1520 (1987).
6. Ashkin, A., Dziedzic, J. M. & Yamane, T. Optical trapping and manipulation of single cells using infrared laser beams. *Nature* **330**, 769–771 (1987).
7. Kepler, J. *Ad Vitellionem Paralipomena* (Francofurti, Apud Claudium Marnium & Haeredes Ioannis Aubrii, Frankfurt, 1604).
8. Kepler, J. *De Cometis Libelli Tres* (Aavgvstae Vindelicorvm, A. Apergeri, Augsburg, 1619).
9. Maxwell, J. C. A dynamical theory of the electromagnetic field. *Philos. Trans. R. Soc. Lond.* **155**, 459–512 (1865).
10. Baxter, C. & Loudon, R. Radiation pressure and the photon momentum in dielectrics. *J. Mod. Opt.* **57**, 830–842 (2010).
11. Block, S. M., Goldstein, L. S. B. & Schnapp, B. J. Bead movement by single kinesin molecules studied with optical tweezers. *Nature* **348**, 348–352 (1990).
12. Yin, H. et al. Transcription against an applied force. *Science* **270**, 1653–1657 (1995).
13. Wang, M. D., Yin, H., Landick, R., Gelles, J. & Block, S. M. Stretching DNA with optical tweezers. *Biophys. J.* **72**, 1335–1346 (1997).
14. Grier, D. G. A revolution in optical manipulation. *Nature* **424**, 810–816 (2003).
15. Grier, D. G. Optical tweezers in colloid and interface science. *Curr. Opin. Colloid Interface Sci.* **2**, 264–270 (1997).
16. MacDonald, M. P., Spalding, G. C. & Dholakia, K. Microfluidic sorting in an optical lattice. *Nature* **426**, 421–424 (2003).
17. Yin, Z. Q., Geraci, A. A. & Li, T. C. Optomechanics of levitated dielectric particles. *Int. J. Mod. Phys. B* **27**, 1330018 (2013).
18. Novotny, L., Bian, R. X. & Xie, X. S. Theory of nanometric optical tweezers. *Phys. Rev. Lett.* **79**, 645–648 (1997).
19. Curtis, J. E., Koss, B. A. & Grier, D. G. Dynamic holographic optical tweezers. *Opt. Commun.* **207**, 169–175 (2002).
20. Ng, J., Lin, Z. F. & Chan, C. T. Theory of optical trapping by an optical vortex beam. *Phys. Rev. Lett.* **104**, 103601 (2010).
21. Arita, Y., Mazilu, M. & Dholakia, K. Laser-induced rotation and cooling of a trapped microgyroscope in vacuum. *Nat. Commun.* **4**, 2374 (2013).
22. Simpson, N. B., Dholakia, K., Allen, L. & Padgett, M. J. Mechanical equivalence of spin and orbital angular momentum of light: an optical spanner. *Opt. Lett.* **22**, 52–54 (1997).
23. Padgett, M. & Bowman, R. Tweezers with a twist. *Nat. Photon.* **5**, 343–348 (2011).
24. Lee, S. H., Roichman, Y. & Grier, D. G. Optical solenoid beams. *Opt. Express* **18**, 6988–6993 (2010).
25. Sukhov, S. & Dogariu, A. On the concept of “tractor beams”. *Opt. Lett.* **35**, 3847–3849 (2010).
26. Chen, J., Ng, J., Lin, Z. F. & Chan, C. T. Optical pulling force. *Nat. Photon.* **5**, 531–534 (2011).
27. Novitsky, A., Qiu, C. W. & Wang, H. Single gradientless light beam drags particles as tractor beams. *Phys. Rev. Lett.* **107**, 203601 (2011).
28. Gao, D. L. et al. Optical manipulation from the microscale to the nanoscale: fundamentals, advances and prospects. *Light Sci. Appl.* **6**, e17039 (2017).
29. Ivinskaya, A. et al. Plasmon-assisted optical trapping and anti-trapping. *Light Sci. Appl.* **6**, e16258 (2016).
30. Bykov, D. S. et al. Long-range optical trapping and binding of microparticles in hollow-core photonic crystal fibre. *Light Sci. Appl.* **7**, 22 (2018).
31. Geraci, A. A., Papp, S. B. & Kitching, J. Short-range force detection using optically cooled levitated microspheres. *Phys. Rev. Lett.* **105**, 101101 (2010).
32. Arvanitaki, A. & Geraci, A. A. Detecting high-frequency gravitational waves with optically levitated sensors. *Phys. Rev. Lett.* **110**, 071105 (2013).
33. Martínez, I. A. et al. Brownian Carnot engine. *Nat. Phys.* **12**, 67–70 (2016).