

Published online: 08 November 2018

OPEN Author Correction: Obstetric outcomes in pregnant women with and without depression: population-based comparison

Hui-Chun Huang 1,2,3, Fung-Chang Sung4,5, Pei-Chun Chen2, Cherry Yin-Yi Chang6, Chih-Hsin Muo⁴, Huei-Sheng Shiue^{2,7,8}, Jian-Pei Huang^{3,9,10}, Tsai-Chung Li¹₁₀, Ya-Ling Tzeng¹¹ & Shu-I Wu^{9,12}

Correction to: Scientific Reports https://doi.org/10.1038/s41598-017-14266-3, published online 24 October 2017

This Article contains errors in the Methods section under subheading 'Data source and study population'.

"From the 2005-2013 claims data, we identified pregnant women who had the diagnosis of depression during pregnancy (ICD-9-CM codes of depression: 296.2, 296.20 to 296.25, 296.2, 296.30 to 296.35, 296.82, 300.4, 309.0, 309.1, 309.28, and 311). Only pregnant women with the depression diagnosis for at least 3 times in the outpatient records or for at least once depression diagnosis in the inpatient records were included in the depression cohort. Depressed women with the history of schizophrenia (ICD-9-CM 295), bipolar disorders (ICD-9-CM 296, 296.2, and 296.3) and/or anxiety (ICD-9-CM 300) were excluded."

should read:

"From the 2005-2013 claims data, we identified pregnant women who had the diagnosis of depression during pregnancy (ICD-9-CM codes of depression: 296.20 to 296.25, 296.30 to 296.35, 296.82, 300.4, 309.0, 309.1, 309.28, and 311). Only pregnant women with the depression diagnosis for at least 3 times in the outpatient records or for at least once depression diagnosis in the inpatient records were included in the depression cohort. Depressed women with the history of schizophrenia (ICD-9-CM 295), bipolar disorders (ICD-9-CM 296, except 296.2, and 296.3) and/or anxiety (ICD-9-CM 300, except 300.4) were excluded."

¹Department of Medical Research, Mackay Memorial Hospital, New Taipei, 251, Taiwan. ²Department of Public Health, China Medical University, Taichung, 404, Taiwan. ³Center for General Education, MacKay Junior College of Medicine, Nursing and Management, Taipei, 112, Taiwan. 4Management Office for Health Data, China Medical University Hospital, Taichung, 404, Taiwan. 5Department of Health Services Administration, China Medical University, Taichung, 404, Taiwan. ⁶Department of Obstetrics and Gynecology, China Medical University Hospital, Taichung, 404, Taiwan. ⁷Division of Physical Medicine and Rehabilitation, E-Da Hospital, Kaohsiung, 824, Taiwan. ⁸School of Medicine for International Students, I-Shou University, Kaohsiung, 824, Taiwan. ⁹Department of Medicine, Mackay Medical College, Taipei, 252, Taiwan. 10 Department of Obstetrics and Gynecology, Mackay Memorial Hospital, Taipei, 104, Taiwan. ¹¹School of Nursing, China Medical University, Taichung, 404, Taiwan. ¹²Department of Psychiatry, Mackay Memorial Hospital, Taipei, 104, Taiwan. Fung-Chang Sung and Shu-I Wu contributed equally. Correspondence and requests for materials should be addressed to F.-C.S. (email: fcsung1008@yahoo.com) or S.-I.W. (email: shuiwu624@gmail.com)

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit https://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2018