

Erratum to: Efficacy and safety of sitagliptin in elderly patients with type 2 diabetes mellitus and comparison of hypoglycemic action of concomitant medications: a subanalysis of the JAMP study

Noriko Ujihara¹ · Hiroshi Sakura² · Naotake Hashimoto³ · Kazuo Sasamoto⁴ · Hiroshi Ohashi⁵ · Sumiko Hasumi⁶ · Tadasu Kasahara⁷ · Osamu Tomonaga⁸ · Hideo Nunome⁹ · Masashi Honda¹⁰ · Yasuhiko Iwamoto¹¹ · for the JAMP Study Investigators

Published online: 14 October 2017
© The Japan Diabetes Society 2017

Erratum to: Diabetol Int
DOI 10.1007/s13340-017-0330-2

In the original publication of the article, a percentage symbol has been included incorrectly in the value of Alcohol consumption (65–74 years) as 45.1% in Table 1. The correct version is published with this erratum.

The online version of the original article can be found under doi:[10.1007/s13340-017-0330-2](https://doi.org/10.1007/s13340-017-0330-2).

✉ Noriko Ujihara
ujihara.noriko@twmu.ac.jp

- ¹ Department of Medicine, Diabetes Center, Institute of Geriatrics, Tokyo Women's Medical University, Shibuya Cross Tower 21F, 2-15-1, Shibuya, Shibuya-ku, Tokyo 150-0002, Japan
- ² Department of Medicine, Tokyo Women's Medical University, Medical Center East, 2-1-10 Nishiogu, Arakawa-ku, Tokyo 116-8567, Japan
- ³ Department of Diabetes, Endocrine and Metabolic Diseases, Tokyo Women's Medical University Yachiyo Medical Center, 477-96, Owada-shinden, Yachiyo-shi, Chiba 276-0046, Japan
- ⁴ Internal Medicine, Suzuki Clinic, 1F, 2-10-14, Koyasu-machi, Hachioji-shi, Tokyo 192-0904, Japan
- ⁵ Internal Medicine, Oyama East Clinic, 1-32-1, Ekihigashi-dori, Oyama-shi, Tochigi 323-0022, Japan
- ⁶ Internal Medicine, Nishiyamado-Keiwa Hospital, 3247-1, Kounosu, Naka-shi, Ibaraki 311-0133, Japan

- ⁷ Josai Hospital, 2-42-11, Kamiogi, Suginami-ku, Tokyo 167-0043, Japan
- ⁸ Diabetes and Lifestyle Center, Tomonaga Clinic, Shinyoncurumu Building 9F, 4-2-23, Shinjuku, Shinjuku-ku, Tokyo 160-0022, Japan
- ⁹ Diabetes Center, Edogawa Hospital, Medical Plaza Shinozaki, SK Building, 7-15-12, Shinozaki-machi, Edogawa-ku, Tokyo 133-0057, Japan
- ¹⁰ Nishikawa Clinic, 2-16-3, Towa, Adachi-ku, Tokyo 120-0003, Japan
- ¹¹ Tokyo Women's Medical University/The Institute for Adult Diseases, Asahi Life Foundation, Asahiseimeisunaga Building 2, 3, 4F, 2-2-6, Nihonbashi Bakuro-cho, Chuo-ku, Tokyo 103-0002, Japan

Table 1 Comparison of patient characteristics between the elderly and non-elderly patients

	≥ 75 years (<i>n</i> = 130)		65–74 years (<i>n</i> = 189)		< 65 years (<i>n</i> = 332)		<i>p</i>
	<i>n</i>	Mean ± SD or %	<i>n</i>	Mean ± SD or %	<i>n</i>	Mean ± SD or %	
Age (years)	130	80.0 ± 4.7	189	69.2 ± 2.9	332	54.5 ± 7.4	0.000*
Height (cm)	127	156.2 ± 9.6	187	161.0 ± 8.6	328	166.2 ± 8.8	0.000*
Waist circumference (cm)	80	85.7 ± 10.5	120	87.3 ± 9.6	193	89.9 ± 12.0	0.009*
Duration of diabetes (days)	111	121.4 ± 94.6	181	114.1 ± 78.4	308	95.4 ± 74.4	0.003*
Sex (male)	67	51.5	117	61.9	250	75.3	0.000*
Smoking	13	10.3	34	18.6	96	29.8	0.000*
Alcohol consumption	39	31.7	82	45.1	180	56.1	0.000*
Retinopathy	8	6.2	8	4.2	32	9.6	0.064
Arteriosclerosis obliterans	6	4.6	21	11.1	28	8.4	0.122
Stroke	0	0.0	1	0.6	0	0.0	0.288
Myocardial infarction	7	5.4	5	2.6	6	1.8	0.108
Angina pectoris	8	6.2	8	4.2	11	3.3	0.387
Cardiac failure	5	3.8	2	1.1	4	1.2	0.102
Atrial fibrillation	4	3.1	5	2.6	7	2.1	0.817
Diet/exercise therapy	49	37.7	48	25.4	92	27.7	0.044*
Low dose of glimepiride	18	13.8	25	13.2	29	8.7	0.153
Medium dose of glimepiride	9	6.9	21	11.1	20	6.0	0.104
BG	9	6.9	25	13.2	65	19.6	0.002*
TZD	7	5.4	12	6.3	19	5.7	0.929
α-GI	5	3.8	3	1.6	10	3.0	0.446
Multidrug therapy	33	25.4	55	29.1	97	29.2	0.692
Antihypertensive drug	81	62.3	106	56.1	144	43.4	0.000*
Antihyperlipidemic drug	62	47.7	94	49.7	156	47.0	0.832
Antihyperuricemic drug	6	4.6	9	4.8	16	4.8	0.996
Antithrombogenic drug	22	16.9	44	23.3	52	15.7	0.088

* *p* < 0.05