



Published in final edited form as:

J Acquir Immune Defic Syndr. 2018 December 15; 79(5): e112–e114. doi:10.1097/QAI.0000000000001863.

Genital HIV-1 Shedding with Dolutegravir (DTG) plus Lamivudine (3TC) Dual Therapy

Sara Gianella¹, Vincent C. Marconi², Baiba Berzins³, Constance A. Benson¹, Paul Sax⁴, Carl J. Fichtenbaum⁵, Timothy Wilkin⁶, Millie Vargas¹, Qianqian Deng¹, Michelli F. Oliveira¹, Carlee Moser⁷, and Babafemi O. Taiwo³

¹Division of Infectious Diseases & Global Public Health, University of California San Diego, La Jolla, California.

²Emory University, School of Medicine and Rollins School of Public Health, Atlanta, Georgia.

³Division of Infectious Diseases, Northwestern University, Chicago.

⁴Division of Infectious Diseases Brigham and Women's Hospital Boston, Massachusetts.

⁵University of Cincinnati Medical Center, Cincinnati

⁶Division of Infectious Diseases, Weill Cornell Medicine, New York.

⁷Department of Biostatistics, Center for Biostatistics and AIDS Research, Harvard TH Chan School of Public Health, Boston, Massachusetts.

Introduction

Detection of HIV RNA in the genital tract is correlated with sexual transmission [1–4], and is best predicted by the degree of plasma viremia [5]. Although there is a near-linear relationship between blood and genital HIV RNA, episodic genital HIV RNA expression occurs in some individuals with suppressed viremia possibly due to genital viral compartmentalization with poor drug penetration [6, 7], or stimulation of virus replication by sexually transmitted infections and genital inflammation [8–10]. Further, discordant viral resistance patterns in blood and genital samples occur in some individuals [11], supporting the possibility of differential virus evolution in these anatomic compartments. While genital HIV RNA shedding has been reported in 2–20% of individuals on standard three-drug ART [8, 9, 12, 13], there is no evidence that such shedding leads to new infections in the context of suppressed viremia, hence the recent consensus statement, “undetectable equals untransmittable” or “U=U” [14].

Corresponding Author: Sara Gianella, University of California San Diego, 9500 Gilman Drive MC 0679, La Jolla, CA 92093-0679, USA, Tel: (858) 552-8585, ext 7193; Fax: (858) 552-7445; gianella@ucsd.edu.

Disclosures

BOT has served as a paid consultant to ViiV, Gilead, and Janssen and on the Clinical Care Options speakers bureau, and has received grant funding to his institution from ViiV/GlaxoSmithKline (GSK). VCM, CJF, CAB, TW, have received grant funding for this study to their institutions through Northwestern University from ViiV/GSK.

BB and SG have no conflicts to declare

Several two-drug regimens are being studied because of potential benefits of reduced antiretroviral exposure and costs. Enthusiasm for two-drug regimens may be limited if they lead to increased genital HIV shedding. Thus, we investigated genital HIV RNA shedding with dolutegravir (DTG) plus lamivudine (3TC), which is under investigation for initial and maintenance therapy.

Methods

Study participant characteristics and clinical sampling.

We recruited participants from two clinical trials: 1) virologically suppressed participants (for >48 weeks at the time of screening) randomized to continuation of three-drug antiretroviral therapy (ART) or switch to DTG+3TC maintenance in ASPIRE (“Antiretroviral Study to Promote Improvement and Reduce Exposure”) [15], and 2) ART-naïve participants, who initiated DTG+3TC in the single-arm AIDS Clinical Trials Group (ACTG) A5353 phase 2 pilot study [16].

At study weeks 24 (or 36) and 48, female genital secretions were collected using self-administered vaginal swabs and immediately frozen at -80°C , while male genital secretions were collected by masturbation. Seminal plasma was processed according to standard procedures and stored at -80°C [5]. The parent studies (i.e. ASPIRE and A5353) and this genital sub-study were approved by the Institutional Review Board at each study site, and each participant provided a written informed consent.

RNA Extraction from genital secretion and HIV quantification

HIV RNA levels were measured in seminal plasma, as previously described [17,18]. For female secretions, RNA was recovered from dry vaginal swabs by suspending with 1ml of RPMI 1640 Medium vortexed and incubated for 10 minutes on ice. Supernatant was centrifuged (23,500xg at 4°C for 1 hour). Concentrated RNA was extracted using High Pure Viral RNA Kit (Roche) and cDNA was generated using the SuperScript III First-Strand Synthesis Kit (Invitrogen) [19, 20]. HIV RNA was quantified by real-time PCR in an ABI 7900HT thermocycler (Applied Biosystems) [17, 21].

DNA Extraction from genital secretion, and herpesvirus DNA quantification

Viral DNA was extracted from 200 μl of seminal plasma and 400 μl dry swab resuspension with 1X PBS using QIAamp DNA Mini Kit (Qiagen) per manufacturer’s protocol. 400 μl DNA from dry swab was further concentrated by standard ethanol-based DNA precipitation. EDTA (50mM) was quickly added to raw semen samples to inhibit DNase activity [18]. Levels of different herpesviruses in semen were measured by real-time PCR in an ABI 7900HT thermocycler (Applied Biosystems) [17, 21].

Genotyping

Extracted RNA was reverse transcribed using the SuperScript III Reverse Transcriptase (Thermo Fisher Scientific). Nested PCR was performed to obtain integrase or Protease/Reverse Transcriptase using Taq HiFi polymerase (Invitrogen) and Sanger sequenced. Sequences were manually edited using BioEdit Sequence Editor (v7.0.5) and evaluated for

the presence of drug resistance mutations using the HIVseq Program from the Stanford University Drug Resistance Database.

Statistical Analysis

Genital HIV RNA above the lower limit of detection (40 copies/mL) was considered detectable. The proportion of participants with detectable HIV RNA was determined in each study arm and 95% confidence intervals were calculated using binomial exact test. No formal statistical comparisons were performed because of the limited sample size.

Results

Study participant characteristics

We enrolled 38 participants from ASPIRE (18 switched to DTG+3TC while 20 had remained on their three-drug regimen), and 13 ART-naïve individuals who initiated DTG+3TC in the ACTG A5353 study. The median ART duration before ASPIRE entry was 5.4 years (IQR: 3.5–7.6) in the DTG+3TC group and 6.2 years (3.8–7.8) in the three-drug group. The pre-randomization regimens included a protease inhibitor (32%), non-nucleoside reverse transcriptase (26%) or integrase inhibitor (42%). Participants' characteristics are summarized in Supplementary Table 1.

HIV RNA shedding

A total of 76 seminal fluid samples were collected from 45 men (20 at week 24, 15 at week 36, 41 at week 48). Twelve vaginal swabs were collected from 6 women (4 at week 24, 2 at week 36, 6 at week 48). All but three participants had undetectable HIV RNA in blood plasma at each time-point. Three male participants, and no female participants, had HIV RNA genital shedding >40 copies/ml (Table 1). They included 1/20 participants (5%, 95%CI [22]) in the ASPIRE three-drug arm with genital HIV RNA of 42 copies/mL at week 24, 1/18 participants (5.6%, 95%CI [0.1%, 27%]) in the ASPIRE DTG+3TC arm with detectable genital HIV RNA at two consecutive timepoints (488 copies/mL at week 36 and 79 copies/mL at week 48), and 1/13 participants (7.7%, 95%CI [0.2%, 36%]) in ACTG A5353 with genital HIV RNA of 48 copies/mL at week 24, see Table 1 and Supplementary Figure 1. Two out of three men with genital HIV shedding also had coincident viremia in plasma.

Gonorrhea and chlamydia RNA were not detected from any of the three HIV genital shedders. One HIV genital shedder (from the ASPIRE DTG+3TC group) had high levels of genital CMV DNA: 314,607 copies/ml and 86,090 copies/ml at the two timepoints where genital HIV RNA was detected. Genotyping of genital HIV isolates was unsuccessful because of the low sample volume and viral load, except integrase sequencing in one ASPIRE DTG+3TC participant (with 488 copies/mL HIV RNA in seminal fluid) that revealed no resistance mutations.

Discussion

There is robust evidence from studies of three-drug regimens that U=U (“undetectable equals untransmittable”) [14]. Less is known about the incidence of genital HIV RNA

shedding and transmission prevention attributes of investigational two-drug regimens. DTG +3TC is of particular interest since it could become a recommended option if its promising preliminary efficacy and safety results are confirmed in ongoing phase 3 trials [23].

In our pilot study of 51 adults living with HIV, the frequency of genital HIV RNA shedding while virologically suppressed in blood was similar between those who were on standard three-drug ART and those who were on DTG+3TC as initial or maintenance therapy. The frequency of genital shedding in each of these groups fell within the lower end of the range reported for three drug regimens in other studies (2–20%) [8, 9, 12, 13]. None of the six women enrolled in our study had genital shedding. As, reported by previous investigators [5], we observed a concordance between viremia and genital shedding as two out of three participants with detectable seminal HIV RNA also had viremia. Taken together, these results suggest that DTG + 3TC is effective in controlling genital HIV shedding, which accounts for the majority of HIV transmission [24]

In a previous study, the triple regimen of DTG+3TC+Abacavir led to more rapid attainment of HIV RNA <40 copies/mL in semen than blood in treatment-naïve individuals [25], likely because of lower baseline HIV RNA levels in semen. It is unknown whether a similar relationship between blood and seminal virus decay exists with DTG+3TC dual therapy, and this could be formally assessed as part of a clinical trial. Nevertheless, good efficacy of DTG + 3TC in the genital compartment is pharmacologically plausible since seminal protein-unbound DTG concentrations exceed the in vitro 50% inhibitory concentration by a median of 214-fold despite lower concentrations in seminal fluid than blood, [25] while seminal fluid concentration of 3TC exceeds the plasma concentration [26].

Our results are limited by the small sample size, the small number of women enrolled as well as by the sparse timing of specimen collection. We also did not assess pre- and post-treatment HIV RNA in each participant hence we are unable to describe the magnitude of genital viral load reduction with DTG + 3TC.

In conclusion, in this small pilot study we did not detect concerning signals about the efficacy of the two-drug regimen of DTG+3TC in controlling genital HIV RNA shedding hence prevention of viral transmission, when HIV RNA is undetectable in blood plasma. These preliminary results suggest that DTG+3TC likely confers similar transmission prevention benefits as triple therapy.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to the study participants and the ASPIRE and A5353 clinical research staff at Northwestern University (Sherrie Wolfe), Emory University (Kishna Outlaw), University of Cincinnati (Eva Whitehead), Cornell (Todd Stroberg), University of California at San Diego (Edward Seefried) whose commitment and cooperation made this study possible.

This work was supported by an Investigator Sponsored Study grant “Virologic Activity of DTG +3TC in the Genital Compartment: Substudy of ASPIRE and A5353” from ViiV Healthcare to Northwestern University. ViiV

Healthcare/GlaxoSmithKline provided funding and study drugs. Additional support was provided from the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under Award Number UM1 AI068634, UM1 AI068636 and UM1 AI106701, the San Diego Center for AIDS Research Translational Virology Core (P30AI036214) as well as the Third Coast Center for AIDS Research (P30AI117943), and Emory Center for AIDS Research (P30AI050409). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or ViiV Healthcare.”

HIV RNA quantification standard was obtained from the DAIDS Virology Quality Assurance (VQA) Program.

References.

1. Attia S, Egger M, Muller M, Zwahlen M, Low N. Sexual transmission of HIV according to viral load and antiretroviral therapy: systematic review and meta-analysis. *AIDS* 2009; 23(11):1397–1404. [PubMed: 19381076]
2. Cohen MS, Chen YQ, McCauley M, Gamble T, Hosseinipour MC, Kumarasamy N, et al. Prevention of HIV-1 infection with early antiretroviral therapy. *N Engl J Med* 2011; 365(6):493–505. [PubMed: 21767103]
3. Donnell D, Baeten JM, Kiarie J, Thomas KK, Stevens W, Cohen CR, et al. Heterosexual HIV-1 transmission after initiation of antiretroviral therapy: a prospective cohort analysis. *Lancet* 2010; 375(9731):2092–2098. [PubMed: 20537376]
4. Reynolds SJ, Makumbi F, Nakigozi G, Kagaayi J, Gray RH, Wawer M, et al. HIV-1 transmission among HIV-1 discordant couples before and after the introduction of antiretroviral therapy. *AIDS* 2011; 25(4):473–477. [PubMed: 21160416]
5. Morris SR, Zhao M, Smith DR, Vargas MV, Little SJ, Gianella S. Longitudinal Viral Dynamics in Semen During Early HIV Infection. *Clin Infect Dis* 2016.
6. Taylor S, Davies S. Antiretroviral drug concentrations in the male and female genital tract: implications for the sexual transmission of HIV. *Curr Opin HIV AIDS* 2010; 5(4):335–343. [PubMed: 20543610]
7. Else LJ, Taylor S, Back DJ, Khoo SH. Pharmacokinetics of antiretroviral drugs in anatomical sanctuary sites: the male and female genital tract. *Antiviral Therapy* 2011; 16(8):1149–1167. [PubMed: 22155899]
8. Sheth PM, Yi TJ, Kovacs C, Kemal KS, Jones RB, Osborne B, et al. Mucosal correlates of isolated HIV semen shedding during effective antiretroviral therapy. *Mucosal immunology* 2012; 5(3):248–257. [PubMed: 22318494]
9. Politch JA, Mayer KH, Welles SL, O'Brien W X, Xu C, Bowman FP, et al. Highly active antiretroviral therapy does not completely suppress HIV in semen of sexually active HIV-infected men who have sex with men. *AIDS* 2012.
10. Cohen MS. Sexually transmitted diseases enhance HIV transmission: no longer a hypothesis. *Lancet* 1998; 351 Suppl 3:5–7.
11. Gianella S, Smith DM, Vargas MV, Little SJ, Richman DD, Daar ES, et al. Shedding of HIV and Human Herpesviruses in the Semen of Effectively Treated HIV-1-Infected Men Who Have Sex With Men. *Clin Infect Dis* 2013; 57(3):441–447. [PubMed: 23595831]
12. Halfon P, Giorgetti C, Khiri H, Penaranda G, Terriou P, Porcu-Buisson G, et al. Semen may harbor HIV despite effective HAART: another piece in the puzzle. *PLoS One* 2010; 5(5):e10569. [PubMed: 20485526]
13. Marcelin AG, Tubiana R, Lambert-Niclot S, Lefebvre G, Dominguez S, Bonmarchand M, et al. Detection of HIV-1 RNA in seminal plasma samples from treated patients with undetectable HIV-1 RNA in blood plasma. *AIDS* 2008; 22(13):1677–1679. [PubMed: 18670231]
14. The Lancet H U=U taking off in 2017. *Lancet HIV* 2017; 4(11):e475. [PubMed: 29096785]
15. Taiwo BO, Marconi VC, Berzins B, Moser CB, Nyaku AN, Fichtenbaum CJ, et al. Dolutegravir Plus Lamivudine Maintains Human Immunodeficiency Virus-1 Suppression Through Week 48 in a Pilot Randomized Trial. *Clin Infect Dis* 2018; 66(11):1794–1797. [PubMed: 29293895]
16. Taiwo BO, Zheng L, Stefanescu A, Nyaku A, Bezins B, Wallis CL, et al. ACTG A5353: A Pilot Study of Dolutegravir Plus Lamivudine for Initial Treatment of Human Immunodeficiency Virus-1

- (HIV-1)-infected Participants With HIV-1 RNA <500000 Copies/mL. *Clin Infect Dis* 2018; 66(11): 1689–1697. [PubMed: 29253097]
17. Gianella S, Strain MC, Rought SE, Vargas M, Little SJ, Richman DD, et al. Associations between the Virologic and Immunologic Dynamics in Blood and in the Male Genital Tract *J Virol* 2012; 86(3):1307–1315. [PubMed: 22114342]
 18. Vargas-Meneses MV, Massanella M, Ignacio CC, Gianella S. Quantification of HIV RNA and Human Herpesvirus DNA in Seminal Plasma. *Bio Protoc* 2015; 5(9).
 19. Althaus CF, Gianella S, Rieder P, von Wyl V, Kouyos RD, Niederost, et al. Rational design of HIV-1 fluorescent hydrolysis probes considering phylogenetic variation and probe performance. *J Virol Methods* 2010; 165(2):151–160. [PubMed: 20116399]
 20. A.R. B, Smith DM, Deutsch R, Ellis RJ, Cherner M, Woods SP, et al. Latent Toxoplasmosis is Associated with Worse Neurocognitive Functioning in HIV-Infected Adults and Correlates with Higher Toxoplasma IgG Levels submitted 2015.
 21. Gianella S, Morris SR, Anderson C, Spina CA, Vargas MV, Young JA, et al. Herpesviruses and HIV-1 Drug Resistance Mutations Influence the Virologic and Immunologic Milieu of the Male Genital Tract. *AIDS* 2013; 27(1):39–47. [PubMed: 22739399]
 22. Zagordi O, Klein R, Daumer M, Beerenwinkel N. Error correction of next-generation sequencing data and reliable estimation of HIV quasispecies. *Nucleic Acids Res* 2010; 38(21):7400–7409. [PubMed: 20671025]
 23. Cahn P, Sierra Madero J, Arribas J, Antinori A, Ortiz R, Clarke A, et al. Non-inferior efficacy of dolutegravir (DTG) plus lamivudine (3TC) versus DTG plus tenofovir/emtricitabine (TDF/FTC) fixed-dose combination in antiretroviral treatment-naïve adults with HIV-1 infection - 48-week results from the GEMINI studies. In: *AIDS Conference Amsterdam*; 2018.
 24. WHO. Prevention and treatment of HIV and other sexually transmitted infections among men who have sex with men and transgender People http://whqlibdoc.who.int/publications/2011/9789241501750_eng.pdf 2011.
 25. Imaz A, Martinez-Picado J, Niubo J, Kashuba AD, Ferrer E, Ouchi D, et al. HIV-1-RNA Decay and Dolutegravir Concentrations in Semen of Patients Starting a First Antiretroviral Regimen. *J Infect Dis* 2016; 214(10):1512–1519. [PubMed: 27578849]
 26. Else LJ, Taylor S, Back DJ, Khoo SH. Pharmacokinetics of antiretroviral drugs in anatomical sanctuary sites: the male and female genital tract. *Antivir Ther* 2011; 16(8):1149–1167. [PubMed: 22155899]

Table 1.

Summary of HIV RNA Shedders (N=3)

Parent study	Study week	ART regimen	Last missed Doses During study	Genital HIV RNA (copies/ml)	Plasma HIV RNA* (copies/ml)	CMV DNA (copies/ml)	HSV DNA (copies/ml)	Gonorrhea RNA	Chlamydia RNA
ASPIRE #1	48	RPV/TDF/FTC	1–2 weeks	42	179	Not detected	Not detected	Not detected	Not detected
ASPIRE #2**	36	DTG+3TC	> 3 months	488	<20	314607	Not detected	Not detected	Not detected
	48	DTG+3TC	Never	79	31	86090	Not detected	Not detected	Not detected
A5353	24	DTG+3TC	Never	48	<40	NA***	NA***	Not detected	Not detected

Legend: RPV: Rilpivirine, TDF: Tenofovir, FTC: Emtricitabine, DTG: Dolutegravir, 3TC: Lamivudine. NA: Not Available.

* Plasma HIV RNA collected at the same time as genital HIV RNA shedding

** ASPIRE participant #2 had detectable HIV RNA at two consecutive time-points (weeks 36 and 48)

*** Genital HSV and CMV testing could not be completed in one participant with HIV genital shedding because there was not enough seminal plasma to perform this analysis.