


## ORIGINAL ARTICLE

# Meridianin C inhibits the growth of YD-10B human tongue cancer cells through macropinocytosis and the down-regulation of Dickkopf-related protein-3

Nam-Sook Park<sup>1</sup> | Yu-Kyoung Park<sup>1</sup> | Mahesh Ramalingam<sup>1</sup> | Anil Kumar Yadav<sup>1</sup> |  
Hyo-Rim Cho<sup>1</sup> | Victor Sukbong Hong<sup>2</sup> | Kunal N. More<sup>2</sup> | Jae-Hoon Bae<sup>3</sup> |  
David Bishop-Bailey<sup>4</sup> | Junko Kano<sup>5</sup> | Masayuki Noguchi<sup>5</sup> | Ik-Soon Jang<sup>6</sup> |  
Kyung-Bok Lee<sup>6</sup> | Jinho Lee<sup>2</sup> | Jong-Soon Choi<sup>6,7</sup> | Byeong-Churl Jang<sup>1</sup> 

<sup>1</sup>Department of Molecular Medicine, College of Medicine, Keimyung University, Daegu, Republic of Korea

<sup>2</sup>Department of Chemistry, College of Natural Sciences, Keimyung University, Daegu, Republic of Korea

<sup>3</sup>Department of Physiology, College of Medicine, Keimyung University, Daegu, Republic of Korea

<sup>4</sup>Comparative Biomedical Sciences, Royal Veterinary College, London, UK

<sup>5</sup>Faculty of Medicine, Department of Pathology, University of Tsukuba, Tsukuba, Japan

<sup>6</sup>Biological Disaster Analysis Group, Division of Convergence Biotechnology, Korea Basic Science Institute, Daejeon, Republic of Korea

<sup>7</sup>Graduate School of Analytical Science and Technology, Chungnam National University, Daejeon, Republic of Korea

**Correspondence:** Jinho Lee, Department of Chemistry, College of Natural Sciences, Keimyung University, 1095 Dalgubeoldae-ro, Dalseogu, Daegu 42601, Republic of Korea (jinho@gw.kmu.ac.kr);  
Jong-Soon Choi, Biological Disaster Analysis Group, Division of Convergence Biotechnology, Korea Basic Science Institute, 169-148 Gwahakro, Yuseonggu, Daejeon 34133, Republic of Korea (jschoi@kbsi.re.kr) and  
Byeong-Churl Jang, Department of Molecular Medicine, College of Medicine, Keimyung University, 1095 Dalgubeoldae-ro, Dalseogu, Daegu 42601, Republic of Korea (jangbc123@gw.kmu.ac.kr)

**Funding information**

Korea Basic Science Institute, Grant/Award Number: D37403; National Research Foundation of Korea, Grant/Award Number: 2014R1A5A2010008

**Abstract**

Meridianin C is a marine natural product known for its anti-cancer activity. At present, the anti-tumour effects of meridianin C on oral squamous cell carcinoma are unknown. Here, we investigated the effect of meridianin C on the proliferation of four different human tongue cancer cells, YD-8, YD-10B, YD-38 and HSC-3. Among the cells tested, meridianin C most strongly reduced the growth of YD-10B cells; the most aggressive and tumorigenic of the cell lines tested. Strikingly, meridianin C induced a significant accumulation of macropinosomes in the YD-10B cells; confirmed by the microscopic and TEM analysis as well as the entry of FITC-dextran, which was sensitive to the macropinocytosis inhibitor amiloride. SEM data also revealed abundant long and thin membrane extensions that resemble lamellipodia on the surface of YD-10B cells treated with meridianin C, pointing out that meridianin C-induced macropinosomes was the result of macropinocytosis. In addition, meridianin C reduced cellular levels of Dickkopf-related protein-3 (DKK-3), a known negative regulator of macropinocytosis. A role for DKK-3 in regulating macropinocytosis in the YD-10B cells was confirmed by siRNA knockdown of endogenous DKK-3, which led to a partial accumulation of vacuoles and a reduction in cell proliferation, and by exogenous DKK-3 overexpression, which resulted in a considerable inhibition of the meridianin C-induced vacuole formation and decrease in cell

Nam-Sook Park and Yu-Kyoung Park contributed equally to this work.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2018 The Authors. Journal of Cellular and Molecular Medicine published by John Wiley & Sons Ltd and Foundation for Cellular and Molecular Medicine.

survival. In summary, this is the first study reporting meridianin C has novel anti-proliferative effects via macropinocytosis in the highly tumorigenic YD-10B cell line and the effects are mediated in part through down-regulation of DKK-3.

#### KEYWORDS

amiloride, DKK-3, macropinocytosis, meridianin C, YD-10B

## 1 | INTRODUCTION

Oral squamous cell carcinoma (OSCC) is the most common malignant tumour of the lip, oral cavity and oropharynx. Moreover, OSCC can be severely disfiguring leading to a poor quality of life, including loss of general cognitive, social, emotional or physical functions as well as social relationships.<sup>1</sup> The primary treatment of OSCC is surgical removal, but radiation therapy and/or chemotherapy have emerged as alternative treatment options for OSCC.<sup>2-4</sup> Characteristically, OSCC cells at the late stage of malignancies are very resistant to cancer therapy-mediated apoptosis. Thus, there is an urgent need for understanding resistance mechanism and identifying new drugs or substances which offer therapeutic efficacy against OSCC.

Macropinocytosis is an endocytic pathway that leads to the formation of single membrane irregular vesicles (macropinosomes) by the internalization of large patches of the plasma membrane along with associated extracellular fluid and solutes.<sup>5</sup> Unlike the canonical and endocytic pathways that depend on coat proteins such as clathrin or caveolin, one of the distinctive hallmarks of macropinocytosis is the formation of expansive membrane ruffles within the plasma membrane.<sup>6</sup> This membrane ruffling is initiated by the rapid polymerization of branching of actin filaments.<sup>7</sup> Multiple proteins and signalling factors are involved in macropinocytosis, including the Rho superfamily of GTPases (Rac, Cdc42), lipid components (phosphoinositides, cholesterol, phosphatidylinositol phosphates), phospholipid kinases, receptor tyrosine kinases and phosphatases, SNX-5 and DKK-3.<sup>8-16</sup> There is also large body of evidence that now shows macropinocytosis can be specifically activated by extracellular stimuli, such as growth factors and natural or synthetic chemicals, and the activated macropinocytosis can have effects on cell survival/proliferation or cell death/growth inhibition on distinct cell types including cancer cells.<sup>17,18</sup>

Meridianin C is one of the marine derived indole alkaloids (meridianin A-G) isolated from the South Atlantic tunicate *Aplidium meridianum*.<sup>19,20</sup> Previously, meridianin C, D or G analogues/derivatives have been shown to inhibit the proliferation of human breast (MCF-7) and cervix (HeLa) cancer and leukaemia (MV4-11) cells.<sup>21-23</sup> At present, neither the anti-tumour effect nor the mode of action of meridianin C in OSCC is known. In this study, we investigated the effect of meridianin C at 1  $\mu$ M concentration on growth of four different human oral carcinoma cell lines (YD-8, YD-10B, YD-38 and HSC-3) that were originated from tongue (YD-8, YD-10B, HSC-3) or lower gingiva (YD-38).<sup>24</sup> It is worth

noting that YD-8 and YD-38 cells have no tumorigenicity, but YD-10B cells are highly tumorigenic and invasive/metastatic.<sup>25,26</sup> In this article, we report for the first time that meridianin C strongly inhibits the growth of YD-10B OSCC cells through macropinocytosis, and its growth inhibitory and macropinocytosis inducing effects on the cells are mediated in part via the reduced expression of DKK-3.

## 2 | MATERIAL AND METHODS

### 2.1 | Cell culture and reagents

Four different human tongue cancer cell lines (YD-8, YD-10B, YD-38, HSC-3) and a human bladder cancer cell line (T24) were purchased from Korean Cell Line Bank (Seoul, Republic of Korea). Normal human gingival fibroblasts (HGF) were obtained from American Type Culture Collection (ATCC, CRL-2014<sup>TM</sup>, Manassas, VA). Cells were cultured in RPMI 1640 medium supplemented with 10% heat-inactivated foetal bovine serum (FBS), 100 units/mL penicillin and 100  $\mu$ g/mL streptomycin and maintained at 37°C in a humidified condition of 95% air and 5% CO<sub>2</sub>. Meridianin C was synthesized as previously described<sup>23</sup> and prepared as a 10 mM stock solution in DMSO. RPMI-1640 (LM011-01), FBS (S001-01) and penicillin-streptomycin (LS202-02) were purchased from Welgene (Deagu, Republic of Korea). All commercial antibodies and chemicals were purchased from the following resources: anti-procaspase-9 (ADI-AAM-139) was from Enzo (Farmingdale, NY, USA); anti-PARP (11835238001) was from Roche Diagnostics (Mannheim, Germany); anti-DR5 (NBP1-45951) was from Novus Biologicals (Littleton, CO, USA); anti-actin (A5441) was from Sigma (St. Louis, MO, USA); anti-DKK-3 (ab187532) was from Abcam (Cambridge, MA, USA); anti-Flag (#2368) from Cell Signalling Technology (Danvers, MA, USA); FITC-dextran (average molecular weight of 70 kDa) (46945) was from Sigma (St. Louis, MO, USA); z-VAD-fmk (627610) was from Calbiochem (Madison, WI, USA); Amiloride (A7410) was from Sigma (St. Louis, MO, USA) and protease inhibitor cocktail (PIC) (100x) (539134) were from Calbiochem (Madison, WI, USA). SuperSignal<sup>TM</sup> West Pico PLUS Enhanced chemiluminescence (ECL, #34080) was purchased from Thermo Scientific (Waltham, MA, USA). Plasticware: 6-well plates, 24-well plates and 60 or 100 mm cell culture dish were purchased from SPL Life Sciences (Gyeonggi-do, Korea). Control siRNA (sc-37007) and DKK-3 siRNA (sc-41102) were purchased from Santa Cruz biotechnology (Delaware, CA, USA).

## 2.2 | Cell count and cell morphology analysis

Respective OSCC cell line (YD-8, YD-10B, YD-38 or HSC-3) was seeded in 24-well plates ( $1 \times 10^5/500 \mu\text{L}/\text{well}$ ) overnight. After cells were treated with meridianin C ( $1 \mu\text{M}$ ) or vehicle control (DMSO; 0.1%) for 48 h, surviving cells, which cannot be stained with trypan blue dye, were counted using phase contrast microscope. Approximately 100 cells were counted for each evaluation. The cell count assay was performed in triplicate. Data are mean  $\pm$  SEM of three independent experiments. Survival is expressed as a percentage of control. In some experiments, YD-10B cells ( $0.5 \times 10^6/2 \text{ mL}/\text{well}$ ) or normal HGFs-1 ( $0.4 \times 10^6/2 \text{ mL}/\text{well}$ ) were also plated in 6-well plates overnight. Cells were then treated with meridianin C ( $1 \mu\text{M}$ ) or DMSO for indicated times (YD-10B cells for 2, 4, 8 or 24 h; normal HGFs for 4, 8, 24 or 48 h). Phase contrast images of the conditioned cells were taken with an Olympus phase contrast microscope equipped with a digital camera (Nikon).

## 2.3 | Transmission electron microscopy (TEM)

YD-10B cells ( $0.5 \times 10^6/2 \text{ mL}/\text{well}$ ) were seeded in 6-well plates the day before meridianin C treatment. Cells were then treated with  $1 \mu\text{M}$  meridianin C for 8 h, while DMSO was used as control. DMSO or meridianin C-treated cells were fixed overnight at  $4^\circ\text{C}$  with 2.5% glutaraldehyde in 0.1 M sodium cacodylate buffer, pH 7.4. Samples were post-fixed for 30 min with 0.5% osmium tetroxide/0.8% potassium ferricyanide, transferred to 1% tannic acid for 1 h and then to 1% uranyl acetate overnight at  $4^\circ\text{C}$ . Next, the samples were dehydrated with a graded ethanol series and embedded in Spurr's resin. Thin sections were cut with a Leica EM UC6 ultramicrotome (Leica), and stained with 1% uranyl acetate and Reynold's lead citrate prior to viewing at 120 kV on a Tecnai BT Spirit transmission electron microscope (FEI). Digital images were acquired with a Hamamatsu XR-100 side mount digital camera system (Advanced Microscopy Techniques) and processed using Adobe Photoshop CS5 (Adobe Systems Inc.).

## 2.4 | Scanning electron microscopy

YD-10B cells were seeded in 6-well plates ( $0.5 \times 10^6 \text{ cells}/2 \text{ mL}/\text{well}$ ) the day before meridianin C treatment. Cells were then treated with vehicle control (DMSO) or  $1 \mu\text{M}$  meridianin C for 2 h. The conditioned cells were centrifuged, fixed in 0.5% glutaraldehyde and 0.5% paraformaldehyde fixative, washed with 0.1 M phosphate-buffered saline (PBS), and post-fixed with 1% osmium tetroxide solution for 1 h. 2% tannic acid was used to conductively stain for 12 h, later washed with PBS, and fixed with 1% osmium tetroxide solution for 1 h. The specimens were dehydrated with ethanol, *t*-butyl alcohol and freeze dryer. Finally, the specimen was sputter coated with platinum-palladium (Pt-Pd) in an ion coater for 2 min, followed by microscopic examinations (S-4200, Hitachi Co., Tokyo, Japan).

## 2.5 | Measurement of DNA fragmentation

YD-10B cells were seeded in 6-well plates at a density of  $0.5 \times 10^6$  cells per well the day before treatment. Cells were incubated with meridianin C or vehicle control (DMSO) for 4 to 48 h, at which point, cells were harvested, washed and lysed in a buffer containing 50 mM Tris (pH 8.0), 0.5% sarkosyl, 0.5 mg/mL proteinase K and 1 mM EDTA at  $55^\circ\text{C}$  for 3 h, followed by addition of RNase A ( $0.5 \mu\text{g}/\text{mL}$ ) for a further 18 h at  $55^\circ\text{C}$ . The lysates were then centrifuged at 10 000 g for 20 min, genomic DNA in the supernatant was extracted with equal volume of neutral phenol-chloroform-isoamyl alcohol mixture (25:24:1), and analysed by electrophoresis on a 1.7% agarose gel. The DNA was visualized and photographed under UV illumination after staining with ethidium bromide ( $0.1 \mu\text{g}/\text{mL}$ ).

## 2.6 | Measurement of the population of sub G1 phase by flow cytometry analysis



After 24- or 48-h treatment with DMSO or meridianin C ( $1 \mu\text{M}$ ), YD-10 B cells were harvested and washed with PBS, fixed in ice-cold 70% ethanol and stored at  $4^\circ\text{C}$ . Cells were then washed once with PBS, suspended in 1 mL of cold propidium iodide (PI) solution containing 100  $\mu\text{g}/\text{mL}$  RNase A, 50  $\mu\text{g}/\text{mL}$  propidium iodide, 0.1% (w/v) sodium citrate and 0.1% (v/v) NP-40 and incubated on ice for additional 30 min in the darkness. Cytometric analyses were carried out with a flow cytometer (FACS Caliber, Becton Dickinson) and CellQuest software. Approximately, 10 000 cells were counted for the analysis.

## 2.7 | Fluorescein isothiocyanate (FITC) staining

To monitor the functionality of meridianin C-induced macropinocytosis (macropinosome formation/internalization),  $0.25 \times 10^5$  YD-10B cells/mL were seeded on coverslips and treated with meridianin C ( $1 \mu\text{M}$ ) and/or FITC-dextran ( $0.5 \text{ mg}/\text{mL}$ ) in the presence or absence of amiloride (4 mM) for 4 h. The cells were washed twice with PBS and mounted onto microscopic glass slides using Permafluor aqueous mounting media (Thermo Scientific, Waltham, MA, USA) media. Bright field and fluorescence were observed using a Zeiss AxioObserver.A1 inverted microscope (Carl Zeiss, Germany) and images acquired using Zen 2 software (Carl Zeiss). Fluorescent intensity was quantified using Image-J software.

## 2.8 | Preparation of whole cell lysates

To see the effect of meridianin C on expression of apoptosis- or macropinocytosis-related proteins, YD-10B cells ( $0.5 \times 10^6/2 \text{ mL}/\text{well}$ ) were seeded in 6-well plates the day before meridianin C treatment. Cells were treated with meridianin C ( $1 \mu\text{M}$ ) or vehicle control (DMSO) for the indicated times. At each time-point, cells were washed twice with PBS and proteins extracted using a modified RIPA buffer (50 mM Tris-Cl (pH 7.4), 150 mM NaCl, 0.1% sodium dodecyl sulphate, 0.25% sodium deoxycholate, 1% Triton X-100, 1%



Nonidet P-40, 1 mM EDTA, 1 mM EGTA, PIC (1×)). The cell lysates were collected and centrifuged at 12 000 rpm for 20 min at 4°C. The supernatants were saved and protein concentrations determined by bicinchoninic acid assay (BCA) protein assay (Pierce).

## 2.9 | Immunoblot analysis

Proteins (50 µg) were separated by SDS-PAGE (10%) and transferred onto nitrocellulose membranes (Millipore, Bedford, MA, USA). The membranes were washed with TBS (10 mM Tris-Cl, 150 mM NaCl, pH 7.5) with 0.05% (v/v) Tween-20 followed by blocking with TBST containing 5% (w/v) non-fat dried milk. The membranes were incubated overnight with antibodies specific for procaspase-9 (1:1000), DR-5 (1:1000), PARP (1:2000), DKK-3 (1:1000), Flag (1:1000) or  $\beta$ -actin (1:10 000) at 4°C. The membranes were then exposed to secondary antibodies conjugated to horseradish peroxidase for 2 h at room temperature and further washed three times with TBST. Immunoreactivity was detected by SuperSignal™ West Pico PLUS enhanced chemiluminescence (ECL) according to manufacturer (Thermo Scientific, Waltham, MA, USA). Equal protein loading was assessed by the expression levels of  $\beta$ -actin.

## 2.10 | Small interfering RNA (siRNA) transfection

YD-10B cells ( $0.5 \times 10^5$ /well) seeded into 6-well plates were transfected for 6 h with control or DKK-3 siRNA (80 pM) using Lipofectamine™ 2000 (Invitrogen, Carlsbad, CA, USA). Culture medium from the transfected cells was removed and refreshed with RPMI containing 10% FBS, followed by incubation for 18 h. The knock-down efficiency of the DKK-3 siRNA was determined by immunoblotting in comparison to  $\beta$ -actin.

## 2.11 | Generation of stable YD-10B cells overexpressing DKK-3

The pcDNA3.1-DKK3-C-Flag or mock vector was constructed as reported previously.<sup>27</sup> Briefly, YD-10B cells were transfected in a stable manner with the pcDNA3.1-DKK-3-Flag or mock vector using LipofectAMINE transfection reagent as prescribed by the manufacturer (Invitrogen, Carlsbad, CA, USA). After 48 h of incubation, transfected cells were selected in primary cell culture medium containing 200 µg/mL G418 (Promega, Madison, WI, USA). After 2 or 3 weeks, to rule out the possibility of clonal differences between the generated stable cell lines, the pooled YD-10B/mock vector and YD-10B/

DKK-3-Flag clones were tested for DKK-3 and Flag expression by immunoblotting and were used in this study.

## 2.12 | Statistical analyses


Cell count analysis was performed in triplicates and repeated three times. Data were expressed as mean  $\pm$  SEM. The significance of difference was determined by one-way ANOVA (Laerd Statistics, Chicago, IL, USA). All significance testing was based upon a  $P < 0.05$ .

# 3 | RESULTS

## 3.1 | Meridianin C has strong growth inhibitory and vacuole-inducing effects on YD-10B cells

We investigated the treatment effect of meridianin C (Figure 1A) at 1 µM concentration for 48 h on the growth of four different human tongue carcinoma cell lines (YD-8, YD-10B, YD-38 and HSC-3) by cell count analysis. Among the cell lines tested, YD-10B cells were the most sensitive to meridianin C (Figure 1B). Meridianin C treatment led to a slight reduction in HSC-3 cells, which was statically insignificant. Both YD-38 and YD-8 and cells were insensitive to meridianin C treatment. Strikingly, YD-10B cells treated with meridianin C were highly vacuolated (Figure 1C). Furthermore, HSC-3 cells treated with meridianin C also formed many vacuoles in much smaller size than those in YD-10B cells. However, both YD-38 and YD-8 cells treated with meridianin C had very low or no cytoplasmic vacuoles. Meridianin C caused a time-dependent increase in cytoplasmic vacuole formation in YD-10B cells, with effects visible by 2 h (Figure 1D). We next examined the effect of meridianin C on growth and accumulation of vacuoles in normal human gingival fibroblast (HGF), but no cytoplasmic vacuoles were observed at 4, 8, 24 or 48 h, and meridianin C was not cytotoxic to HGF (Figure 1E). However, it was shown that treatment with meridianin C at higher concentrations (5 or 10 µM) and different times (2, 4, 8, 24 or 48 h) strongly reduced survival and induced massive vacuoles of both YD-10B and normal HGFs, pointing out that meridianin C at higher concentrations (5 or 10 µM) than 1 µM are cytotoxic to YD-10B cells as well as normal HGFs. These results show that meridianin C has a strong and selective anti-survival and vacuole-inducing effects on the highly invasive and tumorigenic YD-10B cells at the 1 µM concentration. Therefore, we decided to focus on the YD-10B cell line and the 1 µM concentration of meridianin C for further studies.

**FIGURE 1** Meridianin C inhibits YD-10B cell survival and induces cellular vacuole formation. (A) The chemical structure of meridianin C. (B) Four human tongue cancer cell lines (YD-8, YD-10B, YD-38, and HSC-3), respectively, were treated with meridianin C (1 µM) or vehicle control (DMSO; 0.1%) for 48 h. The number of surviving cells was measured by trypan blue dye exclusion. Data are mean  $\pm$  SEM of three independent experiments, each done in triplicate. \* $P < 0.05$  compared to the value of meridianin C free control at the indicated time. (C) YD-8, YD-10B, YD-38 and HSC-3 cells, respectively, were treated with meridianin C (1 µM) or vehicle control (DMSO) for 48 h. Images of the conditioned cells were obtained by phase contrast microscopy, 400 X. Each image is a representative of three independent experiments. (D, E) YD-10B cells or normal human gingival fibroblasts (HGF) were treated with meridianin C (1 µM) or vehicle control (DMSO) for the times indicated. At each time-point, images were obtained by phase contrast microscopy, 400 X. Each image is a representative of three independent experiments



**FIGURE 2** Meridianin C does not induce apoptosis of YD-10B cells. (A) YD-10B cells were treated with meridianin C (1  $\mu$ M) or vehicle control (DMSO) for the times indicated. At each time-point, extra-nuclear fragmented DNA was extracted, and analysed on a 1.7% agarose gel. The images are representative of three independent experiments. SM, DNA size marker. (B) YD-10B cells were treated with meridianin C (1  $\mu$ M) or vehicle control (DMSO) for the times indicated and evaluated for apoptotic subG1 DNA content by flow cytometer. The tables represent the fraction of apoptotic cells. Data are mean  $\pm$  SEM of three independent experiments. (C) YD-10B cells were treated with meridianin C (1  $\mu$ M) or vehicle control (DMSO) for the times designated. At each time-point, whole cell lysates were prepared, and analysed for procaspase-9, DR5, PARP cleavage and  $\beta$ -actin by Western blotting. The images are representative of three independent experiments. (D) YD-10B cells were treated with meridianin C (1  $\mu$ M) in the absence or presence of z-VAD (50  $\mu$ M), a pan-caspase inhibitor, for 48 h, and cell survival counted by the trypan blue dye exclusion assay. Data are mean  $\pm$  SEM of three independent experiments, each performed in triplicate. \* $P < 0.05$  compared to the control at the indicated time. (E) YD-10B cells were treated with meridianin C (1  $\mu$ M) in the absence or presence of z-VAD (50  $\mu$ M), a pan-caspase inhibitor, for 48 h. Images were obtained by phase contrast microscopy, 400 X. Each image is representative of three independent experiments


### 3.2 | Meridianin C does not induce apoptosis in YD-10B cells

We next determined whether meridianin C treatment induces apoptosis in YD-10B cells by nuclear DNA fragmentation assay. As shown in Figure 2A, no apoptotic DNA fragmentation occurred in YD-10B cells when treated with meridianin C for 4, 8, 24 or 48 h. Further to confirm apoptosis, we analysed the amount of subG1 DNA by flow cytometry in YD-10B cells treated with 1  $\mu$ M meridianin C for the indicated times (24, 48 h). As shown in Figure 2B, the addition of meridianin C did not result in increased accumulation of sub-G1 phase, a well-known apoptotic marker. Further, Western blotting analysis revealed that meridianin C treatment did not affect expression levels of apoptosis-related proteins, procaspase-9 and death receptor-5 (DR-5), or PARP cleavage (Figure 2C) until 48 h of treatment. Moreover, treatment with z-VAD-fmk, a pan-caspase inhibitor<sup>28</sup> did not block the meridianin C-induced growth inhibition and accumulation of vacuoles in YD-10B cells (Figure 2D, E).

### 3.3 | Meridianin C induces macropinosomes (macropinosomes) in YD-10B cells

Using transmission electron microscopy (TEM), we investigated whether vacuoles induced by meridianin C in YD-10B cells are macropinosomes. While YD-10B cells treated with vehicle control (DMSO) had no or tiny vacuoles in the cytoplasm (Figure 3A), the meridianin C-treated cells displayed ultrastructural characteristics of macropinosomes (vacuoles) in different sizes (arrows) (Figure 3B). Under high magnification, the meridianin C-treated YD-10B cells showed fusion between large and small vacuoles in the cytoplasm (arrows) and indentation of the nuclear membrane (arrowhead) probably because of enlarged vacuolar structure (Figure 3C). Furthermore, residual materials, including membranous debris (arrowheads), were seen within the cytoplasmic vacuoles (arrows) in YD-10B cells treated with meridianin C (Figure 3D). It is documented that macropinosomes are derived from membrane ruffles and protrusions folding back and fusing with the plasma membrane to form large

**FIGURE 3** Transmission and scanning electron micrographs of meridianin C-induced macropinosomes and macropinocytosis in YD-10B cells. (A–D) YD-10B cells were treated with meridianin C (1  $\mu$ M) or vehicle control (DMSO) for 8 h. (A) Vehicle (DMSO)-treated cells, 2000 X. (B) Meridianin C-treated cells, 2000 X. Arrows indicate the cytoplasmic vacuoles of different sizes. (C) Meridianin C-treated cells, 4000 X. Arrowhead indicates the indented or distorted nucleus because of enlarged vacuole. Arrows indicate fusion between cytoplasmic vacuoles of different sizes. (D) Meridianin C-treated cells, 5000 X. Arrows indicate macropinosomes containing membranous contents in different sizes, which are shown by arrowheads. (E, F) YD-10B cells were treated with meridianin C (1  $\mu$ M) or vehicle control (DMSO) for 2 h. (E) Vehicle control treated cells, which show many short and thick bacilli bound to the cell surface (thin arrows). (F) Meridianin C-treated cells, which exhibit abundant long and thin membrane extensions that resemble lamellipodia on the cells surface (wide white arrows)



vesicles.<sup>29</sup> Further, to confirm that the formation of macropinosomes induced by meridianin C is the result of macropinocytosis, scanning electron microscopy (SEM) was used to see the effect of meridianin C on induction of membrane ruffling or lamellipodia formation in YD-10B cells. Apparently, while vehicle control (DMSO)-treated YD-10B cells had many short and thick bacilli bound to the cell surface (thin arrows) (Figure 3E), there were abundant long and thin membrane extensions that resemble lamellipodia (wide white arrows) on the surface of cells treated with meridianin C for 2 h (Figure 3F).


### 3.4 | Meridianin C-induced macropinocytosis (macropinosome formation/internalization) in YD-10B cells is functional

Amiloride inhibits macropinocytosis by blocking the activity of Na/H exchanger, causing a decrease in pH at the plasma membrane.<sup>30</sup> This amiloride-induced pH drop suppresses the membrane ruffling and membrane extensions that form the macropinosome.<sup>31</sup> Soluble

sugars or proteins, such as dextran or albumin, are useful fluorescent markers specific for macropinocytosis.<sup>32</sup> Using amiloride and fluorescein isothiocyanate-dextran with molecular weight of 70 kDa (FITC-dextran), we sought to explore whether the meridianin C-induced macropinocytosis (macropinosome formation/internalization) in YD-10B cells is functional. Microscopic observations clearly demonstrated that amiloride strongly blocked the meridianin C-induced accumulation of vacuoles in YD-10B cells (Figure 4A). Moreover, meridianin C increased the delivery of FITC-dextran into YD-10B cells while amiloride strongly inhibited it (Figure 4B). Further, fluorescence intensity of FITC-dextran (green colour) was determined by Image-J software (Figure 4C).

### 3.5 | Meridianin C-induced down-regulation of DKK-3 protein in YD-10B cells

DKK-3 plays a role in modulating macropinocytosis in T24 human bladder cancer cells<sup>16</sup> and is also expressed in OSCC-derived cell



**FIGURE 4** Suppression of meridianin C-induced macropinocytosis (macropinosomes) in YD-10B cells by amiloride, a macropinocytosis inhibitor. (A) YD-10B cells were treated with meridianin C (1  $\mu$ M) in the absence or presence of amiloride (4 mM) for 4 h. Images were obtained by phase contrast microscopy, 400 X. Each image is a representative of three independent experiments. (B) YD-10B cells were seeded on coverslips and treated with meridianin C (1  $\mu$ M) and/or FITC-dextran (0.5 mg/ml) in the absence or presence of amiloride (4 mM) for 4 h. The cells were washed and mounted onto microscopic glass slides. Bright field and fluorescence were observed with a Zeiss AxioObserver.A1 inverted microscope and images were acquired using Zen 2 software. (C) Fluorescence intensity of FITC-dextran (green colour) was determined by Image-J software. Each value in the graph represents the mean  $\pm$  SEM of three independent experiments. \* $P < 0.05$  compared to the meridianin C treatment at the indicated time

lines.<sup>33</sup> We therefore sought to explore whether DKK-3 is expressed in YD-10B cells and is regulated by meridianin C treatment. As shown in Figure 5A, high cellular levels of DKK-3 protein were seen in YD-10B cells, and its expression level was even higher relative to  $\beta$ -actin than that in T24 cells. There was no detectable DKK-3 protein expression in YD-8 or YD-38 cells. Time course experiments further revealed that meridianin C treatment led to a rapid time-dependent reduction in DKK-3 protein in YD-10B cells (Figure 5B-D).


### 3.6 | DKK-3 knock-down led to partial accumulation of vacuoles in YD-10B cells along with partial decrease of the cell growth

To evaluate the role of reduced DKK-3 in meridianin C's growth inhibition and/or macropinocytosis in YD-10B cells, we transfected control or DKK-3 siRNA into YD-10B cells and measured the number of surviving cells and accumulation of cytoplasmic vacuoles. There was almost complete knock-down of DKK-3 in the DKK-3 siRNA-transfected YD-10B cells, compared with the control siRNA-transfected ones (Figure 6A), suggesting high DKK-3 siRNA transfection efficiency. YD-10B cells transfected with DKK-3 siRNA had partial accumulation of vacuoles (Figure 6B) accompanied with a partial but significant reduction in cell survival (Figure 6C).

### 3.7 | Overexpression of DKK-3 largely blocks meridianin C-induced accumulation of vacuoles in YD-10B cells along with significant increase of the cell survival

To further clarify that DKK-3 plays an important role in the function of meridianin C, we next performed DKK-3-Flag cDNA transfection experiments to generate stable YD-10B cells that overexpress DKK-3-Flag. Using these stable clones, we tested whether DKK-3 overexpression could interfere with the meridianin C's growth inhibitory and vacuole-inducing activities in YD-10B cells. As shown in Figure 7A, there was a large increase in DKK-3 protein levels in YD-10B<sup>DKK-3</sup> cells (cells overexpressing DKK-3-Flag) compared with YD-10B<sup>mock</sup> cells (cells transfected with mock vector). High levels of Flag expression in YD-10B<sup>DKK-3</sup>, but not in YD-10B<sup>mock</sup>, cells further support the DKK-3 cDNA transfection efficiency. As expected, meridianin C led to a decrease in DKK-3 protein levels in YD-10B<sup>mock</sup> cells. However, meridianin C did not influence DKK-3 protein expression in YD-10B<sup>DKK-3</sup> cells, although it slightly reduced Flag levels in the cells. Importantly, while meridianin C treatment largely induced many vacuoles in YD-10B<sup>mock</sup> cells, it had no effect on the formation of vacuoles in YD-10B<sup>DKK-3</sup> cells (Figure 7B). Furthermore, as shown in Figure 7C, the meridianin C-induced reduction in YD-10B cell survival was considerably blocked by DKK-3 overexpression.






**FIGURE 5** Effect of meridianin C on expression of DKK-3 in YD-10B cells. (A) Whole cell lysates of YD-8, YD-10B, YD-38 or T24 cells were analysed for DKK-3 or  $\beta$ -actin by Western blotting. (B) YD-10B cells were treated with meridianin C (1  $\mu$ M) or vehicle control (DMSO) for the times designated or (C) for 20 min. At each time-point, whole cell lysates were prepared, and analysed for DKK-3 compared to  $\beta$ -actin by Western blotting. (D) The densitometry data of (C). \* $P < 0.05$  compared to the control (-) at the indicated time

### 3.8 | The meridianin C's growth inhibitory and macropinocytosis inducing effects on YD-10B cells are reversible

To determine whether meridianin C exerts its effects on YD-10B cell survival and macropinocytosis in a reversible or irreversible manner, YD-10B cells were incubated with meridianin C or DMSO solvent control at the designated time-points, as illustrated in Figure 8A. YD-10B cells that were incubated with vehicle DMSO control for 4 h and then incubated with DMSO vehicle for an additional 1, 2, 4 and 20 h, showed no vacuole formation (treatment a) (Figure 8B). In contrast, YD-10B cells that were incubated with meridianin C for 4 h and then further incubated with meridianin C for an additional 1, 2, 4 and 20 h, respectively, showed many vacuoles (treatment b). However, YD-10B cells that were incubated with meridianin C for 4 h and then incubated with DMSO vehicle control for an additional 1, 2, 4 and 20 h, respectively, showed a time-dependent reduction in




**FIGURE 6** Effects of DKK-3 knockdown on accumulation of vacuoles and growth of YD-10B cells. (A) YD-10B cells were transfected with 80 pM of control or DKK-3 siRNA for 24 h. Whole cell lysates were prepared, and analysed by Western blot. The pictures are representative of three independent experiments. (B) YD-10B cells were transfected with 80 pM of control or DKK-3 siRNA for 24 h. Images were obtained by phase contrast microscopy, 400 X. Each image is representative of three independent experiments. (C) YD-10B cells were transfected with 80 pM of control or DKK-3 siRNA for 24 h, and cell survival counted by the trypan blue dye exclusion assay. Data are mean  $\pm$  SEM of three independent experiments, each performed in triplicate. \* $P < 0.05$  compared to the value of control siRNA at the indicated time

vacuole number (treatment c). Importantly, the survival of YD-10B cells in treatment c was much higher than that of the cells in treatment b (Figure 8C). These results show that the growth inhibitory and macropinocytosis inducing effects of meridianin C on YD-10B cells do occur in a reversible manner.

### 3.9 | Comparison of the effects of meridianin C and its derivatives on growth and macropinosome (vacuole) accumulation in YD-10B cells

We have recently synthesized a novel series of meridianin C derivatives substituted at C-5 position (7a-j) and reported their anti-



**FIGURE 7** Effects of DKK-3 overexpression on meridianin C-induced accumulation of vacuoles and decrease in survival of YD-10B cells. (A) YD-10B cells were transfected with pcDNA3.1-DKK-3-Flag or mock vector using LipofectAMINE 2000 reagents. Stable clones were selected in the culture media containing 200 μg/ml G418 sulphate for several weeks. The selected clones (DKK-3-Flag or mock vector-transfected YD-10B cells) were seeded overnight. Cells were then treated with or without meridianin C (1 μM) for 20 min. Cell lysates were prepared and analysed by Western blotting for measurement of the expression levels of DKK-3, Flag and β-actin. (B) DKK-3- or mock-transfected YD-10B cells were treated vehicle or meridianin C (1 μM) for 4 h. Images were obtained by phase contrast microscopy, 400 X. Each image is a representative of three independent experiments. (C) DKK-3- or mock-transfected YD-10B cells were treated with or without meridianin C (1 μM) for 24 h, followed measurement of the number of cells survived by the trypan blue dye exclusion assay. Data are mean ± SEM of three independent experiments, each performed in triplicate. \* $P < 0.05$  compared to the control at the indicated time


proliferative activities against human leukaemic cell lines.<sup>23</sup> In this study, we tested these meridianin C derivatives on growth and macropinosome (vacuole) accumulation of YD-10B cells. Meridianin C treatment for 48 h strongly inhibited the survival of YD-10B cells (Figure 9A). However, treatment with 7a, 7b, 7c, 7f, 7e, 7 h or 7i only slightly decreased of the cell survival. Treatment with 7d, 7 g or 7j had no effect on the survival of YD-10B cells. Similarly, meridianin C treatment induced a significant accumulation of vacuoles in YD-10B cells, whereas compounds 7a-j had no or very low vacuole-inducing activity (Figure 9B).

## 4 | DISCUSSION

Macropinocytosis can be activated and exploited in normal and cancer cells to confer a cell growth/survival advantage.<sup>34–36</sup> By contrast, hyper-stimulation of macropinocytosis might play a critical role in the cytotoxicity of normal and cancer cells.<sup>35,37</sup> Notably, there are a number of several natural or synthetic chemicals that have anti-tumour effects through stimulation of macropinocytosis.<sup>38,39</sup> Indeed, it has been suggested that any substance that specifically and massively induces macropinocytosis in cancer cells may have therapeutic potential as an anti-tumour agent. Regulation of human tongue cancer cell growth and macropinocytosis by meridianin C has not been reported previously. In this study, we demonstrate that meridianin C has strong growth suppressive and

macropinocytosis inducing effects on YD-10B human tongue cancer cells and the effects are mediated in part through regulation of DKK-3 expression.

Aforementioned, meridianin C, D or G analogues/derivatives are anti-proliferative in several types of human cancer cells, including breast (MCF-7), cervix (HeLa) and leukaemia (MV4-11).<sup>21–23</sup> Here we initially screened four different human tongue cancer cell lines (YD-8, YD-10B, YD-38 and HSC-3) to see whether meridianin C treatment affects the cell survival. We found that meridianin C strongly and specifically reduced the survival of YD-10B cells. However, the apoptotic parameters, including nuclear DNA fragmentation, population of sub G1, expression of procaspase-9 (no expression of cleaved and active form of caspase-9) and PARP cleavage, did not indicate apoptosis as the mechanism of action of meridianin C. Moreover, the microscopic and TEM analyses in this study showed that meridianin C rapidly induced cytoplasmic vacuoles/macropinosomes in YD-10B cells. The SEM analysis further demonstrated, the ability of meridianin C to rapidly induce abundant long and thin membrane extensions that resemble lamellipodia on the surface of YD-10B cells. These results strongly support the notion that the formation of macropinosomes induced by meridianin C in YD-10B cells is the result of macropinocytosis. Meridianin C has previously been reported to inhibit Pim-1, CDK-1, CDK-5, PKA, PKG, GSK-3β and CK-1 activity with IC<sub>50</sub> values of 1.4, 3, 6, 0.7, 0.4, 2 and 30 μM, respectively.<sup>23,40</sup> There are no reports showing a relationship between cell survival or macropinocytosis induction and those




kinases in cancer cells including YD-10B cells. However, one can speculate that meridianin C's growth inhibitory and/or macropinocytosis inducing effects on YD-10B cells may be through acting upon one or more of these kinases, which yet remain to be elucidated.

A notable finding of the present study is meridianin C regulation of DKK-3 expression in YD-10B cells. DKK-3 is one of the DKK protein families and is known as a Wnt antagonist.<sup>41,42</sup> It is also recognized as a potential tumour suppressor, based on the facts that DKK-3 expression is frequently down-regulated in a wide array of malignancies.<sup>43,44</sup> Recent evidence indicates that DKK-3 may also play additional roles in cancer cell survival. DKK-3 knock-down by siRNA in T24 human bladder cancer cells inhibited cell growth and induced macropinocytosis and autophagy.<sup>16</sup> It also has been recently demonstrated that DKK-3 is expressed in a subset of OSCC-derived cell lines including HSC-4.<sup>33</sup> Moreover, in HSC-4 cells, DKK-3 knock-down inhibited migration and invasion. These results imply that DKK-3 may have a metastasis-related oncogenic function in OSCC. Here we found DKK-3 is highly expressed in YD-10B cells and meridianin C suppresses its expression. To the best of our knowledge, this is the first study showing the ability of meridianin C to down-regulate DKK-3 in cancer cells. Of further importance, the present study revealed that siRNA-based gene silencing of endogenous DKK-3 led to partial accumulation of vacuoles in YD-10B cells along with partial inhibition of

the cell survival. Moreover, we herein have demonstrated that overexpression of DKK-3 blocks the macropinocytosis inducing property of meridianin C and leads to increase in cell survival against meridianin C treatment. Altogether, these findings show that DKK-3 may have a role in controlling survival and macropinocytosis in YD-10B cells and this is inhibited by meridianin C.

The reversibility of cancer cell growth inhibition after drug removal has clinical implications. In this study, we further evaluated the effects of meridianin C withdrawal on cell survival and macropinocytosis in YD-10B cells. We found that meridianin C's anti-survival and macropinocytosis inducing effects on YD-10B cells were reversible. Several reports have recently described that hyper-stimulation of macropinocytosis and/or disruption of normal macropinosome maturation associated with abnormal/massive accumulation of intracellular vacuoles leads to the cytotoxicity and/or nonapoptotic cell death of cancer cells.<sup>45-47</sup> We would similarly speculate that the anti-survival effect of meridianin C on YD-10B cells is closely linked to this drug-induced rapid and sustained accumulation of macropinosomes in the cells. We have similarly found that meridianin C can induce macropinocytosis (macropinosomes) in HSC-3 cells, another human tongue cancer cell line, indicating that the meridianin C's macropinocytosis effects are not limited to YD-10B cells. Further characterization of meridianin C's responses in YD-10B cells compared to YD-8 and



**FIGURE 9** Effects of meridianin C and its derivatives on growth and vacuole formation of YD-10B cells. (A) YD-10B cells were treated with 1  $\mu$ M of meridianin C or each derivative of meridianin C (a to j) for 48 h, and the cell count assay performed in triplicate. Data are mean  $\pm$  SEM of three independent experiments. \* $P$  < 0.05 compared to the control at the indicated time. (B) YD-10B cells were treated with 1  $\mu$ M of meridianin C or each derivative (a to j) for 48 h. Images were obtained by phase contrast microscopy, 400 X. Each image is representative of three independent experiments

YD-38 cells, which are unresponsive to meridianin C, may further help to elucidate its mechanism of action.

We have recently synthesized a series of meridianin C derivatives, and demonstrated their anti-proliferative activity in human leukaemia cell lines including MV4-11.<sup>23</sup> In the current study, we further compared the growth inhibitory and/or macropinocytosis inducing effects of meridianin C and its derivatives on YD-10B cells. Although some of meridianin C derivatives had macropinocytosis inducing and growth inhibitory effects on YD-10 cells, it was evident that their effects were much weaker than those induced by its parental compound meridianin C. These results indicate that meridianin C has a unique structural moiety leading to strong growth suppressive and macropinocytosis inducing activity in YD-10B cells, which is distinct from other cancer cells.

In summary, meridianin C induces a significant level of macropinocytosis and selectively decreases the survival of YD-10B cells in part through a reduction in DKK-3. Although there are still important issues that remain to be resolved, including meridianin C's macropinocytosis inducing and anti-tumour effects on animal models, our findings suggest that meridianin C is a potential novel target for metastatic human tongue cancer.

#### ACKNOWLEDGEMENTS

This work was supported by the Korea Basic Science Institute (grant no. D37403) and the National Research Foundation of Korea (NRF)

grant funded by the Korea Government (MSIP) (no. 2014R1A5A2010008).

#### CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

#### AUTHORS CONTRIBUTION

N.-S.P., Y.-K.P., H.-R.C., J.-H.B., K.N.M., M.R., J.K., M.N. and A.K.Y performed experiments. J.L., J.-S.C. and B.-C.J. designed the research study and supervised the study. V.S.H., D.B.-B. and B.-C.J. wrote the manuscript. J.L., J.K., M.N., K.-B.L., I.-S.J., D.B.-B. and B.-C.J. analysed the data.

#### ORCID

Byeong-Churl Jang  <http://orcid.org/0000-0002-2691-5503>

#### REFERENCES

- Zain RB. Cultural and dietary risk factors of oral cancer and precancer—a brief overview. *Oral Oncol.* 2001;37:205-210.
- Claus F, Duthoy W, Botterberg T, et al. Intensity modulated radiation therapy for oropharyngeal and oral cavity tumors: clinical use and experience. *Oral Oncol.* 2002;38:597-604.

3. Nakazawa M, Ohnishi T, Ohmae M, et al. Phase II study of a novel oral formation of 5-fluorouracil in combination with low-dose cisplatin as preoperative chemotherapy of oral squamous cell carcinoma. *Int J Clin Pharmacol Res.* 2005;25:115-122.
4. Vokes EE, Weichselbaum RR, Lippman SM, Hong WK. Head and neck cancer. *N Engl J Med.* 1993;328:184-194.
5. Falcone S, Cocucci E, Podini P, et al. Macropinocytosis: regulated coordination of endocytic and exocytic membrane traffic events. *J Cell Sci.* 2006;119:4758-4769.
6. Doherty GJ, McMahon HT. Mechanisms of endocytosis. *Annu Rev Biochem.* 2009;78:857-902.
7. Mullins RD, Heuser JA, Pollard TD. The interaction of Arp2/3 complex with actin: nucleation, high affinity pointed end capping, and formation of branching networks of filaments. *Proc Natl Acad Sci U S A.* 1998;95:6181-6186.
8. Grimmer S, van Deurs B, Sandvig K. Membrane ruffling and macropinocytosis in A431 cells require cholesterol. *J Cell Sci.* 2002;115:2953-2962.
9. Mañes S, Ana Lacalle R, Gómez-Moutón C, Martínez-A C. From rafts to crafts: membrane asymmetry in moving cells. *Trends Immunol.* 2003;24:320-326.
10. Innocenti M, Zucconi A, Disanza A, et al. Abi1 is essential for the formation and activation of a WAVE2 signalling complex. *Nat Cell Biol.* 2004;6:319-327.
11. Redelman-Sidi G, Iyer G, Solit DB, Glickman MS. Oncogenic activation of Pak1-dependent pathway of macropinocytosis determines BCG entry into bladder cancer cells. *Cancer Res.* 2013;73:1156-1167.
12. Levin R, Grinstein S, Schlam D. Phosphoinositides in phagocytosis and macropinocytosis. *Biochim Biophys Acta.* 2015;1851:805-823.
13. Viaud J, Mansour R, Antkowiak A, et al. Phosphoinositides: important lipids in the coordination of cell dynamics. *Biochimie.* 2016;125:250-258.
14. Lanzetti L, Palamidessi A, Arecas L, et al. Rab5 is a signalling GTPase involved in actin remodelling by receptor tyrosine kinases. *Nature.* 2004;429:309-314.
15. Lim JP, Teasdale RD, Gleeson PA. SNX5 is essential for efficient macropinocytosis and antigen processing in primary macrophages. *Biol Open.* 2012;1:904-914.
16. Tsujimura N, Yamada NO, Kuranaga Y, et al. Novel role of Dickkopf-related protein 3 in macropinocytosis in human bladder cancer T24 cells. *Int J Mol Sci.* 2016;17:E1846.
17. Li C, MacDonald JI, Talebian A, et al. Unravelling the mechanism of TrkA-induced cell death by macropinocytosis in medulloblastoma daoy cells. *Mol Cell Biol.* 2016;36:2596-2611.
18. Recouvreur MV, Commisso C. Macropinocytosis: a metabolic adaptation to nutrient stress in cancer. *Front Endocrinol (Lausanne).* 2017;8:261.
19. Gompel M, Leost M, De Kier Joffe EB, et al. Meridianins, a new family of protein kinase inhibitors isolated from the ascidian *Aplidium meridianum*. *Bioorg Med Chem Lett.* 2004;14:1703-1707.
20. Bharate SB, Yadav RR, Battula S, Vishwakarma RA. Meridianins: marine derived potent kinase inhibitors. *Mini Rev Med Chem.* 2012;12:618-631.
21. Giraud F, Alves G, Debiton E, et al. Synthesis, protein kinase inhibitory potencies, and in vitro antiproliferative activities of meridianin derivatives. *J Med Chem.* 2011;54:4474-4489.
22. Radwan MA, El-Sherbiny M. Synthesis and antitumor activity of indolylpyrimidines: marine natural product meridianin D analogues. *Bioorg Med Chem.* 2007;15:1206-1211.
23. More KN, Jang HW, Hong VS, et al. Pim kinase inhibitory and antiproliferative activity of a novel series of meridianin C derivatives. *Bioorg Med Chem Lett.* 2014;24:2424-2428.
24. Lee EJ, Kim J, Lee SA, et al. Characterization of newly established oral cancer cell lines derived from six squamous cell carcinoma and two mucoepidermoid carcinoma cells. *Exp Mol Med.* 2005;37:379-390.
25. Lee G, Kim HM. JNK-associated scattered growth of YD-10B oral squamous carcinoma cells while maintaining the epithelial phenotype. *Biochem Biophys Res Commun.* 2017;487:862-867.
26. Kim KR, Park KK, Chung WY, Hwang YS. The inhibitory effect of buddlejasaponin IV on the growth of YD-10B human oral squamous cell carcinoma cells. *J Cancer Prev.* 2013;18:330-336.
27. Qui S, Kano J, Noguchi M. Dickkopf 3 attenuates xanthine dehydrogenase expression to prevent oxidative stress-induced apoptosis. *Genes Cells.* 2017;22:406-417.
28. Rajah T, Chow SC. The inhibition of human T cell proliferation by the caspase inhibitor z-VAD-FMK is mediated through oxidative stress. *Toxicol Appl Pharmacol.* 2014;278:100-106.
29. Kerr MC, Teasdale RD. Defining macropinocytosis. *Traffic.* 2009;10:364-371.
30. Alvarez de la Rosa D, Canessa CM, Fyfe GK, Zhang P. Structure and regulation of amiloride-sensitive sodium channels. *Annu Rev Physiol.* 2000;62:573-594.
31. Koivusalo M, Welch C, Hayashi H, et al. Amiloride inhibits macropinocytosis by lowering submembranous pH and preventing Rac1 and Cdc42 signaling. *J Cell Biol.* 2010;188:547-563.
32. Li L, Wan T, Wan M, Liu B, Cheng R, Zhang R. The effect of the size of fluorescent dextran on its endocytic pathway. *Cell Biol Int.* 2015;39:531-539.
33. Katase N, Lefeuvre M, Tsujigiwa H, et al. Knockdown of Dkk-3 decreases cancer cell migration and invasion independently of the Wnt pathways in oral squamous cell carcinoma derived cells. *Oncol Rep.* 2013;29:1349-1355.
34. Choi J, Kim H, Bae YK, Cheong H. REP1 modulates autophagy and macropinocytosis to enhance cancer cell survival. *Int J Mol Sci.* 2017;18:E1886.
35. Zhu BY, Shang BY, Du Y, et al. A new HDAC inhibitor cinnamoylphenazine shows antitumor activity in association with intensive macropinocytosis. *Oncotarget.* 2017;80:14748-14758.
36. Yoshida S, Pacitto R, Inoki K, Swanson J. Macropinocytosis, mTORC1 and cellular growth control. *Cell Mol Life Sci.* 2017;. <https://doi.org/10.1007/s00018-017-2710-y>.
37. Li C, MacDonald JI, Hryciw T, Meakin SO. Nerve growth factor activation of the TrkA receptor induces cell death, by macropinocytosis, in medulloblastoma Daoy cells. *J Neurochem.* 2010;112:882-899.
38. John S, Sivakumar KC, Mishra R. Bacoside A induces tumor cell death in human glioblastoma cell lines through catastrophic macropinocytosis. *Front Mol Neurosci.* 2017;10:171.
39. Sun L, Li B, Su X, et al. An ursolic acid derived small molecule triggers cancer cell death through hyperstimulation of macropinocytosis. *J Med Chem.* 2017;60:6638-6648.
40. Akue-Gedu R, Debiton E, Ferandin Y, et al. Synthesis and biological activities of aminopyrimidyl-indoles structurally related to meridianins. *Bioorg Med Chem.* 2009;17:4420-4424.
41. Kawano Y, Kypta R. Secreted antagonists of the Wnt signalling pathway. *J Cell Sci.* 2003;116:2627-2634.
42. Niehrs C. Function and biological roles of the Dickkopf family of Wnt modulators. *Oncogene.* 2006;25:7469-7481.
43. Yue W, Sun Q, Dacic S, et al. Downregulation of Dkk3 activates beta-catenin/TCF-4 signaling in lung cancer. *Carcinogenesis.* 2008;29:84-92.
44. Kobayashi K, Ouchida M, Tsuji T, et al. Reduced expression of the REIC/Dkk-3 gene by promoter-hypermethylation in human tumor cells. *Gene.* 2002;282:151-158.

45. Nara A, Aki T, Funakoshi T, et al. Hyperstimulation of macropinocytosis leads to lysosomal dysfunction during exposure to methamphetamine in SH-SY5Y cells. *Brain Res.* 2012;1466:1-14.
46. Maltese WA, Overmeyer JH. Non-apoptotic cell death associated with perturbations of macropinocytosis. *Front Physiol.* 2015;6:38.
47. Kitambi SS, Toledo EM, Usoskin D, et al. Vulnerability of glioblastoma cells to catastrophic vacuolization and death induced by a small molecule. *Cell.* 2017;170:407.

**How to cite this article:** Park N-S, Park Y-K, Ramalingam M, et al. Meridianin C inhibits the growth of YD-10B human tongue cancer cells through macropinocytosis and the down-regulation of Dickkopf-related protein-3. *J Cell Mol Med.* 2018;22:5833–5846. <https://doi.org/10.1111/jcmm.13854>