

Post-prandial Remnant Lipoprotein Metabolism in Sitosterolemia

Hayato Tada¹, Akihiro Nomura¹, Atsushi Nohara¹, Akihiro Inazu², Hiroshi Mabuchi¹, Masakazu Yamagishi¹ and Masa-aki Kawashiri¹

¹Department of Cardiovascular and Internal Medicine, Kanazawa University Graduate School of Medicine, Kanazawa, Japan

²Department of Laboratory Science, Molecular Biochemistry and Molecular Biology, Graduate School of Medical Science, Kanazawa University, Kanazawa, Japan

Aim: We aimed to clarify post-prandial accumulation of remnant-like particles (RLP) in patients with sitosterolemia.

Methods: Oral fat tolerance test cream (Jomo Shokuhin, Takasaki, Japan) 50 g was given per body surface area (m^2); blood sampling was performed at 2 h intervals up to 6 h. Plasma lipoprotein fractions and RLP fractions were determined in four sitosterolemic subjects with double mutations in ATP-binding cassette (ABC) sub-family G member 5 or member 8 (*ABCG5* or *ABCG8*) gene (mean age=18 yr, median low-density lipoprotein cholesterol [LDL-C]=154 mg/dL), six heterozygous carriers (mean age=31 yr, median LDL-C=105 mg/dL), and five subjects with heterozygous familial hypercholesterolemia (FH, mean age=32 yr, median LDL-C=221 mg/dL). The incremental area under curve (iAUC) of lipids, including LDL-C, apolipoprotein B-48 (apoB48), RLP cholesterol (RLP-C), and RLP triglyceride (RLP-TG) were evaluated.

Results: After oral fat load, there was no significant difference of the iAUC of LDL-C between sitosterolemia and heterozygous FH, whereas the iAUC of apoB48 was significantly larger in the sitosterolemic subjects compared with that of heterozygous FH (2.9 μ g/mL \times h vs. 1.3 μ g/mL \times h, $p<0.05$). Under these conditions, the iAUCs of RLP-C and RLP-TG levels were significantly larger in the sitosterolemic subject compared with those of heterozygous FH (9.5 mg/dL \times h vs. 5.7 mg/dL \times h, $p<0.05$; 149 mg/dL \times h vs. 40 mg/dL \times h, $p<0.05$, respectively), whereas those of heterozygous carriers were comparable with those with heterozygous FH.

Conclusions: Post-prandial lipoprotein metabolism in sitosterolemia appeared to be impaired, leading to their elevation in serum sterol levels. (UMIN Clinical Trials Registry number, UMIN000020330)

See editorial vol. 25: 1183-1184

Key words: Sitosterolemia, Remnant, Familial hypercholesterolemia, Remnant-like-particles, OFTT

Introduction

Familial hypercholesterolemia (FH) is a common inherited disorder of plasma lipoprotein metabolism, characterized by an elevated level of low-density lipoprotein cholesterol (LDL-C), tendon xanthomas, and premature coronary artery disease¹⁾. Monogenic causes of FH involve gene mutations such as LDL receptor, apolipoprotein B-100 (*apoB100*), and proprotein convertase subtilisin/kexin type 9 (*PCSK9*)²⁾. Post-prandial accumulation of lipoprotein remnants has been

shown to be related with elevated cardiovascular risk³⁾. Under these conditions, it has been shown that post-prandial lipoprotein metabolism is severely impaired in a dominant form of FH⁴⁾, whereas we have shown that such lipoprotein metabolism is preserved in a recessive form of FH called autosomal recessive hypercholesterolemia caused by mutations in LDL receptor adaptor protein 1 (*LDLRAP1*) gene⁵⁾. Investigating the extreme cases harboring mutations in a specific gene provides an opportunity to directly observe the role of certain molecules in lipoprotein metabolism.

Address for correspondence: Hayato Tada, Department of Cardiovascular and Internal Medicine, Kanazawa University Graduate School of Medicine, 13-1 Takaramachi, Kanazawa, 920-8641, Japan E-mail: ht240z@sa3.so-net.ne.jp

Received: March 8, 2018 Accepted for publication: May 14, 2018

Copyright©2018 Japan Atherosclerosis Society

This article is distributed under the terms of the latest version of CC BY-NC-SA defined by the Creative Commons Attribution License.

Table 1. Characteristics of the study subjects

	Sitosterolemia (n=4)	Single mutation carrier (n=6)	Heterozygous FH (n=5)
Age	18 ± 13	31 ± 7	32 ± 11
Sex (male/female)	2/2	3/3	2/3
BMI (kg/m ²)	23.1 ± 4.1	23.9 ± 6.5	24.5 ± 4.0
TC (mg/dl)	241 [226–275]	223 [189–233]	289 [284–305]
TG (mg/dl)	201 [132–241]	99 [91–138]	101 [94–146]
HDL-C (mg/dl)	50 [47–60]	57 [50–58]	49 [46–55]
LDL-C (mg/dl)	154 [150–156]	105 [89–164]	221 [220–246]
apoB48 (μg/ml)	5.7 [3.6–7.9]	2.8 [1.9–3.4]	1.4 [1.4–2.2]
Lathosterol (μg/ml)	3.0 [2.3–3.6]	1.5 [1.1–1.6]	3.8 [2.6–3.8]
Campesterol (μg/ml)	57 [49–70]	14.9 [13.9–15.5]	4.6 [3.3–4.9]
Sitosterol (μg/ml)	88.5 [73.6–113.2]	6.5 [5.1–8.4]	2.5 [1.9–2.9]
RLP-C (mg/dl)	14.4 [11.8–16.5]	3.6 [3.7–8.4]	9.1 [7.2–10.8]
RLP-TG (mg/dl)	41.5 [37.0–82.6]	14.2 [11.4–40.0]	12.7 [12.5–54.2]

FH: familial hypercholesterolemia, BMI: body mass index, TC: total cholesterol, TG: triglyceride, HDL-C: high-density lipoprotein cholesterol, LDL-C: low-density lipoprotein cholesterol, apoB48: apolipoprotein B-48, RLP-C: remnant-like particle cholesterol, RLP-TG: remnant-like particle triglyceride

Sitosterolemia (OMIM #210250) is a rare, inherited, autosomal recessive disorder of lipid metabolism characterized by increased absorption and decreased biliary excretion of plant sterols and cholesterol, resulting in prominently elevated serum concentrations of plant sterols such as sitosterol and campesterol¹⁾. This disease is caused by mutations in either of the two genes, named ATP-binding cassette (ABC) sub-family G member 5 and member 8 (*ABCG5* and *ABCG8*). Subjects suffering from sitosterolemia present primarily with tendinous and tuberous xanthomas, premature coronary atherosclerosis such as FH^{6–8)}. LDL-C levels are more variable in sitosterolemia than in other genetic hyperlipidemias, but can be extremely elevated in some patients, especially in breastfed infantile patients⁹⁾. This report indicates that patients with sitosterolemia might be vulnerable to post-prandial hyperlipidemia. Remnant-like particles (RLP) are known as a good marker to evaluate lipid metabolism after diet. Also, increased RLP levels are associated with high LDL-C, endothelium dysfunction, and coronary artery disease (CAD). Thus, this post-prandial condition may play an important role of plasma high LDL-C levels in sitosterolemic patients. In this study, we examined post-prandial lipoprotein metabolism in sitosterolemia to determine their vulnerability to oral fat load for the first time¹⁰⁾.

Materials and Methods

Study Design

This study is a single-arm, non-randomized, open-

label, uncontrolled trial. Patients with homozygous sitosterolemia with double mutations, heterozygous carriers with single mutation, and heterozygous FH are enrolled in this study. They receive an oral fat tolerance test (OFTT) cream when they met the inclusion criteria. We then compare post-prandial plasma RLP cholesterol (RLP-C) levels between the patients with sitosterolemia with double mutations in *ABCG5* or *ABCG8* gene, heterozygous mutation carriers with single mutation, and heterozygous FH. This study has been registered at the University Hospital Medical Information Network (UMIN) (UMIN ID: 000020330)¹⁰⁾.

Study Subjects

Four patients with sitosterolemia harboring double mutations in *ABCG5* or *ABCG8* gene, six relatives with a heterozygous mutation, and five unrelated subjects with heterozygous FH were enrolled in this study. None of the subjects had taken medication known to affect plasma lipids for at least 4 weeks before this study was conducted. None had smoking habits and excessive alcohol intake (>ethanol 60 g/day).

Oral Fat Tolerance Test

OFTT cream of 50 g was given per body surface area (squared meter) as described elsewhere (Jomo Shokuhin, Takasaki, Japan)¹¹⁾. The cream consisted of fat 33%, cholesterol 74 mg, and 341 kcal per 100 g and is rich in palmitic and oleic acids. Blood sampling was performed at 2 h intervals up to 6 h.




Fig. 1. Line charts of lipoproteins

Bars indicate standard errors.

Red: Sitosterolemia

Pink: Single mutation carrier

Light blue: Heterozygous FH

Lipoprotein Analysis and ApoE Phenotype

LDL-C values were measured directly using QUALIGENT® (Sekisui Medical, Tokyo, Japan). RLP is estimated as the unbound fraction of plasma after incubation with immunoaffinity gel of apoB100 monoclonal antibody and apolipoprotein A-I monoclonal antibody as described¹²⁾. ApoE phenotype was separated by isoelectric focusing and detected by western blot with apoE polyclonal antibody (phenotyping apoE IEF system, JOKOH, Tokyo, Japan). Apolipoprotein B-48 (apoB48) levels were determined by ELISA method using monoclonal antibody against C-terminal decapeptide of apoB48¹³⁾. Serum levels of sterols, including sitosterol, lathosterol, and campesterol, were determined using gas–liquid chromatography–mass spectrometry as previously described⁹⁾.

Statistical Analysis

Values are expressed as medians (interquartile range [IQR]) unless otherwise stated. Area under curve (AUC) for triglyceride (TG), LDL-C, apoB48, sitosterol, RLP-C, and RLP triglyceride (RLP-TG) at baseline and after fat load were calculated using the trapezoid rule. The differences in the incremental AUC (iAUC) of the plasma variables between the three groups were examined using Kruskal–Wallis test. Post hoc analyses using Scheffé's method were performed when the main effect was significant. All tests of statistical significance were assumed at a level of $p < 0.05$.

Ethical Considerations

This study was approved by the Ethics Committee of Kanazawa University and carried out in accordance with the Declaration of Helsinki (2008) of the




Fig. 2. Line charts of RLP fractions

Bars indicate standard errors.

Red: Sitosterolemia

Pink: Single mutation carrier

Light blue: Heterozygous FH

World Medical Association. All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Declaration of Helsinki of 1975, as revised in 2008. Informed consents were obtained from all subjects for being included in the study.

Results

The baseline characteristics of the three study groups are shown in **Table 1**. Genetic backgrounds of the study subjects are shown in **Supplemental Table 1**. These groups of patients were comparable in terms of age, gender, and body mass index (BMI). Interestingly, the baseline LDL-C level of sitosterolemic patients with double mutations was significantly higher than that of single mutation, whereas baseline LDL-C level of heterozygous FH was significantly higher than that in sitosterolemic subjects. On the other hand, serum levels of sitosterol and campesterol in sitosterolemic patients with double mutations were significantly higher than those with a single mutation as well as those in heterozygous FH. In addition, apoB48, TG, RLP-C, and RLP-TG levels were significantly higher in sitosterolemic subjects than those in heterozygous FH (**Table 1**).

After oral fat load, lipoproteins, other than LDL-C, increased through 2 to 4 h (**Figs. 1** and **2**). There was no significant difference in the iAUC of LDL-C

between sitosterolemia and heterozygous FH, whereas the iAUC of TG was significantly larger in sitosterolemic subjects than that in heterozygous FH ($154 \text{ mg/dL} \times \text{h}$ vs. $50 \text{ mg/dL} \times \text{h}$, $p < 0.05$, **Fig. 3**).

Interestingly, the iAUC of apoB48 was significantly larger in the sitosterolemic subjects compared with that of heterozygous FH ($2.9 \mu\text{g/mL} \times \text{h}$ vs. $1.3 \mu\text{g/mL} \times \text{h}$, $p < 0.05$, **Fig. 3**). In addition to those results, the iAUC of sitosterol in the sitosterolemic subjects was significantly larger than that of heterozygous FH as expected ($66 \mu\text{g/mL} \times \text{h}$ vs. $-0.5 \mu\text{g/mL} \times \text{h}$, $p < 0.05$). Under these conditions, the iAUCs of RLP-C and RLP-TG levels were significantly larger in the sitosterolemic subjects compared with those of heterozygous FH ($9.5 \text{ mg/dL} \times \text{h}$ vs. $5.7 \text{ mg/dL} \times \text{h}$, $p < 0.05$; $149 \text{ mg/dL} \times \text{h}$ vs. $40 \text{ mg/dL} \times \text{h}$, $p < 0.05$, respectively, **Fig. 4**), whereas those of single mutation carrier in *ABCG5* or *ABCG8* gene were comparable with those with heterozygous FH.

Discussion

The main finding of the present study is that the clearance of post-prandial RLP fraction was impaired in sitosterolemia compared with heterozygous FH. This is the first study to demonstrate the impaired post-prandial RLP metabolism in sitosterolemia.

Impaired TG-rich lipoprotein metabolism under the condition of disturbance of *ABCG5/ABCG8* have




Fig. 3. The iAUC of lipoproteins

Boxplots illustrating the iAUC of (A) TG, (B) LDL-C, (C) apoB48, and (D) sitosterol in three groups.
 Red: Sitosterolemia
 Pink: Single mutation carrier
 Light blue: Heterozygous FH

been implicated by the observational and interventional studies of sitosterolemic subjects^{14, 15)} as well as by the experimental study using *ABCG5/ABCG8* knockout mice (sitosterolemic mice)¹⁶⁾. Previously, we demonstrated that breastfed infantile cases with sitosterolemia harboring double mutations in *ABCG5* gene exhibit transient extreme hyper-LDL cholesterolemia⁹⁾. In addition, there have been great diversities in the LDL-C levels among the subjects with sitosterolemia described so far¹⁷⁾. Such observations as well as the facts that *ABCG5/ABCG8* are playing an important role in excretion of sterols in the intestine could lead us to investigate if the patients with sitosterolemia harboring *ABCG5/ABCG8* mutations are vulnerable to diet-induced hyper-

lipidemia. We confirmed that post-prandial RLP metabolism was disturbed in the patients with sitosterolemia for the first time. Moreover, apoB48 level, which has been shown to be associated with atherosclerotic cardiovascular diseases¹⁸⁾, was significantly elevated in sitosterolemic patients both in fasting state, as well as in post-prandial state. In addition, we observed that the peaks of the post-prandial lipoproteins were earlier in sitosterolemic patients than those in heterozygous FH. Moreover, sitosterol level in sitosterolemic patients increased after fat load, although such trends were not observed in heterozygous FH nor in single mutation carriers. Thus, dietary counseling for the patients with sitosterolemia should be one of the reasonable approaches




Fig. 4. The iAUC of RLP fractions

Boxplots illustrating the iAUC of (A) RLP-C and (B) RLP-TG in three groups.

Red: Sitosterolemia

Pink: Single mutation carrier

Light blue: Heterozygous FH

for their risk management.

In this study, we compared post-prandial lipoprotein metabolism in sitosterolemia with that in heterozygous FH, not with normal controls. It is well known that the patients with sitosterolemia exhibit dyslipidemia, including elevation of LDL-C. And RLP-C has been shown to be correlated with LDL-C, especially in cases with elevated LDL-C¹⁹⁾. Accordingly, it is rather fair to compare sitosterolemia and heterozygous FH under the condition of elevated LDL-C in both sides.

This study has several limitations. First, only a small number of subjects were included in this study. It is difficult to enroll more sitosterolemic patients because of the rarity of this disorder. However, our subjects were comparatively uniform in terms of factors potentially affecting the post-prandial lipoprotein metabolism—age, sex, and BMI—among study subjects. Second, we used OFTT cream without plant sterols making it difficult to see the effect on such sterols in this study. However, loading sitosterol on sitosterolemia are considered ethically unacceptable, although the increased plant sterols themselves may not relate with atherosclerosis^{20, 21)}. Third, TG level was significantly higher in sitosterolemia than in other two groups. In this regard, it has been shown that fasting TG level was significantly associated with post-prandial lipemia²²⁾. Accordingly, our results may not be reflected

by actual “post-prandial” lipemia itself. Despite those limitations, we believe that this study provides new insights into the roles of ABCG5/ABCG8 in the post-prandial lipoprotein metabolism.

Acknowledgements and Notice of Grant Support

We express our special thanks to Kazuko Honda and Sachio Yamamoto (staff of Kanazawa University) for their outstanding technical assistance. We also express our special thanks to Drs. Takuwa Nakahashi, Yoshihiro Tanaka, Taro Ichise, Takashi Kobayashi, Azusa Ohbatake, and Akari Wada for their assistance in data collection. We have received the research grants from Sakakibara Memorial Research Grant from the Japan Research Promotion Society for Cardiovascular Diseases and Astellas Foundation for Research on Metabolic Disorders.

COI

Hayato Tada has received a research grant from Sanofi K.K. Atsushi Nohara and Hiroshi Mabuchi have received research grants from MSD K.K., Sanofi K.K., Shionogi & Co., Ltd., Kowa Co., Ltd., Astellas Pharma Inc., AstraZeneca K.K., Keiai-Kai Medical Corp., and

Biopharm of Japan Co. Masakazu Yamagishi has received research grants from MSD K.K., Astellas Pharma Inc., Daiichi-Sankyo Co., Ltd., and Otsuka Pharmaceutical Co., Ltd., and he has received payments for lectures from Astellas Pharma Inc., Daiichi-Sankyo Co., Ltd., Shionogi & Co., Ltd., and Kowa Co., Ltd. Masa-aki Kawashiri has received payments for lectures from Amgen Astellas Biopharma K.K. and Astellas Pharma Inc.

References

- 1) Goldstein JL, Hobbs HH, Brown MS. Familial hypercholesterolemia. In The Metabolic and Molecular Bases of Inherited Diseases. 8th edition. Scriver CR, Beaudet AL, Sly WS, Valle D, editors. McGraw-Hill, New York. 2001; 2863-2913
- 2) Soutar AK, Naoumova RP. Mechanisms of Disease: genetic causes of familial hypercholesterolemia. *Nat Pract Cardiovasc Med.* 2007; 4: 214-225
- 3) Karpe F, Steiner G, Uffelmann K, Olivecrona T, Hamsten A: Postprandial lipoproteins and progression of coronary atherosclerosis. *Atherosclerosis.* 1994; 106: 83-97
- 4) de Sauvage Nolting PR, Twickler MB, Dallinga-Thie GM, Buurma RJ, Hutten BA, Kastelein JJ; Examination of Probands and Relatives in Statin Studies with Familial Hypercholesterolemia (ExPRESS) Study Group. Elevated remnant-like particles in heterozygous familial hypercholesterolemia and response to statin therapy. *Circulation.* 2002; 106: 788-792
- 5) Tada H, Kawashiri MA, Tanaka A, Nakano T, Nakajima K, Inoue T, Noguchi T, Nakanishi C, Konno T, Hayashi K, Nohara A, Inazu A, Kobayashi J, Mabuchi H, Yamagishi M. Post-prandial remnant lipoprotein metabolism in autosomal recessive hypercholesterolemia. *Eur J Clin Invest.* 2012; 42: 1094-1099
- 6) Tada H, Kawashiri MA, Okada H, Endo S, Toyoshima Y, Konno T, Nohara A, Inazu A, Takao A, Mabuchi H, Yamagishi M, Hayashi K. A Rare Coincidence of Sitosterolemia and Familial Mediterranean Fever Identified by Whole Exome Sequencing. *J Atheroscler Thromb.* 2016; 23: 884-890
- 7) Sakuma N, Tada H, Mabuchi H, Hibino T, Kasuga H. Lipoprotein Apheresis for Sitosterolemia. *Ann Intern Med.* 2017; 167: 896-899
- 8) Kawamura R, Saiki H, Tada H, Hata A. Acute myocardial infarction in a 25-year-old woman with sitosterolemia. *J Clin Lipidol.* 2018; 12: 246-249
- 9) Tada H, Kawashiri MA, Takata M, Matsunami K, Imamura A, Matsuyama M, Sawada H, Nunoi H, Konno T, Hayashi K, Nohara A, Inazu A, Kobayashi J, Mabuchi H, Yamagishi M. Infantile Cases of Sitosterolemia with Novel Mutations in the ABCG5 Gene: Extreme Hypercholesterolemia is Exacerbated by Breastfeeding. *JIMD Rep.* 2015; 21: 115-122
- 10) Nomura A, Tada H, Nohara A, Kawashiri MA, Yamagishi M. Oral Fat Tolerance Test for Sitosterolemia and Familial Hypercholesterolemia: A Study Protocol. *J Atheroscler Thromb.* 2018; 25: 741-746
- 11) Inazu A, Nakajima K, Nakano T, Niimi M, Kawashiri MA, Nohara A, Kobayashi J, Mabuchi H. Decreased post-prandial triglyceride response and diminished remnant lipoprotein formation in cholestrylo ester transfer protein (CETP) deficiency. *Atherosclerosis.* 2008; 196: 953-957
- 12) Nakajima K, Saito T, Tamura A, Suzuki M, Nakano T, Adachi M, Tanaka A, Tada N, Nakamura H, Campos E, Havel RJ. Cholesterol in remnant-like lipoproteins in human serum using monoclonal anti apo B-100 and anti apo A-I immunoaffinity mixed gels. *Clin Chim Acta.* 1993; 223: 53-71
- 13) Kinoshita M, Kojima M, Matsushima T, Teramoto T. Determination of apolipoprotein B-48 in serum by a sandwich ELISA. *Clin Chim Acta.* 2005; 351: 115-120
- 14) Salen G, von Bergmann K, Lütjohann D, Kwiterovich P, Kane J, Patel SB, Musliner T, Stein P, Musser B; Multi-center Sitosterolemia Study Group. Ezetimibe effectively reduces plasma plant sterols in patients with sitosterolemia. *Circulation.* 2004; 109: 966-971
- 15) Hidaka H, Nakamura T, Aoki T, Kojima H, Nakajima Y, Kosugi K, Hatanaka I, Harada M, Kobayashi M, Tamura A, Fujii Tatsuzo, Shigeta Yukio. Increased plasma plant sterol levels in heterozygotes with sitosterolemia and xanthomatosis. *J Lipid Res.* 1990; 31: 881-888
- 16) Wang J, Joy T, Mymin D, Frohlich J, Hegele RA. Phenotypic heterogeneity of sitosterolemia. *J Lipid Res.* 2004; 45: 2361-2367
- 17) Méndez-González J, Julve J, Rotllan N, Llaverias G, Blanco-Vaca F, Escolà-Gil JC. ATP-binding cassette G5/G8 deficiency causes hypertriglyceridemia by affecting multiple metabolic pathways. *Biochim Biophys Acta.* 2011; 1811: 1186-1193
- 18) Masuda D, Yamashita S. Postprandial Hyperlipidemia and Remnant Lipoproteins. *J Atheroscler Thromb.* 2017; 24: 95-109
- 19) Nakajima K, Nakajima Y, Takeichi S, Fujita MQ. Plasma remnant-like lipoprotein particles or LDL-C as major pathologic factors in sudden cardiac death cases. *Atherosclerosis.* 2008; 198: 237-246
- 20) Gylling H, Plat J, Turley S, Ginsberg HN, Ellegård L, Jessup W, Jones PJ, Lütjohann D, Maerz W, Masana L, Silbernagel G, Staels B, Borén J, Catapano AL, De Backer G, Deanfield J, Descamps OS, Kovanen PT, Riccardi G, Tokgozoglu L, Chapman MJ; European Atherosclerosis Society Consensus Panel on Phytosterols. Plant sterols and plant stanols in the management of dyslipidaemia and prevention of cardiovascular disease. *Atherosclerosis.* 2014; 232: 346-360
- 21) Silbernagel G, Chapman MJ, Genser B, Kleber ME, Fauler G, Scharnagl H, Grammer TB, Boehm BO, Mäkelä KM, Kähönen M, Carmena R, Rietzschel ER, Bruckert E, Deanfield JE, Miettinen TA, Raitakari OT, Lehtimäki T, März W. High intestinal cholesterol absorption is associated with cardiovascular disease and risk alleles in ABCG8 and ABO: evidence from the LURIC and YFS cohorts and from a meta-analysis. *J Am Coll Cardiol.* 2013; 62: 291-299
- 22) Masuda D, Sakai N, Sugimoto T, Kitazume-Taneike R, Yamashita T, Kawase R, Nakaoka H, Inagaki M, Nakatani K, Yuasa-Kawase M, Tsubakio-Yamamoto K, Ohama T, Nakagawa-Toyama Y, Nishida M, Ishigami M, Masuda Y, Matsuyama A, Komuro I, Yamashita S. Fasting serum apolipoprotein B-48 can be a marker of postprandial hyperlipidemia. *J Atheroscler Thromb.* 2011; 18: 1062-1070

Supplemental Table 1. Genetic backgrounds of the subjects

Sitosterolemia (n=4)	Heterozygous carrier (n=6)	Heterozygous FH (n=5)
c.1256G>A/c.1763-1G>A (ABCG5)	c.1256G>A (ABCG5)	c.1702C>G (LDLR)
c.454C>T/c.1403_1404delTC (ABCG8)	c.1763-1G>A (ABCG5)	c.1845+2T>C (LDLR)
c.1306G>A/c.1813_1817delCTTTT (ABCG5)	c.1306G>A (ABCG5)	c.2054C>T (LDLR)
c.130T>G/c.1306G>A (ABCG5)	c.1813_1817delCTTTT (ABCG5)	c.2431A>T (LDLR)
	c.1306G>A (ABCG5)	c.94G>A (PCSK9)
	c.130T>G (ABCG5)	

FH: familial hypercholesterolemia, ABCG5: ATP-binding cassette (ABC) sub-family G member 5, ABCG8: ATP-binding cassette (ABC) sub-family G member 8, LDLR: low-density lipoprotein receptor, PCSK9: proprotein convertase subtilisin/kexin type 9