- of folate, vitamin B12, and the 5,10-methylenetetrahydrofolate reductase 677C→T variant. Am J Obstet Gynecol 2002;186:499–503.
- Morkbak AL, Hvas AM, Milman N, Nexo E. Holotranscobalamin remains unchanged during pregnancy: longitudinal changes of cobalamins and their binding proteins during pregnancy and postpartum. Haematologica 2007;92:1711–2.
- Pardo J, Peled Y, Bar J, Hod M, Sela BA, Rafael ZB, Orvieto R. Evaluation of low serum vitamin B(12) in the non-anaemic pregnant patient. Hum Reprod 2000;15:224–6.
- Metz J, McGrath K, Bennett M, Hyland K, Bottiglieri T. Biochemical indices of vitamin B12 nutrition in pregnant patients with subnormal serum vitamin B12 levels. Am J Hematol 1995;48:251–5.
- Murphy MM, Scott JM, Arija V, Molloy AM, Fernandez-Ballart JD. Maternal homocysteine before conception and throughout pregnancy predicts fetal homocysteine and birth weight. Clin Chem 2004;50:1406–12.
- Molloy AM, Mills JL, Cox C, Daly SF, Conley M, Brody LC, Kirke PN, Scott JM, Ueland PM. Choline and homocysteine interrelations in umbilical cord and maternal plasma at delivery. Am J Clin Nutr 2005;82: 836–42.

doi:	10.3945/a	icn 113	073593

Erratum

Tate DF, Turner-McGrievy G, Lyons E, Stevens J, Erickson K, Polzien K, Diamond M, Wang X, Popkin B. Replacing caloric beverages with water or diet beverages for weight loss in adults: main results of the Choose Healthy Options Consciously Everyday (CHOICE) randomized clinical trial. Am J Clin Nutr 2012;95:555–63.

In the print version of the article, the second sentence of the Results section of the abstract contains a copyediting error. Negative signs should be included in the DB and Water percent results, so that the sentence reads as follows: "Mean (\pm SEM) weight losses at 6 mo were $-2.5 \pm 0.45\%$ in the DB group, $-2.03 \pm 0.40\%$ in the Water group, and $-1.76 \pm 0.35\%$ in the AC group; there were no significant differences between groups." The online version of this article was corrected at final publication.

doi: 10.3945/ajcn.113.075432.