

Article

# Extraction, Preconcentration and Isolation of Flavonoids from *Apocynum venetum* L. Leaves Using Ionic Liquid-Based Ultrasonic-Assisted Extraction Coupled with an Aqueous Biphasic System

Zhijian Tan \*, Yongjian Yi, Hongying Wang, Wanlai Zhou and Chaoyun Wang \*

Institute of Bast Fiber Crops, Chinese Academy of Agricultural Sciences, Changsha 410205, China; ibfcyyj@163.com (Y.Y.); Cswwhy328@126.com (H.W.); aruofly@126.com (W.Z.)

\* Correspondence: tanzhijian@caas.cn (Z.T.); ibfcwcy@139.com (C.W.);

Tel.: +86-731-8899-8517 (Z.T.); +86-731-8899-8501 (C.W.); Fax: +86-731-8899-8501 (C.W.)

Academic Editor: Derek J. McPhee

Received: 23 December 2015 ; Accepted: 19 February 2016 ; Published: 4 March 2016

**Abstract:** *Background:* Ionic liquids (ILs) are considered as green solvents, and widely applied for the extraction of various compounds. *Methods:* The present research focuses on the extraction of flavonoids from *Apocynum venetum* L. leaves by ultrasound-assisted extraction (UAE). Several major influencing factors were optimized. Then, an aqueous biphasic system (ABS) was applied for further isolation of flavonoids. *Results:* The flavonoids were mainly distributed in the top phase, while impurities were extracted to the bottom phase. The parameters influencing the extraction, namely type and concentration of salt, temperature, and pH, were studied in detail. Under optimized conditions (72.43% IL extract, 28.57% (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 25 °C temperature, pH 4.5), the preconcentration factor and extraction efficiency were found to be 3.78% and 93.35%, respectively. *Conclusions:* This simple and efficient methodology is expected to see great use in the extraction and isolation of pharmaceutically active components from medicinal plant resources.

**Keywords:** ionic liquids; ultrasonic-assisted extraction; flavonoids; aqueous biphasic system; isolation


## 1. Introduction

*Apocynum venetum* L., belonging to the family Apocynaceae, is a traditional medicinal plant distributed throughout mid- and north-western China, mainly in the Xinjiang Uygur Autonomous Region. *A. venetum* has several pharmacological effects such as lowering blood pressure, antidepressant, anti-anxiety, antinephritis and antineurasthenia [1–4]. Tea prepared from *A. venetum* leaves has gained popularity as a healthy beverage that acts as an anti-aging nutritional supplement [5]. The flavonoids present as the major active constituents in *A. venetum* leaf are responsible for the pharmacological effect of lowering blood pressure, scavenging free radicals and anti-inflammatory action [6–8].

Room temperature ionic liquids (ILs) are molten salts with a melting point below 100 °C. Usually, they are composed of organic cations and inorganic anions. As opposed to water and conventional volatile organic solvents, ILs have unique features of high ionic conductivity, wide electrochemical windows, negligible volatility, and high polarity and chemical stability [9–12]. In addition, some other assisted technologies coupled with ILs, such as ionic liquid-based ultrasonic-assisted extraction (IL-UAE) [13], ionic liquid-based microwave-assisted extraction (IL-MAE) [14], and ionic liquid-based negative pressure cavitation-assisted extraction (IL-NPCE) [15], provide remarkable advantages in the extraction and separation field. In particular, IL-UAE is an innovative technology combining IL and

ultrasonic extraction, which can produce clear synergies in some cases [16]. In the field of extraction of medicinal components from different herb sources, IL-UAE has been reported to be a comparatively more effective method for the extraction of various compounds, such as aesculin and aesculetin from *Cortex fraxini* [17], piperine from white pepper [18], ginsenosides from ginseng root [19], alkaloids from *Catharanthus roseus* [20], biphenyl cyclooctene lignans from *Schisandra chinensis* Baill [21], magnolol and honokiol from *Magnoliae officinalis* cortex [22]. Aqueous biphasic systems based on ionic liquids (IL-ABS) were proposed in 2003 by Rogers [23], and have been developed to extract and isolate several bioactive compounds, such as anthraquinones [24], gallic acid [25], puerarin [26], and alkaloids [27].

The analysis and identification of the flavonoids in *A. venetum* leaves using liquid chromatography-tandem mass spectrometry (LC-MS/MS) are well reported in the literature [28–30]. The present article reports the extraction of flavonoids from *A. venetum* into IL aqueous solution by the IL-UAE method, with further application of IL-ABS for preconcentration and purification of the flavonoids. Some factors affecting UAE were optimized by response surface methodology (RSM) to obtain the optimal extraction yields. The influencing parameters in IL-ABS were also investigated to obtain the maximal extraction efficiencies. The process flowchart for extraction and purification of flavonoids from *A. venetum* L. using IL-UAE coupled with IL-ABS and back-extraction is shown in Figure 1.




**Figure 1.** Process flowchart for the extraction and purification of flavonoids from *Apocynum venetum* L. using IL-UAE coupled with IL-ABS and back-extraction.

## 2. Results and Discussion

### 2.1. Selection of Ionic Liquid

Commonly used ILs such as  $[C_n\text{mim}]\text{Br}$  ( $n = 2, 4,$  and  $6$ ),  $[C_4\text{mim}][\text{BF}_4]$ , and  $[C_4\text{mim}][\text{N}(\text{CN})_2]$  were considered as the extractants for the present study. From Figure 2, it can be observed that both  $[C_4\text{mim}][\text{BF}_4]$  and  $[C_4\text{mim}][\text{N}(\text{CN})_2]$  exhibited higher extraction yields than other ILs used in this IL-UAE study. The probable reason for this is that the extraction ability is related to the hydrophobicity of ILs, usually increasing with their increasing hydrophobicity. The order of hydrophobicity for the five studied ILs is as follows:  $[C_2\text{mim}]\text{Br} < [C_4\text{mim}]\text{Br} < [C_6\text{mim}]\text{Br} < [C_4\text{mim}][\text{N}(\text{CN})_2] < [C_4\text{mim}][\text{BF}_4]$ . In addition, it was reported that the factors influencing the extraction ability are very complicated, hydrogen-bonding ability can be the main factor influencing the extraction considering the anions [31],  $\pi$ - $\pi$  and  $n$ - $\pi$  interactions also should not be omitted in view of imidazole ring-containing cations [32]. Although  $[C_4\text{mim}]\text{BF}_4$  was commonly used as the extractant in IL-UAE [18,33–35], some research groups had reported that tetrafluoroborate-based ILs were not water-stable as they could undergo hydrolysis in aqueous solution [36,37]. Thus,  $[C_4\text{mim}][\text{N}(\text{CN})_2]$  was selected for further study. In general,  $[C_4\text{mim}][\text{N}(\text{CN})_2]$  has comparatively lower viscosity than the other ILs used in the present study, about  $29.3 \text{ mPa}\cdot\text{s}$  at  $298.15 \text{ K}$  and  $0.101 \text{ MPa}$  [38].




**Figure 2.** UAE of total flavonoids from *A. venetum* L. leaves using different ILs.

## 2.2. Optimization of UAE

### 2.2.1. Single Factor Analysis

The factors such as IL concentration, liquid/solid ratio, ultrasonic time, extraction temperature, and pH can influence the extraction, hence, we had to optimize these conditions in order to obtain the maximal flavonoids extraction yield. The effect of aqueous solutions with different [C<sub>4</sub>mim][N(CN)<sub>2</sub>] concentrations (from 0.1 to 0.5 g/mL) was studied to find the optimal concentration for IL-UAE. As shown in Figure 3a, the maximal extraction yield was obtained at 0.2 g/mL IL concentration. Thus, the concentration of 0.2 g/mL was selected considering to obtain the maximum extraction yield and reduce operational cost.



**Figure 3.** The effect of (a) IL concentration (the extraction conditions were 20:1 liquid/solid ratio, 30 min ultrasonic time at room temperature with pH being not adjusted); (b) liquid/solid ratio (the extraction conditions were 0.2 g/mL IL concentration, 30 min ultrasonic time at room temperature with pH being not adjusted); (c) ultrasonic time (the extraction conditions were 0.2 g/mL IL concentration, 20:1 liquid/solid ratio at room temperature with pH being not adjusted); (d) extraction temperature (the extraction conditions were 0.2 g/mL IL concentration, 20:1 liquid/solid ratio, 30 min ultrasonic time with pH being not adjusted); and (e) pH value (the extraction conditions were 0.2 g/mL IL concentration, 30 min ultrasonic time, 20:1 liquid/solid ratio at room temperature) on the extraction yields by UAE.

Liquid/solid ratio (volume of extraction solvent/amount of dried *A. venetum* L. leaves) is another factor affecting the extraction. Employing a higher liquid/solid ratio may result in excess consumption of ILs, while a lower ratio may lead to incomplete extraction of flavonoids; hence, it is important to optimize the liquid/solid ratio. For this purpose, the influence of liquid/solid ratios of 10:1–30:1 mL/g on the extraction yield was investigated. The results in Figure 3b revealed that the maximal extraction yield was obtained at the liquid/solid ratio of 20:1. Hence, this value was selected for further study.

Ultrasound exposure time can also affect the UAE to some extent; hence, its optimization is essential to ensure complete extraction. The effect of ultrasonic irradiation times ranging from 10 to 30 min was studied. From Figure 3c, it can be observed that 20 min was sufficient for the extraction of flavonoids.

The effect of extraction temperature in the range of 25–65 °C was investigated. As shown in Figure 3d, the extraction yields varied little at temperatures below 55 °C, thus, the extraction was performed at room temperature.

Na<sub>2</sub>HPO<sub>4</sub>-H<sub>3</sub>PO<sub>4</sub> buffer solution coupled with hydrochloric acid and sodium hydroxide solutions were applied to adjust the pH. The effect of pH in the range of 1.0–9.0 was studied. The results in Figure 3e indicated that the extraction yields varied little at pHs below pH 7.0. Thus, the extraction was performed under acidic conditions.

### 2.2.2. Response Surface Methodology

To study the mutual influence among these factors, a Design-Expert (DE) 7.0 software (Statease Inc., Minneapolis, MN, USA) was applied to optimize the experimental conditions. Three factors with three levels—IL concentration (0.2, 0.3 and 0.4 g/mL), liquid/solid ratio (15:1, 20:1 and 25:1), and ultrasonic time (15, 20 and 25 min)—were developed on the basis of Box-Behnken experimental design. The experimental conditions of different runs and the results for extraction yield were shown in Table 1. The experimental data were analyzed by multiple regressions to fit the second order regression equation. The regression model in the form of marked factors as an equation of independent variables was obtained:  $Y = 8.81 - 0.73A + 0.21B - 0.24C - 0.16AB + 0.26AC + 0.067BC - 1.10A^2 - 0.33B^2 - 0.73C^2$  ( $R^2 = 0.9947$ ),  $Y$  represented the extraction yield, the marked factors A, B, and C represented IL concentration, liquid/solid ratio, and ultrasonic time, respectively.

**Table 1.** Experimental conditions for extraction yield using a three-factors/three-level Box-Behnken design.

Run	Factor A: Concentration of IL (g/mL)	Factor B: Liquid/Solid Ratio	Factor C: Ultrasonic Time (min)	Response Average Extraction Yield (mg/g)
1	0.20	15:1	20	7.84
2	0.40	15:1	20	6.59
3	0.20	25:1	20	8.50
4	0.40	25:1	20	6.61
5	0.20	20:1	15	8.10
6	0.40	20:1	15	6.24
7	0.20	20:1	25	7.21
8	0.40	20:1	25	6.39
9	0.30	15:1	15	7.86
10	0.30	25:1	15	8.23
11	0.30	15:1	25	7.15
12	0.30	25:1	25	7.79
13	0.30	20:1	20	8.77
14	0.30	20:1	20	8.79
15	0.30	20:1	20	8.89
16	0.30	20:1	20	8.76
17	0.30	20:1	20	8.85


Table 2 shows that the coefficient ( $R^2$ ) obtained from the calculated model was 0.9947, which implied that over 99.47% of the variations could be explained by this model, and the value of “Lack of fit” was not significant ( $p > 0.05$ ). The Model  $F$ -value was 147.00 and the  $p$ -value was  $<0.0001$  implied the fact that this model was significant, there was 0.01% probability for “Model  $F$ -value” occurring owing to noise. All of the results above demonstrated that the model was acceptable. The value of “Prob  $> F$ ” was less than 0.05, indicating the significance of the model terms. On this occasion, the

effects of linear terms A, B, and C, all quadratic terms, and two interaction terms (AB and AC) were significant model.


**Table 2.** Analysis of variance for the response surface quadratic regression model.

Source	Degrees of Freedom	Sum of Squares	Mean Square	F-Value	p-Value Prob > F
<b>Model</b>	<b>9</b>	<b>13.90</b>	<b>1.54</b>	<b>147.00</b>	<b>&lt;0.0001</b>
A	1	4.23	4.23	402.94	<0.0001
B	1	0.36	0.36	53.98	0.0006
C	1	0.45	0.45	42.49	0.0003
AB	1	0.1	0.1	9.75	0.0168
AC	1	0.27	0.27	25.73	0.0014
BC	1	0.018	0.018	1.73	0.2293
A <sup>2</sup>	1	5.09	5.09	484.63	<0.0001
B <sup>2</sup>	1	0.45	0.45	42.91	0.0003
C <sup>2</sup>	1	2.23	2.23	211.93	<0.0001
Residual	7	0.074	0.011		
Lack of fit	3	0.061	0.02	6.53	0.0508
Pure Error	4	0.012	$3.12 \times 10^{-3}$		
Cor total	16	13.98			

Figure 4 showed the 3D response surfaces for the mutual interaction between IL concentration, liquid/solid ratio, and ultrasonic time. The optimum conditions proposed by DE software were as follows: IL concentration was 0.26 g/mL; liquid/solid ratio was 21.98:1; and ultrasonic time was 18.93 min. Under the practical conditions for use these values (0.26 g/mL IL concentration, 22:1 liquid/solid ratio; and 19 min ultrasonic irradiation time) in routine applications, the experiments were repeated 3 times. The realistic experimental value (8.92 mg/g) was satisfactorily close to the predicted value (9.02 mg/g). The results of verification experiments proved that this model can well reflect the expected optimization.



**Figure 4.** Cont.



**Figure 4.** The plots for 3D response surface revealed the effect of (a) IL concentration and liquid/solid ratio; (b) IL concentration and ultrasonic time; (c) liquid/solid ratio and ultrasonic time on the extraction yields of total flavonoids.

### 2.3. Comparison of Different Extraction Methods

IL-UAE was compared to other methods used for the extraction of flavonoids, such as heat reflux extraction using ILs  $[C_4mim][N(CN)_2]$  (IL-HRE), Soxhlet extraction using ethanol (Ethanol-SE), and UAE using ethanol (Ethanol-UAE). The liquid/solid ratios for all extraction methods are 30:1. It can be seen in Table 3 that the IL-UAE method gave the highest extraction yield in a shorter time and under milder temperature conditions. Therefore, the replacement of traditional volatile organic solvents by ionic liquid is feasible, moreover, the UAE is an effective auxiliary method in solvent extraction.

**Table 3.** Comparison of IL-UAE to conventional extraction methods.


Methods	Temperature ( $^{\circ}C$ )	Time (h)	Yield (mg/g)	RSD (%)
IL-HRE(40%(w/w) IL)	100	6	7.31	3.5
Ethanol-SE(100%(w/w) ethanol)	80	6	6.45	2.8
Ethanol-UAE(40%(w/w) ethanol)	30	0.5	7.65	2.4
Ethanol-UAE(70%(w/w) ethanol)	30	0.5	7.88	1.8
Ethanol-UAE(100%(w/w) ethanol)	30	0.5	7.14	2.6
IL-UAE(40%(w/w) IL)	30	0.5	8.67	1.7

### 2.4. Extraction and Purification of Flavonoids by IL-ABS

For further extraction and purification of flavonoids, ABS was formed by adding different salts to the IL extract. To obtain the optimum extraction conditions of IL-ABS extraction, the major factors influencing the extraction, such as salt type and concentration, extraction temperature, and system pH were studied. Three types of salts— $NaH_2PO_4$ ,  $(NH_4)_2SO_4$ , and  $K_3PO_4$ —were chosen as the components to form ABS. The extraction efficiencies of ABSs formed by these salts are shown in Figure 5a. It can be seen that the system formed by  $(NH_4)_2SO_4$  had the best extraction ability among the three cases. The optimum extraction efficiency was obtained when the system was formed with 3.0 g IL extract (72.43%, w/w) and 1.2 g  $(NH_4)_2SO_4$  (28.57%, w/w). As the obtained flavonoids were acidic, 28.57%  $(NH_4)_2SO_4$  was chosen as the optimal salt concentration to ensure stability under acidic conditions.

The partitioning of flavonoids in  $[C_4mim][N(CN)_2]/(NH_4)_2SO_4$  ABS was investigated in the 25–45  $^{\circ}C$  temperature range and at a constant pH value. From Figure 5b, it can be seen that the extraction efficiencies decreased as the temperature increased. The maximum extraction efficiency of 93.22% was obtained at 25  $^{\circ}C$ . Thus, the operation can be conducted at room temperature without heating, which benefits both the cost reduction of the process and the stability of the flavonoids.


To study the effect of pH, pH values of 1.5–9.0 were developed in  $[\text{C}_4\text{mim}][\text{N}(\text{CN})_2]/(\text{NH}_4)_2\text{SO}_4$  ABS. As can be seen from Figure 5c, a notable extraction efficiency was obtained under acidic conditions with a maximum of 93.35% at pH 4.5, but no obvious difference was observed at pH 1.5–6.0. As the IL/ $(\text{NH}_4)_2\text{SO}_4$ ABS has an optimum pH of 2.95, the system pH needed not be adjusted. Under the optimized conditions, a preconcentration factor of 3.78 for total flavonoids was obtained.



**Figure 5.** Effect of (a) salt type and concentration (each IL-ABS contains a given mass of IL extract and salt at room temperature); (b) extraction temperature; (c) system pH on the extraction efficiencies.

### 2.5. Isolation of Flavonoids and Recovery of IL

The results (Figure 6) for back-extraction showed that the maximum back-extraction efficiency (83.5% for total flavonoids) was obtained using *n*-butanol; dichloromethane and chloroform had no back-extraction ability, diethyl ether, petroleum ether and *n*-hexane had much lower back-extraction efficiencies. The organic alcohols thus had higher back-extraction efficiencies than other solvents; this is because alcohols have stronger hydrogen bonding interactions with flavonoids and water molecules, resulting in the flavonoids and water migrating to the organic layer from the IL layer. The IL can be reused after removal of water. The recovered IL was used for new extraction, and satisfactory extraction efficiency results were obtained (91.35%), the purity of recovered IL is 92.62% after further treatments.



**Figure 6.** The back-extraction efficiencies for flavonoids using different organic solvents.

### 3. Materials and Methods

#### 3.1. Materials and Reagents

*A. venetum* L. leaves were purchased from Hunan Xin Shen Zhi Lin Chinese Herbal Medicine Co. Ltd., (Changsha, China). The leaves were dried at 50 °C then crushed to powder of 60 mesh (250 µm) particle size. Five imidazolium ILs used in this study of  $[C_n\text{mim}]\text{Br}$  ( $n = 2, 4$ , and 6, 1-alkyl-3-methylimidazolium bromide), 1-butyl-3-methylimidazolium tetrafluoroborate ( $[C_4\text{mim}][\text{BF}_4]$ ), and 1-butyl-3-methylimidazolium dicyanamide ( $[C_4\text{mim}][\text{N}(\text{CN})_2]$ ) were synthesized using reported procedures [39,40]. The rutin standard (HPLC purity > 98%) was acquired from Aladdin Industrial Corporation (Shanghai, China). The phase-forming salts  $\text{NaH}_2\text{PO}_4 \cdot 2\text{H}_2\text{O}$  and  $(\text{NH}_4)_2\text{SO}_4$  (analytical grade) were purchased from Sinopharm Chemical Reagent Co., Ltd. (Shanghai, China). All other chemicals were of reagent-grade quality and used without further treatment.

#### 3.2. Ultrasonic-Assisted Extraction

The IL aqueous solution was used for UAE of flavonoids. To a tube, a certain amount of *A. venetum* leaf powder and IL aqueous solution were added to a certain liquid/solid ratio, and the tube was placed in a KQ-5200DE ultrasonic instrument (Kunshan Ultrasound Co. Ltd., Jiangsu, China) at 30 °C for extraction of flavonoids in a certain time. The electric power and generators frequency is 200 W and 40 kHz, respectively. Then, the tubes were centrifuged at  $10,000 \times g$  for 10 min to remove the sediment. The supernatant was separated for further analysis of the total flavonoids content. The  $\text{NaNO}_2\text{-Al}(\text{NO}_3)_3\text{-NaOH}$  colorimetry method reported by Ozsoy [41] was used to determine the total flavonoids content at 510 nm using UV-2100 spectrophotometer (Unico, Dayton, NJ, USA). The calibration curve for the flavonoids analysis was  $Y = 0.5985X + 0.0575$  with  $R^2 = 0.9997$  (RSD = 1.0%,  $n = 5$ ), where  $Y$  is represented as absorbance,  $X$  is represented as concentration of rutin standard at 0.02–0.1 µg/mL. The extraction yield of total flavonoids was calculated using Equation (1):

$$\text{Extraction yield} \left( \frac{\text{mg}}{\text{g}} \right) = \frac{\text{Mass of total flavonoids determined (mg)}}{\text{Mass of Apocynum venetum L. leaf powder (g)}} \quad (1)$$

#### 3.3. Aqueous Biphasic System

To a tube, a certain mass of salt and IL extract were added. The tube was centrifuged at  $5000 \times g$  for 5 min after complete dissolution of salt. Two clear phases (one top IL-rich phase and the other bottom salt-rich phase) were observed with the volume of each phase being noted. Samples in top phase were withdrawn to another tube. The total flavonoids content in top phase was determined, and the mass in bottom phase was calculated by mass balance. The phase ratio ( $R$ ) was defined according to Equation (2):

$$R = \frac{V_t}{V_b} \quad (2)$$



where  $V_t$  and  $V_b$  were the volumes of the top and bottom phase, respectively. The partition coefficient ( $K$ ) was calculated according to Equation (3):

$$K = \frac{C_t}{C_b} \quad (3)$$

where  $C_t$  was the total flavonoids content in top phase,  $C_b$  was the total flavonoids content in bottom phase. The extraction efficiency ( $E$ ) of total flavonoids in top phase was obtained according to Equation (4):

$$E = \frac{K}{\left(K + \frac{1}{R}\right)} \times 100\% \quad (4)$$

The preconcentration factor ( $F$ ) of the total flavonoids defined as given in Equation (5):

$$F = \frac{C_t}{C_{initial}} \quad (5)$$

where  $C_{initial}$  and  $C_t$  represent the total flavonoids content in the initial aqueous solution before and after ABS extraction in the IL-rich phase, respectively.

#### 3.4. Back-Extraction of Flavonoids and Recycle of IL

The flavonoids in the IL-rich phase were separated by back extraction using water immiscible organic solvents (dichloromethane, chloroform, *n*-butanol, isoamyl alcohol, diethyl ether, petroleum ether, ethyl acetate, and *n*-hexane) as reported by Bogdanov [42]. The flavonoids are back-extracted into the organic phase, while the ILs remain in aqueous phase and can thus be recycled.

## 4. Conclusions

In this work, IL-UAE coupled with ABS were applied for the extraction of flavonoids from *A. venetum* leaf using the IL [C<sub>4</sub>mim][N(CN)<sub>2</sub>] aqueous solution. The extraction conditions of IL concentration, liquid/solid ratio, and ultrasonic time were optimized by RSM, with further purification of the flavonoids by ABS. Using this developed methodology, we could obtain a preconcentration factor of 3.78%, with a remarkable extraction efficiency of 93.35%. After ABS, the back-extraction efficiency of flavonoids reached to 83.5% using *n*-butanol. The obtained results indicated that this IL-ABS coupled with ultrasound had the potentiality in the extraction and isolation of pharmaceutical components from various plant sources.

**Acknowledgments:** This work was financially supported by the National Natural Science Foundation of China (No. 21406262), the earmarked fund for the China Agriculture Research System (No. CARS-19), and the Agricultural Science and Technology Innovation Program (No. ASTIP-IBFC07).

**Author Contributions:** Zhijian Tan and Chaoyun Wang conceived and designed the experiments; Yongjian Yi, Hongying Wang, and Wanlai Zhou performed the experiments; Zhijian Tan analyzed the data; Zhijian Tan wrote and revised the paper.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Irie, K.; Sato, T.; Tanaka, I.; Nakajima, J.-I.; Kawaguchi, M.; Himi, T. Cardiotoxic effect of *Apocynum venetum* L. Extracts on isolated guinea pig atrium. *J. Nat. Med.* **2009**, *63*, 111–116. [[CrossRef](#)] [[PubMed](#)]
- Liang, T.; Yue, W.; Li, Q. Comparison of the phenolic content and antioxidant activities of *Apocynum venetum* L. (luo-bu-ma) and two of its alternative species. *Int. J. Mol. Sci.* **2010**, *11*, 4452–4464. [[CrossRef](#)] [[PubMed](#)]
- Kamata, K.; Seo, S.; Nakajima, J.-I. Constituents from leaves of *Apocynum venetum* L. *J. Nat. Med.* **2008**, *62*, 160–163. [[CrossRef](#)] [[PubMed](#)]
- Xie, W.; Zhang, X.; Wang, T.; Hu, J. Botany, traditional uses, phytochemistry and pharmacology of *Apocynum venetum* L. (luobuma): A review. *J. Ethnopharmacol.* **2012**, *141*, 1–8. [[CrossRef](#)] [[PubMed](#)]

5. Grundmann, O.; Nakajima, J.-I.; Seo, S.; Butterweck, V. Anti-anxiety effects of *Apocynum venetum* L. In the elevated plus maze test. *J. Ethnopharmacol.* **2007**, *110*, 406–411. [[CrossRef](#)] [[PubMed](#)]
6. Lin, X.; Wang, Y.; Liu, X.; Huang, S.; Zeng, Q. IIs-based microwave-assisted extraction coupled with aqueous two-phase for the extraction of useful compounds from Chinese medicine. *Analyst* **2012**, *137*, 4076–4085. [[CrossRef](#)] [[PubMed](#)]
7. Cao, Y.H.; Zhang, X.; Fang, Y.H.; Ye, J.N. Determination of active ingredients of *Apocynum venetum* by capillary electrophoresis with electrochemical detection. *Mikrochim. Acta* **2001**, *137*, 57–62. [[CrossRef](#)]
8. Ma, M.; Hong, C.L.; An, S.Q.; Li, B. Seasonal, spatial, and interspecific variation in quercetin in *Apocynum venetum* and *Poacynum hendersonii*, Chinese traditional herbal teas. *J. Agric. Food Chem.* **2003**, *51*, 2390–2393. [[CrossRef](#)] [[PubMed](#)]
9. Zhao, H.; Xia, S.; Ma, P. Use of ionic liquids as “Green” Solvents for extractions. *J. Chem. Technol. Biotechnol.* **2005**, *80*, 1089–1096. [[CrossRef](#)]
10. Rogers, R.D. Materials science—Reflections on ionic liquids. *Nature* **2007**, *447*, 917–918. [[CrossRef](#)] [[PubMed](#)]
11. Coutinho, J.A.P.; Neves, C.M.S.S.; Ventura, S.P.M.; Freire, M.G.; Marrucho, I.M. Evaluation of cation influence on the formation and extraction capability of ionic-liquid-based aqueous biphasic systems. *J. Phys. Chem. B* **2009**, *113*, 5194–5199.
12. Tang, S.; Baker, G.A.; Zhao, H. Ether- and alcohol-functionalized task-specific ionic liquids: Attractive properties and applications. *Chem. Soc. Rev.* **2012**, *41*, 4030–4066. [[CrossRef](#)] [[PubMed](#)]
13. Albishri, H.M.; Abd El-Hady, D. Eco-friendly ionic liquid based ultrasonic assisted selective extraction coupled with a simple liquid chromatography for the reliable determination of acrylamide in food samples. *Talanta* **2014**, *118*, 129–136. [[CrossRef](#)] [[PubMed](#)]
14. Zhang, Y.; Liu, Z.; Li, Y.; Chi, R. Optimization of ionic liquid-based microwave-assisted extraction of isoflavones from *Radix puerariae* by response surface methodology. *Sep. Purif. Technol.* **2014**, *135*, 285–285. [[CrossRef](#)]
15. Duan, M.-H.; Luo, M.; Zhao, C.-J.; Wang, W.; Zu, Y.-G.; Zhang, D.-Y.; Yao, X.-H.; Fu, Y.-J. Ionic liquid-based negative pressure cavitation-assisted extraction of three main flavonoids from the pigeonpea roots and its pilot-scale application. *Sep. Purif. Technol.* **2013**, *107*, 26–36. [[CrossRef](#)]
16. Chatel, G.; Macfarlane, D.R. Ionic liquids and ultrasound in combination: Synergies and challenges. *Chem. Soc. Rev.* **2014**, *43*, 8132–8149. [[CrossRef](#)] [[PubMed](#)]
17. Yang, L.; Liu, Y.; Zu, Y.-G.; Zhao, C.-J.; Zhang, L.; Chen, X.-Q.; Zhang, Z.-H. Optimize the process of ionic liquid-based ultrasonic-assisted extraction of aesculin and aesculetin from *Cortex fraxini* by response surface methodology. *Chem. Eng. J.* **2011**, *175*, 539–547. [[CrossRef](#)]
18. Cao, X.; Ye, X.; Lu, Y.; Yu, Y.; Mo, W. Ionic liquid-based ultrasonic-assisted extraction of piperine from white pepper. *Anal. Chim. Acta* **2009**, *640*, 47–51. [[CrossRef](#)] [[PubMed](#)]
19. Lin, H.; Zhang, Y.; Han, M.; Yang, L. Aqueous ionic liquid based ultrasonic assisted extraction of eight ginsenosides from ginseng root. *Ultrason. Sonochem.* **2013**, *20*, 680–684. [[CrossRef](#)] [[PubMed](#)]
20. Yang, L.; Wang, H.; Zu, Y.-G.; Zhao, C.; Zhang, L.; Chen, X.; Zhang, Z. Ultrasound-assisted extraction of the three terpenoid indole alkaloids vindoline, catharanthine and vinblastine from *Catharanthus roseus* using ionic liquid aqueous solutions. *Chem. Eng. J.* **2011**, *172*, 705–712. [[CrossRef](#)]
21. Ma, C.-H.; Liu, T.-T.; Yang, L.; Zu, Y.-G.; Wang, S.-Y.; Zhang, R.-R. Study on ionic liquid-based ultrasonic-assisted extraction of biphenyl cyclooctene lignans from the fruit of *Schisandra chinensis* baill. *Anal. Chim. Acta* **2011**, *689*, 110–116. [[CrossRef](#)] [[PubMed](#)]
22. Zhang, L.; Wang, X. Hydrophobic ionic liquid-based ultrasound-assisted extraction of magnolol and honokiol from cortex *Magnoliae officinalis*. *J. Sep. Sci.* **2010**, *33*, 2035–2038. [[CrossRef](#)] [[PubMed](#)]
23. Gutowski, K.E.; Broker, G.A.; Willauer, H.D.; Huddleston, J.G.; Swatloski, R.P.; Holbrey, J.D.; Rogers, R.D. Controlling the aqueous miscibility of ionic liquids: Aqueous biphasic systems of water-miscible ionic liquids and water-structuring salts for recycle, metathesis, and separations. *J. Am. Chem. Soc.* **2003**, *125*, 6632–6633. [[CrossRef](#)] [[PubMed](#)]
24. Tan, Z.; Li, F.; Xu, X. Isolation and purification of aloe anthraquinones based on an ionic liquid/salt aqueous two-phase system. *Sep. Purif. Technol.* **2012**, *98*, 150–157. [[CrossRef](#)]
25. Claudio, A.F.M.; Ferreira, A.M.; Freire, C.S.R.; Silvestre, A.J.D.; Freire, M.G.; Coutinho, J.A.P. Optimization of the gallic acid extraction using ionic-liquid-based aqueous two-phase systems. *Sep. Purif. Technol.* **2012**, *97*, 142–149. [[CrossRef](#)]

26. Fan, J.-P.; Cao, J.; Zhang, X.-H.; Huang, J.-Z.; Kong, T.; Tong, S.; Tian, Z.-Y.; Zhu, J.-H.; Ouyang, X.-K. Extraction of puerarin using ionic liquid based aqueous two-phase systems. *Sep. Sci. Technol.* **2012**, *47*, 1740–1747. [[CrossRef](#)]
27. Freire, M.G.; Neves, C.M.S.S.; Marrucho, I.M.; Lopes, J.N.C.; Rebelo, L.P.N.; Coutinho, J.A.P. High-performance extraction of alkaloids using aqueous two-phase systems with ionic liquids. *Green Chem.* **2010**, *12*, 1715–1718. [[CrossRef](#)]
28. Shi, J.; Li, G.; Zhang, R.; Zheng, J.; Suo, Y.; You, J.; Liu, Y.-J. A validated HPLC-DAD-MS method for identifying and determining the bioactive components of two kinds of luobuma. *J. Liquid Chromatogr. Relat. Technol.* **2011**, *34*, 537–547. [[CrossRef](#)]
29. Zhang, Y.; Liu, C.; Zhang, Z.; Wang, J.; Wu, G.; Li, S. Comprehensive separation and identification of chemical constituents from *Apocynum venetum* leaves by high-performance counter-current chromatography and high performance liquid chromatography coupled with mass spectrometry. *J. Chromatogr. B* **2010**, *878*, 3149–3155. [[CrossRef](#)] [[PubMed](#)]
30. An, H.; Wang, H.; Lan, Y.; Hashi, Y.; Chen, S. Simultaneous qualitative and quantitative analysis of phenolic acids and flavonoids for the quality control of *Apocynum venetum* L. Leaves by HPLC-DAD-ESI-IT-TOF-MS and HPLC-DAD. *J. Pharm. Biomed.* **2013**, *85*, 295–304. [[CrossRef](#)] [[PubMed](#)]
31. Claudio, A.F.M.; Swift, L.; Hallett, J.P.; Welton, T.; Coutinho, J.A.P.; Freire, M.G. Extended scale for the hydrogen-bond basicity of ionic liquids. *Phys. Chem. Chem. Phys.* **2014**, *16*, 6593–6601. [[CrossRef](#)] [[PubMed](#)]
32. Bogdanov, M.G. Ionic liquids as alternative solvents for extraction of natural products. In *Alternative Solvents for Natural Products Extraction*; Chemat, F., Abert-Vian, M., Eds.; Springer: Berlin, Germany; Heidelberg, Germany, 2014; pp. 127–166.
33. Saleh, A.; Yamini, Y.; Faraji, M.; Rezaee, M.; Ghambarian, M. Ultrasound-assisted emulsification microextraction method based on applying low density organic solvents followed by gas chromatography analysis for the determination of polycyclic aromatic hydrocarbons in water samples. *J. Chromatogr. A* **2009**, *1216*, 6673–6679. [[CrossRef](#)] [[PubMed](#)]
34. Han, D.; Zhu, T.; Row, K.H. Ultrasonic extraction of phenolic compounds from *Laminaria japonica* aresch using ionic liquid as extraction solvent. *Bull. Korean Chem. Soc.* **2011**, *32*, 2212–2216. [[CrossRef](#)]
35. Cao, X.; Qiao, J.; Wang, L.; Ye, X.; Zheng, L.; Jiang, N.; Mo, W. Screening of glycoside isomers in *P. Scrophulariiflora* using ionic liquid-based ultrasonic-assisted extraction and ultra-performance liquid chromatography/electrospray ionization quadrupole time-of-flight tandem mass spectrometry. *Rapid Commun. Mass Spectrom.* **2012**, *26*, 740–748. [[CrossRef](#)] [[PubMed](#)]
36. Freire, M.G.; Neves, C.; Marrucho, I.M.; Coutinho, J.A.P.; Fernandes, A.M. Hydrolysis of tetrafluoroborate and hexafluorophosphate counter ions in imidazolium-based ionic liquids. *J. Phys. Chem. A* **2010**, *114*, 3744–3749. [[CrossRef](#)] [[PubMed](#)]
37. Archer, D.G.; Widegren, J.A.; Kirklin, D.R.; Magee, J.W. Enthalpy of solution of 1-octyl-3-methylimidazolium tetrafluoroborate in water and in aqueous sodium fluoride. *J. Chem. Eng. Data* **2005**, *50*, 1484–1491. [[CrossRef](#)]
38. Yoshida, Y.; Baba, O.; Larriba, C.; Saito, G. Imidazolium-based ionic liquids formed with dicyanamide anion: Influence of cationic structure on ionic conductivity. *J. Phys. Chem. B* **2007**, *111*, 12204–12210. [[CrossRef](#)] [[PubMed](#)]
39. Fredlake, C.P.; Crosthwaite, J.M.; Hert, D.G.; Aki, S.N.V.K.; Brennecke, J.F. Thermophysical properties of imidazolium-based ionic liquids. *J. Chem. Eng. Data* **2004**, *49*, 954–964. [[CrossRef](#)]
40. Stoppa, A.; Hunger, J.; Buchner, R. Conductivities of binary mixtures of ionic liquids with polar solvents. *J. Chem. Eng. Data* **2009**, *54*, 472–479. [[CrossRef](#)]
41. Ozsoy, N.; Can, A.; Yanardag, R.; Akev, N. Antioxidant activity of *Smilax excelsa* L. Leaf extracts. *Food Chem.* **2008**, *110*, 571–583. [[CrossRef](#)]
42. Bogdanov, M.G.; Keremedchieva, R.; Svinyarov, I. Ionic liquid-supported solid-liquid extraction of bioactive alkaloids. III. Ionic liquid regeneration and glaucine recovery from ionic liquid-aqueous crude extract of *Glaucium flavum* cr.(papaveraceae). *Sep. Purif. Technol.* **2015**, *155*, 13–19. [[CrossRef](#)]

**Sample Availability:** Samples of the compounds flavonoids from *A. venetum* L. leaves are available from the authors.

