



Minireview

Chemically Induced Cellular Proteolysis: An Emerging Therapeutic Strategy for Undruggable Targets

Seonghyeon Moon¹, and Byung-Hoon Lee^{1,*}

¹Department of New Biology, DGIST, Daegu 42988, Korea
*Correspondence: byung-hoon_lee@dgist.ac.kr
<http://dx.doi.org/10.14348/molcells.2018.0372>
www.molcells.org

Traditionally, small-molecule or antibody-based therapies against human diseases have been designed to inhibit the enzymatic activity or compete for the ligand binding sites of pathological target proteins. Despite its demonstrated effectiveness, such as in cancer treatment, this approach is often limited by recurring drug resistance. More importantly, not all molecular targets are enzymes or receptors with druggable 'hot spots' that can be directly occupied by active site-directed inhibitors. Recently, a promising new paradigm has been created, in which small-molecule chemicals harness the naturally occurring protein quality control machinery of the ubiquitin-proteasome system to specifically eradicate disease-causing proteins in cells. Such 'chemically induced protein degradation' may provide unprecedented opportunities for targeting proteins that are inherently undruggable, such as structural scaffolds and other non-enzymatic molecules, for therapeutic purposes. This review focuses on surveying recent progress in developing E3-guided proteolysis-targeting chimeras (PROTACs) and small-molecule chemical modulators of deubiquitinating enzymes upstream of or on the proteasome.

Keywords: deubiquitinating enzyme, induced proteolysis, PROTAC, small-molecules, ubiquitin-proteasome system, undruggable target

INTRODUCTION

Cellular proteolysis can be achieved through two major pathways: the ubiquitin-proteasome system (UPS) and the lysosomal degradation pathway (Ciechanover, 2005). Ubiquitination, a type of protein post-translational modification, is covalent conjugation of the small, stable protein, ubiquitin through the formation of an isopeptide bond between a Gly of ubiquitin and a Lys on the substrate. This reaction occurs through an enzymatic cascade of ubiquitin-modifying enzymes comprising E1 (ubiquitin activating enzyme), E2 (ubiquitin conjugating enzyme), and E3 (ubiquitin ligase) (Finley and Chau, 1991). Although ubiquitin conjugation may serve non-proteolytic functions, its most recognized function is targeted protein degradation via the 26S proteasome, a multi-protein protease complex (Fig. 1) (Finley, 2009; Komander and Rape, 2012). Ubiquitination process is tightly controlled, reflecting the critical importance of protein turnover rates for cellular function. This requirement strongly suggests that the ubiquitination reaction must be remarkably versatile. Notable in this context, more than 600 E3 ligases are known, helping to explain the observed substrate specificity (Deshaies and Joazeiro, 2009); moreover, deubiquitinating enzymes (DUBs) can reverse this reaction by break-

Received 4 September 2018; revised 28 October 2018; accepted 30 October 2018; published online 7 November, 2018

eISSN: 0219-1032

©The Korean Society for Molecular and Cellular Biology. All rights reserved.

©This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>.

ing down ubiquitin polymers (Wilkinson, 1997). E3 ligases transfer ubiquitin from a ubiquitin-charged E2 complex to the substrate and this reaction can be repeated, producing a polyubiquitinated substrate that is recognized by the proteasome. The specificity of E3 ligases suggests that E3 inhibitors could produce more targeted effects and thus might be less harsh than E1- or E2-targeting drugs. In fact, several studies have shown that specifically targeting E3 ligase may be therapeutically effective against certain cancers (Chan et al., 2013; Shangary et al., 2008; Vassilev et al., 2004). In addition to E3 ligases, the human proteome contains approximately 100 DUBs belonging to at least six subfamilies (Komander et al., 2009), which play key roles in ubiquitin-mediated proteolytic pathways as well as other biological processes, including (a) rescue of protein substrates from degradation by ubiquitin removal, (b) editing of ubiquitin chains to regulate the function or half-lives of protein substrates, and (c) ubiquitin recycling into free chains or ubiquitin monomers.

Since the initial discovery of the proteolytic machinery, a plethora of evidence has shown that the UPS is closely associated with various diseases, such as cancers and neurodegenerative diseases (Popovic et al., 2014). These findings have garnered immense attention in treating such diseases by targeting the UPS. For example, following a path similar to that for kinase inhibitors, bortezomib and its progeny have been successfully developed as FDA-approved anti-cancer drugs for the treatment of multiple myeloma (MM) and other blood cancers, and in fact, these agents are proteasome inhibitors that block the proteolytic reaction (Manasanch and Orlowski, 2017; Richardson et al., 2005; Stewart et al., 2015). Moreover, a number of E3 ligase inhibitors and DUB inhibitors have been tested in preclinical or clinical phase studies, representing novel strategies for target-directed therapies (Gopinath et al., 2016; Farshi et al., 2015; Weathington and Mallampalli, 2014).

Traditional active site-directed therapies, which generally employ small-molecule inhibitors and antibodies, are designed to inhibit the activity of target proteins. Small-molecule chemicals have proven effective for inhibiting several oncogenic enzymes, including protein kinases (e.g., imatinib and erlotinib), histone deacetylases (e.g., belinostat), and poly (ADP-ribose) polymerase (PARP) (e.g., olaparib) by directly targeting their active sites and thereby disrupting the enzymatic function (Salami and Crews, 2017). Despite being promising, these approaches have inherent limitations. First, there are only about 400 human proteins as possible drug targets among ~3000 disease-related proteins (Rask-Andersen et al., 2011). Furthermore, more than 90% of these possible protein targets fall into the categories of GPCRs, enzymes, CD markers, transporters and ion channels, which are considered “druggable”. This suggests that current therapeutics target only a fraction of the disease-causing proteome. Second, small-molecule and antibody binding is predominantly non-covalent and reversible. This indicates that treatment efficacy requires the inhibitors to be maintained at high concentration. Maintaining a high dose of drug is one of the major challenges in modern drug development, and this may also cause unexpected side effects

(Salami and Crews, 2017). Lastly, antibodies induce altered cellular responses by blocking extracellular protein-protein or protein-ligand interactions. Such antibodies, including rituximab, tocilizumab, and belimumab, are effective against tumors and several autoimmune diseases. The main advantage of antibody-based therapy is its high affinity for the target protein and the longer half-life owing to endosomal FcRn-IgG recycling. However, therapeutic antibodies are also highly limited because of their poor cell permeability, oral unavailability, and high cost (Neklesa et al., 2017).

An emerging new paradigm of disease treatment—“targeted protein degradation (TPD)”—promises to help overcoming the limitations of typical active site-directed therapies (Zhou, 2005). TPD aims to induce degradation of target proteins, especially those considered “undruggable”, by harnessing the endogenous protein quality control machinery—the UPS. One elaborate approach is to induce ubiquitin tagging and subsequent target protein degradation through E3-guided proteolysis-targeting chimeras (PROTACs). Small-molecule DUB inhibitors that act upstream of or on the proteasome can also be utilized for antagonizing substrate deubiquitination. This review will focus on recent progress in the development of TPD, specifically PROTACs and DUB inhibitors, and their therapeutic potential. Figure 1 highlights PROTAC and DUB inhibitors as chemical proteolytic inducers in the UPS pathway.

INDUCED PROTEOLYSIS BY PROTACS

PROTAC, an E3-guided proteolysis-targeting chimera, is a bifunctional molecule composed of three parts: an E3 ligase binding ligand, a substrate binding ligand, and a linker between the two. E3 ligase generally requires sequential recognition events through multi-step E1/E2/substrate reactions (Deshaies and Joazeiro, 2009). PROTACs can simplify this process—they transiently recruit target substrate and E3 ligase in close proximity and facilitate the ubiquitination of the substrate (Coleman and Crews, 2018). As a result, target proteins become more efficiently polyubiquitinated and undergo degradation by the 26S proteasome (Fig. 1). This idea originated with the elegant work of Sakamoto and colleagues, in which the researchers developed a chimera linking the E3 ligase SCF- β -TRCP (Skp1/Cullin/F box- β -TRCP) to methionine aminopeptidase 2 (MetAP2) (Sakamoto et al., 2001). This chimeric molecule, PROTAC-1, contains a 10-amino acid I κ B α phosphopeptide that can be recognized by β -TRCP at one end, while the other side is an MetAP1 inhibitor, ovalicin. PROTAC-1 indeed promotes degradation of MetAP1 in a ubiquitination-dependent manner. Since first-generation PROTACs had low cell permeability because of the peptidic sequence, next-generations adopted von Hippel-Lindau (VHL) as an E3 ligase instead of SCF- β -TRCP (Schneekloth et al., 2004). This variant, developed by Crews and colleagues, exploited a short hydroxyproline-containing peptide derived from the VHL substrate, hypoxia-inducible factor 1- α (HIF1 α), and they applied the resulting chimeras for selective degradation of androgen receptor (AR) and FK506-binding protein 12 (FKBP12) in intact cells. Although the hydroxyproline-containing degron displayed improved




Fig. 1. Mechanisms of induced protein degradation by PROTACs and DUB inhibitors. (Left) General scheme of ubiquitin-proteasome system. An enzymatic cascade of E1-E2-E3 transfers ubiquitin to Lys residues on the substrate. As a consequence, the polyubiquitinated substrate is recognized by the 26S proteasome and undergoes degradation. (Right) Induced proteolysis by PROTAC and/or DUB inhibitor. PROTAC links E3 and the target protein, enhances E3-mediated ubiquitination, and promotes degradation of the target molecule. By contrast, DUBs acting upstream of the proteasome (violet) or on the proteasome (olive drab) can inhibit degradation through substrate deubiquitination. Therefore, DUB inhibitors may facilitate the proteasomal degradation pathway by antagonizing deubiquitination.

Table 1. Representative examples of PROTACs

E3 ligase/HyT ^a	E3 binder/HyT (type ^b)	Target binder (type)	Target	Reference
β -TRCP	I κ B α .phosphopeptide (P)	Ovalicin (S)	MetAP2	Sakamoto et al., 2001
cIAP	MeBS (S)	ATRA (S)	CRABP-I, II	Itoh et al., 2010
	BE04 (S)	Ch55 / Estrone / DHT (S)	RAR / ER / AR	Itoh et al., 2011
	BE04 (S)	KHS108 (S)	TACC3	Ohoka et al., 2014
	BE04 (S)	Alkyl chloride analog (S)	HaloTag	Tomoshige et al., 2015
CRBN	Thalidomide (S)	JQ1 / SLF (S)	BET family / FKBP12	Winter et al., 2015
	Pomalidomide (S)	Bosutinib, Dasatinib (S)	BCR-ABL	Lai et al., 2016
	Thalidomide (S)	Aminopyrazole analog (S)	CDK9	Robb et al., 2017
	Pomalidomide (S)	Ceritinib (S)	ALK	Zhang et al., 2018
	Lenalidomide (S)	HJB97 (S)	BET family	Zhou et al., 2018
	Pomalidomide (S)	Ibrutinib derivative (S)	BTK (Bruton's Tyr kinase)	Buhimschi et al., 2018
	Pomalidomide (S)	HDAC inhibiting aldehydes (S)	HDAC6	Yang et al., 2018
	Thalidomide (S)	Sirtuin Rearranging Ligand (SirReal) (S)	Sirtuin 2 (Sirt2)	Schiedel et al., 2018
Keap1	Keap1 binding motif (P)	Tau binding motif (P)	Tau	Lu et al., 2018
MDM2	Nutlin-3 (S)	SARM (S)	AR	Schneekloth et al., 2008
VHL	HIF α degron (P)	DHT / AP21998 (S)	AR / FKBP12	Schneekloth et al., 2004
	HIF α degron (P)	DHT / Estradiol (S)	AR / ER	Rodriguez-Gonzalez et al., 2008
	HIF α degron (P)	TrkA degron / ErbB3 degron (P)	FRS2 α / PI3K	Hines et al., 2013
	HIF α derivative (S)	Vandetanib / Phenoxy TZD analog (S)	RIPK2 / ER α	Bondeson et al., 2015
	HIF α derivative (S)	JQ1 (S)	BRD4	Zengerle et al., 2015
	HIF α derivative (S)	Chloroalkane analog (S)	HaloTag	Buckley et al., 2015
	HIF α derivative (S)	Triazolo-diazepine acetamide (S)	BET family	Raina et al., 2016
	HIF α degron (P)	CPP-tri_a (S)	AKT	Henning et al., 2016
	HIF α derivative (S)	Dasatinib (S)	c-ABL	Lai et al., 2016
	HIF α degron (P)	EN300-72284 (S)	Smad3	Wang et al., 2016a
	HIF α degron (P)	Tau binding motif (P)	Tau	Chu et al., 2016
	HIF α derivative (S)	TBK1 ligand (S)	TBK1 (TANK-Binding Kinase 1)	Crew et al., 2018
	HIF α degron (P)	TD-PERM (P)	ER α	Jiang et al., 2018
	HIF α derivative (S)	Lapatinib, Gefitinib, Afatinib (S)	EGFR, HER2, c-Met	Burslem et al., 2018
HyT	Adamantane (S)	Haloalkane reactive linker (S)	HaloTag	Neklesa et al., 2011
	Boc ₃ Arg (S)	Trimethoprim / EA, Thiobenzofurazan (S)	DHFR / GST	Long et al., 2012
	Adamantane (S)	TX1-85-1 (S)	HER3	Xie et al., 2014
	Adamantane (S)	RU59063 (S)	AR	Gustafson et al., 2015

a: HyT, Hydrophobic tagging; b: P, peptide, S, small molecule

cell applicability, this VHL-recognized peptide still lacked ideal drug-like properties. Recent efforts have further validated MDM2/Nutlins, Cereblon (CRBN)/thalidomide, cIAP ligands, and optimized VHL ligands as E3-recruiting moieties for target protein degradation in PROTAC technology (Table 1). Notably, landmark papers from Bradner and Crews groups demonstrated that small-molecule VHL- and CRBN-directed PROTACs are highly effective in degrading estrogen-related receptor α (ERR α) and Bromodomain-containing protein 4 (BRD4) in animal cancer models (Bondeson et al., 2015; Winter et al., 2015). PROTACs can be designed to behave like traditional small-molecules, yet their distinctive mode of action provides innovative therapeutic opportunities. There has also been considerable effort devoted to PROTACs into commercial drugs, exemplified by the recent foundation of Arvinas and C4 Therapeutics (Neklesa et al., 2017).

PROTACs have several advantages over active site-directed therapies, ideally overcoming the drawbacks of small-molecule inhibitors and therapeutic antibodies. The most prominent feature of PROTAC is that induced protein degradation, in principle, renders conventionally undruggable targets druggable. As previously noted, typical active site-directed drug targets constitute only about 13% of the thousands of disease-related proteins, mostly enzymes or membrane receptors. This clearly indicates that transcription factors, scaffolding proteins, and toxic protein aggregates have remained therapeutically intractable. PROTACs, by linking ligands and appropriate E3 ligases, can induce the proteolysis of traditionally undruggable proteins, including transcriptional regulators (e.g., Myc, Gli, β -catenin, and STAT family) and signaling scaffolds (e.g., ERBB3, KSR, Gab family, β -arrestin, BCL10, and AKAPs) (Coleman and Crews, 2018; Neklesa et al., 2017). Protein aggregates are particularly interesting targets because they are common features of several neurodegenerative diseases, such as tauopathies and amyotrophic lateral sclerosis (ALS) (Ross and Poirier, 2004). Recent studies have shown that the low efficacy problem of early PROTACs can be substantially resolved, as evidenced by the apparent nanomolar range of EC₅₀s of more recently developed ones (Lai and Crews, 2017). Analogous to E3-guided PROTAC, the 'hydrophobic tagging (HyT)' technology has also been developed to destabilize target proteins and/or recruit chaperones for induced protein degradation (Huang and Dixit, 2016; Lai and Crews, 2017). The novel repertoire of E3 ligases (e.g., Keap1), HyT binders, and target ligands has rapidly expanded, strongly suggesting that PROTAC may serve as a platform technology (Chu et al., 2016; Lu et al., 2018). Table 1 summarizes representative examples of PROTACs.

















SMALL-MOLECULE INHIBITORS TARGETING DEUBIQUITINATING ENZYMES UPSTREAM OF OR ON THE PROTEASOME

Reversing the ubiquitination process is exclusively fulfilled by deubiquitinating enzymes (DUBs), numbering nearly 100 in the human genome (Komander et al., 2009). To date, six subfamilies of DUBs have been identified: (1) ubiquitin carboxyl-terminal hydrolases (UCHs), (2) ubiquitin specific pro-














teases (USPs), (3) ovarian tumor like proteases (OTUs), (4) JAMM/MPN metalloproteases, (5) Machado-Jacob-disease proteases (MJDs), and (6) motif interacting with Ub-containing novel DUB family (MINDY) (Abdul Rehman et al., 2016; Clague et al., 2013). DUBs should represent promising drug targets because they become increasingly implicated in various human diseases (Harrigan et al., 2018). For example, a growing number of DUBs (e.g., USP28, JOSD1, UCHL1, CSN5, USP9x, USP10, USP11, USP22, and USP48) were found to be overexpressed in diverse cancer types. Moreover, genetic alterations of DUBs (e.g., USP6/Tre2 and USP28) can be truly oncogenic in certain cancers (Fraile et al., 2012; Sacco et al., 2010). The isopeptidase activity of DUBs can be selectively targeted by inhibiting the catalytic sites with drug-like compounds, and in this sense, they are typical targets of active site-directed inhibitors. In an interesting twist, however, most cellular proteins are degraded by the ubiquitin-mediated proteasome pathway; thus, DUBs in principle should antagonize the destruction of these proteins (Fig. 1). Indeed, deubiquitination can occur on histones, transcription factors, and other regulatory proteins, so the pool of target substrates must far exceed the number of conventionally druggable targets. Therefore, similar to PROTACs, small-molecule DUB inhibitors may also induce the degradation of previously intractable targets. While of great therapeutic interest, only a handful of DUB inhibitors have been reported, and none has reached in advanced clinical trials, providing a valuable opportunity for developing first-in-class therapeutic DUB inhibitors (Farshi et al., 2015)(Table 2).

Analogous to the cell cycle, deubiquitination reactions at two points-upstream of and on the proteasome-may serve as critical checkpoints for ubiquitin-mediated degradation pathways (Fig. 1). DUBs acting upstream of the proteasome may act as a "rescue crew", salvaging specific substrates by deubiquitination before they are engaged in degradation. Accordingly, chemical blockers of the DUB may accelerate the turnover of the specific substrate for proteasome-mediated degradation. For instance, USP7/HAUSP is the most well-recognized drug target because of its cellular role as a regulator of the tumor suppressor p53. USP7 deubiquitinates MDM2, an E3 ligase of p53, and in turn destabilizes p53 (Colland et al., 2009). Thus, by stabilizing p53, a USP7 inhibitor may be beneficial for the treatment of certain cancers. The USP7 inhibitors, HBX 19,818 and P5091, were previously reported to selectively inhibit USP7 *in vitro* and *in vivo*, and importantly, P5091 showed cytotoxicity in relapsed MM cells derived from cancer patients (Chauhan et al., 2012; Reverdy et al., 2012). More recently, a series of USP7 inhibitors have been discovered and among them, a non-covalent inhibitor FT671 and a covalent inhibitor FT827 were identified by molecular-based screening (Turnbull et al., 2017). These compounds were found to be highly selective for USP7 in counter-screening against a panel of DUBs including USP10 and USP47, which are also inhibited by P5091. *In vivo* experiments in cancer cell lines and MM.1s xenograft animal models demonstrated that FT671 induces p53 stabilization and MDM2 degradation, leading to anti-tumor activity via USP7 blocking. Another NMR and structure-based screening study identified the USP7 inhibitors,

Table 2. Representative examples of deubiquitinating enzyme inhibitors

Target DUB	DUB inhibitor	
UCH-L1	 LDN-57444 Liu et al., 2003	
USP1/UAF1	 Pimozide Chen et al., 2011	 GW7647
	 C527 Mistry et al., 2013	 ML323 Dexheimer et al., 2014
	 HBX 41108 Colland et al., 2009	 HBX 19818 Reverdy et al., 2012
	 HBX 28258	
USP7 (USP47 ^a)	 P22077 Tian et al., 2011	 P5091 Chauhan et al., 2012
	 P5091 analogue ^a Weinstock et al., 2012	
	 FT671 Turnbull et al., 2017	 FT827
	 GNE-6640 Kategaya et al., 2017	 GNE-6776
	 XL188 Lamberto et al., 2017	

(continued)

Target DUB	DUB inhibitor
USP7 (USP47 ^a)	 <p>Compound 4 Gavory et al., 2018</p>
USP9x/ USP5/ USP14/ UCH37	 <p>WP1130 Kapuria et al., 2010</p>
USP14 (UCH37 ^b)	<div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">  <p>IU1 Lee et al., 2010</p> </div> <div style="text-align: center;">  <p>IU1-47 Boselli et al., 2017</p> </div> </div> <div style="display: flex; justify-content: space-around; margin-top: 20px;"> <div style="text-align: center;">  <p>CuPT^b Liu et al., 2014a</p> </div> <div style="text-align: center;">  <p>Auranofin^b Liu et al., 2014b</p> </div> </div> <div style="display: flex; justify-content: space-around; margin-top: 20px;"> <div style="text-align: center;">  <p>b-AP15^b D'Arcy et al., 2011</p> </div> <div style="text-align: center;">  <p>VLX1570^b Wang et al., 2016b</p> </div> </div>
USP30	<div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">  <p>15-oxospiramylactone Yue et al., 2014</p> </div> <div style="text-align: center;">  <p>MF-094 Kluge et al., 2018</p> </div> <div style="text-align: center;">  <p>MF-095 Kluge et al., 2018</p> </div> </div>
RPN11	 <p>Capzimin Li et al., 2017</p>
Broad DUB inhibitor	 <p>PR-619 Tian et al., 2011</p>

a: reported to inhibit both USP7 and USP47; b: reported to inhibit both USP14 and UCH37

GNE-6640 and GNE-6776 (Kategaya et al., 2017). These compounds may selectively interfere with K48 linkage-directed ubiquitin chain cleavage mediated by USP7, suggesting that K48-linked substrates such as MDM2 could be

susceptible. More recently, an elegant fragment-based screen combined with structure-guided medicinal chemistry identified a highly potent and selective USP7 inhibitor, "compound 4" (IC₅₀ = 6 nM). This allosteric inhibitor showed

strong anti-proliferative effects against several cancer cell lines with equal or even greater efficacy compared to known clinical MDM2 antagonists (Gavory et al., 2018). A mitochondria-localized DUB, USP30 may also represent a promising therapeutic target due to its involvement in mitophagy-related Parkinson's disease as well as cancers. USP30 antagonizes Parkin-mediated ubiquitination on multiple mitochondrial substrates (Bingol et al., 2014; Liang et al., 2015). Recently, a potent USP30 inhibitor MF-094 was developed through high-throughput screening and subsequent structure-activity relationship (SAR) studies of acyl benzenesulfonamide derivatives, and this compound showed the increased mitophagy in C2C12 cells (Kluge et al., 2018).

Targeting DUBs on the proteasome may also offer an exciting strategy for induced protein degradation. There are three major and distinctive DUBs on human proteasome: USP14, UCH37, and RPN11 (de Poot et al., 2017; Finley, 2009). USP14 and UCH37 may rescue substrates from degradation prior to the proteasome's commitment step, whereas RPN11 is coupled to degradation. Finley and colleagues have screened out highly selective USP14 inhibitors, IU1 and its derivatives, and showed that their treatment promotes the degradation of proteopathic substrates in neurodegenerative disease models (Boselli et al., 2017; Lee et al., 2010; 2016). USP14 inhibitors may uncheck and bypass the deubiquitination-mediated proteolytic checkpoint on the proteasome under certain conditions of proteotoxic stress. By contrast, the proteasome 19S DUB inhibitors, b-AP15 and VLX1570, were reported to suppress tumor progression by inhibiting both USP14 and UCH37 activities (D'Arcy et al., 2011; Wang et al., 2015; 2016b). b-AP15 treatment leads to accumulation of polyubiquitinated conjugates and inhibition of protein degradation. Recently, capzimin was identified as a potent and specific RPN11 inhibitor (Li et al., 2017). Capzimin, a quinoline-8-thiol (8-TQ) derivative, induced the stabilization of proteasome substrates and inhibited cancer cell proliferation probably through the unfolded protein response (UPR). Unlike IU1, the anti-tumor effects of b-AP15 and capzimin might rely on 'restrained protein degradation' rather than induced proteolysis.

FUTURE PERSPECTIVES

Here we described PROTACs and DUB inhibitors—two emerging strategies of chemically induced proteolysis that utilize the endogenous ubiquitin-proteasome system to inhibit previously undruggable targets. While certainly bearing tremendous promise for new therapeutic applications, these approaches could also face several challenges. For example, current PROTACs are orally unavailable, probably due to its relatively large size, typically 700–1000 Da. Their pharmacokinetic properties also need to be improved for better drug metabolism. Besides, only a few E3 ligases have been exploited, and not all E3 ligases might be co-expressed with target proteins in specific tissues, which makes diagnostics arduous (Huang and Dixit, 2016). PROTAC optimization—E3 ligase selection, ligand availability, and linker design—is another challenging issue. In this context, ligand screening can

be performed by advanced screening tools, such as computer-aided drug design and DNA-encoded small molecule libraries, which can be accomplished on the order of $\sim 10^9$ compounds in a single vial (Chan et al., 2015). Although DUB inhibitors might be more orally bioavailable, their specificity and utility still remain to be explored. Given the smaller pool of DUB members compared to over 600 E3 ligases, DUB inhibitors may target only a subset of substrates with limited specificity. Nevertheless, one can envisage that 'degradation inducers' might pioneer the valuable therapeutic strategies and provide more advanced platform technologies, leading to a new era of UPS-related drug development. It will be also interesting to investigate potential combinatorial treatment with E3-guided and DUB-guided degradation inducers or to design new classes of chemically induced proteolysis chimeras recruiting substrates, E3s, DUBs, chaperones, or proteasomes.

ACKNOWLEDGEMENTS

The authors apologize to researchers whose citations are not included due to space limitations. We greatly appreciate HJ Park for help with graphics and other lab members for critical reading. BHL gratefully acknowledges the National Research Foundation of Korea (NRF) grant, DGIST Start-up Fund, and HRHR grant funded by the Ministry of Science, ICT, and Future Planning (2017R1A2B4006671, 2018010095, and 2018010146) for financial support.

REFERENCES

- Abdul Rehman, Syed A., Kristariyanto Yosua, A., Choi, S.Y., Nkosi, P.J., Weidlich, S., Labib, K., Hofmann, K., and Kulathu, Y. (2016). MINDY-1 is a Member of an Evolutionarily Conserved and Structurally Distinct New Family of Deubiquitinating Enzymes. *Mol. Cell* *63*, 146–155.
- Bingol, B., Tea, J.S., Phu, L., Reichelt, M., Bakalarski, C.E., Song, Q., Foreman, O., Kirkpatrick, D.S., and Sheng, M. (2014). The mitochondrial deubiquitinase USP30 opposes parkin-mediated mitophagy. *Nature* *510*, 370–375.
- Bondeson, D.P., Mares, A., Smith, I.E.D., Ko, E., Campos, S., Miah, A.H., Mulholland, K.E., Routly, N., Buckley, D.L., Gustafson, J.L., et al. (2015). Catalytic *in vivo* protein knockdown by small-molecule PROTACs. *Nat. Chem. Biol.* *11*, 611–U120.
- Boselli, M., Lee, B.H., Robert, J., Prado, M.A., Min, S.W., Cheng, C., Silva, M.C., Seong, C., Elsasser, S., Hatle, K.M., et al. (2017). An inhibitor of the proteasomal deubiquitinating enzyme USP14 induces tau elimination in cultured neurons. *J. Biol. Chem.* *292*, 19209–19225.
- Buckley, D.L., Raina, K., Darricarrere, N., Hines, J., Gustafson, J.L., Smith, I.E., Miah, A.H., Harling, J.D., and Crews, C.M. (2015). HaloPROTACs: Use of Small Molecule PROTACs to Induce Degradation of HaloTag Fusion Proteins. *ACS Chem. Biol.* *10*, 1831–1837.
- Buhimschi, A.D., Armstrong, H.A., Toure, M., Jaime-Figueroa, S., Chen, T.L., Lehman, A.M., Woyach, J.A., Johnson, A.J., Byrd, J.C., and Crews, C.M. (2018). Targeting the C481S Ibrutinib-Resistance Mutation in Bruton's Tyrosine Kinase Using PROTAC-Mediated Degradation. *Biochemistry-U.S.* *57*, 3564–3575.
- Burslem, G.M., Smith, B.E., Lai, A.C., Jaime-Figueroa, S., McQuaid, D.C., Bondeson, D.P., Toure, M., Dong, H.Q., Qian, Y.M., Wang, J., et al. (2018). The Advantages of Targeted Protein Degradation Over

Inhibition: An RTK Case Study. *Cell Chemical Biology* 25, 67-77.

Chan, A.I., McGregor, L.M., and Liu, D.R. (2015). Novel selection methods for DNA-encoded chemical libraries. *Curr. Opin. Chem. Biol.* 26, 55-61.

Chan, C.H., Morrow, J.K., Li, C.F., Gao, Y., Jin, G., Moten, A., Stagg, L.J., Ladbury, J.E., Cai, Z., Xu, D., et al. (2013). Pharmacological inactivation of Skp2 SCF ubiquitin ligase restricts cancer stem cell traits and cancer progression. *Cell* 154, 556-568.

Chauhan, D., Tian, Z., Nicholson, B., Kumar, K.G., Zhou, B., Carrasco, R., McDermott, J.L., Leach, C.A., Fulciniti, M., Kodrasov, M.P., et al. (2012). A small molecule inhibitor of ubiquitin-specific protease-7 induces apoptosis in multiple myeloma cells and overcomes bortezomib resistance. *Cancer Cell* 22, 345-358.

Chen, J., Dexheimer, T.S., Ai, Y., Liang, Q., Villamil, M.A., Inglesse, J., Maloney, D.J., Jadhav, A., Simeonov, A. and Zhuang, Z. (2011). Selective and cell-active inhibitors of the USP1/ UAF1 deubiquitinase complex reverse cisplatin resistance in non-small cell lung cancer cells. *Chem. Biol.* 18, 1390-1400.

Chu, T.T., Gao, N., Li, Q.Q., Chen, P.G., Yang, X.F., Chen, Y.X., Zhao, Y.F., and Li, Y.M. (2016). Specific Knockdown of Endogenous Tau Protein by Peptide-Directed Ubiquitin-Proteasome Degradation. *Cell Chem. Biol.* 23, 453-461.

Ciechanover, A. (2005). Proteolysis: from the lysosome to ubiquitin and the proteasome. *Nat. Rev. Mol. Cell Biol.* 6, 79-87.

Clague, M.J., Barsukov, I., Coulson, J.M., Liu, H., Rigden, D.J., and Urbe, S. (2013). Deubiquitylases from genes to organism. *Physiol. Rev.* 93, 1289-1315.

Coleman, K.G., and Crews, C.M. (2018). Proteolysis-Targeting Chimeras: Harnessing the Ubiquitin-Proteasome System to Induce Degradation of Specific Target Proteins. *Annual Review of Cancer Biology* 2, 41-58.

Colland, F., Formstecher, E., Jacq, X., Reverdy, C., Planquette, C., Conrath, S., Trouplin, V., Bianchi, J., Aushev, V.N., Camonis, J., et al. (2009). Small-molecule inhibitor of USP7/HAUSP ubiquitin protease stabilizes and activates p53 in cells. *Mol. Cancer Ther.* 8, 2286-2295.

Crew, A.P., Raina, K., Dong, H., Qian, Y., Wang, J., Vigil, D., Serebrenik, Y.V., Hamman, B.D., Morgan, A., Ferraro, C., et al. (2018). Identification and Characterization of Von Hippel-Lindau-Recruiting Proteolysis Targeting Chimeras (PROTACs) of TANK-Binding Kinase 1. *J. Med. Chem.* 61, 583-598.

D'Arcy, P., Brnjic, S., Olofsson, M.H., Fryknas, M., Lindsten, K., De Cesare, M., Perego, P., Sadeghi, B., Hassan, M., Larsson, R., et al. (2011). Inhibition of proteasome deubiquitinating activity as a new cancer therapy. *Nat. Med.* 17, 1636-1640.

de Poot, S.A.H., Tian, G., and Finley, D. (2017). Meddling with Fate: The Proteasomal Deubiquitinating Enzymes. *J. Mol. Biol.* 429, 3525-3545.

Deshaies, R.J., and Joazeiro, C.A. (2009). RING domain E3 ubiquitin ligases. *Annu. Rev. Biochem.* 78, 399-434.

Dexheimer, T.S., Rosenthal, A.S., Luci, D.K., Liang, Q., Villamil, M.A., Chen, J., Sun, H., Kerns, E.H., Simeonov, A., Jadhav, A., et al. (2014). Synthesis and structure-activity relationship studies of N-benzyl-2-phenylpyrimidin-4-amine derivatives as potent USP1/UAF1 deubiquitinase inhibitors with anticancer activity against nonsmall cell lung cancer. *J. Med. Chem.* 57, 8099-8110.

Farshi, P., Deshmukh, R.R., Nwankwo, J.O., Arkwright, R.T., Cvek, B., Liu, J., and Dou, Q.P. (2015). Deubiquitinases (DUBs) and DUB inhibitors: a patent review. *Expert Opin. Ther. Pat.* 25, 1191-1208.

Finley, D. (2009). Recognition and Processing of Ubiquitin-Protein Conjugates by the Proteasome. *Annual Review of Biochemistry* 78, 477-513.

Finley, D., and Chau, V. (1991). Ubiquitination. *Annu. Rev. Cell. Biol.* 7, 25-69.

Fraille, J.M., Quesada, V., Rodriguez, D., Freije, J.M., and Lopez-Otin, C. (2012). Deubiquitinases in cancer: new functions and therapeutic options. *Oncogene* 31, 2373-2388.

Gavory, G., O'Dowd, C.R., Helm, M.D., Flasz, J., Arkoudis, E., Dossang, A., Hughes, C., Cassidy, E., McClelland, K., Odrzywol, E., et al. (2018). Discovery and characterization of highly potent and selective allosteric USP7 inhibitors. *Nat. Chem. Biol.* 14, 118-125.

Gopinath, P., Ohayon, S., Nawatha, M., and Brik, A. (2016). Chemical and semisynthetic approaches to study and target deubiquitinases. *Chem. Soc. Rev.* 45, 4171-4198.

Gustafson, J.L., Neklesa, T.K., Cox, C.S., Roth, A.G., Buckley, D.L., Tae, H.S., Sundberg, T.B., Stagg, D.B., Hines, J., McDonnell, D.P., et al. (2015). Small-Molecule-Mediated Degradation of the Androgen Receptor through Hydrophobic Tagging. *Angew. Chem. Int. Ed. Engl.* 54, 9659-9662.

Harrigan, J.A., Jacq, X., Martin, N.M., and Jackson, S.P. (2018). Deubiquitylating enzymes and drug discovery: emerging opportunities. *Nat. Rev. Drug Discov.* 17, 57-77.

Henning, R.K., Varghese, J.O., Das, S., Nag, A., Tang, G., Tang, K., Sutherland, A.M., and Heath, J.R. (2016). Degradation of Akt using protein-catalyzed capture agents. *J. Pept. Sci.* 22, 196-200.

Hines, J., Gough, J.D., Corson, T.W., and Crews, C.M. (2013). Posttranslational protein knockdown coupled to receptor tyrosine kinase activation with phosphoPROTACs. *P. Natl. Acad. Sci. USA* 110, 8942-8947.

Huang, X., and Dixit, V.M. (2016). Drugging the undruggables: exploring the ubiquitin system for drug development. *Cell Res.* 26, 484-498.

Itoh, Y., Ishikawa, M., Naito, M., and Hashimoto, Y. (2010). Protein Knockdown Using Methyl Bestatin-Ligand Hybrid Molecules: Design and Synthesis of Inducers of Ubiquitination-Mediated Degradation of Cellular Retinoic Acid-Binding Proteins. *J. Am. Chem. Soc.* 132, 5820-5826.

Itoh, Y., Kitaguchi, R., Ishikawa, M., Naito, M., and Hashimoto, Y. (2011). Design, synthesis and biological evaluation of nuclear receptor-degradation inducers. *Bioorg. Med. Chem.* 19, 6768-6778.

Jiang, Y., Deng, Q., Zhao, H., Xie, M., Chen, L., Yin, F., Qin, X., Zheng, W., Zhao, Y., and Li, Z. (2018). Development of Stabilized Peptide-Based PROTACs against Estrogen Receptor alpha. *ACS Chem. Biol.* 13, 628-635.

Kapur, V., Peterson, L.F., Fang, D., Bornmann, W.G., Talpaz, M., and Donato, N.J. (2010). Deubiquitinase Inhibition by Small-Molecule WP1130 Triggers Aggresome Formation and Tumor Cell Apoptosis. *Cancer Res.* 70, 9265-9276.

Kategaya, L., Di Lello, P., Rouge, L., Pastor, R., Clark, K.R., Drummond, J., Kleinheinz, T., Lin, E., Upton, J.P., Prakash, S., et al. (2017). USP7 small-molecule inhibitors interfere with ubiquitin binding. *Nature* 550, 534-538.

Kluge, A.F., Lagu, B.R., Maiti, P., Jaleel, M., Webb, M., Malhotra, J., Mallat, A., Srinivas, P.A., and Thompson, J.E. (2018). Novel highly selective inhibitors of ubiquitin specific protease 30 (USP30) accelerate mitophagy. *Bioorg. Med. Chem. Letters* 28, 2655-2659.

Komander, D., Clague, M.J., and Urbe, S. (2009). Breaking the chains: structure and function of the deubiquitinases. *Nat. Rev. Mol. Cell Biol.* 10, 550-563.

Komander, D., and Rape, M. (2012). The ubiquitin code. *Annu. Rev. Biochem.* 81, 203-229.

Lai, A.C., and Crews, C.M. (2017). Induced protein degradation: an emerging drug discovery paradigm. *Nat. Rev. Drug Discov.* 16, 101-

114.

- Lai, A.C., Toure, M., Hellerschmied, D., Salami, J., Jaime-Figueroa, S., Ko, E., Hines, J., and Crews, C.M. (2016). Modular PROTAC Design for the Degradation of Oncogenic BCR-ABL. *Angew. Chem. Int. Ed. Engl.* *55*, 807-810.
- Lamberto, I., Liu, X., Seo, H.S., Schauer, N.J., Iacob, R.E., Hu, W., Das, D., Mikhailova, T., Weisberg, E.L., Engen, J.R., et al. (2017). Structure-Guided Development of a Potent and Selective Non-covalent Active-Site Inhibitor of USP7. *Cell Chem. Biol.* *24*, 1490-1500 e1411.
- Lee, B.H., Lee, M.J., Park, S., Oh, D.C., Elsasser, S., Chen, P.C., Gartner, C., Dimova, N., Hanna, J., Gygi, S.P., et al. (2010). Enhancement of proteasome activity by a small-molecule inhibitor of USP14. *Nature* *467*, 179-U163.
- Lee, B.H., Lu, Y., Prado, M.A., Shi, Y., Tian, G., Sun, S., Elsasser, S., Gygi, S.P., King, R.W., and Finley, D. (2016). USP14 deubiquitinates proteasome-bound substrates that are ubiquitinated at multiple sites. *Nature* *532*, 398-401.
- Li, J., Yakushi, T., Parlati, F., Mackinnon, A.L., Perez, C., Ma, Y., Carter, K.P., Colayco, S., Magnuson, G., Brown, B., et al. (2017). Capzimin is a potent and specific inhibitor of proteasome isopeptidase Rpn11. *Nat. Chem. Biol.* *13*, 486-493.
- Liang, J.R., Martinez, A., Lane, J.D., Mayor, U., Clague, M.J., and Urbe, S. (2015). USP30 deubiquitylates mitochondrial Parkin substrates and restricts apoptotic cell death. *Embo. Rep.* *16*, 618-627.
- Liu, N., Liu, C., Li, X., Liao, S., Song, W., Yang, C., Zhao, C., Huang, H., Guan, L., Zhang, P., et al. (2014a). A novel proteasome inhibitor suppresses tumor growth via targeting both 19S proteasome deubiquitinases and 20S proteolytic peptidases. *Sci. Rep.* *4*, 5240.
- Liu, N.N., Li, X.F., Huang, H.B., Zhao, C., Liao, S.Y., Yang, C.S., Liu, S.T., Song, W.B., Lu, X.Y., Lan, X.Y., et al. (2014b). Clinically used antirheumatic agent auranofin is a proteasomal deubiquitinase inhibitor and inhibits tumor growth. *Oncotarget* *5*, 5453-5471.
- Liu, Y.C., Lashuel, H.A., Choi, S., Xing, X.C., Case, A., Ni, J., Yeh, L.A., Cuny, G.D., Stein, R.L., and Lansbury, P.T. (2003). Discovery of inhibitors that elucidate the role of UCH-L1 activity in the H1299 lung cancer cell line. *Chem. Biol.* *10*, 837-846.
- Long, M.J., Gollapalli, D.R., and Hedstrom, L. (2012). Inhibitor mediated protein degradation. *Chem. Biol.* *19*, 629-637.
- Lu, M., Liu, T., Jiao, Q., Ji, J., Tao, M., Liu, Y., You, Q., and Jiang, Z. (2018). Discovery of a Keap1-dependent peptide PROTAC to knockdown Tau by ubiquitination-proteasome degradation pathway. *Eur. J. Med. Chem.* *146*, 251-259.
- Manasanch, E.E., and Orlowski, R.Z. (2017). Proteasome inhibitors in cancer therapy. *Nat. Rev. Clin. Oncol.* *14*, 417.
- Mistry, H., Hsieh, G., Buhrlage, S.J., Huang, M., Park, E., Cuny, G.D., Galinsky, I., Stone, R.M., Gray, N.S., D'Andrea, A.D., et al. (2013). Small-molecule inhibitors of USP1 target ID1 degradation in leukemic cells. *Mol. Cancer Ther.* *12*, 2651-2662.
- Neklesa, T.K., Tae, H.S., Schnekloth, A.R., Stulberg, M.J., Corson, T.W., Sundberg, T.B., Raina, K., Holley, S.A., and Crews, C.M. (2011). Small-molecule hydrophobic tagging-induced degradation of HaloTag fusion proteins. *Nat. Chem. Biol.* *7*, 538-543.
- Neklesa, T.K., Winkler, J.D., and Crews, C.M. (2017). Targeted protein degradation by PROTACs. *Pharmacol. Ther.* *174*, 138-144.
- Ohoka, N., Nagai, K., Hattori, T., Okuhira, K., Shibata, N., Cho, N., and Naito, M. (2014). Cancer cell death induced by novel small molecules degrading the TACC3 protein via the ubiquitin-proteasome pathway. *Cell Death Dis.* *5*.
- Popovic, D., Vucic, D., and Dikic, I. (2014). Ubiquitination in disease pathogenesis and treatment. *Nat. Med.* *20*, 1242-1253.
- Raina, K., Lu, J., Qian, Y.M., Altieri, M., Gordon, D., Rossi, A.M.K., Wang, J., Chen, X., Dong, H.Q., Siu, K., et al. (2016). PROTAC-induced BET protein degradation as a therapy for castration-resistant prostate cancer. *P. Natl. Acad. Sci. USA* *113*, 7124-7129.
- Rask-Andersen, M., Almen, M.S., and Schioth, H.B. (2011). Trends in the exploitation of novel drug targets. *Nat. Rev. Drug Discov.* *10*, 579-590.
- Reverdy, C., Conrath, S., Lopez, R., Planquette, C., Atmanene, C., Collura, V., Harpon, J., Battaglia, V., Vivat, V., Sippl, W., et al. (2012). Discovery of specific inhibitors of human USP7/HAUSP deubiquitinating enzyme. *Chem. Biol.* *19*, 467-477.
- Richardson, P.G., Sonneveld, P., Schuster, M.W., Irwin, D., Stadtmauer, E.A., Facon, T., Harousseau, J.L., Ben-Yehuda, D., Lonial, S., Goldschmidt, H., et al. (2005). Bortezomib or high-dose dexamethasone for relapsed multiple myeloma. *N. Engl. J. Med.* *352*, 2487-2498.
- Robb, C.M., Contreras, J.I., Kour, S., Taylor, M.A., Abid, M., Sonawane, Y.A., Zahid, M., Murry, D.J., Natarajan, A., and Rana, S. (2017). Chemically induced degradation of CDK9 by a proteolysis targeting chimera (PROTAC). *Chem. Commun. (Camb.)* *53*, 7577-7580.
- Rodriguez-Gonzalez, A., Cyrus, K., Salcius, M., Kim, K., Crews, C.M., Deshaies, R.J., and Sakamoto, K.M. (2008). Targeting steroid hormone receptors for ubiquitination and degradation in breast and prostate cancer. *Oncogene* *27*, 7201-7211.
- Ross, C.A., and Poirier, M.A. (2004). Protein aggregation and neurodegenerative disease. *Nat. Med.* *10 Suppl*, S10-17.
- Sacco, J.J., Coulson, J.M., Clague, M.J., and Urbe, S. (2010). Emerging roles of deubiquitinases in cancer-associated pathways. *IUBMB Life* *62*, 140-157.
- Sakamoto, K.M., Kim, K.B., Kumagai, A., Mercurio, F., Crews, C.M., and Deshaies, R.J. (2001). Protacs: Chimeric molecules that target proteins to the Skp1-Cullin-F box complex for ubiquitination and degradation. *P. Natl. Acad. Sci. USA* *98*, 8554-8559.
- Salami, J., and Crews, C.M. (2017). Waste disposal-An attractive strategy for cancer therapy. *Science* *355*, 1163-1167.
- Schiedel, M., Herp, D., Hammelmann, S., Swyter, S., Lehotzky, A., Robaa, D., Olah, J., Ovadi, J., Sippl, W., and Jung, M. (2018). Chemically induced degradation of sirtuin 2 (Sirt2) by a proteolysis targeting chimera (PROTAC) based on sirtuin rearranging ligands (SirReals). *J. Med. Chem.* *61*, 482-491.
- Schnekloth, A.R., Puchault, M., Tae, H.S., and Crews, C.M. (2008). Targeted intracellular protein degradation induced by a small molecule: En route to chemical proteomics. *Bioorg. Med. Chem. Lett.* *18*, 5904-5908.
- Schnekloth, J.S., Jr., Fonseca, F.N., Koldobskiy, M., Mandal, A., Deshaies, R., Sakamoto, K., and Crews, C.M. (2004). Chemical genetic control of protein levels: selective *in vivo* targeted degradation. *J. Am. Chem. Soc.* *126*, 3748-3754.
- Shangary, S., Qin, D., McEachern, D., Liu, M., Miller, R.S., Qiu, S., Nikolovska-Coleska, Z., Ding, K., Wang, G., Chen, J., et al. (2008). Temporal activation of p53 by a specific MDM2 inhibitor is selectively toxic to tumors and leads to complete tumor growth inhibition. *Proc. Natl. Acad. Sci. USA* *105*, 3933-3938.
- Stewart, A.K., Rajkumar, S.V., Dimopoulos, M.A., Masszi, T., Spicka, I., Oriol, A., Hajek, R., Rosinol, L., Siegel, D.S., Mihaylov, G.G., et al. (2015). Carfilzomib, lenalidomide, and dexamethasone for relapsed multiple myeloma. *N. Engl. J. Med.* *372*, 142-152.
- Tian, X., Isamidinova, N.S., Peroutka, R.J., Goldenberg, S.J., Mattem, M.R., Nicholson, B., and Leach, C. (2011). Characterization of selective ubiquitin and ubiquitin-like protease inhibitors using a fluorescence-based multiplex assay format. *Assay. Drug Dev. Technol.*

9, 165-173.

Tomoshige, S., Naito, M., Hashimoto, Y., and Ishikawa, M. (2015). Degradation of HaloTag-fused nuclear proteins using bestatin-HaloTag ligand hybrid molecules. *Org. Biomol. Chem.* *13*, 9746-9750.

Turnbull, A.P., Ioannidis, S., Krajewski, W.W., Pinto-Fernandez, A., Heride, C., Martin, A.C.L., Tonkin, L.M., Townsend, E.C., Buker, S.M., Lancia, D.R., et al. (2017). Molecular basis of USP7 inhibition by selective small-molecule inhibitors. *Nature* *550*, 481-486.

Vassilev, L.T., Vu, B.T., Graves, B., Carvajal, D., Podlaski, F., Filipovic, Z., Kong, N., Kammlott, U., Lukacs, C., Klein, C., et al. (2004). *In Vivo* Activation of the p53 Pathway by Small-Molecule Antagonists of MDM2. *303*, 844-848.

Wang, X., D'Arcy, P., Caulfield, T.R., Paulus, A., Chitta, K., Mohanty, C., Gullbo, J., Chanan-Khan, A., and Linder, S. (2015). Synthesis and evaluation of derivatives of the proteasome deubiquitinase inhibitor b-AP15. *Chem. Biol. Drug Des.* *86*, 1036-1048.

Wang, X., Feng, S., Fan, J., Li, X., Wen, Q., and Luo, N. (2016a). New strategy for renal fibrosis: Targeting Smad3 proteins for ubiquitination and degradation. *Biochem. Pharmacol.* *116*, 200-209.

Wang, X., Mazurkiewicz, M., Hillert, E.K., Olofsson, M.H., Pierrou, S., Hillertz, P., Gullbo, J., Selvaraju, K., Paulus, A., Akhtar, S., et al. (2016b). The proteasome deubiquitinase inhibitor VLX1570 shows selectivity for ubiquitin-specific protease-14 and induces apoptosis of multiple myeloma cells. *Sci Rep-Uk* *6*, 26979.

Weathington, N.M., and Mallampalli, R.K. (2014). Emerging therapies targeting the ubiquitin proteasome system in cancer. *J. Clin. Invest.* *124*, 6-12.

Weinstock, J., Wu, J., Cao, P., Kingsbury, W.D., McDermott, J.L., Kodrasov, M.P., McKelvey, D.M., Suresh Kumar, K.G., Goldenberg, S.J., Mattern, M.R., et al. (2012). Selective Dual Inhibitors of the Cancer-Related Deubiquitylating Proteases USP7 and USP47. *ACS Med. Chem. Lett.* *3*, 789-792.

Wilkinson, K.D. (1997). Regulation of ubiquitin-dependent processes by deubiquitinating enzymes. *FASEB J.* *11*, 1245-1256.

Winter, G.E., Buckley, D.L., Paulk, J., Roberts, J.M., Souza, A., Dhe-Paganon, S., and Bradner, J.E. (2015). Phthalimide conjugation as a strategy for *in vivo* target protein degradation. *Science* *348*, 1376-1381.

Xie, T., Lim, S.M., Westover, K.D., Dodge, M.E., Ercan, D., Ficarro, S.B., Udayakumar, D., Gurbani, D., Tae, H.S., Riddle, S.M., et al. (2014). Pharmacological targeting of the pseudokinase Her3. *Nat. Chem. Biol.* *10*, 1006-1012.

Yang, K., Song, Y., Xie, H., Wu, H., Wu, Y.T., Leisten, E.D., and Tang, W. (2018). Development of the first small molecule histone deacetylase 6 (HDAC6) degraders. *Bioorg. Med. Chem. Lett.* *28*, 2493-2497.

Yue, W., Chen, Z., Liu, H., Yan, C., Chen, M., Feng, D., Yan, C., Wu, H., Du, L., Wang, Y., et al. (2014). A small natural molecule promotes mitochondrial fusion through inhibition of the deubiquitinase USP30. *Cell Res.* *24*, 482-496.

Zengerle, M., Chan, K.H., and Ciulli, A. (2015). Selective Small Molecule Induced Degradation of the BET Bromodomain Protein BRD4. *ACS Chem. Biol.* *10*, 1770-1777.

Zhang, C., Han, X.R., Yang, X., Jiang, B., Liu, J., Xiong, Y., and Jin, J. (2018). Proteolysis Targeting Chimeras (PROTACs) of Anaplastic Lymphoma Kinase (ALK). *Eur. J. Med. Chem.* *151*, 304-314.

Zhou, B., Hu, J., Xu, F., Chen, Z., Bai, L., Fernandez-Salas, E., Lin, M., Liu, L., Yang, C.Y., Zhao, Y., et al. (2018). Discovery of a Small-Molecule Degradator of Bromodomain and Extra-Terminal (BET) Proteins with Picomolar Cellular Potencies and Capable of Achieving Tumor Regression. *J. Med. Chem.* *61*, 462-481.

Zhou, P. (2005). Targeted protein degradation. *Curr. Opin. Chem. Biol.* *9*, 51-55.