



Published in final edited form as:

*Braz J Infect Dis.* 2012 ; 16(3): 226–231.

## Prior history of Sexually transmitted diseases in women living with AIDS in São Paulo, Brazil

Valdir Monteiro Pinto<sup>a,b,\*</sup>, Mariza Vono Tancredi<sup>a</sup>, Jonathan Eric Golub<sup>c</sup>, Ariane de Castro Coelho<sup>a</sup>, Antonio Tancredi Neto<sup>d</sup>, and Angelica Espinosa Miranda<sup>b</sup>

<sup>a</sup>Centro de Referência e Treinamento em Doenças Sexualmente Transmissíveis/AIDS (CRT–DST/AIDS), SP, Brazil

<sup>b</sup>Universidade Federal do Espírito Santo (UFES), ES, Brazil

<sup>c</sup>Johns Hopkins University, Baltimore, USA

<sup>d</sup>Instituto do Coração (InCor), Hospital das Clínicas, Faculdade de Medicina da Universidade de São Paulo (FMUSP), SP, Brazil

### Abstract

**Objectives:** To describe the epidemiological profile, risk behaviors, and the prior history of Sexually transmitted diseases (STDs) in women living with acquired immunodeficiency syndrome (AIDS).

**Methods:** Cross-sectional study, performed at the Centro de Referência e Treinamento em DST/AIDSs of São Paulo. The social, demographic, behavioral, and clinical data such as age, schooling, marital status, age at first Sexual intercourse, number of Sexual partners, parity, use of drugs, time of HIV diagnosis, CD4 count, and viral load determination were abstracted from the medical records of women living with AIDS who had gynecological consultation scheduled in the period from June 2008 to May 2009.

**Results:** Out of 710 women who were scheduled to a gynecological consultation during the period of the study, 598 were included. Previous STD was documented for 364 (60.9%; 95% CI: 56.9%–64.8%) women. The associated factors with previous STDs and their respective risks were: human development index (HDI) < 0.50 (OR<sub>aj</sub> = 5.5; 95% CI: 2.8–11.0); non-white race (OR<sub>aj</sub> = 5.2; 95% CI: 2.5–11.0); first Sexual intercourse at or before 15 years of age (OR<sub>aj</sub> = 4.4; 95% CI: 2.3–8.3); HIV infection follow-up time of nine years or more (OR<sub>aj</sub> = 4.2; 95% CI: 2.3–7.8); number of Sexual partners during the entire life between three and five partners (OR<sub>aj</sub> = 2.2; 95% CI: 1.1–4.6), and six or more Sexual partners (OR<sub>aj</sub> = 3.9; 95% CI: 1.9–8.0%); being a Sex worker (OR<sub>aj</sub> = 1.9; 95% CI: 1.1–3.1).

---

Este é um artigo Open Access sob a licença de CC BY-NC-ND

\*Corresponding author at: rua santa cruz, 81, São Paulo, sP, 04121-000, Brazil, vmpinto@usp.br (Valdir Monteiro Pinto).

Conflict of interest

All authors declare to have no conflict of interest.

**Conclusions:** A high prevalence of a prior history of STDs in the studied population was found. It is essential to find better ways to access HIV infection prevention, so that effective interventions can be more widely implemented.

### Keywords

Sexually transmitted diseases; Women; HIV; Behavior

---

### Introduction

One reason for the increase of Sexually transmitted diseases (STDs) in many developing countries is the lack of access to effective and reliable health care services.<sup>1</sup> factors such as being young and Sexually active, urban migration with sociocultural changes, multiple Sexual partnerships without the use of condoms, and high prevalence of resistance against antimicrobial drugs also contribute to this increase.<sup>2,3</sup>

The analysis of studies performed in several countries<sup>4</sup> showed that people with STDs, even the non-ulcerative kind, presented a risk three to 10 times higher of being infected by the human immunodeficiency virus (HIV), depending on the STD type and etiology. People who live with HIV/AIDS (PLHA) have a high rate of prior STDs<sup>5,6</sup> and, if PLHA acquire any form of STD, the HIV viral load in the genital secretions increases,<sup>7,8</sup> causing its infectivity also to increase considerably. Among ulcerative STDs, it has been recently shown that genital herpes can be considered the main co-factor for the higher proportion of new HIV infections.<sup>9</sup>

In heteroSexual partners, the higher the HIV plasma concentration the higher the risk of HIV transmission,<sup>10</sup> and a study of HIV infected men with urethritis-associated STD showed a HIV average concentration in the seminal plasma eight times higher.<sup>11</sup> However, some studies showed no correlation between the HIV viral load in the plasma and in the semen.<sup>12</sup>

Some STDs, as human papilloma virus (HPV) infection, generally present a higher prevalence among HIV-infected women when compared to HIV-negative women,<sup>13-16</sup> and this persistent infection constitutes a higher risk for the development of cervical intraepithelial neoplasia (CIN).<sup>17</sup>

According to the Brazilian Ministry of Health, 544,846 cases of AIDS were reported from 1980 to June 2009, and HIV infection is having an increasing impact on reproductive health in Brazil. During this time, the male to female ratio has been progressively reduced from 26.7:1 in 1985 to 1.5:1 currently.<sup>18</sup> Although the number of accumulated cases among males is higher than in females, women represent the population where the fastest epidemic growth is observed in the country.<sup>18</sup>

Women experience different constraints for the exercise of Sexuality, making it difficult for them to incorporate protection practices, and health care services are not always able to deal with this situation, which in turn, increases their vulnerability.<sup>19</sup> since women are asymptomatic for most STDs, the percentage in which these infections are involved in the HIV transmission is unknown.

The propose of this study was to describe the epidemiological profile, risk behavior, and frequency of prior STDs among women living with AIDS, aiming at collecting data that could be used in the implementation of prevention and assistance programs for women.

## Methodology

This cross-sectional study was performed in the Centro de Referência e Treinamento em DST/AIDSs in the city of São Paulo, Brazil (CRT-DST/AIDSs). Out of approximately 4,000 HIV/AIDS patients receiving care at this center 1,100 were women.

Data were abstracted from medical records of HIV-infected women. Women scheduled for a gynecological appointment in the outpatient gynecological clinics of the reference center between July 1, 2008 and May 31, 2009 were included in the study.

Social, demographic, behavioral, and clinical characteristics such as age, educational level, marital status, age at first Sexual intercourse, number of Sexual partners, parity, use of drugs, time elapsed since the diagnosis of HIV infection, CD4<sub>+</sub> count, and HIV viral load determination, among others, were selected for analysis.

The study included women who used highly active antiretroviral drugs, and excluded those who did not have the results of CD4<sub>+</sub> count and/or HIV viral load, and those whose gynecological consultation had not been described.

The reports of previous STDs were analyzed, as well as the results of laboratory exams and of cervical oncotoc cytology. The variable race, self-reported, considered as black those women who declared themselves as black or brown. Educational level and the human development index (HDI) were used as indicators of social and economic level due to their association with several lifestyles characteristics. The data collected from the medical records were linked to the HDI database of the city of São Paulo for the year 2000, through the deterministic record linkage method, which made nominal identification between the two databases.

History of prior STDs (syphilis, herpes, gonorrhea, *Chlamydia*, *Trichomonas*, hepatitis b, and HPV) was considered as dependent variable for the purpose of this analysis. The risks of prior STDs were computed for all independent variables of interest mentioned above.

The clinical and epidemiological information and the laboratorial findings were coded and stored in a database created for this purpose. The statistical program STATA 10.0 was used for data storage and analysis.

The analysis was performed using data exploratory techniques to check the distribution patterns and trends of the main variables. Then, univariate analysis was performed to check the association between variables. The chi-square test ( $\chi^2$ ) was used to assess the difference between proportions and odds ratio calculated with the respective 95% confidence intervals. student's t-test and analysis of variance were used to assess differences between means. The variables were selected for stepwise logistic regression analysis based on a p-value equal to

or lower than 0.25 in the likelihood ratio test. The importance of the variables for the final model was assessed with the likelihood ratio test, considering  $p < 0.05$ .

The project was approved by the Ethics committee of Centro de Referência e Treinamento em DST/AIDSs of São Paulo.

## Results

A total of 598 women out of the 710 scheduled for the gynecological consultation in the study period were included. One-hundred and twelve (15.8%) were excluded for the following reasons: 33 (4.6%) women with no information about prior STDs; 29 (4.1%) did not have previous or current gynecological consultations; 23 (3.2%) had never been on antiretroviral drugs; 18 (2.5%) had a negative HIV serology; and nine (1.3%) had acquired HIV through vertical transmission (VT) and never had Sexual activity. Prior STDs was observed in 364 women (60.9%; 95% CI: 56.9%–64.8%).

The proportions of prior STDs were 44.0% (263/598) for HPV; 17.9% (107/598) for herpes; 6.2% (37/598) for syphilis; 3.0% (18/598) for *Trichomonas*; 2.2% (13/598) for hepatitis b; 0.8% (5/598) for gonorrhea; and 0.7% (4/598) for *Chlamydia*. It should be pointed out that 21.7% of the patients mentioned two or more prior STDs, in addition to HIV infection.

The socioeconomical, behavioral and clinical characteristics of the women are presented in Table 1. The majority of women was less than 40 years old (73.6%) at the time of HIV diagnosis, and most of them were white (67.2%). As to their educational level, 50% had completed more than 8 years of schooling, being 56.1% among the white women and 37.6% among the black women. More than 60% were not married and had no stable partner.

Table 2 describes the behavioral and clinical characteristics of the studied population, comparing women with and without a history of prior STDs. More than half of them (53.2%) began their Sexual life after 15 years of age; 35.1% mentioned six or more Sexual partners throughout life; and 2.7% declared themselves as Sex workers. A total of 43.6% had been diagnosed with HIV infection for nine or more years, varying from 1 to 22 years.

Use of intravenous drugs was admitted by 2.2% of the women; 13.2% reported being non-injectable drug users and, among them, 41.8% used more than one drug.

Factors associated to prior STDs included in the logistic regression model are presented in Table 3. These factors and their correspondent risks were: HDI of the place of residence  $< 0.50$  ( $OR_{aj} = 5.5$ ; 95% CI: 2.8–11.0); race other than white ( $OR_{aj} = 5.2$ ; 95% CI: 2.5–11.0); age of the first Sexual intercourse up to 15 years ( $OR_{aj} = 4.4$ ; 95% CI: 2.3–8.3); follow-up time of the HIV infection of nine years or more [ $OR_{aj} = 4.2$  (95% CI: 2.3–7.8)]; number of Sexual partners in lifetime between three and five ( $OR_{aj} = 2.2$ ; 95% CI: 1.1–4.6)], and six or more ( $OR_{aj} = 3.9$ ; 95% CI: 1.9–8.0)]; being a Sex worker ( $OR_{aj} = 1.9$ ; 95% CI: 1.1–3.1).

## Discussion

The purpose of this study was to describe the epidemiological profile, risk behaviors, and the frequency of prior STDs in women living with AIDS receiving care at the Centro de Referência e Treinamento em DST/AIDSs in São Paulo. A high prevalence of prior STDs was found in the population studied. In Brazil, there are few available studies that analyzed STD prevalence in PLHA.

Two studies were carried out in the state of PeRNambuco. One assessed STD prevalence in 399 PLHA in a sample made up of heteroSexuals (33%), homoSexuals (23%), and biSexuals (11%), 75% males, in which syphilis was the most frequent disease, with a prevalence of 8.8%, followed by 5.8% of genital herpes and 4.3% of genital candidiasis.<sup>20</sup> the second study<sup>21</sup> assessed genital infection in women, showed frequencies of HPV, *Chlamydia trachomatis*, and *Trichomonas vaginalis* of 20.0%, 2.2%, and 2.2%, respectively. Grinsztejn et al.<sup>22</sup> found a HPV prevalence of 48% in a sample of 634 HIV-infected women in Rio de Janeiro.

Studies conducted in Jamaica and taiwan<sup>5,6</sup> have shown rates of prior STDs in PLHA of 51.1% and 43.1%, respectively. In the present study, prior STD, in addition to presenting a higher proportion (60.9%), was significantly associated with the district of residence, race other than white, first Sexual intercourse under the age of 16, three or more Sexual partners in life, being a Sex worker, and having been diagnosed with the HIV infection for nine years or more. these results are supported by other studies in Brazil, showing that the AIDS epidemic in the country affects predominantly (80%) individuals at the lower socioeconomic level.<sup>16,23</sup>

The available data provide further evidence that other STDs facilitate HIV transmission through direct biological mechanisms, and that the early treatment of STDs should be a part of high-quality strategies for the prevention of HIV infection.<sup>24</sup> Perhaps a high percentage of the participants of the present study have been infected, or have transmitted the HIV infection, due to a concomitant STD, since the majority of STDs are asymptomatic in women.

The high proportion of women with history of prior HPV in this study (44.0%) draws the attention to the need of preventive actions in relation to gynecological cancers, since many studies have shown a strong association between HIV and HPV co-infection and the development of CIN and genital cancer.<sup>25–27</sup> there is evidence that HIV-infected women have significantly higher CIN rates and a higher probability to evolve to invasive carcinoma than non-infected women.<sup>28–31</sup>

The present study found a history of prior genital herpes in 17.9% of the women, a higher rate than that reported by Saxton et al. (13.7%), in Ukraine.<sup>32</sup> Abu-Raddad et al., showed that herpes simplex virus 2 (HsV-2) infection is a biological co-factor in HIV acquisition and transmission, and can facilitate the spread of HIV infection even among the population with a lower risk of infection who maintain stable Sexual partnerships.<sup>33</sup> History of prior of syphilis was observed in 6.2% of the sample, a similar rate to that found by Hutton-rose et

al. (7.3%),<sup>5</sup> although lower than that found by Saxton et al. (1.9%),<sup>32</sup> which may be explained by the lower age of their study population.

In the present study, history of prior *Chlamydia trachomatis* infection was low (0.7%). Perhaps due to the non-availability of routine diagnostic work-up for this infection in Brazil, and its asymptomatic character in up to 80% of the women. Its importance is not well known by the population. This fact must be considered in HIV acquisition, because of the high prevalence of this infection among the Brazilian female population according to a multicenter trial conducted by the Ministry of Health,<sup>34</sup> and another study with a nationally representative sample,<sup>35</sup> with prevalences of 9.4% and 9.8%, respectively.

The use of retrospective data is one of the limitations of this study. The temporality of the infections could not be assessed. However, both the good quality of the data of the medical records and the low rate of missing information may counteract this limitation. In attempting to give socially accepted answers, another limitation that may have occurred would be the inaccuracy of the information on the use of condoms, age at first Sexual intercourse, and number of Sexual partners.

Health care services for HIV positive women exist, and they are able to control successfully the infection and prevent the disease progression. Although this is promising, there is still much work to be done to identify innovative interventions that target the social, cultural, and environmental influences of the STD presence in this group. It is also necessary to provide better ways to access prevention programs of HIV infection, so that effective interventions can be more broadly used. Sexually active women need confidential, affordable, and supportive services to teach them how to protect themselves against diseases, including STDs and HIV infection.

Considering that STDs are among the most well established risk factors for HIV infection, public health programs should be enforced with the implementation of control actions and tracking, diagnosis, and early treatment strategies, thus avoiding complications, reducing morbidity, and enhancing the Sexual and reproductive health of the population.

## Acknowledgements

Supported by the National Institute of Health grant number 3D43tW000010-21S1 and National Institutes of Health grant AI06699 (JEG).

## REFERENCES

1. Dallabetta GA, Gerbase AC, Holmes KK. Problems, solutions, and challenges in syndromic management of Sexually transmitted diseases. *Sex Transm Infect.* 1998;74(Suppl 1):S1-11. [PubMed: 10023346]
2. Mayaud P, Mabey D. Approaches to the control of Sexually transmitted infections in developing countries: old problems and modern challenges. *Sex Transm Infect.* 2004;80:174-82. [PubMed: 15169997]
3. Ferreira W, Sardinha J, Schettini A, Ferreira C, Benzaken A. susceptibilidade de cepas de *Neisseria gonorrhoeae* aos antibióticos utilizados para o tratamento de uretrites e cervicites gonocócicas em ambulatório de DST de Manaus – brasil. *J bras Doenças Sex transm.* 2001;13(6):36-40.

4. Wasserheit JN. Epidemiological synergy: interrelationships between human immunodeficiency virus infection and other Sexually transmitted diseases. *Sex Transm Dis.* 1992;19:61–77. [PubMed: 1595015]
5. Hutton-Rose n, Blythe C, Ogbonna C, McGrowder D. The prevalence of other Sexually transmitted infections in confirmed HIV cases at a referral clinic in Jamaica. *J R Soc Promot Health.* 2008;128:242–7. [PubMed: 18814406]
6. He Lee, Ko NY Chang CM, Liu SY Ko Wc. trends in Sexually transmitted diseases and risk behaviors among HIV-infected patients at an outpatient clinic in southern Taiwan. *Sex Transm Dis.* 2010;37:86–93. [PubMed: 19901862]
7. Gray RH, Wawer MJ, Brookmeyer R, et al. Probability of HIV-1 transmission per coital act in monogamous, heterosexual, HIV-1 discordant couples in Rakai, Uganda. *Lancet.* 2001;357:1149–53. [PubMed: 11323041]
8. Ghys PD, Fransen K, Diallo MO, et al. The associations between cervicovaginal HIV shedding, Sexually transmitted diseases and immunosuppression in female Sex workers in Abidjan, Cote d'Ivoire. *AIDS.* 1997;11:F85–93. [PubMed: 9342059]
9. consultation on STD interventions for preventing HIV: what is the evidence? UnAIDS/WHO, Isbn 92-9173-137-4. Geneva, 2000 Available from: [http://www.who.int/hiv/pub/sti/who\\_hsi\\_2000\\_02.pdf](http://www.who.int/hiv/pub/sti/who_hsi_2000_02.pdf)
10. Donnell D, Baeten JM, Kiarie J, et al. HeteroSexual HIV-1 transmission after initiation of antiretroviral therapy: a prospective cohort analysis. *Lancet.* 2010;375:2092–8. [PubMed: 20537376]
11. Ms Cohen, Hoffman IF Royce RA, et al. reduction of concentration of HIV-1 in semen after treatment of urethritis: implications for prevention of Sexual transmission of HIV-1. *Lancet.* 1997;349:1868–73. [PubMed: 9217758]
12. Marcelin AG, Tubianar R, Lambert-Niclot S, et al. Detection of HIV-1 RNA in seminal plasma samples from treated patients with undetectable HIV-RNA in blood plasma. *AIDS.* 2008;22:1677–9. [PubMed: 18670231]
13. Ellerbrock tV Chiasson MA, Bush TJ, et al. Incidence of cervical squamous intraepithelial lesions in HIV-infected women. *JAMA.* 2000;283:1031–7. [PubMed: 10697063]
14. Mandelblatt JS, Fahs M, Garibaldi K, Senie RT, Petersen HB. Association between HIV infection and cervical neoplasia: implications for clinical care of women at risk for both conditions. *AIDS.* 1992;6:173–8. [PubMed: 1558715]
15. Palefsky J Human papillomavirus-related disease in people with HIV. *Curr Opin HIV AIDS.* 2009;4:52–6. [PubMed: 19339939]
16. Campos RR, Melo VH, Castillo DM, Nogueira CPF. Prevalência do papilomavirus humano e seus genótipos em mulheres portadoras e não-portadoras do vírus da imunodeficiência humana. *Rev Bras Ginecol Obstet.* 2005;27:248–56.
17. Heard I cervical disease and cancer in HIV positive women. recommendations for screening and diagnosis. *Med Wieku Rozwoj.* 2003;7:479–85. [PubMed: 15010558]
18. Ministério da Saúde (Brasil). Secretaria de Vigilância em Saúde. Departamento de DST, AIDS e Hepatites virais Boletim Epidemiológico - AIDS e Dst. Brasília: Ministério da saúde; 2009.
19. Villela WV, Pinto VM. Atenção às Dst em mulheres In: Compromissos do governo brasileiro com a plataforma conferência inteRNacional sobre população e desenvolvimento: rumos para Cairo +20. Brasília: Presidência da República. Secretaria Especial de Políticas para as Mulheres; 2010 p. 159–179.
20. Rodrigues EH, Abath FG. Sexually transmitted diseases in patients infected with HIV/AIDS in the state of PeRNambuco, Brazil. *Rev Soc Bras Med trop.* 2000;33:47–52. [PubMed: 10881118]
21. Henriques APC, Magalhães V, Tenório T. Infecções genitais em mulheres infectadas pelo vírus da imunodeficiência humana (HIV) acompanhadas no hospital das clínicas da UFPE. *newsLab.* 2005;72:90–102.
22. Grinsztejn B, Veloso VG, Levi JE, et al. factors associated with increased prevalence of human papillomavirus infection in a cohort of HIV-infected Brazilian women. *Int J Infect Dis.* 2009;13:72–80. [PubMed: 18632296]



23. Fonseca MG, Bastos FI. Twenty-five years of the AIDS epidemic in Brazil: principal epidemiological findings, 1980–2005. *Cad Saude Publica*. 2007;23(Suppl 3):S333–44. [PubMed: 17992340]
24. Flemming D, Wasserheit J. From epidemiological synergy to public health policy and practice: the contribution of other Sexually transmitted diseases to Sexual transmission of HIV infection. *Sex Transm Inf*. 1999;75:3–17.
25. Hawes SE, Critchlow CW, Faye Niang MA, et al. Increased risk of high-grade cervical squamous intraepithelial lesions and invasive cervical cancer among African women with human immunodeficiency virus type 1 and 2 infections. *J Infect Dis*. 2003;188:555–63. [PubMed: 12898443]
26. Heard I, Tassie JM, Schmitz V, Mandelbrot L, Kazatchkine MD, Orth G. Increased risk of cervical disease among human immunodeficiency virus infected women with severe immunosuppression and high human papillomavirus load. *Obstet Gynecol*. 2000;96:403–9. [PubMed: 10960634]
27. de Sanjose S, Valls I, Paz Cañadas M, et al. Human papillomavirus and human immunodeficiency virus infections as risk factors for cervix cancer in women prisoners. *Med clin (barc)*. 2000;115:81–4. [PubMed: 10965480]
28. Nyagol J, Leucci E, Onnis A, et al. The effects of HIV-1 Tat protein on cell cycle during cervical carcinogenesis. *cancer biol ther*. 2006;5:684–90. [PubMed: 16855377]
29. Gichangi PB, Bwayo J, Estambale B, et al. Impact of HIV infection on invasive cervical cancer in Kenyan women. *AIDS*. 2003;17(13):1963–1968. [PubMed: 12960829]
30. Clarke B, Chetty R. Postmodern cancer: the role of human immunodeficiency virus in uterine cervical cancer. *Mol Pathol*. 2002;55:19–24. [PubMed: 11836442]
31. Ng'andwe, Lowe JL, Richards PJ, Hause L, Wood C, Angeletti PC. The distribution of Sexually transmitted human papillomaviruses in HIV positive and negative patients in Zambia, Africa. *BMC Infect Dis*. 2007;7:77. [PubMed: 17634108]
32. Saxton J, Malyuta R, Semenko I, et al. Previous reproductive history and post-natal family planning among HIV-infected women in Ukraine. *Hum Reprod*. 2010;25:2366–73. [PubMed: 20643695]
33. Abu-Raddad LJ, Magaret As, Celum C, et al. Genital herpes has played a more important role than any other Sexually transmitted infection in driving HIV prevalence in Africa. *PLoS OnE*. 2008;3:e2230. [PubMed: 18493617]
34. Jalil EM, Pinto VM, Benzaken As, et al. Prevalência da infecção por clamídia e gonococo em gestantes de seis cidades brasileiras. *Rev Bras Ginecol Obstet*. 2008;30:614–9. [PubMed: 19219343]
35. Pinto VM, Szwarcwald CL, Baroni C, Stringari LL, Inocêncio LA, Miranda AE. Chlamydia trachomatis prevalence and risk behaviors in parturient women aged 15 to 24 in Brazil. *Sex Transm Dis*. 2011;38:957–61. [PubMed: 21934572]



Distribution of the sociodemographic characteristics of women living with AIDS according to history of prior STD at the Centro de Referência e Treinamento em DST/AIDS, São Paulo, Brazil, 2008 to 2009

**Table 1 -**

| Characteristics               | With STD (n = 364) |      | Without STD (n = 234) |      | Total (598) |      | P       |
|-------------------------------|--------------------|------|-----------------------|------|-------------|------|---------|
|                               | n                  | %    | n                     | %    | n           | %    |         |
| Age at AIDS diagnosis         |                    |      |                       |      |             |      |         |
| up to 40 years                | 265                | 72.8 | 175                   | 74.8 | 440         | 73.6 | 0.591   |
| > 40 years                    | 99                 | 27.2 | 59                    | 25.2 | 158         | 26.4 |         |
| Educational level (years)     |                    |      |                       |      |             |      |         |
| None                          | 4                  | 1.1  | 2                     | 0.9  | 6           | 1.0  | 0.956   |
| 1 to 4 years                  | 29                 | 8.0  | 19                    | 8.1  | 48          | 8.0  |         |
| 5 to 8 years                  | 151                | 41.5 | 91                    | 38.9 | 242         | 40.5 |         |
| 9 to 11 years                 | 135                | 37.1 | 87                    | 37.2 | 222         | 37.1 |         |
| 12 or more                    | 42                 | 11.5 | 33                    | 14.1 | 75          | 12.5 |         |
| Unknown                       | 3                  | 0.8  | 2                     | 0.9  | 5           | 0.8  |         |
| Race (self-referred)          |                    |      |                       |      |             |      |         |
| White                         | 202                | 55.5 | 200                   | 85.5 | 402         | 67.2 | < 0.001 |
| Not white                     | 162                | 44.5 | 32                    | 13.7 | 194         | 32.4 |         |
| Unknown                       | 0                  | 0.0  | 2                     | 0.9  | 2           | 0.3  |         |
| Marital status*               |                    |      |                       |      |             |      |         |
| Single                        | 102                | 28.0 | 63                    | 26.9 | 165         | 27.6 | 0.909   |
| Married/lives with partner    | 128                | 35.2 | 76                    | 32.5 | 204         | 34.1 |         |
| Separated/divorced            | 62                 | 17.0 | 46                    | 19.7 | 108         | 18.1 |         |
| Widowed                       | 69                 | 19.1 | 47                    | 20.1 | 116         | 19.4 |         |
| Unknown                       | 3                  | 0.8  | 2                     | 0.9  | 5           | 0.8  |         |
| HDI of the residence district |                    |      |                       |      |             |      |         |
| > 0.550                       | 172                | 47.3 | 213                   | 91.0 | 385         | 64.4 | < 0.001 |
| 0–0.550                       | 192                | 52.7 | 21                    | 9.0  | 213         | 35.6 |         |

STD, sexually transmitted diseases; HDI, human development index;

\* five women with unknown data were excluded.

Distribution of behavioral characteristics of women living with AIDS according to history of prior STD at the Centro de Referência e Treinamento em DST/AIDS, São Paulo, Brazil, 2008 to 2009

**Table 2 -**

| Characteristics                 | With STD (n = 364) |      | Without STD (n = 234) |       | Total (598) |      | P       |
|---------------------------------|--------------------|------|-----------------------|-------|-------------|------|---------|
|                                 | n                  | %    | n                     | %     | n           | %    |         |
| Age at first sexual intercourse |                    |      |                       |       |             |      | < 0.001 |
| Older than 15 years             | 138                | 37.9 | 180                   | 76.9  | 318         | 53.2 |         |
| Up to 15 years                  | 190                | 52.2 | 33                    | 14.1  | 223         | 37.3 |         |
| Unknown                         | 36                 | 9.9  | 21                    | 9.0   | 57          | 9.5  |         |
| No. of partners in lifetime     |                    |      |                       |       |             |      | < 0.001 |
| 1 to 2                          | 34                 | 9.3  | 68                    | 29.1  | 102         | 17.1 |         |
| 3 to 5                          | 104                | 28.6 | 38                    | 16.2  | 142         | 23.7 |         |
| 6 or more                       | 180                | 49.5 | 30                    | 12.8  | 210         | 35.1 |         |
| Unknown                         | 46                 | 12.6 | 98                    | 41.9  | 144         | 24.1 |         |
| Use of non-intravenous drugs    |                    |      |                       |       |             |      | 0.001   |
| No                              | 299                | 82.1 | 218                   | 93.2  | 517         | 86.5 |         |
| Yes                             | 63                 | 17.3 | 16                    | 6.8   | 79          | 13.2 |         |
| Unknown                         | 2                  | 0.5  | 0                     | 0.0   | 2           | 0.3  |         |
| Use of intravenous drugs        |                    |      |                       |       |             |      | 0.007   |
| No                              | 349                | 95.9 | 234                   | 100.0 | 583         | 97.5 |         |
| Yes                             | 13                 | 3.6  | 0                     | 0.0   | 13          | 2.2  |         |
| Unknown                         | 2                  | 0.5  | 0                     | 0.0   | 2           | 0.3  |         |

STD, Sexually transmitted diseases.

Table 3 -

Univariate and multivariate analysis of the factors associated with history of prior STD in women living with AIDS at the Centro de Referência e Treinamento em DST/AIDS, São Paulo, Brazil, 2008 to 2009

| Characteristics                  | OR <sub>br</sub> | 95%CI (OR <sub>br</sub> ) | p      | OR <sub>aj</sub> | 95%CI (OR <sub>aj</sub> ) | p      |
|----------------------------------|------------------|---------------------------|--------|------------------|---------------------------|--------|
| HDI of the residence district    |                  |                           |        |                  |                           |        |
| > 0.550                          | 1                | -                         | -      | 1                | -                         | -      |
| 0-0.550                          | 11.3             | 6.9-18.5                  | <0.001 | 5.5              | 2.8-11.0                  | <0.001 |
| Race (self-referred)             |                  |                           |        |                  |                           |        |
| White                            | 1                | -                         | -      | 1                | -                         | -      |
| Not white                        | 5.0              | 3.3-7.7                   | <0.001 | 5.2              | 2.5-11.0                  | <0.001 |
| Age at first sexual intercourse  |                  |                           |        |                  |                           |        |
| Older than 15 years              | 1                | -                         | -      | 1                | -                         | -      |
| Up to 15 years                   | 7.5              | 4.9-11.5                  | <0.001 | 4.4              | 2.3-8.3                   | <0.001 |
| Time of HIV diagnosis (in years) |                  |                           |        |                  |                           |        |
| 1 to 8 years                     | 1                | -                         | -      | 1                | -                         | -      |
| 9 years and more                 | 4.4              | 3.0-6.3                   | <0.001 | 4.2              | 2.3-7.8                   | <0.001 |
| No. of sexual partners in life   |                  |                           |        |                  |                           |        |
| 1 to 2                           | 1                | -                         | -      | 1                | -                         | -      |
| 3 to 5                           | 5.5              | 3.1-9.5                   | <0.001 | 2.2              | 1.1-4.6                   | 0.045  |
| 6 and more                       | 12.0             | 6.8-21.1                  | <0.001 | 3.9              | 1.9-8.0                   | <0.001 |

STD, sexually transmitted diseases; HDI, human development index.