



Published in final edited form as:

Mov Disord. 2017 March ; 32(3): 332–345. doi:10.1002/mds.26944.

Overcoming the Divide Between Ataxias and Spastic Paraplegias: Shared Phenotypes, Genes, and Pathways

Matthis Synofzik, MD^{1,2,*} and Rebecca Schüle, MD^{1,2}

¹Department of Neurodegeneration, Hertie Institute for Clinical Brain Research (HIH), University of Tübingen, Tübingen, Germany

²German Center for Neurodegenerative Diseases (DZNE), Tübingen, Germany

Abstract

Autosomal-dominant spinocerebellar ataxias, autosomal-recessive spinocerebellar ataxias, and hereditary spastic paraplegias have traditionally been designated in separate clinicogenetic disease classifications. This classification system still largely frames clinical thinking and genetic workup in clinical practice. Yet, with the advent of next-generation sequencing, phenotypically unbiased studies have revealed the limitations of this classification system. Various genes (eg, *SPG7*, *SYNE1*, *PNPLA6*) traditionally rooted in either the ataxia or hereditary spastic paraplegia classification system have now been shown to cause ataxia on the one end of the disease continuum and hereditary spastic paraplegia on the other. Other genes such as *GBA2* and *KIF1C* were almost simultaneously published as both a hereditary spastic paraplegia and an ataxia gene. The variability and fluidity of observed phenotypes along the ataxia-spasticity spectrum warrants a rethinking of the traditional classification system. We propose to replace this divisive diagnosis-driven ataxia and hereditary spastic paraplegia classification system by a descriptive, unbiased approach of *modular phenotyping*. This approach is also open to expansion of the phenotype beyond ataxia and spasticity, which often occur as part of broader multisystem neuronal dysfunction. The concept of a continuous ataxia-spasticity disease spectrum is further supported by ataxias and hereditary spastic paraplegias sharing not only overlapping phenotypes and underlying genes, but also common cellular pathways and disease mechanisms. This suggests a shared vulnerability of cerebellar and corticospinal neurons for common pathophysiological processes. It might be this *mechanistic* overlap that drives their *clinical* overlap. A mechanistically inspired classification system will help to pave the way for mechanism-based strategies for drug development.

Keywords

spinocerebellar ataxia; recessive ataxia; hereditary spastic paraplegia; genetics; classification; pathway; molecular

*Correspondence to: Dr. Matthis Synofzik, Department of Neurodegenerative Diseases, Hertie-Institute for Clinical Brain Research, University of Tübingen, Hoppe-Seyler-Str. 3, 72076 Tübingen, Germany; matthis.synofzik@uni-tuebingen.de.

Supporting Data

Additional Supporting Information may be found in the online version of this article at the publisher's website.

Hereditary spinocerebellar ataxias and hereditary spastic paraplegias (HSPs) each define a genetically heterogeneous group of rare degenerative disorders characterized by progressive degeneration of the cerebellar Purkinje cells and spinocerebellar tracts (ataxias) and corticospinal tracts (HSPs), respectively. They were traditionally designated in separate clinicogenetic disease classifications, according to the predominant disease phenotype on first gene locus description and to the mode of inheritance:

1. Neurodegenerative diseases first conceptualized as *autosomal-dominant spinocerebellar ataxias (SCAs)* were classified in the SCA classification, which entails 43 SCA subtypes to date.¹
2. Neurodegenerative diseases first described as *autosomal-recessive spinocerebellar ataxias (SCARs)* were classified in the SCAR classification, comprising 24 subtypes.¹ This SCAR classification is partly paralleled and duplicated, yet with different numbers, by another autosomal-recessive cerebellar ataxia (ARCA) classification, the ARCA classification.¹
3. Neurodegenerative diseases first reported with *spastic paraplegia* were classified in the spastic paraplegia gene (SPG) classification irrespective of mode of inheritance. Seventy-eight distinct SPG loci are currently reported by OMIM.¹
4. A small number of genes presenting with combined ataxia and spasticity were somewhat arbitrarily also categorized as *spastic ataxia* genes (SPAX/SAX). The 7 loci listed in these classifications are mostly duplicate entries also contained in either the HSP or ataxia classification systems.

Each of these classification systems bears in itself the same problems known from similar classification systems of other movement disorders (for a broader discussion, see the analysis by the International Parkinson and Movement Disorder Society Task Force²). These include (1) erroneously assigned loci, (2) duplicated loci, (3) missing symbols or loci, and (4) unconfirmed loci and genes.² For example, some recessive ataxias are not contained in the SCAR or the ARCA list (eg, Friedreich's ataxia or AOA1), and some recessive ataxias are listed only in one of them (eg, AOA2 only in SCAR classification). Moreover, some dominant ataxias can also be inherited in a recessive manner and vice versa (*GRID2*,³ *AFG3L2*,⁴ *SPTBN2*), making it difficult to designate them as either on the SCA or the SCAR/ARCA list (or both). Most important, the systematic value of each of these classification systems is also very limited. Numbers in the SCA/ARCA/SCAR/SPG lists are assigned in the order in which the disease was identified (initially by linkage analysis and more recently by gene discovery). Yet these numbers do not carry any systematic information in themselves that might help to facilitate clinical diagnostics, to understand the disease etiology, or to devise treatment strategies.

In addition to these shortcomings, each of these classification systems carries in itself the classification systems for ataxias and HSPs that also bear a particular limitation when seen together. They suggest a conceptual and classificatory divide between ataxias and HSPs, when in fact there exists a large phenotypic, genetic, and pathophysiological overlap. This intersection between ataxias and HSPs has been increasingly acknowledged throughout the last decade,⁶ but its appreciation was notably facilitated by recent next-generation

sequencing (NGS) studies. Classic clinical and genetic strategies were largely constrained by preexisting clinical conceptions, classifications, and diagnostic workflows, leading to confirmation bias in genotype-phenotype correlation studies. In contrast, NGS has facilitated gene discoveries and phenotypic classifications unbiased from prior clinical and diagnostic preconceptions. This development has led to further weakening and even partial removal of the defined boundaries between ataxias and HSPs. As we show here, recent NGS and related genomic studies have demonstrated

1. a rapidly increasing number of both *novel* genes and *long-established* “ataxia genes” and “HSP genes,” causing a phenotypic spectrum ranging from ataxia to HSP as 2 extremes on a continuous spectrum;
2. shared pathways and mechanisms between ataxias and HSPs.

We thus argue to move on from the linkage-inspired divisive classifications system of largely distinct ataxias and HSP categories toward a more *modular understanding of phenotypes* that reflects the increasingly complex relationship between genotype, neuronal system damage, and phenotypic expression. The frequent co-occurrence of ataxia and spasticity might thereby be driven by shared vulnerability of corticospinal tract axons and cerebellar circuits toward disturbances of the same molecular pathways (for graphical overview of the main hypothesis and concept proposed here, see Fig. 1). A mechanistically inspired classification system will prioritize research on shared pathways and pave the way for mechanism-based strategies for drug development.

Discovering the Phenotypic and Genetic Spectrum From the Extremes

Discovery of an increasing number of genes causing both prominent cerebellar and predominantly pyramidal phenotypes over the past few years has raised awareness of the substantial overlap between these 2 disease classifications. Thereby, the “divide” was closed from both sides: classical “HSP genes” were discovered to cause ataxia as well as classical “ataxia genes” were recognized to result in HSP phenotypes.

For genes discovered in the pre-NGS era, it commonly took years (and, in some cases, the phenotypically unbiased screening approaches enabled by NGS application) to overcome the preconception of the predominant phenotype associated with a gene. *SPG7*, identified as a cause of HSP in 1998,⁷ was not systematically considered a cause of predominant (and even pure) cerebellar ataxia until 15 years later.⁸ Yet, within the past 2 years, it has been appreciated as one of the most common causes of autosomal-recessive cerebellar ataxia,^{9,10} and the cerebellar features may be even more pronounced than spasticity in some cohorts.¹⁰ Mutations in *PNPLA6* were identified as a cause of autosomal-recessive HSP complicated by motor axonal neuropathy in 2008, leading to the designation SPG39.¹¹ However, it was not before 2014 that mutations in *PNPLA6* were also appreciated as a cause of predominant cerebellar ataxia,^{12,13} and it has now been shown that *PNPLA6* mutations can even cause pure cerebellar ataxia.¹⁴ In light of these observations of patients with predominant or pure cerebellar disease, the terms “SPG7” and “SPG39” reflect the historical meaning at best — and appear to be misnomers for these patients and phenotypes. The fatty acid 2-hydroxylase gene (*FA2H*) is even part of multiple classification systems. After initially being discovered

as the causative gene for leukodystrophy associated with spastic paraparesis and dystonia,¹⁵ it was published 2 years later as a novel HSP gene (*SPG35*),¹⁶ only to be recognized to cause a novel form of neurodegeneration with brain iron accumulation (NBIA) termed FAHN — *FA2H*-associated neurodegeneration — a few months later.¹⁷ Not until recently, the substantial cerebellar ataxia present in many patients with *FA2H* mutations was systematically recognized.^{18,19}

Likewise, HSP phenotypes were often recognized belatedly for “traditional” ataxia genes. Recessive mutations in *SYNE1* were identified as a cause of cerebellar ataxia in 2007²⁰ and consequently designated ARCA1 and SCAR8. For almost a decade mutations in *SYNE1* were thought to cause a slowly progressive, largely pure cerebellar ataxia,^{21,22} before it was realized in 2016 that they are in fact causative for a broad pleiotropic phenotypic spectrum, with corticospinal tract damage and even predominant complicated HSP presentations among the most frequent features.^{23,24} Recessive mutations in *PLA2G6* were found in 2006 to cause, among others, a childhood-onset ataxia cluster (termed infantile neuroaxonal dystrophy).²⁵ Although concomitant corticospinal tract features have already been described in several reports in recent years, it was not until recently that complicated HSP has been acknowledged as one of the main phenotypic presentations of *PLA2G6* (Ozes et al, submitted). Biallelic *STUB1* mutations were first published as a cause of recessive ataxia (eg, as part of a Gordon Holmes syndrome).^{26,27} Later studies then revealed that corticospinal tract damage is a frequent concomitant feature²⁸ and sometimes even is predominate in the clinical presentation.²⁹ Examples can also be found for recently identified *autosomal-dominant* disease genes. Dominant mutations in *KCNA2* were first reported as a cause of (early-onset) cerebellar ataxia in 2015,^{30,31} before it was shown in 2016 that dominant *KCNA2* mutations can also cause HSP phenotypes,³² with both phenotypes occurring on a phenotypic continuum.

NGS has sped up not only disease gene discovery but also the time span from disease gene discovery until a broadened phenotypic spectrum can be appreciated. In some cases, this has led to the almost simultaneous “discovery” of one and the same gene as a novel ataxia gene and an HSP gene. Autosomal-recessive mutations in the nonlysosomal glucosylceramidase gene *GBA2*, for example, were designated SPG46 because of the predominant lower-limb spasticity noted by the European team of researchers.³³ In the same journal issue, however, *GBA2* was published as a novel gene for “cerebellar ataxia with spasticity” because of the initial disease manifestation as cerebellar ataxia in this independent patient cohort.³⁴ Similarly, *KIF1C* mutations were discovered to cause autosomal-recessive HSP complicated by ataxia features, termed SPG58.³⁵ At the same time, however, it was discovered that *KIF1C* mutations can also cause predominant cerebellar ataxia (with variable spasticity of the lower limbs).³⁶

These recent examples underscore the value of unbiased screening approaches enabled by NGS technology that — when combined with a modular phenotyping approach — enable rapid and comprehensive delineation of phenotypic spectra associated with Mendelian disease genes. Moreover, they illustrate that cerebellar and pyramidal disease manifestations commonly cooccur and can vary considerably in predominance and phenotypic expression along a continuous spectrum. This variable phenotypic presentation therefore does not

justify the classification of these ataxia-spasticity spectrum genes as SPG versus SCA/SCAR/ARCA genes. The distinct SPG-versus-SCA/SCAR/ARCA classification system fails to capture this inherent phenotypic fluidity, rendering it in part arbitrary, and is therefore of limited systematic value for clinic and research.

Large Common Genetic Basis of Ataxias and HSPs

The aforementioned examples of ataxia-spasticity spectrum (ASS) genes are part of a larger, rapidly growing list of genes causing ataxia and HSP on a phenotypic continuum. Based on review of the literature and our own experience with whole-exome sequencing (WES) and whole-genome sequencing of large cohorts of cases with ataxia and/or HSP, we have compiled an extendable list of 69 genes that we consider relevant in the differential diagnosis of ASS disease (Table 1). We included only genes the phenotypic descriptions of which included both ataxia and spasticity (rather than merely pyramidal signs) in subjects from at least 2 different families (rather than merely single cases). The majority of these ASS genes cause autosomal-recessive disease (n = 49), but autosomal-dominant (n = 16) and X-linked recessive (n = 3) modes of inheritance also occur. For mutations in *AFG3L2*, autosomal-recessive (SCAR5) and autosomal-dominant (SCA28) modes of inheritance have been established. Notably, only 29 genes (42%) are part of either of the HSP or ataxia classification systems mentioned above (SCA/SCAR/ARCA/SPG). Consequently, even combining disease genes contained in either of the HSP or ataxia classifications is insufficient to capture the relevant disease genes for the ASS. The implications for clinical and genetic diagnostic practice are apparent: NGS-based approaches to test for mutations in ataxia genes (“ataxia panels”) need to also comprise HSP genes and vice versa to do the overlapping disease spectra justice; in addition, both ataxia and HSP gene panels should be expanded to cover not only the relevant genes “by classification,” but need to go beyond classification systems to cover also genes not included in any of the classification systems.

Common Pathophysiological Pathways and Mechanisms in Ataxias and HSPs

Under the surface of the seemingly disparate clinical syndromic and diagnostic classifications between ataxias and HSPs lurk not only shared allelic genes, but also common mechanisms and pathways. In this respect, the overlap between ataxias and HSPs resembles the well-established gene and pathway overlap between amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD). Like HSP and ataxias, these 2 conditions have long been considered clinically disparate syndromes. Yet, over the past decade, we have increasingly recognized that they co-occur within families and even within individuals and largely share the same genes. Consequently, ALS and FTD are now usually studied jointly as a disease spectrum. Overcoming the diagnostic divide between ALS and FTD and focusing on shared pathways instead have led to identification of major shared mechanism hubs. For example, dysfunctional nuclear-cytoplasmic transport has emerged as a common mechanistic denominator uniting not only the different clinical conditions, but also various ALS/FTD genes like *C9orf72*, *FUS*, and *TARDBP*.³⁷⁻³⁹

Similarly, HSP and ataxias, which share a substantial number of genes, might also be connected on a functional level via shared cellular pathways and pathomechanisms. A protein-protein interaction network using known ASS genes as seeds (Table 1, n = 63, here excluding the dominant repeat ataxias) reveals that the proteins encoded by these genes share a multitude of physical interactions and form several highly connected “protein communities” that are visualized by different colors shown in Figure 2. Functional annotation of these genes using GO terms and subsequent gene set enrichment analysis highlight functional clusters that are enriched in these proteins (Fig. 3, Supplementary Table). The 3 major functional clusters are: (1) lipid metabolic processes, (2) acid metabolic processes, and (3) cytoskeleton or dendritic intracellular transport processes. These 3 clusters represent only a small subset of molecular pathways known to be involved in HSPs or cerebellar ataxias individually. This supports the hypothesis that pathways affected in ASS reflect shared selective vulnerabilities of corticospinal and cerebellar neurons. The *clinical* overlap of ASS spectrum diseases might thus be driven by underlying *mechanistic* overlaps (for an illustration of the relation between genetic, pathway, and clinical overlaps, see Fig. 1).

Some exemplary clusters of shared or interacting pathways underlying ASS diseases are:

- *Phospholipid metabolism*, including the genes *PNPLA6*,^{12,40,41} *PLA2G6*, *DDHD1* (SPG 28), *DDHD2* (SPG54⁴²), *CYP2U1* (SPG49), and *ABHD12*⁴³ (for further overview, see references ⁴⁰ and ⁴⁴).
- *Sphingolipid metabolism*, including the genes *FA2H*,¹⁵ *GBA2*,^{33,45} *GALC*, *HEXA*, *ASA*, *PSAP*, and *GLB1*.
- *Autophagy-lysosomal activity*, including the genes *SPG15*, *SPG11*,^{46,47} *ATP13A2* (SPG78),^{48,49} *NPC1*, and *NPC2* disease.⁵⁰⁻⁵⁵

Toward a Mechanism-Based Classification of Ataxia-Spasticity Spectrum Diseases

As our concepts of cellular pathways involved in ASS diseases grow, a mechanism-based classification system of the ASS comes into reach. Classification of genetically defined disorders by shared affected pathways rather than the perceived predominant phenotype will allow overcoming the classic SCA/SCAR/ARCA and HSP/SPG divide and appreciation of a more systematic, pathophysiological perspective. Other than the resolution of multiple inconsistencies of the traditional classification system which we have detailed above, a mechanistically inspired classification system of ASS diseases offers key advantages in therapeutic respects. Such a classification system will prioritize research on shared pathways and might pave the way for mechanism-based strategies for drug development.

Hypothetically, compounds targeting dysfunctional pathways rather than single genes have the potential to address groups of genetically defined diseases rather than single ataxia or HSP subtypes (for graphical illustration of this idea, see label “causal treatment strategies targeting pathways” in Fig. 1). For example, one class of drugs might target ASS diseases with abnormal cholesterol processing and cholesterol sequestration such as *CYP7B1* (SPG5), *NPC1*, *NPC2*, or *SERAC1* by exploiting cholesterol-depleting agents.⁵⁶ Another

class of drugs might aim at ASS diseases with defective autophagy-lysosomal activity (eg, *SPG11*, *ZFYVE26*, *ATP13A2*), using an autophagy inducer.⁵⁶ A mechanism-based disease classification might thus facilitate the translation of the giant genetic progress rendered possible by NGS over the past 5 years into first targeted molecular therapies.

Conclusions for Clinical Practice

In conclusion, we suggest to give up the classificatory divide between ataxias and HSPs in favor of a concept of a clinical, genetic, and pathophysiological ASS. From this inclusive rather than discriminatory approach, a number of advantages can be inferred for current clinical practice:

1. *Increased precision of phenotypic description and improved efficiency of diagnostic workup.* Early discriminatory classification of patients into fixed diagnostic categories potentially introduces bias into the clinical and diagnostic workup. We suggest taking a modular approach to phenotyping that allows the appreciation of nuanced individual phenotypic expression along the spectrum of ataxia and spasticity. This descriptive, unbiased approach of *modular phenotyping* would also be open to expansion of the phenotype beyond ataxia and HSP, as ataxia and spasticity often occur not in isolation, but as part of multisystem neuronal dysfunction. It thus allows for a more comprehensive, dynamic and systematic perspective than the traditional SCA/SCAR/ARCA and HSP/SPG classifications. Avoidance of narrow-minded ataxia and HSP clinical engrams will ultimately facilitate diagnosis in so-far unexplained complex neurodegenerative disease.
2. *Individualized treatment.* Following the idea of individualized medicine, modular phenotyping allows for individualized clinical treatment and management according to each individual's particular phenotypic spectrum (rather than by the overall clinical diagnosis or SPG/SCA/ARCA classification) (for a graphic illustration of the role of symptomatic treatment according to individual phenotype, see Figure 1). For example, patients with a major ataxia component due to *PNPLA6* or *SPG7* mutations will be clinically managed according to their individual ataxia, receiving, for example, physiotherapy exercises specifically targeting ataxia dysfunctions,^{57,58} even if these genes are traditionally grouped in the HSP/SPG classification (*SPG39* and *SPG7*, respectively). Vice versa, patients with pronounced spasticity because of *SYNE1* or *STUB1* mutations will be clinically managed according to their spasticity, receiving, for example, antispastic drugs, even if these genes are traditionally grouped in ARCA classifications.
3. *Efficient diagnostic testing.* Given the variability of phenotypes across the ASS and the sheer number of ASS genes, genetic testing on a gene-by-gene basis or relying on small gene panels is inefficient and mostly obsolete. Instead, genetic testing needs to resort to large gene panels or WES covering all ASS genes. Single-gene testing in ataxia spasticity spectrum diseases should be largely reserved for a few exceptions, for example, genotyping the *FRDA* repeat in

patients with afferent ataxia and pyramidal tract damage without major cerebellar atrophy, or the *SACS* gene in patients with the characteristic hypointense pontine stripes on T2-MRI imaging.⁵⁹

4. *Aggregated ASS gene panels and gene lists.* In NGS diagnostics, the design of separate ataxia and HSP NGS gene panels and of separate ataxia and HSP gene lists, respectively, for WES analyses is not productive. NGS gene panels and lists need to aggregate *all* ASS genes.

Limitations and Future Challenges

The proposed approach of modular phenotyping bears several limitations. Patients might prefer to have a clear-cut clinical label for their disease (eg, HSP or spinocerebellar ataxia) rather than an open and dynamic broad descriptive phenotypic description of the individually affected neurological systems. A clear label might yet be given the name of the underlying gene and/or the pathway cluster. However, sporadic ASS patients without monogenic disease causation or obvious hit in one of the pathway clusters will escape classification by the proposed pathway-driven classification system.

The suggested pathway-driven classification is also limited by it requiring the affected cellular pathways to be known. For the large majority of ASS diseases, however, the pathway implications of the respective disease genes have yet to be identified. Future basic research now has to move on from NGS genetics to functional pathway explorations, both for each specific ASS gene and for possible shared pathway hubs, identifying in particular those pathway hubs that might be druggable.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding agencies: This research was supported by the E-RARE-3 Joint Transnational Call grant “Preparing therapies for autosomal recessive ataxias” (PREPARE; BMBF, 01GM1607 to M.S.), the E-Rare network grant NEUROLIPID (BMBF, 01GM1408B to R.S.), the Marie Curie International Outgoing Fellowship (grant PIOF-GA-2012-326681 to R.S.), and by the National Institute of Neurological Disorders and Stroke of the National Institutes of Health under award number R01NS072248 (to R.S.).

References

1. Man OMI. Online Mendelian Inheritance in Man, OMIM. Baltimore, MD: McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University Available at: <http://omim.org/> Accessed October 30, 2016.
2. Marras C, Lang A, van de Warrenburg BP, et al. Nomenclature of genetic movement disorders: Recommendations of the international Parkinson and movement disorder society task force. *Mov Disord* 2016;31(4):436–457. [PubMed: 27079681]
3. Coutelier M, Burglen L, Mundwiller E, et al. GRID2 mutations span from congenital to mild adult-onset cerebellar ataxia. *Neurology* 2015;84(17):1751–1759. [PubMed: 25841024]
4. Pierson TM, Adams D, Bonn F, et al. Whole-exome sequencing identifies homozygous AFG3L2 mutations in a spastic ataxianeuropathy syndrome linked to mitochondrial m-AAA proteases. *PLoS Genet* 2011;7(10):e1002325. [PubMed: 22022284]

5. Lise S, Clarkson Y, Perkins E, et al. Recessive mutations in SPTBN2 implicate beta-III spectrin in both cognitive and motor development. *PLoS Genet* 2012;8(12):e1003074. [PubMed: 23236289]
6. de Bot ST, Willemsen MA, Vermeer S, Kremer HP, van de Warrenburg BP. Reviewing the genetic causes of spastic-ataxias. *Neurology* 2012;79(14):1507–1514. [PubMed: 23033504]
7. Casari G, De Fusco M, Ciarmatori S, et al. Spastic paraplegia and OXPHOS impairment caused by mutations in paraplegin, a nuclear-encoded mitochondrial metalloprotease. *Cell* 1998;93(6):973–983. [PubMed: 9635427]
8. van Gassen KL, van der Heijden CD, de Bot ST, et al. Genotype-phenotype correlations in spastic paraplegia type 7: a study in a large Dutch cohort. *Brain* 2012;135(Pt 10):2994–3004. [PubMed: 22964162]
9. Pfeffer G, Pyle A, Griffin H, et al. SPG7 mutations are a common cause of undiagnosed ataxia. *Neurology* 2015;84(11):1174–1176. [PubMed: 25681447]
10. Choquet K, Tetreault M, Yang S, et al. SPG7 mutations explain a significant proportion of French Canadian spastic ataxia cases. *Eur J Hum Genet* 2016;24(7):1016–1021. [PubMed: 26626314]
11. Rainier S, Bui M, Mark E, et al. Neuropathy target esterase gene mutations cause motor neuron disease. *Am J Hum Genet* 2008; 82(3):780–785. [PubMed: 18313024]
12. Synofzik M, Gonzalez MA, Lourenco CM, et al. PNPLA6 mutations cause Boucher-Neuhauser and Gordon Holmes syndromes as part of a broad neurodegenerative spectrum. *Brain* 2014; 137(Pt 1):69–77. [PubMed: 24355708]
13. Synofzik M, Hufnagel R, Zuchner S. PNPLA6-Related Disorders In: Pagon RA, Adam MP, Ardinger HH, et al., eds. *GeneReviews*. Seattle, WA: University of Washington; 1993–2014 Available at: <http://www.ncbi.nlm.nih.gov/books/NBK247161/2014>. Accessed August 18, 2016.
14. Wiethoff S, Bettencourt C, Paudel R, et al. Pure cerebellar ataxia with homozygous mutations in the PNPLA6 Gene. *Cerebellum* 2016;16(1): 262–267.
15. Edvardson S, Hama H, Shaag A, et al. Mutations in the fatty acid 2-hydroxylase gene are associated with leukodystrophy with spastic paraparesis and dystonia. *Am J Hum Genet* 2008;83(5): 643–648. [PubMed: 19068277]
16. Dick KJ, Eckhardt M, Paisan-Ruiz C, et al. Mutation of FA2H underlies a complicated form of hereditary spastic paraplegia (SPG35). *Hum Mutat* 2010;31(4):E1251–E1260. [PubMed: 20104589]
17. Kruer M, Gregory A, Hayflick SJ. Fatty Acid Hydroxylase-Associated Neurodegeneration In: Pagon RA, Bird TD, Dolan CR, Stephens K, eds. *GeneReviews*. Seattle, WA; 1993.
18. Pierson TM, Simeonov DR, Sincan M, et al. Exome sequencing and SNP analysis detect novel compound heterozygosity in fatty acid hydroxylase-associated neurodegeneration. *Eur J Hum Genet* 2012;20(4):476–479. [PubMed: 22146942]
19. Soehn AS, Rattay TW, Beck-Wodl S, et al. Uniparental disomy of chromosome 16 unmasks recessive mutations of FA2H/SPG35 in 4 families. *Neurology* 2016;87(2):186–191. [PubMed: 27316240]
20. Gros-Louis F, Dupre N, Dion P, et al. Mutations in SYNE1 lead to a newly discovered form of autosomal recessive cerebellar ataxia. *Nat Genet* 2007;39(1):80–85. [PubMed: 17159980]
21. Noreau A, Bourassa CV, Szuto A, et al. SYNE1 mutations in autosomal recessive cerebellar ataxia. *JAMA Neurol* 2013;70(10): 1296–1231. [PubMed: 23959263]
22. Dupre N, Gros-Louis F, Chrestian N, et al. Clinical and genetic study of autosomal recessive cerebellar ataxia type 1. *Ann Neurol* 2007;62(1):93–98. [PubMed: 17503513]
23. Synofzik M, Smets K, Mallaret M, et al. SYNE1 ataxia is a common recessive ataxia with major non-cerebellar features: a large multi-centre study. *Brain* 2016;139(Pt 5):1378–1393. [PubMed: 27086870]
24. Mademan I, Harmuth F, Giordano I, et al. Multisystemic SYNE1 ataxia: confirming the high frequency and extending the mutational and phenotypic spectrum. *Brain* 2016;139(Pt 8):e46. [PubMed: 27197992]
25. Khateeb S, Flusser H, Ofir R, et al. PLA2G6 mutation underlies infantile neuroaxonal dystrophy. *Am J Hum Genet* 2006;79(5): 942–948. [PubMed: 17033970]
26. Shi Y, Wang J, Li JD, et al. Identification of CHIP as a novel causative gene for autosomal recessive cerebellar ataxia. *PLoS One* 2013;8(12):e81884. [PubMed: 24312598]

27. Shi CH, Schisler JC, Rubel CE, et al. Ataxia and hypogonadism caused by the loss of ubiquitin ligase activity of the U box protein CHIP. *Hum Mol Genet* 2014;23(4):1013–1024. [PubMed: 24113144]
28. Synofzik M, Schule R, Schulze M, et al. Phenotype and frequency of STUB1 mutations: next-generation screenings in Caucasian ataxia and spastic paraplegia cohorts. *Orphanet J Rare Dis* 2014; 9(1):57. [PubMed: 24742043]
29. Hayer SN, Deconinck T, Bender B, et al. STUB1/CHIP mutations cause Gordon Holmes syndrome as part of a widespread multisystemic neurodegeneration: evidence from four novel mutations. *Orphanet Journal of Rare Diseases*: Epub ahead of print.
30. Syrbe S, Hedrich UB, Riesch E, et al. De novo loss- or gain-of-function mutations in KCNA2 cause epileptic encephalopathy. *Nat Genet* 2015;47(4):393–399. [PubMed: 25751627]
31. Pena SD, Coimbra RL. Ataxia and myoclonic epilepsy due to a heterozygous new mutation in KCNA2: proposal for a new channelopathy. *Clin Genet* 2015;87(2):e1–e3. [PubMed: 25477152]
32. Helbig KL, Hedrich UB, Shinde DN, et al. A recurrent mutation in KCNA2 as a novel cause of hereditary spastic paraplegia and ataxia. *Ann Neurol* 2016;80(4).
33. Martin E, Schule R, Smets K, et al. Loss of Function of Glucocerebrosidase GBA2 Is Responsible for Motor Neuron Defects in Hereditary Spastic Paraplegia. *Am J Hum Genet* 2013;92(2):238–244. [PubMed: 23332916]
34. Hammer MB, Eleuch-Fayache G, Schottlaender LV, et al. Mutations in GBA2 cause autosomal-recessive cerebellar ataxia with spasticity. *Am J Hum Genet* 2013;92(2):245–251. [PubMed: 23332917]
35. Novarino G, Fenstermaker AG, Zaki MS, et al. Exome sequencing links corticospinal motor neuron disease to common neurodegenerative disorders. *Science* 2014;343(6170):506–511. [PubMed: 24482476]
36. Dor T, Cinnamon Y, Raymond L, et al. KIF1C mutations in two families with hereditary spastic paraparesis and cerebellar dysfunction. *J Med Genet* 2014;51(2):137–142. [PubMed: 24319291]
37. Jovicic A, Paul JW, 3rd, Gitler AD. Nuclear transport dysfunction: a common theme in amyotrophic lateral sclerosis and frontotemporal dementia. *J Neurochem* 2016;138(Suppl 1):134–144. [PubMed: 27087014]
38. Boeynaems S, Bogaert E, Van Damme P, Van Den Bosch L. Inside out: the role of nucleocytoplasmic transport in ALS and FTLD. *Acta Neuropathol* 2016;132(2):159–173. [PubMed: 27271576]
39. Prpar Mihevc S, Darovic S, Kovanda A, Bajc Cesnik A, Zupunski V, Rogelj B. Nuclear trafficking in amyotrophic lateral sclerosis and frontotemporal lobar degeneration. *Brain* 2017;140(Pt 1):13–26. [PubMed: 27497493]
40. Tesson C, Nawara M, Salih MA, et al. Alteration of fatty-acid-metabolizing enzymes affects mitochondrial form and function in hereditary spastic paraplegia. *Am J Hum Genet* 2012;91(6):1051–1064. [PubMed: 23176821]
41. Kmoch S, Majewski J, Ramamurthy V, et al. Mutations in PNPLA6 are linked to photoreceptor degeneration and various forms of childhood blindness. *Nat Commun* 2015;6:5614. [PubMed: 25574898]
42. Schuurs-Hoeijmakers JH, Geraghty MT, Kamsteeg EJ, et al. Mutations in DDHD2, encoding an intracellular phospholipase A(1), cause a recessive form of complex hereditary spastic paraplegia. *Am J Hum Genet* 2012;91(6):1073–1081. [PubMed: 23176823]
43. Fiskerstrand T, H'Mida-Ben Brahim D, Johansson S, et al. Mutations in ABHD12 cause the neurodegenerative disease PHARC: An inborn error of endocannabinoid metabolism. *Am J Hum Genet* 2010;87(3):410–417. [PubMed: 20797687]
44. Lamari F, Mochel F, Saudubray JM. An overview of inborn errors of complex lipid biosynthesis and remodelling. *J Inherit Metab Dis* 2015;38(1):3–18. [PubMed: 25238787]
45. Sultana S, Reichbauer J, Schule R, Mochel F, Synofzik M, van der Spoel AC. Lack of enzyme activity in GBA2 mutants associated with hereditary spastic paraplegia/cerebellar ataxia (SPG46). *Biochem Biophys Res Commun* 2015;465(1):35–40. [PubMed: 26220345]
46. Chang J, Lee S, Blackstone C. Spastic paraplegia proteins spastizin and spatacsin mediate autophagic lysosome reformation. *J Clin Invest* 2014;124(12):5249–5262. [PubMed: 25365221]


47. Varga RE, Khundadze M, Damme M, et al. In vivo evidence for lysosome depletion and impaired autophagic clearance in hereditary spastic paraplegia type SPG11. *PLoS Genet* 2015;11(8): e1005454. [PubMed: 26284655]
48. Estrada-Cuzcano A, Martin S, Chamova T, et al. Loss-of-function mutations in the ATP13A2/PARK9 gene cause complicated hereditary spastic paraplegia (SPG78). *Brain* [in press].
49. Bento CF, Ashkenazi A, Jimenez-Sanchez M, Rubinsztein DC. The Parkinson's disease-associated genes ATP13A2 and SYT11 regulate autophagy via a common pathway. *Nat Commun* 2016; 7:11803. [PubMed: 27278822]
50. Guo H, Zhao M, Qiu X, et al. Niemann-Pick type C2 deficiency impairs autophagy-lysosomal activity, mitochondrial function, and TLR signaling in adipocytes. *J Lipid Res* 2016;57(9):1644–1658. [PubMed: 27402802]
51. Tamura A, Yui N. Beta-cyclodextrin-threaded biocleavable polyrotaxanes ameliorate impaired autophagic flux in Niemann-Pick type C disease. *J Biol Chem* 2015;290(15):9442–9454. [PubMed: 25713067]
52. Sarkar S, Carroll B, Buganim Y, et al. Impaired autophagy in the lipid-storage disorder Niemann-Pick type C1 disease. *Cell Rep* 2013;5(5):1302–1315. [PubMed: 24290752]
53. Schicks J, Muller Vom Hagen J, Bauer P, et al. Niemann-Pick type C is frequent in adult ataxia with cognitive decline and vertical gaze palsy. *Neurology* 2013;80(12):1169–1170. [PubMed: 23427322]
54. Synofzik M, Harmuth F, Stampfer M, Muller Vom Hagen J, Schols L, Bauer P. NPC1 is enriched in unexplained early onset ataxia: a targeted high-throughput screening. *J Neurol* 2015; 262(11): 2557–2563. [PubMed: 26338816]
55. Synofzik M, Fleszar Z, Schols L, et al. Identifying Niemann-Pick type C in early-onset ataxia: two quick clinical screening tools. *J Neurol* 2016;263(10):1911–1918. [PubMed: 27314965]
56. Sarkar S, Maetzel D, Korolchuk VI, Jaenisch R. Restarting stalled autophagy a potential therapeutic approach for the lipid storage disorder, Niemann-Pick type C1 disease. *Autophagy* 2014;10(6): 1137–1140. [PubMed: 24879158]
57. Ilg W, Synofzik M, Brotz D, Burkard S, Giese MA, Schols L. Intensive coordinative training improves motor performance in degenerative cerebellar disease. *Neurology* 2009;73(22):1823–1830. [PubMed: 19864636]
58. Ilg W, Schatton C, Schicks J, Giese MA, Schols L, Synofzik M. Video game-based coordinative training improves ataxia in children with degenerative ataxia. *Neurology* 2012;79(20):2056–2060. [PubMed: 23115212]
59. Synofzik M, Soehn AS, Gburek-Augustat J, et al. Autosomal recessive spastic ataxia of Charlevoix Saguenay (ARSACS): expanding the genetic, clinical and imaging spectrum. *Orphanet J Rare Dis* 2013;8(1):41. [PubMed: 23497566]
60. Shannon P, Markiel A, Ozier O, et al. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* 2003;13(11):2498–2504. [PubMed: 14597658]
61. Razick S, Magklaras G, Donaldson IM. iRefIndex: a consolidated protein interaction database with provenance. *BMC Bioinformatics* 2008;9:405. [PubMed: 18823568]
62. Razick S, Mora A, Michalickova K, Boddie P, Donaldson IM. iRefScape. A Cytoscape plug-in for visualization and data mining of protein interaction data from iRefIndex. *BMC Bioinformatics* 2011;12:388. [PubMed: 21975162]
63. Huang da W, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 2009;4(1):44–57. [PubMed: 19131956]
64. Huang da W, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 2009;37(1):1–13. [PubMed: 19033363]
65. Gene Ontology Consortium. Gene Ontology Consortium: going forward. *Nucleic Acids Res* 2015;43(Database issue):D1049–D1056. [PubMed: 25428369]
66. Merico D, Isserlin R, Bader GD. Visualizing gene-set enrichment results using the Cytoscape plug-in enrichment map. *Methods Mol Biol* 2011;781:257–277. [PubMed: 21877285]

67. Mosser J, Douar AM, Sarde CO, et al. Putative X-linked adrenoleukodystrophy gene shares unexpected homology with ABC transporters. *Nature* 1993;361(6414):726–730. [PubMed: 8441467]
68. Rauschka H, Colsch B, Baumann N, et al. Late-onset metachromatic leukodystrophy: genotype strongly influences phenotype. *Neurology* 2006;67(5):859–863. [PubMed: 16966551]
69. Nagafuchi S, Yanagisawa H, Sato K, et al. Dentatorubral and pallidoluysian atrophy expansion of an unstable CAG trinucleotide on chromosome 12p. *Nat Genet* 1994;6(1):14–18. [PubMed: 8136826]
70. Estrada-Cuzcano A, Shaun M, Chamova T, et al. Loss-of-function mutations in the ATP13A2/PARK9 gene cause complicated hereditary spastic paraplegia (SPG78). *Brain* 140:287–305. [PubMed: 28137957]
71. Banfi S, Servadio A, Chung MY, et al. Identification and characterization of the gene causing type 1 spinocerebellar ataxia. *Nat Genet* 1994;7(4):513–520. [PubMed: 7951322]
72. Pulst SM, Nechiporuk A, Nechiporuk T, et al. Moderate expansion of a normally biallelic trinucleotide repeat in spinocerebellar ataxia type 2. *Nat Genet* 1996;14(3):269–276. [PubMed: 8896555]
73. Kawaguchi Y, Okamoto T, Taniwaki M, et al. CAG expansions in a novel gene for Machado-Joseph disease at chromosome 14q32.1. *Nat Genet* 1994;8(3):221–228. [PubMed: 7874163]
74. Koob MD, Moseley ML, Schut LJ, et al. An untranslated CTG expansion causes a novel form of spinocerebellar ataxia (SCA8). *Nat Genet* 1999;21(4):379–384. [PubMed: 10192387]
75. L II, Loupatty FJ, Ruitter JP, Duran M, Lehnert W, Wanders RJ. 3-Methylglutaconic aciduria type I is caused by mutations in AUH. *Am J Hum Genet* 2002;71(6):1463–1466. [PubMed: 12434311]
76. Gan-Or Z, Bouslam N, Birouk N, et al. Mutations in CAPN1 cause autosomal-recessive hereditary spastic paraplegia. *Am J Hum Genet* 2016;98(5):1038–1046. [PubMed: 27153400]
77. Cali JJ, Hsieh CL, Francke U, Russell DW. Mutations in the bile acid biosynthetic enzyme sterol 27-hydroxylase underlie cerebrotendinous xanthomatosis. *J Biol Chem* 1991;266(12):7779–7783. [PubMed: 2019602]
78. Salen G, Berginer V, Shore V, et al. Increased concentrations of cholestanol and apolipoprotein B in the cerebrospinal fluid of patients with cerebrotendinous xanthomatosis. Effect of chenodeoxycholic acid. *N Engl J Med* 1987;316(20):1233–1238. [PubMed: 3106810]
79. Tsaousidou MK, Ouahchi K, Warner TT, et al. Sequence alterations within CYP7B1 implicate defective cholesterol homeostasis in motor-neuron degeneration. *Am J Hum Genet* 2008;82(2):510–515. [PubMed: 18252231]
80. Schule R, Siddique T, Deng HX, et al. Marked accumulation of 27-hydroxycholesterol in SPG5 patients with hereditary spastic paresis. *J Lipid Res* 2010;51(4):819–823. [PubMed: 19812052]
81. Scheper GC, van der Kloek T, van Andel RJ, et al. Mitochondrial aspartyl-tRNA synthetase deficiency causes leukoencephalopathy with brain stem and spinal cord involvement and lactate elevation. *Nat Genet* 2007;39(4):534–539. [PubMed: 17384640]
82. Wan J, Yourshaw M, Mamsa H, et al. Mutations in the RNA exosome component gene EXOSC3 cause pontocerebellar hypoplasia and spinal motor neuron degeneration. *Nat Genet* 2012;44(6):704–708. [PubMed: 22544365]
83. Campuzano V, Montermini L, Molto MD, et al. Friedreich's ataxia: autosomal recessive disease caused by an intronic GAA triplet repeat expansion. *Science* 1996;271(5254):1423–1427. [PubMed: 8596916]
84. Tappino B, Biancheri R, Mort M, et al. Identification and characterization of 15 novel GALC gene mutations causing Krabbe disease. *Hum Mutat* 2010;31(12):E1894–1914. [PubMed: 20886637]
85. Zemmouri R, Azzedine H, Assami S, et al. Charcot-Marie-Tooth 2-like presentation of an Algerian family with giant axonal neuropathy. *Neuromuscul Disord* 2000;10(8):592–598. [PubMed: 11053687]
86. Li R, Johnson AB, Salomons G, et al. Glial fibrillary acidic protein mutations in infantile, juvenile, and adult forms of Alexander disease. *Ann Neurol* 2005;57(3):310–326. [PubMed: 15732097]
87. Orthmann-Murphy JL, Salsano E, Abrams CK, et al. Hereditary spastic paraplegia is a novel phenotype for GJA12/GJC2 mutations. *Brain* 2009;132(Pt 2):426–438. [PubMed: 19056803]

88. Chakraborty S, Rafi MA, Wenger DA. Mutations in the lysosomal beta-galactosidase gene that cause the adult form of GM1 gangliosidosis. *Am J Hum Genet* 1994;54(6):1004–1013. [PubMed: 8198123]
89. Baker PR, 2nd, Friederich MW, Swanson MA, et al. Variant non ketotic hyperglycinemia is caused by mutations in LIAS, BOLA3 and the novel gene GLRX5. *Brain* 2014;137(Pt 2):366–379. [PubMed: 24334290]
90. Hills LB, Masri A, Konno K, et al. Deletions in GRID2 lead to a recessive syndrome of cerebellar ataxia and tonic upgaze in humans. *Neurology* 2013;81(16):1378–1386. [PubMed: 24078737]
91. Neudorfer O, Pastores GM, Zeng BJ, Gianutsos J, Zaroff CM, Kolodny EH. Late-onset Tay-Sachs disease: phenotypic characterization and genotypic correlations in 21 affected patients. *Genet Med* 2005;7(2):119–123. [PubMed: 15714079]
92. Lee YC, Durr A, Majczenko K, et al. Mutations in KCND3 cause spinocerebellar ataxia type 22. *Ann Neurol* 2012;72(6):859–869. [PubMed: 23280837]
93. Klebe S, Lossos A, Azzedine H, et al. KIF1A missense mutations in SPG30, an autosomal recessive spastic paraplegia: distinct phenotypes according to the nature of the mutations. *Eur J Hum Genet* 2012;20(6):645–649. [PubMed: 22258533]
94. Caballero Oteyza A, Battaloglu E, Ocek L, et al. Motor protein mutations cause a new form of hereditary spastic paraplegia. *Neurology* 2014;82(22):2007–2016. [PubMed: 24808017]
95. Bayat V, Thiffault I, Jaiswal M, et al. Mutations in the mitochondrial methionyl-tRNA synthetase cause a neurodegenerative phenotype in flies and a recessive ataxia (ARSAL) in humans. *PLoS Biol* 2012;10(3):e1001288. [PubMed: 22448145]
96. Christodoulou J, Ho G. MECP2-Related Disorders In: Pagon RA, Adam MP, Ardinger HH, et al., eds. *GeneReviews*. Seattle, WA; 1993.
97. Coelho D, Suormala T, Stucki M, et al. Gene identification for the cblD defect of vitamin B12 metabolism. *N Engl J Med* 2008; 358(14):1454–1464. [PubMed: 18385497]
98. Crosby AH, Patel H, Chioza BA, et al. Defective mitochondrial mRNA maturation is associated with spastic ataxia. *Am J Hum Genet* 2010;87(5):655–660. [PubMed: 20970105]
99. Patterson M Niemann-Pick Disease Type C In: Pagon RA, Adam MP, Ardinger HH, et al., eds. *GeneReviews*. Seattle, WA; 1993.
100. Yu-Wai-Man P, Griffiths PG, Gorman GS, et al. Multi-system neurological disease is common in patients with OPA1 mutations. *Brain* 2010;133(Pt 3):771–786. [PubMed: 20157015]
101. Bonifert T, Karle KN, Tonagel F, et al. Pure and syndromic optic atrophy explained by deep intronic OPA1 mutations and an intralocus modifier. *Brain* 2014;137(Pt 8):2164–2177. [PubMed: 24970096]
102. Gunay-Aygun M, Huizing M, Anikster Y. OPA3-Related 3-Methylglutaconic Aciduria In: Pagon RA, Adam MP, Ardinger HH, et al., eds. *GeneReviews*. Seattle, WA; 1993.
103. Schiff M, Mine M, Brivet M, et al. Leigh's disease due to a new mutation in the PDHX gene. *Ann Neurol* 2006;59(4):709–714. [PubMed: 16566017]
104. Ebberink MS, Csanyi B, Chong WK, et al. Identification of an unusual variant peroxisome biogenesis disorder caused by mutations in the PEX16 gene. *J Med Genet* 2010;47(9):608–615. [PubMed: 20647552]
105. Ozes B, Karagoz N, Schule R, et al. New families with PLA2G6 gene mutations suggest a continuous clinical spectrum from neuroaxonal dystrophy to hereditary spastic paraplegia. [under revision].
106. Synofzik M, Gasser T. Moving beyond syndromic classifications in neurodegenerative disease: the example of PLA2G6. *Movement Disorders Clinical Practice* 2017;4:8–11.
107. Inoue K PLP1-related inherited dysmyelinating disorders: Pelizaeus-Merzbacher disease and spastic paraplegia type 2. *Neurogenetics* 2005;6(1):1–16. [PubMed: 15627202]
108. Bernard G, Chouery E, Putorti ML, et al. Mutations of POLR3A encoding a catalytic subunit of RNA polymerase Pol III cause a recessive hypomyelinating leukodystrophy. *Am J Hum Genet* 2011;89(3):415–423. [PubMed: 21855841]
109. Minnerop M, Kurzwelly D, Wagner H, et al. Intronic mutations in POLR3A are a major cause of sporadic and recessive spastic ataxia [under review].

110. Saitsu H, Osaka H, Sasaki M, et al. Mutations in POLR3A and POLR3B encoding RNA Polymerase III subunits cause an autosomal-recessive hypomyelinating leukoencephalopathy. *Am J Hum Genet* 2011;89(5):644–651.
111. Zarranz JJ, Digon A, Atares B, et al. Phenotypic variability in familial prion diseases due to the D178N mutation. *J Neurol Neurosurg Psychiatry* 2005;76(11):1491–1496. [PubMed: 16227536]
112. Synofzik M, Bauer P, Schols L. Prion mutation D178N with highly variable disease onset and phenotype. *J Neurol Neurosurg Psychiatry* 2009;80(3):345–346. [PubMed: 19228673]
113. Kretz KA, Carson GS, Morimoto S, Kishimoto Y, Fluharty AL, O'Brien JS. Characterization of a mutation in a family with saposin B deficiency: a glycosylation site defect. *Proc Natl Acad Sci U S A* 1990;87(7):2541–2544. [PubMed: 2320574]
114. Dolzhanskaya N, Gonzalez MA, Sperziani F, et al. A novel p.Leu(381)Phe mutation in presenilin 1 is associated with very early onset and unusually fast progressing dementia as well as lysosomal inclusions typically seen in Kufs disease. *J Alzheimers Dis* 2014;39(1):23–27. [PubMed: 24121961]
115. Engert JC, Berube P, Mercier J, et al. ARSACS, a spastic ataxia common in northeastern Quebec, is caused by mutations in a new gene encoding an 11.5-kb ORF. *Nat Genet* 2000;24(2):120–125. [PubMed: 10655055]
116. Hammer MF, Wagnon JL, Mefford HC, Meisler MH. SCN8A-Related Epilepsy With Encephalopathy In: Pagon RA, Adam MP, Ardinger HH, et al., eds. *GeneReviews*. Seattle, WA; 1993.
117. Bourgeron T, Rustin P, Chretien D, et al. Mutation of a nuclear succinate dehydrogenase gene results in mitochondrial respiratory chain deficiency. *Nat Genet* 1995;11(2):144–149. [PubMed: 7550341]
118. Anheim M, Monga B, Fleury M, et al. Ataxia with oculomotor apraxia type 2: clinical, biological and genotype/phenotype correlation study of a cohort of 90 patients. *Brain* 2009;132(Pt 10):2688–2698. [PubMed: 19696032]
119. Verheijen FW, Verbeek E, Aula N, et al. A new gene, encoding an anion transporter, is mutated in sialic acid storage diseases. *Nat Genet* 1999;23(4):462–465. [PubMed: 10581036]
120. Camacho JA, Obie C, Biery B, et al. Hyperornithinaemia-hyper-ammonaemia-homocitrullinuria syndrome is caused by mutations in a gene encoding a mitochondrial ornithine transporter. *Nat Genet* 1999;22(2):151–158. [PubMed: 10369256]
121. Weber YG, Kamm C, Suls A, et al. Paroxysmal choreoathetosis/ spasticity (DYT9) is caused by a GLUT1 defect. *Neurology* 2011; 77(10):959–964. [PubMed: 21832227]
122. Stevanin G, Santorelli FM, Azzedine H, et al. Mutations in SPG11, encoding spatacsin, are a major cause of spastic paraplegia with thin corpus callosum. *Nat Genet* 2007;39(3):366–372. [PubMed: 17322883]
123. Hanein S, Martin E, Boukhris A, et al. Identification of the SPG15 gene, encoding spastizin, as a frequent cause of complicated autosomal-recessive spastic paraplegia, including Kjellin syndrome. *Am J Hum Genet* 2008;82(4):992–1002. [PubMed: 18394578]
124. Bonafe L, Thony B, Penzien JM, Czarniecki B, Blau N. Mutations in the sepiapterin reductase gene cause a novel tetrahydrobiopterin-dependent monoamine-neurotransmitter deficiency without hyperphenylalaninemia. *Am J Hum Genet* 2001; 69(2):269–277. [PubMed: 11443547]
125. Synofzik M, Schule R, Schulze M, et al. Phenotype and frequency of STUB1 mutations: next-generation screenings in Caucasian ataxia and spastic paraplegia cohorts. *Orphanet J Rare Dis* 2014; 9:57. [PubMed: 24742043]
126. Koide R, Kobayashi S, Shimohata T, et al. A neurological disease caused by an expanded CAG trinucleotide repeat in the TATA-binding protein gene: a new polyglutamine disease? *Hum Mol Genet* 1999;8(11):2047–2053. [PubMed: 10484774]
127. Ghezzi D, Arzuffi P, Zordan M, et al. Mutations in TTC19 cause mitochondrial complex III deficiency and neurological impairment in humans and flies. *Nat Genet* 2011;43(3):259–263. [PubMed: 21278747]
128. Ouahchi K, Arita M, Kayden H, et al. Ataxia with isolated vitamin E deficiency is caused by mutations in the alpha-tocopherol transfer protein. *Nat Genet* 1995;9(2):141–145. [PubMed: 7719340]

129. Miyatake S, Osaka H, Shiina M, et al. Expanding the phenotypic spectrum of TUBB4A-associated hypomyelinating leukoencephalopathies. *Neurology* 2014;82(24):2230–2237. [PubMed: 24850488]
130. Bilguvar K, Tyagi NK, Ozkara C, et al. Recessive loss of function of the neuronal ubiquitin hydrolase UCHL1 leads to early-onset progressive neurodegeneration. *Proc Natl Acad Sci U S A* 2013; 110(9):3489–3494. [PubMed: 23359680]
131. Bourassa CV, Meijer IA, Merner ND, et al. VAMP1 mutation causes dominant hereditary spastic ataxia in Newfoundland families. *Am J Hum Genet* 2012;91(3):548–552. [PubMed: 22958904]
132. Kawarai T, Tajima A, Kuroda Y, et al. A homozygous mutation of VWA3B causes cerebellar ataxia with intellectual disability. *J Neurol Neurosurg Psychiatry* 2016;87(6):656–662. [PubMed: 26157035]

**FIG. 1.**

From clinical diagnosis to modular phenotyping and underlying shared genes and pathways in ataxia-spasticity spectrum diseases. Ataxias and HSPs have traditionally been designated in separate clinicogenetic disease classifications, depending on the first phenotypic descriptions and pattern of inheritance, namely, either in autosomal-dominant spinocerebellar ataxias (SCAs) and autosomal-recessive spinocerebellar ataxias (SCARs) or hereditary spastic paraplegias (HSPs/SPGs). However, the molecular etiologies of these 2 disease groups overlap greatly, based on manifold shared disease genes (top). Moreover, proteins encoded by HSP and ataxia genes closely interact physically as well as functionally. The heterogeneous genetic etiology of HSPs and ataxias thus converges into a small number of cellular pathways that are dysregulated in both diseases (right). Selective vulnerability of specific neuronal cell types that can be modified by additional genetic, epigenetic and environmental factors ultimately determines which neuronal systems and circuits will be affected by the pathway dysfunction (bottom). In ataxia-spasticity spectrum diseases, cerebellar and corticospinal tract neurons share selective vulnerabilities. The individual phenotypic expression (left) is a result of the pattern of neuronal system affection. It is essential to appreciate these 4 aspects, not only to understand an individual's disease, but also to use all therapeutic routes, whether they be symptomatic or causal/disease modifying. Pathway-based treatment approaches are hereby particularly promising, as (1) they offer the potential to cure, not only to modify the disease condition, with the (2) pathway-based etiologies partly converging from the vastly heterogeneous genetic etiology. Targeting dysfunctional pathways rather than single genes or disease conditions thus has the potential to address whole groups of genetically defined ASS diseases. [Color figure can be viewed at wileyonlinelibrary.com]




FIG. 2.

Protein-protein-interaction network for ataxia-HSP spectrum disease genes. To generate a protein-protein-interaction (PPI) network, interactors (iteration 1) of 63 spastic ataxia genes listed in Table 1 were extracted using the iRefScape plugin in Cytoscape v2.8.3.⁶⁰⁻⁶² Autosomal-dominant repeat-expansion genes were removed from the seed list, as their binding properties appear to be largely shaped by their polyglutamine tracts rather than the properties of the wild-type protein. The PPI network was then imported into Gephi v0.9.1 and filtered, whereby nonhuman interactions, predicted interactions, interactions based solely on high-throughput experiments, self-loops, and nodes with a degree < 2 were removed, retaining only nodes that were maximum 1 degree removed from the input spastic ataxia seed genes. The resulting network contained 389 nodes and 2582 undirected edges. We then applied the Louvain method to detect communities, neighborhoods of highly connected nodes. A total of 8 communities were detected, represented by differently colored nodes and edges in the figure. Ataxia-spasticity spectrum seed genes are represented by larger dots and labeled with the respective gene name. [Color figure can be viewed at wileyonlinelibrary.com]




FIG. 3.

Pathway enrichment map of ataxia-HSP spectrum gene sets. Gene-set enrichment analysis reveals several functional clusters associated with ataxia-spasticity spectrum genes. To generate the pathway, enrichment map spastic ataxia genes (Table 1, $n = 63$ genes; dominant repeat genes excluded) were uploaded to DAVID Bioinformatics Resources 6.8^{63,64} and annotated with gene ontology terms (GOTERM_BP_FAT, GOTERM_MF_FAT).⁶⁵ The fully annotated gene list is provided in the Supplementary Table. A gene enrichment map was then generated using the Enrichment Map plugin⁶⁶ in Cytoscape v3.2 with the following parameters: P cutoff, 0.0001; FDR Q cutoff, 0.05; similarity cutoff overlap, 0.4. Three major enrichment clusters can be appreciated: (1) lipid metabolic processes (blue), (2) acid metabolic processes (orange), and (3) cytoskeleton or dendritic intracellular transport processes (green). Each major network contains several subnetworks highlighting a specific cellular process underlying ataxia-spasticity spectrum disease. The size of the nodes reflects the number of ataxia-spasticity spectrum genes represented in the respective functional cluster; the number of genes is also indicated by the number in each of the nodes. [Color figure can be viewed at wileyonlinelibrary.com]

TABLE 1.

Genes causing ataxia and HSP (ataxia-HSP spectrum disease genes)

Gene	Locus	Protein name (UniProt)	Inheritance	OMIM	Remarks	Key references
<i>ABCD1</i>		ATP-binding cassette subfamily D member 1	XR	#300100	Adrenoleukodystrophy (ALD), adrenomyeloneuropathy (AMN); increased very long-chain fatty acids in plasma	67
<i>ABHD12</i>		Monoacylglycerol lipase ABHD12	AR	#612674	Peripheral neuropathy, hearing loss, retinitis pigmentosa, cataract (PHARC)	43
<i>AFG3L2</i>	SPAX5 (AR), SCA28 (AD)	AFG3-like protein 2	AR/AD	#614487	Catalytic subunit of the m-AAA protease (like paraplegin/SPG7); ophthalmoparesis, slow saccades, ptosis	4
<i>ARSA</i>		Arylsulfatase A	AR	#250100	Metachromatic leukodystrophy (MLD); reduced arylsulfatase A activity in leukocytes	68
<i>ATN1</i>		Atrophin-1	AD	#125370	Dentatorubral-pallidolysian atrophy (DRPLA); CAG trinucleotide expansion; myoclonic epilepsy, dementia, ataxia, and choreoathetosis; rare outside Japan	69
<i>ATP13A2</i>	SPG78	Probable cation-transporting ATPase 13A2	AR	#606693	Juvenile parkinsonism, vertical gaze palsy, cognitive deficits	70
<i>ATXN1</i>	SCA1	Ataxin-1	AD	#164400	CAG trinucleotide expansion	71
<i>ATXN2</i>	SCA2	Ataxin-2	AD	#183090	CAG trinucleotide expansion; slow horizontal saccades	72
<i>ATXN3</i>	SCA3	Ataxin-3	AD	#109150	CAG trinucleotide expansion; Machado-Joseph disease; frequent SCA in central Europe	73
<i>ATXN8/ATXN8OS</i>	SCA8	Ataxin-8; putative protein ATXN8OS	AD	#608768	Expanded CTG trinucleotide repeat in ATXN8OS gene and complementary CAG repeat in ATXN8 gene	74
<i>AUH</i>		Methylglutaconyl-CoA hydratase, mitochondrial	AR	#250950	3-Methylglutaconic aciduria type 1 (MCGA1); elevated levels of 3-methylglutaconic acid (3-MGA), 3-methylglutaric acid (3-MG) and 3-hydroxyisovaleric acid (3-HIVA) in urine; cognitive deficits	75
<i>CAPN1</i>	SPG76	Calpain-1 catalytic subunit	AR	#616907	Upper limb involvement, foot deformities, dysarthria	76
<i>CYP27A1</i>		Sterol 26-hydroxylase, mitochondrial	AR	#213700	Cerebrotendinous xanthomatosis (CTX); juvenile cataract, lipid deposits i. a. in brain, lungs, and Achilles tendons, chronic diarrhea, early atherosclerosis, elevated levels of cholesterol in plasma	77,78
<i>CYP7B1</i>	SPG5	25-Hydroxycholesterol 7-alpha-hydroxylase	AR	#270800	Afferent ataxia due to dorsal column dysfunction, elevated levels of 27-hydroxycholesterol, 25-hydroxycholesterol, and cholestanic acid in plasma and CSF	79,80
<i>DARS2</i>		Aspartate-tRNA ligase, mitochondrial	AR	#611105	Leukoencephalopathy with brain stem and spinal cord involvement and lactate elevation (LBSL)	81
<i>EXOSC3</i>	PCH1B	Exosome complex component RRP40	AR	#614678	Pontocerebellar hypoplasia, type IB	82
<i>FA2H</i>	SPG35	Fatty acid 2-hydroxylase	AR	#612319	Spastic paraplegia, leukodystrophy, and/or brain iron deposition	18,19
<i>FXN</i>		Frxatin	AR	#229300	Friedreich ataxia (FRDA); predominant afferent ataxia with pyramidal tract signs	83
<i>GALC</i>		Galactocerebrosidase	AR	#245200	Krabbe disease; infantile forms with extreme irritability, spasticity, and developmental delay; late-adult forms: spasticity, ataxia; reduced GALC enzyme activity	84
<i>GAN</i>		Gigaxonin	AR	#256850	Giant axonal neuropathy (GAN1); infantile form: kinky hair and unique posture of legs	85
<i>GBA2</i>	SPG46	Nonlysosomal glucosylesteramidase	AR	#614409	Mental impairment, cataracts, cerebral, cerebellar, and corpus callosum atrophy	33
<i>GFAP</i>		Glial fibrillary acidic protein	AD	#203450	Alexander disease; infantile form: leukoencephalopathy with macrocephaly, seizures, and psychomotor retardation; adult form: bulbar signs and spasticity, more slowly progressive	86
<i>GJC2</i>	SPG44, HLD2	Gap junction gamma-2 protein	AR	#613206, #608804	Hypomyelinating leukodystrophy; Pelizaeus-Merzbacher-like disease (PMLD)	87

Gene	Locus	Protein name (UniProt)	Inheritance	OMIM	Remarks	Key references
<i>GLBI</i>		Beta-galactosidase	AR	#230500, #230600, #230650	GM1-gangliosidosis, type I-III (GLB1); mucopolysaccharidosis type IVB (Morquio syndrome B); reduced beta-galactosidase-1 enzyme activity	88
<i>GLRX5</i>		Glutaredoxin-related protein 5, mitochondrial	AR	#616859	Increased serum glycine; leukodystrophy and/or lesions in the upper spinal cord	89
<i>GRID2</i>	SCAR18	Glutamate receptor ionotropic, delta-2	AR	#616204	Early-onset cerebellar ataxia, intellectual disability; occasional or persistent tonic upgaze	90
<i>HEXA</i>		Beta-hexosaminidase subunit alpha	AR	#272800	Tay-Sachs disease/GM2-gangliosidosis; infantile: developmental retardation, followed by paresis, cognitive decline, and blindness; adult: lower motor neuron damage, psychosis, dementia; reduced hexosaminidase A enzyme activity	91
<i>KCNA2</i>		Potassium voltage-gated channel subfamily A member 2	AD	#616366	Early infantile epileptic encephalopathy; ataxia, spasticity; often de novo	30,32
<i>KCND3</i>	SCA19	Potassium voltage-gated channel subfamily D member 3	AD	#607346	Allelic with Brigada syndrome 9	92
<i>KIF1A</i>	SPG30, HSN2C	Kinesin-like protein KIF1A	AR	#610357, #614213	Allelic with autosomal-dominant mental retardation 9 (MRD9)	93
<i>KIF1C</i>	SPG58, SPAX2	Kinesin-like protein KIF1C	AR	#611302	Cerebellar ataxia and variable spasticity of the lower limbs in first 2 decades of life	36,94
<i>MARS2</i>	SPAX3	Methionine-tRNA ligase, mitochondrial	AR	#611390	Often deletions or duplications; decreased activity of mitochondrial complexes I and IV	95
<i>MECP2</i>		Methyl-CpG-binding protein 2	XR	#312750	Atypical Rett syndrome	96
<i>MMADHC</i>		Methylmalonic aciduria and homocystinuria type D protein, mitochondrial	AR	#277410	Homocystinuria and/or methylmalonic aciduria; usually additional complicating features like developmental delay	97
<i>MTPAP</i>	SPAX4	Poly(A) RNA polymerase, mitochondrial	AR	#613672	Childhood-onset cerebellar ataxia, spastic paraparesis, dysarthria, and optic atrophy	98
<i>NPCI</i>		Niemann-Pick C1 protein	AR	#257220	Niemann-Pick disease, type C1/D; infantile: often accompanying neurovisceral phenotype; juvenile and adult: often vertical supranuclear gaze palsy or cognitive decline	55,99
<i>NPC2</i>		Epididymal secretory protein E1	AR	#607625	Niemann-Pick disease, type C2; infantile: often accompanying neurovisceral phenotype; juvenile and adult: often vertical supranuclear gaze palsy or cognitive decline	55,99
<i>OPA1</i>		Dynammin-like 120-kDa protein, mitochondrial	AD	#125250	Optic atrophy plus syndrome, in particular, in cases of biallelic OPA1 mutations	100,101
<i>OPA3</i>		Optic atrophy 3 protein	AR	#258501	Optic atrophy plus syndrome, 3-methylglutaconic aciduria, type III	102
<i>PDHX</i>		Pyruvate dehydrogenase protein X component, mitochondrial	AR	#245349	Lacticacidemia from PDX1 deficiency; often additional mental retardation, delayed psychomotor development and/or seizures	103
<i>PEX16</i>		Peroxisomal membrane protein PEX16	AR	#614877	Peroxisome biogenesis disorder 8B; white matter abnormalities; increased VLCFA	104
<i>PLA2G6</i>	NBIA2A	85/88-kDa calcium-independent phospholipase A2	AR	#256600, #610217	Infantile neuroaxonal dystrophy 1 (INAD); neurodegeneration with brain iron accumulation 2B (NBIA2B), autosomal recessive Parkinson's disease 14 (PARK14)	105,106
<i>PLP1</i>	SCA2	Myelin proteolipid protein	XR	#312920, #312080	Pelizaeus-Merzbacher disease (PMD); X-linked recessive hypomyelinated leukodystrophy (HLD1)	107
<i>PNPLA6</i>	SPG39	Neuropathy target esterase	AR	#612020, #215470, #275400, #245800	Boucher-Neuhauser syndrome, Gordon Holmes syndrome; Oliver-McFarlane syndrome, Laurence-Moon syndrome	12
<i>POLR3A</i>	HLD7	DNA-directed RNA polymerase III subunit RPC1	AR	#607694	Hypomyelinating leukodystrophy 7 (HLD7) with or without oligodontia and/or hypogonadotropic hypogonadism	108,109
<i>POLR3B</i>	HLD8	DNA-directed RNA polymerase III subunit RPC2	AR	#614381	Hypomyelinating leukodystrophy 8 (HLD8) with or without oligodontia and/or hypogonadotropic hypogonadism, but leukodystrophy might also be missing	110
<i>PRNP</i>		Major prion protein	AD	#137440	Familial Creutzfeldt-Jakob disease (CJD), Gerstmann-Strausler disease (GSD), fatal familial insomnia (FFI), but also complicated HSP	111,112

Gene	Locus	Protein name (UniProt)	Inheritance	OMIM	Remarks	Key references
<i>PSAP</i>		Prosaposin	AR	#249900	Metachromatic leukodystrophy from SAP-b deficiency; atypical Krabbe disease; atypical Gaucher disease	113
<i>PSEN1</i>		Presenilin-1	AD	#607822	Early-onset Alzheimer's disease, sometimes complicated by spastic paraparesis and/or ataxia	114
<i>SACS</i>	SPAX6	Sacsin	AR	#270550	Autosomal-recessive spastic ataxia Charlevoix-Saguenay (ARSA), early-onset ataxia with spastic paraparesis and axonal-demyelinating sensorimotor neuropathy; hypointense pontine stripes on T2-MRI	59,115
<i>SCN8A</i>		Sodium channel protein type 8 subunit alpha	AD	#614306	Early infantile epileptic encephalopathy 13 (EIEE13)	116
<i>SDHA</i>		Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial	AR	#252011	Mitochondrial complex II deficiency	117
<i>SETX</i>	SCAR1	Probable helicase senataxin	AR	#606002	AD mutations in SETX are associated with ALS4 (#602433); AR mutations with early-onset ataxia with elevated alpha-fetoprotein	118
<i>SLC17A5</i>		Sialin	AR	#604369, #269920	Sialic acid storage disorder	119
<i>SLC25A15</i>		Mitochondrial ornithine transporter 1	AR	#238970	Hyperornithinemia-hyperammonemia/homocitrullinemia syndrome	120
<i>SLC2A1</i>	DYT9	Solute carrier family 2, facilitated glucose transporter member 1	AD	#612126	GLUT1 deficiency syndrome 1 (GLUT1); DYT9	121
<i>SPG11</i>	SPG11	Spatacsin	AR	#604360	cHSP with thin corpus callosum; juvenile-onset amyotrophic lateral sclerosis-5	122
<i>ZFYVE26</i>	SPG15	Zinc finger FYVE domain-containing protein 26	AR	#270700	cHSP with variable mental retardation, hearing and visual defects, and thin corpus callosum	123
<i>SPG7</i>	SPG7	Paraplegin	AR	#607259	Variable spasticity and cerebellar ataxia	7
<i>SPR</i>		Septapterin reductase	AR	#612716	Dopa-responsive dystonia because of septapterin reductase deficiency	124
<i>STUB1</i>	SCAR16	E3 ubiquitin-protein ligase CHIP	AR	#615768	Spasticity and ataxia can be part of a broader multisystemic neurodegeneration, including hypogonadotropic hypogonadism, and cognitive decline	29,125
<i>SYNE1</i>	SCAR8	Nesprin-1	AR	#610743	Cerebellar ataxia and variable spasticity and further multisystemic neurologic damage	23,24
<i>TBP</i>	SCA17	TATA box-binding protein	AD	#607136	CAG repeat expansion; Huntington's disease-like 4; ataxia, pyramidal and extrapyramidal signs, cognitive impairments, psychosis, and seizures	126
<i>TTC19</i>		Tetratricopeptide repeat protein 19, mitochondrial	AR	#615157	Mitochondrial complex III deficiency nuclear type 2 (MC3DN2); sometimes abnormal signals putamen, caudate, and brain stem on T2-MRI	127
<i>TTPA</i>		Alpha-tocopherol transfer protein	AR	#277460	Ataxia with isolated vitamin E deficiency	128
<i>TUBB4A</i>	HLD6, DYT4	Tubulin beta-4A chain	AD	#612438, #128101	Hypomyelinating leukodystrophy (HLD6); autosomal-dominant dystonia-4 (DYT4)	129
<i>UCHL1</i>	PARK5	Ubiquitin carboxyl-terminal hydrolase isozyme L1	AR	#615491	Childhood-onset neurodegeneration with optic atrophy	130
<i>VAMP1</i>	SPAX1	Vesicle-associated membrane protein 1	AD	#108600	Newfoundland families	131
<i>VWA3B</i>	SCAR22	von Willebrand factor A domain-containing protein 3B	AR	#616948	Intellectual disability associated with adult-onset cerebellar ataxia and spasticity	132

List of genes causing ataxia-spasticity spectrum disease; we anticipate this list to grow considerably in the future (ie, dynamic, extendable list). The selection contains only genes whose phenotypic descriptions include both manifest ataxia and spasticity (rather than merely pyramidal signs) in subjects from at least 2 different families (rather than merely single cases).

OMIM, Online Mendelian Inheritance in Man; AD, autosomal-dominant; AR, autosomal-recessive; XR, x-chromosomal recessive.