

**Original article:**

**EVEROLIMUS, A MAMMALIAN TARGET OF RAPAMYCIN  
INHIBITOR, AMELIORATED STREPTOZOTOCIN-INDUCED  
LEARNING AND MEMORY DEFICITS VIA NEUROCHEMICAL  
ALTERATIONS IN MALE RATS**

Sahar Fanoudi<sup>a,b</sup>, Mahmoud Hosseini<sup>c</sup>, Mohaddeseh Sadat Alavi<sup>a,b</sup>,  
Mohammad Taher Boroushaki<sup>a,b</sup>, Azar Hosseini<sup>b</sup>, Hamid R. Sadeghnia<sup>a-c\*</sup>

<sup>a</sup> Department of Pharmacology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>b</sup> Pharmacological Research Center of Medicinal Plants, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>c</sup> Division of Neurocognitive Sciences, Psychiatry and Behavioral Sciences Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

\* **Corresponding author:** Hamid R. Sadeghnia, Department of Pharmacology, Pharmacological Research Center of Medicinal Plants, Division of Neurocognitive Sciences, Psychiatry and Behavioral Sciences Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, PO Box 99199-91766, Iran, Tel: +98 513 8828566, Fax: +98 513 8828567, E-mail: [sadeghniahr@mums.ac.ir](mailto:sadeghniahr@mums.ac.ir)

<http://dx.doi.org/10.17179/excli2018-1626>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

**ABSTRACT**

Everolimus (EVR), as a rapamycin analog, is a selective inhibitor of the mammalian target of rapamycin (mTOR) kinase and its associated signaling pathway. mTOR is a serine/threonine protein kinase and its hyperactivity is involved in the pathophysiology of Alzheimer's disease (AD) and associated cognitive deficits. The present study evaluated the impact of EVR, on cognitive functions, hippocampal cell loss, and neurochemical parameters in the intracerebroventricular streptozotocin (icv-STZ) model of AD rats. EVR (1 and 5 mg/kg) was administered for 21 days following the single administration of STZ (3 mg/kg, icv) or for 7 days on days 21-28 post-STZ injection after establishment of cognitive dysfunction. Cognitive deficits (passive avoidance and spatial memory), oxidative stress parameters, acetylcholinesterase (AChE) activity, and percentage of cell loss were evaluated in the hippocampus. Chronic administration (1 and 5 mg/kg for 21 days from the day of surgery and icv-STZ infusion) or acute injection (5 mg/kg for 7 days after establishment of cognitive impairment) of EVR significantly attenuated cognitive dysfunction, neuronal loss, oxidative stress and AChE activity in the hippocampus of STZ-AD rats. In conclusion, our study showed that EVR could prevent or improve deteriorations in behavioral, biochemical and histopathological features of the icv-STZ rat model of AD. Therefore, inhibition of the hyperactivated mTOR may be an important therapeutic target for AD.

**Keywords:** Everolimus, mTOR, Alzheimer's disease (AD), oxidative stress, acetylcholinesterase, streptozotocin

## INTRODUCTION

The most prevalent type of dementia is Alzheimer's disease (AD), which is considered as a debilitating neurodegenerative illness, characterized by the progressive cognitive impairments and extensive death of neurons mainly of the cholinergic system. The presence of senile plaques derived from amyloid beta (A $\beta$ ) protein and neurofibrillary tangles (NFT), particularly in the hippocampus or cerebral cortex, are the hallmarks of AD (Ebrahimi et al., 2017). At the present time, the most commonly available medications for AD treatment are acetylcholinesterase (AChE) inhibitors (tacrine, donepezil, rivastigmine and galantamine), which are aimed to improve cognition without any disease modifying properties (Casey et al., 2010; Stuchbury and Münch, 2005). Therefore, identifying novel therapeutic strategies that prevent or reverse the underlying pathological mechanisms of AD are needed.

The mammalian target of rapamycin (mTOR) is a serine/threonine protein kinase that has a key role in various cellular functions. mTOR signaling dysfunction has been involved in the pathophysiology of several neurodegenerative, neurological, and neurodevelopmental disorders (Perluigi et al., 2015; Russo et al., 2012; Wong, 2013). mTOR signaling is necessary for memory retention, synaptic plasticity, neuroendocrine regulation, and neuronal recovery in the central nervous system (CNS) (Blagosklonny, 2012; Chong et al., 2010; Pei and Hugon, 2008; Troca-Marín et al., 2012). However, mTOR hyperactivity is closely associated with the presence of two hallmarks of AD (A $\beta$  plaques and NFT) and cognitive deficits in clinical presentation (Cai et al., 2012; Lafay-Chebassier et al., 2005; Pozueta et al., 2013). The intracerebroventricular (icv) injection of the sub-diabetogenic dose of streptozotocin (STZ) (at a dose of 3 mg/kg) provides a related model for sporadic dementia of AD (Ramezani et al., 2016). It has been reported that icv-STZ induced brain insulin resistance, reduced brain glucose metabolism, oxidative

stress, cholinergic deficits, gliosis, accumulation of tau and A $\beta$  proteins, and learning and memory impairments in rats (Kamat, 2015; Salkovic-Petrisic et al., 2013). In addition, it has been shown that the activity of the hippocampal mTOR system increased two weeks post icv-STZ injection (An et al., 2003; Wang et al., 2014). Therefore, central administration of STZ in rats caused behavioral, histological, and neurochemical changes like human AD features (Dhull et al., 2012; Ponce-Lopez et al., 2011).

Everolimus (EVR), as a rapamycin analog, is a selective mTOR inhibitor, mainly mTORC1 and likely mTORC2, which are two main complexes of mTOR (Hasskarl, 2014; Sarbassov et al., 2006). EVR is clinically used as an immunosuppressive drug in organ transplantation and also as a medication in human cancer treatment (Richardson et al., 2015; Yates, 2016). In a human study on recipients of heart transplant, EVR improved psychiatric symptoms and memory (Lang et al., 2009). Furthermore, rapamycin administration improved memory functions in murine models of AD (Caccamo et al., 2010; Majumder et al., 2012; Spilman et al., 2010). The present study was designed to investigate, for the first time, the effects of EVR as a selective mTOR inhibitor on learning and memory deficits, hippocampal oxidative stress, AChE activity, and neuronal loss in STZ-induced AD rats.

## MATERIAL AND METHODS

### Chemicals

5,5'-Dithiobis-2-nitrobenzoic acid (DTNB), 2-thiobarbituric acid (TBA), HCl, trichloroacetic acid (TCA), ethylenediaminetetraacetic acid disodium salt (Na<sub>2</sub>EDTA), tris (hydroxymethyl) aminomethane (Trizma base), phosphate buffered saline (PBS), and dimethyl sulfoxide (DMSO) were purchased from Merck (Darmstadt, Germany). Everolimus (EVR), streptozotocin, and acetylthiocholine iodide were purchased from Sigma (St. Louis, USA).

## Animals

In this study, male Wistar rats weighing  $200 \pm 20$  g (6-8 weeks of age) at the beginning of the experiments were used. The animals were obtained from the Animal House of the Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran. The animals were kept 4 to 5 per cage under standard temperature ( $22 \pm 2$  °C), and lighting (12 h light/dark cycle) conditions for all the trials. The rats had *ad libitum* access to food and water. All animals were treated in accordance with the National Institutes of Health Guidance for the Care and Use of Laboratory Animals and the Animal Ethics Committee of the Mashhad University of Medical Sciences approved the experimental protocols.

## Experimental protocols

In the present study, two experimental protocols were used (Figure 1). Experiment I was done to determine the therapeutic effect of EVR on STZ-induced AD (groups 1-4). To do so, EVR was intraperitoneally (ip) injected to AD rats on days 21-28 post-STZ injection, one hour before the behavioral study. As our preliminary experiments, along with other studies, showed that learning and memory impairments were established at day 21 post-STZ administration (Ashrafpour et al., 2015; Mehla et al., 2013; Salkovic-Petrisic et al., 2013). Experiment II was carried out to evaluate the effect of the mTORC1 inhibitor on the progress of Alzheimer's disease (groups 1, 2, 5-7). Therefore, EVR was administered

from day 0 [day of icv-STZ injection] to day 20 (Veerendra Kumar and Gupta, 2003).

After acclimatization, the rats were randomly divided into 7 groups (n = 9-12):

Group 1 - sham (icv administration of normal saline + ip injection of vehicle solution)

Group 2 - STZ (STZ 3 mg/kg was administered once, icv)

Group 3 - STZ + EVR 1 mg/kg (for 7 days)


Group 4 - STZ + EVR 5 mg/kg (for 7 days)

Group 5 - STZ + EVR 1 mg/kg (for 21 days)

Group 6 - STZ + EVR 5 mg/kg (for 21 days)

Group 7 - EVR 5 mg/kg (for 21 days).

Passive avoidance paradigm on days 21-22 and Morris water maze test on days 23-28 were done. Twenty-four hours after probe test, the animals were euthanized and their brains were quickly removed. Then the hippocampus was separated and dissected on a glass plate located on ice and stored at  $-80$  °C for biochemical assays. The samples were thawed and homogenized (10 % w/v) in ice-cold 0.1 M phosphate buffer (pH 7.4) to determine of oxidative stress parameters and acetylcholinesterase enzyme (AChE) activity. Hematoxylin and eosin (H & E) staining were done for histopathological examinations.



**Figure 1:** Diagrammatic sketch for the behavioral, biochemical and histopathological experiments. EVR: everolimus; icv-STZ: intracerebroventricular-streptozotocin; PA: passive avoidance; MWM: Morris water maze; SAC: sacrificed for biochemical or histopathological experiments. Day 0 refers to the day of surgery (icv-STZ infusion).

***Surgical procedure: Intracerebroventricular (icv) administration of streptozotocin***

For surgical procedure, the rats were anaesthetized using a combination of ketamine (100 mg/kg, i.p.) and xylazine (20 mg/kg, i.p.). Then, the head of the animal was restrained onto a stereotaxic frame, and a midline sagittal incision was made in the scalp and the skull was exposed. Coordinates for the lateral ventricles (LV) were -0.8 mm posterior and 1.5 mm lateral from the bregma and -4.0 mm below (Paxinos and Watson, 1986). Through these skull holes, a 28-gauge stainless steel needle was lowered into each LV, and STZ was injected slowly and bilaterally at a dose of 3 mg/kg (5  $\mu$ L/side). To prepare the STZ solution, it was freshly dissolved in sterile 0.9 % saline before the injection. The animals in the STZ and STZ + EVR groups received an icv injection of STZ and the sham group received the same volume of sterile saline, icv. After the surgery, the animals were kept separately in individual cages. The stereotaxic coordinates for icv administration was confirmed using an injection of 5  $\mu$ l methylene blue and anatomical observation 24 h after the injection.

***Behavioral study***

Twenty-one days after icv-STZ injection, behavioral tests were started. The experiments were performed at standard optimal conditions between 9:00 am and 4:00 pm in the laboratory (Bassani et al., 2017).

***Passive avoidance test***

The animals underwent a passive avoidance learning test to evaluate their memory retention deficit (Reeta et al., 2009). In this test, the rat was placed in the apparatus, consisting of a light and a dark compartment separated by a guillotine door. On the acquisition trial, the rat was accustomed to the light chamber for 60 s. After that, the guillotine was opened and the initial latency to enter the dark compartment was recorded. Rats that exhibited an initial latency time of more than 60 s were excluded from further experiments. When the rat entered the dark chamber, the

guillotine door was closed and an electric foot shock (1 mA, 2 s duration) was delivered through the grid floor. Then, the rat was removed from the dark chamber and returned to the home cage. Three and 24 hour later, retention latency time was measured in the same way, without the foot shock, and the latency time to enter the dark chamber was recorded up to a maximum of 300 s.

***Morris water maze (MWM) test***

After the passive avoidance test, the rats were submitted to training in the Morris water maze (MWM) apparatus to test their spatial learning and memory (Morris, 1984). It consisted of a black circular tank (136 cm diameter, 60 cm height) filled with water (depth 30 cm;  $24 \pm 1$  °C) and a circular platform (10 cm diameter) was hidden 2 cm below the surface of the water in the center of the southwest quadrant of the pool. Movement of the animals was recorded by a camera above the pool that was connected to a computer. The pool was divided into four equal quadrants, as four different starting positions, labeled North (N), South (S), East (E), and West (W). All four starting points were utilized once in a random sequence. Before the training, each rat was habituated to MWM for 30 s without the platform. For each animal four trials were carried out daily, for 5 consecutive days. Each trial was started when the rat was released into the tank, facing the wall of tank at one of the four random starting positions. The rat was allowed a maximal time of 60 s for swimming to find the hidden platform and to remain on the platform for 15 s. If the rat failed to find the hidden platform within 60 s, it was gently placed on the platform and remained there for 15 s. After each trial, the rat was dried and at the end of the session it was returned to its cage. The time to reach the platform (latency time) and the length of the swimming path (traveled distance) were recorded in training trials by a video tracking system. At 24 h after the acquisition phase, the platform was removed and probe test was conducted, and the rats were given 60 s to swim to explore the

platform. The time spent in the target quadrant and the original position of the platform served as the degree of memory consolidation.

### **Biochemical measurements**

#### *Tissue preparation*

Twenty-four hours after the probe test, the animals were euthanized under deep anesthesia, the brains were quickly taken out, and the hippocampus was separated and dissected on a glass plate located on ice and stored at -80 °C for biochemical assays. Then samples were thawed and homogenized (10 % w/v) in ice-cold 0.1 M phosphate buffer (pH 7.4) to determine oxidative stress parameters, as well as acetylcholinesterase enzyme (AChE) activity.

#### *Oxidative stress markers*

##### *Estimation of lipid peroxidation*

The level of malondialdehyde (MDA) as a lipid peroxidation marker was estimated as described previously (Vafaei et al., 2015; Zabihi et al., 2014). The reaction of MDA with TBA leads to the production of a red complex with a peak absorbance at 535 nm. Briefly, 2 mL of TBA/trichloroacetic acid (TCA)/hydrochloric acid (HCL) reagent was added to 1 mL of the hippocampus homogenate. Then, the solution was placed in a boiling water bath for 40 min. After cooling and centrifugation (1000 g for 10 minutes), the absorbance was measured at 535 nm. The MDA concentration (C) was calculated based on this equation:

$$C (M) = \text{Absorbance} / 1.56 \times 10^5.$$

##### *Estimation of total thiol group*

Total thiol groups were estimated using DTNB (2,2'-dinitro-5,5'-dithiodibenzoic acid), a thiol specific reagent. DTNB reacts with the SH groups and generates a yellow mixture with a peak absorbance at 412 nm. Briefly, 50 µl of the homogenate sample was mixed with 1 ml of Tris-EDTA buffer (pH = 8.6) and the absorbance was measured at 412 nm against the Tris-EDTA buffer alone (A1). Then, 20 µl of the DTNB reagents (10 mM in

methanol) was added to the mixture and after 15 min (while being kept in laboratory temperature), the sample absorbance was read again (A2). The absorbance of the DTNB reagent was also measured as a blank (B). Total thiol concentration (mM) was calculated according to the following equation (Hosseinzadeh and Sadeghnia, 2005):

$$\text{Total thiol concentration (mM)} = (A-A1-B) \times 1.07/0.05 \times 13.6.$$

##### *Cholinesterase activity measurement*

The AChE activity was measured in rat hippocampus using the method of Ellman et al. (1961). Briefly, 0.1 mL of 10 mM DTNB was added to 0.04 mL of homogenate sample and 2.55 mL of PBS. This solution was stored for 5 min at 37 °C and the absorbance was measured at 412 nm. Then, 0.02 mL of 75 mM acetylthiocholine iodide was added, and further incubated at 37 °C for 5 min. Then, the alteration in the absorbance was read at 412 nm. AChE activity was calculated and reported as µmol/g tissue/min (Ellman et al., 1961; Isomae et al., 2002; Zhong et al., 2009).

##### *Histopathological examinations*

The animals were deeply anaesthetized and transcardially perfused with 100 mL of heparinized phosphate buffered saline (PBS), followed by 100 mL of 4 % paraformaldehyde in phosphate buffer (pH 7.4). Brains were carefully removed and postfixed in the same fixative for 24 h, dehydrated, and embedded in paraffin using an automated tissue processor. A total number of six coronal brain sections (5 µm) per rat were used. The sections were then stained with H&E and examined under a Leica DMRB microscope (Leica, India). Photographs were then taken using a Canon PowerShot S70 digital camera (Canon, Japan) at 400x magnification, and cell counts in both the ipsilateral and contralateral sides of the hippocampal areas were scored in a blinded manner (Sadeghnia et al., 2017).

##### *Statistical analysis*

GraphPad Prism (version 6.01, La Jolla, CA) software was used for statistical evaluation. All the data were expressed as mean ±

SEM. Statistical analysis of the results during the five days of the MWM and passive avoidance tests were carried out using repeated measures analysis of variance (ANOVA) with Tukey's post hoc testing. Moreover, statistical analysis of the probe trial and swimming speed in the MWM, MDA, and total thiol contents and AChE activity were performed using one-way ANOVA followed by Tukey's post hoc testing. A statistical  $p$  value  $< 0.05$  was considered significant.

## RESULTS

### *EVR improved STZ-induced memory deficits in the passive avoidance test*

As shown in Figure 2A-B, at baseline, there was no significant difference in the mean initial latency time among all groups. However, the retention latency time was significantly decreased ( $p < 0.001$ ) in the STZ group compared with the sham animals 3 and 24 h after the shock.

Short-term treatment with EVR (5 mg/kg for 7 days) significantly improved retention latency time, as compared with the STZ group ( $p < 0.001$ ). There was no significant difference in the latency time of the EVR group (1 mg/kg for 7 days) when compared with the STZ group (Figure 2A).

However, retention latency was significantly altered towards the control value after the STZ rats were chronically administered with EVR at the doses of 1 mg/kg ( $p < 0.001$ ) and 5 mg/kg ( $p < 0.001$ ) for 21 days (Figure 2B).

EVR improved STZ-induced learning and memory deficits in the Morris water maze test.


Figure 3 shows the decrease in latency time to find the hidden platform in all groups during the five-day training trials in the MWM task. According to Figures 3A-B, STZ rats showed significant higher escape latency than the sham-operated animals from day 2 ( $p < 0.001$ ) to day 5 ( $p < 0.001$ ). Results also revealed that short-term treatment with a high dose of EVR (5 mg/kg for 7 days), and not a low dose (1 mg/kg for 7 days), significantly improved learning performance as compared

to the STZ group (Figure 3A,  $p < 0.05$ ). On the other hand, chronic administration of EVR (1 and 5 mg/kg) for 21 days significantly decreased the mean latency time to find the hidden platform, as compared with the STZ group ( $p < 0.001$ ) (Figure 3B). According to Figs. 4A-B on the probe trial, the STZ rats failed to remember the platform location and spent less time in the target quadrant than the sham group ( $p < 0.001$ ). Figure 4A shows that short-term treatment with EVR (5 mg/kg for 7 days) significantly increased the time spent in the target quadrant, as compared to the STZ group ( $p < 0.001$ ). However, treatment with a low dose of EVR (1 mg/kg for 7 days) failed to improve STZ-induced memory deficits. In the same way, the time spent in the target quadrant was significantly improved following chronic administration of EVR (at both low and high doses), as compared to the STZ group, validating a marked cognitive improvement (Figure 4B,  $p < 0.001$ ). Furthermore, there were no significant differences among the swimming speeds of all groups during the experiments, indicating no sensorimotor effect (Figure 5A-B,  $p > 0.05$ ).


### *Effect of EVR on the hippocampal MDA level*

The degree of free radical damage following the STZ injection was evaluated via lipid peroxidation, which was measured as MDA levels. According to Figure 6, the hippocampal MDA level was significantly increased in the STZ group ( $p < 0.001$ ), in comparison to the sham group.

Treatment with a high dose of EVR (5 mg/kg for 7 days) resulted in a significant reduction in the lipid peroxidation as illustrated by a decrease in the MDA level, in comparison with the STZ group (Figure 6A,  $p < 0.05$ ). Chronic administration of EVR also significantly attenuated the increased MDA levels induced by STZ, at both doses of 1 and 5 mg/kg ( $p < 0.01$  and  $p < 0.001$ , respectively) in the hippocampus (Figure 6B). MDA levels were not changed significantly when EVR (5 mg/kg for 21 days) was given alone, as compared to the sham group (Figure 6B,  $p > 0.05$ ).




**Figure 2:** Effects of 7 days (**A**, on days 21-28 post-STZ infusion, after establishment of cognitive dysfunctions) or 21 days (**B**, from day 0 which refers to the day of surgery and icv-STZ infusion) administration of EVR on passive avoidance memory of STZ-induced AD rats. The latency to enter the dark chamber of the shuttle box apparatus was acquired before, 3 h and 24 h after delivering electrical foot shock (1 mA, 2 s duration). Values are means  $\pm$  SEM. \*\*\* $p < 0.001$  vs. STZ group.



**Figure 3:** Effects of 7 days (A, on days 21-28 post-STZ infusion, after establishment of cognitive dysfunction) or 21 days (B, from day 0 which refers to the day of surgery and icv-STZ infusion) administration of EVR on escape latency of STZ-induced AD rats in Morris water maze (MWM) task. Values are means  $\pm$  SEM. \*\* $p < 0.01$  and \*\*\* $p < 0.001$  vs. STZ group.






**Figure 4:** Effects of 7 days (A, on days 21-28 post-STZ infusion, after establishment of cognitive dysfunction) or 21 days (B, from day 0 which refers to the day of surgery and icv-STZ infusion) administration of EVR on time spent in target quadrant of STZ-induced AD rats in Morris water maze (MWM) task. Values are means  $\pm$  SEM. \*\*\* $p$  < 0.001 vs. STZ group.


#### Effect of EVR on hippocampal thiol levels

The total thiol concentration was measured to evaluate the non-enzymatic defense potential against the oxidative stress. As shown in Figure 7, following AD induction by icv-STZ, the hippocampal thiol concentrations were significantly decreased, as compared to the sham group ( $p$  < 0.001). As illustrated in Figure 7A, treatment with EVR 5

mg/kg ( $p$  < 0.05) significantly restored the decreased sulfhydryl groups in the hippocampus, when compared with the STZ group.




**Figure 5:** Effects of 7 days (A, on days 21-28 post-STZ infusion, after establishment of cognitive dysfunction) or 21 days (B, from day 0 which refers to the day of surgery and icv-STZ infusion) administration of EVR on swimming speed of STZ-induced AD rats in Morris water maze (MWM) task. Values are means  $\pm$  SEM. No significant differences were seen between sham-, STZ and EVR-treated rats.



**Figure 6:** Effects of 7 days (A, on days 21-28 post-STZ infusion, after establishment of cognitive dysfunction) or 21 days (B, from day 0 which refers to the day of surgery and icv-STZ infusion) administration of EVR on malondialdehyde (MDA) level in the hippocampus of STZ-induced AD rats. Values are means  $\pm$  SEM. \* $p < 0.05$ , \*\* $p < 0.01$  and \*\*\* $p < 0.001$  vs. STZ group.

Also, chronic administration of EVR (1 and 5 mg/kg) resulted in a significant increment of total thiol levels, as compared to the STZ group ( $p < 0.01$ ,  $p < 0.001$ , respectively). The levels of thiol were not significantly ( $p >$


0.05) altered by the administration of EVR (5 mg/kg for 21 days) alone, as shown by the absence of significant differences between the EVR group and the sham group (Figure 7B).



**Figure 7:** Effects of 7 days (A, on days 21-28 post-STZ infusion, after establishment of cognitive dysfunction) or 21 days (B, from day 0 which refers to the day of surgery and icv-STZ infusion) administration of EVR on total thiol level in the hippocampus of STZ-induced AD rats. Values are means  $\pm$  SEM. \* $p < 0.05$ , \*\* $p < 0.01$  and \*\*\* $p < 0.001$  vs. STZ group.

### Effect of EVR on activity of acetylcholinesterase

Figure 8A-B shows that there was a significant ( $p < 0.001$ ) increase in the AChE activity in the hippocampus of STZ injected rats, as compared to the sham group. Treatment with 5 mg/kg EVR (for 7 days) significantly decreased the AChE activity vs the STZ group ( $p < 0.01$ ) (Figure 8A).



**Figure 8:** Effects of 7 days (A, on days 21-28 post-STZ infusion, after establishment of cognitive dysfunction) or 21 days (B, from day 0 which refers to the day of surgery and icv-STZ infusion) administration of EVR on acetylcholinesterase (AChE) activity in the hippocampus of STZ-induced AD rats. Values are means  $\pm$  SEM. \*\* $p < 0.01$ , \*\*\* $p < 0.001$  vs. STZ group.

The increase in the activity of AChE was also significantly attenuated by the chronic administration of EVR (1 and 5 mg/kg) when compared to the STZ group (Figure 8B,  $p < 0.01$ ,  $p < 0.001$ , respectively).

### Effect of EVR on hippocampal cell loss


The representative H&E staining of the Cornu Ammonis (CA1) region of the hippocampus in different treatment groups has been shown in Figure 9. The STZ group exhibited greater cell nuclei shrinkage than those of the EVR-treated rats. The STZ group revealed a large number of degenerated cells compared with the STZ-EVR group. Statistical analysis revealed significant neuronal cell loss in the hippocampal CA1 subfield of the AD animals compared with the sham group. Acute injection of EVR at a high dose (5 mg/kg) for 7 days or chronic administration for 21 days significantly attenuated hippocampal cell loss in comparison with the STZ AD rats (Figure 10). In addition, there was no significant difference in the CA1 neuronal cell loss between animals that received EVR (5 mg/kg) alone and the sham group (data not shown).

## DISCUSSION


The present study demonstrated that icv-STZ administration impaired learning and memory, increased oxidative stress, hippocampal cell loss, and induced cholinergic dysfunction in the rat hippocampus, which are consistent with previous studies (Ishrat et al., 2006; Nazifi et al., 2018; Nitsch and Hoyer, 1991; Sharma and Gupta, 2001; Shoham et al., 2007). Our findings revealed that short-term treatment with a high dose of EVR (5 mg/kg for 7 days on days 21-28 post STZ injection when the learning and memory impairments were established) or chronic administration of low (1 mg/kg) or high (5 mg/kg) doses of EVR from day 0 (day of STZ icv injection) to day 20, were able to attenuate STZ-induced learning and memory deficits by inhibiting oxidative stress, amelioration of neuronal injury, and restoration of AChE activity in the hippocampus. The doses and time schedule of the administration of EVR were

selected based on preliminary experiments, as well as, previous studies (Russo et al., 2016;

Sanchez-Fructuoso, 2008; Yokomasu et al., 2009).



**Figure 9:** Effects of 7 days (on days 21-28 post-STZ infusion, after establishment of cognitive dysfunction) or 21 days (from day 0 which refers to the day of surgery and icv-STZ infusion) administration of EVR on cell density of degenerated cells in the hippocampal CA1 region of STZ-induced AD rats.



**Figure 10:** Effects of 7 days or 21 days administration of EVR on percentage of degenerated cells in the hippocampal CA1 region of STZ-induced AD rats

The passive avoidance test was performed for the assessment of learning and memory deficits in the rats. In this task, the cognitive ability of the animals was indicated by avoiding entry into the dark chamber due to the painful experience of the electric foot shock. STZ-administered rats showed reduction in retention latencies in passive avoidance behavior, demonstrating learning and memory impairments. These results are consistent with previous investigations that indicated cognitive impairment after icv-STZ in rats (Javed et al., 2012; Mansouri et al., 2013).

On the other hand, treatment with EVR significantly increased the latency time in two retention trials of the test, which means STZ-induced decreased retention latency was reversed by EVR. The MWM performed in this study is a well-validated behavioral model to evaluate spatial memory and learning in rodents (D'Hooze and De Deyn, 2001). As such, a marked decline in latency during the trials (4 trials/day for 5 days) shows intact acquisition or learning ability. Furthermore, a normal retrieval memory was indicated by increased time spent in the target quadrant. STZ-injected rats showed a significant increase in the latency time and a decrease in time spent in

the target quadrant, whereas improvement of spatial memory by the EVR treatment in STZ-injected rats was confirmed by shorter escape latencies and longer durations in the target quadrant, suggesting enhanced learning ability and memory consolidation by EVR. Our results were supported by recent reports in which EVR improved learning and memory in a mouse model of vascular dementia (Chen et al., 2016) and depression (Russo et al., 2016).

It has been well documented that oxidative stress is an important factor in the pathogenesis of AD and cognitive deficits (Hamilton and Holscher, 2012; Su et al., 2008; Sultana and Butterfield, 2013; Texel and Mattson, 2011). It has been shown that icv-STZ administration caused oxidative stress via an increase in the MDA level and decrease in the glutathione level (Agrawal et al., 2009; Kamat et al., 2016), which were similar to our results. Our findings, along with a previous report, revealed an imbalance in the oxidative and antioxidative system parallel with cognitive dysfunction in STZ rats, indicated that learning and memory impairments are related to oxidative stress in this model (Saxena et al., 2008). In the present study, treatment with EVR attenuated the STZ-induced increase in lipid peroxidation and decrease in the total thiol level, proposing that the protective effects of EVR against STZ might be partly related to increases in antioxidant defense. The mTOR signaling pathway is closely related to oxidative stress or antioxidant capacity. mTOR not only modulates oxidative stress, but is also affected by reactive oxygen species (ROS) (Zhao et al., 2017).

It has been shown that mTOR promotes oxidative stress-induced apoptosis of rat mesangial cells exposed to high glucose, while pretreatment of the cells with rapamycin, as an mTOR inhibitor, ameliorated oxidative stress and reduced the number of apoptotic cells (Lu et al., 2018). Hyperactivation of mTORC1 eventually induces endoplasmic reticulum stress, mitochondrial oxidative stress and activation of apoptosis process

(Wang et al., 2016). Moreover, the higher activity of glutathione reductase has been reported in everolimus-treated mice (Kezic et al., 2013). Pharmacologic inhibition of TORC1 signaling by rapamycin also restored the elevated levels of MDA and 4-hydroxyalkenals in a *Drosophila* model of Friedreich's Ataxia (Calap-Quintana et al., 2015). Our results are also concordant with an earlier report, which revealed the protective effects of rapamycin on A $\beta$ 1-42-induced AD via its antioxidation function (Singh et al., 2017).

Furthermore, we showed that the administration of EVR reversed STZ-induced increase in the AChE activity in the hippocampus. It has been established that the cholinergic system has an essential role in the regulation of learning and memory and alters during aging and AD progression (Anand and Singh, 2013; Lagarde et al., 2017). AChE inhibitors, by inhibiting the hydrolysis of acetylcholine, enhance the cholinergic transmission and temporarily decrease the AD-induced cognitive deficits (Darreh-Shori et al., 2004). In line with our finding, previous studies also showed increased AChE activity and concurrent memory impairments in STZ-induced AD rats (Agrawal et al., 2009; Gutierrez et al., 2014; Saxena et al., 2008; Sharma et al., 2012; Tiwari et al., 2009).

Moreover, it has been shown that activation of autophagy following the administration of rapamycin restored the elevated AChE activity in AD induced by A $\beta$ 1-42 (Singh et al., 2017). mTOR signaling is a pivotal regulator for neural cell metabolism, growth and proliferation, redox status, and autophagy (Maiese, 2014). mTOR complexes (mTORC1 and mTORC2) are directly and indirectly involved in the regulation of autophagy (Abada and Elazar, 2014; Kim et al., 2011). Autophagy is defined as a major degradation pathway in eukaryotic cells that is vital for eliminating damaged organelles and macromolecules from the cytoplasm and reprocessing amino acids in periods of starvation (Mizushima et al., 2008). In fact, autophagy controls cellular stress conditions such as oxidative load, and improves neuronal survival (Damme et al.,

2015). Thus, the imperfect autophagy leads to accumulation of damaged organelles and toxic protein aggregates that contribute to neuronal death and pathogenesis of several neurodegenerative diseases including AD (Li et al., 2010). It has been revealed that autophagy may be a part of pro-survival (PI3K/Akt/mTOR/CREB) signaling, and the activation of autophagy results in the restoration of oxidative defense mechanisms and neuronal damages as well as conserving the integrity of synapse and neurotransmission in the rat model of AD (Singh et al., 2017).

Besides, the continuous activation of the neuronal PI3K/Akt/mTOR pathway has been related with the induction of insulin resistance in the AD brain. It has been indicated that the effect of glucose on the augmentation of memory can be regulated via mTOR signaling (Dash et al., 2006) and in the insulin resistance condition the mTOR pathway is dysregulated (Dann et al., 2007; Zick, 2005). It was also confirmed that the hyperactivation of mTOR causes the inhibition of insulin receptor signaling (Gupta and Dey, 2012). Therefore, there is a molecular link between AD pathology and insulin resistance via mTOR hyperactivation (Barone et al., 2016; Caccamo et al., 2010; Perluigi et al., 2014; Tramutola et al., 2015). mTOR hyperactivation, by blocking autophagy, is involved in the pathologic accumulation of tau and A $\beta$  proteins and associated neurodegeneration (Caccamo et al., 2013; Khurana et al., 2006; Tang et al., 2015).

mTOR is also essential for synaptic plasticity and memory formation (Braak and Braak, 1991), and mTOR signaling hyperactivity has detrimental effects on learning and memory functions (Bolduc et al., 2008). On the other hand, mTOR inhibition could improve learning and memory functions, reduce tau pathology (Caccamo et al., 2010), increase autophagy and decrease the amyloid  $\beta$  level in animal models of AD (Spilman et al., 2010). mTOR inhibition can also increase lifespan and delay age-related cognitive impairments (Johnson et al., 2013; Maiese et al., 2013).

The beneficial effects of EVR seen in this study are in accordance with some previous results from rapamycin, which is able to ameliorate age-dependent learning and memory impairments (Majumder et al., 2012) and reduce cognitive deficits related to neurological disorders such as AD (Spilman et al., 2010) and status epilepticus (Brewster et al., 2013). It has been previously shown that EVR acted as a protector against cognitive decline by increasing neurogenesis in an animal model of depression (Russo et al., 2016). Therefore, mTOR inhibitors might be considered as modifying agents for AD.

In conclusion, our study demonstrated, for the first time, that chronic or acute administration of EVR protected against cognitive impairments and deleterious alterations in oxidative status and AChE activity in the hippocampus of the STZ-AD rats. It presents new opportunities for drug development against AD.

### **Acknowledgment**

This study was financially supported by the Vice Chancellery for Research and Technology, Mashhad University of Medical Sciences (under grant number: 941697). The authors are grateful for this financial support.

### **Conflict of interest**

The authors declare that they have no conflict of interest.

## **REFERENCES**

- Abada A, Elazar Z. Getting ready for building: signaling and autophagosome biogenesis. *EMBO Rep.* 2014; e201439076.
- Agrawal R, Tyagi E, Shukla R, Nath C. A study of brain insulin receptors, AChE activity and oxidative stress in rat model of ICV STZ induced dementia. *Neuropharmacology.* 2009;56:779-87.
- An W-L, Cowburn RF, Li L, Braak H, Alafuzoff I, Iqbal K, et al. Up-regulation of phosphorylated/activated p70 S6 kinase and its relationship to neurofibrillary pathology in Alzheimer's Disease. *Am J Pathol.* 2003; 163:591-607.
- Anand P, Singh B. A review on cholinesterase inhibitors for Alzheimer's disease. *Arch Pharm Res.* 2013; 36:375-99.
- Ashrafpour M, Parsaei S, Sepehri H. Quercetin improved spatial memory dysfunctions in rat model of intracerebroventricular streptozotocin-induced sporadic Alzheimer's disease. *Natl J Physiol Pharm Pharmacol.* 2015;5:411-5.
- Barone E, Di Domenico F, Cassano T, Arena A, Tramutola A, Lavecchia MA, et al. Impairment of biliverdin reductase-A promotes brain insulin resistance in Alzheimer disease: A new paradigm. *Free Radic Biol Med.* 2016;91:127-42.
- Bassani TB, Turnes JM, Moura EL, Bonato JM, C oppola-Segovia V, Zanata SM, et al. Effects of curcumin on short-term spatial and recognition memory, adult neurogenesis and neuroinflammation in a streptozotocin-induced rat model of dementia of Alzheimer's type. *Behav Brain Res.* 2017;335:41-54.
- Blagosklonny MV. Prospective treatment of age-related diseases by slowing down aging. *Am J Pathol.* 2012;181:1142-6.
- Bolduc FV, Bell K, Cox H, Broadie KS, Tully T. Excess protein synthesis in *Drosophila* fragile X mutants impairs long-term memory. *Nat Neurosci.* 2008;11: 1143-5.
- Braak H, Braak E. Neuropathological staging of Alzheimer-related changes. *Acta Neuropathol.* 1991;82: 239-59.
- Brewster AL, Lugo JN, Patil VV, Lee WL, Qian Y, Vanegas F, et al. Rapamycin reverses status epilepticus-induced memory deficits and dendritic damage. *PLoS One.* 2013;8:e57808.
- Caccamo A, Majumder S, Richardson A, Strong R, Oddo S. Molecular interplay between mammalian target of rapamycin (mTOR), amyloid-beta, and Tau: effects on cognitive impairments. *J Biol Chem.* 2010; 285:13107-20.
- Caccamo A, Magri A, Medina DX, Wisely EV, Lopez-Aranda MF, Silva AJ, et al. mTOR regulates tau phosphorylation and degradation: implications for Alzheimer's disease and other tauopathies. *Aging Cell.* 2013; 12:370-80.
- Cai Z, Zhao B, Li K, Zhang L, Li C, Quazi SH, et al. Mammalian target of rapamycin: a valid therapeutic target through the autophagy pathway for Alzheimer's disease? *J Neurosci Res.* 2012;90:1105-18.

- Calap-Quintana P, Soriano S, Llorens JV, Al-Ramahi I, Botas J, Molto MD, et al. TORC1 Inhibition by Rapamycin Promotes Antioxidant Defences in a Drosophila Model of Friedreich's Ataxia. *PLoS One*. 2015; 10:e0132376.
- Casey DA, Antimisiaris D, O'Brien J. Drugs for Alzheimer's disease: are they effective? *P & T*. 2010;35: 208.
- Chen L, Zhang Y, Li D, Zhang N, Liu R, Han B, et al. Everolimus (RAD001) ameliorates vascular cognitive impairment by regulating microglial function via the mTORC1 signaling pathway. *J Neuroimmunol*. 2016; 299:164-71.
- Chong ZZ, Shang YC, Zhang L, Wang S, Maiese K. Mammalian target of rapamycin: hitting the bull's-eye for neurological disorders. *Oxid Med Cell Longev*. 2010;3:374-91.
- D'Hooge R, De Deyn PP. Applications of the Morris water maze in the study of learning and memory. *Brain Res Brain Res Rev*. 2001;36:60-90.
- Damme M, Suntio T, Saftig P, Eskelinen E-L. Autophagy in neuronal cells: general principles and physiological and pathological functions. *Acta Neuropathol*. 2015;129:337-62.
- Dann SG, Selvaraj A, Thomas G. mTOR Complex1-S6K1 signaling: at the crossroads of obesity, diabetes and cancer. *Trends Mol Med*. 2007;13:252-9.
- Darreh-Shori T, Hellstrom-Lindahl E, Flores-Flores C, Guan ZZ, Soreq H, Nordberg A. Long-lasting acetylcholinesterase splice variations in anticholinesterase-treated Alzheimer's disease patients. *J Neurochem*. 2004;88:1102-13.
- Dash PK, Orsi SA, Moore AN. Spatial memory formation and memory-enhancing effect of glucose involves activation of the tuberous sclerosis complex-mammalian target of rapamycin pathway. *J Neurosci*. 2006;26:8048-56.
- Dhull DK, Jindal A, Dhull RK, Aggarwal S, Bhateja D, Padi SS. Neuroprotective effect of cyclooxygenase inhibitors in ICV-STZ induced sporadic Alzheimer's disease in rats. *J Mol Neurosci*. 2012;46:223-35.
- Ebrahimi K, Majdi A, Baghaiee B, Hosseini SH, Sadigh-Eteghad S. Physical activity and beta-amyloid pathology in Alzheimer's disease: A sound mind in a sound body. *EXCLI J*. 2017;16:959-72.
- Ellman GL, Courtney KD, Andres V, Featherstone RM. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem Pharmacol*. 1961; 7:88-95.
- Gupta A, Dey CS. PTEN, a widely known negative regulator of insulin/PI3K signaling, positively regulates neuronal insulin resistance. *Mol Biol Cell*. 2012; 23:3882-98.
- Gutierrez JM, Carvalho FB, Schetinger MR, Marisco P, Agostinho P, Rodrigues M, et al. Anthocyanins restore behavioral and biochemical changes caused by streptozotocin-induced sporadic dementia of Alzheimer's type. *Life Sci*. 2014;96:7-17.
- Hamilton A, Holscher C. The effect of ageing on neurogenesis and oxidative stress in the APP(swe)/PS1(deltaE9) mouse model of Alzheimer's disease. *Brain Res*. 2012;1449:83-93.
- Hasskarl J. Everolimus. *Recent Results Cancer Res*. 2014;201:373-92.
- Hosseinzadeh H, Sadeghnia HR. Safranal, a constituent of *Crocus sativus* (saffron), attenuated cerebral ischemia induced oxidative damage in rat hippocampus. *J Pharm Pharm Sci*. 2005;8:394-9.
- Ishrat T, Khan MB, Hoda MN, Yousuf S, Ahmad M, Ansari MA, et al. Coenzyme Q10 modulates cognitive impairment against intracerebroventricular injection of streptozotocin in rats. *Behav Brain Res*. 2006;171:9-16.
- Isomae K, Ishikawa M, Ohta M, Ogawa Y, Hasegawa H, Kohda T, et al. Effects of T-82, a new quinoline derivative, on cholinesterase activity and extracellular acetylcholine concentration in rat brain. *Jpn J Pharmacol*. 2002;88:206-12.
- Javed H, Khan M, Ahmad A, Vaibhav K, Ahmad M, Khan A, et al. Rutin prevents cognitive impairments by ameliorating oxidative stress and neuroinflammation in rat model of sporadic dementia of Alzheimer type. *Neuroscience*. 2012;210:340-52.
- Johnson SC, Rabinovitch PS, Kaeblerlein M. mTOR is a key modulator of ageing and age-related disease. *Nature*. 2013;493:338-45.
- Kamat PK. Streptozotocin induced Alzheimer's disease like changes and the underlying neural degeneration and regeneration mechanism. *Neural Regen Res*. 2015; 10:1050-2.
- Kamat PK, Kalani A, Rai S, Tota SK, Kumar A, Ahmad AS. Streptozotocin Intracerebroventricular-Induced Neurotoxicity and Brain Insulin Resistance: a Therapeutic Intervention for Treatment of Sporadic Alzheimer's Disease (sAD)-Like Pathology. *Mol Neurobiol*. 2016;53:4548-62.



- Kezic A, Thaïss F, Becker JU, Tsui TY, Bajcetic M. Effects of everolimus on oxidative stress in kidney model of ischemia/reperfusion injury. *Am J Nephrol*. 2013;37:291-301.
- Khurana V, Lu Y, Steinhilb ML, Oldham S, Shulman JM, Feany MB. TOR-mediated cell-cycle activation causes neurodegeneration in a *Drosophila* tauopathy model. *Curr Biol*. 2006;16:230-41.
- Kim J, Kundu M, Viollet B, Guan K-L. AMPK and mTOR regulate autophagy through direct phosphorylation of Ulk1. *Nat Cell Biol*. 2011;13:132-41.
- Lafay-Chebassier C, Paccalin M, Page G, Barc-Pain S, Perault-Pochat MC, Gil R, et al. mTOR/p70S6k signaling alteration by A $\beta$  exposure as well as in APP-PS1 transgenic models and in patients with Alzheimer's disease. *J Neurochem*. 2005;94:215-25.
- Lagarde J, Sarazin M, Chauvire V, Stankoff B, Kas A, Lacomblez L, et al. Cholinergic changes in aging and Alzheimer Disease: An [18F]-F-A-85380 exploratory PET study. *Alzheimer Dis Assoc Disord*. 2017;31:8-12.
- Lang UE, Heger J, Willbring M, Domula M, Matschke K, Tugtekin SM. Immunosuppression using the mammalian target of rapamycin (mTOR) inhibitor everolimus: pilot study shows significant cognitive and affective improvement. *Transplant Proc*. 2009;41:4285-8.
- Li L, Zhang X, Le W. Autophagy dysfunction in Alzheimer's disease. *Neurodegener Dis*. 2010;7:265-71.
- Lu Q, Zhou Y, Hao M, Li C, Wang J, Shu F, et al. The mTOR promotes oxidative stress-induced apoptosis of mesangial cells in diabetic nephropathy. *Mol Cell Endocrinol*. 2018;473:31-43.
- Maiese K. Driving neural regeneration through the mammalian target of rapamycin. *Neural Regen Res*. 2014;9:1413.
- Maiese K, Chong ZZ, Shang YC, Wang S. mTOR: on target for novel therapeutic strategies in the nervous system. *Trends Mol Med*. 2013;19:51-60.
- Majumder S, Caccamo A, Medina DX, Benavides AD, Javors MA, Kraig E, et al. Lifelong rapamycin administration ameliorates age-dependent cognitive deficits by reducing IL-1 $\beta$  and enhancing NMDA signaling. *Aging Cell*. 2012;11:326-35.
- Mansouri MT, Naghizadeh B, Ghorbanzadeh B, Farbood Y, Sarkaki A, Bavarsad K. Gallic acid prevents memory deficits and oxidative stress induced by intracerebroventricular injection of streptozotocin in rats. *Pharmacol Biochem Behav*. 2013;111:90-6.
- Mehla J, Pahuja M, Gupta YK. Streptozotocin-induced sporadic Alzheimer's disease: selection of appropriate dose. *J Alzheimers Dis*. 2013;33:17-21.
- Mizushima N, Levine B, Cuervo AM, Klionsky DJ. Autophagy fights disease through cellular self-digestion. *Nature*. 2008;451:1069.
- Morris R. Developments of a water-maze procedure for studying spatial learning in the rat. *J Neurosci Meth*. 1984;11:47-60.
- Nazifi M, Oryan S, Eslimi Esfahani D, Moghadamnia A, Ashrafpour M. Chronic administration of piperine attenuates behavioural, biochemical and histological alterations induced by intracerebroventricular streptozotocin in rats. *Basic Clin Pharmacol Toxicol*. 2018; epub ahead of print.
- Nitsch R, Hoyer S. Local action of the diabetogenic drug, streptozotocin, on glucose and energy metabolism in rat brain cortex. *Neurosci Lett*. 1991;128:199-202.
- Paxinos G, Watson C. The rat brain in stereotaxic coordinates. New York: Academic Press, 1986.
- Pei JJ, Hugon J. mTOR-dependent signalling in Alzheimer's disease. *J Cell Mol Med*. 2008;12:2525-32.
- Perluigi M, Pupo G, Tramutola A, Cini C, Coccia R, Barone E, et al. Neuropathological role of PI3K/Akt/mTOR axis in Down syndrome brain. *Biochim Biophys Acta*. 2014;1842:1144-53.
- Perluigi M, Di Domenico F, Butterfield DA. mTOR signaling in aging and neurodegeneration: At the crossroad between metabolism dysfunction and impairment of autophagy. *Neurobiol Dis*. 2015;84:39-49.
- Ponce-Lopez T, Liy-Salmeron G, Hong E, Meneses A. Lithium, phenserine, memantine and pioglitazone reverse memory deficit and restore phospho-GSK3 $\beta$  decreased in hippocampus in intracerebroventricular streptozotocin induced memory deficit model. *Brain Res*. 2011;1426:73-85.
- Pozueta J, Lefort R, Shelanski ML. Synaptic changes in Alzheimer's disease and its models. *Neuroscience*. 2013;251:51-65.
- Ramezani M, Darbandi N, Khodaghali F, Hashemi A. Myricetin protects hippocampal CA3 pyramidal neurons and improves learning and memory impairments in rats with Alzheimer's disease. *Neural Regen Res*. 2016;11:1976-80.
- Reeta KH, Mehla J, Gupta YK. Curcumin is protective against phenytoin-induced cognitive impairment and oxidative stress in rats. *Brain Res*. 2009;1301:52-60.

- Richardson A, Galvan V, Lin A-L, Oddo S. How longevity research can lead to therapies for Alzheimer's disease: The rapamycin story. *Exp Gerontol.* 2015;68:51-8.
- Russo E, Citraro R, Constanti A, De Sarro G. The mTOR signaling pathway in the brain: focus on epilepsy and epileptogenesis. *Mol Neurobiol.* 2012;46:662-81.
- Russo E, Leo A, Crupi R, Aiello R, Lippiello P, Spiga R, et al. Everolimus improves memory and learning while worsening depressive- and anxiety-like behavior in an animal model of depression. *J Psychiatr Res.* 2016;78:1-10.
- Sadeghnia HR, Shaterzadeh H, Forouzanfar F, Hosseinzadeh H. Neuroprotective effect of safranal, an active ingredient of *Crocus sativus*, in a rat model of transient cerebral ischemia. *Folia Neuropathol.* 2017;55:206-13.
- Salkovic-Petrisic M, Knezovic A, Hoyer S, Riederer P. What have we learned from the streptozotocin-induced animal model of sporadic Alzheimer's disease, about the therapeutic strategies in Alzheimer's research. *J Neural Transm.* 2013;120:233-52.
- Sanchez-Fructuoso AI. Everolimus: an update on the mechanism of action, pharmacokinetics and recent clinical trials. *Expert Opin Drug Metab Toxicol.* 2008;4:807-19.
- Sarbasov DD, Ali SM, Sengupta S, Sheen JH, Hsu PP, Bagley AF, et al. Prolonged rapamycin treatment inhibits mTORC2 assembly and Akt/PKB. *Mol Cell.* 2006;22:159-68.
- Saxena G, Singh SP, Agrawal R, Nath C. Effect of donepezil and tacrine on oxidative stress in intracerebral streptozotocin-induced model of dementia in mice. *Eur J Pharmacol.* 2008;581:283-9.
- Sharma M, Gupta YK. Intracerebroventricular injection of streptozotocin in rats produces both oxidative stress in the brain and cognitive impairment. *Life Sci.* 2001;68:1021-9.
- Sharma V, Bala A, Deshmukh R, Bedi KL, Sharma PL. Neuroprotective effect of RO-20-1724-a phosphodiesterase4 inhibitor against intracerebroventricular streptozotocin induced cognitive deficit and oxidative stress in rats. *Pharmacol Biochem Behav.* 2012;101:239-45.
- Shoham S, Bejar C, Kovalev E, Schorer-Apelbaum D, Weinstock M. Ladostigil prevents gliosis, oxidative-nitrative stress and memory deficits induced by intracerebroventricular injection of streptozotocin in rats. *Neuropharmacology.* 2007;52:836-43.
- Singh AK, Kashyap MP, Tripathi VK, Singh S, Garg G, Rizvi SI. Neuroprotection through rapamycin-induced activation of autophagy and PI3K/Akt1/mTOR/CREB signaling against amyloid- $\beta$ -induced oxidative stress, synaptic/neurotransmission dysfunction, and neurodegeneration in adult rats. *Mol Neurobiol.* 2017;54:5815-28.
- Spilman P, Podlutskaya N, Hart MJ, Debnath J, Gorostiza O, Bredesen D, et al. Inhibition of mTOR by rapamycin abolishes cognitive deficits and reduces amyloid- $\beta$  levels in a mouse model of Alzheimer's disease. *PLoS One.* 2010;5:e9979.
- Stuchbury G, Münch G. Alzheimer's associated inflammation, potential drug targets and future therapies. *J Neural Transm.* 2005;112:429-53.
- Su B, Wang X, Nunomura A, Moreira PI, Lee HG, Perry G, et al. Oxidative stress signaling in Alzheimer's disease. *Curr Alzheimer Res.* 2008;5:525-32.
- Sultana R, Butterfield DA. Oxidative modification of brain proteins in Alzheimer's disease: perspective on future studies based on results of redox proteomics studies. *J Alzheimers Dis.* 2013;33(Suppl 1):S243-51.
- Tang Z, Ioja E, Bereczki E, Hultenby K, Li C, Guan Z, et al. mTOR mediates tau localization and secretion: Implication for Alzheimer's disease. *Biochim Biophys Acta.* 2015;1853:1646-57.
- Texel SJ, Mattson MP. Impaired adaptive cellular responses to oxidative stress and the pathogenesis of Alzheimer's disease. *Antioxid Redox Signal.* 2011;14:1519-34.
- Tiwari V, Kuhad A, Bishnoi M, Chopra K. Chronic treatment with tocotrienol, an isoform of vitamin E, prevents intracerebroventricular streptozotocin-induced cognitive impairment and oxidative-nitrosative stress in rats. *Pharmacol Biochem Behav.* 2009;93:183-9.
- Tramutola A, Triplett JC, Di Domenico F, Niedowicz DM, Murphy MP, Coccia R, et al. Alteration of mTOR signaling occurs early in the progression of Alzheimer disease (AD): analysis of brain from subjects with pre-clinical AD, amnesic mild cognitive impairment and late-stage AD. *J Neurochem.* 2015;133:739-49.
- Troca-Marín JA, Alves-Sampaio A, Montesinos ML. Deregulated mTOR-mediated translation in intellectual disability. *Prog Neurobiol.* 2012;96:268-82.

- Vafae F, Hosseini M, Hassanzadeh Z, Edalatmanesh MA, Sadeghnia HR, Seghatoleslam M, et al. The effects of *Nigella sativa* hydro-alcoholic extract on memory and brain tissues oxidative damage after repeated seizures in rats. *Iranian journal of pharmaceutical research: IJPR*. 2015;14:547.
- Veerendra Kumar M, Gupta Y. Effect of *Centella asiatica* on cognition and oxidative stress in an intracerebroventricular streptozotocin model of Alzheimer's disease in rats. *Clin Exp Pharmacol Physiol*. 2003;30:336-42.
- Wang C, Yu JT, Miao D, Wu ZC, Tan MS, Tan L. Targeting the mTOR signaling network for Alzheimer's disease therapy. *Mol Neurobiol*. 2014;49:120-35.
- Wang J, Yang X, Zhang J. Bridges between mitochondrial oxidative stress, ER stress and mTOR signaling in pancreatic beta cells. *Cell Signal*. 2016;28:1099-104.
- Wong M. Mammalian target of rapamycin (mTOR) pathways in neurological diseases. *Biomed J*. 2013;36:40-50.
- Yates DH. mTOR treatment in lymphangioleiomyomatosis: the role of everolimus. *Expert Rev Respir Med*. 2016;10:249-60.
- Yokomasu A, Yano I, Sato E, Masuda S, Katsura T, Inui K. Effect of itraconazole on the pharmacokinetics of everolimus administered by different routes in rats. *Biopharm Drug Dispos*. 2009;30:517-23.
- Zabihi H, Hosseini M, Pourganji M, Oryan S, Soukhtanloo M, Niazmand S. The effects of tamoxifen on learning, memory and brain tissues oxidative damage in ovariectomized and naive female rats. *Adv Biomed Res*. 2014;3:219.
- Zhao D, Yang J, Yang L. Insights for oxidative stress and mTOR signaling in myocardial ischemia/reperfusion injury under diabetes. *Oxid Med Cell Longev*. 2017;2017:6437467.
- Zhong S-Z, Ge Q-H, Qu R, Li Q, Ma S-P. Paeonol attenuates neurotoxicity and ameliorates cognitive impairment induced by d-galactose in ICR mice. *J Neurol Sci*. 2009;277:58-64.
- Zick Y. Ser/Thr phosphorylation of IRS proteins: a molecular basis for insulin resistance. *Sci STKE*. 2005;2005:pe4.