



HHS Public Access

Author manuscript

J Clin Psychiatry. Author manuscript; available in PMC 2018 December 17.

Published in final edited form as:

J Clin Psychiatry. 2016 October ; 77(10): 1413–1419. doi:10.4088/JCP.15m09963.

Psychiatric disorders among patients seeking treatment for co-occurring chronic pain and opioid use disorder

Declan T. Barry, Ph.D.^{a,b}, Christopher J. Cutter, Ph.D.^{a,b}, Mark Beitel, Ph.D.^{a,b}, Robert D. Kerns, Ph.D.^{a,c}, Christopher Liang, B.A.^b, and Richard S. Schottenfeld, M.D.^a

^aYale University School of Medicine, New Haven, CT 06519 USA

^bThe APT Foundation Pain Treatment Services, New Haven, CT 06519 USA

^cVA Connecticut Healthcare System, West Haven, CT 06516 USA

Abstract

Objective: Psychiatric comorbidities complicate treatment of patients with chronic pain and opioid use disorder, but the prevalence of specific comorbid psychiatric disorders in this population has not been systematically investigated.

Methods: 170 consecutive participants entering a treatment research program for co-occurring chronic pain and opioid use disorder between March 2009 and July 2013 were evaluated with the Structured Clinical Interview for DSM-IV-TR Axis I Disorders (SCID-I/P) and the Diagnostic Interview for DSM-IV Personality Disorders (DIPD-IV).

Results: The prevalence of any lifetime (and current) comorbid Axis I disorder was 91% (75%); 52% met criteria for lifetime anxiety disorder (48% current), 57% for lifetime mood disorder (48% current), and 78% for lifetime non-opioid substance use disorder (34% current). Common current anxiety diagnoses were posttraumatic stress disorder (21%), generalized anxiety disorder (16%), and panic disorder without agoraphobia (16%). Common current mood diagnoses were major depressive disorder (40%) and dysthymia (11%). A majority of patients had a personality disorder (52%).

Conclusion: High rates and persistence of co-occurring psychiatric disorders, including anxiety or mood disorders, may explain in part the difficulty providers have treating patients with co-occurring opioid use disorder and chronic pain, and suggest possible targets for improving treatment.

Keywords

chronic pain; opioid-related disorders; mental disorders

Correspondence concerning this article should be addressed to: Declan T. Barry, Ph.D., Yale University School of Medicine, CMHC/SAC Room 220, 34 Park Street, New Haven, CT 06519-1187 USA. Phone: (203) 285-2708. Fax: (203) 781-4861. declan.barry@yale.edu.

Role of the sponsors: The sponsors had no role in the design or conduct of the study. The views expressed in this article are those of the authors and do not necessarily represent the position or policy of the Department of Veterans Affairs or NIDA.

INTRODUCTION

The use of opioid analgesics to treat chronic non-cancer pain (i.e., pain lasting at least three months) has increased three-fold since the early 1990s and has brought with it an epidemic of nonmedical opioid use, opioid overdose, and opioid use disorder¹⁻⁵. Patients with co-occurring chronic pain and opioid use disorder present considerable management challenges: Unrelieved pain is associated with anxiety and depression, sleep problems, and opioid or other substance misuse, often to self-medicate⁶⁻⁹. Clinicians report difficulty managing patients with chronic pain and opioid use disorder because of the perceived high rates of accompanying psychopathology, a dearth of evidence-based treatments for these co-existing conditions, and a paucity of providers who specialize in their treatment¹⁰⁻¹². Given their expertise in psychiatric assessment and treatment, prescribing adjuvant psychotropic pain medications (e.g., antidepressants), and the biopsychosocial model, psychiatrists can perform a pivotal role in the treatment of chronic pain and opioid use disorder¹³⁻¹⁵. However, many may feel ill-equipped to do so; medical schools and psychiatry residency programs in the U.S. provide limited training on managing chronic pain, especially in the context of an opioid use disorder^{13,16}. Increased knowledge about the psychiatric comorbidity of co-occurring opioid use disorder and chronic pain might assist psychiatrists in treating or consulting on this vulnerable patient population.

High rates of psychiatric disorders have been found among patients with chronic pain¹⁷⁻²² and with opioid use disorder²³⁻²⁵, and co-occurring psychopathology in patients with these disorders has important treatment implications^{26,27}. Relatively few studies have investigated psychopathology among patients with co-occurring opioid use disorder and chronic pain, however. Elevated levels of anxiety and depression have been reported in surveys of methadone-maintained patients with chronic pain^{6,9,28,29}, and one small study found high rates of Axis-I disorders among 40 patients with chronic pain referred from multidisciplinary pain clinics who also had either opioid abuse (n=12) or dependence (n=28)³⁰. Although anxiety and depression can exacerbate pain in non-addicted patients with chronic pain^{26,31}, the association of pain with anxiety disorders or with mood disorders has not been systematically examined in those with co-occurring opioid use disorder and chronic pain.

Consequently, this study evaluated the prevalence of DSM-IV-TR³² Axis I and Axis II disorders and the association between psychiatric disorders and pain intensity or pain interference (i.e., decrement in functioning resulting from pain) among patients with co-occurring opioid use disorder and chronic pain entering either of two clinical trials combining behavioral interventions and opioid agonist maintenance treatment with buprenorphine/naloxone or methadone. We anticipated that rates of Axis I psychiatric disorders would be even higher among patients with co-occurring opioid use disorder and chronic pain than those previously reported among patients with either condition alone. Finally, we hypothesized that in comparison to patients with neither an anxiety disorder nor a mood disorder, those with an anxiety disorder, a mood disorder, or both would have higher pain intensity and pain interference.

METHOD

Participants

Participants were a consecutive series of 170 adults who completed evaluations for enrollment in a treatment research program for co-occurring chronic pain and opioid dependence in New Haven, Connecticut, between March 2009 and July 2013. Participants were recruited through referrals from providers and intake workers at primary care centers, pain management clinics, and opioid treatment programs; advertisements (e.g., radio, Internet, posters in the community); and word-of-mouth. One hundred and thirteen participants were evaluated for a research protocol involving office-based buprenorphine/naloxone treatment (NCT006348030), while the remaining 57 were assessed for a research protocol involving opioid treatment program-based methadone maintenance treatment (NCT00727675). Participants were administered the diagnostic interviews following medication induction and initial stabilization (generally one to two weeks following initiation of either buprenorphine/naloxone or methadone). The study received appropriate Institutional Review Board and institutional approval (at Yale University School of Medicine and at APT Foundation). Participants provided voluntary, written informed consent. The study design and conduct complied with the Declaration of Helsinki.

To be eligible for study inclusion, participants needed to: 1) be at least 18 years old; 2) be able to read and write in English; 3) experience moderate to severe chronic low back pain (ratings of pain at its worst or peak in past week 4 on a 0 [no pain]-10 [worst pain imaginable] numeric rating scale) of at least six months duration and classified as non-specific in origin, according to the guidelines issued by the American College of Physicians (ACP) and the American Pain Society (APS)³³; and 4) meet DSM-IV-TR criteria for opioid dependence³² and joint American Academy of Pain Medicine-American Pain Society-American Society of Addiction Medicine (AAPM-APS-ASAM) criteria for opioid addiction, which require compulsive use, continued use despite harm, or craving³⁴. Participants could have multiple sites of pain, but low back pain was required to be a primary pain site. Exclusion criteria included drug treatment in the prior 30 days and current suicide or homicide risk. Participants received a comprehensive evaluation from a study physician prior to admission to ensure study eligibility.

Demographic and Clinical Characteristics

Participants provided information on demographics (age, gender, race, ethnicity, education, employment, marital status) and clinical characteristics (HIV status, primary drug of abuse [heroin, opioid medication, other], outpatient mental health visit in the past month [yes/no] and whether they had taken prescribed psychiatric medication in the past month [yes/no]).

Pain Characteristics

Participants provided information on current pain location and genesis, and the duration of chronic pain. Participants also completed the Brief Pain Inventory (BPI)^{35,36}, where they rated four facets of pain experienced in the past seven days (i.e., “pain at its worst,” “pain at its least,” “pain on average,” “pain right now”) using 0 (“no pain at all”) to 10 (“pain as bad as you can imagine”) scales, and seven pain interference items that assessed the extent to

which their pain in the last seven days had interfered with their “general activity,” “mood,” “walking ability,” “normal work (includes both outside the home and housework),” “relations with other people,” “sleep,” and “enjoyment of life” using 0 (“does not interfere”) to 10 (“completely interferes”) scales. The mean of the four pain intensity ratings was calculated to find the “pain severity score,” while the mean of the seven interference items was calculated to determine the “pain interference score” (both scored on 0 to 10 scales)³⁷.

Diagnostic Interviews

The Structured Clinical Interview for DSM-IV-TR Axis I Disorders (SCID-I/P)³⁸ includes modules on current (last 30 days) and lifetime (either current or past) mood, anxiety, and substance use disorders, while the Diagnostic Interview for DSM-IV Personality Disorders (DIPD-IV)³⁹ collects diagnostic information pertaining to the last 2 years (i.e., current only) on Cluster A (paranoid, schizoid, schizotypal), Cluster B (antisocial, borderline, histrionic, narcissistic), and Cluster C (avoidant, dependent, obsessive-compulsive) personality disorders.

Interviewers

The diagnostic interviews were conducted by addiction psychiatry and psychology fellows as well as clinical and counseling psychologists. A standardized training protocol was followed: research clinicians attended didactic workshops; observed two batteries performed by experienced assessors; conducted two supervised batteries; and subsequently conducted the diagnostic evaluations independently, receiving ongoing supervision as needed.

Data Analysis

Descriptive statistics were used to calculate demographic characteristics and the prevalence of psychiatric disorders. T-tests and chi-square tests were performed to compare baseline characteristics between patients from the buprenorphine/naloxone and methadone maintenance trials. Subsequent analyses examined the overall sample, using univariate analysis of variance (ANOVA) tests to compare pain severity scores and pain interference scores in patients with (a) co-occurring anxiety and mood disorders, (b) anxiety or mood disorder (but not both), and (c) no anxiety or mood disorders. Separate 2X2 chi-square tests were performed for participants in the buprenorphine/naloxone and methadone maintenance trials to determine whether the presence of any current or lifetime anxiety, mood, or non-opioid substance use disorder, or any current personality disorder was associated with study completion or clinically significant changes in pain severity or pain interference (defined as 2 decrease on 0–10 scales from baseline to end of treatment). Chi-square tests were also performed to examine differences between participants whose reported primary drug of abuse was heroin or prescription opioids on the presence of lifetime and current psychiatric disorders. The Fisher’s Exact test was used in cases where the expected cell frequencies for the chi-square tests was less than 5. Statistical significance was set at $p < 0.05$. All statistics were performed on IBM® SPSS Statistics Version 19.

RESULTS

Demographic and Clinical Characteristics

The 170 participants ranged in age from 19 to 64 years old ($M=36.0$, $SD=10.0$); 71% were men; 87% were white; 47% were employed; 87% had greater than or equal to a high school education; 20% were married; and 51% reported opioid medications as their primary drug of abuse (Table 1). The average ages of onset of chronic pain and opioid dependence were 23.7 and 25.3, respectively. On average, participants in the buprenorphine/naloxone trial, in comparison to those in the methadone maintenance trial, were ($p<0.05$) younger (34.6 vs. 38.8 years old); more likely to be employed (56% vs. 28%) and report that opioid medication was their primary drug of abuse (61% vs. 30%); and less likely to report that they had been prescribed psychiatric medication in the past month (11% vs. 23%).

Pain Characteristics

The duration of pain ranged from <1 to 58 years ($M=13.2$, $SD=10.6$). Causes of current pain included accident (63%), nerve pain (22%), arthritis (18%), surgery (17%), opioid withdrawal (16%), work (14%), HIV (0.6%), and “other” (12%). Inclusive of low back, the number of reported pain sites ranged from 1 to 10 ($M=3.0$, $SD=1.8$), and comprised legs (48%), shoulder (29%), feet (21%), neck (20%), hands (17%), head (15%), stomach (14%), knees (14%), arms (13%), hips (9%), pelvis (9%), teeth (2%), face (1%), and “other” (6%). Participants’ average pain severity and pain interference scores were 5.6 ($SD=1.5$) and 6.0 ($SD=2.3$), respectively.

Psychiatric Disorder Prevalence, Persistence, Comorbidity, and Treatment

The prevalence of any lifetime (and current) comorbid Axis I disorder was 91% (75%). As summarized in Table 2, 52% met criteria for lifetime anxiety disorder (48% current), 57% for lifetime mood disorder (48% current), and 78% for lifetime non-opioid substance use disorder (34% current). A majority of patients had an Axis II disorder (52%). As summarized in Table 3, patients whose primary reported drug of abuse was heroin (compared to prescription opioids) were more likely to meet lifetime criteria for any Axis I disorder and any non-opioid substance use disorder as well as any current Axis II disorder.

The proportion of participants who met lifetime criteria for zero, one, two, or at least three comorbid psychiatric disorders (i.e., any anxiety disorder, any mood disorder, any non-opioid substance use disorder, any Axis II disorder) was as follows: 9%, 20%, 20%, and 51%, respectively, while the proportion who met current criteria were: 19%, 22%, 25% and 34%, respectively. About two-thirds of participants met criteria for current co-occurring anxiety or mood disorders (65%), including 33% who met criteria for both disorders. Participants’ average ages of onset for anxiety disorders and mood disorders were 16.9 ($SD=10.8$) and 23.3 ($SD=10.3$) years, respectively; the average durations of current anxiety disorders and current mood disorders were 12.3 ($SD=10.7$) and 18.2 ($SD=12.5$) years, respectively. Of participants meeting criteria for a lifetime anxiety disorder, 93% also met criteria for a current anxiety disorder; 84% of patients meeting criteria for a lifetime mood disorder also met criteria for a current mood disorder. In the month prior to baseline, 4% of participants reported attending a mental health appointment, 15% reported that they had

taken a prescribed psychiatric medication, and 16% reported either having a mental health visit and/or taking a prescribed psychiatric medication.

Anxiety and Mood Disorders, Pain Severity, Pain Interference

Among participants with current co-occurring anxiety and mood disorders, either a current anxiety or mood disorder (but not both), or no current anxiety or mood disorders, there were no significant differences found in pain severity score (5.7 vs. 5.7 vs. 5.5, respectively) ($F=0.53$, $df=2$, 167 , $p=0.58$) or pain interference score (6.3 vs. 6.0 vs. 5.7, respectively) ($F=1.14$, $df=2$, 167 , $p=0.32$). Treatment retention and clinically significant changes in pain severity or pain interference were not significantly associated with the presence of current or lifetime psychiatric disorders (p values ranged from 0.12 to 1.00 and 0.06 to 1.00, respectively for participants in the buprenorphine/naloxone and methadone maintenance trials).

DISCUSSION

This study found a very high prevalence of DSM-IV-TR Axis-I and Axis-II disorders among patients seeking treatment for co-occurring opioid use disorder and chronic pain. Most participants (81%) met current criteria for at least one—and the majority (59%) for at least two—of the following psychiatric comorbidities: anxiety disorder, mood disorder, non-opioid substance use disorder, or personality disorder. Notably, a high proportion of participants with lifetime psychiatric disorders also met current diagnostic criteria. Despite the high prevalence and persistence of psychopathology, few participants had received either mental health treatment (3.5%) or a psychiatric medication in the past month (14.8%). Contrary to findings among patients with chronic pain but without co-occurring opioid use disorder^{26,31}, in this study, the presence of an anxiety disorder, mood disorder, or both, was not associated with higher pain severity or pain interference scores.

To put the findings of the current study in context, the prevalence of current anxiety disorders (48%) among study participants was higher than found in prior studies^a of patients with opioid dependence (5–17%)^{23,24} or in studies of patients with chronic pain (11–25%)^{19–21}. (Because of differences in study methodology, we did not make direct statistical comparisons between the current and prior studies.) Similarly, the prevalence of current mood disorders (48%) among study patients was higher than that found in studies of patients with opioid dependence (4–24%)^{23–25} and at the higher end of the range of those reported in studies of patients with chronic pain (34–56%)^{18,40}. The prevalence of current non-opioid substance use disorders (34%) was in the range found in previous studies of patients with opioid use disorders (40–68%)^{24,25,41}, but substantially higher than the estimated rates of substance use disorders reported in studies of patients with chronic pain (11–19%)^{20–22,42}. The prevalence of personality disorders (52%) was also in the range reported in prior studies of patients with opioid use disorders (35–57%)^{23–25,43} or chronic pain (41–69%)^{18,20–22,44,45}. Our findings are consistent with a previous investigation of a group of 40 patients with co-occurring opioid use disorders and chronic pain, which reported

^aIn these comparisons, we looked at studies using either DSM-III-R, DSM-IV, or DSM-IV-TR diagnostic criteria for the SCID (Axis I/Axis II) and DIPD (Axis II) to determine diagnostic prevalence.

numerically even higher rates of lifetime anxiety (93%) and mood disorders (85%) than those found in this study³⁰.

This study is among the first to document higher rates of any lifetime Axis I disorder, any lifetime non-opioid substance use disorder, and any current Axis II disorder among patients with co-occurring chronic pain and opioid dependence reporting heroin (compared to prescription opioids) as their primary drug of abuse, and this finding merits further investigation. While patients with an anxiety and/or mood disorder had numerically higher pain intensity and interference than their counterparts without these disorders, group differences did not reach the level of statistical significance, but may have done so with a larger sample size. However, it is also possible that the presence of an opioid (or non-opioid substance) use disorder attenuates the relationship previously reported between pain intensity or interference and the presence of anxiety or mood disorders among individuals with chronic pain^{26,31}. One striking finding in this study was the high rate of non-opioid substance-related disorders; the extent to which the order of onset of chronic pain and opioid use disorder is associated with the prevalence of psychopathology, including non-opioid substance use disorders is currently unclear and should be further investigated.

Another important finding of the current study is the high persistence of lifetime anxiety or mood disorders into the present: 93% of patients meeting criteria for a lifetime anxiety disorder also met criteria for a current anxiety disorder, and 84% of patients meeting criteria for a lifetime mood disorder met criteria for a current mood disorder. The high prevalence of current psychiatric comorbidity at treatment entry as well as the persistence of these disorders, as indicated by the prolonged average duration of anxiety disorders (more than 12 years) and mood disorders (more than 18 years), may explain in part the clinical management problems experienced by providers treating these patients^{10,11}. The findings suggest targets for potential pharmacological or psychosocial treatments (e.g., duloxetine and other antidepressant medications⁴⁶ and cognitive-behavioral therapy⁴⁷⁻⁴⁹ have demonstrated efficacy in treating anxiety disorders, mood disorders, and chronic pain). It is notable, however, that neither lifetime nor current psychiatric disorders in the buprenorphine/naloxone or methadone maintenance trials were associated with study completion or clinically significant changes in pain severity or pain interference. This supports the conceptualization of these patients as persons with multiple interrelated morbidities who may benefit from the development and evaluation of integrated patient-centered interventions targeting chronic pain, opioid dependence, and other psychopathology.

Considering the high level of psychiatric comorbidity, it is noteworthy that a minority of participants (16%) reported past-month receipt of a psychiatric medication or a mental health treatment episode. This low treatment rate may be related to multiple factors, including: a) failure of healthcare professionals to identify and treat psychiatric disorders in this patient group; b) appropriate concerns about the use of benzodiazepines to treat anxiety disorders in this population; c) lack of access to psychiatrists or psychologists with expertise in treating co-occurring opioid use disorder and chronic pain in addition to other psychiatric disorders; d) logistical barriers (e.g., financial cost); and e) patient-level factors (e.g., lack of awareness of having a potentially treatable psychiatric disorder rather than simply

experiencing symptoms of anxiety or depression related to chronic pain or opioid use disorder).

Several potential study limitations are worth noting. We did not collect data on the reliability of the study diagnoses (e.g., test-retest, inter-rater), but use of doctoral-level experienced clinicians as interviewers mitigates this concern. We used DSM-IV-TR³² (and not DSM-5⁵⁰) diagnoses. Given the importance of the construct of craving in both DSM-5 criteria for opioid use disorder⁵⁰ and joint AAPM-APS-ASAM criteria for opioid addiction³⁴, future research might benefit from a more detailed assessment of opioid craving^{51,52}. The extent to which anxiety disorders and mood disorders among participants were primary, or a function of chronic pain or substance use, is unclear. Since participants were seeking treatment for co-occurring chronic pain and opioid dependence, it is unclear whether findings generalize to non-treatment-seeking individuals with these co-existing conditions.

Among patients seeking treatment for opioid use disorder and co-occurring chronic pain, the high rates of current and lifetime Axis I and Axis II disorders, the apparent absence of remission of anxiety disorders and mood disorders and the low rates of current mental health treatment are salient. Overall, our findings suggest that effective treatments for co-occurring chronic pain and opioid dependence may need to address co-occurring psychiatric disorders, especially given their apparent chronicity, and that strategies for improving access to psychiatric and mental health services for these patients are needed.

Acknowledgments

Conflicts of Interest and Source of Funding: This research was supported by funding from the National Institute on Drug Abuse to Dr. Barry (K23 DA024050) and Dr. Schottenfeld (K24 DA000445, R01DA024695). Dr. Kerns was supported by the Veterans Health Administration Health Services Research and Development Service Center of Innovation (CIN 13–407). None of the authors are affiliated with the tobacco, alcohol, pharmaceutical or gambling industry in a manner that we believe represents a conflict of interest with the current manuscript. The findings of this study were presented in part at the 75th Annual Scientific Meeting of the College on Problems of Drug Dependence—San Diego, CA, June 2013.

REFERENCES

1. Reuben DB, Alvanzo AA, Ashikaga T, et al. National institutes of health pathways to prevention workshop: the role of opioids in the treatment of chronic pain. *Ann Intern Med* 2015; 162(4): 295–300. [PubMed: 25581341]
2. Okie S A flood of opioids, a rising tide of deaths. *N Engl J Med* 2010; 363(21): 1981–5. [PubMed: 21083382]
3. Dart RC, Surratt HL, Cicero TJ, et al. Trends in opioid analgesic abuse and mortality in the United States. *N Engl J Med* 2015; 372(3): 241–8. [PubMed: 25587948]
4. Dunn KM, Saunders KW, Rutter CM, et al. Opioid prescriptions for chronic pain and overdose: a cohort study. *Ann Intern Med* 2010; 152(2): 85–92. [PubMed: 20083827]
5. Gomes T, Mamdani MM, Dhalla IA, Cornish S, Paterson JM, Juurlink DN. The burden of premature opioid-related mortality. *Addiction* 2014; 109(9): 1482–8. [PubMed: 25041316]
6. Barry DT, Beitel M, Garnet B, Joshi D, Rosenblum A, Schottenfeld RS. Relations among psychopathology, substance use, and physical pain experiences in methadone-maintained patients. *J Clin Psychiatry* 2009; 70(9): 1213–8. [PubMed: 19607760]
7. Barry DT, Beitel M, Joshi D, Schottenfeld RS. Pain and substance-related pain reduction behaviors among opioid dependent individuals seeking methadone maintenance treatment. *Am J Addict* 2009; 18(2): 117–21. [PubMed: 19283562]

8. Peles E, Schreiber S, Adelson M. Documented poor sleep among methadone-maintained patients is associated with chronic pain and benzodiazepine abuse, but not with methadone dose. *Eur Neuropsychopharmacol* 2009; 19(8): 581–8. [PubMed: 19411166]
9. Rosenblum A, Joseph H, Fong C, Kipnis S, Cleland C, Portenoy RK. Prevalence and characteristics of chronic pain among chemically dependent patients in methadone maintenance and residential treatment facilities. *JAMA* 2003; 289(18): 2370–8. [PubMed: 12746360]
10. Barry DT, Bernard MJ, Beitel M, Moore BA, Kerns RD, Schottenfeld RS. Counselors' experiences treating methadone-maintained patients with chronic pain: A needs assessment study. *J Addict Med* 2008; 2(2): 108–11. [PubMed: 21768980]
11. Barry DT, Irwin KS, Jones ES, et al. Opioids, chronic pain, and addiction in primary care. *J Pain* 2010; 11(12): 1442–50. [PubMed: 20627817]
12. Berg K, Arnsten J, Sacajiu G, Karasz A. Providers' experiences treating chronic pain among opioid-dependent drug users. *J Gen Intern Med* 2009; 24(4): 482–8. [PubMed: 19189194]
13. Elman I, Zubietta JK, Borsook D. The missing p in psychiatric training: why it is important to teach pain to psychiatrists. *Arch Gen Psychiatry* 2011; 68(1): 12–20. [PubMed: 21199962]
14. Krashin D, Murinova N, Ballantyne J. Management of pain with comorbid substance abuse. *Curr Psychiatry Rep* 2012; 14(5): 1–7. [PubMed: 22076689]
15. Cheatle MD, Gallagher RM. Chronic pain and comorbid mood and substance use disorders: A biopsychosocial treatment approach. *Curr Psychiatry Rep* 2006; 8(5): 371–6. [PubMed: 16968617]
16. Mezei L, Murinson BB. Pain education in North American medical schools. *J Pain* 2011; 12(12): 1199–208. [PubMed: 21945594]
17. McWilliams LA, Cox BJ, Enns MW. Mood and anxiety disorders associated with chronic pain: an examination in a nationally representative sample. *Pain* 2003; 106(1–2): 127–33. [PubMed: 14581119]
18. Gatchel RJ, Garofalo J, Ellis E, Holt C. Major psychological disorders in acute and chronic TMD: an initial examination. *J Am Dent Assoc* 1996; 127(9): 1365–70. [PubMed: 8854613]
19. Knaster P, Karlsson H, Estlander A-M, Kalso E. Psychiatric disorders as assessed with SCID in chronic pain patients: the anxiety disorders precede the onset of pain. *Gen Hosp Psychiatry* 2012; 34(1): 46–52. [PubMed: 22001549]
20. Dersh J, Gatchel RJ, Mayer T, Polatin P, Temple OR. Prevalence of psychiatric disorders in patients with chronic disabling occupational spinal disorders. *Spine* 2006; 31(10): 1156–62. [PubMed: 16648753]
21. Kinney RK, Gatchel RJ, Polatin PB, Fogarty WT, Mayer TG. Prevalence of psychopathology in acute and chronic low back pain patients. *J Occup Rehabil* 1993; 3(2): 95–103. [PubMed: 24243229]
22. Polatin PB, Kinney RK, Gatchel RJ, Lillo E, Mayer TG. Psychiatric illness and chronic low-back pain. The mind and the spine--which goes first? *Spine* 1993; 18(1): 66–71. [PubMed: 8434327]
23. Abbott PJ, Weller SB, Walker SR. Psychiatric disorders of opioid addicts entering treatment. *J Addict Dis* 1995; 13(3): 1–11.
24. Brooner RK, King VL, Kidorf M, Schmidt CW, Jr., Bigelow GE. Psychiatric and substance use comorbidity among treatment-seeking opioid abusers. *Arch Gen Psychiatry* 1997; 54(1): 71–80. [PubMed: 9006403]
25. Kidorf M, Disney ER, King VL, Neufeld K, Beilenson PL, Brooner RK. Prevalence of psychiatric and substance use disorders in opioid abusers in a community syringe exchange program. *Drug Alcohol Depend* 2004; 74(2): 115–22. [PubMed: 15099655]
26. Bair MJ, Wu J, Damush TM, Sutherland JM, Kroenke K. Association of depression and anxiety alone and in combination with chronic musculoskeletal pain in primary care patients. *Psychosom Med* 2008; 70(8): 890–7. [PubMed: 18799425]
27. Cacciola JS, Alterman AI, Rutherford MJ, McKay JR, Mulvaney FD. The relationship of psychiatric comorbidity to treatment outcomes in methadone maintained patients. *Drug Alcohol Depend* 2001; 61(3): 271–80. [PubMed: 11164691]
28. Barry DT, Beitel M, Cutter CJ, et al. Exploring relations among traumatic, posttraumatic, and physical pain experiences in methadone-maintained patients. *J Pain* 2011; 12(1): 22–8. [PubMed: 20646965]

29. Jamison RN, Kauffman J, Katz NP. Characteristics of methadone maintenance patients with chronic pain. *J Pain Symptom Manage* 2000; 19(1): 53–62. [PubMed: 10687327]
30. Haller DL, Acosta MC. Characteristics of pain patients with opioid-use disorder. *Psychosomatics* 2010; 51(3): 257–66. [PubMed: 20484724]
31. Ligthart L, Gerrits MM, Boomsma DI, Penninx BW. Anxiety and depression are associated with migraine and pain in general: an investigation of the interrelationships. *J Pain* 2013; 14.
32. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4th ed., text revision ed. Washington, DC: American Psychiatric Association; 2000.
33. Chou R, Qaseem A, Snow V, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med* 2007; 147(7): 478–91. [PubMed: 17909209]
34. AAPM, APS, ASAM. Consensus document from the American Academy of Pain Medicine, American Pain Society, and the American Society of Addiction Medicine. Glenview: The American Academy of Pain Medicine; 2001.
35. Cleeland CS. Pain assessment in cancer In: Osaba D, ed. Effect of cancer on quality of life. Boca Raton, FL: CRC Press; 1991: 293–305.
36. Cleeland CS, Ryan KM. Pain assessment: Global use of the Brief Pain Inventory. *Ann Acad Med Singapore* 1994; 23(2): 129–38. [PubMed: 8080219]
37. Cleeland C The Brief Pain Inventory User Guide. Houston, TX: The University of Texas MD Anderson Cancer Center, 2009.
38. First MB, Spitzer RL, Gibbon M, Williams JBW. Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Patient Edition (SCID-I/P). New York: Biometrics Research, New York State Psychiatric Institute; 2002.
39. Zanarini M, Frankenburg F, Sickel A, Yong L. The Diagnostic Interview for DSM-IV Personality Disorders (DIPD-IV). Belmont, Mass: McLean Hospital; 1996.
40. Fishbain DA, Goldberg M, Robert Meagher B, Steele R, Rosomoff H. Male and female chronic pain patients categorized by DSM-III psychiatric diagnostic criteria. *Pain* 1986; 26(2): 181–97. [PubMed: 3763232]
41. Strain EC, Brooner RK, Bigelow GE. Clustering of multiple substance use and psychiatric diagnoses in opiate addicts. *Drug Alcohol Depend* 1991; 27(2): 127–34. [PubMed: 2055161]
42. Mayer TG, Towns BL, Neblett R, Theodore BR, Gatchel RJ. Chronic widespread pain in patients with occupational spinal disorders: prevalence, psychiatric comorbidity, and association with outcomes. *Spine* 2008; 33(17): 1889–97. [PubMed: 18670343]
43. Rounsaville BJ, Kranzler HR, Ball S, Tennen H, Poling J, Triffleman E. Personality disorders in substance abusers: relation to substance use. *J Nerv Ment Dis* 1998; 186(2): 87–95. [PubMed: 9484308]
44. Monti DA, Herring CL, Schwartzman RJ, Marchese M. Personality assessment of patients with complex regional pain syndrome type I. *Clin J Pain* 1998; 14(4): 295–302. [PubMed: 9874007]
45. Conrad R, Schilling G, Bausch C, et al. Temperament and character personality profiles and personality disorders in chronic pain patients. *Pain* 2007; 133(1): 197–209. [PubMed: 17964076]
46. Mancini M, Perna G, Rossi A, Petralia A. Use of duloxetine in patients with an anxiety disorder, or with comorbid anxiety and major depressive disorder: a review of the literature. *Expert Opin Pharmacother* 2010; 11(7): 1167–81. [PubMed: 20402555]
47. Cuijpers P, Berking M, Andersson G, Quigley L, Kleiboer A, Dobson KS. A meta-analysis of cognitive-behavioural therapy for adult depression, alone and in comparison with other treatments. *Can J Psychiatry* 2013; 58(7): 376–85. [PubMed: 23870719]
48. Hofmann SG, Smits JA. Cognitive-behavioral therapy for adult anxiety disorders: a meta-analysis of randomized placebo-controlled trials. *J Clin Psychiatry* 2008; 69(4): 621–32. [PubMed: 18363421]
49. Tolin DF. Is cognitive-behavioral therapy more effective than other therapies?: a meta-analytic review. *Clin Psychol Rev* 2010; 30(6): 710–20. [PubMed: 20547435]
50. American Psychiatric Association. Diagnostic and Statistical Manual for Mental Disorders (Fifth Edition). Arlington, VA: American Psychiatric Association; 2013.

51. McHugh RK, Fitzmaurice GM, Carroll KM, et al. Assessing craving and its relationship to subsequent prescription opioid use among treatment-seeking prescription opioid dependent patients. *Drug Alcohol Depend* 2014; 145: 121–6. [PubMed: 25454409]
52. Wasan AD, Butler SF, Budman SH, et al. Does report of craving opioid medication predict aberrant drug behavior among chronic pain patients? *Clin J Pain* 2009; 25(3): 193–8. [PubMed: 19333168]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Clinical Points:

- This study found high rates of current and lifetime Axis I and Axis II psychiatric disorders in patients with co-occurring chronic pain and opioid dependence.
- This study also found that despite the long duration of current Axis I psychiatric disorders and the persistence of these disorders, very few participants had received mental health treatment or a psychiatric medication in the past month.
- The high rates and persistence of psychiatric disorders and low rates of current mental health treatment among participants with co-occurring chronic pain and opioid use disorder suggest important clinical targets in developing treatments for of these co-occurring chronic medical conditions.

Table 1. Demographic and clinical characteristics of patients with chronic pain and opioid dependence

	Buprenorphine (n=113)	Metadone (n=57)	Total (n=170)
Age, mean (SD) [†]	34.6 (9.3)	38.8 (10.6)	36.0 (10.0)
Gender, % (n)			
Male	70.8 (80)	71.9 (41)	71.2 (121)
Female	29.2 (33)	28.1 (16)	28.8 (49)
Race, % (n)			
White	86.7 (98)	87.7 (50)	87.1 (148)
African-American	4.4 (5)	5.3 (3)	4.1 (7)
Other	8.9 (10)	7.0 (4)	8.2 (14)
Ethnicity, % (n)			
Hispanic	9.7 (11)	10.6 (6)	10.0 (17)
Non-Hispanic	90.3 (102)	89.4 (51)	90.0 (153)
Education, % (n)			
Less than high school	12.4 (14)	14.0 (8)	12.9 (22)
High school graduates	86.7 (98)	86.0 (49)	86.5 (147)
Employment status, % (n) [*]			
Employed (full time, part time, student)	55.8 (63)	28.1 (16)	46.5 (79)
Unemployed (not employed, retired)	44.2 (50)	71.9 (41)	53.5 (91)
Marital status, % (n)			
Married	21.2 (24)	17.5 (10)	20.0 (34)
Single/divorce/separated	78.8 (89)	82.5 (47)	80.0 (136)
HIV status, % (n)			
Positive	1.8 (2)	0.0 (0)	1.2 (2)
Negative	96.5 (109)	96.5 (55)	96.5 (164)
Never tested/uncertain	1.8 (2)	3.5 (2)	2.4 (4)
Primary drug of abuse, % (n) [*]			
Heroin	38.1 (43)	59.6 (34)	45.3 (77)
Opioid medication	61.1 (69)	29.8 (17)	50.6 (86)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

	Buprenorphine (n=113)	Metadone (n=57)	Total (n=170)
Other	0.9 (1)	10.5 (6)	4.1 (7)
Outpatient mental health visit in past month, % (n)	3.5 (4)	3.6 (2)	3.5 (6)
Prescribed psychiatric medication in past month, % (n)*	10.6 (12)	23.2 (13)	14.8 (25)

[†] t-test, $p < 0.05$

* chi-square, $p < 0.05$

Table 2.

Psychiatric diagnoses of patients with chronic pain and opioid dependence

DSM-IV Psychiatric Diagnosis	(n=170)	
	Lifetime % (n)	Current % (n)
Axis I Diagnosis:		
Any Axis I disorder [†]	90.6 (154)	75.3 (128)
Anxiety disorders		
Any anxiety disorder	51.8 (88)	48.2 (82)
Posttraumatic stress disorder	23.5 (40)	20.6 (35)
Generalized anxiety disorder	15.9 (27)	15.9 (27)
Panic disorder (without agoraphobia)	15.9 (27)	15.9 (27)
Social Phobia	11.2 (19)	9.4 (16)
Specific phobias	5.9 (10)	5.3 (9)
Obsessive-compulsive disorder	7.1 (12)	6.5 (11)
Panic disorder (with agoraphobia)	3.5 (6)	2.9 (5)
Substance induced anxiety disorder	2.4 (4)	2.4 (4)
Agoraphobia (no history of panic disorder)	1.2 (2)	1.2 (2)
Anxiety Disorder, NOS	1.2 (2)	1.2 (2)
Mood disorders		
Any mood disorder	57.1 (97)	48.2 (82)
Major depressive disorder, recurrent	25.3 (43)	19.4 (33)
Major depressive disorder, single	23.5 (40)	20.6 (35)
Dysthymia	10.6 (18)	10.6 (18)
Bipolar disorder	4.7 (8)	4.7 (8)
Substance induced mood disorder	2.4 (4)	1.8 (3)
Depressive disorder, NOS	0.0 (0)	0.0 (0)
Cyclothymia	0.0 (0)	0.0 (0)
Adjustment disorder	0.0 (0)	0.0 (0)
Opioid dependence	100.0 (170)	100.0 (170)
Non-opioid substance use disorder		
Any non-opioid substance use disorder	77.6 (132)	33.5 (57)
Alcohol dependence	37.6 (64)	4.7 (8)
Alcohol abuse	19.4 (33)	3.5 (6)
Cocaine dependence	31.8 (54)	14.1 (24)
Cocaine abuse	11.4 (20)	3.5 (6)
Cannabis dependence	22.9 (39)	10.0 (17)
Cannabis abuse	18.8 (32)	7.6 (13)
Sedative/hypnotic dependence	6.5 (11)	1.8 (3)
Sedative/hypnotic abuse	8.8 (15)	4.7 (8)
Other drug dependence	7.1 (12)	1.2 (2)
Other drug abuse	5.7 (10)	2.4 (4)

DSM-IV Psychiatric Diagnosis	(n=170)	
	Lifetime % (n)	Current % (n)
Axis II Diagnosis:		
Any axis II disorder	N/A	51.8 (88)
Paranoid personality disorder	N/A	16.5 (28)
Schizoid personality disorder	N/A	8.2 (14)
Schizotypal personality disorder	N/A	0.0 (0)
Antisocial personality disorder	N/A	21.8 (37)
Borderline personality disorder	N/A	12.4 (21)
Histrionic personality disorder	N/A	0.0 (0)
Narcissistic personality disorder	N/A	0.0 (0)
Avoidant personality disorder	N/A	18.8 (32)
Dependent personality disorder	N/A	7.6 (13)
Obsessive-compulsive personality disorder	N/A	11.2 (19)
Depressive personality disorder	N/A	11.8 (20)
Axis IV:		
Financial problems	N/A	59.4 (101)
Problems with primary support group	N/A	51.2 (87)
Occupational problems	N/A	32.4 (55)
Lack of social support	N/A	8.2 (14)
Relationship problems	N/A	11.2 (19)
Unemployment	N/A	13.5 (23)
Problems with access to medical care	N/A	2.9 (5)
Homelessness	N/A	1.8 (3)
Financial stress	N/A	4.1 (7)
Problems with transportation	N/A	1.2 (2)
Work-related problems	N/A	1.8 (3)
Limited social network	N/A	1.8 (3)
Family stress	N/A	0.6 (1)
Housing problems	N/A	1.8 (3)
Axis V:		
GAF score, mean (SD)	N/A	59.4 (4.8)

[†]Does not include diagnosis of opioid dependence

Comparison of psychiatric diagnoses in patients with chronic pain and opioid dependence by self-reported primary drug of abuse⁷

Table 3.

DSM-IV Psychiatric Diagnosis	Heroin		PO		p	X ² (df=1)	p	Heroin		PO	
	Lifetime %	%	X ² (df=1)	p				Current %	%		
Axis I Diagnosis:											
Any Axis I disorder	97.4	84.9	7.620	0.005	80.5	68.6	3.015	0.059			
Anxiety disorders											
Any anxiety disorder	53.2	50.0	0.171	0.399	53.2	43.0	1.702	0.126			
Posttraumatic stress disorder	27.3	22.1	0.589	0.279	27.3	16.3	2.912	0.065			
Generalized anxiety disorder	16.9	12.8	0.542	0.303	16.9	12.8	0.542	0.303			
Panic disorder (without agoraphobia)	14.3	17.4	0.302	0.370	14.3	17.4	0.302	0.370			
Social Phobia	14.3	9.3	0.980	0.228	11.7	8.1	0.578	0.309			
Specific phobias	5.2	5.8	0.030*	1.000	5.2	4.7	0.026*	1.000			
Obsessive-compulsive disorder	9.1	5.8	0.640	0.308	7.8	5.8	0.253	0.423			
Panic disorder (with agoraphobia)	3.9	3.5	0.019*	1.000	3.9	2.3	0.337*	0.668			
Substance induced anxiety disorder	2.6	1.2	0.463*	0.603	2.6	1.2	0.463*	0.603			
Agoraphobia (no history of panic disorder)	1.3	1.2	0.006*	1.000	1.3	1.2	0.006*	1.000			
Anxiety Disorder, NOS	2.6	0.0	2.262*	0.222	2.6	0.0	2.262*	0.222			
Mood disorders											
Any mood disorder	59.7	53.5	0.646	0.259	48.1	46.5	0.039	0.484			
Major depressive disorder, recurrent	35.1	16.3	7.615	0.005	26.0	12.8	4.584	0.026			
Major depressive disorder, single	16.9	30.2	3.978	0.034	14.3	26.7	3.820	0.038			
Dysthymia	6.5	12.8	1.820	0.139	6.5	12.8	1.820	0.139			
Bipolar disorder	3.9	4.7	0.056*	1.000	3.9	4.7	0.056*	1.000			
Substance induced mood disorder	2.6	2.3	0.013*	1.000	2.6	1.2	0.463*	0.603			
Depressive disorder, NOS	0.0	0.0	N/A	N/A	0.0	0.0	N/A	N/A			
Cyclothymia	0.0	0.0	N/A	N/A	0.0	0.0	N/A	N/A			
Adjustment disorder	0.0	0.0	N/A	N/A	0.0	0.0	N/A	N/A			
Opioid dependence	100.0	100.0	N/A	N/A	100.0	100.0	N/A	N/A			

DSM-IV Psychiatric Diagnosis	Heroin		PO		P	Heroin		PO		X ² (df=1)	P
	Lifetime %	%	Lifetime %	%		Current %	%	Current %	%		
Non-opioid substance use disorder											
Any non-opioid substance use disorder	88.3	69.8	8.285	0.003	39.0	27.9	2.241	0.092			
Alcohol dependence	45.5	31.4	3.407	0.046	6.5	2.3	1.717*	0.257			
Alcohol abuse	16.9	22.1	0.699	0.262	2.6	4.7	0.483*	0.685			
Cocaine dependence	41.6	23.3	6.264	0.010	18.2	11.6	1.390	0.169			
Cocaine abuse	15.6	8.1	2.187	0.109	5.2	2.3	0.943*	0.423			
Cannabis dependence	27.3	17.4	2.282	0.093	10.4	8.1	0.246	0.410			
Cannabis abuse	23.4	16.3	1.297	0.173	9.1	7.0	0.247	0.416			
Sedative/hypnotic dependence	10.4	2.3	4.588*	0.048	2.6	0.0	2.262*	0.222			
Sedative/hypnotic abuse	9.1	9.3	0.002	0.590	5.2	4.7	0.026*	1.000			
Other drug dependence	7.8	5.8	0.253	0.423	1.3	1.2	0.006*	1.000			
Other drug abuse	9.1	3.5	2.214*	0.193	2.6	2.3	0.013*	1.000			
Axis II Diagnosis:											
Any axis II disorder	N/A	N/A	N/A	N/A	61.0	44.2	4.624	0.023			
Paranoid personality disorder	N/A	N/A	N/A	N/A	20.8	12.8	1.876	0.123			
Schizoid personality disorder	N/A	N/A	N/A	N/A	10.4	5.8	1.159	0.216			
Schizotypal personality disorder	N/A	N/A	N/A	N/A	0.0	0.0	N/A	N/A			
Antisocial personality disorder	N/A	N/A	N/A	N/A	26.0	18.6	1.282	0.173			
Borderline personality disorder	N/A	N/A	N/A	N/A	11.7	12.8	0.046	0.511			
Histrionic personality disorder	N/A	N/A	N/A	N/A	0.0	0.0	N/A	N/A			
Narcissistic personality disorder	N/A	N/A	N/A	N/A	0.0	0.0	N/A	N/A			
Avoidant personality disorder	N/A	N/A	N/A	N/A	18.2	18.6	0.005	0.554			
Dependent personality disorder	N/A	N/A	N/A	N/A	5.2	9.3	1.005	0.243			
Obsessive-compulsive personality disorder	N/A	N/A	N/A	N/A	14.3	8.1	1.562	0.159			
Depressive personality disorder	N/A	N/A	N/A	N/A	9.1	14.0	0.933	0.236			

[†] Primary drug of abuse was either heroin or PO (prescription opioids)

* Fisher's Exact Test used due to insufficient cell size

Bolded p-values were significant at p<.05.