



HHS Public Access

Author manuscript

Eur J Neurosci. Author manuscript; available in PMC 2020 August 01.

Published in final edited form as:

Eur J Neurosci. 2019 August ; 50(3): 2054–2064. doi:10.1111/ejn.13941.

Varying the rate of intravenous cocaine infusion influences the temporal dynamics of both drug and dopamine concentrations in the striatum

Ellie-Anna Minogianis¹, Waqqas M. Shams², Omar S. Mabrouk³, Jenny-Marie T. Wong⁴, Wayne G. Brake², Robert T. Kennedy^{3,4}, Patrick du Souich¹, Anne-Noël Samaha^{1,5}

¹Department of Pharmacology and Physiology, Faculty of Medicine, Université de Montréal;

²Department of Psychology, Center for Studies in Behavioral Neurobiology (CSBN), Concordia University;

³Department of Pharmacology, University of Michigan;

⁴Department of Chemistry, University of Michigan;

⁵Groupe de recherche sur le système nerveux central, Faculty of Medicine, Université de Montréal.

Abstract

The faster drugs of abuse reach the brain, the greater is the risk of addiction. Even small differences in the rate of drug delivery can influence outcome. Infusing cocaine intravenously over 5 versus 90–100 seconds promotes sensitization to the psychomotor and incentive motivational effects of the drug and preferentially recruits mesocorticolimbic regions. It remains unclear whether these effects are due to differences in how fast and/or how much drug reaches the brain. Here, we predicted that varying the rate of intravenous cocaine infusion between 5–90 seconds produces different rates of rise of brain drug concentrations, while producing similar peak concentrations. Freely-moving male Wistar rats received acute intravenous cocaine infusions (2.0 mg/kg/infusion) over 5, 45 and 90 seconds. We measured cocaine concentrations in the dorsal striatum using rapid-sampling microdialysis (1 sample/minute) and high-performance liquid chromatography-tandem mass spectrometry. We also measured extracellular concentrations of dopamine and other neurochemicals. Regardless of infusion rate, acute cocaine did not change concentrations of non-dopaminergic neurochemicals. Infusion rate did not significantly influence peak concentrations of cocaine or dopamine, but concentrations increased faster following 5-

Corresponding Author: Anne-Noël Samaha, Department of Pharmacology and Physiology, Université de Montréal, C.P. 6128, Succursale Centre-ville, Montreal, Quebec, H3C 3J7, Canada, Tel: 514 343 6111 x. 32788, Fax: 514 343 2291, Anna.samaha@umontreal.ca.

Authors contribution

ANS, EAM, WGB and WMS designed the study. EAM and WMS performed the *in vivo* microdialysis experiments. OSM and JTW analysed microdialysis samples with HPLC-MS/MS. EAM performed the psychomotor activity experiment. EAM, WMS, PS, OSM, JTW, RTK and ANS analyzed the data and interpreted the findings. EAM and ANS drafted the manuscript. PS, RTK, WMS, OSM, JTW and WGB critically revised the manuscript. All authors critically reviewed content and approved final version for publication.

Conflict of Interest


The authors declare no conflicts of interest.

Data accessibility

Raw data generated by these experiments will be stored on Figshare. Data and statistical analyses will be made available upon request.

second infusions. We also assessed psychomotor activity as a function of cocaine infusion rate. Infusion rate did not significantly influence total locomotion, but locomotion increased earlier following 5-second infusions. Thus, small differences in the rate of cocaine delivery influence both the rate of rise of drug and dopamine concentrations and psychomotor activity. A faster rate of rise of drug and dopamine concentrations might be an important issue in making rapidly delivered cocaine more addictive.

Graphical Abstract



Varying the rate of i.v. cocaine delivery between 5–90 seconds determines the drug's effects on brain and behaviour. We show that injecting cocaine between 5 and 90 seconds in rats alters the rates of rise of cocaine and dopamine in the dorsal striatum, without significantly changing peak concentrations. Faster injections also increase locomotor behaviour earlier than slower injections. Thus, beyond achieved dose, differences in the rates of rise of cocaine and dopamine can determine outcome.

Keywords

Cocaine addiction; *In vivo* microdialysis; Locomotor activity; Male Wistar rats; Pharmacokinetics

Introduction

The rate of drug delivery to the brain is important in determining the risk of developing addiction. For instance, smoking cocaine or injecting the drug intravenously is associated with a greater vulnerability to addiction than intranasal use (Gossop *et al.*, 1994; Hatsukami & Fischman, 1996). After smoking or intravenous (i.v.) infusion, plasma cocaine concentrations rise more rapidly and reach higher peaks than after intranasal administration (Javaid *et al.*, 1978; Jeffcoat *et al.*, 1989; Jones, 1990). Such pharmacokinetic differences are important in determining outcome. For instance, different pharmacokinetic profiles are one reason why the same drug can lead to addiction when taken by one route (nicotine inhaled from tobacco smoke) but can treat addiction when taken by another [nicotine administered orally from a gum or through the skin from a patch (Henningfield & Keenan, 1993)].

Varying the rate of drug delivery to the brain influences the psychological and behavioural effects of drugs. In humans, cocaine (Resnick *et al.*, 1977) and heroin (Comer *et al.*, 1999) evoke more immediate and stronger pleasurable effects when administered i.v. rather than intranasally. Similarly, increasing the rate of i.v. cocaine (Fischman & Schuster, 1984; Abreu *et al.*, 2001) or morphine (Marsch *et al.*, 2001) infusion produces greater self-reports of euphoria. Early studies in laboratory animals show that relatively large variations in the rate of i.v. drug administration [i.e., injecting a dose between 5–600 seconds (s)] influence cocaine self-administration behaviour (Balster & Schuster, 1973; Woolverton & Wang, 2004). More recent experiments have explored the effects of varying the rate of i.v. infusion over a smaller range of time (5–100 s). Doing so is important for three main reasons. First, experienced i.v. cocaine users report that they inject the drug rapidly and over a narrow range [3–10 s (Zernig *et al.*, 2003)]. Second, the time difference in the onset of subjective drug effects in i.v. versus intranasal cocaine users is also narrow [2–9 s (Zernig *et al.*, 2003) and 30–120 s (Jones, 1984; Weiss R.D., 1993), respectively]. Finally, injecting cocaine i.v. over 2 versus 60 s influences the magnitude of subjective cocaine effects in humans (Abreu *et al.*, 2001). Such small differences in the rate of drug delivery can have large effects on behaviour in laboratory rats. For instance, injecting cocaine or nicotine i.v. over 5 versus 90–100 s promotes sensitization to the psychomotor (Samaha *et al.*, 2002; Samaha *et al.*, 2004; Samaha & Robinson, 2005; Allain *et al.*, 2017) and incentive motivational effects of these drugs (Liu *et al.*, 2005; Minogianis *et al.*, 2013; Bouayad-Gervais *et al.*, 2014; Allain *et al.*, 2017). Rats that self-administer i.v. cocaine injections delivered over 5 versus 90 s also take more drug (Wakabayashi *et al.*, 2010; Minogianis *et al.*, 2013; Bouayad-Gervais *et al.*, 2014), and are more susceptible to relapse following extended abstinence (Wakabayashi *et al.*, 2010).

Variation in the rate of drug delivery influences behaviour presumably because it changes the neurobiological impact of drugs. Studies in rats show that increasing the rate of cocaine delivery to the brain enhances cellular activity in mesocorticolimbic areas [reviewed in

(Samaha & Robinson, 2005)]. A first study showed that compared to intraperitoneal cocaine, i.v. cocaine enhances glucose utilization in corticolimbic regions (Porrino, 1993). More recent work shows that injecting cocaine i.v. over 5 versus 90–100 s increases drug-induced immediate early gene expression (Samaha *et al.*, 2004), heat-producing metabolic activity (Brown & Kiyatkin, 2005), and regulation of the growth factor, brain-derived neurotrophic factor and its receptor TrkB (Bouayad-Gervais *et al.*, 2014). Chronic intake of i.v. infusions of cocaine delivered over 5 s but not over 90 s also increases the function of metabotropic group II receptors in the prelimbic cortex and nucleus accumbens (Allain *et al.*, 2017).

Thus, small differences in the rate of i.v. cocaine delivery can predict outcome but is this due to how fast cocaine enters the brain, peak achieved concentration, or both? This question remains unanswered because brain concentrations of cocaine have been *modelled* (Samaha *et al.*, 2002; Allain *et al.*, 2017), but not actually *measured* as a function of the rate of i.v. drug delivery. It is possible that varying the rate of i.v. drug infusion, even over a small range, produces differences in achieved concentration, and this could contribute to the effects seen on brain and behaviour. However, beyond achieved dose, temporal pharmacokinetic variables such as how fast drug levels rise can determine behavioural and neurobiological effects relevant to addiction (Allain *et al.*, 2015; Allain *et al.*, 2017). The brain is sensitive to the temporal pattern of cellular stimulation. For instance, in a context outside of drugs of abuse, different temporal patterns of electrical stimulation are differentially effective in producing long-term potentiation (Larson & Lynch, 1986) and dopamine-evoked synaptic plasticity (Wieland *et al.*, 2015).

Here, we hypothesized that varying the rate of i.v. cocaine delivery between 5 and 90 s produces differences in the rate of rise of striatal cocaine and dopamine concentrations, without producing large effects on peak concentrations. We make this prediction first because it is supported by pharmacokinetic modeling [(Samaha *et al.*, 2002; Allain *et al.*, 2017); where the pharmacokinetic model was based on (Pan *et al.*, 1991)]. Second, dopamine and cocaine concentrations are tightly correlated in the brain (Nicolaysen *et al.*, 1988; Shou *et al.*, 2006), and injecting cocaine i.v. over 5 versus 100 s produces differences in the rate of rise of extracellular dopamine concentrations in the striatum, without affecting peak concentrations [(Ferrario *et al.*, 2008) but see (Samaha *et al.*, 2004), where brief differences in the half-life of electrically-stimulated extracellular dopamine were observed]. To test our hypothesis, we injected freely-moving rats with i.v. infusions of cocaine delivered over 5, 45 and 90 s, and we used rapid-sampling (1 sample/minute) microdialysis coupled with high performance liquid chromatography-tandem mass spectrometry (HPLC-MS/MS) to simultaneously measure cocaine and dopamine concentrations in the dorsal striatum. We measured dopamine concentrations because cocaine-induced increases in dopamine overflow regulate the incentive motivational effects of the drug. The microdialysis samples were also used to determine cocaine effects on 20 other neurochemicals in the striatum. As an additional functional measure, we assessed the psychomotor response to acute cocaine infused over 5, 45 and 90 seconds, in a separate cohort of rats. Based on prior work, we predicted that varying the rate of cocaine delivery over this range would not influence the magnitude of the locomotor response (Samaha *et al.*, 2002), but that rate could influence the time course of cocaine-induced locomotion.

Materials and Methods

Animals and Housing

Twenty male Wistar rats (7 for microdialysis and 13 for cocaine-induced psychomotor activity; Charles River Laboratories, St-Constant, QC) weighing between 225–300 g upon arrival were housed individually in a climate-controlled colony room maintained on a reverse 12 h/12 h light/dark cycle (lights off at 8:30 – 9:00 am). Experiments were conducted during the dark phase of the rats' circadian cycle. Food and water were available *ad libitum*. The animal care committees of the Université de Montréal (CDEA 14–149 and 17–095) and of Concordia University approved all procedures involving animals. These procedures complied with guidelines of the Canadian Council on Animal Care.

Surgery

Following 1–4 weeks of acclimatization to the vivarium, a custom-made, indwelling catheter was implanted into the jugular vein of rats anaesthetized with isoflurane (CDMV; St-Hyacinthe, QC). The other end of the catheter was set to exit between the scapulae. Animals to be used for *in vivo* microdialysis were then placed in a stereotaxic apparatus and a cannula (21-gauge; HRS Scientific, Montreal, QC) was implanted into the dorsal striatum of one hemisphere (counterbalanced across animals; coordinates relative to Bregma; anteroposterior, +1.6 mm, mediolateral, \pm 2.5 mm, and dorsoventral, –3.0 mm). Cannulae were anchored to the skull with jeweller's screws, secured in place with dental cement and sealed with obturators (22-gauge; HRS Scientific). We targeted the dorsal striatum because it is rich in dopamine transporters and it mediates the expression of addiction-related behaviours (Everitt & Robbins, 2005). At the time of surgery, rats received a subcutaneous injection of 5 mg/kg Carprofen (Rimadyl; 50 mg/ml; CDMV, Saint-Hyacinthe, QC) and an intramuscular injection of 0.02 ml of a penicillin G procaine solution (Procillin; 300 000 IU/ml; CDMV, Saint-Hyacinthe, QC). Intravenous catheters were flushed on alternate days with either 0.1 mL physiological saline or a solution containing 0.2 mg/mL Heparin (Sigma-Aldrich, Oakville, ON) and 2 mg/mL of the antibiotic Baytril (CDMV, St-Hyacinthe, QC). Rats recovered in their home cages for 7 days prior to further manipulation.

Determination of microdialysis probe delay time *in vitro*

A first goal was to measure the delay time between placement of our microdialysis probes into a cocaine solution, and detection of the drug in a collected sample. This is important, as it provides a measure of the inevitable delay time that a sample needs to travel from the microdialysis probe, through the tubing set-up and finally into the sample collection vials. Two custom-made microdialysis probes (described below) were perfused at 2 μ L/minute with artificial cerebrospinal fluid (aCSF) containing 145 mM NaCl, 2.68 mM KCl, 1.40 mM $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 1.01 mM $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 1.55 mM Na_2HPO_4 , 0.45 mM $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ in HPLC-grade water (pH = 7.4; chemicals from Fisher Scientific, Saint-Laurent, Qc; water from Sigma Aldrich, Oakville, ON). Probes were first placed into a beaker containing a stirred solution of aCSF and ascorbic acid (0.25 mM) at room temperature. After 20 minutes, 10 samples were collected at 1-minute intervals. Probes were then transferred to a second beaker containing aCSF + ascorbic acid solution and 1 μ M cocaine, from which 26 samples were taken. Finally, the probes were returned to the aCSF + ascorbic acid solution

and 16 additional samples were collected. Samples were collected in 300- μ L microcentrifuge vials (VWR, Montreal, QC), and immediately placed in dry ice and stored at -80°C until analysis by HPLC-MS/MS.

***In vivo* microdialysis probes**

In vivo microdialysis was conducted in four custom-made hexagonal chambers ($42 \times 39 \times 33.5$ cm) each placed within a larger sound- and light-attenuating cabinet. Microdialysis probes were custom-made. They consisted of a 1 cm length of semi-permeable dialysis membrane (200 μm ID, 216 μm OD, with a molecular weight cut-off of 13 kDa; VWR, Montreal, QC) that extended 4 mm below the tip of the guide cannula. The membrane was inserted into a 20-mm length of 26G stainless steel tubing. The outer end of the membrane was occluded with super glue to create a closed system for dialysate flow. The stainless-steel shaft was inserted into one end of polyethylene (PE) 20 tubing (0.381 mm ID, 1.0922 mm OD; 65 cm long; Scientific Commodities Inc., Lake Havasu City, AZ), while the other end was connected to a dual-channel liquid swivel (Lomir, Notre-Dame-de-l'Île-Perrot, Qc). The swivel was in turn connected to a syringe pump (Harvard Apparatus, Saint-Laurent, Canada) with PE-20 tubing. Small-diameter fused silica tubing (41 μm ID, 110 μm OD; 2 cm long; HRS Scientific, Montreal, QC) extended into the probe 0.5 mm from the glued tip of the semi-permeable membrane. The other end of the fused silica was glued to PE-1 tubing (0.127 mm ID, 0.254 mm OD; 60 cm long) that extended out of the PE-20 tubing, serving as an outlet for the probe. The microdialysis probe was secured to the cannula with a stainless-steel collar. The relative recovery rates from the microdialysis probes were 10% for cocaine and 8–15% for dopamine.

***In vivo* Microdialysis Experiment**

Figure 1A illustrates the sequence of experimental events. Following recovery from surgery, microdialysis rats were placed in the test chambers for two, 3.5-hour (h) habituation sessions (one/day). Their catheters were tethered to the i.v. drug infusion line and their cannulae were tethered to the steel spring casing used to protect the microdialysis tubing during sampling. During each habituation session, each rat received 78 μL of saline i.v. over 5, 45 and 90 s, with one infusion given every 90 minutes in counter-balanced order. We injected a volume of 78 μL because for cocaine injections, we would inject 34 μL of saline to account for average catheter volume + 10 μL cocaine solution + 34 μL of saline to ensure that none of the cocaine remains in the catheter. Catheter patency was verified on the second habituation day by i.v. infusion of the short-acting barbiturate, sodium thiopental (20 mg/mL in sterile water, 0.1–0.2 mL/rat; CDMV, St-Hyacinthe, QC). All rats became ataxic within 5 s of the infusion, confirming catheter patency. In rats, sodium thiopental has a $T_{1/2}$ of ~ 4 –5 h (Shideman *et al.*, 1953), thus we do not expect it to influence microdialysis measurements performed on the next day. On the following day, rats were placed in the test chambers and dialysis probes were lowered into the striatum for a 5-h habituation period. During this period, aCSF was perfused through the microdialysis probe at a rate of 2.0 $\mu\text{L}/\text{minute}$. Rats were awake and freely moving during the experiment. Three hours and fifty minutes into the habituation period, animals were tethered to the cocaine infusion line. It contained 10 μL of cocaine solution (Medisca Pharmaceutique Inc, St-Laurent, QC; 2 mg/kg/infusion in 0.9% physiological saline) separated from additional saline by a small air bubble. The other end of

the line was attached to a 3 c.c. BD syringe placed on a syringe pump. A 2 mg/kg dose of cocaine is similar to doses used in prior studies that have measured cocaine or dopamine concentrations in the striatum using *in vivo* microdialysis (Hurd *et al.*, 1988; Hurd & Ungerstedt, 1989; Ferrario *et al.*, 2008). This dose also evokes robust immediate early gene expression in the dorsal striatum (Samaha *et al.*, 2004).

Over the last 10 minutes of the 5-h habituation period, 10 baseline dialysate samples were collected, at 1-minute intervals and at a flow rate of 2.0 $\mu\text{L}/\text{minute}$, yielding 2.0 $\mu\text{L}/\text{sample}$. Next, each animal received the first of three i.v. cocaine infusions (2.0 mg/kg/10 $\mu\text{L}/\text{infusion}$), delivered over 5, 45 and 90 s, injected 90 minutes apart, in counter-balanced order. Cocaine infusions were spaced 90 minutes apart because this is 3–4 times longer than cocaine's $T_{1/2}$ in rat brain (Nayak *et al.*, 1976; Hurd *et al.*, 1988). Thus, the 90-minute inter-infusion interval reduces the possibility of carry-over effects between infusions [see also (Hurd *et al.*, 1988)]. Following each cocaine infusion, 15 samples were collected, at 1-minute intervals. Five 1-minute baseline samples were also collected prior to the 2nd and 3rd cocaine infusions. Samples were collected in a 300- μL polypropylene microsampling vial placed on the end of the outlet line. Samples were immediately placed in dry ice and stored at -80°C until processing. At the end of sampling, each microdialysis probe was visually inspected for leaks or breakage. None were detected.

High performance liquid chromatography-tandem mass spectrometry (HPLC-MS/MS)

Small molecule neurochemical analysis using a triple quadrupole (QQQ) MS.

—All chemicals were from Sigma Aldrich (St. Louis, MO) unless noted otherwise. Water and acetonitrile for mobile phases were Burdick & Jackson HPLC grade (VWR, Radnor, PA). Artificial CSF consisted of 145 mM NaCl, 2.68 mM KCl, 1.40 mM CaCl_2 , 1.01 mM MgSO_4 , 1.55 mM Na_2HPO_4 , and 0.45 mM NaH_2PO_4 , adjusted pH to 7.4 with NaOH. Cocaine standards (Mallinckrodt Inc., St. Louis, MO) were spiked into a standard mixture for a six-point calibration curve. A modified LC-MS method (Song *et al.*, 2012; Wong *et al.*, 2016) was used to quantify concentrations of extracellular cocaine and dopamine, as well as acetylcholine, adenosine, aspartate, 3,4-dihydroxyphenylacetic acid, γ -aminobutyric acid, glutamate, glutamine, glucose, glycine, histamine, 5-hydroxyindoleacetic acid, homovanillic acid, 3-methoxytyramine, norepinephrine, normetanephrine, phenylalanine, serine, serotonin, taurine, and tyrosine. Samples were thawed and derivatized with 1.5 μL sodium carbonate, 100 mM; 1.5 μL BzCl, 2% (v/v) BzCl in acetonitrile; 1.5 μL isotopically labeled internal standard mixture diluted in 50% (v/v) acetonitrile containing 1% (v/v) sulfuric acid, and spiked with deuterated acetylcholine and choline (C/D/N isotopes, Pointe-Claire, Canada) to a final concentration of 20 nM. Derivatized samples were analyzed using Thermo Scientific Accela UHPLC system interfaced to a Thermo Scientific TSQ Quantum Ultra triple quadrupole mass spectrometer fitted with a HESI II ESI probe, operating in multiple reaction monitoring. Five- μL samples were injected onto a Phenomenex core-shell biphenyl Kinetex HPLC column (1.7 μm particles, 2.1 mm x 100 mm). Mobile phase A was 10 mM ammonium formate with 0.15% formic acid, and mobile phase B was acetonitrile. Mass transitions for these derivatized analytes were reported in Wong *et al.* (2016). Cocaine was detected in its native form with a MS/MS transition of 304 \rightarrow 182 m/z. The mobile phase was delivered through an elution gradient at 450 $\mu\text{L}/\text{minute}$ as follows: initial, 0% B; 0.01

minutes, 19% B; 1 minute, 26% B; 1.5 minutes, 75% B; 2.5 minutes, 100% B; 3 minutes, 100% B; 3.1 minutes, 5% B; and 3.5 minutes, 5% B. Thermo Xcalibur QuanBrowser (Thermo Fisher Scientific) software automatically processed and integrated peaks. Each peak was visually inspected to ensure proper integration. The limits of detection for cocaine and dopamine after derivatization were 0.291 and 0.086 nM, respectively. Limits of detection for the other analytes were 0.045–141 nM.

Histology

Following sampling, animals were anaesthetized with isoflurane and decapitated. Brains were then extracted, frozen and stored at -80°C . The neuroanatomical location of the probes was located on 20- μm thick coronal brain sections, according to the rat brain atlas of Paxinos and Watson (1986).

Psychomotor Activity Experiment

Figure 1B shows the sequence of experimental events. Locomotor activity was assessed in four chambers ($31.8 \times 25.4 \times 26.7$ cm; Med Associates, St-Albans, VT), each containing 4 infrared photocells aligned horizontally at the bottom of each cage. The chambers were run by a computer using Med Associates Med-PC version IV software (Med Associates, ST-Albans, VT). Following recovery from surgery, rats were placed in the test chambers for two, 2-h habituation sessions (one/day). Their catheters were tethered to the i.v. infusion line and a steel spring casing used to protect the line during testing. In the first habituation session, no i.v. infusion was given. During the second habituation session, rats received saline i.v. over 5, 45 or 90 s, thirty minutes into the session. Following the end of this second habituation session, catheter patency was verified by i.v. infusion of Propofol (10 mg/mL; 0.1 mL/rat; CDMV, St-Hyacinthe, QC), a short-acting anaesthetic [$T_{1/2} \sim 27$ minutes in Wistar rats; (Dutta *et al.*, 1997)]. Sodium thiopental was not used in this 2nd experiment because it was no longer available from the manufacturer. All rats became ataxic within 5 s of the Propofol infusion, confirming catheter patency. On the next day, animals were placed in the test chambers and tethered to an infusion line containing the cocaine solution (2.0 mg/kg in 10 μL). The other end of the line was attached to a 1 c.c. BD syringe and placed on a microsyringe pump (HAVARD PHD, 2000; HARVARD Apparatus, Saint-Laurent, Canada). Following a 30-minute habituation period, each animal received the first of three experimenter-administered i.v. cocaine infusions delivered over 5, 45 or 90 s, injected 90 minutes apart and in counter-balanced order. The test session lasted five hours, and locomotor activity was recorded as photocell beam breaks, computed over 10-s intervals. At the end of the study, catheter patency was verified once again with Propofol and animals were immediately euthanized by decapitation while still under anaesthesia. One rat did not become ataxic and was eliminated from the study.

Statistical analysis

Data was analyzed with Prism 7 software (GraphPad Software Inc., La Jolla, CA). Changes in cocaine concentration as a function of time during the *in vitro* assay were analyzed using one-way analysis of variance (ANOVA). Repeated measures two-way ANOVA was used to analyze the effects of cocaine infusion rate on average extracellular concentrations of cocaine and dopamine over time, and on locomotor activity over time ('Infusion Rate' and

'Time' as within-subjects variables). Repeated measures one-way ANOVA, followed by Tukey's multiple comparison tests or two-tailed paired *t*-tests, were used to analyse the effects of cocaine infusion rate on cocaine and dopamine C_{max} (the highest value for each analyte obtained from each rat after cocaine, averaged by infusion rate), T_{max} (time to reach C_{max} in each rat, averaged by infusion rate), and time to first significant increase (the first value > 2 standard deviations above baseline in each rat, averaged by infusion rate). One-way ANOVA was also used to assess the effects of cocaine infusion rate on both total locomotor activity (from minute '0' to minute '30') and locomotor activity within the first minute after injection. Linear regression was used for the correlations and Fisher's *r*-to-*z* transformation was then used to compare correlation coefficients (*r*). Effects were considered statistically significant when *P* < 0.05.

Results

Determination of microdialysis probe delay time *in vitro*

Figure 2 shows cocaine concentrations at 1-minute intervals when microdialysis probes were placed in an aCSF/ascorbic acid solution, then in an aCSF/ascorbic acid/1 μM cocaine solution, and back again. Of note, apparent cocaine concentrations do not start at or return to '0' when the probes are placed in the aCSF/ascorbic acid solution, before and after probe immersion in the cocaine solution. This background signal is likely produced by a contaminant with a similar mass transition to cocaine. We overcame this issue by taking the background contaminant signal into account when assigning the lower limit of quantification (LLOQ) of the LC-MS assay. This LLOQ was estimated as being 3X greater than the signal generated in aCSF/ascorbic acid alone, or 23 nM. The baseline values in Figure 2 are all below 23 nM and thus these values can be considered to be a background signal. It is only when the dialysis probe is placed in the cocaine-containing solution that cocaine concentrations rise above the LLOQ. Once the microdialysis probes were placed into the cocaine solution, cocaine concentrations increased significantly from baseline levels (One-way ANOVA on minutes -10 to 26; $F(35,36) = 26.08$; $P < 0.0001$). Cocaine concentrations began to increase 5–6 minutes after probe immersion into the cocaine solution, and reached near-maximum concentrations 1–2 minutes later. Thus, the delay time between the microdialysis probe and the collection vial is 5–6 minutes, and after this delay, near peak concentrations of cocaine were detected within 1–2 minutes. When the microdialysis probes were placed back into the solution that did not contain cocaine, drug concentrations significantly decreased (One-way ANOVA on minutes 0 to 42; $F(42,43) = 17.51$; $P < 0.0001$). Cocaine concentrations began to decrease 5–6 minutes after removal of the probes from the cocaine solution, and they returned to baseline levels 1–2 minutes later. Thus, there was a 5–6-minute delay time to detect the presence and then absence of cocaine in a solution. This delay is the time it takes for aCSF to be transported across the microdialysis probe membrane, through the probe outlet tubing and into the collection vial. Residual cocaine is also removed from the microdialysis probes within 2 minutes after this delay time. Based on these findings, the curves illustrating brain cocaine and dopamine concentrations *in vivo* (Figure 3) were corrected for a 5-minute delay time.

Varying the rate of i.v. cocaine delivery between 5 and 90 seconds significantly influences striatal cocaine and dopamine Tmax, but not Cmax

In all 7 rats, the unilateral microdialysis probe lay in the dorsal striatum between 2.2 and 1.2 mm anterior to Bregma (Figure 3a). Some probe tips extended slightly into the nucleus accumbens core. However, there would be little to no sampling from the probe tips as tips were occluded with super glue to create a closed system for dialysate flow. All animals received all three intravenous cocaine infusions administered over 5, 45 and 90 s, in counter-balanced order. Baseline levels of cocaine and dopamine during the 5 minutes prior to each infusion were comparable (Cocaine: $F(2, 18) = 1.42$, $P = 0.27$; Dopamine: $F(2, 18) = 0.15$, $P = 0.86$; data not shown). Thus, cocaine and dopamine concentrations returned to pre-cocaine baseline levels before each infusion, and there were no significant carry-over effects from one infusion to the next. At all rates, i.v. cocaine administration increased extracellular concentrations of both cocaine and dopamine above baseline (Figures 3b and 3c, respectively). The rate of i.v. cocaine infusion significantly influenced cocaine and dopamine concentrations over time (Infusion rate \times Time interaction effect; Figure 3b, $F_{36, 216} = 4.38$; Figure 3c, $F_{36, 216} = 3.56$; all P s < 0.0001). There was no significant effect of the rate of i.v. cocaine infusion on peak concentrations (Cmax values) of cocaine (Figure 4a; $F_{1.61, 9.67} = 1.71$, $P = 0.23$; 5 s: 267.82 ± 52.11 ; 45 s: 223.69 ± 41.19 ; 90 s: 185.09 ± 27.18 nM) or dopamine (Figure 4b $F_{1.41, 8.44} = 0.87$, $P = 0.42$; 5 s: 3.50 ± 0.49 , 45 s: 3.79 ± 0.82 ; 90 s: 3.06 ± 0.53 nM). However, as hypothesized, peak concentrations of cocaine (Figure 4c; $F_{1.5, 9} = 16.23$, $P = 0.0016$; 5–45 s < 90 s; 5 s: 4.33 ± 0.86 ; 45 s: 6.86 ± 1.28 ; 90 s: 11.71 ± 0.47 minutes) and dopamine (Figure 4d; $F_{1.53, 9.17} = 7.34$, $P = 0.02$; 5 < 90 s; 5 s: 4.33 ± 0.42 ; 45 s: 6.71 ± 0.89 ; 90 s: 9.29 ± 1.49 minutes) were reached earlier following faster i.v. infusions. To further examine this effect, we analyzed the effects of infusion rate on the time interval before cocaine and dopamine concentrations were greater than baseline concentrations by two standard deviations. This further confirmed that a 5-s infusion led to the fastest increases in extracellular concentrations of cocaine and dopamine (Figure 4e; $F_{2, 12} = 6.42$, $P = 0.01$; 5 versus 90 s: $t_6 = 2.74$, $P = 0.03$; 45 versus 90 s: $t_6 = 2.59$, $P = 0.04$; Figure 4f; $F_{2, 12} = 4.31$, $P = 0.04$; 5 versus 90 s: $t_6 = 2.54$, $P = 0.04$). In parallel, we observed the animals during testing and we noted that locomotor activity increased earlier following rapid (5 s) versus more sustained (45–90 s) i.v. cocaine infusions.

In summary, at all cocaine i.v. infusion rates, we measured a significant increase in drug and dopamine concentrations in the dorsal striatum. Increasing the rate of i.v. cocaine delivery between 90 and 5 s did not significantly influence peak concentrations of the drug or of dopamine. However, cocaine and dopamine concentrations reached peak levels earlier following more rapid infusions.

Cocaine and dopamine concentrations are tightly coupled across time

Data from representative rats show that, at all infusion rates, cocaine and dopamine concentrations were very closely linked across time (Figures 5a–c). Across all experimental rats and infusion rates, there was a significant positive correlation between extracellular cocaine and dopamine concentrations at each sampling time after drug infusion (Figure 5d; $r^2 = 0.77$, $P < 0.0001$). There was also a significant positive correlation between extracellular cocaine and dopamine concentrations at each infusion rate (5 s, $r^2 = 0.61$, $P = 0.0006$; 45 s,

$r^2 = 0.88$, $P < 0.0001$; 90 s, $r^2 = 0.91$, $P < 0.0001$, Data not shown). This correlation was stronger following a 90-s infusion compared to a 5-s infusion ($Z = -2.02$, $P = 0.04$; no other comparisons were statistically significant). The linear relationship between cocaine and dopamine was observed up to 187–220 nM cocaine, the highest concentrations measured following i.v. infusion of the drug over 5–90 s, respectively.

I.v. cocaine infusion does not significantly influence striatal levels of other neurochemicals

In addition to dopamine, we also measured extracellular concentrations of 20 other neurochemicals in the striatum after cocaine infusion. These were acetylcholine, adenosine, aspartate, γ -aminobutyric acid, glucose, glutamate, glutamine, glycine, histamine, norepinephrine, phenylalanine, serine, serotonin, taurine, tyrosine and the metabolites 3,4-dihydroxyphenylacetic acid, homovanillic acid, 3-methoxytyramine, 5-hydroxyindoleacetic acid and normetanephrine in the dorsal striatum. There were no significant effects of cocaine or of the rate of infusion on the concentrations of any of these (see supplementary figures S1–S3).

Locomotor activity increases earlier when cocaine is infused over 5 seconds

I.v. infusion rate did not influence the locomotor response to saline (Saline; $F_{2,9} = 0.21$, $P = 0.81$). Thus, saline-induced locomotion was pooled across rates and this served as the control condition. At all infusion rates, cocaine increased locomotor activity relative to saline (Figure 6; $F_{3,33} = 6.80$, $P = 0.001$; 5 s vs. Saline: $F_{1,11} = 11.43$, $P = 0.006$; 45 s vs. Saline: $F_{1,11} = 16.24$, $P = 0.002$; 90 s vs. Saline: $F_{1,11} = 27.74$, $P = 0.0003$), and all rates evoked a comparable increase in locomotion (One-way ANOVA on total locomotor counts over the 30 min after cocaine injection; $F_{1,10,12,13} = 2.17$, $P = 0.17$). However, infusion rate significantly influenced cocaine-induced locomotion across time (Infusion rate \times Time interaction effect; $F_{68, 748} = 1.47$, $P = 0.01$). To analyse this further, we compared locomotor activity in the first minute following cocaine injection. This showed that injecting cocaine i.v. over 5 versus 90 s evokes greater locomotor activity in the first minute post-injection ($F_{1,57,17,24} = 10.87$, $P = 0.002$; 5 vs. 90 s, $P = 0.004$).

Discussion

To our knowledge, the present experiment is the first to simultaneously measure brain concentrations of extracellular cocaine and dopamine in awake, freely-moving rats, and to also assess the influence of the rate of i.v. cocaine infusion. We found that injecting cocaine i.v. between 5 and 90 s robustly increases cocaine and dopamine concentrations in the dorsal striatum, without significantly altering peak concentrations of either analyte [see also (Ferrario *et al.*, 2008)]. However, drug and dopamine concentrations rose faster when cocaine was administered over 5 s, such that peak concentrations were reached earlier. Previous work shows that compared to slower i.v. cocaine infusions (25–100 s), rapid infusions (5–16 s) promote the development of sensitization to both the psychomotor activating and incentive motivational effects of cocaine (Samaha *et al.*, 2002; Samaha *et al.*, 2004; Liu *et al.*, 2005; Minogianis *et al.*, 2013; Bouayad-Gervais *et al.*, 2014; Allain *et al.*, 2017), lead to greater drug intake (Wakabayashi *et al.*, 2010; Minogianis *et al.*, 2013; Bouayad-Gervais *et al.*, 2014), and increase the risk of cocaine-primed relapse following

extended abstinence (Wakabayashi *et al.*, 2010). Rapid cocaine delivery to the brain also preferentially engages mesocorticolimbic cells (Porrino, 1993; Samaha *et al.*, 2004; Brown & Kiyatkin, 2005; Ferrario *et al.*, 2008). If rapid i.v. cocaine administration promotes these addiction-relevant effects by changing dopamine neurotransmission, our findings suggest that the critical factor is likely the time it takes to transition from low/baseline to high extracellular dopamine concentrations rather than any large differences in peak concentrations (Ferrario *et al.*, 2008).

Our results confirm predictions derived from pharmacokinetic modeling. Samaha *et al.* (2002) used a pharmacokinetic model validated by Pan *et al.* (1991) and estimated that injecting cocaine i.v. over 5, 50 or 100 s would not significantly influence peak brain concentrations of the drug, but would change the rate of rise of drug concentrations. Our findings confirm this to be the case. The temporal profile of brain cocaine concentrations we report also matches that in Samaha *et al.* (2002). However, Samaha *et al.* (2002) estimated higher peak cocaine values in the brain than measured here (~4 μM after 1 mg/kg i.v. cocaine compared to 0.2–0.26 μM after 2 mg/kg i.v. cocaine here). Samaha *et al.* (2002) used a pharmacokinetic model that estimates whole brain concentrations corrected for probe recovery rate, while here we report uncorrected concentrations from dialysate samples. Uncorrected concentration values in dialysate samples would be lower. In accordance with this, in the present study, an *in vitro* test showed that peak concentrations of cocaine detected by the probes were ten-fold lower than the concentration in the prepared solution (0.1–0.14 μM were detected in a 1- μM cocaine solution). This represents a 10% probe recovery at the 2 $\mu\text{L}/\text{minute}$ flow rate used here. As such, keeping in mind the 10% recovery rate from the microdialysis probe, the striatal cocaine and dopamine values we report can be used to select behaviourally-relevant cocaine and dopamine concentrations in *in vitro* experiments (Nicolaysen *et al.*, 1988). Indeed, the cocaine dose we used (2 mg/kg) also produced a robust increase in psychomotor activity (detailed below).

Variation in the rate of i.v. cocaine infusion significantly influenced the temporal dynamics of cocaine and dopamine levels in the dorsal striatum. By sampling at 1-minute intervals, we found that peak brain concentrations of cocaine were reached 4 minutes after a 5-s infusion, 7 minutes after a 45-s infusion and 12 minutes after a 90-s infusion. Similarly, Hurd *et al.* (1988), who sampled every 10 minutes, found that following an 85-s i.v. cocaine infusion, peak drug concentrations in the striatum are seen within the first 10-minute sample. The cocaine T_{max} values we observed here after a 5-versus 90-s i.v. infusion are also generally similar to those seen in rats following i.v. (1–2 minutes) and intranasal (15 minutes) cocaine administration, respectively (Chow *et al.*, 1999). This suggests that injecting cocaine i.v. over 5 versus 90 s in rats can to some extent model the temporal kinetics of i.v. versus intranasal cocaine. Our findings also agree with those of Ferrario *et al.* (2008) showing that infusing cocaine i.v. between 5 and 90 s does not produce large effects on peak dopamine concentrations in the striatum (or on area under the curve values for dopamine), but it produces significant differences in dopamine T_{max} . Ferrario *et al.* (2008) also reported that dopamine concentrations start to rise 1–2 minutes after a 5-s i.v. cocaine infusion, and 3–3.5 minutes after a 100-s infusion. Similarly, here we found that dopamine concentrations begin to increase 1 minute after a 5-s cocaine infusion, and 5 minutes after a 90-s infusion. We have previously used *in vivo* voltammetry techniques in urethane-anesthetized rats and

found that peak levels of dopamine reuptake inhibition occur within 1 minute after a 5-s i.v. cocaine infusion (Samaha *et al.*, 2004). This is much sooner than the time to reach peak brain concentrations of dopamine we observed here using awake, freely-moving animals and *in vivo* microdialysis (4 minutes after a 5-s infusion). However, our findings agree with those of others using *in vivo* voltammetry techniques in awake, freely-moving rats, and showing that peak dopamine inhibition does not occur until 6 minutes after a 12–15 s i.v. cocaine infusion (Kiyatkin *et al.*, 2000). The reasons for the discrepancy between these findings could involve the use of anesthetized versus awake animals. It is also possible that there is a non-linear relationship between the ability of an i.v. infusion of cocaine to occupy dopamine transporters and its ability to block the reuptake of dopamine, and thus increase dopamine overflow (Brodnik *et al.*, 2017).

Brain cocaine and dopamine concentrations were tightly coupled across time, at all infusion rates. This agrees with previous studies where cocaine was given via the intraperitoneal or i.v. routes (Nicolaysen *et al.*, 1988; Shou *et al.*, 2006). Thus, once in the brain, cocaine quickly occupies dopamine transporters, producing rapid dopamine reuptake blockade and overflow into the extracellular space. The cocaine-induced increases in dopamine concentrations we measured here are likely largely due to blockade of the dopamine transporter, but cocaine could also be altering dopamine release (Venton *et al.*, 2006; Mejias-Aponte *et al.*, 2015). As this matter is resolved, the very close temporal relationship between brain cocaine and dopamine concentrations we observed suggests that the pharmacokinetic profile of one compound can be used as a proxy for the other.

In contrast to what we observed with dopamine, acute i.v. cocaine injections did not change the concentrations of other neurochemicals in the striatum. We tested only one (albeit relatively high) dose of cocaine. In addition, our rats received a total of 3 cocaine infusions. This might not be sufficient to reliably alter extracellular concentrations of some of the compounds we measured. For instance, several exposures to cocaine are generally needed to significantly increase striatal glutamate concentrations (Zhang *et al.*, 2001). Interestingly, here and in prior work (Hurd & Ungerstedt, 1989), cocaine did not significantly change dopamine metabolite concentrations (DOPAC, HVA and 3-MT), even though the drug robustly increased extracellular dopamine concentrations. This suggests that the relationship between dopamine release and metabolism is not simple. Cocaine might not significantly change dopamine metabolite concentrations in the dorsal striatum because the drug potentially blocks dopamine reuptake, thus preventing dopamine metabolism in the cell terminal (Hurd & Ungerstedt, 1989). Although acute cocaine can significantly increase norepinephrine overflow in the ventral striatum (Reith *et al.* 1997), we found no effects on norepinephrine overflow in the dorsal striatum. To our knowledge, this is the first report on how systemic cocaine administration influences norepinephrine overflow in the dorsal striatum. The dorsal striatum contains a limited density of scattered noradrenergic axons and very little norepinephrine (Fornai *et al.*, 1996). We also saw no effect of i.v. cocaine on serotonin concentrations in the dorsal striatum. This was surprising since cocaine has similar affinities for brain serotonin and dopamine transporters (Miller *et al.*, 2001; Rothman *et al.*, 2001). Others have reported that acute cocaine increases extracellular concentrations of serotonin in the dorsal (Bradberry *et al.*, 1993) and ventral (Essman *et al.*, 1994) striatum. The discrepancy between our findings and these reports could be due to the use of anaesthetized

rats (Bradberry *et al.*, 1993), the route of cocaine administration (Bradberry *et al.*, 1993), and the striatal subregion sampled (Essman *et al.*, 1994).

The rate of cocaine infusion also influenced the temporal dynamics of psychomotor activity. While i.v. infusion rate did not significantly influence total levels of cocaine-induced locomotion, faster i.v. infusions evoked a greater locomotor response in the first minute after infusion. Thus, the rate of cocaine infusion influenced the temporal dynamics of both dopamine concentrations and drug-induced psychomotor activity. However, the relationship between these two measures is not straightforward. First, at all infusion rates, cocaine-induced locomotion both peaked earlier and returned to baseline levels sooner than drug or dopamine concentrations (compare Figures 6 and 3). We measured cocaine/dopamine concentrations and cocaine-induced locomotor activity in different animals. However, others have taken the two measures in the same animals and also found that in previously drug-naive rats, cocaine-induced locomotion peaks earlier than cocaine-induced dopamine concentrations [(Kalivas & Duffy, 1990) but see (Hemby *et al.*, 1994)]. Other studies with methylphenidate (Gerasimov *et al.*, 2000) or nicotine (Benwell & Balfour, 1992) also show a similar effect. Combined, the present study and these prior reports have measured drug-induced dopamine overflow in both the ventral and dorsal striatum. Thus, it is possible that drug-induced dopamine accumulation outside of these brain regions also significantly contributes to the psychomotor response (Kalivas & Duffy, 1990). It is also possible that cocaine increases dopamine concentrations in the synapse to induce psychomotor activity, without proportionate or immediate diffusion of dopamine outside of the synapse, to the microdialysis probe (Gonon, 1988; Kalivas & Duffy, 1990). This would explain why cocaine-induced dopamine concentrations detected by the probe peak later in time compared to cocaine-induced locomotion.

There are elements to consider in interpreting our findings. First, we do not know how the rate of i.v. cocaine infusion influences cocaine/dopamine pharmacokinetics in extra-striatal brain regions. We would predict that the cocaine pharmacokinetics we measured in the dorsal striatum likely reflect those in the rest of the brain. Indeed, cocaine is distributed uniformly in the brain following i.v. or intranasal administration (Reed & Spiehler, 1985). A Positron Emission Tomography study in humans also suggests that drug pharmacokinetics in individual brain regions are very similar to those in whole brain (Berridge *et al.*, 2010). Second, the neurobehavioural effects of cocaine that are relevant to addiction come about following chronic exposure to the drug. We do not know how our measurements would change with more extensive drug exposure. Of note, brain and blood concentrations of cocaine do not significantly change following repeated i.v. administration (Pan *et al.*, 1991). However, the ability of i.v. cocaine to inhibit dopamine reuptake can increase with repeated exposure (Brodnik *et al.*, 2017). This being said, studying brain cocaine and dopamine pharmacokinetics after a single cocaine exposure is important. A single exposure to psychostimulant drugs like cocaine or d-amphetamine can produce effects that are relevant to the addiction process, in both laboratory rats and humans. These effects include psychomotor sensitization (Robinson *et al.*, 1982; Lin-Chu *et al.*, 1985; Strakowski & Sax, 1998; Samaha *et al.*, 2002) and changes in spine density on medium spiny neurons of the nucleus accumbens (Kolb *et al.*, 2003). Finally, we only tested a single cocaine dose, but we

would predict that, across a range of cocaine doses, varying the rate of i.v. drug infusion changes the rate of rise of drug concentrations in the brain.

In summary, we measured striatal concentrations of cocaine and dopamine, as well as cocaine-induced locomotion across a range of i.v. infusion rates that significantly influences brain and behaviour [Reviewed in (Samaha & Robinson, 2005; Allain *et al.*, 2015)]. We found that increasing the rate of i.v. drug delivery increases the rate of rise of cocaine and dopamine concentrations in the dorsal striatum, without producing large effects in peak concentrations. This was mirrored by a more immediate increase in locomotor activity following rapid versus slower i.v. cocaine infusions. Thus, our results raise the possibility that differences in the rate of cocaine rise alone can determine outcome, perhaps by influencing the temporal dynamics of dopamine accumulation in the synapse, and the temporal pattern of dopamine receptor occupancy. In support of this, work in humans shows that smoked and intranasal cocaine can produce equivalent levels of dopamine transporter blockade, but smoked cocaine produces a stronger self-reported high (Volkow *et al.*, 2000). Thus, we conclude that differences in the rate of rise of drug and dopamine levels in the brain might be an important issue in thinking about why drugs, formulations, and routes of administration that achieve a rapid drug onset are the most addictive.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This research was supported by grants from the Canadian Institutes of Health Research (Grant No. 157572) and the Canada Foundation for Innovation (Grant No. 24326) to ANS, as well as a discovery grant from the Natural Sciences and Engineering Research Council (NSERC) of Canada (Grant No. RGPIN-2016-06653) awarded to WGB. ANS holds a salary award from the Fonds de la Recherche du Québec - Santé (Grant No. 28988). EAM holds a graduate fellowship from the Fonds de la Recherche du Québec - Santé (Grant No. 29651). WMS holds a graduate fellowship from the Fonds de Recherche du Québec - Nature et Technologies. OSM was supported by the Michael J Fox Foundation for Parkinson's Disease Research (BioFind and Dyskinesia Challenge). RTK was supported by NIH R37 EB003320.

Abbreviations

aCSF	artificial cerebrospinal fluid
ANOVA	analysis of variance
C_{max}	maximal concentration
DS	dorsal striatum
h	hours
i.v.	intravenous
μM	micromoles per liter
nM	nanomoles per liter

NacC	nucleus accumbens core
NacSh	nucleus accumbens shell
s	seconds
SEM	standard error of the mean
Tmax	time to reach maximal concentration

References

- Abreu ME, Bigelow GE, Fleisher L & Walsh SL (2001) Effect of intravenous injection speed on responses to cocaine and hydromorphone in humans. *Psychopharmacology*, 154, 76–84. [PubMed: 11292009]
- Allain F, Minogianis E-A, Roberts DC & Samaha A-N (2015) How fast and how often: The pharmacokinetics of drug use are decisive in addiction. *Neuroscience & Biobehavioral Reviews*, 56, 166–179. [PubMed: 26116543]
- Allain F, Roberts DC, Levesque D & Samaha AN (2017) Intermittent intake of rapid cocaine injections promotes robust psychomotor sensitization, increased incentive motivation for the drug and mGlu2/3 receptor dysregulation. *Neuropharmacology*, 117, 227–237. [PubMed: 28137451]
- Balster RL & Schuster CR (1973) FIXED-INTERVAL SCHEDULE OF COCAINE REINFORCEMENT: EFFECT OF DOSE AND INFUSION DURATION1. *Journal of the experimental analysis of behavior*, 20, 119–129. [PubMed: 4197505]
- Benwell ME & Balfour DJ (1992) The effects of acute and repeated nicotine treatment on nucleus accumbens dopamine and locomotor activity. *British journal of pharmacology*, 105, 849–856. [PubMed: 1504716]
- Berridge MS, Apana SM, Nagano KK, Berridge CE, Leisure GP & Boswell MV (2010) Smoking produces rapid rise of [¹¹C]nicotine in human brain. *Psychopharmacology*, 209, 383–394. [PubMed: 20232056]
- Bouayad-Gervais K, Minogianis E-A, Lévesque D & Samaha A-N (2014) The self-administration of rapidly delivered cocaine promotes increased motivation to take the drug: contributions of prior levels of operant responding and cocaine intake. *Psychopharmacology*, 231, 4241–4252. [PubMed: 24752656]
- Bradberry CW, Nobiletti JB, Elsworth JD, Murphy B, Jatlow P & Roth RH (1993) Cocaine and Cocaethylene: Microdialysis Comparison of Brain Drug Levels and Effects on Dopamine and Serotonin. *Journal of Neurochemistry*, 60, 1429–1435. [PubMed: 8455033]
- Brodnik ZD, Ferris MJ, Jones SR & España RA (2017) Reinforcing Doses of Intravenous Cocaine Produce Only Modest Dopamine Uptake Inhibition. *ACS Chemical Neuroscience*, 8, 281–289. [PubMed: 27936579]
- Brown PL & Kiyatkin EA (2005) Brain temperature change and movement activation induced by intravenous cocaine delivered at various injection speeds in rats. *Psychopharmacology*, 181, 299–308. [PubMed: 15778873]
- Chow HHS, Chen Z & Matsuura GT (1999) Direct transport of cocaine from the nasal cavity to the brain following intranasal cocaine administration in rats. *Journal of Pharmaceutical Sciences*, 88, 754–758. [PubMed: 10430537]
- Comer SD, Collins ED, MacArthur RB & Fischman MW (1999) Comparison of intravenous and intranasal heroin self-administration by morphine-maintained humans. *Psychopharmacology*, 143, 327–338. [PubMed: 10367549]
- Dutta S, Matsumoto Y, Gothgen NU & Ebling WF (1997) Concentration–EEG Effect Relationship of Propofol in Rats. *Journal of Pharmaceutical Sciences*, 86, 37–43. [PubMed: 9002457]
- Essman WD, Singh A & Lucki I (1994) Serotonergic properties of cocaine: Effects on a 5-HT₂ receptor-mediated behavior and on extracellular concentrations of serotonin and dopamine. *Pharmacology Biochemistry and Behavior*, 49, 107–113.

- Everitt BJ & Robbins TW (2005) Neural systems of reinforcement for drug addiction: from actions to habits to compulsion. *Nature neuroscience*, 8, 1481–1489. [PubMed: 16251991]
- Ferrario C, Shou M, Samaha A, Watson C, Kennedy R & Robinson T (2008) The rate of intravenous cocaine administration alters c-fos mRNA expression and the temporal dynamics of dopamine, but not glutamate, overflow in the striatum. *Brain research*, 1209, 151–156. [PubMed: 18396266]
- Fischman M & Schuster C (1984) Injection duration of cocaine in humans. *Federation proceedings*, 43, 570.
- Fornai F, Torracca MT, Bassi L, D'Errigo DA, Scalori V & Corsini GU (1996) Norepinephrine loss selectively enhances chronic nigrostriatal dopamine depletion in mice and rats. *Brain Research*, 735, 349–353. [PubMed: 8911678]
- Gerasimov MR, Franceschi M, Volkow ND, Gifford A, Gatley SJ, Marsteller D, Molina PE & Dewey SL (2000) Comparison between intraperitoneal and oral methylphenidate administration: a microdialysis and locomotor activity study. *Journal of Pharmacology and Experimental Therapeutics*, 295, 51–57. [PubMed: 10991960]
- Gonon FG (1988) Nonlinear relationship between impulse flow and dopamine released by rat midbrain dopaminergic neurons as studied by in vivo electrochemistry. *Neuroscience*, 24, 19–28. [PubMed: 3368048]
- Gossop M, Griffiths P, Powis B & Strang J (1994) Cocaine: patterns of use, route of administration, and severity of dependence. *The British Journal of Psychiatry*, 164, 660–664. [PubMed: 7921717]
- Hatsukami DK & Fischman MW (1996) Crack cocaine and cocaine hydrochloride: Are the differences myth or reality? *JAMA*, 276, 1580–1588. [PubMed: 8918856]
- Hemby S, Jones G, Hubert G, Neill D & Justice J (1994) Assessment of the relative contribution of peripheral and central components in cocaine place conditioning. *Pharmacology Biochemistry and Behavior*, 47, 973–979.
- Henningfield JE & Keenan RM (1993) Nicotine delivery kinetics and abuse liability. *Journal of consulting and clinical psychology*, 61, 743–750. [PubMed: 8245272]
- Hurd YL, Kehr J & Ungerstedt U (1988) In Vivo Microdialysis as a Technique to Monitor Drug Transport: Correlation of Extracellular Cocaine Levels and Dopamine Overflow in the Rat Brain. *Journal of Neurochemistry*, 51, 1314–1316. [PubMed: 3418351]
- Hurd YL & Ungerstedt U (1989) Cocaine: an in vivo microdialysis evaluation of its acute action on dopamine transmission in rat striatum. *Synapse*, 3, 48–54. [PubMed: 2537539]
- Javaid J, Fischman M, Schuster CR, Dekirmenjian H & Davis J (1978) Cocaine plasma concentration: relation to physiological and subjective effects in humans. *Science*, 202, 227–228. [PubMed: 694530]
- Jeffcoat AR, Perez-Reyes M, Hill JM, Sadler BM & Cook CE (1989) Cocaine disposition in humans after intravenous injection, nasal insufflation (snorting), or smoking. *Drug Metabolism and Disposition*, 17, 153–159. [PubMed: 2565204]
- Jones RT (1984) The pharmacology of cocaine. *Cocaine: Pharmacology, Effects and Treatment of Abuse*. National Institute on Drug Abuse Research Monograph, 50, 34–53.
- Jones RT (1990) The pharmacology of cocaine smoking in humans. *Research findings on smoking of abused substances*. National Institute on Drug Abuse Research Monograph, 99, 30–41.
- Kalivas PW & Duffy P (1990) Effect of acute and daily cocaine treatment on extracellular dopamine in the nucleus accumbens. *Synapse*, 5, 48–58. [PubMed: 2300906]
- Kiyatkin E, Kiyatkin D & Rebec G (2000) Phasic inhibition of dopamine uptake in nucleus accumbens induced by intravenous cocaine in freely behaving rats. *Neuroscience*, 98, 729–741. [PubMed: 10891616]
- Kolb B, Gorny G, Li Y, Samaha A-N & Robinson TE (2003) Amphetamine or cocaine limits the ability of later experience to promote structural plasticity in the neocortex and nucleus accumbens. *Proceedings of the National Academy of Sciences*, 100, 10523–10528.
- Larson J & Lynch G (1986) Induction of synaptic potentiation in hippocampus by patterned stimulation involves two events. *Science*, 232, 985–988. [PubMed: 3704635]
- Lin-Chu G, Robinson TE & Becker JB (1985) Sensitization of rotational behavior produced by a single exposure to cocaine. *Pharmacology Biochemistry and Behavior*, 22, 901–903.

- Liu Y, Roberts D & Morgan D (2005) Sensitization of the reinforcing effects of self-administered cocaine in rats: effects of dose and intravenous injection speed. *European Journal of Neuroscience*, 22, 195–200. [PubMed: 16029209]
- Marsch LA, Bickel WK, Badger GJ, Rathmell JP, Swedberg MD, Jonzon B & Norsten-Höög C (2001) Effects of infusion rate of intravenously administered morphine on physiological, psychomotor, and self-reported measures in humans. *Journal of Pharmacology and Experimental Therapeutics*, 299, 1056–1065. [PubMed: 11714895]
- Mejias-Aponte CA, Ye C, Bonci A, Kiyatkin EA & Morales M (2015) A subpopulation of neurochemically-identified ventral tegmental area dopamine neurons is excited by intravenous cocaine. *Journal of Neuroscience*, 35, 1965–1978. [PubMed: 25653355]
- Miller GM, Yatin SM, De La Garza Ii, Goulet R, M. & Madras BK (2001) Cloning of dopamine, norepinephrine and serotonin transporters from monkey brain: relevance to cocaine sensitivity. *Molecular Brain Research*, 87, 124–143. [PubMed: 11223167]
- Minogianis E-A, Lévesque D & Samaha A-N (2013) The Speed of Cocaine Delivery Determines the Subsequent Motivation to Self-Administer the Drug. *Neuropsychopharmacology*, 38, 2644–2656. [PubMed: 23921953]
- Nayak P, Misra A & Mule S (1976) Physiological disposition and biotransformation of (3H) cocaine in acutely and chronically treated rats. *Journal of Pharmacology and Experimental Therapeutics*, 196, 556–569. [PubMed: 1263111]
- Nicolaysen LC, Pan H-T & Justice JB Jr (1988) Extracellular cocaine and dopamine concentrations are linearly related in rat striatum. *Brain research*, 456, 317–323. [PubMed: 3208083]
- Pan HT, Menacherry S & Justice J (1991) Differences in the Pharmacokinetics of Cocaine in Naive and Cocaine-Experienced Rats. *Journal of neurochemistry*, 56, 1299–1306. [PubMed: 2002342]
- Paxinos G & Watson C (1986) *The rat brain in stereotaxic coordinates*. Academic Press, San Diego, 119.
- Porrino LJ (1993) Functional consequences of acute cocaine treatment depend on route of administration. *Psychopharmacology (Berl)*, 112, 343–351. [PubMed: 7871040]
- Reed D & Spiehler VR (1985) Brain concentrations of cocaine and benzoylecgonine in fatal cases. *Journal of Forensic Science*, 30, 1003–1011.
- Resnick R, Kestenbaum R & Schwartz L (1977) Acute systemic effects of cocaine in man: a controlled study by intranasal and intravenous routes. *Science*, 195, 696–698. [PubMed: 841307]
- Robinson TE, Becker JB & Presty SK (1982) Long-term facilitation of amphetamine-induced rotational behavior and striatal dopamine release produced by a single exposure to amphetamine: Sex differences. *Brain Research*, 253, 231–241. [PubMed: 6891283]
- Rothman RB, Baumann MH, Dersch CM, Romero DV, Rice KC, Carroll FI & Partilla JS (2001) Amphetamine-type central nervous system stimulants release norepinephrine more potently than they release dopamine and serotonin. *Synapse*, 39, 32–41. [PubMed: 11071707]
- Samaha A-N, Li Y & Robinson TE (2002) The rate of intravenous cocaine administration determines susceptibility to sensitization. *The Journal of neuroscience*, 22, 3244–3250. [PubMed: 11943825]
- Samaha A-N, Mallet N, Ferguson SM, Gonon F & Robinson TE (2004) The rate of cocaine administration alters gene regulation and behavioral plasticity: implications for addiction. *The Journal of neuroscience*, 24, 6362–6370. [PubMed: 15254092]
- Samaha A-N & Robinson TE (2005) Why does the rapid delivery of drugs to the brain promote addiction? *Trends in Pharmacological Sciences*, 26, 82–87. [PubMed: 15681025]
- Shideman F, Winters WD, Peterson RC, Wilner WK & Gould TC (1953) The distribution and in vivo rate of metabolism of thiopental. *Journal of Pharmacology and Experimental Therapeutics*, 107, 368–378. [PubMed: 13035675]
- Shou M, Ferrario CR, Schultz KN, Robinson TE & Kennedy RT (2006) Monitoring Dopamine in Vivo by Microdialysis Sampling and On-Line CE-Laser-Induced Fluorescence. *Analytical Chemistry*, 78, 6717–6725. [PubMed: 17007489]
- Song P, Mabrouk OS, Hershey ND & Kennedy RT (2012) In Vivo Neurochemical Monitoring using Benzoyl Chloride Derivatization and Liquid Chromatography – Mass Spectrometry. *Analytical Chemistry*, 84, 412–419. [PubMed: 22118158]

- Strakowski SM & Sax KW (1998) Progressive behavioral response to repeated d-amphetamine challenge: further evidence for sensitization in humans. *Biological psychiatry*, 44, 1171–1177. [PubMed: 9836021]
- Venton BJ, Seipel AT, Phillips PE, Wetsel WC, Gitler D, Greengard P, Augustine GJ & Wightman RM (2006) Cocaine increases dopamine release by mobilization of a synapsin-dependent reserve pool. *Journal of Neuroscience*, 26, 3206–3209. [PubMed: 16554471]
- Volkow ND, Wang GJ, Fischman MW, Foltin R, Fowler JS, Franceschi D, Franceschi M, Logan J, Gatley SJ, Wong C, Ding Y-S, Hitzemann R & Pappas N (2000) Effects of route of administration on cocaine induced dopamine transporter blockade in the human brain. *Life Sciences*, 67, 1507–1515. [PubMed: 10983846]
- Wakabayashi KT, Weiss MJ, Pickup KN & Robinson TE (2010) Rats markedly escalate their intake and show a persistent susceptibility to reinstatement only when cocaine is injected rapidly. *The Journal of Neuroscience*, 30, 11346–11355. [PubMed: 20739555]
- Weiss RD, Bartel R. MSM (1993) Cocaine. *American Psychiatric Assoc Pub.*
- Wieland S, Schindler S, Huber C, Kohr G, Oswald MJ & Kelsch W (2015) Phasic Dopamine Modifies Sensory-Driven Output of Striatal Neurons through Synaptic Plasticity. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 35, 9946–9956.
- Wong J-MT, Malec PA, Mabrouk OS, Ro J, Dus M & Kennedy RT (2016) Benzoyl chloride derivatization with liquid chromatography–mass spectrometry for targeted metabolomics of neurochemicals in biological samples. *Journal of Chromatography A*, 1446, 78–90. [PubMed: 27083258]
- Woolverton WL & Wang Z (2004) Relationship between injection duration, transporter occupancy and reinforcing strength of cocaine. *European Journal of Pharmacology*, 486, 251–257. [PubMed: 14985046]
- Zernig G, Giacomuzzi S, Riemer Y, Wakonigg G, Sturm K & Saria A (2003) Intravenous drug injection habits: drug users' self-reports versus researchers' perception. *Pharmacology*, 68, 49–56. [PubMed: 12660479]
- Zhang Y, Loonam TM, NOAILLES PA & Angulo JA (2001) Comparison of Cocaine-and Methamphetamine-Evoked Dopamine and Glutamate Overflow in Somatodendritic and Terminal Field Regions of the Rat Brain during Acute, Chronic, and Early Withdrawal Conditions. *Annals of the New York Academy of Sciences*, 937, 93–120. [PubMed: 11458542]




Figure 1. The sequence of experimental events.

Panel a illustrates the timeline of experimental events for the *in vivo* microdialysis study. Rats were implanted with a unilateral cannula into the dorsal striatum and a catheter into the jugular vein. Following recovery, the rats were habituated to the *in vivo* microdialysis apparatus and to the i.v. infusion procedure on 2 daily sessions. On the following test day, microdialysis probes were inserted into the dorsal striatum and i.v. catheters were tethered to the cocaine infusion set up. Each rat received an i.v. infusion of 2 mg/kg cocaine, delivered over 5, 45 or 90 s, in counterbalanced order, with infusions administered 90 minutes apart. Dialysate samples were collected every minute for 5–10 minutes before each infusion and for 15 minutes thereafter. Panel b shows the sequence of experimental events for the psychomotor activity study. Rats were implanted with an intrajugular catheter and allowed to recover for 7 days. Rats were then habituated to the psychomotor activity cages and i.v. infusion lines on 2 daily sessions. On the following test day, rats were tethered to the cocaine infusion lines and locomotor activity was measured. Each rat received i.v. cocaine (2.0 mg/kg/infusion) delivered over 5, 45 and 90 s, in counterbalanced order, with infusions administered 90 minutes apart.




Figure 2. Cocaine concentrations detected when microdialysis probes were placed into a solution containing 1 μM cocaine.

Two probes were each transferred from a solution containing aCSF/ascorbic acid to a solution containing aCSF/ascorbic acid/cocaine (1 μM), and back again. All solutions were stirred and tested at room temperature. The data shown are averages (\pm SEM) from two independent tests carried out with two different probes. Samples were collected at 1-minute intervals, at a flow rate of 2 $\mu\text{L}/\text{minute}$, and analyzed by HPLC-MS/MS. $n = 2$. μM , micromoles/liter. $n\text{M}$, nanomoles/liter.




Figure 3. Varying the rate of i.v. infusion between 5–90 s influences the temporal kinetics of extracellular cocaine and dopamine concentrations in the dorsal striatum.

The location of microdialysis probes in the striatum is shown in panel (a). The distance in millimeters (mm) from Bregma is given for each coronal rat brain section. The white boxes at the tips of each probe indicate the segment where glue was applied, and where no exchange is possible. Panels (b) and (c) show striatal cocaine and dopamine concentrations over time, respectively, as a function of i.v. drug infusion rate. All values are mean \pm SEM. $n = 7$ rats/infusion rate. *DS*, Dorsal Striatum, *NacC*, Nucleus Accumbens Core, *NacSh*, Nucleus Accumbens Shell, *nM*, nanomoles/liter. *s*, seconds.




Figure 4. Varying the rate of i.v. infusion between 5–90 s influences the rate of rise of cocaine and dopamine concentrations (Tmax) in the dorsal striatum without producing large effects on maximum concentrations (Cmax).

Panels (a) and (b) show average Cmax values for cocaine and dopamine, respectively, as a function of the rate of i.v. cocaine infusion. The time to reach peak concentrations (Tmax) of cocaine (c) and dopamine (d) in the dorsal striatum is shown as a function of the rate of i.v. cocaine delivery. Panels (e) and (f) show the time interval before cocaine and dopamine concentrations were significantly greater than baseline levels (> 2 standard deviations above baseline), as a function of the rate of i.v. cocaine delivery. All values are mean \pm SEM. $n = 7$ rats/infusion rate. * $P < 0.05$ compared to 90 s. *nM*, nanomoles/liter. *s*, seconds.




Figure 5. Extracellular dopamine and cocaine concentrations in the dorsal striatum are linearly correlated.

Panels (a-c) show extracellular concentrations of dopamine and cocaine over time from representative rats, for each i.v. infusion rate. Panel (d) shows a significant positive correlation between dopamine and cocaine concentrations in the dorsal striatum, at each sampling time, across all rats and infusion rates. For this analysis, we used linear regression to model the relationship between dopamine concentrations and cocaine concentrations at each of the 15 post-cocaine samples. $n = 7$ rats/infusion rate. *nM*, nanomoles/liter. *s*, seconds.




Figure 6. The influence of the rate of i.v. cocaine infusion on locomotor activity. Locomotor activity (beam breaks per minute) as a function of i.v. drug infusion rate. Saline-induced locomotor activity is also shown for comparison. The shading highlights the first minute after cocaine, where locomotor activity was greater after a 5-s injection than after a 90-s injection. All values are mean \pm SEM. $n = 12$ rats/infusion rate. s, seconds.