

A Journey to Understand Glucose Homeostasis: Starting from Rat Glucose Transporter Type 2 Promoter Cloning to Hyperglycemia

Yong Ho Ahn

Department of Biochemistry and Molecular Biology, Integrated Genomic Research Center for Metabolic Regulation, Yonsei University College of Medicine, Seoul, Korea

My professional journey to understand the glucose homeostasis began in the 1990s, starting from cloning of the promoter region of glucose transporter type 2 (*GLUT2*) gene that led us to establish research foundation of my group. When I was a graduate student, I simply thought that hyperglycemia, a typical clinical manifestation of type 2 diabetes mellitus (T2DM), could be caused by a defect in the glucose transport system in the body. Thus, if a molecular mechanism controlling glucose transport system could be understood, treatment of T2DM could be possible. In the early 70s, hyperglycemia was thought to develop primarily due to a defect in the muscle and adipose tissue; thus, muscle/adipose tissue type glucose transporter (*GLUT4*) became a major research interest in the diabetology. However, glucose utilization occurs not only in muscle/adipose tissue but also in liver and brain. Thus, I was interested in the hepatic glucose transport system, where glucose storage and release are the most actively occurring.


Keywords: Adipogenesis; Diabetes mellitus, type 2; Glucokinase; Gluconeogenesis; Glucose transporter type 2; Glycolysis; Transcription factors

INTRODUCTION

In β -cells of pancreas and liver, glucose transport machinery consists of glucose transporter type 2 (*GLUT2*) and glucokinase (*Gck*), which is also called a glucose sensor [1]. But in the 1980s, the mechanism of the regulation of gene expression of hepatic glucose transport system in the insulin resistant, diabetic states was largely unknown. Thus, we started to clone upstream regions (promoters) of *GLUT2*, *Gck*, and glucose-6-phosphatase (*G6Pase*) and searched for potential trans-acting factors regulating the expression of these genes. In this review, I will introduce my research path which has mainly focused on the characterization of the transcriptional network of genes involved in the glucose homeostasis.

STARTING FROM RAT GLUT2 PROMOTER

In the 1990s, cloning of genes with upstream regulatory elements was the mainstream biochemistry field. Because it was the previous era of the Genome Project we had to manually sequence a lot of short gene fragments using Sanger's dideoxy method to clone the whole rat *GLUT2* gene, which was laborious and time consuming. Dr. Gil-Soo Han, my first research associate, started the project, but left my Lab in 1993 before the work was completed. So, we had to determine sequence of exon 1 with 5'-flanking region. Repeated screenings of lambda phage clones finally revealed the exon 1 and upstream sequences, and furthermore, using rapid amplification of 5' complementary DNA ends (5'-RACE) and primer extension,

Corresponding author: Ahn Yong Ho  <https://orcid.org/0000-0002-4133-0757>
Department of Biochemistry and Molecular Biology, Integrated Genomic Research Center for Metabolic Regulation, Yonsei University College of Medicine, 50 Yonsei-ro, Seodaemun-gu, Seoul 03722, Korea
E-mail: yha111@yuhs.ac

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received: Jul. 30, 2018; Accepted: Aug. 8, 2018

Copyright © 2018 Korean Diabetes Association <http://e-dmj.org>

we were able to identify complete structure of rat *GLUT2* gene, which has four split exon 1 (1a to 1d). This discovery let us to obtain the rat *GLUT2* promoter region, which conveys many potential transcription regulatory elements [2]. Next, we tried to find the mechanism of regulation of *GLUT2* gene expression in liver. *GLUT2* is a major glucose transporter in liver and pancreatic β -cells, and we initially focused to liver because nuclear extract from this organ is easier to prepare to study DNA-protein interaction, including DNase I footprinting and electrophoretic mobility shift assay (EMSA). The DNase I footprinting assay was a difficult and laborious experiment and took a long time, but finally we demonstrated CCAAT/enhancer binding protein (C/EBP) elements in the *GLUT2* promoter. Furthermore, it was also identified that the regulation by C/EBPs were gradually attenuated in the primary cultured hepatocytes, providing an evidence why primary hepatocytes on culture dishes lose their ability to express *GLUT2* [3]. These earlier studies became foundation stone of my research.

EXPANSION TO HUMAN *GLUT2* PROMOTER

GLUT2 gene is mainly expressed in liver and pancreatic β -cells; however, the transcription factors and cis-elements regulating the tissue-specific expression of the *GLUT2* were largely unknown. Transfection studies using pancreatic β -cells (HIT-T15, MIN6), hepatoma cell (HepG2), and fibroblast cells (NIH3T3, HeLa) revealed the differences in the promoter activities and site C (87 to 132) responsible for tissue-specific expression of the human *GLUT2* gene [4]. We identified that both hepatocyte nuclear factor 1 (HNF1) and HNF3 function as transcriptional activators in *GLUT2* gene expression. The *GLUT2* mRNA level was well correlated with the status of HNF1 and HNF3 mRNA levels regardless of the origins or types of cells. Furthermore, we showed that the promoter activity of the mutG construct containing the 103A→G mutation, which was found in a Korean non-insulin-dependent diabetes mellitus (NIDDM) patients, was in part compensated by the binding of nuclear factor Y (NF-Y) instead of HNF1 or HNF3 [5].

NUCLEAR RECEPTOR AND GLUCOSE SENSOR REGULATION IN LIVER AND β -CELLS

In the late 1990s, peroxisome proliferator-activated receptor γ

(PPAR γ) was identified as a target for thiazolidinedione (TZD), the first insulin sensitizing agent [6]. Anti-diabetic action of TZD was thought to be mediated by PPAR γ in adipose tissues until we identified that TZD restores the *GLUT2* expression in pancreatic β -cells of type 2 diabetes mellitus (T2DM) model mice. While searching for transcription factors regulating *GLUT2* gene, we noticed the presence of PPAR γ response element (PPRE) in 5' untranslated region of the *GLUT2* gene. Although PPAR γ in β -cells is not abundant, TZD treatment directly up-regulated PPAR γ -dependent *GLUT2* expression and restored glucose sensing ability of β -cell in diabetic condition [7]. Subsequently, we identified PPRE in β -cell specific glucokinase (β Gck) promoter and showed its functional relevance in β -cells [8].

After that, we studied liver glucokinase (L-Gck) promoter, because we wanted to stay focused in the hepatic glucose sensor in connection with glucose homeostasis. In liver, we were able to identify and characterize the role of PPAR γ in the regulation of L-Gck [9,10]. Next, we demonstrated the interrelationship among PPAR γ , liver X-receptor α (LXR α), sterol regulatory element binding protein-1c (SREBP-1c), and small heterodimer partner (SHP) in the regulation of hepatic glucokinase expression [11].

SREBP-1c AND GLUCOSE METABOLISM

The regulation of hepatic glucose metabolism is important in glucose homeostasis and L-Gck plays a central role in this process. L-Gck expression is known to be regulated by insulin and SREBP-1c was suggested to be a transcription factor that mediates the action of insulin on Gck transcription in liver. However, the mechanism how SREBP-1c induce hepatic Gck expression in response to insulin is not well characterized. Using DNA-protein interactions techniques, we identified sterol regulatory elements (SREs) in rat L-Gck promoter and present evidences that insulin increases the binding of SREBP-1c on L-Gck promoter resulting in the increase of the L-Gck transcription [12].

In 2004, we performed an experiment whether SREBP-1c could be a potential trans-acting factor regulating *GLUT2* in liver. At that time, *GLUT2* gene expression was not known to be regulated by insulin. And thus, SREBP-1c could not be a transcriptional coactivator of *GLUT2* gene expression. However, we observed that both *GLUT2* and *SREBP-1c* gene expression were increased in hyperglycemic states. Moreover, we

demonstrated that SREBP-1c binding region (SRE) is present in the mouse GLUT2 promoter. With these data, we concluded that glucose-stimulated activation of *GLUT2* gene expression might be induced by SREBP-1c in primary cultured hepatocytes. Adenoviral expression of the dominant negative form of SREBP-1c suppressed glucose-stimulated GLUT2 mRNA level in primary cultured hepatocytes [13].

We also showed regulatory mechanism of *GLUT4* gene expression in adipocytes. With increased GLUT4 expression, mRNA levels of SREBP-1c in adipose tissue were proportionally increased by refeeding. Insulin treatment increased mRNA levels of GLUT4 in adipose tissue. Because *SREBP-1c* gene expression is regulated by insulin, it is thought that GLUT4 might be regulated by SREBP-1c. On this basis, we examined presence of a potential site binding SBREP-1c in the GLUT4 promoter and found highly conserved SRE. Through this study, we published a paper entitled, "Insulin-mediated activation of human GLUT4 promoter is directly induced by SREBP-

1c in adipocytes" [14].

Subsequent to this study, we hypothesized that transcription factor E3 (TFE3) could be able to up-regulate *Gck* because TFE3 was shown to up-regulate genes of insulin signaling pathway [15]. Using adenoviral transduction of TFE3, we demonstrated the presence of TFE3 binding site (E-box) in the *Gck* and TFE3 up-regulates *Gck* with amelioration of hyperglycemia in the *db/db* and *ob/ob* mice [16]. Fig. 1 shows a summary of *cis*-elements and trans-acting factors acting on the glucose sensor genes in liver explored by my team.

ROLE OF PPAR α AND PROLACTIN REGULATORY ELEMENT-BINDING ON THE GLUCONEOGENESIS

When glucose level is low, blood glucose level is maintained by hepatic gluconeogenesis. Because increase in a hepatic glucose production occurs in the post-absorptive state of T2DM, it is




Fig. 1. Schematic summary of the regulation of liver type glucose transporter (glucose transporter type 2 [GLUT2]) and glucokinase (Gck) promoter. During the long journey to understand glucose homeostasis, main interest of my group was how glucose sensors, GLUT2 and glucokinase, are regulated by transcription factors. Tissue-specific study, and disease-specific alteration as well, revealed that many transcriptional factors are interconnected in the regulation of glucose sensor in liver. C/EBP, CCAAT/enhancer binding protein; PPAR γ , peroxisome proliferator-activated receptor γ ; RXR α , retinoid X receptor α ; PPRE, PPAR γ response element; SREBP-1c, sterol regulatory element binding protein-1c; SRE, sterol regulatory element; HNF1 α , hepatocyte nuclear factor 1 α ; HNF, hepatocyte nuclear factor; FOXA2, forkhead box A2; LXR α , liver X-receptor α ; SHP, small heterodimer partner; L-GcK, liver glucokinase; TFE3, transcription factor E3; LXRE, liver X-receptor response element.

required to unveil the mechanisms of how hyperglycemia develops. To explore a pathogenesis, we are interested in a role of PPAR α and prolactin regulatory element-binding (PREB). First, we have performed experiments based on the previous observation that PPAR α -null mice shows severe hypoglycemia following 24-hour fasting, characterized by a 50% drop in blood glucose level, suggesting a potential role of PPAR α in glucose metabolism [17]. Using PPAR α knockout (KO) mice as an animal model, we were able to demonstrate presence of PPRE and indicated that PPAR α might be responsible for glucose production through the regulation of hepatic *G6pase* expression [18]. Most recently, we observed that PREB protein is down-regulated in the liver of *db/db*, *ob/ob*, and high-fat diet-induced obese mice. This observation led us to explore a possibility that PREB could be a transacting factor down-regulating gluconeogenic. Indeed, we found a *cis*-element, prolactin core-binding element in the promoters of gluconeogenic genes, which is responsible for down-regulation of these genes, that is, decreased PREB in the liver of insulin resistant animal cannot repress the expression of gluconeogenic genes (*G6Pase* and phosphoenolpyruvate carboxykinase [*Pck*]), resulting in the hyperglycemia, a typical manifestation of the diabetic model mice or high fat induced obesity mice [19].

ROLE OF PROTEIN INTERACTIONS AND GLUCOSE HOMEOSTASIS

Glucose homeostasis is of prime importance in maintaining whole body metabolism. In the insulin resistant states, a balance in the intermediary metabolism may be disturbed. During prolonged starvation, hepatic gluconeogenesis is a major pathway that maintains normal blood glucose levels. In the postprandial state, temporarily increased glucose needs to be disposed by liver to prevent the cells from glucotoxicity. In this process, Gck is known to play the most important role [20].

Although we have been stayed focus on the transcriptional aspects of metabolic genes in liver, we have also searched for the possibility that post-translational modification of metabolic enzymes/protein could also be a novel regulatory mechanism. Using current proteome technology, we were able to demonstrate that glucokinase regulatory protein (GKRP) in liver is acetylated by acetyltransferase p300. Acetylated GKRP decreases ubiquitination of the protein itself with an increase in its affinity to Gck, resulting in increased nuclear retention of Gck and decreased glycolytic flux [21].

To understand this, we have done couple of studies to elucidate mechanisms of how hepatic gluconeogenesis occurs. From the beginning, we showed that insulin-mediated repression of *G6pase* and *Pck* gene expression is restored by resveratrol, which might be associated with sirtuin 1 (Sirt1) mediated forkhead box O1 (FOXO1) deacetylation [22]. And then, we found a role of thioredoxin-interacting protein (TXNIP) in the hyperglycemia. The increased TXNIP in the hyperglycemic mice model, impairs glucose and insulin tolerance in mice by up-regulating *G6pase* through interaction with SHP. Binding of TXNIP to SHP resulted in the degradation of SHP through ubiquitination, allowing FOXO1 to up-regulate gluconeogenic gene expression [23].

ROLE OF CREB3L4 ON THE ADIPOGENESIS

Another field of our research was transcriptional connections in adipogenesis. We were interested in obesity problem, which became a worldwide epidemic. To this end, we wanted to explore signaling networks governed by transcription factors involved in the adipogenesis, ultimately to understand the pathophysiology of obesity and T2DM. Using cyclic AMP-responsive element-binding protein 3 (CREB3)-like 4 (*CREB3L4*) KO mice, we demonstrated that CREB3L4 acts as a negative regulator of adipogenesis [24]. In this paper we demonstrated that CREB3L4 binds C/EBP β , promoting its ubiquitination, resulting in the inhibition of adipogenesis and obesity in mice (Fig. 2A). Thus, if a measure to trigger *CREB3L4* expression is taken, it could be one of novel way to treat obesity or T2DM. In connection with this work, we published a paper entitled, "The effects of low fat diet (LFD) or aging on the metabolic profiles of *Creb3l4* KO mice," which showed significant weight gain and adiposity with impaired glucose tolerance and decreased insulin sensitivity in *Creb3l4* KO mice [25]. Furthermore, we demonstrated that CREB3L4 is required for proliferation of prostate cancer cells and CREB3L4 is a crucial activator of androgen receptor (AR) function. CREB3L4 directly interacts with, and facilitates, AR recruitment to the androgen responsive elements (AREs) of AR target genes. In addition, indirectly, through inositol requiring enzyme 1 α signaling, a distinct AR-endoplasmic reticulum stress-CREB3L4 regulatory axis also plays a role in prostate cancer proliferation (Fig. 2B) [26]. Table 1 shows a summary of major works done by my team for the last 25 years.




Fig. 2. Role of cyclic AMP-responsive element-binding protein 3 (CREB3)-like 4 (CREB3L4) on the adipogenesis. (A) In preadipocytes, CREB3L4 acts as a negative regulator of adipogenesis by both regulating the stability of CCAAT/enhancer binding protein β (C/EBP β) protein and increasing GATA binding protein 3 (GATA3) expression. (B) Proposed mechanism of action of CREB3L4 in prostate cancer progression. Adapted from Kim et al. [24,26]. IBMX, 3-isobutyl-1-methylxanthine; AR, androgen receptor; IRE1 α , inositol requiring enzyme 1 α .

Table 1. Summary of major research papers of my team

Year	Journal	Research subject	Contributor(s)
1995	Arch Biochem Biophys	L-GLUT2 promoter cloning	Ahn et al. [2]
1998	Biochem J	L-GLUT2 promoter, C/EBP	Kim et al. [3]
2000	J Biol Chem	L-GLUT2 liver specific expression, HNF1&3	Cha et al. [4]
2000	Diabetes	L-GLUT2 promoter, PPRE	Kim et al. [7]
2002	Diabetes	B-Glucokinase, PPAR γ	Kim et al. [8]
2004	Diabetes	L-GcK promoter, PPAR γ	Kim et al. [10]
2004	J Biol Chem	L-GcK promoter, SREBP-1c	Kim et al. [12]
2005	Diabetes	L-GLUT2 promoter, SREBP-1c	Im et al. [13]
2006	Biochem J	GLUT4 promoter, SREBP-1c adipocytes	Im et al. [14]
2009	J Biol Chem	L-GcK promoter, LXR, SREBP-1c, PPAR γ	Kim et al. [11]
2011	J Biol Chem	G6Pase promoter, PPAR α	Im et al. [18]
2013	Diabetologia	L-GcK gene, TFE3	Kim et al. [16]
2013	Diabetologia	Liver gluconeogenic gene, Txnip	Jo et al. [23]
2014	Cell Death Dis	Adipogenesis and CREB3L4	Kim et al. [24]
2015	Sci Rep	L-GcK and GKRP acetylation	Park et al. [21]
2017	Sci Rep	Prostate cancer and CREB3L4	Kim et al. [26]
2018	BBA Mol Basis Dis	Liver PREB and glucose homeostasis	Park et al. [19]

L-GLUT2, liver-glucose transporter type 2; C/EBP, CCAAT/enhancer binding protein; HNF, hepatocyte nuclear factor; PPRE, PPAR γ response element; PPAR, peroxisome proliferator-activated receptor; L-GcK, liver glucokinase; SREBP-1c, sterol regulatory element binding protein-1c; LXR, liver X-receptor; G6Pase, glucose-6-phosphatase; TFE3, transcription factor E3; CREB3L4, cyclic AMP-responsive element-binding protein 3 (CREB3)-like 4; GKRP, glucokinase regulatory protein; PREB, prolactin regulatory element-binding.

SUMMARY

My team has been exploring the roles of transcription factors regulating the genes involved in glucose metabolism for the last 30 years. The main field of study can be summarized as (1) identification of promoters of genes of glucose homeostasis, (2) exploration of interaction between promoters and transcription factors leading to tissue-specific gene expression, (3) modification and interactions between transcription factors in metabolic diseases such as diabetes or obesity, and (4) role of a transcription factor in the adipocyte differentiation and development of obesity. While studying transcriptional regulation, my team members were interconnected as if we were transcription factors coordinating with each other. As an analogy, I might act as a basal transcription factor and my team-mates, wherever they are, played co-activator role. Without their devotion in science, the works of my group could not be made possible.

CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

ACKNOWLEDGMENTS

I would like to thank all the people who worked with me at the same time or at different times. Also, I would like to apologize that I could not mention all the researches in this article, in which every single study carried out was meaningful. The manuscript was mainly prepared by Drs, Mi Young Kim, Seung Soon Im, Ha Il Kim, Ji Young Cha, and Jae Woo Kim. We thank Dong-Su Jang for the schematic representation of the model of CREB3L4 action.

REFERENCES

- Schuit FC, Huypens P, Heimberg H, Pipeleers DG. Glucose sensing in pancreatic beta-cells: a model for the study of other glucose-regulated cells in gut, pancreas, and hypothalamus. *Diabetes* 2001;50:1-11.
- Ahn YH, Kim JW, Han GS, Lee BG, Kim YS. Cloning and characterization of rat pancreatic beta-cell/liver type glucose transporter gene: a unique exon/intron organization. *Arch Biochem Biophys* 1995;323:387-96.
- Kim JW, Ahn YH. CCAAT/enhancer binding protein regulates the promoter activity of the rat GLUT2 glucose transporter gene in liver cells. *Biochem J* 1998;336(Pt 1):83-90.
- Cha JY, Kim H, Kim KS, Hur MW, Ahn Y. Identification of transacting factors responsible for the tissue-specific expression of human glucose transporter type 2 isoform gene. Cooperative role of hepatocyte nuclear factors 1alpha and 3beta. *J Biol Chem* 2000;275:18358-65.
- Cha JY, Kim HS, Kim HI, Im SS, Kim SY, Kim JW, Yeh BI, Ahn YH. Analysis of polymorphism of the GLUT2 promoter in NIDDM patients and its functional consequence to the promoter activity. *Ann Clin Lab Sci* 2002;32:114-22.
- Nolan JJ, Ludvik B, Beerdsen P, Joyce M, Olefsky J. Improvement in glucose tolerance and insulin resistance in obese subjects treated with troglitazone. *N Engl J Med* 1994;331:1188-93.
- Kim HI, Kim JW, Kim SH, Cha JY, Kim KS, Ahn YH. Identification and functional characterization of the peroxisomal proliferator response element in rat GLUT2 promoter. *Diabetes* 2000;49:1517-24.
- Kim HI, Cha JY, Kim SY, Kim JW, Roh KJ, Seong JK, Lee NT, Choi KY, Kim KS, Ahn YH. Peroxisomal proliferator-activated receptor-gamma upregulates glucokinase gene expression in beta-cells. *Diabetes* 2002;51:676-85.
- Kim HI, Ahn YH. Role of peroxisome proliferator-activated receptor-gamma in the glucose-sensing apparatus of liver and beta-cells. *Diabetes* 2004;53 Suppl 1:S60-5.
- Kim SY, Kim HI, Park SK, Im SS, Li T, Cheon HG, Ahn YH. Liver glucokinase can be activated by peroxisome proliferator-activated receptor-gamma. *Diabetes* 2004;53 Suppl 1:S66-70.
- Kim TH, Kim H, Park JM, Im SS, Bae JS, Kim MY, Yoon HG, Cha JY, Kim KS, Ahn YH. Interrelationship between liver X receptor alpha, sterol regulatory element-binding protein-1c, peroxisome proliferator-activated receptor gamma, and small heterodimer partner in the transcriptional regulation of glucokinase gene expression in liver. *J Biol Chem* 2009;284:15071-83.
- Kim SY, Kim HI, Kim TH, Im SS, Park SK, Lee IK, Kim KS, Ahn YH. SREBP-1c mediates the insulin-dependent hepatic glucokinase expression. *J Biol Chem* 2004;279:30823-9.
- Im SS, Kang SY, Kim SY, Kim HI, Kim JW, Kim KS, Ahn YH. Glucose-stimulated upregulation of GLUT2 gene is mediated by sterol response element-binding protein-1c in the hepatocytes. *Diabetes* 2005;54:1684-91.
- Im SS, Kwon SK, Kang SY, Kim TH, Kim HI, Hur MW, Kim

- KS, Ahn YH. Regulation of GLUT4 gene expression by SREBP-1c in adipocytes. *Biochem J* 2006;399:131-9.
15. Nakagawa Y, Shimano H, Yoshikawa T, Ide T, Tamura M, Furusawa M, Yamamoto T, Inoue N, Matsuzaka T, Takahashi A, Hasty AH, Suzuki H, Sone H, Toyoshima H, Yahagi N, Yamada N. TFE3 transcriptionally activates hepatic IRS-2, participates in insulin signaling and ameliorates diabetes. *Nat Med* 2006;12:107-13.
 16. Kim MY, Jo SH, Park JM, Kim TH, Im SS, Ahn YH. Adenovirus-mediated overexpression of Tcf3 ameliorates hyperglycaemia in a mouse model of diabetes by upregulating glucokinase in the liver. *Diabetologia* 2013;56:635-43.
 17. Kersten S, Seydoux J, Peters JM, Gonzalez FJ, Desvergne B, Wahli W. Peroxisome proliferator-activated receptor alpha mediates the adaptive response to fasting. *J Clin Invest* 1999;103:1489-98.
 18. Im SS, Kim MY, Kwon SK, Kim TH, Bae JS, Kim H, Kim KS, Oh GT, Ahn YH. Peroxisome proliferator-activated receptor {alpha} is responsible for the up-regulation of hepatic glucose-6-phosphatase gene expression in fasting and db/db mice. *J Biol Chem* 2011;286:1157-64.
 19. Park JM, Kim MY, Kim TH, Min DK, Yang GE, Ahn YH. Pro-lactin regulatory element-binding (PREB) protein regulates hepatic glucose homeostasis. *Biochim Biophys Acta* 2018;1864(6 Pt A):2097-107.
 20. Niswender KD, Shiota M, Postic C, Cherrington AD, Magnusson MA. Effects of increased glucokinase gene copy number on glucose homeostasis and hepatic glucose metabolism. *J Biol Chem* 1997;272:22570-5.
 21. Park JM, Kim TH, Jo SH, Kim MY, Ahn YH. Acetylation of glucokinase regulatory protein decreases glucose metabolism by suppressing glucokinase activity. *Sci Rep* 2015;5:17395.
 22. Park JM, Kim TH, Bae JS, Kim MY, Kim KS, Ahn YH. Role of resveratrol in FOXO1-mediated gluconeogenic gene expression in the liver. *Biochem Biophys Res Commun* 2010;403:329-34.
 23. Jo SH, Kim MY, Park JM, Kim TH, Ahn YH. Txnip contributes to impaired glucose tolerance by upregulating the expression of genes involved in hepatic gluconeogenesis in mice. *Diabetologia* 2013;56:2723-32.
 24. Kim TH, Jo SH, Choi H, Park JM, Kim MY, Nojima H, Kim JW, Ahn YH. Identification of Creb3l4 as an essential negative regulator of adipogenesis. *Cell Death Dis* 2014;5:e1527.
 25. Kim TH, Park JM, Jo SH, Kim MY, Nojima H, Ahn YH. Effects of low-fat diet and aging on metabolic profiles of Creb3l4 knockout mice. *Nutr Diabetes* 2015;5:e179.
 26. Kim TH, Park JM, Kim MY, Ahn YH. The role of CREB3L4 in the proliferation of prostate cancer cells. *Sci Rep* 2017;7:45300.