

Funding

This work was supported by the British Heart Foundation [FS/10/62/28409 to S.P.] and Dutch ZonMw [104003001 to J.W].

Conflict of interest: None declared.

References

- Vaitkus PT, Barnathan ES. Embolic potential, prevention and management of mural thrombus complicating anterior myocardial infarction: a meta-analysis. *J Am Coll Cardiol* 1993; **22**:1004–9.
- Solheim S, Seljeflot I, Lunde K, Bjørnerheim R, Aakhus S, Forfang K et al. Frequency of left ventricular thrombus in patients with anterior wall acute myocardial infarction treated with percutaneous coronary intervention and dual anti-platelet therapy. *Am J Cardiol* 2010; **106**:1197–200.
- Visser CA, Kan G, Meltzer RS, Dunning AJ, Roelandt J. Embolic potential of left ventricular thrombus after myocardial infarction: a two-dimensional echocardiographic study of 119 patients. *J Am Coll Cardiol* 1985; **5**:1276–80.
- Celik S, Baykan M, Erdöl C, Gökce M, Durmus İ, Örem C et al. Doppler-derived mitral deceleration time as an early predictor of left ventricular thrombus after first anterior acute myocardial infarction. *Am Heart J* 2000; **140**:772–6.
- Maze SS, Kotler MN, Parry WR. Flow characteristics in the dilated left ventricle with thrombus: qualitative and quantitative Doppler analysis. *J Am Coll Cardiol* 1989; **13**:873–81.
- Delemarre BJ, Visser CA, Bot H, Dunning AJ. Prediction of apical thrombus formation in acute myocardial infarction based on left ventricular spatial flow pattern. *J Am Coll Cardiol* 1990; **15**:355–60.
- Eriksson J, Bolger AF, Ebbers T, Carlsson C-J. Four-dimensional blood flow-specific markers of LV dysfunction in dilated cardiomyopathy. *Eur Heart J Cardiovasc Imaging* 2013; **14**:417–24.
- Arvidsson PM, Töger J, Heiberg E, Carlsson M, Arheden H. Quantification of left and right atrial kinetic energy using four-dimensional intracardiac magnetic resonance imaging flow measurements. *J Appl Physiol* 2013; **114**:1472–81.
- Eriksson J, Dyverfeldt P, Engvall J, Bolger AF, Ebbers T, Carlsson C-J. Quantification of presystolic blood flow organization and energetics in the human left ventricle. *Am J Physiol Heart Circ Physiol* 2011; **300**:H2135–41.
- Wong J, Chabiniok R, deVecchi A, Dedieu N, Sammut E, Schaeffter T et al. Age-related changes in intraventricular kinetic energy: a physiological or pathological adaptation? *Am J Physiol Heart Circ Physiol* 2016; **310**:H747–55.
- Svalbring E, Fredriksson A, Eriksson J, Dyverfeldt P, Ebbers T, Bolger AF et al. Altered diastolic flow patterns and kinetic energy in subtle left ventricular remodeling and dysfunction detected by 4D Flow MRI. *PLoS One* 2016; **11**:e0161391.
- Hussain SF, Roldán-Alzate A, Francois CJ. Left and right ventricular kinetic energy using time-resolved versus time-average ventricular volumes. *J Cardiovasc Magn Reson* 2015; **17**:P67.
- Kanski M, Arvidsson PM, Töger J, Borgquist R, Heiberg E, Carlsson M et al. Left ventricular fluid kinetic energy time curves in heart failure from cardiovascular magnetic resonance 4D flow data. *J Cardiovasc Magn Reson* 2015; **17**:111.
- Garg P, Westenberg JJM, van den Boogaard PJ, Swoboda PP, Aziz R, Foley JRJ et al. Comparison of fast acquisition strategies in whole-heart four-dimensional flow cardiac MR: two-center, 1.5 Tesla, phantom and in vivo validation study. *J Magn Reson Imaging* 2018; **47**:272–81.
- Elbaz MSM, van der Geest RJ, Calkoen EE, Roos A. D, Lelieveldt BP, Roest AA et al. Assessment of viscous energy loss and the association with three-dimensional vortex ring formation in left ventricular inflow: in vivo evaluation using four-dimensional flow MRI. *Magn Reson Med* 2017; **77**:794–805.
- Bissell MM, Hess AT, Biasioli L, Glaze SJ, Loudon M, Pitcher A et al. Aortic dilation in bicuspid aortic valve disease: flow pattern is a major contributor and differs with valve fusion type. *Circ Cardiovasc Imaging* 2013; **6**:499–507.
- Hathcock JJ. Flow effects on coagulation and thrombosis. *Arterioscler Thromb Vasc Biol* 2006; **26**:1729.
- Asinger RW, Mikell FL, Elsperger J, Hodges M. Incidence of left-ventricular thrombosis after acute transmural myocardial infarction. *N Engl J Med* 1981; **305**:297–302.
- Chiarella F, Santoro E, Domenicucci S, Maggioni A, Vecchio C. Predischarge two-dimensional echocardiographic evaluation of left ventricular thrombosis after acute myocardial infarction in the GISSI-3 study. *Am J Cardiol* 1998; **81**:822–7.
- Keren A, Goldberg S, Gottlieb S, Klein J, Schuger C, Medina A et al. Natural history of left ventricular thrombi: their appearance and resolution in the posthospitalization period of acute myocardial infarction. *J Am Coll Cardiol* 1990; **15**:790–800.
- McManus DD, Gore J, Yarzebski J, Spencer F, Lessard D, Goldberg RJ. Recent trends in the incidence, treatment, and outcomes of patients with STEMI and NSTEMI. *Am J Med* 2011; **124**:40–7.

Corrigendum

Corrigendum to: Left ventricular thrombus formation in myocardial infarction is associated with altered left ventricular blood flow energetics [*Eur Heart J Cardiovasc Imaging* 2019; **20**:108–17]

There were three errors in the originally published version:

- Affiliation 3 was incorrect in the original version and should have read ‘Radboudumc, Department of Cardiology, Geert Grooteplein Zuid 10, 6525 GA Nijmegen, The Netherlands.’
- Under the heading ‘CMR examination’, the abbreviation ‘MR’ has been spelt out as ‘mitral regurgitation’ but should have read ‘magnetic resonance’.
- Under the heading ‘Haemodynamic analysis’, the abbreviation ‘MR’ stands for ‘mitral regurgitation’.

doi:10.1093/ehjci/jey148

Online publish-ahead-of-print 1 October 2018

© The Author(s) 2018. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.