



Published in final edited form as:

Methods Mol Biol. 2018 ; 1766: 231–238. doi:10.1007/978-1-4939-7768-0_13.

Genome-wide Mapping of Protein-DNA Interactions on Nascent Chromatin

Chenhuan Xu and Victor G. Corces

Department of Biology, Emory University, 1510 Clifton Rd NE, Atlanta, GA 30322, USA

Abstract

Chromatin Immunoprecipitation (ChIP) is the most widely used method to analyze protein-DNA interactions *in vivo*. Coupled with next generation sequencing, ChIP-seq experiments map protein-DNA interactions in a genome-wide fashion. Here we describe a novel method called nasChIP-seq for mapping genome-wide occupancy of post-translationally modified histones or transcription factors on newly replicated DNA.

Keywords

Chromatin Immunoprecipitation; ChIP-seq; DNA replication; nascent chromatin; Click chemistry

1 Introduction

ChIP assays utilize the specific interaction between an antibody and a protein of interest to pull down the fraction of genomic DNA occupied by this protein, achieving an enrichment of regions occupied by the protein out of all randomly sheared genomic regions [1]. The isolated DNA can be used to prepare a library and sequenced under high-throughput conditions to reveal the underlying sequence content. These sequences can then be mapped to a reference genome to locate the genomic regions harboring the sequence, which can be visualized in a genome browser to show the locations of the enriched occupied regions [2–4]. Tandem ChIP assays (re-ChIP or sequential ChIP) have been used to map the location of protein co-occupancy sites [5, 6], or the location of protein occupied sites on chromatin under a certain context, such as nascent (newly replicated) chromatin [7]. Petruk *et al* used BrdU to label newly replicated DNA in fly embryos, followed by ChIP with antibodies targeting Trithorax group proteins and BrdU sequentially. Quantitative PCR (qPCR) allowed these authors to measure the occupancy of proteins on nascent chromatin [7]. Here we describe a method utilizing EdU to label newly replicated DNA followed by click chemistry to transform EdU into biotin on nascent DNA [8–11]. The replacement of BrdU with EdU enabled us to take advantage of the strong interaction between biotin and streptavidin for further isolation of labeled DNA. Furthermore, we carried out the click chemistry in a facilitative aqueous phase of purified ChIP DNA molecules, instead of the classical way to “click” in a suspension of formaldehyde-crosslinked nuclei [8–11], which severely limits the reaction components to access the EdU-labeled DNA. These key improvements allowed us

to achieve very high labeling efficiency and pull down of the nascent chromatin fraction. Here we describe two examples in which this technique is used for genome-wide mapping of histone modifications or protein occupancy on nascent chromatin. In one example we use 20 min of EdU labeling with 10 million synchronized *Drosophila* Kc167 cells to map the distribution of H3K4me3 on nascent chromatin (Figure 1A). In the second example, we describe the mapping of CTCF occupancy on nascent chromatin with 20 min labeling of 10 million asynchronous H9 human embryonic stem cells (Figure 1B).

2 Materials

Prepare all solutions every three months unless otherwise noted. Molecular biology grade water is the solvent for all solutions unless otherwise noted.

2.1 Cell Synchronization, EdU Labeling, and Chromatin Immunoprecipitation

1. Kc167 cells (ATCC) or H9 cells (WiCell)
2. SFX-Insect cell culture media (HyClone) or StemPro hESC SFM (ThermoFisher)
3. 2 M hydroxyurea solution. Aliquot and store at -20°C (see Note 1).
4. 40 mM EdU (5-ethynyl-2'-deoxyuridine) solution. Aliquot and store at -20°C (see Note 2).
5. 1x DPBS (Dulbecco's Phosphate Buffered Saline).
6. Formaldehyde solution (37%). Store in the dark.
7. 1.25 M glycine solution. Store at 4°C .
8. anti-H3K4me3 antibody. Store at 4°C .
9. anti-CTCF antibody. Store at 4°C .
10. Refer to [14] for other reagents for CHIP.

2.2 Click Chemistry

1. 1 mM Biotin azide in DMSO. Aliquot and store at -20°C .
2. 100 mM CuSO_4 solution. Store at 4°C .
3. 100 mM sodium ascorbate solution. Aliquot and store at -20°C .
4. Microcentrifuge tubes.
5. Ethanol. Store at -20°C .

¹-Hydroxyurea will decompose in water. Make the stock solution monthly and aliquot into small volumes for single use and store at -20°C .

²-Short-term 37°C incubation and vigorous vortexing are sometimes necessary to fully dissolve the 40 mM EdU stock solution stored at -20°C .

6. 5 M NaCl solution.
7. 20 mg/ml glycogen solution. Aliquot and store at -20°C .

2.3 Nascent DNA Pull Down

1. 70% ethanol solution. Store at 4°C .
2. Dynabeads® MyOne™ Streptavidin C1. Store at 4°C .
3. 2x binding and washing buffer: 10 mM Tris-HCl (pH 7.5), 1 mM EDTA, 2 M NaCl. Store at 4°C .
4. 10 mM Tris-HCl (pH 8.0). Store at 4°C .

2.4 Library Preparation

1. NEB End Repair Enzyme Mix. Store at -20°C .
2. Klenow Fragment ($3' \rightarrow 5'$ exo⁻). Store at -20°C .
3. 1 mM dATP. Store at -20°C .
4. T4 DNA Ligase. Store at -20°C .
5. KAPA SYBR FAST qPCR Kit Master Mix (2x). Store at -20°C .
6. Refer to [15] for sequence information on adaptors and PCR primers.
7. Agencourt AMPure XP beads

3 Methods

All steps were performed at room temperature (25°C) unless otherwise noted.

3.1 Cell Synchronization, EdU Labeling, and Chromatin Immunoprecipitation

1. Kc167: Cells were grown to 50% confluence in SFX medium at 25°C in an incubator.

H9: Cells were grown to 60–70% confluence in StemPro hESC medium at 37°C in an incubator under 5% CO_2 .

2. Kc167: Add hydroxyurea to cells at a final concentration of 2 mM, and culture cells in an incubator for 20 hr (see Note 3).

3. Kc167: Pre-mix EdU with fresh SFX medium to obtain a final concentration of $40\ \mu\text{M}$. Discard hydroxyurea-containing medium and wash the cells gently with DPBS once. Add

³Usually the cells will reach 70%–80% confluence after synchronization with hydroxyurea for 20 hr. It should be noted that synchronization is only necessary for DNA replication timing-specific mapping. Our scheme of 20 min labeling after synchronization and release preferentially maps the genomic regions replicated during first 20 min after entry into S phase.

EdU-containing medium to cells (see Note 4). Leave cells in the incubator for 20 min (see Note 5).

H9: Pre-mix EdU with fresh StemPro medium to obtain a final concentration of 50 μM . Pre-warm the medium to 37 $^{\circ}\text{C}$. Add EdU-containing medium to cells. Leave cells in the incubator for 20 min (see Note 6).

4. Pre-mix formaldehyde with DPBS to obtain a final concentration of 1%. Discard EdU-containing medium, add 1% formaldehyde to cells.

5. After 10 min, add glycine solution to reach 0.125 M final concentration.

6. Proceed to ChIP with H3K4me3 or CTCF antibody respectively, and refer to [14] for details.

3.2 Click Chemistry

1. Centrifuge ethanol precipitated ChIP DNA at 13,200 rpm for 30 min at 4 $^{\circ}\text{C}$.

2. Discard supernatant. Add 70% ice-cold ethanol. Invert the microcentrifuge tube several times to wash the pellet (see Note 7).

3. Centrifuge at 13,200 rpm for 5 min.

4. Discard supernatant. Air dry the pellet (see Note 8).

5. Add 174 μl 1xDPBS to the pellet. Incubate at 37 $^{\circ}\text{C}$ for 10 min.

6. Add these reagents to 174 μl DNA solution: 2 μl 1 mM biotin-azide (or DMSO, as control, see Note 9), 4 μl 100 mM CuSO_4 solution. Vortex briefly.

7. Add 20 μl 100 mM sodium ascorbate solution. Immediately vortex for 3 s (see Note 10).

8. Incubate at 37 $^{\circ}\text{C}$ for 1 hr (see Note 11).

⁴. Kc167 cells attach loosely after synchronization with hydroxyurea. Be cautious not to perturb the cells when washing and changing the medium. Always pipette the medium through the side surface of the cultureware.

⁵. The labeling time is highly tunable. For high temporal resolution (the degree of nascentness), choose the shortest labeling time that generates a library with good signal-to-noise ratio after sequencing. For high yield of final DNA product, elongate the labeling time appropriately. Also, cell harvesting can be done after a “chase” period following EdU labeling to map dynamic changes in protein occupancy on nascent chromatin.

⁶. Pre-warm the EdU medium to 37 $^{\circ}\text{C}$ to ensure efficient labeling. When labeling cells in multiple pieces of cultureware at the same time, always place individual cultureware back in the incubator immediately after the EdU medium is added.

⁷. Washing with 70% ethanol helps to remove residual SDS (sodium dodecyl sulfate) carried over from elution buffer in ChIP. SDS is a strong inhibitor of click chemistry. Repeat this step if necessary.

⁸. 5–10 min are optimal for air drying the DNA pellet. Over-drying will lead to low solubility of the pellet and overall reduction of final DNA yield.

⁹. Setting aside a DMSO reaction and making a mock library serves as a negative control for the nascent DNA pull down step. In our hands, amplification of the DMSO library with the same number of PCR cycles yields very little DNA, ruling out the possibility that the nascent DNA after pull down contains some carry-over ChIP DNA, proving the nascent ChIP DNA faithfully reflects the enrichment of protein occupancy on nascent chromatin.

¹⁰. Sodium ascorbate is highly prone to oxidation. Always keep caps on when not using it. Protect from light. Make fresh stock solution monthly. Vortex immediately after adding it to the mixture, ensuring timely reaction with Cu^{2+} before any putative oxidation.

¹¹. Oxygen interferes with click chemistry. Choose microcentrifuge tubes with good airtightness as reaction containers. Place tubes in an airtight container if necessary.

9. Add 9 μ l 5 M NaCl solution and 2 μ l glycogen. Vortex briefly. Add 500 μ l ice-cold ethanol. Vortex vigorously. Incubate at -80°C for 2 hr.

3.3 Nascent DNA Pull Down

1. Centrifuge ethanol precipitated DNA at 13,200 rpm for 30 min at 4°C .
2. Discard supernatant. Add 70% ice-cold ethanol. Invert the microcentrifuge tube several times to wash the pellet (see Note 12).
3. Centrifuge at 13,200 rpm for 5 min.
4. Discard supernatant. Air dry the pellet (see Note 8).
5. Add 100 μ l 10 mM Tris-HCl (pH 8.0) to the pellet. Incubate at 37°C for 10 min.
6. Wash 5 μ l Streptavidin C1 beads with 1 ml 1x binding and washing buffer. Remove supernatant on a magnetic stand. Resuspend the beads in 100 μ l 2x binding and washing buffer. Mix 100 μ l DNA solution with 100 μ l beads suspension. Rotate on a rotor for 30 min.
7. Remove supernatant on a magnetic stand. Wash the beads with 1 ml 1x binding and washing buffer on a rotor for 10 min. Repeat this step once.
8. Remove supernatant on a magnetic stand. Wash the beads with 1 ml 10 mM Tris-HCl (pH 8.0) on a rotor for 5 min.
9. Gently suspend the beads in 24 μ l dH_2O .

3.4 Library Preparation

1. Add 3 μ l reaction buffer and 3 μ l NEB End Repair Enzyme Mix to the bead suspension. Gently mix. Rotate on a rotor for 30 min (see Note 13).
2. Remove supernatant on a magnetic stand. Wash the beads with 1 ml 1x binding and washing buffer on a rotor for 10 min.
3. Remove supernatant on a magnetic stand. Wash the beads with 1 ml 10 mM Tris-HCl (pH 8.0) on a rotor for 5 min.
4. Gently resuspend the beads in 19 μ l dH_2O . Add 3 μ l reaction buffer, 6 μ l 1 mM dATP, and 2 μ l Klenow Fragment ($3' \rightarrow 5'$ exo⁻). Gently mix. Rotate on a rotor at 37°C for 30 min.
5. Repeat steps 2 & 3.
6. Gently resuspend the beads in 19 μ l dH_2O . Add 3 μ l reaction buffer, 5 μ l 10 μM adaptors, and 3 μ l T4 DNA ligase. Gently mix. Rotate on a rotor for 4 hr.

¹²Washing with 70% ethanol helps to remove residual biotin azide carried over from click chemistry. Free biotin will reduce the chance of biotin-labeled DNA to bind to the streptavidin beads. Repeat this step if necessary.

¹³We determined 30–40 μ l as the optimal reaction volume in a 1.5 ml microcentrifuge tube. Volumes lower than this will lead to higher density of beads and higher chance of bead aggregation. Higher volumes will lead to spill of beads to the side surface of the microcentrifuge tube during rotation. Both conditions reduce the overall yield of final DNA.

7. Repeat steps 2 & 3.
8. Gently resuspend the beads in 21 μ l dH₂O. Add 2 μ l each of 10 mM primers, and 25 μ l KAPA SYBR FAST qPCR Kit Master Mix (2x). Gently mix. Amplify on a qPCR instrument. Monitor the real-time amplification curve and stop the amplification appropriately (see Note 14).
9. Size-select and clean the DNA using Agencourt AMPure XP beads (Refer to product manuals.).
10. Send the library to sequencing service providers for quality assessment and high throughput sequencing.

Acknowledgements

Work in the authors' laboratory was supported by U.S. Public Health Service Award R01 GM035463 from the National Institutes of Health. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

1. Struhl K (2007) Interpreting Chromatin Immunoprecipitation Experiments In Evaluating Techniques in Biochemical Research, Zuk D, ed. (Cambridge, MA: Cell Press)
2. Barski A, Cuddapah S, Cui K, Roh TY, Schones DE, Wang Z et al. (2007) High-resolution profiling of histone methylations in the human genome. *Cell* 129:823–837 [PubMed: 17512414]
3. Johnson DS, Mortazavi A, Myers RM, Wold B (2007) Genome-wide mapping of in vivo protein-DNA interactions. *Science* 316:1497–1502 [PubMed: 17540862]
4. Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz G et al. (2011) Integrative genomics viewer. *Nat Biotechnol* 29:24–26 [PubMed: 21221095]
5. Chaya D, Hayamizu T, Bustin M, Zaret KS (2001) Transcription factor FoxA (HNF3) on a nucleosome at an enhancer complex in liver chromatin. *J Biol Chem* 276:44385–44389 [PubMed: 11571307]
6. Metivier R, Penot G, Hübner MR, Reid G, Brand H, Kos M et al. (2003) Estrogen receptor-alpha directs ordered, cyclical, and combinatorial recruitment of cofactors on a natural target promoter. *Cell* 115:751–763 [PubMed: 14675539]
7. Petruk S, Sedkov Y, Johnston DM, Hodgson JW, Black KL, Kovermann SK et al. (2012) TrxG and PcG Proteins but Not Methylated Histones Remain Associated with DNA through Replication. *Cell* 150:922–933 [PubMed: 22921915]
8. Sirbu BM, Couch FB, Feigerle JT, Bhaskara S, Hiebert SW, Cortez D (2011) Analysis of protein dynamics at active, stalled, and collapsed replication forks. *Genes Dev* 25:1320–1327 [PubMed: 21685366]
9. Sirbu BM, Couch FB, Cortez D (2012) Monitoring the spatiotemporal dynamics of proteins at replication forks and in assembled chromatin using isolation of proteins on nascent DNA. *Nat Protoc* 7:594–605 [PubMed: 22383038]
10. Lopez-Contreras AJ, Ruppen I, Nieto-Soler M, Murga M, Rodriguez-Acebes S, Remeseiro S et al. (2013) A proteomic characterization of factors enriched at nascent DNA molecules. *Cell Rep* 3:1105–1116 [PubMed: 23545495]

¹⁴PCR amplification with optimal cycle numbers is critical for the quality of the library. Insufficient amplification will lead to insufficient input material for sequencing and failure of the experiment. Over-amplification will lead to higher PCR duplication level (piling-up of clonally identical sequence) and reduce the overall cost-efficiency of sequencing.

11. Aranda S, Rutishauser D, Ernfors P (2014) Identification of a large protein network involved in epigenetic transmission in replicating DNA of embryonic stem cells. *Nucleic Acids Res* 42:6972–6986 [PubMed: 24852249]
12. Li L, Lyu X, Hou C, Takenaka N, Nguyen HQ, Ong CT et al. (2015) Widespread Rearrangement of 3D Chromatin Organization Underlies Polycomb-Mediated Stress-Induced Silencing. *Mol Cell* 58:216–231 [PubMed: 25818644]
13. Gertz JI, Savic D, Varley KE, Partridge EC, Safi A, Jain P et al. (2013) Distinct properties of cell-type-specific and shared transcription factor binding sites. *Mol Cell* 52:25–36 [PubMed: 24076218]
14. Lee TI, Johnstone SE, Young RA (2006) Chromatin immunoprecipitation and microarray-based analysis of protein location. *Nat Protoc* 1:729–748 [PubMed: 17406303]
15. Bowman SK, Simon MD, Deaton AM, Tolstorukov M, Borowsky ML, Kingston RE (2013) Multiplexed Illumina sequencing libraries from picogram quantities of DNA. *BMC Genomics* 14:466. doi: 10.1186/1471-2164-14-466 [PubMed: 23837789]




Figure 1.

Results from ChIP-seq on nascent chromatin. (A) A 75 kb genomic region of the *Drosophila* genome showing a published H3K4me3 ChIP-seq data in asynchronous Kc167 cells (upper track) [12], and our H3K4me3 nasChIP-seq data in synchronized Kc167 cells with 20 min of EdU labeling (lower track). The sequence read density was normalized to *per* one million reads (RPM). The left half of the genomic region shows full restoration of H3K4me3 on nascent chromatin within 20 min, while H3K4me3 is only partially restored in the right half of the region on nascent chromatin. (B) A 5 kb genomic region of the human genome showing a published CTCF ChIP-seq data in H1 hES cells (upper track) [13], and our CTCF nasChIP-seq data in H9 hES cells (lower track). The sequence read density was normalized to *per* one million reads (RPM). One conserved CTCF occupied site and one nascent chromatin-specific site are shown around the transcription start site of the TPBG gene.