



CORRIGENDUM

CHCHD2 accumulates in distressed mitochondria and facilitates oligomerization of CHCHD10

Xiaoping Huang^{1,†}, Beverly P. Wu^{1,†}, Diana Nguyen^{1,†}, Yi-Ting Liu¹, Melika Marani¹, Jürgen Hench², Paule Bénit³, Vera Kozjak-Pavlovic⁵, Pierre Rustin^{3,4}, Stephan Frank² and Derek P. Narendra^{1,*}

¹Inherited Movement Disorders Unit, Neurogenetics Branch, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD 20892, USA, ²Division of Neuropathology, Institute of Pathology, University Hospital Basel, Basel, Switzerland, ³INSERM UMR 1141, Hôpital Robert Debré, Paris, France, ⁴Université Paris Diderot, Paris, France and ⁵Department of Microbiology, Biocenter, University of Wuerzburg, Am Hubland, D-97074 Wuerzburg, Germany

*To whom correspondence should be addressed at: 35 Convent Drive, Bldg 35 Rm 2A215, Bethesda, MD, 20892, USA. Tel: 1301944737; Fax: 13014803365; Email: derek.narendra@nih.gov

Human Molecular Genetics, 2018; doi: 10.1093/hmg/ddy270.

Upon initial publication, Figure 4 incorrectly stated that the mitochondrial marker used was Tom70. This should have been Tom20. This has now been corrected.

The authors regret the error.