

Ketogenic diet for weight loss

Rhonda Ting PharmD MSc Nicolas Dugré PharmD MSc G. Michael Allan MD CCFP Adrienne J. Lindblad ACPH PharmD

Clinical question

Is the ketogenic diet effective for weight loss?

Bottom line

Ketogenic diets can help patients lose about 2 kg more than low-fat diets do at 1 year, but higher-quality studies show no difference. Weight loss peaks at about 5 months but is often not sustained. Individual weight change can vary from losing 30 kg to gaining 10 kg with any diet.

Evidence

- In a systematic review of 13 RCTs of ketogenic versus low-fat diets (N=1577, 61% women, BMI 30 to 43 kg/m²), those on ketogenic diets lost 0.9 kg more than those on low-fat diets at 12 to 24 months (statistically different).¹
 - There were statistically significant but likely clinically meaningless changes in surrogate markers.
 - The dropout rate was 13% to 84% across studies.
- A systematic review of 11 RCTs (N=1369, 71% women, BMI 30 to 36 kg/m²) found at 6 to 24 months² that the ketogenic diet group lost 2.2 kg more than the low-fat diet group (statistically different but results were inconsistent). Higher-quality studies showed no difference.
 - Surrogate marker changes were similar to those above.^{1,2}
- Other systematic reviews (5 to 24 RCTs) were confounded by low-carbohydrate diets that were likely not ketogenic. Results ranged from no difference³⁻⁵ to a 3.6-kg loss.⁶⁻⁸
- No systematic reviews or RCTs examined mortality or cardiovascular disease.²
- An RCT (N=609) found weight loss at 1 year for low-carbohydrate diets (<20 g/d to start) of 6.0 kg compared with 5.3 kg for low-fat diets (not statistically different).⁹
 - Patient genotypes (favouring 1 diet type) had no effect.
 - Weight change varied from losing 30 kg to gaining 10 kg in either group.

Context

- A typical Canadian diet contains 48% carbohydrates, 32% fat, and 17% protein.¹⁰
- Most ketogenic diets start with carbohydrate restriction of less than 20 to 50 g/d (10% of energy intake) for about 2 months before slow reintroduction.^{1,11}
- Weight loss peaks at about 5 months, then weight is slowly regained.¹²
- Ketogenic diets tend to decrease caloric intake.^{9,12}

- Observational data suggest long-term low carbohydrate intake might be associated with increased mortality.¹³

Implementation

Ketogenic diets imply minimizing carbohydrate intake and maximizing protein intake to induce ketosis.¹² Adverse effects are common, including constipation (33%), halitosis (30%), muscle cramps (28%) (numbers needed to harm of 3 to 4 compared with low-fat diets), headache, diarrhea, weakness, and rash (numbers needed to harm of 5 to 7).¹⁴ Urine ketone monitoring is advocated in the lay press, but it is not consistently reported in RCTs and its benefit is unknown. 🌿

Ms Ting is a doctoral student in the Faculty of Pharmacy and Pharmaceutical Sciences at the University of Alberta in Edmonton. **Dr Dugré** is Clinical Assistant Professor in the Faculty of Pharmacy at the University of Montreal in Quebec. **Dr Allan** is Professor with the PEER Group in the Department of Family Medicine at the University of Alberta and the Alberta College of Family Physicians. **Dr Lindblad** is Knowledge Translation and Evidence Coordinator for the Alberta College of Family Physicians and Associate Clinical Professor in the Department of Family Medicine at the University of Alberta.

Competing interests

None declared

The opinions expressed in Tools for Practice article are those of the authors and do not necessarily mirror the perspective and policy of the Alberta College of Family Physicians.

References

- Bueno NB, de Melo IS, de Oliveira SL, da Rocha Ataide T. Very-low-carbohydrate ketogenic diet v. low-fat diet for long-term weight loss: a meta-analysis of randomised controlled trials. *Br J Nutr* 2013;110(7):1178-87.
- Mansoor N, Vinknes KJ, Veierød MB, Retterstøl K. Effects of low-carbohydrate diets v. low-fat diets on body weight and cardiovascular risk factors: a meta-analysis of randomised controlled trials. *Br J Nutr* 2016;115(3):466-79.
- Meng Y, Bai H, Wang S, Li Z, Wang Q, Chen L. Efficacy of low carbohydrate diet for type 2 diabetes mellitus management: a systematic review and meta-analysis of randomized controlled trials. *Diabetes Res Clin Pract* 2017;131:124-31.
- Hu T, Mills KT, Yao L, Demanelis K, Eloustaz M, Yancy WS Jr, et al. Effects of low-carbohydrate diets versus low-fat diets on metabolic risk factors: a meta-analysis of randomized controlled clinical trials. *Am J Epidemiol* 2012;176(Suppl 7):S44-54.
- Huntriss R, Campbell M, Bedwell C. The interpretation and effect of a low-carbohydrate diet in the management of type 2 diabetes: a systematic review and meta-analysis of randomised controlled trials. *Eur J Clin Nutr* 2018;72(3):311-25.
- Nordmann AJ, Nordmann A, Briel M, Keller U, Yancy WS Jr, Brehm BJ, et al. Effects of low-carbohydrate vs low-fat diets on weight loss and cardiovascular risk factors: a meta-analysis of randomized controlled trials. *Arch Intern Med* 2006;166(3):285-93. Erratum in: *Arch Intern Med* 2006;166(8):932.
- Sackner-Bernstein J, Kanter D, Kaul S. Dietary intervention for overweight and obese adults: comparison of low-carbohydrate and low-fat diets. A meta-analysis. *PLoS One* 2015;10(10):e0139817.
- Bravata DM, Sanders L, Huang J, Krumholz HM, Olkin I, Gardner CD, et al. Efficacy and safety of low-carbohydrate diets: a systematic review. *JAMA* 2003;289(14):1837-50.
- Gardner CD, Trepanowski JF, Del Gobbo LC, Hauser ME, Rigdon J, Ioannidis JPA, et al. Effect of low-fat vs low-carbohydrate diet on 12-month weight loss in overweight adults and the association with genotype pattern or insulin secretion. The DIETFITS randomized controlled trial. *JAMA* 2018;319(7):667-79. Errata in: *JAMA* 2018;319(13):1386, *JAMA* 2018;319(16):1728.
- Statistics Canada. *Canadian Community Health Survey – nutrition: nutrient intakes from food and nutritional supplements*. Ottawa, ON: Statistics Canada; 2017. Available from: <https://www150.statcan.gc.ca/n1/daily-quotidien/170620/dq170620b-eng.htm>. Accessed 2018 Sep 20.
- Moreno B, Crujeiras AB, Bellido D, Sajoux I, Casanueva FF. Obesity treatment by very low-calorie-ketogenic diet at two years: reduction in visceral fat and on the burden of disease. *Endocrine* 2016;54(3):681-90.
- Johnstone AM, Horgan GW, Murison SD, Bremner DM, Lobley GE. Effects of a high-protein ketogenic diet on hunger, appetite, and weight loss in obese men feeding ad libitum. *Am J Clin Nutr* 2008;87(1):44-55.
- Seidemann SB, Claggett B, Cheng S, Henglin M, Shah A, Steffen LM, et al. Dietary carbohydrate intake and mortality: a prospective cohort study and meta-analysis. *Lancet Public Health* 2018;3(9):e419-28.
- Yancy WS Jr, Olsen MK, Guyton JR, Bakst RP, Westman EC. A low-carbohydrate, ketogenic diet versus a low-fat diet to treat obesity and hyperlipidemia: a randomized, controlled trial. *Ann Intern Med* 2004;140(10):769-77.

This article is eligible for Mainpro+ certified Self-Learning credits. To earn credits, go to www.cfp.ca and click on the Mainpro+ link.

La traduction en français de cet article se trouve à www.cfp.ca dans la table des matières du numéro de décembre 2018 à la page e529.

Tools for Practice articles in *Canadian Family Physician* are adapted from articles published on the Alberta College of Family Physicians (ACFP) website, summarizing medical evidence with a focus on topical issues and practice-modifying information. The ACFP summaries and the series in *Canadian Family Physician* are coordinated by Dr G. Michael Allan, and the summaries are co-authored by at least 1 practising family physician and are peer reviewed. Feedback is welcome and can be sent to toolsforpractice@cfpc.ca. Archived articles are available on the ACFP website: www.acfp.ca.