

CORRECTION

# Correction: Fecal microbiome of periparturient dairy cattle and associations with the onset of *Salmonella* shedding

The *PLOS ONE* Staff

## Notice of republication

This article was republished on February 06, 2019 to correct for errors in the Data Availability statement introduced during the typesetting process. The publisher apologizes for the errors. Please download this article again to view the correct version. The originally published, uncorrected article and the republished, corrected article are provided here for reference.

## Supporting information

**S1 File. Originally published, uncorrected article.**

(PDF)

**S2 File. Republished, corrected article.**

(PDF)

## Reference

1. Muñoz-Vargas L, Opiyo SO, Digianantonio R, Williams ML, Wijeratne A, Habing G (2018) Fecal microbiome of periparturient dairy cattle and associations with the onset of *Salmonella* shedding. *PLoS ONE* 13(5): e0196171. <https://doi.org/10.1371/journal.pone.0196171> PMID: 29750790



## OPEN ACCESS

**Citation:** The *PLOS ONE* Staff (2019) Correction: Fecal microbiome of periparturient dairy cattle and associations with the onset of *Salmonella* shedding. *PLoS ONE* 14(2): e0212528. <https://doi.org/10.1371/journal.pone.0212528>

**Published:** February 13, 2019

**Copyright:** © 2019 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.