

# Plasmid-mediated colistin resistance gene *mcr-1* in *Escherichia coli* and *Klebsiella pneumoniae* isolated from market retail fruits in Guangzhou, China

Fan Yang,<sup>1,\*</sup> Cong Shen,<sup>2,3,\*</sup>  
Xiaobin Zheng,<sup>4</sup> Yan Liu,<sup>5</sup>  
Mohamed Abd El-Gawad  
El-Sayed Ahmed,<sup>2,3,6</sup> Zihan  
Zhao,<sup>2,3</sup> Kang Liao,<sup>7</sup> Yaling Shi,<sup>8</sup>  
Xin Guo,<sup>2,3</sup> Ruoxuan Zhong,<sup>2,3</sup>  
Zhimin Xu,<sup>2,3</sup> Guo-Bao Tian<sup>2,3</sup>

<sup>1</sup>Department of Microbiology, School of Basic Medical Science, Xinxiang Medical University, Xinxiang, China;

<sup>2</sup>Department of Microbiology, Zhongshan School of Medicine, Sun Yat-sen University, Guangzhou, China;

<sup>3</sup>Key Laboratory of Tropical Diseases Control (Sun Yat-sen University), Ministry of Education, Guangzhou, China;

<sup>4</sup>Department of Respiratory Medicine, The Fifth Affiliated Hospital of Sun Yat-sen University, Zhuhai, China;

<sup>5</sup>Department of Laboratory, The Fifth Affiliated Hospital of Sun Yat-sen University, Zhuhai, China;

<sup>6</sup>Department of Microbiology and Immunology, Faculty of Pharmacy, Misr University for Science and Technology (MUST), Cairo, 6th of October City, Egypt;

<sup>7</sup>Department of Clinical Laboratory, The First Affiliated Hospital of Sun Yat-Sen University, Guangzhou, China;

<sup>8</sup>Department of Clinical Laboratory, Guangzhou Eighth People's Hospital, Guangzhou Medical University, Guangzhou, China

\*These authors contributed equally to this work

Correspondence: Guo-Bao Tian  
Zhongshan School of Medicine, Sun Yat-sen University, 74 Zhongshan 2nd Road, Guangzhou 510080, China  
Tel +86 020 8733 5387  
Fax +86 020 8733 5387  
Email tiangb@mail.sysu.edu.cn

**Background:** As a result of the growing prevalence of the plasmid-mediated mobile colistin resistance gene *mcr-1* among Gram-negative bacteria, the surveillance of *mcr-1* has been globally applied. In our study, we aimed to shed light on the possibility of transmission of *mcr-1*-resistant isolates through market retail fruits.

**Methods and results:** Herein, 133 different fruit surface samples were collected and screened for the different MCR variants (*mcr-1* to *mcr-8*) using PCR and confirmed with sequencing. We identify for the first time *mcr-1*-carrying *Escherichia coli* and *Klebsiella pneumoniae* from market retail fruits in Guangzhou, China. Minimum inhibitory concentrations were detected by the broth microdilution method. Liquid mating was performed to check the transferability of the *mcr-1* gene. Pulsed field gel electrophoresis analysis of S1 nuclease-digested DNA and Southern blotting were performed to check the location of the *mcr-1* gene. Then, whole-genome sequencing and in silico multilocus sequence typing analysis were performed.

**Conclusion:** We showed that *E. coli* GB110 can mediate the spreading of antibiotic resistance genes through the food chain, while *K. pneumoniae* GB015 was considered to be the progenitor of the most successful multidrug-resistant clone. Since fruits are usually consumed fresh, this may serve as a direct source of *mcr-1*-producing bacteria in humans that requires prompt surveillance and intervention to limit the spread of resistance.

**Keywords:** colistin, *mcr-1*, *Escherichia coli*, *Klebsiella pneumoniae*, fruit

## Introduction

Colistin is a polymyxin antibiotic that has been used for many years in veterinary medicine. Nowadays, a need for using colistin in human medicine has evolved as the last resort drug for the treatment of infections caused by multidrug-resistant bacteria, especially carbapenem-resistant Enterobacteriaceae. However, the use of colistin as a last resort antibiotic is seriously threatened by the rise of plasmid-borne mobile colistin resistance (*mcr* family) genes (*mcr-1*, -2, -3, -4, -5, -6, -7 and -8),<sup>1-3</sup> which can spread rapidly via horizontal gene transfer between bacterial strains and species.<sup>4</sup> In August 2016, a survey reported detection of *mcr-1*-positive Enterobacteriaceae from farming soil in Shandong province, China, suggesting that Enterobacteriaceae harboring *mcr-1* can contaminate agricultural products.<sup>5</sup> Here, we report the identification of *mcr-1* in *Escherichia coli* and *Klebsiella pneumoniae* isolated from market retail fruits in Guangzhou, China.

## Methods and results

A total of 133 fruit surface samples were collected from market retail fruits in Guangzhou, China, from June to November 2016, at various intervals. Different fruits were

collected, including apples (n=52), oranges (n=31), tangerines (n=16), pears (n=15), bananas (n=11), pomegranates (n=6) and grapes (n=2). Initially, surface samples from the fruits were wiped with sterile cotton swabs moistened with saline. Each swab contained one fruit surface sample and was transported to the laboratory immediately. The swabs were cultured on to lysogeny broth (LB) liquid medium (Oxoid, Basingstoke, UK), then DNA extraction was performed using the boiling method. All samples were screened for *mcr-1* to *mcr-8* using PCR and confirmed by Sanger sequencing. The primers and conditions used for the PCR assays are listed in Table S1.<sup>3</sup>

Out of 133 fruit samples, *mcr-1* was detected in two samples (1.5%). No *mcr-2* to *mcr-8* genes were identified among these samples. To screen for colistin resistance, the two *mcr-1*-positive samples were directly seeded on LB agar plates containing 4 µg/mL colistin and incubated at 37°C for 24 hours. The species of selected colonies from the LB agar plates were identified using the API 20E system (bioMérieux, Marcy l'Etoile, France) and confirmed by 16S rRNA sequencing. As a result, *mcr-1*-harboring *E. coli* GB110 and *K. pneumoniae* GB015 were identified from apple and orange samples, respectively. Minimum inhibitory concentrations (MICs) were detected by the broth microdilution method and interpreted according to the Clinical and Laboratory Standards Institute guidelines (CLSI, 2017). It was found that *E. coli* GB110 was susceptible to all tested agents except

colistin and polymyxin B; and *K. pneumoniae* GB015 was resistant to colistin, polymyxin B and ampicillin (Table 1).


A conjugation experiment was performed using streptomycin-resistant *E. coli* C600 as the recipient. The *mcr-1*-producing isolates and recipient were mixed (ratio of 1:9) in LB and incubated overnight at 37°C. The mixture was then spread on LB agar plates containing colistin (2 µg/mL) and streptomycin (2,000 µg/mL). The transconjugants were confirmed for *mcr-1* by PCR and Sanger sequencing. The results showed that *mcr-1* was successfully transferred to streptomycin-resistant *E. coli* C600 through conjugation in both isolates, suggesting that *mcr-1* was located on transferable plasmids. Plasmid incompatibility groups were identified by PCR assay as described previously.<sup>6</sup> Pulsed field gel electrophoresis analysis of S1 nuclease-digested DNA (S1-PFGE), followed by Southern blotting, showed that *mcr-1* was located on ~62.9 kb IncFIA and ~204.2 kb IncHI1 plasmids for the transconjugants of *E. coli* GB110 and *K. pneumoniae* GB015, respectively (Figure 1).

The genome DNA of *E. coli* GB110 and *K. pneumoniae* GB015 were extracted using a Qiagen Blood & Tissue kit (Qiagen, Hilden, Germany) and DNA libraries were constructed with 350 bp paired-end fragments. In total, 9,367,660 and 10,515,584 paired-end 150 bp reads were produced by the Illumina HiSeq2000 platform for *E. coli* GB110 and *K. pneumoniae* GB015, respectively. Reads were assembled using SPAdes

**Table 1** Characteristics of *mcr-1*-producing *Escherichia coli* GB110, *Klebsiella pneumoniae* GB015 and their transconjugants isolated from market retail fruits in China

Isolate code	Source	Isolation site	MLST	Plasmid replicon type	Encoding genes of resistance	MICs
<i>E. coli</i> GB110	Apple	Market retail fruits	ST189	IncHI1, IncFIA	<i>aadA2</i> , <i>aadA1</i> , <i>mcr-1</i> , <i>floR</i> , <i>cmiA1</i> , <i>sul2</i> , <i>sul3</i> , <i>tetA</i> , <i>tetM</i> , <i>dfrA12</i> , <i>mdfA</i>	CL (16), PB (8), TGC (≤1), AMP (4), AMC (8), CTX (≤0.125), CAZ (≤0.125), FEP (0.25), GEN (≤1), AMK (4), ETP (≤0.125), IPM (≤0.125), MEM (≤0.125), FOS (≤8), NIT (≤8), CIP (≤0.03)
<i>K. pneumoniae</i> GB015	Orange	Market retail fruits	ST442	IncHI1, IncFIB	<i>bla<sub>SHV-110</sub></i> , <i>mcr-1</i> , <i>qnrS1</i> , <i>oqxA</i> , <i>oqxB</i> , <i>fosA6</i> , <i>sul1</i> , <i>tetA</i> , <i>dfrA1</i>	CL (64), PB (64), TGC (≤1), AMP (64), AMC (8), CTX (≤0.125), CAZ (0.5), FEP (≤0.125), GEN (≤1), AMK (4), ETP (≤0.125), IPM (0.5), MEM (≤0.125), FOS (16), NIT (64), CIP (1)
<i>E. coli</i> C600 (transconjugant of <i>E. coli</i> GB110)	–	–	–	IncFIA	<i>mcr-1</i>	CL (16), PB (8), TGC (≤1), AMP (4), AMC (8), CTX (≤0.125), CAZ (≤0.125), FEP (≤0.125), GEN (≤1), AMK (4), ETP (≤0.125), IPM (≤0.125), MEM (≤0.125), FOS (≤8), NIT (8), CIP (≤0.03)
<i>E. coli</i> C600 (transconjugant of <i>K. pneumoniae</i> GB015)	–	–	–	IncHI1	<i>mcr-1</i>	CL (16), PB (16), TGC (≤1), AMP (4), AMC (8), CTX (≤0.125), CAZ (≤0.125), FEP (≤0.125), GEN (≤1), AMK (4), ETP (≤0.125), IPM (≤0.125), MEM (≤0.125), FOS (≤8), NIT (≤8), CIP (≤0.03)

**Abbreviations:** AMC, amoxicillin–clavulanic acid; AMK, amikacin; AMP, ampicillin; CAZ, ceftazidime; CIP, ciprofloxacin; CL, colistin; CTX, cefotaxime; ETP, ertapenem; FEP, cefepime; FOS, fosfomycin; GEN, gentamicin; IPM, imipenem; MEM, meropenem; MIC, minimum inhibitory concentration; MLST, multilocus sequence typing; NIT, nitrofurantoin; PB, polymyxin B; TGC, tigecycline.



**Figure 1** SI-PFGE and Southern hybridization with the *mcr-1* probe.

**Abbreviations:** *E. coli*, *Escherichia coli*; *K. pneumoniae*, *Klebsiella pneumoniae*; SI-PFGE, pulsed field gel electrophoresis analysis of S1 nuclease-digested DNA.

v3.12.0.<sup>7</sup> This Whole Genome Shotgun project has been deposited at GenBank under the accession number PRJNA482733. The genome annotation was performed using PATRICK (<https://www.patricbrc.org/>) and ISFinder (<https://www-is.biotoul.fr/>). The assembled genomes showed that IS*Apl1* was located upstream of *mcr-1* for *E. coli* GB110 and downstream of *mcr-1* for *K. pneumoniae* GB015, consistent with reported variability in the location of IS*Aba1*, which was probably involved in the original mobilization of *mcr-1* from *Moraxella* spp.<sup>4,8</sup>

Multilocus sequence typing (MLST), serotyping, antimicrobial resistance genes and virulence factors were annotated by the Center for Genomic Epidemiology website (<http://www.genomicepidemiology.org/>).

MLST analysis of *E. coli* GB110 (<http://mlst.warwick.ac.uk/mlst/dbs/Ecoli>) showed that it belonged to sequence type 189 (165 cplx), which was reported from poultry retail meat mediating the spread of extended-spectrum  $\beta$ -lactamase (ESBL) genes in Colombia,<sup>9</sup> while *K. pneumoniae* GB015 belonged to ST442. ST442 and ST11, the most common carbapenem-resistant clones in China, are considered to be the progenitors of *K. pneumoniae* ST258, which is widespread worldwide as the most successful multidrug-resistant clone of *K. pneumoniae*.<sup>10</sup>

The serotype of *E. coli* GB110 was identified as O38:H26, which can cause diarrhea in humans.<sup>11</sup> The *E. coli* heat-stable enterotoxin-1 (EAST-1)-encoding gene *astA* was found in *E.*

*coli* GB110, which is associated with human diseases. These results indicate the risk that eating contaminated fruits could lead to diarrheal diseases.

The *E. coli* GB110 was found to harbor ten resistance genes, including *mcr-1* and aminoglycoside resistance genes, *aadA2* and *aadA1*, while *K. pneumoniae* GB015 was found to possess nine resistance genes, including *mcr-1*, *bla*<sub>SHV-110</sub>, *qnrS1* and *fosA6* (Table 1).

## Conclusion

In our study, we reported the detection of *mcr-1* from market retail fruits, which is significant since, unlike meat and vegetables, fruits are usually consumed without cooking or processing, making them a potentially high-risk source of *mcr-1* acquisition and infection in humans. This work sheds light on the urgent need for continued surveillance and prompt intervention in China to guard against the worldwide distribution of *mcr-1*, which threatens the use of colistin as a last resort antibiotic.

## Acknowledgments

This work was supported by the Science and Technology Research Project of Henan Province (grant 182102310553) and the National Natural Science Foundation of China (grant numbers 81722030 and 81830103).

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Mendes AC, Novais Â, Campos J, et al. *mcr-1* in carbapenemase-producing *Klebsiella pneumoniae* with hospitalized patients, Portugal, 2016-2017. *Emerg Infect Dis*. 2018;24(4):762-766.
- Liu YY, Wang Y, Walsh TR, et al. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis*. 2016;16(2):161-168.
- Wang X, Wang Y, Zhou Y, et al. Emergence of a novel mobile colistin resistance gene, *mcr-8*, in NDM-producing *Klebsiella pneumoniae*. *Emerg Microbes Infect*. 2018;7(1):122.
- Wang R, van Dorp L, Shaw LP, et al. The global distribution and spread of the mobilized colistin resistance gene *mcr-1*. *Nat Commun*. 2018;9(1):1179.
- Zheng B, Huang C, Xu H, et al. Occurrence and Genomic Characterization of ESBL-Producing, MCR-1-Harboring *Escherichia coli* in Farming Soil. *Front Microbiol*. 2017;8:2510.
- Zhong LL, Zhang YF, Doi Y, et al. Coproduction of MCR-1 and NDM-1 by colistin-resistant *Escherichia coli* isolated from a healthy individual. *Antimicrob Agents Chemother*. 2016;61(1).
- Nurk S, Bankevich A, Antipov D, et al. Assembling single-cell genomes and mini-metagenomes from chimeric MDA products. *J Comput Biol*. 2013;20(10):714-737.
- Snesrud E, McGann P, Chandler M. The birth and demise of the IS*Apl1*-*mcr-1*-IS*Apl1* composite transposon: the vehicle for transferable colistin resistance. *MBio*. 2018;9(1):e02381-e02317.
- Castellanos LR, Donado-Godoy P, León M, et al. High heterogeneity of *Escherichia coli* sequence types harbouring ESBL/AmpC genes on IncI1 plasmids in the Colombian Poultry Chain. *PLoS One*. 2017;12(1):e0170777.
- Pitout JD, Nordmann P, Poirel L. Carbapenemase-producing *Klebsiella pneumoniae*, a key pathogen set for global nosocomial dominance. *Antimicrob Agents Chemother*. 2015;59(10):5873-5884.
- Buvs G, Lauwers S, Piérard D. Prevalence of subtilase cytotoxin in verocytotoxin-producing *Escherichia coli* isolated from humans and raw meats in Belgium. *Eur J Clin Microbiol Infect Dis*. 2010;29(11):1395-1399.

## Supplementary material

**Table S1** Primers and PCR conditions used in this study

Purpose	Primer	Nucleotide sequence (5'→3')	Amplicon size (bp)	Annealing temperature (°C)
Amplification of <i>mcr-1</i>	MCR-1F	ATCAGCCAAACCTATCCCATCG	1,257	55
	MCR-1R	GCAGACGCACAGCAATGCCTAT		
Amplification of <i>mcr-2</i>	MCR-2F	GCGATGGCGGTCTATCCTGTAT	378	55
	MCR-2R	TGCGATGACATGGGGTGTGAGC		
Amplification of <i>mcr-3</i>	MCR-3F	TATGGGTTACTATTGCTGG	814	55
	MCR-3R	CTACCCTGATGCTCATCG		
Amplification of <i>mcr-4</i>	MCR-4F	GTCATAGTGGTATAAAAAGTACAG	669	55
	MCR-4R	CCACCGTCTATCAGAGCCAAC		
Amplification of <i>mcr-5</i>	MCR-5F	GCGGTTGTCTGCATTTATCAC	1,042	50
	MCR-5R	CTTTGAAAACCTGTCTTCGGCA		
Amplification of <i>mcr-6</i>	MCR-6F	GTCCGGTCAATCCCTATCTGT	556	55
	MCR-6R	ATCACGGGATTGACATAGCTAC		
Amplification of <i>mcr-7</i>	MCR-7F	TGCTCAAGCCCTTCTTTTCGT	892	55
	MCR-7R	TTCATCTGCGCCACCTCGT		
Amplification of <i>mcr-8</i>	MCR-8F	AACCGCCAGAGCACAGAATT	667	60
	MCR-8R	TTCCCCAGCGATTCTCCAT		

### Infection and Drug Resistance

#### Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic

resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>

Dovepress