

Received:
31 January 2019
Accepted:
31 January 2019

Cite as: Mai Yokoji-Takeuchi,
Koichi Tabeta,
Naoki Takahashi,
Kei Arimatsu,
Haruna Miyazawa,
Yumi Matsuda-Matsukawa,
Keisuke Sato, Miki Yamada,
Kazuhisa Yamazaki.
Corrigendum to “Indirect
regulation of PCSK9 gene in
inflammatory response by
Porphyromonas gingivalis
infection” [Heliyon 5 (1)
(January 2019) e011111].
Heliyon 5 (2019) e01210.
doi: [10.1016/j.heliyon.2019.e01210](https://doi.org/10.1016/j.heliyon.2019.e01210)



Corrigendum

Corrigendum to “Indirect regulation of PCSK9 gene in inflammatory response by *Porphyromonas gingivalis* infection” [Heliyon 5 (1) (January 2019) e01111]

Mai Yokoji-Takeuchi^{a,b}, Koichi Tabeta^{a,*}, Naoki Takahashi^a, Kei Arimatsu^a,
Haruna Miyazawa^a, Yumi Matsuda-Matsukawa^a, Keisuke Sato^a, Miki Yamada^{a,b},
Kazuhisa Yamazaki^{b,**}

^a Division of Periodontology, Department of Oral Biological Science, Niigata University Graduate School of Medical and Dental Sciences, Niigata, Japan

^b Research Unit for Oral-Systemic Connection, Division of Oral Science for Health Promotion, Niigata University Graduate School of Medical and Dental Sciences, Niigata, Japan

DOI of original article: [10.1016/j.heliyon.2018.e01111](https://doi.org/10.1016/j.heliyon.2018.e01111)

* Corresponding author.

** Corresponding author.

E-mail addresses: koichi@dent.niigata-u.ac.jp (K. Tabeta), kaz@dent.niigata-u.ac.jp (K. Yamazaki).

In the original published version of this article, an error was present in figure 3. During the typesetting stage, an incorrect version of figure 3 was uploaded which was missing several figure panels which had been present at acceptance. The authors apologize for this mistake. A corrected version of Figure 3 is displayed below. Both the HTML and PDF versions of the article have been updated to correct the error.




Fig. 3. Changes in serum lipid levels following the administration of *P. gingivalis* and nucleic acid antigens: (A) total cholesterol (TC), (B) low-density lipoprotein cholesterol (LDL-C), (C) high-density lipoprotein cholesterol (HDL-C), and (D) triglycerides (TG). $N = 5$ mice per group. Results are expressed as means \pm SEM. *, $P < 0.05$; **, $P < 0.01$. Analysis of serum cholesterol levels.: Correlations between the serum lipids and serum PCSK9 level or hepatic $TNF-\alpha$ gene expression in wild-type mice. (E) LDL-C versus PCSK9, (F) HDL-C levels versus PCSK9, (G) TC versus PCSK9, (H) LDL-C versus *Tnfα*, (I) HDL-C versus *Tnfα*, and (J) TC versus *Tnfα*. The ratio of each LDL-C fraction level to the total: (K) large ($\phi 28.6$ nm), (L) medium ($\phi 25.5$ nm), (M) small ($\phi 23.0$ nm), or (N) very small ($\phi 16.7-20.7$ nm). $N = 5$ mice per group. Results are expressed as means \pm SEM. *, $P < 0.05$; **, $P < 0.01$.