


TOPICAL REVIEW

Mechanoadaptation: articular cartilage through thick and thin

Tonia L. Vincent  and Angus K. T. Wann 

Arthritis Research UK Centre for OA Pathogenesis, Kennedy Institute of Rheumatology, University of Oxford, Oxford, UK

Edited by: Ole Petersen & Maike Glitsch



Tonia L. Vincent is an academic rheumatologist and Director of the Centre for OA Pathogenesis in Oxford. She works on the molecular response of cartilage to injury and how this relates to the development of osteoarthritis. She did her PhD with Professor Jeremy Saklatvala and has been supported by Wellcome Trust and Arthritis Research UK fellowships. She continues to be clinically active. Following PhD training under Professor J. Rodney Levick at St George's, University of London, studying the mechanobiology of the synovial joint, **Angus Wann** moved to Bioengineering at Queen Mary University of London to work with Professor Martin Knight and became fascinated by the primary cilium. He received a fellowship jointly funded by the Kennedy Trust for Rheumatology Research (KTRR) and the Arthritis Research UK Centre for OA Pathogenesis and now runs his own lab focusing on pathophysiological roles for ciliary proteins in musculoskeletal disease.



understood despite the identification of multiple mechanosensing mechanisms in cartilage. Moreover, there has been a tendency to equate cartilage loss with osteoarthritic degeneration. Here, we review the historic literature and clarify the structural, metabolic and clinical features that clearly distinguish cartilage loss due to disuse atrophy and those due to osteoarthritis. We speculate on the molecular sensing pathways in cartilage that may be responsible for cartilage mechanoadaptation.

(Received 4 April 2018; accepted after revision 1 May 2018; first published online 19 June 2018)

Corresponding author T. L. Vincent: Arthritis Research UK Centre for OA Pathogenesis, Kennedy Institute of Rheumatology, University of Oxford, Roosevelt Drive, Oxford OX3 7FY, UK. Email: tonia.vincent@kennedy.ox.ac.uk


Abstract figure legend Cartilage atrophy is distinct to osteoarthritic damage. In normal full-thickness cartilage (left panel) chondrocytes are diffusely distributed through a proteoglycan-rich (light pink) articular cartilage matrix separated from calcified cartilage beneath by a tidemark. Cartilage atrophy (middle panel) is associated with thinning of the cartilage and reduced proteoglycan content but the articular surface is intact. In contrast, osteoarthritic tissue (right hand panel), whilst also associated with thinning, has marked surface damage and patchy proteoglycan staining.

The mechanosensitivity of musculoskeletal tissues is extremely well documented but for the most part this has focused on muscle and bone rather than articular cartilage (Ziaaldini *et al.* 2017; Galea *et al.* 2017; Uda *et al.* 2017). Part of the reason for this lies in the fact that the cartilage is not easily visualised; its thickness is inferred from the gap between the ends of the bone (the joint space) on a plain radiograph. Cartilage has only relatively recently been directly visualised by magnetic resonance imaging (MRI).

Articular cartilage is adapted to withstand mechanical stress

Articular cartilage develops and operates under very high compressive and shear stresses throughout its lifetime, and has only modest purported reparative capability, particularly in the elderly. Cartilage is an extracellular matrix-rich tissue where only 5–10% of the tissue volume is cell (Hunziker *et al.* 2002). The principal fibrillar component is type II collagen, forming staggered arrangements of triple helical molecules that are highly cross-linked and organised in such a way that they lie parallel with the surface layer superficially but perpendicular to it in the deep regions (Fig. 1). The complex organisation of primary and secondary structures provides cartilage with high tensile strength and the orientation of fibres allows mechanical loads to be dissipated over the surface, so as to minimise impact on the focal 'load-bearing' regions of the joint. The tissue holds water by means of fixed negative charges offered by the proteoglycan aggrecan, itself aggregated on hyaluronan chains in the tissue (Dudhia, 2005). Aggrecan has attachment sites for many chondroitin sulfate and keratan sulfate glycosaminoglycan (GAG) chains that lie within the territorial and inter-territorial parts of the matrix. The hydration allowed by the fixed

negative charge of these chains provides the tissue with a swelling pressure enabling absorption of, and resistance to, mechanical shock. The chondrocyte is separated from the territorial matrix by a specialised pericellular matrix (PCM). Together, the chondrocyte and the PCM form the 'chondron'. Chondrons are themselves orientated



The Journal of
Physiology

Figure 1. Schematic diagram of articular cartilage

Illustration of tertiary structure of fibrillar collagens in articular cartilage showing parallel arrangement in superficial layer and perpendicular orientation in deeper layers. Arrows show direction of mechanical load and how it is dissipated across the articulating surface.

relative to local mechanical strain, exhibiting a polarised morphology that varies with tissue depth (Poole, 1997).

Loss of proteoglycans by cleavage of the core molecule by aggrecanases occurs in osteoarthritis (OA) and is regarded as a key pathological feature of disease (Glasson *et al.* 2005). Tissue catabolism is also mediated by other members of the matrix metalloproteinase (MMP) family. MMP13, one of the mammalian collagenases, has a proven role in osteoarthritis (Mitchell *et al.* 1996), whilst MMP3 (stromelysin) does not appear to drive disease and may be more important in normal tissue turnover (Clements *et al.* 2003; Troeberg & Nagase, 2012).

Articular chondrocytes are highly mechanosensitive

For articular cartilage to be able to respond to its mechanical environment, it must be able to sense a change in the magnitude and type of mechanical load and respond to this change. Historically, the chondrocyte was regarded as a largely quiescent cell, one that rarely proliferated and was responsible for slow turnover of the matrix. Early studies by Mankin showed that proteoglycan synthesis was moderately active (with a half-life of around 8 days) (Mankin & Lippiello, 1969), and synthetic activity increased substantially following injury (Meachim, 1963).

Multiple mechanisms by which chondrocytes sense mechanical load have been identified. Cellular mechanosensors include cell surface integrins, stretch and cation-sensitive ion channels (Ross *et al.* 2013; Lee *et al.* 2014; O'Connor *et al.* 2014; see also review by Drexler *et al.* 2014). The chondrocyte primary cilium has also been implicated in mechanotransduction, as it has in many other cell types (Wann *et al.* 2012). Beyond the chondrocyte another important mechanism is release of sequestered heparan sulfate-bound molecules of the PCM in response to compression or cutting injury (Vincent *et al.* 2002, 2004; Vincent, 2013) (Fig. 2). The best described of these is fibroblast growth factor 2 (FGF2), which is chondroprotective *in vivo* in models of osteoarthritis (Chia *et al.* 2009). Other molecules include connective tissue growth factor (CTGF) which is covalently bound to latent transforming growth factor beta (TGF β). Release of the complex upon cutting injury results in activation of TGF β at the cell surface in a CTGF-dependent manner (Tang *et al.* 2018). Activation of a number of intracellular signalling pathways occurs rapidly upon injury including TGF β -activated kinase 1 (TAK1), β -catenin, Src, Smads, mitogen activated kinases (ERK, p38 and JNK) and nuclear factor κ B (NF κ B) (Vincent *et al.* 2002; Clements *et al.* 2003, 2011; Gruber *et al.* 2004; Dell'Accio *et al.* 2006, 2008; Troeberg & Nagase, 2012; Watt *et al.* 2013; Ismail *et al.* 2017). The receptor proximal mechanism of activation for most of these remains unknown.




Figure 2. Schematic diagram of a chondron

Illustration of the chondrocyte sitting within its pericellular matrix (PCM), together forming the chondron. Note absence of fibrillar collagens and aggrecan in the PCM, but abundance in the adjacent territorial matrix. Multiple heparan sulfate-bound growth factors are bound within the PCM and released upon tissue compression or injury.

Interestingly, both experimental data and *in silico* models show that the forces experienced by an individual chondrocyte are largely the same irrespective of the mass of the animal (Simon, 1970). This is most likely explained by 'isometric' scaling of the skeleton, where the joint surface area increases in proportion to the mass of the animal. Cartilage thickness does increase with animal size, but exhibits allometric rather than isometric scaling. Indeed, it has been suggested that there is an optimal maximum thickness of cartilage, beyond which perfusion of the chondrocytes deep in the tissue would be compromised (Malda *et al.* 2013). One might thus hypothesise the existence of an evolutionary 'mechanostat' for chondrocytes that has been tuned to an optimal mechanical input. The mechanical input itself, being related to the properties (stiffness) of the surrounding matrix.

Evidence for disuse cartilage atrophy in humans

In 1974, Winford Pool reviewed 200 cases of flaccid paralysis of the lower extremities. Twenty-five cases had demonstrable narrowing of the hip joint space by plain radiograph of at least 50% (Pool, 1974). His conclusions at the time were that flaccid paralysis caused significant reduction of all types of load on the joint (compared with spastic paralysis where there is increased static load by action of muscle). He proposed that this reduced the

Table 1. Causes of thinning of the articular cartilage

Cause	Mechanism	Reversible?
Osteoarthritis	Proteolytic degradation of matrix	No, except by regeneration
Atrophy	Reduced synthesis of proteoglycan; a mechanoadaptive response to reduced load	Yes, weeks–months
Cartilage compression	Water extrusion from matrix due to compression	Yes, minutes–hours
Apparent reduction in joint space ('pseudo thinning')	Migration of tidemark or meniscal extrusion	Unknown

nutrient supply to the cartilage, which is dependent upon pumping action of load-bearing activity. His findings were supported by those of Anderson and Breidahl who performed a similar study in 71 patients with either flaccid or spastic paralysis due to spinal cord damage. Their results confirmed the strong association between paralysis and joint space narrowing at the hip which was more common in flaccid rather than spastic paralysis (Anderson & Breidahl, 1981). Similar results were obtained when lower limb amputees were examined. Importantly, in this paper they also comment that features of OA were never seen on the amputated side (Benichou & Wirotius, 1982). This is one of the earliest clarifications regarding cartilage atrophy and OA. Despite both conditions leading to thinning of articular cartilage (see Table 1), OA develops as a result of increased mechanical load whereas atrophy results from reduced load.

The subsequent development in the early 1990s of non-invasive methodologies to assess the volume and proteoglycan content of articular cartilage by MRI allowed much more sensitive assessments of change with disuse and disease (Recht *et al.* 1993; Eckstein *et al.* 1997; Haubner *et al.* 1997; Löscher *et al.* 1997). A small number of cross-sectional and longitudinal studies have been carried out in spinal cord injury patients since then. Cartilage volume loss was seen to decrease by around 6% in the first 6 months from baseline (baseline was on average 4 weeks post-spinal injury), resulting in a loss of around 25% of the medial tibial thickness by 24 months after injury when compared with normal controls (Vanwanseele *et al.* 2002, 2003). In individuals who are temporarily immobilised, through ankle fracture, a 6.6% loss of cartilage thickness was measured in the medial tibia over 7 weeks following fracture. Sensitivity of these techniques continues to increase and it is now possible to detect very small strain levels (around 3%) in the tissue that occur on normal weight-bearing activity (Eckstein *et al.* 2000; Sutter *et al.* 2015). No human study to our knowledge has demonstrated recovery of cartilage volume after remobilisation but importantly, moderate programmed exercise was shown to have a positive effect on increasing cartilage GAG content (assessed by delayed

gadolinium-enhanced MRI of cartilage, dGEMRIC) in individuals at risk of OA due to previous meniscal injury (Roos & Dahlberg, 2005).

Disuse atrophy in large animal models

The greatest body of literature on cartilage atrophy stems from *in vivo* studies, mainly carried out in large animals (dog and rabbit). These studies have done much to provide definitive evidence for the importance of mechanics to joint homeostasis and disuse atrophy, greatly enhancing our understanding of the metabolic processes that lead to the condition. In 1979, Pamoski *et al.* produced a series of interesting studies in which they immobilised healthy adult dogs using a light cast that fixed the knee joint at 90°. The cast was strapped to the trunk preventing weight bearing or active flexion/extension. These studies revealed that 6 days of immobilisation was sufficient to observe a reduction in proteoglycan staining in the tissue and by 8 weeks there was a 30–50% reduction in cartilage thickness with almost complete loss of proteoglycan. Although the authors did not describe whether the changes occurred in the calcified or non-calcified cartilage, others have suggested that the calcified cartilage layer (below the tidemark) is more sensitive to atrophic changes (Kiviranta *et al.* 1987). Interestingly, despite loss of proteoglycan there was an increase in the water content in the tissue, a feature that was confirmed by other similar studies (Pamoski *et al.* 1979; Behrens *et al.* 1989), and a change in the proteoglycan produced by the tissue such that it failed to aggregate with hyaluronan *in vitro*. This aggregation defect suggests a change in the GAG rather than the hyaluronan itself, although a reduction in hyaluronan tissue content has also been described with disuse (Haapala *et al.* 1996). Hyaluronan synthesis itself was later shown to be mechanosensitive in the synovial joint (Ingram *et al.* 2009). The hyaluronan aggregation defect was reversed upon voluntary remobilisation (Pamoski *et al.* 1979) and may be related to restoration of normal hyaluronan synthesis (Müller *et al.* 1994). Recovery from atrophy has been documented consistently in other studies, although

is affected by the severity of atrophy attained and may be incomplete when immature joints are immobilised (Palmoski & Brandt, 1981; Behrens *et al.* 1989; Kiviranta *et al.* 1992).

In a follow-up study in 1981, Palmoski & Brandt showed forced treadmill exercise prevented recovery of the tissue after immobilisation. Newly synthesised proteoglycans measured by ³⁵S incorporation demonstrated that there was a decrease in proteoglycan synthesis upon disuse, consistent with later studies examining aggrecan gene expression (Djurasovic *et al.* 1998). Proteoglycan synthesis increased upon remobilisation in both the free and forced remobilised groups. The authors concluded that during forced remobilisation there must also be increased proteolytic activity to account for why proteoglycan failed to recover despite increased synthesis (Palmoski & Brandt, 1981). Whether increased proteolysis contributes to atrophic changes in the cartilage has been an area of speculation but to date there is little evidence to support this despite some efforts (Grumbles *et al.* 1995; Haapala *et al.* 2001).

The focus on proteoglycan metabolism in these studies is justified. There is significantly less information available to suggest a change in the collagenous matrix in atrophic cartilage apart from some modest reduction in collagen crosslinking which recovers after remobilisation (Haapala *et al.* 1999). This would be consistent with the known stability of fibrillar collagens and evidence for minimal turnover in human cartilage over the lifetime (Heinemeier *et al.* 2016). The biomechanical consequences of matrix imbalance in atrophic tissue have been studied, *ex vivo*, by mechanical testing. These studies collectively show some site-specific changes in shear and compressive properties of the cartilage, although they show some inconsistencies (Jurvelin *et al.* 1986*b*; Leroux *et al.* 2001).

Disuse atrophy in small animal models

Generally speaking all types of joint immobilisation lead to atrophy although differences in how immobilisation is achieved, e.g. by rigid external fixation or cast immobilisation, and at which sites the effects are measured, does affect the type of outcome measures (Säämänen *et al.* 1990) and magnitude of change (Behrens *et al.* 1989). Two studies performed in the last decade have studied immobilisation in rodents. In the first, immobilisation was achieved by spinal cord injury in which the animals drag the affected limb. The study revealed changes in the thickness of cartilage across different regions of the joint at baseline with predicted weight-bearing regions having higher recorded thickness. Upon spinal cord injury a 30% decrease in cartilage thickness was measured in off-loaded areas (Moriyama

et al. 2008). The second study compared rigid (by externally fixed Kirschner wires) and 'tail suspension' immobilisation in mice. Both led to a reduction in cartilage thickness, largely due to a decrease in the calcified cartilage, and were associated with loss of proteoglycan in the non-calcified layer particularly. Atrophy was slightly enhanced in the rigid fixation group, presumably because the joint was not subjected to movements associated with muscle activity (which are present during tail suspension) (Nomura *et al.* 2017).

Increased thickness of cartilage upon exercise

If disuse leads to cartilage thinning then it follows that exercise training should result in thicker cartilage, so-called cartilage hypertrophy. This has been shown in a number of large animal studies and is reflected by changes in the same biochemical metrics as measured in atrophy. Increased synthesis of the guanidine-extractable proteoglycan was documented in exercised rabbits (Tammi *et al.* 1983; Säämänen *et al.* 1988) and in dogs, there was an increase in extractable GAGs that were less able to aggregate with hyaluronan *in vitro*. These GAGs also had increased ratios of chondroitin-6 to chondroitin-4 sulfate (Tammi *et al.* 1983; Säämänen *et al.* 1989). Treadmill running of 4 km a day for 15 weeks in dogs was associated with an increase in cartilage thickness (11% in loaded areas) and mechanical stiffness (6%) (Jurvelin *et al.* 1986*a*). Treadmill training for up to 40 weeks at 5 h/week was associated with an increase in the non-calcified cartilage thickness by 19–23% and glycosaminoglycans by 28% (Kiviranta *et al.* 1988).

Exercise dosage has been examined in rats running on a treadmill. Speeds of 15 m/min with no incline, and 19 m/min with a 5° incline for 5 h per week for 6 weeks were associated with increased cartilage thickness and proteoglycan synthesis compared with non-exercised control animals (Ni *et al.* 2013). Similar to studies of atrophy there are site-specific changes in cartilage that reflect the type of exercise undertaken, e.g. downhill treadmill running (Hamann *et al.* 2014).

In contrast to exercise-induced adaptation of muscle and bone in adult human joints, there is much less evidence for cartilage thickness adaptation. While associations exist between rates of cartilage accrual and physical activity during childhood (Jones *et al.* 2003), studies comparing athletes and inactive cohorts found the variance in cartilage thickness was not clearly attributable to activity levels (Eckstein *et al.* 2002). However, an association between exercise and joint size (cartilage surface area) did exist, consistent with the isometric relationship described for body mass (or by extrapolation, load) and surface area in other mammals (Malda *et al.* 2013).

Table 2. Clinical distinguishing features between cartilage atrophy and osteoarthritis

Clinical features	Cartilage atrophy	Osteoarthritis
Joint space narrowing (JSN)	Yes	Yes
History of immobilisation	Yes	No, increased load
Pain	No	Yes
Synovial hypertrophy/inflammation	No	Yes
Bone changes	Local osteoporosis	Subchondral bone sclerosis; osteophytes

Molecular mechanisms underlying mechanoadaptation in cartilage

The evidence thus far points to cartilage being highly 'mechanoadaptable' in contrast to its somewhat 'inert' reputation in the post-developmental context. Early speculation suggested that this was due to altered nutrition of the cartilage due to its dependence on mechanical load to deliver nutrients and oxygen to the chondrocytes. However, the relatively recent appreciation that chondrocytes can sense their mechanical environment and respond differently to distinct mechanical cues, amplitudes and frequencies, suggests that these mechano-sensing mechanisms will turn out to be responsible for driving mechanoadaptation. Although, to our knowledge, this has not been directly addressed, careful scrutiny of the literature does provide us with potential clues as to the pathways that may be responsible. For instance, increased cartilage thickness has been described in mice that have been subjected to a transient burst of Wnt signalling (by tamoxifen-induced transgene activation) (Yuasa *et al.* 2009). In this case there was an immediate loss of proteoglycan preceding the 'rebound' hypertrophy of cartilage. Wnt signalling occurs rapidly upon mechanical injury of cartilage *in vitro* and a number of Wnt ligands are matrix bound and present in cartilage (Dell'Accio *et al.* 2006, 2008; Berendsen *et al.* 2011). Wnts are also involved in joint cavitation during development, a process that is critically dependent upon mechanical force by associated muscle action (Kahn *et al.* 2009; Shea *et al.* 2015).

Vincent has also observed an increase in cartilage thickness associated with constitutive activation of Smad2 signalling suggesting that TGF β family members may drive hypertrophic cartilage responses (Tang *et al.* 2018). As TGF β is one of the growth factors present in the pericellular matrix and decreases in cartilage with age (Madej *et al.* 2016; van Caam *et al.* 2016), it is tempting to speculate that its release upon tissue compression could be relevant to reduced mechanoadaptation over the life course, a process described in other skeletal tissues (Greig *et al.* 2011; Brook *et al.* 2016). Another molecule that is involved in chondrocyte mechanosensing is FGF2. FGF2 is an important mechanosensitive mediator of joint cavitation during development (Kavanagh *et al.* 2006; Shea *et al.* 2015) and is also released by cartilage

compression (Vincent *et al.* 2007). However, we found no evidence of a change in cartilage thickness in male or female naive FGF2 knockout mice (Chia *et al.* 2009). Activation of cell surface mechanoreceptors piezo 1 and Trpv4 in chondrocytes leads to anabolic responses *in vitro* and chondroprotection *in vivo*. Moreover, their activity is associated with proteoglycan synthesis, making these interesting molecular candidates for further investigation (Clark *et al.* 2010; Lee *et al.* 2014; O'Connor *et al.* 2014, 2016).

What is the relationship between atrophy and osteoarthritis?

While mechanoadaptation of cartilage has become a somewhat neglected area of study, interest in molecular pathogenesis of osteoarthritis has greatly increased (see Tables 2 and 3 for comparisons). Cartilage loss in OA has completely different molecular and chemical drivers; matrix catabolism is highly protease dependent but is associated with a strong anabolic reaction suggesting an unsuccessful repair response. Increased mechanical load is a critical aetiological factor that is strongly supported by epidemiological studies (Brandt *et al.* 2009), and by showing that regulation of proteases *in vivo* upon induction of disease (by surgical joint destabilisation) is abrogated by joint immobilisation (Burleigh *et al.* 2012). OA is also distinguished from atrophy by loss of integrity of the tissue starting with changes at the articular surface, which spread throughout the tissue as the disease progresses and are accompanied by aberrant bone remodelling. From a molecular and structural point of view, the process of atrophy and OA could not be further apart. The only feature they share is thinning of the cartilage and therefore an apparent decrease in the joint space on plain radiograph.

But is there a relationship between atrophy and OA? And why does exercise sometimes result in thickening of the cartilage and sometimes OA? The answer likely relates back to the defined 'mechanostat' within the tissue. The authors speculate that if mechanical stimuli exceed a given threshold, or fall outside certain parameters, they are perceived as injurious and trigger pathways associated with rapid matrix remodelling and attempted repair. TAK1

Table 3. Biochemical changes in atrophic and osteoarthritic cartilage

Tissue characteristic	Cartilage atrophy	Recovery with <i>ad libitum</i> remobilisation?	Osteoarthritis
Proteoglycan synthesis	Decreased	Yes	Increased
Total tissue proteoglycan	Decreased	Yes	Decreased
Evidence of proteolysis?	No	N/A	Yes, of proteoglycan and collagen
Proteoglycan aggregation defect	Yes	Yes	Probably not
Chondroitin-6 sulfate	Not changed	Reduced relative to chondroitin-4-sulfate (Säämänen <i>et al.</i> 1990)	Increased (Ratcliffe <i>et al.</i> 1993)
Water content	Increased	Yes	Increased early in disease
Collagen cross-linking	Decreased	Yes	Unknown
Appearance	Smooth, shiny, healthy	N/A	Roughened, dull

activation appears to be critical in this response. TAK1 is a pivotal upstream regulator of inflammatory signalling, is induced by cartilage trauma and drives inflammatory gene regulation and matrix proteolysis (Ismail *et al.* 2015, 2016). The upstream activator of TAK1 remains elusive and does not appear to be mediated by a soluble factor (Ismail *et al.* 2017). Exceeding the injury threshold could occur either when the mechanical load is increased, or when the load is moderate but the joint has lost its mechanoprotective mechanisms. Examples of the latter include loss of stability of the joint and weakening of the cartilage matrix by atrophy. This would provide a plausible explanation for why *ad libitum* remobilisation leads to recovery after disuse but forced remobilisation leads to OA-like changes in the cartilage (Palmoski & Brandt, 1981). Similarly in joints weakened by papain injection, moderate exercise leads to rapid degeneration of the tissue (Siebelt *et al.* 2014). A systematic review of experimental daily exercise in animals also suggests a non-linear dose response with medium loads appearing good for cartilage but high loads precipitating degeneration (Bricca *et al.* 2017).

We presume that the ‘injury’ threshold changes as the cartilage mechanoadapt and will be different for each individual according to genetics, the mechanical integrity of the tissue, the amount of load usually experienced by the joint and the ‘pre-tuned’ sensing mechanisms of their chondrocytes. Nonetheless, it is highly likely that mechanical training (graded exercise rather than overnight athleticism) will better protect the cartilage from load-induced activation of matrix remodelling and subsequent OA. Aiming for thicker cartilage is probably a good start. Whether it is possible to optimise cartilage thickness during skeletal development to prevent OA later in life, remains an important unanswered question.

Conclusions

Mechanoadaptation in cartilage is rapid and reversible, and potentially of a similar scale to that seen in muscle and bone. Comprehensive studies of cartilage atrophy *in vivo* have been carried out although largely during the ‘pre-molecular’ era and prior to the discovery of direct chondrocyte mechanosensing mechanisms. Advances in newly available glycobiology and proteomic techniques, in combination with genetic modification in rodents, will add considerable value to future studies. Cartilage atrophy is readily distinguished from osteoarthritis at the clinical, tissue and molecular level but cannot be discerned on a plain radiograph except by noting the absence of bone remodelling. As joint space narrowing is typically used to diagnose OA, it is important to consider atrophy as a differential diagnosis. Harnessing the molecules that drive mechanoadaptation in articular cartilage may provide novel strategies to prevent or treat OA.

References

- Anderson J & Bredahl P (1981). Cartilage atrophy following spinal cord damage. *Australas Radiol* **25**, 98–103.
- Behrens F, Kraft EL & Oegema TR (1989). Biochemical changes in articular cartilage after joint immobilization by casting or external fixation. *J Orthop Res* **7**, 335–343.
- Benichou C & Wirocius JM (1982). Articular cartilage atrophy in lower limb amputees. *Arthritis Rheum* **25**, 80–82.
- Berendsen AD, Fisher LW, Kilts TM, Owens RT, Robey PG, Gutkind JS & Young MF (2011). Modulation of canonical Wnt signaling by the extracellular matrix component biglycan. *Proc Natl Acad Sci U S A* **108**, 17022–17027.
- Brandt KD, Dieppe P & Radin EL (2009). Commentary: is it useful to subset “primary” osteoarthritis? A critique based

- on evidence regarding the etiopathogenesis of osteoarthritis. *Semin Arthritis Rheum* **39**, 81–95.
- Bricca A, Juhl CB, Grodzinsky AJ & Roos EM (2017). Impact of a daily exercise dose on knee joint cartilage – a systematic review and meta-analysis of randomized controlled trials in healthy animals. *Osteoarthritis Cartilage* **25**, 1223–1237.
- Brook MS, Wilkinson DJ, Phillips BE, Perez-Schindler J, Philp A, Smith K & Atherton PJ (2016). Skeletal muscle homeostasis and plasticity in youth and ageing: impact of nutrition and exercise. *Acta Physiol (Oxf)* **216**, 15–41.
- Burleigh A, Chanalaris A, Gardiner MD, Driscoll C, Boruc O, Saklatvala J & Vincent TL (2012). Joint immobilization prevents murine osteoarthritis and reveals the highly mechanosensitive nature of protease expression in vivo. *Arthritis Rheum* **64**, 2278–2288.
- Chia S-L, Sawaji Y, Burleigh A, McLean C, Inglis J, Saklatvala J & Vincent T (2009). Fibroblast growth factor 2 is an intrinsic chondroprotective agent that suppresses ADAMTS-5 and delays cartilage degradation in murine osteoarthritis. *Arthritis Rheum* **60**, 2019–2027.
- Clark AL, Votta BJ, Kumar S, Liedtke W & Guilak F (2010). Chondroprotective role of the osmotically sensitive ion channel transient receptor potential vanilloid 4: age- and sex-dependent progression of osteoarthritis in *Trpv4*-deficient mice. *Arthritis Rheum* **62**, 2973–2983.
- Clements KM, Flannelly JK, Tart J, Brockbank SMV, Wardale J, Freeth J, Parker AE & Newham P (2011). Matrix metalloproteinase 17 is necessary for cartilage aggrecan degradation in an inflammatory environment. *Ann Rheum Dis* **70**, 683–689.
- Clements KM, Price JS, Chambers MG, Visco DM, Poole AR & Mason RM (2003). Gene deletion of either interleukin-1 β , interleukin-1 β -converting enzyme, inducible nitric oxide synthase, or stromelysin 1 accelerates the development of knee osteoarthritis in mice after surgical transection of the medial collateral ligament and partial medial meniscectomy. *Arthritis Rheum* **48**, 3452–3463.
- Dell'Accio F, De Bari C, Eltawil NM, Vanhummelen P & Pitzalis C (2008). Identification of the molecular response of articular cartilage to injury, by microarray screening: Wnt-16 expression and signaling after injury and in osteoarthritis. *Arthritis Rheum* **58**, 1410–1421.
- Dell'Accio F, De Bari C, El Tawil NMF, Barone F, Mitsiadis TA, O'Dowd J & Pitzalis C (2006). Activation of WNT and BMP signaling in adult human articular cartilage following mechanical injury. *Arthritis Res Ther* **8**, R139.
- Djurasovic M, Aldridge JW, Grumbles R, Rosenwasser MP, Howell D & Ratcliffe A (1998). Knee joint immobilization decreases aggrecan gene expression in the meniscus. *Am J Sports Med* **26**, 460–466.
- Drexler S, Wann A & Vincent TL (2014). Are cellular mechanosensors potential therapeutic targets in osteoarthritis? *Int J Clin Rheumatol* **9**, 155–167.
- Dudhia J (2005). Aggrecan, aging and assembly in articular cartilage. *Cell Mol Life Sci* **62**, 2241–2256.
- Eckstein F, Adam C, Sittke H, Becker C, Milz S, Schulte E, Reiser M & Putz R (1997). Non-invasive determination of cartilage thickness throughout joint surfaces using magnetic resonance imaging. *J Biomech* **30**, 285–289.
- Eckstein F, Faber S, Mühlbauer R, Hohe J, Englmeier KH, Reiser M & Putz R (2002). Functional adaptation of human joints to mechanical stimuli. *Osteoarthritis Cartilage* **10**, 44–50.
- Eckstein F, Lemberger B, Stammberger T, Englmeier KH & Reiser M (2000). Patellar cartilage deformation in vivo after static versus dynamic loading. *J Biomech* **33**, 819–825.
- Galea GL, Lanyon LE & Price JS (2017). Sclerostin's role in bone's adaptive response to mechanical loading. *Bone* **96**, 38–44.
- Glasson SS, Askew R, Sheppard B, Carito B, Blanchet T, Ma H-L, Flannery CR, Peluso D, Kanki K, Yang Z, Majumdar MK & Morris EA (2005). Deletion of active ADAMTS5 prevents cartilage degradation in a murine model of osteoarthritis. *Nature* **434**, 644–648.
- Greig CA, Gray C, Rankin D, Young A, Mann V, Noble B & Atherton PJ (2011). Blunting of adaptive responses to resistance exercise training in women over 75y. *Exp Gerontol* **46**, 884–890.
- Gruber J, Vincent TL, Hermansson M, Bolton M, Wait R & Saklatvala J (2004). Induction of interleukin-1 in articular cartilage by explantation and cutting. *Arthritis Rheum* **50**, 2539–2546.
- Grumbles RM, Howell DS, Howard GA, Roos BA, Setton LA, Mow VC, Ratcliffe A, Müller FJ & Altman RD (1995). Cartilage metalloproteinases in disuse atrophy. *J Rheumatol Suppl* **43**, 146–148.
- Haapala J, Arokoski JP, Hyttinen MM, Lammi M, Tammi M, Kovanen V, Helminen HJ & Kiviranta I (1999). Remobilization does not fully restore immobilization induced articular cartilage atrophy. *Clin Orthop Relat Res* **218**–229.
- Haapala J, Arokoski JP, Rönkkö S, Agren U, Kosma VM, Lohmander LS, Tammi M, Helminen HJ & Kiviranta I (2001). Decline after immobilisation and recovery after remobilisation of synovial fluid IL1, TIMP, and chondroitin sulphate levels in young beagle dogs. *Ann Rheum Dis* **60**, 55–60.
- Haapala J, Lammi MJ, Inkinen R, Parkkinen JJ, Agren UM, Arokoski J, Kiviranta I, Helminen HJ & Tammi MI (1996). Coordinated regulation of hyaluronan and aggrecan content in the articular cartilage of immobilized and exercised dogs. *J Rheumatol* **23**, 1586–1593.
- Hamann N, Zaucke F, Heilig J, Oberländer KD, Brüggemann G-P & Niehoff A (2014). Effect of different running modes on the morphological, biochemical, and mechanical properties of articular cartilage. *Scand J Med Sci Sports* **24**, 179–188.
- Haubner M, Eckstein F, Schnier M, Löscher A, Sittke H, Becker C, Kolem H, Reiser M & Englmeier KH (1997). A non-invasive technique for 3-dimensional assessment of articular cartilage thickness based on MRI. Part 2: Validation using CT arthrography. *Magn Reson Imaging* **15**, 805–813.
- Heinemeier KM, Schjerling P, Heinemeier J, Møller MB, Krosgaard MR, Grum-Schwensen T, Petersen MM & Kjaer M (2016). Radiocarbon dating reveals minimal collagen turnover in both healthy and osteoarthritic human cartilage. *Sci Transl Med* **8**, 346ra90.

- Hunziker EB, Quinn TM & Häuselmann H-J (2002). Quantitative structural organization of normal adult human articular cartilage. *Osteoarthritis Cartilage* **10**, 564–572.
- Ingram KR, Wann AKT, Wingate RM, Coleman PJ, McHale N & Levick JR (2009). Signal pathways regulating hyaluronan secretion into static and cycled synovial joints of rabbits. *J Physiol* **587**, 4361–4376.
- Ismail HM, Didangelos A, Vincent TL & Saklatvala J (2017). Rapid activation of transforming growth factor β -activated kinase 1 in chondrocytes by phosphorylation and K⁶³-linked polyubiquitination upon injury to animal articular cartilage. *Arthritis Rheumatol* **69**, 565–575.
- Ismail HM, Miotla-Zarebska J, Troeberg L, Tang X, Stott B, Yamamoto K, Nagase H, Fosang AJ, Vincent TL & Saklatvala J (2016). Brief report: JNK-2 controls aggrecan degradation in murine articular cartilage and the development of experimental osteoarthritis. *Arthritis Rheumatol* **68**, 1165–1171.
- Ismail HM, Yamamoto K, Vincent TL, Nagase H, Troeberg L & Saklatvala J (2015). Interleukin-1 acts via the JNK-2 signaling pathway to induce aggrecan degradation by human chondrocytes. *Arthritis Rheumatol* **67**, 1826–1836.
- Jones G, Ding C, Glisson M, Hynes K, Ma D & Cicuttini F (2003). Knee articular cartilage development in children: a longitudinal study of the effect of sex, growth, body composition, and physical activity. *Pediatr Res* **54**, 230–236.
- Jurvelin J, Kiviranta I, Tammi M & Helminen HJ (1986a). Effect of physical exercise on indentation stiffness of articular cartilage in the canine knee. *Int J Sports Med* **7**, 106–110.
- Jurvelin J, Kiviranta I, Tammi M & Helminen HJ (1986b). Softening of canine articular cartilage after immobilization of the knee joint. *Clin Orthop Relat Res* **246–252**.
- Kahn J, Shwartz Y, Blitz E, Krief S, Sharir A, Breitel DA, Rattenbach R, Relaix F, Maire P, Rountree RB, Kingsley DM & Zelzer E (2009). Muscle contraction is necessary to maintain joint progenitor cell fate. *Dev Cell* **16**, 734–743.
- Kavanagh E, Church VL, Osborne AC, Lamb KJ, Archer CW, Francis-West PH & Pitsillides AA (2006). Differential regulation of GDF-5 and FGF-2/4 by immobilisation in ovo exposes distinct roles in joint formation. *Dev Dyn* **235**, 826–834.
- Kiviranta I, Jurvelin J, Tammi M, Säämänen A-M & Helminen HJ (1987). Weight bearing controls glycosaminoglycan concentration and articular cartilage thickness in the knee joints of young beagle dogs. *Arthritis Rheum* **30**, 801–809.
- Kiviranta I, Tammi M, Jurvelin J, Arokoski J, Säämänen A-M & Helminen HJ (1992). Articular cartilage thickness and glycosaminoglycan distribution in the canine knee joint after strenuous running exercise. *Clin Orthop Relat Res* **302–308**.
- Kiviranta I, Tammi M, Jurvelin J, Säämänen A-M & Helminen HJ (1988). Moderate running exercise augments glycosaminoglycans and thickness of articular cartilage in the knee joint of young beagle dogs. *J Orthop Res* **6**, 188–195.
- Lee W, Leddy HA, Chen Y, Lee SH, Zelenski NA, McNulty AL, Wu J, Becker KN, Coles J, Zauscher S, Grandl J, Sachs F, Guilak F & Liedtke WB (2014). Synergy between Piezo1 and Piezo2 channels confers high-strain mechanosensitivity to articular cartilage. *Proc Natl Acad Sci U S A* **111**, E5114–E5122.
- Leroux MA, Cheung HS, Bau JL, Wang JY, Howell DS & Setton LA (2001). Altered mechanics and histomorphometry of canine tibial cartilage following joint immobilization. *Osteoarthritis Cartilage* **9**, 633–640.
- Lösch A, Eckstein F, Haubner M & Englmeier KH (1997). A non-invasive technique for 3-dimensional assessment of articular cartilage thickness based on MRI. Part 1: Development of a computational method. *Magn Reson Imaging* **15**, 795–804.
- Madej W, van Caam A, Blaney Davidson EN, Hannink G, Buma P & van der Kraan PM (2016). Ageing is associated with reduction of mechanically-induced activation of Smad2/3P signaling in articular cartilage. *Osteoarthritis Cartilage* **24**, 146–157.
- Malda J, de Grauw JC, Benders KEM, Kik MJL, van de Lest CHA, Creemers LB, Dhert WJA & van Weeren PR (2013). Of mice, men and elephants: the relation between articular cartilage thickness and body mass. *PLoS One* **8**, e57683.
- Mankin HJ & Lippiello L (1969). The turnover of adult rabbit articular cartilage. *J Bone Joint Surg Am* **51**, 1591–1600.
- Meachim G (1963). The effect of scarification on articular cartilage in the rabbit. *J Bone Joint Surg Br* **45B**, 150–161.
- Mitchell PG, Magna HA, Reeves LM, Lopresti-Morrow LL, Yocum SA, Rosner PJ, Geoghegan KF & Hambor JE (1996). Cloning, expression, and type II collagenolytic activity of matrix metalloproteinase-13 from human osteoarthritic cartilage. *J Clin Invest* **97**, 761–768.
- Moriyama H, Yoshimura O, Kawamata S, Takayanagi K, Kurose T, Kubota A, Hosoda M & Tobimatsu Y (2008). Alteration in articular cartilage of rat knee joints after spinal cord injury. *Osteoarthritis Cartilage* **16**, 392–398.
- Müller FJ, Setton LA, Manicourt DH, Mow VC, Howell DS & Pita JC (1994). Centrifugal and biochemical comparison of proteoglycan aggregates from articular cartilage in experimental joint disuse and joint instability. *J Orthop Res* **12**, 498–508.
- Ni G-X, Liu S-Y, Lei L, Li Z, Zhou Y-Z & Zhan L-Q (2013). Intensity-dependent effect of treadmill running on knee articular cartilage in a rat model. *Biomed Res Int* **2013**, 172392.
- Nomura M, Sakitani N, Iwasawa H, Kohara Y, Takano S, Wakimoto Y, Kuroki H & Moriyama H (2017). Thinning of articular cartilage after joint unloading or immobilization. An experimental investigation of the pathogenesis in mice. *Osteoarthritis Cartilage* **25**, 727–736.
- O'Connor CJ, Leddy HA, Benefield HC, Liedtke WB & Guilak F (2014). TRPV4-mediated mechanotransduction regulates the metabolic response of chondrocytes to dynamic loading. *Proc Natl Acad Sci U S A* **111**, 1316–1321.
- O'Connor CJ, Ramalingam S, Zelenski NA, Benefield HC, Rigo I, Little D, Wu C-L, Chen D, Liedtke W, McNulty AL & Guilak F (2016). Cartilage-specific knockout of the mechanosensory ion channel TRPV4 decreases age-related osteoarthritis. *Sci Rep* **6**, 29053.
- Palmoski M, Perricone E & Brandt KD (1979). Development and reversal of a proteoglycan aggregation defect in normal canine knee cartilage after immobilization. *Arthritis Rheum* **22**, 508–517.

- Palmoski MJ & Brandt KD (1981). Running inhibits the reversal of atrophic changes in canine knee cartilage after removal of a leg cast. *Arthritis Rheum* **24**, 1329–1337.
- Pool WH (1974). Cartilage atrophy. *Radiology* **112**, 47–50.
- Poole CA (1997). Articular cartilage chondrons: form, function and failure. *J Anat* **191**, 1–13.
- Ratcliffe A, Shurety W & Caterson B (1993). The quantitation of a native chondroitin sulfate epitope in synovial fluid lavages and articular cartilage from canine experimental osteoarthritis and disuse atrophy. *Arthritis Rheum* **36**, 543–551.
- Recht MP, Kramer J, Marcelis S, Pathria MN, Trudell D, Haghighi P, Sartoris DJ & Resnick D (1993). Abnormalities of articular cartilage in the knee: analysis of available MR techniques. *Radiology* **187**, 473–478.
- Roos EM & Dahlberg L (2005). Positive effects of moderate exercise on glycosaminoglycan content in knee cartilage: a four-month, randomized, controlled trial in patients at risk of osteoarthritis. *Arthritis Rheum* **52**, 3507–3514.
- Ross TD, Coon BG, Yun S, Baeyens N, Tanaka K, Ouyang M & Schwartz MA (2013). Integrins in mechanotransduction. *Curr Opin Cell Biol* **25**, 613–618.
- Säämänen A-M, Tammi M, Jurvelin J, Kiviranta I & Helminen HJ (1990). Proteoglycan alterations following immobilization and remobilization in the articular cartilage of young canine knee (stifle) joint. *J Orthop Res* **8**, 863–873.
- Säämänen A-M, Tammi M, Kiviranta I & Helminen HJ (1988). Running exercise as a modulator of proteoglycan matrix in the articular cartilage of young rabbits. *Int J Sports Med* **9**, 127–133.
- Säämänen A-M, Tammi M, Kiviranta I, Jurvelin J & Helminen HJ (1989). Levels of chondroitin-6-sulfate and nonaggregating proteoglycans at articular cartilage contact sites in the knees of young dogs subjected to moderate running exercise. *Arthritis Rheum* **32**, 1282–1292.
- Shea CA, Rolfe RA & Murphy P (2015). The importance of foetal movement for co-ordinated cartilage and bone development in utero: clinical consequences and potential for therapy. *Bone Joint Res* **4**, 105–116.
- Siebelt M, Groen HC, Koelewijn SJ, de Blois E, Sandker M, Waarsing JH, Müller C, van Osch GJVM, de Jong M & Weinans H (2014). Increased physical activity severely induces osteoarthritic changes in knee joints with papain induced sulfate-glycosaminoglycan depleted cartilage. *Arthritis Res Ther* **16**, R32.
- Simon WH (1970). Scale effects in animal joints. I. Articular cartilage thickness and compressive stress. *Arthritis Rheum* **13**, 244–256.
- Sutter EG, Widmyer MR, Utturkar GM, Spritzer CE, Garrett WE & Defrate LE (2015). In vivo measurement of localized tibiofemoral cartilage strains in response to dynamic activity. *Am J Sports Med* **43**, 370–376.
- Tammi M, Säämänen A-M, Jauhiainen A, Malminen O, Kiviranta I & Helminen H (1983). Proteoglycan alterations in rabbit knee articular cartilage following physical exercise and immobilization. *Connect Tissue Res* **11**, 45–55.
- Tang X, Muhammad H, McLean C, Miotla-Zarebska J, Flemming J, Didangelos A, Onnerfjord P, Leask A, Saklatvala J & Vincent TL (2018). Connective tissue growth factor contributes to joint homeostasis and osteoarthritis severity by controlling the matrix sequestration and activation of latent TGF β . *Ann Rheum Dis*. <http://doi.org/10.1136/annrheumdis-2018-212964>
- Troeberg L & Nagase H (2012). Proteases involved in cartilage matrix degradation in osteoarthritis. *Biochim Biophys Acta* **1824**, 133–145.
- Uda Y, Azab E, Sun N, Shi C & Pajevic PD (2017). Osteocyte mechanobiology. *Curr Osteoporos Rep* **15**, 318–325.
- van Caam A, Madej W, Thijsen E, Garcia de Vinuesa A, van den Berg W, Goumans M-J, ten Dijke P, Blaney-Davidson E & van der Kraan PM (2016). Expression of TGF β -family signalling components in ageing cartilage: Age-related loss of TGF β and BMP receptors. *Osteoarthritis Cartilage* **24**, 1235–1245.
- Vanwanseele B, Eckstein F, Knecht H, Spaepen A & Stüssi E (2003). Longitudinal analysis of cartilage atrophy in the knees of patients with spinal cord injury. *Arthritis Rheum* **48**, 3377–3381.
- Vanwanseele B, Eckstein F, Knecht H, Stüssi E & Spaepen A (2002). Knee cartilage of spinal cord-injured patients displays progressive thinning in the absence of normal joint loading and movement. *Arthritis Rheum* **46**, 2073–2078.
- Vincent T, Hermansson M, Bolton M, Wait R & Saklatvala J (2002). Basic FGF mediates an immediate response of articular cartilage to mechanical injury. *Proc Natl Acad Sci U S A* **99**, 8259–8264.
- Vincent TL (2013). Targeting mechanotransduction pathways in osteoarthritis: a focus on the pericellular matrix. *Curr Opin Pharmacol* **13**, 449–454.
- Vincent TL, Hermansson MA, Hansen UN, Amis AA & Saklatvala J (2004). Basic fibroblast growth factor mediates transduction of mechanical signals when articular cartilage is loaded. *Arthritis Rheum* **50**, 526–533.
- Vincent TL, McLean CJ, Full LE, Peston D & Saklatvala J (2007). FGF-2 is bound to perlecan in the pericellular matrix of articular cartilage, where it acts as a chondrocyte mechanotransducer. *Osteoarthritis Cartilage* **15**, 752–763.
- Wann AKT, Zuo N, Haycraft CJ, Jensen CG, Poole CA, McGlashan SR & Knight MM (2012). Primary cilia mediate mechanotransduction through control of ATP-induced Ca²⁺ signaling in compressed chondrocytes. *FASEB J* **26**, 1663–1671.
- Watt FE, Ismail HM, Didangelos A, Peirce M, Vincent TL, Wait R & Saklatvala J (2013). Src and fibroblast growth factor 2 independently regulate signaling and gene expression induced by experimental injury to intact articular cartilage. *Arthritis Rheum* **65**, 397–407.
- Yuasa T, Kondo N, Yasuhara R, Shimono K, Mackem S, Pacifici M, Iwamoto M & Enomoto-Iwamoto M (2009). Transient activation of Wnt/ β -catenin signaling induces abnormal growth plate closure and articular cartilage thickening in postnatal mice. *Am J Pathol* **175**, 1993–2003.
- Ziaaldini MM, Marzetti E, Picca A & Murlasits Z (2017). Biochemical pathways of sarcopenia and their modulation by physical exercise: a narrative review. *Front Med (Lausanne)* **4**, 167.

Additional information

Competing interests

None declared.

Author contributions

Both authors have approved the final version of the manuscript and agree to be accountable for all aspects of

the work. All persons designated as authors qualify for authorship, and all those who qualify for authorship are listed.

Funding

We acknowledge salary support from Arthritis Research UK (grant no. 20205) and the Kennedy Trust for Rheumatology Research.