



Published in final edited form as:

*Microbiol Spectr.* 2019 March ; 7(2): . doi:10.1128/microbiolspec.PSIB-0009-2018.

## Type VI secretion systems and the gut microbiota

**Michael J. Coyne and Laurie E. Comstock**

Division of Infectious Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, MA

### Abstract

The human colonic microbiota is a dense ecosystem comprised of numerous microbes including bacteria, phage, fungi, archaea, and protozoa that compete for nutrients and space. Studies are beginning to reveal the antagonistic mechanisms that gut bacteria use to compete with other members of this ecosystem. In the healthy human colon, the majority of the Gram negative bacteria are of the order Bacteroidales. Proteobacteria, such as *E. coli*, are numerically fewer, but confer important properties to the host, such as colonization resistance. Several enteric pathogens use T6SSs to antagonize symbiotic gut *E. coli* facilitating colonization and disease progression. T6SSs loci are also widely distributed in human gut Bacteroidales, which includes three predominant genera: *Bacteroides*, *Parabacteroides*, and *Prevotella*. There are three distinct genetic architectures of T6SS loci among the gut Bacteroidales, termed GA1, GA2, and GA3. GA1 and GA2 T6SS loci are contained on integrative and conjugative elements and are the first T6SS loci shown to be readily transferred in the human gut between numerous species and families of Bacteroidales. In contrast, the GA3 T6SSs are present exclusively in *Bacteroides fragilis*. There are divergent regions in all three T6SS GAs that contain genes encoding effector and immunity proteins, many of which function by unknown mechanisms. To date, only the GA3 T6SSs have been shown to antagonize bacteria, and they target nearly all gut Bacteroidales species analyzed. This chapter will delve more deeply into properties of the T6SSs of these human gut bacteria and the ecological outcomes of their synthesis *in vivo*. Type VI secretion systems were first identified and characterized in pathogenic bacteria of the Proteobacterial phylum (1, 2). The discovery in 2010 that these secretion systems can target and intoxicate not only eukaryotic cells, but also other bacteria (3), revealed that some T6SSs help bacteria compete with other bacteria in a community setting. Indeed, many Proteobacterial symbionts, including the plant symbiont *Pseudomonas putida*, the bumble bee gut symbiont *Snodgrassella alvi*, and the squid symbiont *Vibrio fischeri*, all have T6SSs that provide a competitive advantage in their natural ecosystems (4–6). An early *in silico* analysis using COG models of Proteobacterial T6SS proteins against primary sequence databases suggested that T6SSs are largely confined to Proteobacterial species, which are minor members of some human-associated microbial communities such as the gut microbiota (7, 8)

In the dense microbial ecosystem of the human colon, contact-dependent mechanisms of antagonism, such as a T6SS, should be an effective means of thwarting competitors. In this microbial ecosystem, the predominant Gram negative bacteria are of the phylum Bacteroidetes, comprising approximately half of the colonic bacteria in most people, and vastly outnumbering

commensal Proteobacteria. Most human gut Bacteroidetes species are contained within the order Bacteroidales, which includes several different families, each with one predominant genus in the human gut: *Bacteroides* (family Bacteroidaceae), the *Parabacteroides* (family Tannerellaceae), *Prevotella* (family Prevotellaceae), and *Alistipes* (family Rickenellaceae), as well as families with more minor representatives. Humans are colonized at high density with numerous Bacteroidales species simultaneously (9–11) and the abundant species in one individual may not be the same as in another (9, 10). T6SSs were not discovered in species of the phylum Bacteroidetes until recently (12–14), mostly due to the lack of primary sequence or profile-sequence similarity of the 13 core Proteobacterial T6SS proteins with Bacteroidetes proteins. The use of profile-profile analyses revealed that Bacteroidales species of the human gut are a rich source of diverse T6SSs (15). As structural and mechanistic properties of T6SSs are discussed in the preceding chapter, here we will focus on unique properties of the T6SS of the human gut Bacteroidales, the ecological and functional properties of these T6SSs that are known to date, and the ecological advantages conferred by the T6SSs of human gut Proteobacterial strains *in vivo*.

---

## GENETIC CHARACTERISTICS AND PROPERTIES OF THE T6SS OF GUT BACTEROIDALES

An analysis of 205 human gut Bacteroidales genomes including 35 different species of four genera, *Bacteroides*, *Parabacteroides*, *Prevotella*, and *Alistipes*, revealed the presence of 130 T6SS loci in 115 of these strains with 15 strains containing two different T6SS loci (16). These T6SS loci were found in the genomes of 19 different species of *Bacteroides*, *Parabacteroides*, and *Prevotella*, but were not found in any of the nine *Alistipes* genomes analyzed. A notable feature is that these T6SSs segregated into three very distinct genetic architectures (GA), termed GA1, GA2 and GA3 (16) (Fig 1A). Within a given GA, the majority of the T6SS genes are of high DNA identify, interspersed with small regions that are variable (Fig 1A). The variable regions of all three T6SS GAs contain genes encoding effector and immunity proteins, some of which are similar to those previously described in other organisms (16, 17). Most of the GA3 T6SS effector and immunity proteins are unlike those previously described and function by mechanisms currently unknown. By comparing the divergent regions within a GA, there are predicted to be at least 30 different variable regions in the 48 GA1 loci analyzed, 21 in the 9 GA2 regions analyzed, and 17 in the 56 GA3 regions analyzed, with each divergent region likely encoding at least one effector immunity pair. Therefore, the T6SSs of gut Bacteroidales encode numerous distinct toxins, many of which operate *via* unknown mechanisms.

An analysis of the predicted proteins produced by these loci showed that four of the conserved T6SS proteins of Proteobacterial species are missing in gut Bacteroidales T6SS: TssA, TssJ, TssL, and TssM. TssJ, L, and M are components of the transmembrane complex (18) and TssA binds this complex and likely recruits the baseplate assemblage and coordinates tail tube and sheath biogenesis (19, 20). Instead of encoding these proteins, all three GAs of Bacteroidales T6SS loci encode four conserved proteins not present in Proteobacterial T6SS loci, namely TssO-TssR (13, 16). These likely serve as functional orthologs of the Proteobacterial proteins comprising the transmembrane complex.

All three T6SS GAs encode multiple TssD (Hcp) needle proteins, with GA2 and GA3 loci encoding six and five distinct TssD proteins, respectively. In the GA2 loci, one of the six TssD proteins has a C-terminal extension, likely conferring toxin activity to this protein (Fig 1A). In the GA3 locus, the main structural TssD protein was identified among the five TssD proteins (Fig 1A)(21). The function of the accessory TssD proteins may be to bind effector proteins and incorporate them into the needle structure, as in both the GA2 and GA3 loci, four of the *tssD* genes flank the effector and immunity gene variable regions (Fig 1A).

An important feature of the three different GAs is their distribution among gut Bacteroidales species. GA1 and GA2 T6SS loci are present in numerous species of *Bacteroides*, *Parabacteroides*, and *Prevotella*, whereas GA3 T6SSs are present exclusively in *B. fragilis*. The wide distribution of GA1 and GA2 in at least three distinct families of Bacteroidales is due to their presence on integrative and conjugative elements (ICE) (12, 15) (Fig 1B). ICE containing GA1 T6SSs are in the range of 120 kb and are extremely similar along their length to ICE containing distinct GA1 T6SS. Excluding the T6SS divergent regions, the DNA of ICE containing distinct GA1 T6SS loci are approximately 95% similar over their entire length. Although very dissimilar to the GA1 ICE, GA2 containing ICE are on the order of 100 kb and share approximately 75% - 99% DNA identity over their length with other GA2 containing ICE. As ICE are chromosomal self-transmissible mobile elements (22, 23), they have the ability to move between species. In fact, the GA1 containing ICE have been found to be transferred extensively between Bacteroidales species that are co-resident in the human intestine (12, 16)(Fig 1C). Examination of hundreds of genomes of gut Bacteroidales has indicated that GA2 T6SS are not present in a strain that has either a GA1 or GA3 T6SS locus. However, *B. fragilis* strains can have both GA1 and GA3 T6SS loci in their genome. The reason for the apparent lack of GA2 T6SS loci in the same chromosome with GA1 or GA3 T6SS loci is currently unknown. Bacteroidales species present in the gut of an individual can, however, collectively contain all three different T6SS GAs (16) (Fig 1C).

## ECOLOGICAL CONSEQUENCES OF GUT BACTEROIDALES T6SS ANTAGONISM

The fitness benefits of the T6SSs of the gut Bacteroidales are still incompletely understood. The best studied are the GA3 T6SSs of *B. fragilis*. GA3 T6SSs were found to in approximately 86% of *B. fragilis* strains based on genome or metagenome analyses including hundreds of strains (16, 24), making them widely distributed in the species. The GA3 T6SSs of two different *B. fragilis* strains were found to antagonize all gut Bacteroidales species tested including *Bacteroides* and *Parabacteroides* species, but not the one *Prevotella copri* strain analyzed nor other *B. fragilis* with the same GA3 T6SS region or immunity genes (21, 25). In addition, no killing was evident against any gut Proteobacterial species analyzed (21), suggesting specificity for Bacteroidales. The TssD needle protein of the *B. fragilis* 638R GA3 T6SS is present in the supernatant of actively growing cells *in vitro* as well as in the feces of monoassociated gnotobiotic mice (21), suggesting this antagonistic system is constitutively synthesized and firing, rather than only responding to specific external signals or threats. The number of GA3 T6SS transmission events in a

human gut colonized at typical levels with *B. fragilis* was predicted to be on the order of  $6 \times 10^{10}$ - $10^{11}$  per day (25).

Effector and immunity proteins were identified in three different GA3 T6SS loci (21, 25, 26). These toxic effector proteins are not similar to other known proteins and therefore intoxicate by as yet unidentified mechanisms. The toxicity of two of these effectors requires an added N-terminal periplasmic targeting sequence when they are produced inside a sensitive cell from an inducible promoter (27), suggesting that they may need to be localized to the periplasm for toxicity. In a few strains, genes encoding immunity proteins to GA3 effectors were found outside of the T6SS regions, in some cases in strains that did not have a GA3 T6SS locus (25). These immunity genes were found to confer protection from attack by a *B. fragilis* strain synthesizing the T6SS effectors to which the immunity proteins are directed, suggesting that acquisition of these immunity genes confers an advantage to an organisms by protecting it from GA3 T6SS-mediated antagonism (25). A recent study analyzing human gut metagenomic data revealed arrays of orphan immunity genes, which were termed Acquired Interbacterial Defense (AID) gene clusters (28). These orphan immunity islands reside on predicted mobile elements and include immunity genes likely derived from *B. fragilis* GA3 T6SS loci and disseminated by lateral transfer.

Although the GA3 T6SSs are very effective at antagonizing Bacteroidales species *in vitro*, the effects *in vivo* are more variable. As *B. fragilis* are known to co-exist in the human gut with numerous other Bacteroidales species that are susceptible to their GA3 T6SS, the spatial organization of the microbiota likely dictates the effectiveness of this weapon. In a gnotobiotic mouse competitive colonization model, an isogenic T6SS<sup>+</sup> wild type strain outcompetes an isogenic mutant strain lacking the T6SS effector and immunity genes (21, 25). As isogenic strains should share the same spatial and nutritional niche and therefore should make frequent contacts, a strong antagonistic effect is expected. The ability of GA3 T6SSs of *B. fragilis* to antagonize other wild type non-isogenic *B. fragilis* strains *in vivo* was also demonstrated (25, 26). However, the effectiveness of the GA3 T6SS was less when analyzing antagonism of different *Bacteroides* species (Fig 2B). A significant effect was observed with a *B. vulgatus* strain (25), but not with *B. thetaiotaomicron* (21, 25). It may be that in this model, where the mice are provided ample nutrients and likely utilize different nutrients in the gut (29, 30), *B. fragilis* and *B. thetaiotaomicron* would make infrequent contacts (Fig 2B).

Analysis of metagenomic datasets of human gut samples revealed a link between the presence of Bacteroidales T6SSs and the composition of the microbiota, especially in regard to the GA3 T6SSs of *B. fragilis* (24). The presence of GA3 T6SS genes correlated significantly with an abundance of *Bacteroides* and a decrease in specific Firmicutes. Moreover, the microbiota of the developing infant gut are significantly more likely to contain GA3 T6SSs than those of adults. Also, *B. fragilis* strain replacement in the infant microbiota is more pronounced than in adults, which appear to be dominated by a single strain of *B. fragilis*. These findings suggest that competition amongst *B. fragilis* strains for dominance is fiercest early in life, and that the ultimate microbiota composition may be influenced by GA3 T6SSs (24).

The functions of the GA1 and GA2 T6SSs have not yet been elucidated. There has been no demonstration that these T6SSs target bacteria as do the GA3 T6SSs. Many of the identifiable effectors are predicted nucleases or other nucleic acid targeting enzymes (16) that could function in bacteria, archaea, or eukaryotes. The fact that the ICE containing these T6SSs are shared between different co-resident species of Bacteroidales in the human gut suggests they likely do not have a Bacteroidales target and may instead provide defense against a common enemy. It is also possible that these T6SSs may allow for nutrient acquisition, or protection from environmental stressors, as has been shown for T6SSs in other organisms (31, 32). The prevalence and transfer of these systems among human gut Bacteroidales species makes them intriguing secretion systems for continued analysis.

## T6SS EFFECTS OF ENTERIC PROTEOBACTERIA IN THE MAMMALIAN GUT

Although less abundant in the healthy human gut microbiota than Bacteroidales, *E. coli* gut symbionts play a crucial role in colonization resistance against enteric pathogens of the Proteobacterial phylum. Enteric pathogens such as *Vibrio cholerae*, *Shigella sonnei*, and *Salmonella enterica*, have all been shown to utilize T6SSs to overcome colonization resistance by antagonizing resident gut *E. coli* (33–37) (Fig 2A). What is less studied is whether the symbiotic gut *E. coli* have T6SSs that function in colonization resistance against enteric pathogens. Antagonism by the diffusible microcin toxin produced by *E. coli* Nissle was shown to function as a colonization barrier to enteric pathogens (38) providing precedent for such an effect. Queries of a set of 1267 human gut metagenomes consolidated into the “three cohorts gene catalog (3CGC)” (39) for matches to Pfam models identifying the four conserved Proteobacterial T6SS proteins absent in Bacteroidales T6SSs (TssA, J, L, and M) revealed that 174 of these metagenomes encoded proteins with motifs that met or exceeded the gathering threshold of all four of the Proteobacterial models used (Coyne and Comstock, unpublished). These findings suggest that resident gut *E. coli* strains likely harbor T6SSs, the *in vivo* effects of which remain to be determined.

## CONCLUDING REMARKS

The discovery of T6SS loci and their prevalence in diverse human gut Bacteroidales species has revealed that the composition of the human gut microbiota is likely significantly influenced by these secretion systems. We still know very little about these secretion systems, including the targets of the GA1 and GA2 T6SSs, the advantage of transfer of these regions to co-resident Bacteroidales species, and the mechanisms of action of many of the toxic effectors. In addition, it will be interesting to determine the prevalence of T6SSs in human gut *E. coli* strains and whether these antagonistic systems contribute to their ability to affect colonization by enteric pathogens.

## Acknowledgements

The Comstock lab is funded to study T6SSs of gut Bacteroidales by Public Health Service grant 01AI120633 from the NIH/National Institute of Allergy and Infectious Diseases.


## References

1. Mougous JD, Cuff ME, Raunser S, Shen A, Zhou M, Gifford CA, Goodman AL, Joachimiak G, Ordonez CL, Lory S, Walz T, Joachimiak A, Mekalanos JJ. 2006 A virulence locus of *Pseudomonas aeruginosa* encodes a protein secretion apparatus. *Science* 312:1526–1530. [PubMed: 16763151]
2. Pukatzki S, Ma AT, Sturtevant D, Krastins B, Sarracino D, Nelson WC, Heidelberg JF, Mekalanos JJ. 2006 Identification of a conserved bacterial protein secretion system in *Vibrio cholerae* using the *Dictyostelium* host model system. *Proc Natl Acad Sci U S A* 103:1528–1533. [PubMed: 16432199]
3. Hood RD, Singh P, Hsu F, Guvener T, Carl MA, Trinidad RR, Silverman JM, Ohlson BB, Hicks KG, Plemel RL, Li M, Schwarz S, Wang WY, Merz AJ, Goodlett DR, Mougous JD. 2010 A type VI secretion system of *Pseudomonas aeruginosa* targets a toxin to bacteria. *Cell Host Microbe* 7:25–37. [PubMed: 20114026]
4. Bernal P, Allsopp LP, Filloux A, Llamas MA. 2017 The *Pseudomonas putida* T6SS is a plant warden against phytopathogens. *ISME J* 11:972–987. [PubMed: 28045455]
5. Steele MI, Kwong WK, Whiteley M, Moran NA. 2017 Diversification of Type VI Secretion System Toxins Reveals Ancient Antagonism among Bee Gut Microbes. *MBio* 8.
6. Speare L, Cecere AG, Guckes KR, Smith S, Wollenberg MS, Mandel MJ, Miyashiro T, Septer AN. 2018 Bacterial symbionts use a type VI secretion system to eliminate competitors in their natural host. *Proc Natl Acad Sci U S A* doi:10.1073/pnas.1808302115.
7. Human Microbiome Project C. 2012 Structure, function and diversity of the healthy human microbiome. *Nature* 486:207–214. [PubMed: 22699609]
8. Grice EA, Segre JA. 2012 The human microbiome: our second genome. *Annu Rev Genomics Hum Genet* 13:151–170. [PubMed: 22703178]
9. Hayashi H, Sakamoto M, Benno Y. 2002 Phylogenetic analysis of the human gut microbiota using 16S rDNA clone libraries and strictly anaerobic culture-based methods. *Microbiol Immunol* 46:535–548. [PubMed: 12363017]
10. Eckburg PB, Bik EM, Bernstein CN, Purdom E, Dethlefsen L, Sargent M. 2005 Diversity of the human intestinal microbial flora. *Science* 308.
11. Zitomersky NL, Coyne MJ, Comstock LE. 2011 Longitudinal analysis of the prevalence, maintenance, and IgA response to species of the order Bacteroidales in the human gut. *Infect Immun* 79:2012–2020. [PubMed: 21402766]
12. Coyne M, Zitomersky N, McGuire A, Earl A, Comstock L. 2014 Evidence of extensive DNA transfer between Bacteroidales species within the human gut. *mBio* 5.
13. Russell AB, Wexler AG, Harding BN, Whitney JC, Bohn AJ, Goo YA, Tran BQ, Barry NA, Zheng H, Peterson SB, Chou S, Gonen T, Goodlett DR, Goodman AL, Mougous JD. 2014 A Type VI Secretion-Related Pathway in Bacteroidetes Mediates Interbacterial Antagonism. *Cell Host Microbe* 16:227–236. [PubMed: 25070807]
14. Wilson MM, Anderson DE, Bernstein HD. 2015 Analysis of the outer membrane proteome and secretome of *Bacteroides fragilis* reveals a multiplicity of secretion mechanisms. *PLoS One* 10.
15. Coyne MJ, Roelofs KG, Comstock LE. 2016 Type VI secretion systems of human gut Bacteroidales segregate into three genetic architectures, two of which are contained on mobile genetic elements. *BMC Genomics* 17:1–21. [PubMed: 26818753]
16. Coyne M, Roelofs KG, Comstock LE. 2016 Type VI secretion systems of human gut Bacteroidales segregate into three genetic architectures, two of which are contained on mobile genetic elements. *BMC Genomics* 17.
17. Zhang D, de Souza RF, Anantharaman V, Iyer LM, Aravind L. 2012 Polymorphic toxin systems: Comprehensive characterization of trafficking modes, processing, mechanisms of action, immunity and ecology using comparative genomics. *Biol Direct* 7:18. [PubMed: 22731697]
18. Durand E, Nguyen VS, Zoued A, Logger L, Pehau-Arnaudet G, Aschtgen MS, Spinelli S, Desmyter A, Bardiaux B, Dujeancourt A, Roussel A, Cambillau C, Cascales E, Fronzes R. 2015 Biogenesis and structure of a type VI secretion membrane core complex. *Nature* 523:555–560. [PubMed: 26200339]

19. Zoued A, Durand E, Brunet YR, Spinelli S, Douzi B, Guzzo M, Flaugnatti N, Legrand P, Journet L, Fronzes R, Mignot T, Cambillau C, Cascales E. 2016 Priming and polymerization of a bacterial contractile tail structure. *Nature* 531:59–63. [PubMed: 26909579]
20. Zoued A, Durand E, Santin YG, Journet L, Roussel A, Cambillau C, Cascales E. 2017 TssA: The cap protein of the Type VI secretion system tail. *Bioessays* 39.
21. Chatzidaki-Livanis M, Geva-Zatorsky N, Comstock LE. 2016 *Bacteroides fragilis* type VI secretion systems use novel effector and immunity proteins to antagonize human gut Bacteroidales species. *Proc Natl Acad Sci U S A* 113:3627–3632. [PubMed: 26951680]
22. Waters JL, Salyers AA. 2013 Regulation of CTnDOT conjugative transfer is a complex and highly coordinated series of events. *MBio* 4:e00569–00513.
23. Wozniak RA, Waldor MK. 2010 Integrative and conjugative elements: mosaic mobile genetic elements enabling dynamic lateral gene flow. *Nat Rev Microbiol* 8:552–563. [PubMed: 20601965]
24. Verster AJ, Ross BD, Radey MC, Bao Y, Goodman AL, Mougous JD, Borenstein E. 2017 The Landscape of Type VI Secretion across Human Gut Microbiomes Reveals Its Role in Community Composition. *Cell Host Microbe* 22:411–419 e414. [PubMed: 28910638]
25. Wexler AG, Bao Y, Whitney JC, Bobay LM, Xavier JB, Schofield WB, Barry NA, Russell AB, Tran BQ, Goo YA, Goodlett DR, Ochman H, Mougous JD, Goodman AL. 2016 Human symbionts inject and neutralize antibacterial toxins to persist in the gut. *Proc Natl Acad Sci U S A* 113:3639–3644. [PubMed: 26957597]
26. Hecht AL, Casterline BW, Earley ZM, Goo YA, Goodlett DR, Bubeck Wardenburg J. 2016 Strain competition restricts colonization of an enteric pathogen and prevents colitis. *EMBO Rep* 17:1281–1291. [PubMed: 27432285]
27. Lim B, Zimmermann M, Barry NA, Goodman AL. 2017 Engineered Regulatory Systems Modulate Gene Expression of Human Commensals in the Gut. *Cell* 169:547–558 e515. [PubMed: 28431252]
28. Ross BD, Verster AJ, Radey MC, Schmidtke DT, Pope CE, Hoffman LR, Hajjar A, Peterson SB, Borenstein E, Mougous J. 2018 Acquired interbacterial defense systems protect against interspecies antagonism in the human gut microbiome. *bioRxiv* doi:10.1101/471110:471110.
29. Pudlo NA, Urs K, Kumar SS, German JB, Mills DA, Martens EC. 2015 Symbiotic Human Gut Bacteria with Variable Metabolic Priorities for Host Mucosal Glycans. *MBio* 6:e01282–01215. [PubMed: 26556271]
30. Tuncil YE, Xiao Y, Porter NT, Reuhs BL, Martens EC, Hamaker BR. 2017 Reciprocal Prioritization to Dietary Glycans by Gut Bacteria in a Competitive Environment Promotes Stable Coexistence. *MBio* 8.
31. Si M, Zhao C, Burkinshaw B, Zhang B, Wei D, Wang Y, Dong TG, Shen X. 2017 Manganese scavenging and oxidative stress response mediated by type VI secretion system in *Burkholderia thailandensis*. *Proc Natl Acad Sci U S A* 114:E2233–E2242. [PubMed: 28242693]
32. Lin J, Zhang W, Cheng J, Yang X, Zhu K, Wang Y, Wei G, Qian PY, Luo ZQ, Shen X. 2017 A *Pseudomonas* T6SS effector recruits PQS-containing outer membrane vesicles for iron acquisition. *Nat Commun* 8:14888. [PubMed: 28348410]
33. Zhao W, Caro F, Robins W, Mekalanos JJ. 2018 Antagonism toward the intestinal microbiota and its effect on *Vibrio cholerae* virulence. *Science* 359:210–213. [PubMed: 29326272]
34. Fast D, Kostiuk B, Foley E, Pukatzki S. 2018 Commensal pathogen competition impacts host viability. *Proc Natl Acad Sci U S A* 115:7099–7104. [PubMed: 29915049]
35. Anderson MC, Vonaesch P, Saffarian A, Marteyn BS, Sansonetti PJ. 2017 *Shigella sonnei* Encodes a Functional T6SS Used for Interbacterial Competition and Niche Occupancy. *Cell Host Microbe* 21:769–776 e763. [PubMed: 28618272]
36. Sana TG, Flaugnatti N, Lugo KA, Lam LH, Jacobson A, Baylot V, Durand E, Journet L, Cascales E, Monack DM. 2016 *Salmonella Typhimurium* utilizes a T6SS-mediated antibacterial weapon to establish in the host gut. *Proc Natl Acad Sci U S A* 113:E5044–5051. [PubMed: 27503894]
37. Pezoa D, Blondel CJ, Silva CA, Yang HJ, Andrews-Polymenis H, Santiviago CA, Contreras I. 2014 Only one of the two type VI secretion systems encoded in the *Salmonella enterica* serotype Dublin genome is involved in colonization of the avian and murine hosts. *Vet Res* 45:2. [PubMed: 24405577]

38. Sassone-Corsi M, Nuccio SP, Liu H, Hernandez D, Vu CT, Takahashi AA, Edwards RA, Raffatellu M. 2016 Microcins mediate competition among Enterobacteriaceae in the inflamed gut. *Nature* 540:280–283. [PubMed: 27798599]
39. Li J, Jia H, Cai X, Zhong H, Feng Q, Sunagawa S, Arumugam M, Kultima JR, Prifti E, Nielsen T, Juncker AS, Manichanh C, Chen B, Zhang W, Levenez F, Wang J, Xu X, Xiao L, Liang S, Zhang D, Zhang Z, Chen W, Zhao H, Al-Aama JY, Edris S, Yang H, Wang J, Hansen T, Nielsen HB, Brunak S, Kristiansen K, Guarner F, Pedersen O, Dore J, Ehrlich SD, Bork P, Wang J, Meta HITC. 2014 An integrated catalog of reference genes in the human gut microbiome. *Nat Biotechnol* 32:834–841. [PubMed: 24997786]





**Figure 1.**

**A.** ORF maps of one representative locus of each of the three genetic architectures (GA) of T6SS loci of gut Bacteroidales. T6SS loci of GA1 and GA2 are present in diverse Bacteroidales species whereas GA3 T6SS loci are confined to *B. fragilis*. T6SS loci of a given GA are extremely similar to each other except for the divergent regions noted by lines above the genes, which encode known or putative effector and immunity proteins. The major TssD protein of GA3 is noted as is the TssD protein of the GA2 loci that have C-terminal extensions likely conferring toxin activity. The ends of the GA1 and GA2 loci have not been precisely determined. **B.** ORF maps of ICE containing GA1 and GA2 T6SS loci of two *Bacteroides* species. The T6SS loci are designated by a line above the map. Genes involved in conjugative transfer (*tra* genes) are colored green (15). **C.** The abundant fecal gut Bacteroidales from three different healthy humans (CL02, CL09, and CL03) were analyzed for the presence of T6SSs. Seven Bacteroidales species were isolated and sequenced from subject CL02 and from subject CL09. In each subject, four of the seven species harbor nearly identical GA1 T6SSs loci within a subject, demonstrating transfer of these ICE between these strains in their gut (12, 15). In contrast, of the eight species isolated and sequenced from human subject CL03, two contain GA2 T6SS loci, albeit with different divergent regions. Therefore, these GA2 ICE were not transferred between these species. In addition, one species contains a GA1 T6SS locus and the *B. fragilis* strain from this


individual contains a GA3 T6SS locus (15). Red, green and yellow dots represent the GA1, GA2, and GA3 T6SS loci.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Figure 2.**

T6SS mediated antagonism in the mammalian gut. **A.** Three different Proteobacterial enteric pathogens, *Vibrio cholerae*, *Salmonella enterica* Typhimurium, and *Shigella sonnei* use T6SSs to target resident gut *E. coli* to overcome colonization resistance and cause disease in animal models (33–37). In the case of *V. cholerae*, the lysed *E. coli* initiate innate immune responses that upregulate virulence factors and increase dissemination (33). **B.** *Bacteroides fragilis* GA3 T6SS antagonize nearly all gut Bacteroidales species *in vitro*. *In vivo*, strong antagonistic effects are seen between two distinct *B. fragilis* strains likely due to their localization at the mucosal surface where they will make frequent contacts. This intraspecies antagonism may lead to the dominance of one strain. *B. vulgatus* was also significantly antagonized by a *B. fragilis* GA3 T6SS, possibly due to overlapping nutritional niches. In contrast, a significant antagonistic effect by the GA3 T6SS of *B. fragilis* was not observed when co-inoculated with *B. thetaiotaomicron*. These varied effects may be due to the substrate preferences of these species that may spatially segregate them under normal dietary conditions.