



HHS Public Access

Author manuscript

Trends Cogn Sci. Author manuscript; available in PMC 2020 March 01.

Published in final edited form as:

Trends Cogn Sci. 2019 March ; 23(3): 200–212. doi:10.1016/j.tics.2019.01.003.

An Evolutionary Perspective on Why Food Overconsumption Impairs Cognition

Mark P. Mattson

Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, MD 21205.

Laboratory of Neurosciences, National Institute on Aging Intramural Research Program, Baltimore, MD 21224.

Abstract

Brain structures and neuronal networks that mediate spatial navigation, decision-making, sociality and creativity evolved, in part, to enable success in food acquisition. Here I discuss evidence suggesting that the reason that overconsumption of energy-rich foods negatively impacts cognition is that signaling pathways that evolved to respond adaptively to food scarcity are relatively disengaged in the setting of continuous food availability. Obesity impairs cognition and increases the risk for some psychiatric disorders and dementias. Moreover, maternal and paternal obesity predispose offspring to poor cognitive outcomes by epigenetic molecular mechanisms. Neural signaling pathways that evolved to bolster cognition in settings of food insecurity can be stimulated by intermittent fasting and exercise to support the cognitive health of current and future generations.

Keywords

brain evolution; epigenetic modifications; hippocampus; intermittent fasting; ketogenic state; obesity; prefrontal cortex; spatial navigation

Setting the Stage

Eating three energy-rich meals plus snacks every day with no physical exertion required is unusual when viewed in the light of evolution. Novel insights into mechanisms of human cognition emerge when information on how food scarcity drove brain evolution is considered in the light of emerging data on how dietary energy intake impacts cognitive trajectories. These insights include: 1) An ecological factor that played a major role in the evolution of advanced cognitive capabilities (food scarcity) has been largely eliminated from day-to-day experience of modern-day humans and domesticated animals. 2) Overindulgent sedentary lifestyles negatively impact cognition, and the underlying metabolic state and its

Correspondence: mmattso2@jhmi.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

associated poor cognitive outcomes can be transmitted epigenetically to offspring. The possibility that this state of excessive energy intake will continue has important implications for brain evolution and the cognitive trajectories of future generations; 3) The cellular and molecular signaling pathways by which the challenge of intermittent food deprivation bolsters cognition have been elucidated in animal studies of intermittent food deprivation and exercise/running; and 4) Incorporation of feeding patterns that include intermittent fasting (IF) into the lifestyles of humans (and domesticated animals) can augment their cognitive capabilities and neuronal resilience.

The Cognitive Repertoire as Evolved Adaptations to Food Scarcity

Many animals in the wild and our human ancestors evolved in environments with sporadic food availability such that they commonly experienced extended periods of many days or more without acquiring food. Accordingly, natural selection favored individuals capable of outperforming their competitors, cognitively and/or physically when in a food-deprived state. The wide range of cognitive capabilities throughout the animal kingdom – spatial navigation, decision-making, sociality and creativity – are largely concerned with food acquisition and reproduction. Success in food acquisition is tightly linked with reproductive success, a topic reviewed elsewhere [1] and not considered further here. From a bioenergetic perspective a major conserved adaptation to food scarcity was metabolic switching from utilization of liver-derived glucose to a ketogenic state in which adipose cell-derived fatty acids and ketones are utilized by neurons and muscle cells as cellular fuels to sustain cognitive performance and physical endurance, respectively [2]. As discussed later in this article, such metabolic switching also stimulates neural signaling pathways that bolster cognition.

Lessons From Rodents, Corvids and Non-Human Primates

Accurate navigation in complex environments is fundamental for success in food acquisition. The neuronal networks and the cellular and molecular mechanisms that control spatial navigation and decision-making in rodents are being elucidated [3]. This research has established a fundamental role for the hippocampus and its functional connections with the entorhinal cortex, prefrontal cortex (PFC), cingulate cortex and associated cerebral cortical networks involved in processing and responding to neural representations of objects and sounds in the environment. Evidence that individual neurons encode the current position and orientation of the animal, as well as ‘imagined’ future navigation paths emerged from electrophysiological recordings and brain imaging studies of rodents navigating through space [4, 5] or in virtual reality scenarios [6–8]. This research established that neurons in the hippocampus called “place cells” encode an animal’s current location within an environment, neurons called “grid cells” in the entorhinal cortex have hexagonally arranged firing fields, and other neurons encode head orientation [9]. Similar to findings from electrophysiological recordings in rodents, fMRI analyses of humans navigating in virtual environments reveal grid-like signals in the entorhinal cortex during imagined navigation, with activity patterns exhibiting six-fold rotational symmetry similar to rodents [10]. In a recent study, food deprived rats were trained to run laps on a linear track to acquire a food reward while activity of scores of neurons in the CA1 region of the hippocampus were

recorded [11]. The study focused on ‘hippocampal replays’ which are episodes of sequential activity in place cells during sharp-wave ripple oscillations. The data reveal both forward and reverse replays of food locations. Interestingly, reverse replays are increased with increasing food rewards, whereas forward replays are unchanged [12]. In the natural environment, reverse replays might therefore be expected to play a key role in food patch-leaving decisions.

Studies of food caching behaviors in corvid species (members of the crow family) reveal a remarkable breadth of cognitive abilities that include remembering the past, planning for the future, and interpreting the behaviors of others. Corvids and parrots have forebrain neuron counts even greater than primates with much larger brains, which may contribute to the advanced intelligence of the birds [13, 14]. As with primates, corvids have a prominent hippocampal formation that plays key roles in the generation of ‘cognitive maps’ and episodic memory, and interconnected brain structures involved in executive functions (goal-oriented planning, strategizing and self-monitoring) and decision-making. Moreover, corvids exhibit “theory of mind”, in that they understand that other individuals have mental states and intents similar to their own [15]. Studies of scrub jays demonstrate that they routinely use mental time travel, problem solving and social cognition to acquire, hide and recover foods (caching behavior) [16]. These birds remember what happened when and where based on a single past experience so that they can discriminate between many similar previous episodes. Thus, they can learn the “shelf life” of foods and eat the more perishable foods first.

There is also evidence that expansion of the PFC during the evolution of non-human primates enabled the critical decision-making required for success in foraging in arboreal canopies [17]. Food sources are usually distributed in patches, which forces the animal to decide between staying in a patch in which food is depleted and leaving that patch to search for a different food-rich patch. It’s been reported that travel time between patches informs leaving decisions by changing the threshold and/or the rate at which the decision variable grows [18]. In this study, activity of neurons in the anterior cingulate cortex was recorded while the monkeys experienced diminishing amounts of a juice reward and were faced with the decision of “leaving the patch” to obtain a larger juice reward after a time delay. Neurons responded every time the monkeys made a choice and the firing frequency increased with time in the current patch. The monkeys abandoned the patch when a threshold level of neuronal activity was reached, and the threshold was associated with a particular travel time. How information encoded in cingulate cortex is integrated with neuronal networks in the visual cortex, prefrontal cortex, hippocampus and motor cortex to regulate foraging decisions remains to be determined. However, such findings in non-human primates support the notion that the increase in overall brain size and expansion of the prefrontal, visual and parietal cortices during primate evolution was driven in part by food scarcity [17].

Food-Centric Origins of Human Creativity, Language and Culture

Here I encapsulate evidence supporting the notion that advanced cognitive capabilities of the human brain – tool manufacture, creativity, social intelligence and language – originally evolved, at least in part, as adaptations that enabled successful food acquisition. As reviewed

by Chang et al., major expansion of the human brain coincided with new behaviors that enabled acquisition of higher calorie diets (e.g., meat and cooking). I quote: “These observations suggest the possibility that social information and information about primary motivators like food are translated into a common framework or currency that drives both learning and decision making.” [19]. However, in addition to cooperation in food acquisition and sharing the evolution of sociality and large group living is associated with several other factors that may have contributed to the evolution of large brains with advanced cognitive repertoires. For example, cooperation amongst individuals in large groups reduces the risk of predation [20]. For readers interested in the importance of social pressures for the evolution of large brain size, several excellent review articles are available [19, 21].

Findings from the field of anthropology establish that most if not all of the early tools used by our human ancestors were created for purposes of food acquisition and processing. These include not only flaked stones, spears, and bows and arrows, but also fire, the wheel, and animal and plant domestication [22, 23]. Advances in the design and production of tools for hunting and foraging and food preparation set the stage for the agricultural revolution, a remarkable period of cultural evolution that facilitated the diversification of human ingenuity and the emergence of philosophy, science and advanced technologies. Brain regions involved in toolmaking expanded prominently during human evolution including visual association cortices, PFC, posterior parietal cortex and insular cortex. These brain regions also play critical roles in communication and complex social interactions. Indeed, it is believed that toolmaking was a precursor to the development of language and cooperation in food acquisition and distribution in large societies [24, 25]. Even human artistic expression has roots in mental representations of food-centric objects and events as demonstrated by the earliest known cave drawings that depict scenes with game animals or domesticated food animals. Therefore, it appears that the neuronal network-based mechanisms of human creativity that evolved, in part, to cope with food scarcity were sufficient to enable creative cognition in realms unrelated to food acquisition and processing.

In sum, there is ample evidence that the need to acquire food played a major role in the evolution of the cognitive repertoire of many species including the superior creativity and decision-making capabilities of humans. A question that arises from the evolutionary perspective is whether the food-replete environments of modern-day humans and domesticated animals affect their cognitive capabilities, and transgenerational trajectories of brain evolution in their descendants.

Use it or Lose it: Food Overabundance, Suboptimal Cognition and the Shrinking Brain

An understanding how food scarcity-based adaptations drove human brain evolution provides insight into how removal of this environmental challenge affects cognition and cellular neuroplasticity. One clue comes from data showing that the overall brain size of domesticated animals is reduced compared to the wild species from which they originated, including farm animals raised for meat production (chickens, pigs and cattle) and pets (dogs and cats) [26–28]. A common environmental factor that might account for the reduction in

brain size is that domesticated animals no longer have to devote cognitive and physical exertion to acquire food. It would therefore be of interest to compare cognitive capabilities of different breeds within a species of domesticated animals in relation to their ad libitum food intake. However, factors other than or in addition to living in a food replete environment may also have contributed to the reduced brain size of domesticated animals perhaps including selection for docility. Nevertheless, empirical evidence from human cross sectional studies and human and animal interventional studies which is summarized below demonstrates that excessive energy intake reduces, while intermittent energy restriction increases, neuroplasticity, cognitive performance in multiple domains, and brain regional gray matter volumes. Similar evidence linking other traits of domesticated animals (docility, floppy ears, fur features, etc.) to cognition is lacking.

Remarkably, measurements of cranial volumes suggest that there has been a ~10% reduction in brain volume in humans during the past 10,000 years which corresponds to the time period after the agricultural revolution and the development of effort-sparing technologies [29]. It has been suggested that humans have undergone “self-domestication”, a process of cultural evolution which selected for the trait of cooperativity in food acquisition, processing and distribution [30]. Moreover, whereas being fluent in both spoken and written language is a requirement for essentially all occupations in modern societies, being adept at critical decision-making while moving through complex physical environments is largely unnecessary. Although we cannot know for certain because intact brains are lacking from humans who died prior to 10,000 years ago, it might be expected that brain regions previously used extensively on a day-to-day basis for navigation and associated cognitive processing required for foraging and hunting (motor cortex, temporal and frontal lobes) have decreased, whereas those regions used for language and abstract thought have increased in size and/or synaptic complexity. Consistent with the latter possibility, cross sectional MRI-based studies of children and adolescents have shown that temporal and frontal cortex volumes are positively correlated with performance IQ, but not with verbal IQ [31].

Removal of a major driving force for brain evolution from modern societies might be expected to influence brain development and plasticity within an individual, as well as in their descendants. Data from brain imaging studies of human subjects and interventional studies of laboratory animals support the latter possibilities. Obesity and insulin resistance increase the risk for cognitive impairment and Alzheimer’s disease as individuals transit their 6th – 8th decades of life [32]. Moreover, chronic excessive energy intake is associated with impaired cognition and reduced temporal lobe gray matter volume and poorer clinical outcomes in patients with major psychiatric disorders including depression [33]. Abdominal obesity is associated with lower total gray matter volume and reduced hippocampal volume in adults with type 2 diabetes [34]. Of particular concern is the negative impact of metabolic morbidity resulting from overindulgent sedentary lifestyles on the brains of children. Young children and adolescents who are obese exhibit impaired cognitive function in multiple domains, and have poorer academic and occupational achievement compared to their normal weight classmates [35–39]. In the United States, the states with the highest prevalence of childhood obesity also have the lowest percentages of high school and college graduates. Interestingly, emerging evidence suggests that a chronic positive energy balance has contributed to the recent increase in the incidence of autism. The increase in childhood

obesity during the past 40 years tracks remarkably closely with the increase in autism incidence. Moreover, obesity reduces brain-derived neurotrophic factor (BDNF) production, and children with a genetic reduction (haploinsufficiency) of BDNF expression score lower on tests of cognition and exhibit more autistic behaviors compared to age-matched children with two functional BDNF genes [40]. Exercise is effective in reducing behavioral symptoms and improving academic performance in children with autism [41, 42].

The association of obesity with poorer cognitive outcomes in humans begs the question of whether food overconsumption is sufficient to impair cognition. Studies of rodents support this possibility. Hippocampal volume is reduced in overfed sedentary rodents, in part by reductions in neurogenesis, dendritic arborization and synaptic density [43]. Mice genetically engineered to eat excessively (leptin receptor mutant mice) exhibit impaired hippocampus-dependent spatial learning and memory, and impaired recall of novel objects [43]. Their cognitive deficits are associated with impaired long-term potentiation at perforant path – hippocampal dentate granule neuron synapses, and reduced hippocampal neurogenesis. The mechanisms by which excessive energy intake and being overweight compromise cognition and neuroplasticity have been elucidated in animal studies (Figure 1). As reviewed in detail elsewhere, these mechanisms include: accumulation of oxidative damage to proteins, lipids and DNA in brain cells; inflammation (microglial activation and pro-inflammatory cytokine production); impaired mitochondrial function; impaired synaptic plasticity; reduced neurogenesis; impaired autophagy; and reduced cellular stress resistance [2, 32]. Cell membrane-associated oxidative stress impairs neuronal ion-motive ATPases (Na⁺ and Ca²⁺ pumps) and glucose transport [44] which likely contribute to the dysregulation of neuronal network activity and reduced neuronal glucose utilization observed in overweight human subjects [45]. Chronic overeating also impairs signaling pathways involved in synaptic plasticity and learning memory. Notably, BDNF expression is reduced in the hippocampus and cerebral cortex of overfed and diabetic laboratory animals [2, 46]. The mechanism involves glucocorticoid-mediated suppression of BDNF expression [47, 48]. In addition, diet-induced obesity was reported to cause synaptic stripping by microglia consistent with a role for abnormal activation of the innate immune system [49].

While the impact of sedentary overindulgent lifestyles on brain development and cognitive trajectories in children is clearly an issue of great concern, so too are emerging findings suggesting that the offspring of overnourished mothers and fathers are predisposed to obesity and poorer cognitive outcomes (Box 1). The underlying molecular mechanisms involve epigenetic modifications such as DNA methylation and histone acetylation that alter the expression of genes involved in neuroplasticity (Box 1; Figure 2). An analysis of demographic data from the Centers for Disease Control [50] reveals a significant positive correlation between childhood obesity and academic achievement amongst different states in America. While socioeconomic factors undoubtedly influence academic performance, it should also be considered that being overweight and sedentary contributes to suboptimal cognition in those children. Indeed, as demonstrated in interventional studies, regular exercise and dietary counseling can improve academic outcomes [51].

The correlations between obesity and cognitive outcomes in humans, together with the evidence that excessive food intake impairs synaptic plasticity and cognition in animal

models, suggests that a return to more evolutionarily normal intermittent eating patterns might enhance neuroplasticity and cognition. Recent findings support this possibility.

Intermittent Metabolic Challenges Enhance Cognition and Neuroplasticity

The typical eating pattern in modern societies of three meals plus snacks every day belies the fact that humans are adapted over millions of years of evolutionary history to sporadic eating patterns. The evolutionary pressure of food scarcity resulted in selection for individuals whose cognitive capabilities were heightened in a food deprived state, suggesting that cognition of modern day humans might be enhanced by a change from eating three meals plus snacks every day to an intermittent fasting (IF) eating pattern. IF eating patterns incorporate periods of time with little or no food intake that typically range from 16 – 48 hours with a frequency of the fasting period ranging from daily to once weekly [52, 53]. From physiological and neurobiological perspectives, such fasting periods are of sufficient length to deplete liver glycogen (glucose) stores. When liver glycogen stores are depleted, fatty acids are released from adipose cells into the circulation, and the fatty acids are then metabolized to ketone bodies in the liver. The two ketones produced, β -hydroxybutyrate (BHB) and acetoacetate, are then transported into the brain where they provide an energy source for neurons. An elevation of circulating ketones provides evidence that the metabolic switch from glucose to ketones has occurred. It typically takes at least 12 hours to deplete liver glycogen stores in someone who is relatively sedentary. However, vigorous exercise can accelerate the onset of ketogenesis and can increase the magnitude of ketone production when the exercise is initiated after the metabolic switch has occurred [54]. As reviewed elsewhere [2], multiple signaling pathways are engaged by intermittent switching between fasted and fed states that together enhance neuroplasticity, cognition and neuronal resilience.

Studies of rodents have shown that IF and running enhance cognition and motor system function by mechanisms involving structural and functional synaptic plasticity, and activation of signaling pathways that bolster neuronal bioenergetics and cellular stress resistance (Box 2) [2, 55, 56]. Learning and remembering previously traveled routes to a food source is associated with structural changes in neuronal circuits, most notably the formation of new synapses and neurogenesis in the hippocampus [57, 58]. Both IF and running enhance spatial learning and memory which is associated with increased density of dendritic spines in hippocampal dentate granule neurons and increased expression of BDNF [43, 46, 59]. Running and IF also stimulate hippocampal neurogenesis resulting in the production of new granule neurons that integrate into the existing hippocampal circuitry [59, 60]. This overlap in the cellular and molecular mechanisms by which exercise and IF enhance neuroplasticity and neuronal resilience likely stems from the fact that they are both bioenergetic challenges that occur coincidentally in food-sparse environments. Both cognitive and physical exertion in the food deprived/fasted state were required for survival. The findings from animal studies suggest that neuroplasticity is enhanced by exposure to conditions that mimic an environment with sparse food sources and the need to travel considerable distances to acquire the food.

Emerging findings suggest that IF enhances neuroplasticity and cognition via signals emanating from the periphery, as well as by activating brain-intrinsic signaling pathways

(Figure 1). One prominent peripheral signal produced during fasting is the ketone BHB which stimulates the production of BDNF by neurons. BDNF plays critical roles in learning and memory, synaptic plasticity and hippocampal neurogenesis, and enhances neuronal stress resistance [61, 62]. Mice adapted to IF exhibit improved regulation of neuronal network activity, effectively enhancing GABAergic tone [63, 64]. BHB may play a role in the latter effect of IF because ketogenic diets can prevent seizures in epilepsy patients [86]. IF and running may also bolster neuronal bioenergetics by stimulating mitochondrial biogenesis resulting in an increased number of healthy mitochondria in neurons [65]. Interestingly, the combination of IF with running wheel exercise results in elevations of BDNF expression and enhances synaptic plasticity by amounts beyond that which occurs with IF or exercise alone [46].

Controlled trials aimed at determining whether IF improves cognition in human subjects have not yet been reported. However, daily calorie restriction by an amount that triggers the metabolic switch to ketones has been reported to improve several cognitive domains (Box 2). In addition, IF results in significant elevations of circulating BHB levels on the fasting days in human subjects [66, 67]. Assuming that BHB induces cerebral BDNF expression in humans, it would be expected that IF enhances synaptic plasticity and cognition in humans. An ongoing clinical trial of IF (fasting 2 days each week) in subjects at risk for cognitive impairment (obese, insulin-resistant subjects between the ages of 55 and 70) may provide initial answers regarding the translatability of findings in animals described above [68].

While enhancement of neuroplasticity by patchy food sources and physical exertion may enable success in food acquisition, these bioenergetic challenges can also enhance neuronal resilience and protect the brain against injury and disease. As evidence, IF reduces neuronal degeneration and improves functional outcome in rodent models of ischemic stroke, spinal cord injury and traumatic brain injury [69–73]. In rats, IF lessens hippocampal neuron dysfunction and degeneration, and preserves hippocampus-dependent learning and memory in an experimental model of epilepsy [74]. Rodents maintained on IF exhibit reduced neuronal degeneration and improved functional outcomes in animal models of Alzheimer's, Parkinson's and Huntington's diseases [75–77]. Daily energy restriction also attenuates depletion of dopamine and improves functional outcome in a rhesus monkey model of Parkinson's disease [78]. Although controlled trials of IF in humans with or at risk for neurological disorders are as yet lacking, there is evidence that individuals with high energy intakes are at risk for Alzheimer's disease, perhaps because of a relative lack of activation of pathways involved in adaptive neuroplasticity [32]. The potential benefits of IF on cognition, mood and academic achievement in obese and autistic children also remain to be tested.

Concluding Comments

It has become clear that being overweight and sedentary adversely affects essentially all organ systems including the brain. As briefly described above and recently reviewed in detail [32, 79], an overarching reason for this is that cells, tissues and organs become 'complacent' when they are not subjected to intermittent metabolic challenges. Signaling pathways involved in adaptive cellular plasticity are down-regulated, organ function becomes impaired and disease processes are enabled. As briefly reviewed herein, much of

the neuronal cytoarchitecture of the brains of animals and humans was ‘sculpted’ by the evolutionary ‘chisel’ of competition for limited and sporadic food availability.

A better understanding of the bioenergetic processes and regulatory signaling pathways that mediate the enhancement of cognitive performance in response to dietary energy restriction and exercise will likely lead to the development of novel approaches for promoting optimal cognition throughout the life course. The metabolic shift to ketogenesis is one such evolutionarily ancient adaptation that can enhance cognition [2]. Although there has been a recent flurry of translational research on IF and general health (reviewed in [53]), its impact on brain health in humans remains to be interrogated in randomized controlled trials. The question of what regimens of IF eating patterns and exercise promote optimal cognition might be answered by such trials.

Finally, it is of more than of academic interest to consider how living in the food-replete niches of modern societies will impact the future evolution of the human brain. Throughout most of human evolution critical thinking and decision-making occurred as individuals navigated through a heterogenous environment hunting and foraging in a fasted state. The latter conditions might be expected to engage more brain regions compared to modern day occupations that are relatively devoid of body movement and multi-modal sensory input, and are performed in the fed state. The reduction of the cognitive load required for successful hunting or foraging is likely a major factor in the reduction in overall brain size in domesticated animals, and a similar scenario may be occurring in humans. While increased specialization among and within occupations may enable more rapid progress and efficiency of very large societies, it might also foster, over a period of relatively few generations, a reduction in the cognitive repertoire of each individual person. If and to what extent “renaissance men and women” bolstered by lifestyles that include IF and exercise will shape our future remains to be determined.

Acknowledgements:

This work was supported by the Intramural Research Program of the National Institute on Aging.

Glossary

Brain-derived neurotrophic factor (BDNF)

a protein that is produced by neurons in response to bioenergetic challenges, and that plays key roles in synaptic plasticity and learning and memory.

Corvid

a species of bird in the crow family.

Creativity

the ability to mentally manipulate information in ways that generate new ideas that can then be implemented to produce novel objects or plans of action.

Epigenetic modifications

molecular modifications of DNA, histones and other chromatin-associated proteins that can result in changes in gene expression; examples include methylation, acetylation and ubiquitination.

Grid cells

neurons in the entorhinal cortex with firing fields that form a regularly spaced hexagonal or triangular grid pattern.

Intelligence

the ability to acquire or infer, and retain information, and then apply it towards adaptive behaviors within an environment or occupation.

Intermittent fasting (IF)

a feeding pattern that includes periods of time of sufficient length to deplete liver glycogen stores and elevate blood ketone levels.

Ketogenic state

a metabolic state that occurs during extended food deprivation or fasting in which liver glycogen stores have been depleted, and fatty acids mobilized from adipose cells are used to produce ketones (β -hydroxybutyrate and acetoacetate); the ketones are used by neurons to sustain their bioenergetic demands.

Mental time travel

the ability to recount one's past and plan for the future.

Neurogenesis

the process by which new neurons are generated from self-renewing stem cells.

Place cells

neurons in the hippocampus that fire selectively when an animal is at one or only a few locations in its environment; place cells are prominent in the hippocampus.

Prefrontal cortex

the most anterior region of the frontal cortex which plays a major role in decision making; it uses information about the current behavioral context to rapidly generate goals based on the current biological needs.

Self-domestication

an evolutionary process that selects for individuals that cooperate with others to enhance group cohesiveness and the allocation of food and other vital resources.

Social intelligence

the awareness and active interpretation of the ongoing and likely future behaviors of oneself and others.

References

1. Lemaître JF et al. (2015) Early-late life trade-offs and the evolution of ageing in the wild. *Proc Biol Sci.* 282, 20150209. [PubMed: 25833848]

2. Mattson MP et al. (2018) Intermittent metabolic switching, neuroplasticity and brain health. *Nat Rev Neurosci.* 19, 63–80.
3. Sanders H et al. (2015) Grid Cells and Place Cells: An Integrated View of their Navigational and Memory Function. *Trends Neurosci.* 38, 763–775. [PubMed: 26616686]
4. Rickgauer JP et al. (2014) Simultaneous cellular-resolution optical perturbation and imaging of place cell firing fields. *Nat Neurosci.* 17, 1816–1824. [PubMed: 25402854]
5. Schmidt-Hieber C and Häusser M (2013) Cellular mechanisms of spatial navigation in the medial entorhinal cortex. *Nat Neurosci.* 16, 325–331. [PubMed: 23396102]
6. Harvey CD et al. (2009) Intracellular dynamics of hippocampal place cells during virtual navigation. *Nature* 461, 941–946. [PubMed: 19829374]
7. Youngstrom IA and Stowbridge BW (2012) Visual landmarks facilitate rodent spatial navigation in virtual reality environments. *Learn Mem.* 19, 84–90. [PubMed: 22345484]
8. Sato M et al. (2017) Hippocampus-dependent goal localization by head-fixed mice in virtual reality. *eNeuro.* 2017; 4(3).
9. Moser M et al. (2015) Place cells, grid cells, and memory. *Cold Spring Harbor Perspect Biol.* 7, a021808.
10. Horner AJ et al. (2016) Grid-like processing of imagined navigation. *Curr Biol.* 26, 842–847. [PubMed: 26972318]
11. Pfeiffer BE and Foster DJ (2013) Hippocampal place-cell sequences depict future paths to remembered goals. *Nature* 497, 74–79. [PubMed: 23594744]
12. Ambrose RE et al. (2016) Reverse replay of hippocampal place cells is uniquely modulated by changing reward. *Neuron* 91, 1124–1136. [PubMed: 27568518]
13. Iwaniuk AN (2005) Interspecific allometry of the brain and brain regions in parrots (psittaciformes): comparisons with other birds and primates. *Brain Behav Evol.* 65, 40–59. [PubMed: 15467290]
14. Olkowitz S et al. (2016) Birds have primate-like numbers of neurons in the forebrain. *Proc Natl Acad Sci U S A.* 113, 7255–7260. [PubMed: 27298365]
15. Clayton NS (2015) Ways of thinking: from crows to children and back again. *Q J Exp Psychol (Hove).* 68, 209–241. [PubMed: 25207993]
16. Clayton NS, and Emery NJ (2015) Avian models for human cognitive neuroscience: a proposal. *Neuron* 86, 1330–1342. [PubMed: 26087161]
17. Passingham RE and Wise SP (2012) *The Neurobiology of the Prefrontal Cortex.* Oxford University Press 399 pp.
18. Hayden BY et al. (2011) Neuronal basis of sequential foraging decisions in a patchy environment. *Nat Neurosci.* 14, 933–939. [PubMed: 21642973]
19. Chang SW et al. (2013) Neuroethology of primate social behavior. *Proc Natl Acad Sci U S A.* 110 Suppl 2, 10387–10394. [PubMed: 23754410]
20. van der Bijl W, Kolm N (2016) Why direct effects of predation complicate the social brain hypothesis: And how incorporation of explicit proximate behavioral mechanisms might help. *Bioessays.* 38, 568–577. [PubMed: 27174816]
21. Tremblay S, Sharika KM and Platt ML (2017) Social decision-making and the brain: a comparative perspective. *Trends Cogn Sci.* 21, 265–276. [PubMed: 28214131]
22. Plummer T (2004) Flaked stones and old bones: biological and cultural evolution at the dawn of technology. *Am J Phys Anthropol. Suppl* 39, 118–164. [PubMed: 15605391]
23. Vigne JD (2011) The origins of animal domestication and husbandry: a major change in the history of humanity and the biosphere. *C R Biol.* 334, 171–181. [PubMed: 21377611]
24. Ruck L (2014) Manual praxis in stone tool manufacture: implications for language evolution. *Brain Lang.* 139, 68–83. [PubMed: 25463818]
25. Kolodny O and Edelman S (2018) The evolution of the capacity for language: the ecological context and adaptive value of a process of cognitive hijacking. *Philos Trans R Soc Lond B Biol Sci.* 373 (1743).
26. Schleifenbaum C (1973) The postnatal development of the brain of poodles and wolves (author's transl). *Z Anat Entwicklungsgesch.* 141, 179–205. [PubMed: 4769553]

27. Hemmer H (1990). *Domestication. The Decline of Environmental Appreciation*. Cambridge: Cambridge University Press Cambridge, UK.
28. Kawabe S et al. (2017) Morphological variation in brain through domestication fowl. *J Anat.* 231, 287–297. [PubMed: 28542781]
29. Henneberg M and Steyn M (1993) Trends in cranial capacity and cranial index in Subsaharan Africa during the Holocene. *Am J Hum Biol.* 5, 473–479. [PubMed: 28548397]
30. Hare B (2017) Survival of the friendliest: *Homo sapiens* evolved via selection for prosociality. *Annu Rev Psychol.* 68, 155–186. [PubMed: 27732802]
31. Lange N et al. (2010) Associations between IQ, total and regional brain volumes and demography in a large normative sample of healthy children and adolescents. *Dev Neuropsychol.* 35, 296–317. [PubMed: 20446134]
32. Mattson MP and Arumugam TV (2018) Hallmarks of brain aging: adaptive and pathological modification by metabolic states. *Cell Metab.* 27, 1176–1199. [PubMed: 29874566]
33. Hidese S et al. (2018) Association of obesity with cognitive function and brain structure in patients with major depressive disorder. *J Affect Disord.* 225, 188–194. [PubMed: 28837952]
34. Climie RE et al. (2015) Abdominal Obesity and Brain Atrophy in Type 2 Diabetes Mellitus. *PLoS One.* 10, e0142589. [PubMed: 26560876]
35. Kamijo K et al. (2014) The negative association of childhood obesity to cognitive control of action monitoring. *Cereb Cortex.* 24, 654–662. [PubMed: 23146965]
36. Liang J et al. (2014) Neurocognitive correlates of obesity and obesity-related behaviors in children and adolescents. *Int J Obes (Lond).* 38, 494–506. [PubMed: 23913029]
37. Kennedy JT et al. (2016) Higher adolescent body mass index is associated with lower regional gray and white matter volumes and lower levels of positive emotionality. *Front Neurosci.* 10, 413. [PubMed: 27660604]
38. Akin O et al. (2017) Neurocognitive functions, particularly verbal abilities, were impaired in obese children with IR. *J Pediatr Endocrinol Metab.* 30, 1027–1032. [PubMed: 28841575]
39. Mestre ZL et al. (2017) Hippocampal atrophy and altered brain responses to pleasant tastes among obese compared with healthy weight children. *Int J Obes (Lond).* 2017; 41:1496–1502. [PubMed: 28572588]
40. Han JC et al. (2013) Association of brain-derived neurotrophic factor (BDNF) haploinsufficiency with lower adaptive behaviour and reduced cognitive functioning in WAGR/11p13 deletion syndrome. *Cortex* 49, 2700–2710. [PubMed: 23517654]
41. Oriol KN et al. (2011) The effects of aerobic exercise on academic engagement in young children with autism spectrum disorder. *Pediatr Phys Ther.* 23, 187–193. [PubMed: 21552085]
42. Pan CY et al. (2017) The impacts of physical activity intervention on physical and cognitive outcomes in children with autism spectrum disorder. *Autism.* 21, 190–202. [PubMed: 27056845]
43. Stranahan AM, Norman ED, Lee K, Cutler RG, Telljohann RS, Egan JM, Mattson MP (2008) Diet-induced insulin resistance impairs hippocampal synaptic plasticity and cognition in middle-aged rats. *Hippocampus.* 18, 1085–1088. [PubMed: 18651634]
44. Mattson MP (2009) Roles of the lipid peroxidation product 4-hydroxynonenal in obesity, the metabolic syndrome, and associated vascular and neurodegenerative disorders. *Exp Gerontol.* 44, 625–633. [PubMed: 19622391]
45. Guzzardi MA and Iozzo P (2018) Brain functional imaging in obese and diabetic patients. *Acta Diabetol.* 2018 Jun 29. doi: 10.1007/s00592-018-1185-0. [Epub ahead of print].
46. Stranahan AM et al. (2009) Voluntary exercise and caloric restriction enhance hippocampal dendritic spine density and BDNF levels in diabetic mice. *Hippocampus.* 19, 951–961. [PubMed: 19280661]
47. Stranahan AM, Arumugam TV, Cutler RG, Lee K, Egan JM, Mattson MP (2008) Diabetes impairs hippocampal function through glucocorticoid-mediated effects on new and mature neurons. *Nat Neurosci.* 11, 309–317. [PubMed: 18278039]
48. Wosiski-Kuhn M, Erion JR, Gomez-Sanchez EP, Gomez-Sanchez CE, Stranahan AM (2014) Glucocorticoid receptor activation impairs hippocampal plasticity by suppressing BDNF expression in obese mice. *Psychoneuroendocrinology.* 42, 165–177. [PubMed: 24636513]

49. Hao S, Dey A, Yu X, Stranahan AM (2016) Dietary obesity reversibly induces synaptic stripping by microglia and impairs hippocampal plasticity. *Brain Behav Immun.* 51, 230–239. [PubMed: 26336035]
50. <https://www.cdc.gov/healthyschools/obesity/obesity-youth.htm>
51. Bustamante EE, Williams CF, Davis CL (2016) Physical activity interventions for neurocognitive and academic performance in overweight and obese youth: a systematic review. *Pediatr Clin North Am.* 63, 459–480. [PubMed: 27261545]
52. Anton SD et al. (2018) Flipping the metabolic switch: understanding and applying the health benefits of fasting. *Obesity (Silver Spring).* 26, 254–268. [PubMed: 29086496]
53. Mattson MP, Longo VD, Harvie M (2017) Impact of intermittent fasting on health and disease processes. *Ageing Res Rev.* 39, 46–58. [PubMed: 27810402]
54. Marosi K et al. (2018) Metabolic and molecular framework for the enhancement of endurance by intermittent food deprivation. *FASEB J.* 32, 3844–3858. [PubMed: 29485903]
55. Ingram DK et al. (1987) Dietary restriction benefits learning and motor performance of aged mice. *J Gerontol* 42, 78–81. [PubMed: 3794202]
56. Mattson MP (2012) Energy intake and exercise as determinants of brain health and vulnerability to injury and disease. *Cell Metab.* 16, 706–722. [PubMed: 23168220]
57. Leuner B and Shors TJ (2004) New spines, new memories. *Mol Neurobiol.* 29, 117–130. [PubMed: 15126680]
58. Segal M (2017) Dendritic spines: Morphological building blocks of memory. *Neurobiol Learn Mem.* 138, 3–9. [PubMed: 27311757]
59. Vivar C and van Praag H (2017) Running changes the brain: the long and the short of it. *Physiology (Bethesda).* 32, 410–424. [PubMed: 29021361]
60. Lee J et al. (2002) Evidence that brain-derived neurotrophic factor is required for basal neurogenesis and mediates, in part, the enhancement of neurogenesis by dietary restriction in the hippocampus of adult mice. *J Neurochem.* 2002; 82:1367–1375. [PubMed: 12354284]
61. Marosi K et al. (2016) 3-Hydroxybutyrate regulates energy metabolism and induces BDNF expression in cerebral cortical neurons. *J Neurochem.* 139, 769–781. [PubMed: 27739595]
62. Sleiman SF et al. (2016) Exercise promotes the expression of brain derived neurotrophic factor (BDNF) through the action of the ketone body β -hydroxybutyrate. *Elife.* 2016 Jun 2; 5 pii: e15092.
63. Liu Y et al. (2018) under review.
64. Neal EG et al. (2008) The ketogenic diet for the treatment of childhood epilepsy: a randomised controlled trial. *Lancet Neurol.* 7, 500–506. [PubMed: 18456557]
65. Cheng A et al. (2012) Involvement of PGC-1 α in the formation and maintenance of neuronal dendritic spines. *Nat Commun.* 3, 1250. [PubMed: 23212379]
66. Johnson JB et al. (2007) Alternate day calorie restriction improves clinical findings and reduces markers of oxidative stress and inflammation in overweight adults with moderate asthma. *Free Radic Biol Med.* 42, 665–674. [PubMed: 17291990]
67. Harvie M et al. (2013) The effect of intermittent energy and carbohydrate restriction v. daily energy restriction on weight loss and metabolic disease risk markers in overweight women. *Br J Nutr.* 110, 1534–1547. [PubMed: 23591120]
68. Kapogiannis, and Mattson MP (2018) (<https://clinicaltrials.gov/ct2/show/NCT02460783?term=kapogiannis&rank=1>)
69. Yu ZF and Mattson MP (1999) Dietary restriction and 2-deoxyglucose administration reduce focal ischemic brain damage and improve behavioral outcome: evidence for a preconditioning mechanism. *J Neurosci Res.* 57, 830–839. [PubMed: 10467254]
70. Arumugam TV et al. (2010) Age and energy intake interact to modify cell stress pathways and stroke outcome. *Ann Neurol.* 67, 41–52. [PubMed: 20186857]
71. Plunet WT et al. (2008) Dietary restriction started after spinal cord injury improves functional recovery. *Exp Neurol.* 213, 28–35. [PubMed: 18585708]
72. Davis LM et al. (2008) Fasting is neuroprotective following traumatic brain injury. *J Neurosci Res.* 86, 1812–1822. [PubMed: 18241053]

73. Rich NJ et al. (2010) Chronic caloric restriction reduces tissue damage and improves spatial memory in a rat model of traumatic brain injury. *J Neurosci Res.* 88, 2933–2939. [PubMed: 20544832]
74. Bruce-Keller AJ et al. (1999) Food restriction reduces brain damage and improves behavioral outcome following excitotoxic and metabolic insults. *Ann Neurol.* 45, 8–15. [PubMed: 9894871]
75. Halagappa VK et al. (2007) Intermittent fasting and caloric restriction ameliorate age-related behavioral deficits in the triple-transgenic mouse model of Alzheimer's disease. *Neurobiol Dis.* 26, 212–220. [PubMed: 17306982]
76. Duan W and Mattson MP (1999) Dietary restriction and 2-deoxyglucose administration improve behavioral outcome and reduce degeneration of dopaminergic neurons in models of Parkinson's disease. *J Neurosci Res.* 57, 195–206. [PubMed: 10398297]
77. Duan W et al. (2003) Dietary restriction normalizes glucose metabolism and BDNF levels, slows disease progression, and increases survival in huntingtin mutant mice. *Proc Natl Acad Sci U S A.* 100, 2911–2916. [PubMed: 12589027]
78. Maswood N et al. (2004) Caloric restriction increases neurotrophic factor levels and attenuates neurochemical and behavioral deficits in a primate model of Parkinson's disease. *Proc Natl Acad Sci U S A.* 101, 18171–18176. [PubMed: 15604149]
79. Stranahan AM, Mattson MP (2012) Recruiting adaptive cellular stress responses for successful brain ageing. *Nat Rev Neurosci.* 13, 209–216. [PubMed: 22251954]
80. Dunn GA, Bale TL (2009) Maternal high-fat diet promotes body length increases and insulin insensitivity in second-generation mice. *Endocrinology* 150, 4999–5009. [PubMed: 19819967]
81. Bruce-Keller AJ, Fernandez-Kim SO, Townsend RL, Kruger C, Carmouche R, Newman S, Salbaum JM, Berthoud HR (2017) Maternal obese-type gut microbiota differentially impact cognition, anxiety and compulsive behavior in male and female offspring in mice. *PLoS One.* 12, e0175577. [PubMed: 28441394]
82. Kim TW, Park HS (2018) Physical exercise improves cognitive function by enhancing hippocampal neurogenesis and inhibiting apoptosis in male offspring born to obese mother. *Behav Brain Res.* 347, 360–367. [PubMed: 29551732]
83. Wolfrum C, Peleg-Raibstein D (2018) Maternal overnutrition leads to cognitive and neurochemical abnormalities in C57BL/6 mice. *Nutr Neurosci.* 1, 1–12.
84. Kang SS, Kurti A, Fair DA, Fryer JD (2014) Dietary intervention rescues maternal obesity induced behavior deficits and neuroinflammation in offspring. *J Neuroinflammation.* 11, 156. [PubMed: 25212412]
85. Tozuka Y, Kumon M, Wada E, Onodera M, Mochizuki H, Wada K (2010) Maternal obesity impairs hippocampal BDNF production and spatial learning performance in young mouse offspring. *Neurochem Int.* 57, 235–247. [PubMed: 20538025]
86. Ravelli AC et al. (1998) Glucose tolerance in adults after prenatal exposure to famine. *Lancet* 351, 173–177. [PubMed: 9449872]
87. Neri C, Edlow AG (2015) Effects of maternal obesity on fetal programming: molecular approaches. *Cold Spring Harb Perspect Med.* 6, a026591. [PubMed: 26337113]
88. Sullivan EL et al. (2010) Chronic consumption of a high-fat diet during pregnancy causes perturbations in the serotonergic system and increased anxiety-like behavior in nonhuman primate offspring. *J Neurosci* 30, 3826–3830. [PubMed: 20220017]
89. Vucetic Z, Kimmel J, Totoki K, Hollenbeck E, Reyes TM (2010) Maternal high-fat diet alters methylation and gene expression of dopamine and opioid-related genes. *Endocrinology* 151, 4756–4764. [PubMed: 20685869]
90. Bilbo SD, Tsang V (2010) Enduring consequences of maternal obesity for brain inflammation and behavior of off-spring. *FASEB J* 24, 2104–2115. [PubMed: 20124437]
91. Sullivan EL, Smith MS, Grove KL (2011) Perinatal exposure to high-fat diet programs energy balance, metabolism and behavior in adulthood. *Neuroendocrinology* 93, 1–8. [PubMed: 21079387]
92. Basatemur E, Gardiner J, Williams C, Melhuish E, Barnes J, Sutcliffe A (2013) Maternal prepregnancy bmi and child cognition: A longitudinal cohort study. *Pediatrics* 131, 56–63. [PubMed: 23230067]

93. Casas M et al. (2013) Maternal pre-pregnancy overweight and obesity, and child neuropsychological development: Two southern european birth cohort studies. *Int. J. Epidemiol* 42, 506–517. [PubMed: 23569191]
94. Bliddal M et al. (2014) Maternal pre-pregnancy bmi and intelligence quotient (IQ) in 5-year-old children: A cohort based study. *PLoS ONE* 2014, 9, e94498. [PubMed: 24727836]
95. Tanda R, Salsberry PJ, Reagan PB, Fang MZ. 2012 The impact of prepregnancy obesity on children's cognitive test scores. *Matern Child Health J* 17: 222–229.
96. Rodriguez A, Miettunen J, Henriksen TB, Olsen J, Obel C, Taanila A, Ebeling H, Linnet KM, Moilanen I, Jarvelin MR. 2008 Maternal adiposity prior to pregnancy is associated with ADHD symptoms in offspring: Evidence from three prospective pregnancy cohorts. *Int J Obes (Lond)* 32: 550–557. [PubMed: 17938639]
97. Krakowiak P, Walker CK, Bremer AA, Baker AS, Ozonoff S, Hansen RL, Hertz-Picciotto I. 2012 Maternal metabolic conditions and risk for autism and other neurodevelopmental disorders. *Pediatrics* 129: e1121–e1128. [PubMed: 22492772]
98. Tanne JH. 2012 Maternal obesity and diabetes are linked to children's autism and similar disorders. *BMJ* 344: e2768. [PubMed: 22511212]
99. Reynolds LC, Inder TE, Neil JJ, Pineda RG, Rogers CE. 2014 Maternal obesity and increased risk for autism and developmental delay among very preterm infants. *J Perinatol* 34: 688–692. [PubMed: 24811227]
100. Edlow AG. (2017) Maternal obesity and neurodevelopmental and psychiatric disorders in offspring. *Prenat Diagn.* 37, 95–110. [PubMed: 27684946]
101. Contu L, Hawkes CA (2017) A Review of the impact of maternal obesity on the Cognitive function and mental health of the offspring. *Int J Mol Sci.* 18(5), E1093. [PubMed: 28534818]
102. Surén P et al. (2014) Parental obesity and risk of autism spectrum disorder. *Pediatrics.* 133, e1128–138. [PubMed: 24709932]
103. Zhou Y et al. (2018) Diet-induced paternal obesity impairs cognitive function in offspring by mediating epigenetic modifications in spermatozoa. *Obesity (Silver Spring).* 26, 1749–1757. [PubMed: 30358144]
104. Zhang TY et al. (2018) Environmental enrichment increases transcriptional and epigenetic differentiation between mouse dorsal and ventral dentate gyrus. *Nat Commun.* 9, 298. [PubMed: 29352183]
105. Moon HY et al. (2016) Running-Induced Systemic Cathepsin B Secretion Is Associated with Memory Function. *Cell Metab.* 24, 332–340. [PubMed: 27345423]
106. Wrann CD et al. (2013) Exercise induces hippocampal BDNF through a PGC-1 α /FNDC5 pathway. *Cell Metab.* 18, 649–59. [PubMed: 24120943]
107. Cheng A et al. (2016) Mitochondrial SIRT3 mediates adaptive responses of neurons to exercise and metabolic and excitatory challenges. *Cell Metab.* 23, 128–142. [PubMed: 26698917]
108. Fedewa AL and Ahn S (2011) The effects of physical activity and physical fitness on children's achievement and cognitive outcomes: a meta-analysis. *Res Q Exerc Sport.* 82, 521–535. [PubMed: 21957711]
109. Makharia A et al. (2016) Effect of environmental factors on intelligence quotient of children. *Ind Psychiatry J.* 25, 189–194. [PubMed: 28659699]
110. Buck SM et al. (2008) The relation of aerobic fitness to stroop task performance in preadolescent children. *Med Sci Sports Exerc.* 40, 166–172. [PubMed: 18091008]
111. Gow AJ et al. (2012) Reverse causation in activity-cognitive ability associations: the Lothian Birth Cohort 1936. *Psychol Aging* 27, 250–255. [PubMed: 21644808]
112. Steinberg H et al. (1997) Exercise enhances creativity independently of mood. *Br J Sports Med.* 31, 240–245. [PubMed: 9298561]
113. Witte AV et al. (2009) Caloric restriction improves memory in elderly humans. *Proc Natl Acad Sci U S A.* 106, 1255–1260. [PubMed: 19171901]
114. Prehn K et al. (2017) Caloric restriction in older adults-differential effects of weight loss and reduced weight on brain structure and function. *Cereb Cortex.* 27, 1765–1778. [PubMed: 26838769]

115. Horie NC et al. (2016) Cognitive effects of intentional weight loss in elderly obese individuals with mild cognitive impairment. *J Clin Endocrinol Metab.* 101, 1104–1112. [PubMed: 26713821]
116. Keifer J (2017) Primetime for learning genes. *Genes* 11, 8(2), E69. [PubMed: 28208656]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Box 1.**Transgenerational Epigenetic Impact of Excessive Food Intake on Cognition**

Findings from recent animal studies in which food intake and exercise were precisely controlled, and associational data from studies of humans, provide evidence that the metabolic status of parents can influence their offspring's risk for obesity and impaired glucose regulation. When female mice maintained on a high fat diet to induce obesity and insulin resistance are mated with healthy males, their offspring exhibit insulin resistance and obesity, and this phenotype persists through at least a second generation in both maternal and paternal lineages [80]. The adverse effect of maternal obesity on the metabolic phenotype of offspring in mice and rats is associated with impaired cognition in the offspring in reference memory and associative learning [81–83]. Spatial learning and memory deficits in male offspring of obese dams can be reversed when the offspring exercise (run) regularly [82]. Interestingly, moderate food restriction/daily IF of nursing dams that were obese during pregnancy can ameliorate behavioral deficits caused by the maternal obesity, suggesting that the early postnatal metabolic state of the mother influences cognitive trajectories of offspring [84]. Several studies have reported that hippocampal BDNF expression is reduced in offspring of overnourished dams suggesting one underlying abnormality in gene expression [82, 85]. It should be noted, however, that starvation in humans has been reported to adversely affect the metabolic outcomes of offspring [86]. Thus, moderate levels of intermittent energy restriction and physical activity may best promote healthy transgenerational metabolic phenotypes.

The adverse impact of overweight and obesity in parents on metabolic outcomes of their descendants is believed to result from epigenetic molecular changes occurring *in utero* that are then propagated transgenerationally [see 87 for review]. The *in utero* environment can have enduring effects on the regulation of gene expression resulting from epigenetic molecular mechanisms which encompass heritable modifications of the genome without changes in the DNA sequence. Prominent among such genome modifications are DNA methylation, posttranslational modifications of chromatin/histone proteins (acetylation, methylation, phosphorylation, ubiquitination) and microRNA alterations [87]. It has been reported that offspring of obese females exhibit DNA hypomethylation in the gene promoters including those encoding proteins involved in dopamine uptake, and serotonergic and opioid signaling [88, 89]. The proximate cause of such epigenetic modifications in brain cells of the developing embryo may include hyperglycemia, insulin resistance and a proinflammatory lipotoxic environment [84, 90, 91].

In utero epigenetic modifications could predispose the offspring of obese mothers to some developmental brain disorders. Multiple studies reveal a negative association of pre-pregnancy maternal obesity and child intelligence quotient (IQ) [92–94]. Children born to obese mothers perform more poorly on reading and math tests than do children born to normal weight mothers [95]. Children born to women who are obese and/or diabetic may be at increased risk for autism spectrum disorder and attention deficit hyperactivity disorder [96–99]. Although not yet conclusive, increasing evidence

suggests that maternal obesity increases the risk of several psychiatric disorders in adulthood, including depression, anxiety disorders and schizophrenia [100]. Consistent with the latter possibility, animal studies have shown that pups born to obese dams exhibit heightened and anxiety and depression-like behaviors [101]. Paternal obesity may also adversely impact cognitive and behavioral outcomes in offspring by epigenetic mechanisms [102, 103]. Collectively, the emerging data suggest that overconsumption of high-energy foods by conceiving mothers and fathers increase the risk of poor or suboptimal cognitive outcomes of their offspring.

Box 2.**Intermittent metabolic challenges boost the cellular engines of cognition**

Studies of rodents have shown that enriched environments, food deprivation and running result in increased activity in neuronal circuits involved in cognition including those in the hippocampus [56, 57]. Such increases in excitatory synaptic (glutamatergic) activity impose a mild metabolic stress on the neurons as a consequence of Na^+ and Ca^{2+} influx, increased mitochondrial metabolism and the activation of ion-motive ATPases. The Ca^{2+} influx and mitochondrial reactive oxygen species activate transcription factors that induce the expression of genes that encode proteins involved in adaptive cellular stress responses and synaptic plasticity [2, 79, 104]. Consequently, synapses involved in learning and memory are potentiated, synapse numbers increase, and neurons are better able to cope with a wide range of environmental challenges including metabolic, excitatory and oxidative stress. In response to the systemic bioenergetic challenges of fasting and exercise peripheral organs produce and release into the circulation molecules that affect neuroplasticity. These include liver-derived ketones, and muscle-derived cathepsin B and irisin [56, 105, 106]. Interestingly, all three of these peripheral signals stimulate the production of BDNF which, in turn, stimulates synaptogenesis and neurogenesis [2]. By pathways involving BDNF, the transcription factor PGC-1 α and the mitochondrial deacetylase SIRT3, exercise and fasting can increase the numbers of well-functioning stress-resistant mitochondria in hippocampal and cerebral cortical neurons to enhance neuroplasticity and resilience [65, 107]. The increased number of healthy mitochondria in neurons is believed to be critical for the enhancement of cognitive performance that occurs in response to fasting and aerobic exercise [2, 65].

Human studies have shown that cognition and intellectual achievement can be enhanced by exercise and energy restriction [108]. In a study of over 1000 schoolchildren between the ages of 12 and 16 it was found that those who engaged in physical activity more than 5 hours/week had a higher IQ compared to their more sedentary classmates [109]. There is also a positive association of aerobic fitness and cognitive processing speed in preadolescent children [110]. In a birth cohort study, general cognitive ability and processing speed at age 70 was positively associated with physical activity regardless of IQ at age 11 [111]. In a study in which adult subjects were randomly assigned to either aerobic exercise or control (video watching) groups, only those in the exercise group exhibited improvements in cognitive flexibility/creativity [112]. Dietary energy restriction can also improve cognition. Compared to their baseline scores, and to subjects in two different control diet groups, those who reduced their daily calorie intake by 30% for three months exhibited significant increases in verbal memory scores [113]. Recognition memory improved, hippocampal and inferior frontal gyrus gray matter volumes increased, and functional connectivity of hippocampal and parietal lobe networks increased in older women during 12 weeks on a very low calorie diet/daily fasting diet [114]. Similarly, improvements in verbal memory and fluency and executive function were improved in response to dietary energy restriction in overweight patients with mild cognitive impairment [115].

Highlights

- Neuronal networks in brain regions critical for spatial navigation and decision-making evolved to enable success in competition for limited food availability in hazardous environments.
- A major ecological factor that drove the evolution of cognition, namely food scarcity, has been largely eliminated from day-to-day experience of modern-day humans and domesticated animals.
- Continuous availability and consumption of energy-rich food in relatively sedentary modern-day humans negatively impacts the lifetime cognitive trajectories of themselves and their children.
- Epigenetic molecular DNA and chromatin protein modifications are impacted by energy intake and can be propagated to future generations.
- The cellular and molecular mechanisms by which intermittent food deprivation enhances cognition and overfeeding impairs cognition are being elucidated.
- A better understanding of the food-centric evolutionary foundations of human brain neuroplasticity is leading to the development of novel bioenergetic challenge-based patterns of eating and exercise aimed at improving cognitive health and resilience.

Outstanding Questions

- How might a better understanding of the impact of food scarcity on brain evolution help conceptualize human cognition in general, and spatial learning and memory and decision-making in particular?
- Did social intelligence and cooperation within large societies evolve as an adaption to overcome food scarcity as the size of societies increased? Can differences in brain structure and/or functional connectivity between current-day hunter - gatherers and individuals in modern food-replete societies be discerned?
- During evolution intense cognitive processing occurred while animals, including our human ancestors, were moving through their territories. Does running and other types physical exertion affect cognition? Which domains are affected and by what mechanisms? Does exercise in the fasted state, a condition common during the evolution of many species, amplify the effects of exercise on cognition?
- What are the cellular and molecular mechanisms by which intermittent fasting enhances brain resilience in the contexts of aging and vulnerability to neurological disorders that impact cognition? Can prescriptions for intermittent eating patterns be implemented as an approach to promoting brain health?
- How will the brain evolve as future generations of humans live in the setting of food overabundance and with sedentary occupations?
- To what extent can pharmacological interventions be developed that activate the same cognition-enhancing signaling pathways that are activated by intermittent fasting and exercise?




Figure 1. Cellular and molecular mechanisms by which food intake impacts neuroplasticity and cognition.

Left) Adaptive responses of neuronal networks to intermittent food deprivation or fasting. Extended periods with no or little energy intake triggers a metabolic shift from utilization of liver glycogen-derived glucose to adipose cell-derived fatty acids and ketone bodies (BHB, β -hydroxybutyrate; AcAc, acetoacetate) generated therefrom. In addition to serving as a source of acetyl CoA for mitochondrial ATP production, ketone bodies can activate signaling pathways involved in synaptic plasticity and cellular stress resistance, including those involving the transcription factors CREB (cyclic AMP response element binding protein) and NF- κ B (nuclear factor kappa B), and neurotrophic factors such as BDNF (brain-derived neurotrophic factor). The increased activity in neuronal networks involved in cognitive processing during food seeking (navigation, decision-making, etc.) engages adaptive signaling pathways that bolster mitochondrial function and up-regulate neurotrophic factors, GABAergic tone, antioxidant defenses and DNA repair, while

suppressing inflammation. These adaptive responses promote synaptic plasticity, neurogenesis and cellular stress resistance which, in turn enhances cognition and resistance of the brain to injury and disease. **Right** Excessive food intake as occurs in laboratory animals fed ad libitum and most humans in modernized countries impairs neuroplasticity. Consumption of food throughout the waking hours results in little or no metabolic switching which can result in insulin resistance and a relative lack of engagement of neuronal networks involved in navigation and critical decision-making. As a consequence, signaling pathways that promote neuroplasticity and resilience are disengaged with the result being suboptimal cognitive abilities and vulnerability of the brain to stress and neurodegenerative disorders. Animal studies have shown that high energy diets and diabetes accelerate cognitive decline and motor dysfunction in models of Alzheimer's disease (AD) and Parkinson's disease (PD), respectively. Excessive energy intake accelerates the underlying accumulation of amyloid β -peptide ($A\beta$) and hyperphosphorylated Tau (pTau) in the brain in AD, and α -synuclein in PD. NRF2, nuclear regulatory factor 2; PGC-1 α , peroxisome proliferator-activated receptor γ coactivator 1 α .




Figure 2. Epigenetic impact of food overabundance and a sedentary lifestyle on the metabolism and cognitive outcomes of offspring.

A. Under conditions of moderate intermittent food intake and regular physical activity female mice give birth to offspring that inherit epigenetic DNA and chromatin modifications that result in gene expression profiles that promote a healthy metabolic phenotype, stress resistance, optimal cognitive function and resistance to neurological disorders. Conversely, females and males that are obese as the result of overconsumption of food and a sedentary lifestyle generate offspring that inherit epigenetic DNA and chromatin modifications and gene expression profiles that render them prone to insulin resistance and obesity, stress susceptibility, impaired cognitive function and vulnerability to neurological disorders. B. Intermittent food restriction and exercise promote epigenetic DNA and chromatin modifications (DNA and histone methylation, and chromatin protein acetylation and ubiquitination) that induce the expression of genes encoding proteins that enhance neuroplasticity and neuronal stress resistance (e.g., neurotrophic factors such as BDNF, antioxidant enzymes, protein deacetylases such as SIRT1 and SIRT3, and DNA repair enzymes such as APE1). Pro-inflammatory gene expression is suppressed by intermittent energy restriction and exercise. Food overconsumption and a sedentary lifestyle shift epigenetic modifications and consequent changes in gene expression in a manner that impairs neuroplasticity and increases vulnerability of the brain to stress. See Box 1 and references 1, 32 and 116 for details.