

Published in final edited form as:

Nat Rev Neurol. 2018 September ; 14(9): 544–558. doi:10.1038/s41582-018-0047-2.

C9orf72-mediated ALS and FTD: multiple pathways to disease

Rubika Balendra^{1,2} and Adrian M. Isaacs^{1,3,*}

¹Department of Neurodegenerative Disease, UCL Institute of Neurology, London, UK

²Department of Genetics, Evolution and Environment, Institute of Healthy Ageing, University College London, London, UK

³UK Dementia Research Institute at UCL, UCL Institute of Neurology, London, UK

Abstract

The discovery that repeat expansions in the *C9orf72* gene are a frequent cause of amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD) has revolutionized our understanding of these diseases. Substantial headway has been made in characterizing *C9orf72*-mediated disease and unravelling its underlying aetiopathogenesis. Three main disease mechanisms have been proposed: loss of function of the *C9orf72* protein, toxic gain of function from *C9orf72* repeat RNA or from dipeptide repeat proteins produced by repeat-associated non-ATG translation. Several downstream processes across a range of cellular functions have also been implicated. In this article, we review the pathological and mechanistic features of *C9orf72*-associated FTD and ALS (collectively termed C9FTD/ALS), the model systems used to study these conditions, and the probable initiators of downstream disease mechanisms. We suggest that a combination of upstream mechanisms involving both loss and gain of function, and downstream cellular pathways involving both cell-autonomous and non-cell-autonomous effects, contribute to disease progression.

Introduction

Amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD) are devastating and fatal neurodegenerative diseases. In common with other neurodegenerative diseases, progress towards finding disease-modifying therapies in ALS and FTD has been slow, in large part owing to an incomplete understanding of disease aetiopathogenesis. In 2011, ground-breaking progress was made with the discovery that a hexanucleotide GGGGCC repeat expansion in the *C9orf72* gene (Fig. 1) is the most frequent genetic cause of both diseases in Europe and North America^{1,2}. Interestingly, C9FTD/ALS — the collective term for *C9orf72*-associated diseases with clinical features of FTD, ALS or both — is extremely rare in Asia and the Middle East^{3,4}, indicating a different genetic architecture underlying

* a.isaacs@ucl.ac.uk.

Author contributions

Both authors researched data for the article, discussed the content, wrote the article, and reviewed and edited the manuscript before submission.

Competing interests statement

The authors declare no competing interests.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

FTD and ALS in these populations. The age of onset of C9FTD/ALS ranges from 27–83 years of age^{3,4}, and the disease duration ranges from 1–22 years^{3,4}. *C9orf72* repeat expansions have also been identified as a rare cause of other neurodegenerative diseases⁴, including Parkinson disease, progressive supranuclear palsy, ataxia, corticobasal syndrome, Huntington disease-like syndrome, Creutzfeldt–Jakob disease and Alzheimer disease.

The vast majority (>95%) of neurologically healthy individuals have 11 hexanucleotide repeats in the *C9orf72* gene⁵. The pathological repeat-length threshold has not been clearly defined; an arbitrary cut-off of 30 repeats is used in most studies, but larger expansions ranging from hundreds to thousands of repeats are most commonly observed in patients with C9FTD/ALS^{5–10}. Importantly, the discovery of C9FTD/ALS has heightened the realization that ALS and FTD are intimately linked on a clinical, genetic, pathological and mechanistic spectrum.

The expanded GGGGCC repeats are bidirectionally transcribed into repetitive RNA, which forms sense and antisense RNA foci^{11–15}. Remarkably, despite being within a non-coding region of *C9orf72*, these repetitive RNAs can be translated in every reading frame to form five different dipeptide repeat proteins (DPRs) — poly-GA, poly-GP poly-GR, poly-PA and poly-PR — via a non-canonical mechanism known as repeat-associated non-ATG (RAN) translation (Fig. 2)^{14–18}. Although *C9orf72* mutations are a relatively recent discovery, progress in understanding their pathogenic effects has been rapid. Three competing but non-exclusive mechanisms have arisen: loss of function of *C9orf72* protein, and toxic gain of function from sense and antisense *C9orf72* repeat RNA or from DPRs. These mechanisms are all likely to contribute to disease to some extent, but it is crucial to determine their relative importance at various disease stages, so as to inform therapeutic strategies. The differential involvement of these mechanisms might also explain clinical, pathological and prognostic heterogeneity that is observed in patients with C9FTD/ALS.

In this article, we review the current understanding of the mechanisms underlying C9FTD/ALS, and the questions that remain unanswered. The clinical and genetic aspects of C9FTD/ALS have been extensively reviewed elsewhere^{19,20} and will not be discussed in detail here.

Repeat size and somatic mosaicism

Defining the minimum number of hexanucleotide repeats in *C9orf72* that cause disease would be invaluable for genetic counselling and to guide disease modelling, but this issue is currently unresolved. Neurologically healthy individuals and patients with ALS or FTD can all have 20–30 repeats, so whether repeat lengths within this range can drive disease is unclear. For example, a screen of control post-mortem brains identified an individual with 30 repeats, sparse RNA foci and DPR inclusions, but no neurological symptoms up until death at 84 years of age²¹.

One confounding factor for accurate repeat sizing is somatic instability of the mutation. Individuals have been identified with large expansions within the CNS but an intermediate repeat length in DNA extracted from blood^{22–24}. These studies show that >50 repeats in

blood-derived DNA can be associated with large CNS expansions, suggesting that 50 repeats is a useful cut-off when analysing blood DNA. However, the current data do not support a single precise cut-off; for instance, 70 repeats were insufficient to cause disease in an 89-year-old whose children inherited much larger expansions and went on to develop C9FTD/ALS²⁵. Furthermore, defining a cut-off for blood-derived DNA does not help answer the key question of the minimum repeat size required in the CNS to cause disease.

A further confounder is that repeat size varies between different brain regions^{6,10}. This so-called somatic mosaicism within the CNS might explain some of the clinical heterogeneity that is observed between patients, but is currently not well understood.

Pathological features of C9FTD/ALS

TDP-43 inclusions

The overwhelming majority of ALS cases and approximately 50% of FTD cases are characterized by inclusions consisting of the RNA-binding protein TDP-43 (TAR DNA-binding protein 43) in neurons and glia^{26,27}. The fact that *C9orf72* mutations can lead to TDP-43 inclusions (Fig. 3) in both ALS and FTD implies a final common pathway in these diseases²⁸. This pathology is evident in various brain regions, including the frontal, temporal and primary motor cortices, hippocampus, basal ganglia, amygdala, thalamus and midbrain^{29–33}. Patients with *C9orf72* mutations who have a predominant ALS syndrome can still exhibit extramotor pathological features consistent with FTD and, conversely, those with a predominant FTD syndrome can show pathology in the motor system³¹.

RNA foci

Sense and antisense RNA foci comprising *C9orf72* repeat RNA are widely distributed across the CNS in patients with C9FTD/ALS (Fig. 3). These foci are found predominantly within neuronal nuclei in the frontal and motor cortices, hippocampus, cerebellum and in the spinal cord, in motor neurons and occasionally in interneurons, and sporadically in the cytoplasm^{11,13–15,34}. Less frequently, foci are detected in glia (astrocytes, microglia, astrocytes and oligodendrocytes^{11,13,15}. Sense RNA foci occur in ~37% and antisense foci in ~26% of neurons in the frontal cortex, respectively^{11,34}, and they co-occur in ~14% of frontal cortex neurons¹¹.

DPR pathology

DPR inclusions are p62-positive and TDP-43-negative and can consist of more than one DPR (Fig. 3)^{14,15,17,18,35}. Several studies have examined DPRs in patients with *C9orf72* repeat expansions, using immunohistochemistry^{14–18,29,35–42}, immunoblotting^{14,16,17,38,41} or immunoassays⁴³, and have reached similar conclusions. DPRs most commonly form neuronal cytoplasmic inclusions^{15–17,29}, but can also exist as neuritic inclusions^{29,44} or as 'pre-inclusions'^{15,29}, which appear as diffuse cytoplasmic staining. Neuronal intranuclear inclusions are sometimes observed^{15–17,29}, and are occasionally paranucleolar³⁶. Sense-derived poly-GA is the most frequent form of DPR^{16,18,38–41}, followed by poly-GP then poly-GR. The antisense-derived DPRs poly-PA and poly-PR are the least frequent forms^{15,18,35,36,38–40}. Although sense-derived DPRs seem

to be more prevalent than antisense-derived DPRs in patients with *C9orf72* repeat expansions, both sense-derived and antisense-derived poly-GP are detected in hippocampal neurons in these individuals, with strand-specific antibodies suggesting a preponderance of antisense-derived poly-GP14. Staining of cortical tissue with antibodies specific to the carboxy-terminal region of the translated DPRs revealed that translation occurs beyond the 3' end of the repeats in the cortex in *C9orf72* repeat expansion carriers14,18.

DPR pathology is most prominent in the cerebellum, hippocampus and neocortex, is less frequent in subcortical regions, and is rarely observed in the brainstem and spinal cord. In addition, all DPRs can be detected in insoluble fractions from patient frontal cortex or cerebellum, as high-molecular-weight species, indicating they are aggregation prone 14,16,17,38. A Meso Scale Discovery immunoassay has quantitatively detected poly-GP and poly-GA — but not, to date, any of the other DPRs — in both CNS tissue43 and cerebrospinal fluid (CSF)45. CSF poly-GP levels might have utility as a biomarker for both diagnosis and pharmacodynamic response46–48.

Pathogenesis of C9FTD/ALS

The complementary use of human tissue and *in vitro* and *in vivo* models, including illuminating mouse models examining both loss-of-function and gain-of-function mechanisms, has informed our current understanding of the contribution of *C9orf72* loss of function, *C9orf72* repeat RNA and DPRs to pathogenesis. These studies also have implicated several downstream mechanisms resulting from *C9orf72* expansions (Fig. 4).

Loss-of-function mechanisms

***C9orf72* transcription and splicing**—In *C9orf72* transcript variant 2, the repeats are located within the promoter region (Fig. 1), so are not incorporated into variant 2 pre-mRNA but have the potential to affect the expression of this variant. By contrast, in variants 1 and 3, the repeats are within intron 1, so are included in the respective pre-mRNAs. Variant 2 is expressed at higher levels than variants 1 and 3 in CNS tissue138,139.

Studies have demonstrated reduced levels of one or more of the *C9orf72* transcript variants in blood lymphocytes1,132,140, induced pluripotent stem cell (iPSC)-derived neurons12,90,94,138, frontal cortex1,132,141–145, cerebellum12,16,143–145, motor cortex12 and cervical spinal cord12 from *C9orf72* expansion carriers compared with controls. The findings are particularly robust for variants 1 and 2145. *C9orf72* protein levels might be correspondingly reduced in the frontal cortex110,144. Transcripts upstream of the repeat are increased relative to downstream transcripts in blood lymphocytes and brain and spinal cord tissue66 from *C9orf72* expansion carriers, possibly owing to abortive transcription in the presence of the repeat expansion66. Raised levels of variant 1 in the frontal cortex and cerebellum are associated with increased survival145 — an important consideration when developing therapies that affect transcript levels.

The intronic location of the repeats in variants 1 and 3 means they should be spliced out of the transcript, but the fact that they are translated into DPRs implies that they are either retained in the transcript or that the spliced intron is translated. Although levels of mature

spliced *C9orf72* mRNA are reduced in the brains of individuals with C9FTD/ALS, levels of sense and antisense transcripts containing intron 1, where the repeats are located, are increased^{14,16}, suggesting stabilization of repeat RNAs. Mature *C9orf72* transcripts with correct splicing or with retention of intron 1 are both detected in C9FTD/ALS lymphoblasts and brain tissue^{138,146}, indicating that both species contribute to RAN translation.

The expression of *C9orf72* is also modified by epigenetic effects (Box 1).

Autophagy and lysosomal function—Bioinformatic analysis shows that *C9orf72* is structurally related to the differentially expressed in normal and neoplastic cells (DENN) guanine nucleotide exchange factor (GEF) proteins, which activate Rab proteins. Rabs are crucial for a wide range of vesicular trafficking events, and multiple lines of evidence from several independent groups point to a role for *C9orf72* in autophagy and endolysosomal trafficking and function.

Knockdown of *C9orf72* in human cell lines and primary neurons specifically inhibits autophagy induction, but not later stages of the autophagy pathway^{91,96}, leading to accumulation of p62^{91,96} and cytoplasmic aggregation of TDP-43⁹⁶. Consistent with these findings, accumulation of autophagy substrates, including p62, is observed in the spleens of *C9orf72*-knockout mice^{54,56}. Conversely, overexpression of *C9orf72* can activate autophagy, leading to an increase in autophagosomes in cell lines⁹¹. The role in autophagy seems to be mediated by the long *C9orf72* protein isoform (*C9orf72-L*) rather than the short isoform (*C9orf72-S*)⁹⁶. The mechanism underlying these changes involves Rab proteins, although no consensus has been reached on which are the most important, with Rab1a, Rab8a and Rab39b all being implicated^{56,78,91,96,154}. *C9orf72* interacts with guanine nucleotide exchange protein SMCR8 and WD repeat-containing protein 41 (WDR41)^{56,58,96,101,122,154}, and the effect on autophagy is generally agreed to be mediated through an interaction with the serine/threonine-protein kinase ULK1 complex^{56,91,96,100,154}, a key initiator of autophagy. Potentially, this interaction can also occur via TBK1⁹⁶, another serine/threonine-protein kinase implicated in autophagy. This finding is particularly intriguing, as loss-of-function mutations in *TBK1* cause ALS and FTD¹⁵⁵.

Another link to known ALS-associated genes is through ataxin-2 (*ATXN2*), in which intermediate expansions of polyglutamine increase the risk of ALS¹⁵⁶. *C9orf72* knockdown specifically increases the aggregation and toxicity of ataxin-2 protein with intermediate polyglutamine repeats⁹⁶.

The relevance of the role of *C9orf72* role in autophagy for disease pathogenesis is unclear, but neurons from patients with C9FTD/ALS have impaired basal autophagy^{91,93} and increased sensitivity to autophagy inhibition⁹⁰, suggesting that reductions in *C9orf72* levels contribute to cellular distress. Furthermore, a Src-c-Abl pathway inhibitor, which increases autophagic processes, rescues survival defects in neurons derived from patients with ALS⁹².

In addition to effects on autophagy, reduced endocytosis was reported in *C9orf72* knockdown cell lines⁷⁸, and impaired endosomal and lysosomal trafficking were observed

in bone marrow-derived macrophages and microglia from homozygous *C9orf72* knockout mice⁵⁴, as well as in patient-derived fibroblasts and neurons⁹³. Moreover, *C9orf72* has been shown to reside on lysosomes and can directly affect lysosomal function, which might also explain the effects of *C9orf72* loss of function on both endolysosomal trafficking and autophagy¹⁰¹. In iPSC-derived motor neurons, *C9orf72* primarily localizes to early endosomes, and iPSC-derived motor neurons from patients with C9FTD/ALS have fewer lysosomes⁹⁴. Both patient neurons and CRISPR–Cas9 *C9orf72*-knockout iPSC-derived neurons have reduced vesicular trafficking, which can be rescued by *C9orf72* overexpression⁹⁴. These cells also have elevated glutamate receptor levels and increased sensitivity to excitotoxicity^{94,127}. Consistent with this observation, increased glutamate receptor levels were found in spinal cord tissue in a *C9orf72*-knockout mice⁹⁴, and in spinal cord^{94,127} and cortical tissue⁹⁴ from patients with C9FTD/ALS.

Taken together, these data suggest that *C9orf72* is involved in multiple cellular trafficking events, and that loss of *C9orf72* in both microglia and neurons can sensitize cells to other insults, thereby contributing to neurodegeneration in C9FTD/ALS.

Further insights from loss-of-function mouse models—The mouse *C9orf72* orthologue shares 98% homology with human *C9orf72* and is expressed in embryonic and early postnatal neurons, various regions of the adult brain and spinal cord, glia, and non-neuronal tissues, including muscle, spleen, kidney and testes^{50–54,157}. Several *C9orf72* knockout or knockdown models have now been reported (Table 1)^{51–58}.

Transient reduction of *C9orf72* expression in the CNS by antisense oligonucleotides (ASOs)¹³ and conditional homozygous knockouts of *C9orf72* in neurons and glia⁵¹ do not lead to motor or behavioural phenotypes. By contrast, ubiquitous knockouts of *C9orf72*^{52–55,57,58} or CRISPR–Cas9-mediated knockouts of *C9orf72* isoforms^{56,57} throughout development led to immune system dysregulation in homozygous mice. The phenotypes included changes in myeloid and/or lymphoid cell populations in the spleen and lymph nodes, increased levels of inflammatory cytokines, and cervical or systemic lymphadenopathy and splenomegaly, sometimes with reduced body weight^{52,53,55,57}, neoplasia⁵⁵ or increased autoimmune antibody titres^{53,57}. In comparison, haploinsufficiency of *C9orf72* does not lead to severe phenotypes.

Although some studies reported mild motor or cognitive phenotypes^{52,53} or reduced lifespan^{52,53,55,57,58} in homozygous *C9orf72*-knockout mice, none reported neuronal loss. Transcriptomic analysis confirmed changes in immune pathways^{53,54}, similar to those observed in CNS tissue from patients with *C9orf72* repeat expansions⁵⁴. Transcriptomic analysis in human tissue has shown that *C9orf72* transcripts are particularly prevalent in CD14⁺ myeloid cells, which are involved in innate and adaptive immunity¹³⁹. Overall, in line with cellular studies of *C9orf72*, these findings suggest an important role for *C9orf72* in immune regulation, possibly through its effects on autophagosome and lysosome function and/or microglial activity, or through alteration of autoimmune responses. However, in contrast to *C9orf72* loss of function, knockout of other autophagy-related genes, including *ATG7158* and *ATG5159*, in neurons in mice leads to neurodegeneration. These findings

suggest that *C9orf72* is not an essential component of the autophagy pathway that mediates neuronal survival.

Crucially, none of the mouse *C9orf72* knockouts recapitulate ALS or FTD, suggesting that *C9orf72* loss of function is insufficient to precipitate disease. However, given the role of *C9orf72* in pathways previously implicated in FTD and ALS160, haploinsufficiency might contribute to the disease process in combination with gain-of-function mechanisms, and an interesting approach will be to breed loss-of-function mouse models with gain-of-function models.

Gain-of-function mechanisms

The question of whether *C9orf72* repeat RNA or DPRs produced by RAN translation are the toxic species in aetiopathogenesis is hotly debated in the field. Various approaches, each of which has limitations, have been used to address this issue. Post-mortem studies generally do not capture the earliest pathogenic events. *In vitro* and *in vivo* models often do not feature the long repeats that are found in patients, owing to methodological difficulties in cloning GC-rich repeats of this length. Therefore, these models might not fully recapitulate the disease mechanisms. Overexpression models do not necessarily reflect endogenous expression levels in patients. Many models express expanded repeats, but as these repeats can go on to produce both *C9orf72* repeat RNA and DPRs, attribution of downstream mechanisms to either entity is challenging. Despite these uncertainties, however, clear mechanistic pathways have emerged (Table 3).

C9orf72* repeat RNA—*In vitro, GGGGCC repeat RNA forms secondary structures, including hairpins^{45,66} and highly stable G-quadruplexes^{45,66,161–163}. Other secondary structures, including DNA–RNA heteroduplexes, RNA duplexes and i-motifs^{164–167}, might arise from the sense and antisense repeat RNA and DNA sequences. *In vivo*, such secondary structures are likely to mediate the sequestration — and, as a consequence, depletion — of RNA-binding proteins (RBPs)^{66,161} (Box 2), thus providing a clear potential route to RNA toxicity (reviewed extensively elsewhere^{164,168}).

DPRs: insights from post-mortem studies—The results of post-mortem studies have raised suspicions that DPR inclusions are not the primary culprit in C9FTD/ALS pathogenesis^{29,35–40,43,181–183}. TDP-43 pathology and neurodegeneration co-occur in affected regions of the CNS in ALS and FTD^{29,35,37–39,43}. By contrast, DPR pathology does not coincide neuroanatomically with TDP-43 pathology⁴³ and is generally not found in the same neurons as TDP-43 inclusions^{16,29,35,40}. Furthermore, DPR inclusions do not differ in neuroanatomical distribution between FTD and ALS cases^{29,37,38} and are rare in the spinal cord in *C9orf72*-associated ALS, whereas TDP-43 pathology is common^{14,17,18,29,36–40}. DPR inclusions are frequent across several brain regions^{17,35–37,39}, including structures that are thought to be minimally affected in ALS and FTD, such as the cerebellum and occipital and parietal lobes. However, one study showed that poly-GR inclusions — but not the other DPRs — correlated with areas of neurodegeneration in *C9orf72*-associated ALS, and also, of all the DPRs, uniquely colocalized with TDP-43 pathology in a small sample of brains that were obtained shortly after death⁴⁴. These data

suggest that further investigation in larger, deep-phenotyped post-mortem cohorts will provide important insights.

Despite these observations, strong counterarguments to support a pathogenic role for DPRs in C9FTD/ALS have been put forward. Post-mortem studies tend to represent the final stages of the disease process and might not reflect the early pathogenicity of DPRs. Aggregates observed at post-mortem could represent protective species, and correlations with inclusions might be misleading if soluble species mediate neurotoxicity. CNS regions that have extensive DPR pathology but are unaffected by neurodegeneration might contain protective factors; indeed, selective vulnerability is a frequent observation in neurodegenerative diseases¹⁸⁴.

TDP-43 pathology is likely to be downstream of DPR pathology, probably explaining why it correlates more closely with neurodegeneration. This idea is consistent with downstream effects on TDP-43 in some experimental models expressing expanded repeats^{109,185} or pure DPRs^{106,185}, suggesting that one or both of these gain-of-function mechanisms are linked to TDP-43. Affected individuals with DPR pathology but relatively mild or absent TDP-43 pathology have been reported^{16,21,43,141,182,186,187}. These individuals include a young patient with evidence of intellectual disability¹⁸⁶, a patient with *C9orf72*-associated ALS who had little extramotor TDP-43 pathology but showed evidence of cognitive impairment and high cerebellar DPR levels⁴³, and patients with pathological or clinical diagnoses of FTLD or FTD^{16,141,182,186,187}, some of whom died prematurely from other causes¹⁸². Therefore, DPR pathology without substantial TDP-43 pathology seems to be sufficient for disease to develop in some cases. Furthermore, as discussed below, considerable evidence from model systems indicates that DPRs can cause neurodegeneration.

Insights from gain-of-function models

Evidence for *C9orf72* repeat RNA toxicity—The effects of *C9orf72* repeat RNA were modelled in primary cortical and motor neurons transfected with expanded GGGGCC repeats within an artificial intronic region of the green fluorescent protein gene, reflecting the intronic human genomic context of *C9orf72* expansions⁴². These neurons demonstrated nuclear RNA foci and reduced survival. Dot blots and immunocytochemistry revealed no DPRs in these cells, suggesting that the reduced survival was attributable to repeat RNA. Interestingly, however, co-expression of poly-PR and the intronic expanded GGGGCC repeats had a synergistic detrimental effect on neuronal survival.

RNA toxicity has also been implicated in eye and motor neuron degeneration in a *Drosophila* model that expresses 30 GGGGCC repeats with a 6 bp (CTCGAG) interruption in the middle of the repeats^{109,172}. DPRs were not detected in the eyes or neurons, and were only detected when the repeats were strongly induced in all tissues¹⁰⁹.

One caveat for the interpretation of both studies is that the inability to detect DPRs is not sufficient to exclude a role for these proteins. In our experience, poly-GR can be difficult to detect even in flies that express this protein at high levels and show overt toxicity. Therefore,

more sensitive detection assays for DPRs will be required to unpick the relative contributions of RNA and DPRs in these models.

A study in developing zebrafish found that both sense and antisense RNA repeats could mediate toxicity, leading to a motor axonopathy phenotype¹³⁴. No DPRs were detected in this model. In a *Drosophila* model, however, neither sense nor antisense repeats with similar lengths to those found in patients with C9FTD/ALS led to degeneration of adult neurons¹⁸⁹. One potential explanation for this difference is that developing neurons are more susceptible to RNA toxicity than adult neurons. Another possibility is that RNA foci in humans sequester RBPs that are not present in *Drosophila*, thereby limiting the utility of *Drosophila* as a model for RNA toxicity.

Evidence for DPR toxicity—Two studies in *Drosophila* models have provided evidence that neurotoxicity is attributable to DPRs rather than repeat RNA^{138,188}. In the first study, overexpression of expanded GGGGCC repeats in *Drosophila* eyes or adult neurons led to neurodegeneration¹⁸⁸. This effect was inhibited when the repeats were interrupted by stop codons in each reading frame that prevented translation of the repeats into DPRs. The second study involved ubiquitous overexpression of 160 GGGGCC repeats in an intronic context, flanked by human *C9orf72* sequence¹³⁸. The intron containing the repeat was spliced out and formed large numbers of sense RNA foci in neurons and glia, without production of DPRs. This model showed no evidence of neurodegeneration, reduced survival or widespread transcriptomic changes. Increasing transgene expression in this model led to DPR production and lifespan reduction, although the frequency of sense foci remained the same¹³⁸, supporting the idea that DPRs rather than sense foci mediate neurodegeneration.

It should be noted that although the RNA sequence interrupted by stop codons in the first study forms the same G-quadruplex secondary structure as uninterrupted *C9orf72* repeat RNA¹⁸⁸, the tertiary and quaternary structures are not necessarily identical, which might affect the dynamics of RBP sequestration. However, the intronic model did not have an interrupted repeat sequence and exhibited multiple sense foci, yet still showed no evidence of toxicity¹³⁸.

Toxicity of individual DPRs—In numerous studies in cell models^{14,42,69,87,104–106,190}, *Drosophila*^{42,72,108,115,123}, zebrafish^{133,134,191} and mice^{63,64}, the repetitive GGGGCC sequence was altered to generate coding sequences that expressed each DPR in isolation, which was often sufficient to produce neurotoxicity and implicate several downstream mechanisms. The main limitation of these models is that DPR overexpression might not reflect the endogenous mechanisms that are seen in patients.

Studies in cultured neuronal or non-neuronal cell lines or primary neuronal cultures indicate that poly-GR and poly-PR are the most toxic of the DPRs^{42,69,87,106}. These arginine-rich DPRs — in particular, poly-PR — are toxic to yeast.⁸⁹ Poly-GA and poly-PA are also toxic, but to a lesser extent⁸⁹. Synthetic poly-PR and poly-GR are highly toxic when exogenously applied to cultured human astrocytes. Poly-PR has a longer half-life than poly-GR, and is especially potent in this context⁶⁸. Synthetic poly-GR and poly-GA are also toxic to

primary neurons¹⁹². Furthermore, overexpression of poly-PR was found to be toxic to control iPSC-derived neurons⁴².

Of all the DPRs, poly-GR had the greatest detrimental effect on development, locomotor activity and survival in a zebrafish model¹³³. In *Drosophila* models, poly-GR and poly-PR were neurotoxic when expressed in the eyes^{42,72,108,115,123,188}. In addition, flies expressing these proteins exhibited reduced survival^{42,108,123,188} and locomotor phenotypes^{42,123,135}. Most of these studies found that poly-GA^{42,115,123}, poly-PA^{42,188} and poly-GP¹⁰⁸ were not toxic in *Drosophila*, although one study reported a mild reduced survival phenotype when poly-GA was expressed in adult neurons¹⁸⁸. Consistent with this finding, poly-GA overexpression in cultured cells^{104,105}, primary neurons¹⁰⁴, zebrafish^{133,191} or mouse brain⁶³ leads to toxicity, and synthetic poly-GA exogenously applied to human cells¹⁹⁰ or primary neurons¹⁹² is also toxic. In addition, cryo-electron tomography revealed that poly-GA forms twisted ribbon structures that sequester the 26S proteasome¹⁰⁷.

Overall, these studies show that poly-GR and poly-PR are potently neurotoxic and poly-GA also exerts toxicity. The remaining DPRs, poly-PA and poly-GP, are unlikely to be toxic species. As poly-GR, poly-PR and poly-GA can all be damaging when overexpressed, a key aim is to establish the levels of each of these species in patient tissue, particularly at early disease stages. New techniques to extract and measure DPRs from patient tissue and assess their toxicity will be essential for this endeavour.

Mouse gain-of-function models—Several mouse models of *C9orf72* gain of function have been characterized. Adeno-associated virus-mediated CNS overexpression of 66 GGGGCC repeats leads to motor and behavioural phenotypes by 6 months, with RNA foci, DPRs, phospho-TDP-43 inclusions and neuronal loss being observed⁵⁹. The same approach was used to generate mice specifically overexpressing poly-GA in the CNS⁶³. These mice developed neuronal cytoplasmic inclusions of fibrillar poly-GA, as well as neurodegeneration and motor, cognitive and behavioural phenotypes. However, these effects were not observed when the poly-GA sequence was mutated to a sequence that was unable to aggregate. Phospho-TDP-43 inclusions were rarely found; therefore, this model does not fully recapitulate the (GGGGCC)₆₆ repeat mouse phenotype, indicating that species other than poly-GA are the main drivers of phospho-TDP-43 accumulation. A further poly-GA model, with expression levels more comparable to those in the cortex in patients with C9FTD/ALS, demonstrated motor deficits, but overall a less severe phenotype than the viral poly-GA model⁶⁴.

Bacterial artificial chromosome transgenic mouse models—Bacterial artificial chromosome (BAC) transgenic mice have the advantages of expressing the human *C9orf72* gene, with surrounding regulatory regions and flanking sequences, at more physiological levels. Three BAC transgenic models have produced similar findings. Two of these models used BAC constructs^{52,60}, containing exons 1–5 of the gene and 300–500 repeats (Table 2). The third BAC model expressed the full gene and protein with 100–1000 repeats⁶¹. All three models developed RNA foci and DPRs in the CNS, but none demonstrated TDP-43 inclusion pathology, neuronal loss or reduced survival. One model developed memory

impairment and loss of hippocampal neurons, with an increase in levels of phosphorylated TDP-43, but was not fully reflective of ALS or FTD⁵². In this model, a single injection of an ASO targeting *C9orf72* RNA led to sustained reductions in RNA foci and DPR pathology, and improved cognition. Repeat length had a strong influence on the formation of RNA foci, with 100-repeat mice developing no foci and 400-repeat mice developing many foci, despite considerably higher transgene expression in the 100-repeat mice⁵².

Only one study has reported BAC transgenic mice with a striking neurodegenerative phenotype⁶². The mice expressed the full gene and different repeat lengths in different lines. Three independent lines, two with ~500 repeats and one with high expression levels of 36 repeats, developed RNA foci, DPR aggregates, TDP-43 pathology and neurodegeneration. A subset of female mice developed an acute motor phenotype, with paralysis, weight loss and decreased survival, with other female and male mice showing slower progressive neurodegeneration. Antisense transcripts and foci were observed at especially high levels in some CNS regions that showed neurodegeneration, whereas sense foci were more evenly distributed. These data show that high expression of short repeats can be toxic, indicating that large repeats are not the critical factor for toxicity in this mouse model. The reason why female mice are particularly susceptible and only a subset develops acute disease is currently unclear, although factors such as methylation might be important.

In-depth molecular and phenotypic comparisons between all the BAC models should further our understanding of *C9orf72*-associated pathogenic mechanisms. One model that expresses the full-length human gene with the repeat expansion recapitulates disease⁶² whereas the other does not⁶¹. These mice feature different genetic backgrounds and flanking *C9orf72* sequences (Table 2), which might be contributory factors. Backcrossing the models that did not show a phenotype into different genetic backgrounds, so as to determine whether certain backgrounds facilitate disease, would provide new insight.

Downstream mechanisms

A range of downstream mechanisms in C9FTD/ALS have been validated across multiple human and non-human model systems and different laboratories (Table 3). Dysfunctional nucleocytoplasmic transport (NCT) is a prominent mechanism that has been identified in genome-wide screens in yeast⁸⁹ and *Drosophila*^{108,109,115}, and in CRISPR–Cas9 screens in human cells and primary neurons⁸⁸ (reviewed elsewhere^{168,193}).

Poly-PR and poly-GR have been shown to interact with proteins that contain low complexity domains (LCDs), which include many RBPs^{72,194}. LCD proteins can undergo liquid–liquid phase separation (LLPS) to form droplets. Through this process, the proteins become compartmentalized in the cell, forming membrane-less organelles such as nucleoli and stress granules. These organelles facilitate the assembly of RNA and RBPs into ribonucleoproteins, and also aid subsequent RNA metabolism. Furthermore, these LCDs can form reversible hydrogels, which have the propensity to fibrillize into irreversible hydrogel-like structures. Mutations frequently occur in the LCD domains in several RBPs that are involved in ALS, including FUS and TDP-43¹⁹⁵. These mutations can alter the LLPS dynamics of RBPs, leading to their fibrillization and aggregation^{196–198}. Poly-GR and poly-PR interact with LCD proteins in nucleoli and stress granules, thereby impairing LLPS,

disrupting the dynamics of assembly of these organelles, and affecting mRNA translation and NCT72,73,82. Aliphatic alcohols, which disrupt phase separation and hydrogel formation, can reduce poly-PR's protein interactions¹⁹⁴ and disrupt the nucleolar localization of poly-PR and poly-GR⁷². Knockdown of several of these LCD proteins modifies the eye degeneration phenotype in a *Drosophila* poly-GR model⁷². Poly-PR also interacts with LCDs in intermediate filament proteins¹⁹⁴. Interestingly, arginine-rich DPRs at high concentrations can undergo LLPS⁷³. In addition, GGGGCC repeat RNA can undergo gel transition^{81,199} and induce phase transition of RNA granule proteins⁸¹ in the absence of other LCD proteins. These data further implicate altered LLPS in C9FTD/ALS pathogenesis.

Effects on membrane-less organelles have been observed in several other studies. Primary cortical neurons that overexpress poly-PR, leading to the formation of nuclear poly-PR aggregates, show a reduction in cytoplasmic P-bodies and an increase in stress granule formation⁴². In cultured cells treated with arsenite, a stress granule inducer, overexpression of poly-PR and poly-GR reduced stress granule formation, whereas poly-GA, poly-GP and poly-PA induced stress granule formation⁶⁹.

Poly-PR interacts with translation initiation and elongation factors, and overexpression of poly-GR and poly-PR, but not poly-GA, inhibited translation *in vitro* and in cell lines⁸⁷. Translational inhibition was attributed to direct binding of poly-GR and poly-PR to mRNA, thereby blocking access to the translation machinery⁸⁷. A second study also observed translational inhibition when (GGGGCC)₃₁ was expressed in cell lines⁷⁹. This inhibition was accompanied by an increase in stress granule formation (a marker of translational arrest) and nuclear accumulation of poly-A mRNAs and PABPc, a protein that facilitates mRNA translation in the cytoplasm⁷⁹. GGGGCC repeat-induced stress granule formation was also observed in subsequent studies^{80,81}. The translational inhibition might be explained by sequestration of PABPc by GGGGCC repeat RNA, but could also be due to the DPR-mediated mechanisms described above. Interestingly, two studies that identified translation inhibition suggested that this process was independent of eukaryotic translation initiation factor 2 subunit 1 (eIF2 α) phosphorylation, an important master regulator of translation that has been implicated in other neurodegenerative diseases²⁰⁰. However, stress granule induction secondary to poly-PR has been shown to require eIF2 α phosphorylation⁷³.

A study published in 2018 showed that poly-GR and poly-PR induced stress granule assembly and localization of NCT factors into these stress granules, thereby mediating NCT dysfunction⁸². Inhibition of stress granule assembly abrogated NCT dysfunction and neurodegeneration in patient-derived neurons and *in vivo*⁸². Stress granules and translation inhibition have been implicated in FTD and ALS associated with TDP-43 and FUS inclusions²⁰¹, and they provide an interesting potential common theme across the disease spectrum.

In three elegant studies, GGGGCC repeat overexpression constructs were used to investigate the mechanisms underlying RAN translation. Two of these studies showed that the integrated stress response, via eIF2 α phosphorylation, selectively increased RAN translation^{80,83}. Therefore, repeat-induced cellular stress could lead to both impaired

translation and enhanced RAN translation, causing a negative feedback loop. In addition, two of the studies showed that RAN translation occurred on unspliced, capped mRNA (in which the repeat-containing intron is retained), and was initiated by an upstream CUG acting as a start codon^{80,202}. By contrast, the third study found that RAN translation occurred on the spliced intronic RNA and was cap-independent, although cap-dependent RAN translation was more efficient⁸³. These mechanisms now need to be investigated in the context of endogenous RAN translation.

When applied exogenously to astrocyte cultures, poly-GR and poly-PR 20mers accumulate in the nucleolus, leading to splicing changes and impaired ribosomal RNA (rRNA) maturation⁶⁸. This finding led to the suggestion that these DPRs bind to and impair nuclear RBP complexes that are involved in mRNA splicing, which could in turn affect ribosome biogenesis and other important genes⁶⁸. Interestingly, given the effect of poly-PR and poly-GR on NCT, splicing of RanGAP, an essential regulator of NCT, was altered⁶⁸. Poly-PR and poly-GR also colocalize with nucleoli in cultured cells, primary neurons, iPSC-derived neurons and *Drosophila*, leading to abnormal nucleolar morphology^{36,42,67,69,71}. Importantly, the brains of individuals with *C9orf79*-associated FTLD exhibit bidirectional nucleolar volume changes, with smaller neuronal nucleoli overall but enlarged nucleoli in neurons containing poly-GR inclusions⁶⁷. Other interactome studies suggest that poly-PR and poly-GR associate with the spliceosome component U2 snRNP in cell nuclear extracts, block spliceosome assembly, and disrupt splicing when applied to cell nuclear extracts⁸⁵. These proteins also interact with mitochondrial ribosomal proteins and other ribosomal proteins¹¹⁷, and evidence of impaired mitochondrial function has been observed in iPSC-derived neurons from patients with C9FTD/ALS^{75,117}. Ribosomal proteins, hnRNPs, nucleolar proteins and RBPs associated with RNA granules have been shown to interact with poly-GR, and overexpression of poly-GR and poly-PR led to decreased levels of rRNA in human cell lines⁶⁹. In agreement with these data, poly-PR was shown to interact with proteins involved in mRNA splicing and ribosome assembly, and with ribosomal proteins⁸⁷. Three studies have shown that in addition to effects on RNA, *C9orf72* repeats can induce DNA damage, probably mediated by DPRs^{117–119}.

Overall, many cellular pathways have been implicated in gain-of-function toxicity, with the majority of studies focused on poly-GR and poly-PR. Given the highly toxic nature of these DPRs, they are likely to influence multiple pathways. Links between the downstream mechanisms and specific DPRs or repeat RNA have not yet been established, and sensitive techniques to measure both repeat RNA and DPRs in models expressing pure GGGGCC repeats will be required to enable the effects of individual molecular species to be distinguished.

Another important issue that remains to be addressed is whether different DPRs or repeat RNA and DPRs act synergistically, potentially in conjunction with loss of function of *C9orf72*, to elicit downstream effects. Crossing of models that express different DPR species should help to address this question. A further priority is to develop physiologically relevant models to reflect endogenous levels of *C9orf72* repeat RNA and DPRs, and human iPSC-derived neuronal models will be a key tool in this regard. Furthermore, *C9orf72* protein depletion, repeat RNA and DPRs might have non-cell-autonomous effects, and iPSC models

of neuronal, glial and muscle co-culture should increase our understanding of these complex interactions.

Therapeutic strategies

Targeting *C9orf72* RNA or DNA

ASOs that target *C9orf72* RNA can rescue *C9orf72*-specific pathologies^{12,13,46,126}, downstream gene expression changes^{12,126}, NCT defects¹⁰⁹ and TDP-43 mislocalization¹⁰⁹ in *C9orf72* fibroblasts or iPSC-derived neurons. These ASOs also reduce NCT defects and neurodegeneration in *Drosophila*¹⁰⁹ and diminish sense RNA foci and DPRs in mice^{46,52}. ASO trials in humans are planned, and the feasibility of this strategy has a precedent in studies of superoxide dismutase 1 (*SOD1*)-targeting ASOs in patients with ALS²⁰³.

An alternative strategy is to use compounds that target the secondary structure of *C9orf72* repeat RNA^{45,109,204–206}. These compounds could prevent sequestration of RBPs and/or interfere with the RNA structure to prevent RAN translation. Such molecules have been shown to affect the secondary structure of *C9orf72* repeat RNA *in vitro*^{45,204,206}, to reduce the production of RNA foci and DPRs when applied to patient neurons^{45,205} and *Drosophila*²⁰⁵, and to rescue nuclear import defects and neurodegeneration in a *Drosophila* model¹⁰⁹.

Targeting of the transcription elongation factor SPT4 reduces levels of sense and antisense *C9orf72* repeat transcripts and ameliorates disease phenotypes *in vitro* and *in vivo*, including in *C9orf72* iPSC-derived neurons²⁰⁷. Single-stranded small inhibitory RNAs have also been proposed as a strategy to silence the *C9orf72* repeat RNA²⁰⁸. In addition, two studies have shown that use of the CRISPR–Cas9 system to target either GGGGCC repeat DNA²⁰⁹ — thus reducing repeat transcription — or repeat RNA²¹⁰ can reduce RNA foci and DPR levels in cell lines. On the basis of these studies, targeting of *C9orf72* RNA or DNA are promising strategies, and ASOs are currently the most advanced in terms of clinical development.

Targeting DPRs and TDP-43

In other neurodegenerative diseases, active and passive immunological approaches have been used to target toxic proteins such as amyloid- β ²¹¹, tau^{212,213} and α -synuclein^{214,215}, leading to improved pathology and phenotypes in model systems. With the ongoing development of specific DPR antibodies, passive immunization to the DPRs could present a novel therapeutic approach for C9FTD/ALS, and poly-GA-specific antibodies have been shown to reduce intracellular poly-GA aggregation and seeding activity of C9FTD/ALS brain extracts²¹⁶. In addition, given that C9FTD/ALS pathogenesis converges on TDP-43, anti-TDP-43 immunotherapy would be a compelling strategy.

Current strategies in other neurodegenerative diseases are based on targeting of the extracellular pool of protein, which is presumably involved in cell-to-cell transmission. The success of such an approach in the case of DPRs and TDP-43 would depend on whether there is a disease-relevant extracellular pool to target. If not, methods to target antibodies

intracellularly would be required. Other important factors to consider are the ideal timing of these treatments, which DPRs and conformations to target, the specificity and safety of the treatment, and the risks of precipitating an autoimmune response. As short repetitive sequences similar to DPRs are present across the proteome, specificity might present a barrier. Increasing clearance of DPRs by other mechanisms could also be effective; for example, the small heat shock protein HSPB8 was shown to reduce DPR levels, probably via the autophagy pathway⁹⁸.

Targeting downstream mechanisms

Targeting of downstream mechanisms might represent a useful therapeutic strategy for C9FTD/ALS. Reducing nuclear export by targeting the nuclear export factors SRSF1 or exportin 1 ameliorates toxicity in *C9orf72*-repeat *Drosophila*^{84,109}; this was suggested to be due to either by reducing the levels of cytoplasmic repeat RNA (and, thus, DPRs) or by a more general mechanism to alterations in NCT. Importantly, in addition to the proof of concept genetic knockdown approaches that were used to reduce SRSF1 and exportin 1, small molecule exportin 1 inhibitors were also used, indicating a potential route to the clinic. It will now be important to address whether there is a large enough therapeutic window when targeting nuclear export, which is a fundamental cellular process. Inhibition of stress granule formation using either ASOs targeting Ataxin-2 or small molecules, also prevents NCT defects in C9-ALS iPSC-neurons and neurodegeneration in a *C9orf72* repeat fly model⁸². Given the success of Ataxin-2-targeting ASOs in extending lifespan in an ALS mouse model overexpressing TDP-43, this strategy and the mechanisms underlying it, are of clear interest. A Src-c-Abl pathway inhibitor that augments autophagy, improved the survival of iPSC-derived neurons from patients with ALS, including those with C9FTD/ALS, suggesting this approach could be beneficial for several forms of FTD/ALS⁹². Knockdown of TMX2, an endoplasmic reticulum protein that was identified in CRISPR-Cas9 screens in human cells and primary neurons, modulated endoplasmic reticulum stress in primary neurons overexpressing DPRs and increased survival of neurons derived from patients with *C9orf72*-associated ALS⁸⁸. Whether TMX2 is druggable awaits further study. High-throughput drug screens — for example, using iPSC-derived neuronal models^{92,94,217} — will be of great importance in establishing treatments to reduce levels of repeat RNA and DPRs, repair dysfunctional cellular mechanisms and rescue disease phenotypes⁸⁸.

Conclusions

Our knowledge of C9FTD/ALS has increased exponentially within a relatively short time period. Dissection of the disease mechanisms has not been straightforward, and the emerging picture is one of a combination of a diverse range of factors that lead to neurodegeneration. Given the relative importance of *C9orf72* repeat expansions as a causative factor for ALS and FTD, rapid translation of the accumulated knowledge into therapeutic strategies would have a substantial impact on patients with these devastating neurodegenerative diseases.

Acknowledgements

R.B. is a Leonard Wolfson Clinical Research Training Fellow and is funded by a Wellcome Trust Research Training Fellowship (107196/Z/14/Z). A.M.I. is funded by the Motor Neuron Disease Association, Alzheimer's Research UK, the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (648716 — C9ND) and the UK Dementia Research Institute. We thank Dr Sarah Mizielinska for reviewing the manuscript and assistance with figures, and Mr Martino Guadalupi and Dr Rachele Saccon for assistance with and design of figures.

References

- DeJesus-Hernandez M, et al. Expanded GGGGCC hexanucleotide repeat in noncoding region of *C9ORF72* causes chromosome 9p-linked FTD and ALS. *Neuron*. 2011; 72:245–256. [PubMed: 21944778]
- Renton AE, et al. A hexanucleotide repeat expansion in *C9ORF72* is the cause of chromosome 9p21-linked ALS-FTD. *Neuron*. 2011; 72:257–268. [PubMed: 21944779]
- Majounie E, et al. Frequency of the *C9orf72* hexanucleotide repeat expansion in patients with amyotrophic lateral sclerosis and frontotemporal dementia: a cross-sectional study. *Lancet Neurol*. 2012; 11:323–330. [PubMed: 22406228]
- Woolacott IO, Mead S. The *C9ORF72* expansion mutation: gene structure, phenotypic and diagnostic issues. *Acta Neuropathol*. 2014; 127:319–332. [PubMed: 24515836]
- Rutherford NJ, et al. Length of normal alleles of *C9ORF72* GGGGCC repeat do not influence disease phenotype. *Neurobiol Aging*. 2012; 33:2950.e5–2950.e7.
- Beck J, et al. Large *C9orf72* hexanucleotide repeat expansions are seen in multiple neurodegenerative syndromes and are more frequent than expected in the UK population. *Am J Hum Genet*. 2013; 92:345–353. [PubMed: 23434116]
- Simón-Sánchez J, et al. The clinical and pathological phenotype of *C9ORF72* hexanucleotide repeat expansions. *Brain*. 2012; 135:723–735. [PubMed: 22300876]
- Harms MB, et al. Lack of *C9ORF72* coding mutations supports a gain of function for repeat expansions in amyotrophic lateral sclerosis. *Neurobiol Aging*. 2013; 34:2234.e13–2234.e19.
- van der Zee J, et al. A pan-European study of the *C9orf72* repeat associated with FTLT: geographic prevalence, genomic instability, and intermediate repeats. *Hum Mutat*. 2013; 34:363–373. [PubMed: 23111906]
- van Blitterswijk M, et al. Association between repeat sizes and clinical and pathological characteristics in carriers of *C9ORF72* repeat expansions (Xpansize-72): a cross-sectional cohort study. *Lancet Neurol*. 2013; 12:978–988. [PubMed: 24011653]
- Mizielinska S, et al. *C9orf72* frontotemporal lobar degeneration is characterised by frequent neuronal sense and antisense RNA foci. *Acta Neuropathol*. 2013; 126:845–857. [PubMed: 24170096]
- Donnelly CJ, et al. RNA toxicity from the ALS/FTD *C9ORF72* expansion is mitigated by antisense intervention. *Neuron*. 2013; 80:415–428. [PubMed: 24139042]
- Lagier-Tourenne C, et al. Targeted degradation of sense and antisense *C9orf72* RNA foci as therapy for ALS and frontotemporal degeneration. *Proc Natl Acad Sci USA*. 2013; 110:E4530–E4539. [PubMed: 24170860]
- Zu T, et al. RAN proteins and RNA foci from antisense transcripts in *C9ORF72* ALS and frontotemporal dementia. *Proc Natl Acad Sci USA*. 2013; 110:E4968–E4977. [PubMed: 24248382]
- Gendron TF, et al. Antisense transcripts of the expanded *C9ORF72* hexanucleotide repeat form nuclear RNA foci and undergo repeat-associated non-ATG translation in c9FTD/ALS. *Acta Neuropathol*. 2013; 126:829–844. [PubMed: 24129584]
- Mori K, et al. The *C9orf72* GGGGCC repeat is translated into aggregating dipeptide-repeat proteins in FTLT/ALS. *Science*. 2013; 339:1335–1338. [PubMed: 23393093]
- Ash PE, et al. Unconventional translation of *C9ORF72* GGGGCC expansion generates insoluble polypeptides specific to c9FTD/ALS. *Neuron*. 2013; 77:639–646. [PubMed: 23415312]

18. Mori K, et al. Bidirectional transcripts of the expanded *C9orf72* hexanucleotide repeat are translated into aggregating dipeptide repeat proteins. *Acta Neuropathol.* 2013; 126:881–893. [PubMed: 24132570]
19. Rohrer JD, et al. *C9orf72* expansions in frontotemporal dementia and amyotrophic lateral sclerosis. *Lancet Neurol.* 2015; 14:291–301. [PubMed: 25638642]
20. Renton AE, Chio A, Traynor BJ. State of play in amyotrophic lateral sclerosis genetics. *Nat Neurosci.* 2014; 17:17–23. [PubMed: 24369373]
21. Gami P, et al. A 30-unit hexanucleotide repeat expansion in *C9orf72* induces pathological lesions with dipeptide-repeat proteins and RNA foci, but not TDP-43 inclusions and clinical disease. *Acta Neuropathol.* 2015; 130:599–601. [PubMed: 26347457]
22. Gijssels I, et al. The *C9orf72* repeat size correlates with onset age of disease, DNA methylation and transcriptional downregulation of the promoter. *Mol Psychiatry.* 2016; 21:1112–1124. [PubMed: 26481318]
23. Nordin A, et al. Extensive size variability of the GGGGCC expansion in *C9orf72* in both neuronal and non-neuronal tissues in 18 patients with ALS or FTD. *Hum Mol Genet.* 2015; 24:3133–3142. [PubMed: 25712133]
24. Fratta P, et al. Screening a UK amyotrophic lateral sclerosis cohort provides evidence of multiple origins of the *C9orf72* expansion. *Neurobiol Aging.* 2015; 36:546.e1–546.e7.
25. Xi Z, et al. Jump from pre-mutation to pathologic expansion in *C9orf72*. *Am J Hum Genet.* 2015; 96:962–970. [PubMed: 26004200]
26. Neumann M, et al. Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Science.* 2006; 314:130–133. [PubMed: 17023659]
27. Arai T, et al. TDP-43 is a component of ubiquitin-positive tau-negative inclusions in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Biochem Biophys Res Commun.* 2006; 351:602–611. [PubMed: 17084815]
28. Ling SC, Polymenidou M, Cleveland DW. Converging mechanisms in ALS and FTD: disrupted RNA and protein homeostasis. *Neuron.* 2013; 79:416–438. [PubMed: 23931993]
29. Mackenzie IR, et al. Dipeptide repeat protein pathology in *C9ORF72* mutation cases: clinico-pathological correlations. *Acta Neuropathol.* 2013; 126:859–879. [PubMed: 24096617]
30. Murray ME, et al. Clinical and neuropathologic heterogeneity of c9FTD/ALS associated with hexanucleotide repeat expansion in *C9ORF72*. *Acta Neuropathol.* 2011; 122:673–690. [PubMed: 22083254]
31. Hsiung GY, et al. Clinical and pathological features of familial frontotemporal dementia caused by *C9ORF72* mutation on chromosome 9p. *Brain.* 2012; 135:709–722. [PubMed: 22344582]
32. Mahoney CJ, et al. Frontotemporal dementia with the *C9ORF72* hexanucleotide repeat expansion: clinical, neuroanatomical and neuropathological features. *Brain.* 2012; 135:736–750. [PubMed: 22366791]
33. Irwin DJ, et al. Cognitive decline and reduced survival in *C9orf72* expansion frontotemporal degeneration and amyotrophic lateral sclerosis. *J Neurol Neurosurg Psychiatry.* 2013; 84:163–169. [PubMed: 23117491]
34. DeJesus-Hernandez M, et al. In-depth clinico-pathological examination of RNA foci in a large cohort of *C9ORF72* expansion carriers. *Acta Neuropathol.* 2017; 134:255–269. [PubMed: 28508101]
35. Mann DM, et al. Dipeptide repeat proteins are present in the p62 positive inclusions in patients with frontotemporal lobar degeneration and motor neurone disease associated with expansions in *C9ORF72*. *Acta Neuropathol Commun.* 2013; 1:68. [PubMed: 24252525]
36. Schludi MH, et al. Distribution of dipeptide repeat proteins in cellular models and *C9orf72* mutation cases suggests link to transcriptional silencing. *Acta Neuropathol.* 2015; 130:537–555. [PubMed: 26085200]
37. Davidson YS, et al. Brain distribution of dipeptide repeat proteins in frontotemporal lobar degeneration and motor neurone disease associated with expansions in *C9ORF72*. *Acta Neuropathol Commun.* 2014; 2:70. [PubMed: 24950788]

38. Mackenzie IR, et al. Quantitative analysis and clinico-pathological correlations of different dipeptide repeat protein pathologies in *C9ORF72* mutation carriers. *Acta Neuropathol.* 2015; 130:845–861. [PubMed: 26374446]
39. Davidson Y, et al. Neurodegeneration in frontotemporal lobar degeneration and motor neurone disease associated with expansions in *C9orf72* is linked to TDP-43 pathology and not associated with aggregated forms of dipeptide repeat proteins. *Neuropathol Appl Neurobiol.* 2016; 42:242–254. [PubMed: 26538301]
40. Gomez-Deza J, et al. Dipeptide repeat protein inclusions are rare in the spinal cord and almost absent from motor neurons in *C9ORF72* mutant amyotrophic lateral sclerosis and are unlikely to cause their degeneration. *Acta Neuropathol Commun.* 2015; 3:38. [PubMed: 26108573]
41. Liu EY, et al. *C9orf72* hypermethylation protects against repeat expansion-associated pathology in ALS/FTD. *Acta Neuropathol.* 2014; 128:525–541. [PubMed: 24806409]
42. Wen X, et al. Antisense proline-arginine RAN dipeptides linked to *C9ORF72*-ALS/FTD form toxic nuclear aggregates that initiate in vitro and in vivo neuronal death. *Neuron.* 2014; 84:1213–1225. [PubMed: 25521377]
43. Gendron TF, et al. Cerebellar c9RAN proteins associate with clinical and neuropathological characteristics of *C9ORF72* repeat expansion carriers. *Acta Neuropathol.* 2015; 130:559–573. [PubMed: 26350237]
44. Saberi S, et al. Sense-encoded poly-GR dipeptide repeat proteins correlate to neurodegeneration and uniquely co-localize with TDP-43 in dendrites of repeat-expanded *C9orf72* amyotrophic lateral sclerosis. *Acta Neuropathol.* 2018; 135:459–474. [PubMed: 29196813]
45. Su Z, et al. Discovery of a biomarker and lead small molecules to target r(GGGGCC)-associated defects in c9FTD/ALS. *Neuron.* 2014; 83:1043–1050. [PubMed: 25132468]
46. Gendron TF, et al. Poly(GP) proteins are a useful pharmacodynamic marker for *C9ORF72*-associated amyotrophic lateral sclerosis. *Sci Transl Med.* 2017; 9
47. Lehmer C, et al. Poly-GP in cerebrospinal fluid links *C9orf72*-associated dipeptide repeat expression to the asymptomatic phase of ALS/FTD. *EMBO Mol Med.* 2017; 9:859–868. [PubMed: 28408402]
48. Balendra R, Moens TG, Isaacs AM. Specific biomarkers for *C9orf72* FTD/ALS could expedite the journey towards effective therapies. *EMBO Mol Med.* 2017; 9:853–855. [PubMed: 28533210]
49. Panda SK, et al. Highly efficient targeted mutagenesis in mice using TALENs. *Genetics.* 2013; 195:703–713. [PubMed: 23979585]
50. Suzuki N, et al. The mouse *C9ORF72* ortholog is enriched in neurons known to degenerate in ALS and FTD. *Nat Neurosci.* 2013; 16:1725–1727. [PubMed: 24185425]
51. Koppers M, et al. *C9orf72* ablation in mice does not cause motor neuron degeneration or motor deficits. *Ann Neurol.* 2015; 78:426–438. [PubMed: 26044557]
52. Jiang J, et al. Gain of toxicity from ALS/FTD-linked repeat expansions in *C9ORF72* is alleviated by antisense oligonucleotides targeting GGGGCC-containing RNAs. *Neuron.* 2016; 90:535–550. [PubMed: 27112497]
53. Atanasio A, et al. *C9orf72* ablation causes immune dysregulation characterized by leukocyte expansion, autoantibody production, and glomerulonephropathy in mice. *Sci Rep.* 2016; 6
54. O'Rourke JG, et al. *C9orf72* is required for proper macrophage and microglial function in mice. *Science.* 2016; 351:1324–1329. [PubMed: 26989253]
55. Sudria-Lopez E, et al. Full ablation of *C9orf72* in mice causes immune system-related pathology and neoplastic events but no motor neuron defects. *Acta Neuropathol.* 2016; 132:145–147. [PubMed: 27206760]
56. Sullivan PM, et al. The ALS/FTLD associated protein *C9orf72* associates with SMCR8 and WDR41 to regulate the autophagy-lysosome pathway. *Acta Neuropathol Commun.* 2016; 4:51. [PubMed: 27193190]
57. Burberry A, et al. Loss-of-function mutations in the *C9ORF72* mouse ortholog cause fatal autoimmune disease. *Sci Transl Med.* 2016; 8:347ra93.
58. Ugolino J, et al. Loss of *C9orf72* enhances autophagic activity via deregulated mTOR and TFEB signaling. *PLoS Genet.* 2016; 12:e1006443. [PubMed: 27875531]

59. Chew J, et al. *C9ORF72* repeat expansions in mice cause TDP-43 pathology, neuronal loss, and behavioral deficits. *Science*. 2015; 348:1151–1154. [PubMed: 25977373]
60. Peters OM, et al. Human *C9ORF72* hexanucleotide expansion reproduces RNA foci and dipeptide repeat proteins but not neurodegeneration in BAC transgenic mice. *Neuron*. 2015; 88:902–909. [PubMed: 26637797]
61. O'Rourke JG, et al. *C9orf72* BAC transgenic mice display typical pathologic features of ALS/FTD. *Neuron*. 2015; 88:892–901. [PubMed: 26637796]
62. Liu Y, et al. *C9orf72* BAC mouse model with motor deficits and neurodegenerative features of ALS/FTD. *Neuron*. 2016; 90:521–534. [PubMed: 27112499]
63. Zhang YJ, et al. *C9ORF72* poly(GA) aggregates sequester and impair HR23 and nucleocytoplasmic transport proteins. *Nat Neurosci*. 2016; 19:668–677. [PubMed: 26998601]
64. Schludi MH, et al. Spinal poly-GA inclusions in a *C9orf72* mouse model trigger motor deficits and inflammation without neuron loss. *Acta Neuropathol*. 2017; 134:241–254. [PubMed: 28409281]
65. Herranz-Martin S, et al. Viral delivery of *C9orf72* hexanucleotide repeat expansions in mice leads to repeat-length-dependent neuropathology and behavioural deficits. *Dis Model Mech*. 2017; 10:859–868. [PubMed: 28550099]
66. Haeusler AR, et al. *C9orf72* nucleotide repeat structures initiate molecular cascades of disease. *Nature*. 2014; 507:195–200. [PubMed: 24598541]
67. Mizielinska S, et al. Bidirectional nucleolar dysfunction in *C9orf72* frontotemporal lobar degeneration. *Acta Neuropathol Commun*. 2017; 5:29. [PubMed: 28420437]
68. Kwon I, et al. Poly-dipeptides encoded by the *C9orf72* repeats bind nucleoli, impede RNA biogenesis, and kill cells. *Science*. 2014; 345:1139–1145. [PubMed: 25081482]
69. Tao Z, et al. Nucleolar stress and impaired stress granule formation contribute to *C9orf72* RAN translation-induced cytotoxicity. *Hum Mol Genet*. 2015; 24:2426–2441. [PubMed: 25575510]
70. Stopford MJ, et al. *C9ORF72* hexanucleotide repeat exerts toxicity in a stable, inducible motor neuronal cell model, which is rescued by partial depletion of Pten. *Hum Mol Genet*. 2017; 26:1133–1145. [PubMed: 28158451]
71. Bennion Callister J, Ryan S, Sim J, Rollinson S, Pickering-Brown SM. Modelling *C9orf72* dipeptide repeat proteins of a physiologically relevant size. *Hum Mol Genet*. 2016; 25:5069–5082. [PubMed: 27798094]
72. Lee KH, et al. *C9orf72* dipeptide repeats impair the assembly, dynamics, and function of membrane-less organelles. *Cell*. 2016; 167:774–788.e17. [PubMed: 27768896]
73. Boeynaems S, et al. Phase separation of *C9orf72* dipeptide repeats perturbs stress granule dynamics. *Mol Cell*. 2017; 65:1044–1055.e5. [PubMed: 28306503]
74. Maharjan N, Kunzli C, Buthey K, Saxena S. *C9ORF72* regulates stress granule formation and its deficiency impairs stress granule assembly, hypersensitizing cells to stress. *Mol Neurobiol*. 2017; 54:3062–3077. [PubMed: 27037575]
75. Dafinca R, et al. *C9orf72* hexanucleotide expansions are associated with altered ER calcium homeostasis and stress granule formation in iPSC-derived neurons from patients with amyotrophic lateral sclerosis and frontotemporal dementia. *Stem Cells*. 2016; 34:2063–2078. [PubMed: 27097283]
76. McGurk L, et al. Poly-A binding protein-1 localization to a subset of TDP-43 inclusions in amyotrophic lateral sclerosis occurs more frequently in patients harboring an expansion in *C9orf72*. *J Neuropathol Exp Neurol*. 2014; 73:837–845. [PubMed: 25111021]
77. Daigle JG, et al. Pur-alpha regulates cytoplasmic stress granule dynamics and ameliorates FUS toxicity. *Acta Neuropathol*. 2016; 131:605–620. [PubMed: 26728149]
78. Farg MA, et al. *C9ORF72*, implicated in amyotrophic lateral sclerosis and frontotemporal dementia, regulates endosomal trafficking. *Hum Mol Genet*. 2014; 23:3579–3595. [PubMed: 24549040]
79. Rossi S, et al. Nuclear accumulation of mRNAs underlies G4C2-repeat-induced translational repression in a cellular model of *C9orf72* ALS. *J Cell Sci*. 2015; 128:1787–1799. [PubMed: 25788698]
80. Green KM, et al. RAN translation at *C9orf72*-associated repeat expansions is selectively enhanced by the integrated stress response. *Nat Commun*. 2017; 8:2005. [PubMed: 29222490]

81. Fay MM, Anderson PJ, Ivanov P. ALS/FTD-associated *C9ORF72* repeat RNA promotes phase transitions in vitro and in cells. *Cell Rep.* 2017; 21:3573–3584. [PubMed: 29262335]
82. Zhang K, et al. Stress granule assembly disrupts nucleocytoplasmic transport. *Cell.* 2018; 173:958–971.e17. [PubMed: 29628143]
83. Cheng W, et al. C9ORF72 GGGGCC repeat-associated non-AUG translation is upregulated by stress through eIF2 α phosphorylation. *Nat Commun.* 2018; 9:51. [PubMed: 29302060]
84. Hautbergue GM, et al. SRSF1-dependent nuclear export inhibition of *C9ORF72* repeat transcripts prevents neurodegeneration and associated motor deficits. *Nat Commun.* 2017; 8
85. Yin S, et al. Evidence that C9ORF72 dipeptide repeat proteins associate with U2 snRNP to cause mis-splicing in ALS/FTD patients. *Cell Rep.* 2017; 19:2244–2256. [PubMed: 28614712]
86. Prudencio M, et al. Distinct brain transcriptome profiles in *C9orf72*-associated and sporadic ALS. *Nat Neurosci.* 2015; 18:1175–1182. [PubMed: 26192745]
87. Kanekura K, et al. Poly-dipeptides encoded by the C9ORF72 repeats block global protein translation. *Hum Mol Genet.* 2016; 25:1803–1813. [PubMed: 26931465]
88. Kramer NJ, et al. CRISPR–Cas9 screens in human cells and primary neurons identify modifiers of C9ORF72 dipeptide-repeat-protein toxicity. *Nat Genet.* 2018; 50:603–612. [PubMed: 29507424]
89. Jovicic A, et al. Modifiers of *C9orf72* dipeptide repeat toxicity connect nucleocytoplasmic transport defects to FTD/ALS. *Nat Neurosci.* 2015; 18:1226–1229. [PubMed: 26308983]
90. Almeida S, et al. Modeling key pathological features of frontotemporal dementia with *C9ORF72* repeat expansion in iPSC-derived human neurons. *Acta Neuropathol.* 2013; 126:385–399. [PubMed: 23836290]
91. Webster CP, et al. The C9orf72 protein interacts with Rab1a and the ULK1 complex to regulate initiation of autophagy. *EMBO J.* 2016; 35:1656–1676. [PubMed: 27334615]
92. Imamura K, et al. The Src/c–Abl pathway is a potential therapeutic target in amyotrophic lateral sclerosis. *Sci Transl Med.* 2017; 9
93. Aoki Y, et al. *C9orf72* and *RAB7L1* regulate vesicle trafficking in amyotrophic lateral sclerosis and frontotemporal dementia. *Brain.* 2017; 140:887–897. [PubMed: 28334866]
94. Shi Y, et al. Haploinsufficiency leads to neurodegeneration in *C9ORF72* ALS/FTD human induced motor neurons. *Nat Med.* 2018; 24:313–325. [PubMed: 29400714]
95. Onesto E, et al. Gene-specific mitochondria dysfunctions in human *TARDBP* and *C9ORF72* fibroblasts. *Acta Neuropathol Commun.* 2016; 4:47. [PubMed: 27151080]
96. Sellier C, et al. Loss of C9ORF72 impairs autophagy and synergizes with polyQ Ataxin-2 to induce motor neuron dysfunction and cell death. *EMBO J.* 2016
97. Busch JI, et al. Increased expression of the frontotemporal dementia risk factor TMEM106B causes C9orf72-dependent alterations in lysosomes. *Hum Mol Genet.* 2016; 35:1276–1297.
98. Cristofani R, et al. The small heat shock protein B8 (HSPB8) efficiently removes aggregating species of dipeptides produced in C9ORF72-related neurodegenerative diseases. *Cell Stress Chaperones.* 2018; 23:1–12. [PubMed: 28608264]
99. Cristofani R, et al. Inhibition of retrograde transport modulates misfolded protein accumulation and clearance in motoneuron diseases. *Autophagy.* 2017; 13:1280–1303. [PubMed: 28402699]
100. Jung J, et al. Multiplex image-based autophagy RNAi screening identifies SMCR8 as ULK1 kinase activity and gene expression regulator. *Elife.* 2017; 6
101. Amick J, Roczniak-Ferguson A, Ferguson SM. C9orf72 binds SMCR8, localizes to lysosomes, and regulates mTORC1 signaling. *Mol Biol Cell.* 2016; 27:3040–3051. [PubMed: 27559131]
102. Gupta R, et al. The proline/arginine dipeptide from hexanucleotide repeat expanded *C9ORF72* inhibits the proteasome. *eNeuro.* 2017; 4
103. Lee S, et al. Activation of HIPK2 promotes ER stress-mediated neurodegeneration in amyotrophic lateral sclerosis. *Neuron.* 2016; 91:41–55. [PubMed: 27321923]
104. Zhang YJ, et al. Aggregation-prone c9FTD/ALS poly(GA) RAN-translated proteins cause neurotoxicity by inducing ER stress. *Acta Neuropathol.* 2014; 128:505–524. [PubMed: 25173361]
105. May S, et al. C9orf72 FTD/ALS-associated Gly–Ala dipeptide repeat proteins cause neuronal toxicity and Unc119 sequestration. *Acta Neuropathol.* 2014; 128:485–503. [PubMed: 25120191]

106. Yamakawa M, et al. Characterization of the dipeptide repeat protein in the molecular pathogenesis of c9FTD/ALS. *Hum Mol Genet.* 2015; 24:1630–1645. [PubMed: 25398948]
107. Guo Q, et al. In situ structure of neuronal C9orf72 poly-GA aggregates reveals proteasome recruitment. *Cell.* 2018; 172:696–705.e12. [PubMed: 29398115]
108. Freibaum BD, et al. GGGGCC repeat expansion in *C9orf72* compromises nucleocytoplasmic transport. *Nature.* 2015; 525:129–133. [PubMed: 26308899]
109. Zhang K, et al. The *C9orf72* repeat expansion disrupts nucleocytoplasmic transport. *Nature.* 2015; 525:56–61. [PubMed: 26308891]
110. Xiao S, et al. Isoform-specific antibodies reveal distinct subcellular localizations of C9orf72 in amyotrophic lateral sclerosis. *Ann Neurol.* 2015; 78:568–583. [PubMed: 26174152]
111. Chou CC, et al. TDP-43 pathology disrupts nuclear pore complexes and nucleocytoplasmic transport in ALS/FTD. *Nat Neurosci.* 2018; 21:228–239. [PubMed: 29311743]
112. Shani T, Levy M, Israelson A. Assay to measure nucleocytoplasmic transport in real time within motor neuron-like NSC-34 cells. *J Vis Exp.* 2017; doi: 10.3791/55676
113. Shi KY, et al. Toxic PRn poly-dipeptides encoded by the *C9orf72* repeat expansion block nuclear import and export. *Proc Natl Acad Sci USA.* 2017; 114:E1111–E1117. [PubMed: 28069952]
114. Khosravi B, et al. Cytoplasmic poly-GA aggregates impair nuclear import of TDP-43 in *C9orf72* ALS/FTLD. *Hum Mol Genet.* 2017; 26:790–800. [PubMed: 28040728]
115. Boeynaems S, et al. *Drosophila* screen connects nuclear transport genes to DPR pathology in c9ALS/FTD. *Sci Rep.* 2016; 6
116. Coyne AN, et al. Post-transcriptional inhibition of Hsc70-4/HSPA8 expression leads to synaptic vesicle cycling defects in multiple models of ALS. *Cell Rep.* 2017; 21:110–125. [PubMed: 28978466]
117. Lopez-Gonzalez R, et al. Poly(GR) in *C9ORF72*-related ALS/FTD compromises mitochondrial function and increases oxidative stress and DNA damage in iPSC-derived motor neurons. *Neuron.* 2016; 92:383–391. [PubMed: 27720481]
118. Walker C, et al. *C9orf72* expansion disrupts ATM-mediated chromosomal break repair. *Nat Neurosci.* 2017; 20:1225–1235. [PubMed: 28714954]
119. Farg MA, Konopka A, Ying Soo K, Ito D, Atkin JD. The DNA damage response (DDR) is induced by the C9orf72 repeat expansion in amyotrophic lateral sclerosis. *Hum Mol Genet.* 2017; 26:2882–2896. [PubMed: 28481984]
120. Schweizer Burguete A, et al. GGGGCC microsatellite RNA is neuritically localized, induces branching defects, and perturbs transport granule function. *Elife.* 2015; 4
121. Ishiguro A, Kimura N, Watanabe Y, Watanabe S, Ishihama A. TDP-43 binds and transports G-quadruplex-containing mRNAs into neurites for local translation. *Genes Cells.* 2016; 21:466–481. [PubMed: 26915990]
122. Blokhuis AM, et al. Comparative interactomics analysis of different ALS-associated proteins identifies converging molecular pathways. *Acta Neuropathol.* 2016; 132:175–196. [PubMed: 27164932]
123. Yang D, et al. FTD/ALS-associated poly(GR) protein impairs the Notch pathway and is recruited by poly(GA) into cytoplasmic inclusions. *Acta Neuropathol.* 2015; 130:525–535. [PubMed: 26031661]
124. Devlin AC, et al. Human iPSC-derived motoneurons harbouring *TARDBP* or *C9ORF72* ALS mutations are dysfunctional despite maintaining viability. *Nat Commun.* 2015; 6
125. Wainger BJ, et al. Intrinsic membrane hyperexcitability of amyotrophic lateral sclerosis patient-derived motor neurons. *Cell Rep.* 2014; 7:1–11. [PubMed: 24703839]
126. Sareen D, et al. Targeting RNA foci in iPSC-derived motor neurons from ALS patients with a *C9ORF72* repeat expansion. *Sci Transl Med.* 2013; 5:208ra149.
127. Selvaraj BT, et al. *C9ORF72* repeat expansion causes vulnerability of motor neurons to Ca²⁺-permeable AMPA receptor-mediated excitotoxicity. *Nat Commun.* 2018; 9:347. [PubMed: 29367641]
128. Schanz O, et al. Cortical hyperexcitability in patients with *C9orf72* mutations: relationship to phenotype. *Muscle Nerve.* 2016; 54:264–269. [PubMed: 26799151]

129. Geevasinga N, et al. Cortical function in asymptomatic carriers and patients with *C9orf72* amyotrophic lateral sclerosis. *JAMA Neurol.* 2015; 72:1268–1274. [PubMed: 26348842]
130. Williams KL, et al. Pathophysiological insights into ALS with *C9ORF72* expansions. *J Neurol Neurosurg Psychiatry.* 2013; 84:931–935. [PubMed: 23463871]
131. Benussi A, et al. Impaired long-term potentiation-like cortical plasticity in presymptomatic genetic frontotemporal dementia. *Ann Neurol.* 2016; 80:472–476. [PubMed: 27438089]
132. Ciura S, et al. Loss of function of *C9orf72* causes motor deficits in a zebrafish model of amyotrophic lateral sclerosis. *Ann Neurol.* 2013; 74:180–187. [PubMed: 23720273]
133. Swaminathan A, et al. Expression of *C9orf72*-related dipeptides impairs motor function in a vertebrate model. *Hum Mol Genet.* 2018; 27:1754–1762. [PubMed: 29528390]
134. Swinnen B, et al. A zebrafish model for *C9orf72* ALS reveals RNA toxicity as a pathogenic mechanism. *Acta Neuropathol.* 2018; 135:427–443. [PubMed: 29302778]
135. Baldwin KR, Godena VK, Hewitt VL, Whitworth AJ. Axonal transport defects are a common phenotype in *Drosophila* models of ALS. *Hum Mol Genet.* 2016; 25:2378–2392. [PubMed: 27056981]
136. Sivadasan R, et al. *C9ORF72* interaction with cofilin modulates actin dynamics in motor neurons. *Nat Neurosci.* 2016; 19:1610–1618. [PubMed: 27723745]
137. Ferraiuolo L, et al. Oligodendrocytes contribute to motor neuron death in ALS via SOD1-dependent mechanism. *Proc Natl Acad Sci USA.* 2016; 113:E6496–E6505. [PubMed: 27688759]
138. Tran H, et al. Differential toxicity of nuclear RNA foci versus dipeptide repeat proteins in a *Drosophila* model of *C9ORF72* FTD/ALS. *Neuron.* 2015; 87:1207–1214. [PubMed: 26402604]
139. Rizzu P, et al. *C9orf72* is differentially expressed in the central nervous system and myeloid cells and consistently reduced in *C9orf72*, *MAPT* and *GRN* mutation carriers. *Acta Neuropathol Commun.* 2016; 4:37. [PubMed: 27079381]
140. Xi Z, et al. Hypermethylation of the CpG island near the G4C2 repeat in ALS with a *C9orf72* expansion. *Am J Hum Genet.* 2013; 92:981–989. [PubMed: 23731538]
141. Gijssels I, et al. A *C9orf72* promoter repeat expansion in a Flanders-Belgian cohort with disorders of the frontotemporal lobar degeneration–amyotrophic lateral sclerosis spectrum: a gene identification study. *Lancet Neurol.* 2012; 11:54–65. [PubMed: 22154785]
142. Fratta P, et al. Homozygosity for the *C9orf72* GGGGCC repeat expansion in frontotemporal dementia. *Acta Neuropathol.* 2013; 126:401–409. [PubMed: 23818065]
143. Belzil VV, et al. Reduced *C9orf72* gene expression in c9FTD/ALS is caused by histone trimethylation, an epigenetic event detectable in blood. *Acta Neuropathol.* 2013; 126:895–905. [PubMed: 24166615]
144. Waite AJ, et al. Reduced *C9orf72* protein levels in frontal cortex of amyotrophic lateral sclerosis and frontotemporal degeneration brain with the *C9ORF72* hexanucleotide repeat expansion. *Neurobiol Aging.* 2014; 35:1779.e5–1779.e13.
145. van Blitterswijk M, et al. Novel clinical associations with specific *C9ORF72* transcripts in patients with repeat expansions in *C9ORF72*. *Acta Neuropathol.* 2015; 130:863–876. [PubMed: 26437865]
146. Niblock M, et al. Retention of hexanucleotide repeat-containing intron in *C9orf72* mRNA: implications for the pathogenesis of ALS/FTD. *Acta Neuropathol Commun.* 2016; 4:18. [PubMed: 26916632]
147. Belzil VV, et al. Characterization of DNA hypermethylation in the cerebellum of c9FTD/ALS patients. *Brain Res.* 2014; 1584:15–21. [PubMed: 24530272]
148. Xi Z, et al. The *C9orf72* repeat expansion itself is methylated in ALS and FTLT patients. *Acta Neuropathol.* 2015; 129:715–727. [PubMed: 25716178]
149. Esanov R, et al. A *C9ORF72* BAC mouse model recapitulates key epigenetic perturbations of ALS/FTD. *Mol Neurodegener.* 2017; 12:46. [PubMed: 28606110]
150. Zeier Z, et al. Bromodomain inhibitors regulate the *C9ORF72* locus in ALS. *Exp Neurol.* 2015; 271:241–250. [PubMed: 26099177]
151. Bauer PO. Methylation of *C9orf72* expansion reduces RNA foci formation and dipeptide-repeat proteins expression in cells. *Neurosci Lett.* 2016; 612:204–209. [PubMed: 26690922]

152. McMillan CT, et al. *C9orf72* promoter hypermethylation is neuroprotective: neuroimaging and neuropathologic evidence. *Neurology*. 2015; 84:1622–1630. [PubMed: 25795648]
153. Russ J, et al. Hypermethylation of repeat expanded *C9orf72* is a clinical and molecular disease modifier. *Acta Neuropathol*. 2015; 129:39–52. [PubMed: 25388784]
154. Yang M, et al. A C9ORF72/SMCR8-containing complex regulates ULK1 and plays a dual role in autophagy. *Sci Adv*. 2016; 2:e1601167. [PubMed: 27617292]
155. Freischmidt A, et al. Haploinsufficiency of *TBK1* causes familial ALS and fronto-temporal dementia. *Nat Neurosci*. 2015; 18:631–636. [PubMed: 25803835]
156. Elden AC, et al. Ataxin-2 intermediate-length polyglutamine expansions are associated with increased risk for ALS. *Nature*. 2010; 466:1069–1075. [PubMed: 20740007]
157. Ferguson R, Serafeimidou-Pouliou E, Subramanian V. Dynamic expression of the mouse orthologue of the human amyotrophic lateral sclerosis associated gene *C9orf72* during central nervous system development and neuronal differentiation. *J Anat*. 2016; 229:871–891. [PubMed: 27476503]
158. Komatsu M, et al. Loss of autophagy in the central nervous system causes neurodegeneration in mice. *Nature*. 2006; 441:880–884. [PubMed: 16625205]
159. Hara T, et al. Suppression of basal autophagy in neural cells causes neurodegenerative disease in mice. *Nature*. 2006; 441:885–889. [PubMed: 16625204]
160. Robberecht W, Philips T. The changing scene of amyotrophic lateral sclerosis. *Nat Rev Neurosci*. 2013; 14:248–264. [PubMed: 23463272]
161. Fratta P, et al. *C9orf72* hexanucleotide repeat associated with amyotrophic lateral sclerosis and frontotemporal dementia forms RNA G-quadruplexes. *Sci Rep*. 2012; 2:1016. [PubMed: 23264878]
162. Reddy K, Zamiri B, Stanley SY, Macgregor RB Jr, Pearson CE. The disease-associated r(GGGGCC)_n repeat from the *C9orf72* gene forms tract length-dependent uni- and multimolecular RNA G-quadruplex structures. *J Biol Chem*. 2013; 288:9860–9866. [PubMed: 23423380]
163. Zhou B, et al. Characterizations of distinct parallel and antiparallel G-quadruplexes formed by two-repeat ALS and FTD related GGGGCC sequence. *Sci Rep*. 2018; 8
164. Vatovec S, Kovanda A, Rogelj B. Unconventional features of *C9ORF72* expanded repeat in amyotrophic lateral sclerosis and frontotemporal lobar degeneration. *Neurobiol Aging*. 2014; 35:2421.e1–2421.e12.
165. Kovanda A, Zalar M, Sket P, Plavec J, Rogelj B. Anti-sense DNA d(GGCCCC)_n expansions in *C9ORF72* form i-motifs and protonated hairpins. *Sci Rep*. 2015; 5
166. Zhang Y, Roland C, Sagui C. Structure and dynamics of DNA and RNA double helices obtained from the GGGGCC and CCCC GG hexanucleotide repeats that are the hallmark of C9FTD/ALS diseases. *ACS Chem Neurosci*. 2017; 8:578–591. [PubMed: 27933757]
167. Zamiri B, et al. Stress-induced acidification may contribute to formation of unusual structures in *C9orf72*-repeats. *Biochim Biophys Acta*. 2018; 1862:1482–1491.
168. Haeusler AR, Donnelly CJ, Rothstein JD. The expanding biology of the *C9orf72* nucleotide repeat expansion in neurodegenerative disease. *Nat Rev Neurosci*. 2016; 17:383–395. [PubMed: 27150398]
169. Lee YB, et al. Hexanucleotide repeats in ALS/FTD form length-dependent RNA foci, sequester RNA binding proteins, and are neurotoxic. *Cell Rep*. 2013; 5:1178–1186. [PubMed: 24290757]
170. Cooper-Knock J, et al. Sequestration of multiple RNA recognition motif-containing proteins by *C9orf72* repeat expansions. *Brain*. 2014; 137:2040–2051. [PubMed: 24866055]
171. Cooper-Knock J, et al. Antisense RNA foci in the motor neurons of *C9ORF72*-ALS patients are associated with TDP-43 proteinopathy. *Acta Neuropathol*. 2015; 130:63–75. [PubMed: 25943887]
172. Xu Z, et al. Expanded GGGGCC repeat RNA associated with amyotrophic lateral sclerosis and frontotemporal dementia causes neurodegeneration. *Proc Natl Acad Sci USA*. 2013; 110:7778–7783. [PubMed: 23553836]
173. Davidson YS, et al. Heterogeneous ribonuclear protein A3 (hnRNP A3) is present in dipeptide repeat protein containing inclusions in frontotemporal lobar degeneration and motor neurone

- disease associated with expansions in *C9orf72* gene. *Acta Neuropathol Commun.* 2017; 5:31. [PubMed: 28431575]
174. Conlon EG, et al. The *C9ORF72* GGGGCC expansion forms RNA G-quadruplex inclusions and sequesters hnRNP H to disrupt splicing in ALS brains. *Elife.* 2016; 5:e17820. [PubMed: 27623008]
 175. Mori K, et al. hnRNP A3 binds to GGGGCC repeats and is a constituent of p62-positive/TDP43-negative inclusions in the hippocampus of patients with *C9orf72* mutations. *Acta Neuropathol.* 2013; 125:413–423. [PubMed: 23381195]
 176. Haas S, et al. A 39-kD DNA-binding protein from mouse brain stimulates transcription of myelin basic protein gene in oligodendrocytic cells. *J Cell Biol.* 1995; 130:1171–1179. [PubMed: 7657701]
 177. Gallia GL, Johnson EM, Khalili K. Puralpha: a multifunctional single-stranded DNA- and RNA-binding protein. *Nucleic Acids Res.* 2000; 28:3197–3205. [PubMed: 10954586]
 178. Ohashi S, et al. The single-stranded DNA- and RNA-binding proteins pur alpha and pur beta link BC1 RNA to microtubules through binding to the dendrite-targeting RNA motifs. *J Neurochem.* 2000; 75:1781–1790. [PubMed: 11032866]
 179. Kanai Y, Dohmae N, Hirokawa N. Kinesin transports RNA: isolation and characterization of an RNA-transporting granule. *Neuron.* 2004; 43:513–525. [PubMed: 15312650]
 180. Celona B, et al. Suppression of *C9orf72* RNA repeat-induced neurotoxicity by the ALS-associated RNA-binding protein Zfp106. *Elife.* 2017; 6:e19032. [PubMed: 28072389]
 181. Mackenzie IR. The role of dipeptide-repeat protein pathology in *C9orf72* mutation cases. *Neuropathol Appl Neurobiol.* 2016; 42:217–219. [PubMed: 26577056]
 182. Baborie A, et al. Accumulation of dipeptide repeat proteins predates that of TDP-43 in frontotemporal lobar degeneration associated with hexanucleotide repeat expansions in *C9ORF72* gene. *Neuropathol Appl Neurobiol.* 2015; 41:601–612. [PubMed: 25185840]
 183. Mann DM. Dipeptide repeat protein toxicity in frontotemporal lobar degeneration and in motor neurone disease associated with expansions in *C9ORF72* — a cautionary note. *Neurobiol Aging.* 2015; 36:1224–1226. [PubMed: 25457023]
 184. Mattsson N, Schott JM, Hardy J, Turner MR, Zetterberg H. Selective vulnerability in neurodegeneration: insights from clinical variants of Alzheimer's disease. *J Neurol Neurosurg Psychiatry.* 2016; 87:1000–1004. [PubMed: 26746185]
 185. Mori K, et al. Reduced hnRNPA3 increases *C9orf72* repeat RNA levels and dipeptide-repeat protein deposition. *EMBO Rep.* 2016; 17:1314–1325. [PubMed: 27461252]
 186. Proudfoot M, et al. Early dipeptide repeat pathology in a frontotemporal dementia kindred with *C9ORF72* mutation and intellectual disability. *Acta Neuropathol.* 2014; 127:451–458. [PubMed: 24445903]
 187. Vatsavayai SC, et al. Timing and significance of pathological features in *C9orf72* expansion-associated frontotemporal dementia. *Brain.* 2016; 139:3202–3216. [PubMed: 27797809]
 188. Mizielinska S, et al. *C9orf72* repeat expansions cause neurodegeneration in *Drosophila* through arginine-rich proteins. *Science.* 2014; 345:1192–1194. [PubMed: 25103406]
 189. Moens TG, et al. Sense and antisense RNA are not toxic in *Drosophila* models of *C9orf72*-associated ALS/FTD. *Acta Neuropathol.* 2018; 135:445–457. [PubMed: 29380049]
 190. Chang YJ, Jeng US, Chiang YL, Hwang IS, Chen YR. The glycine–alanine dipeptide repeat from *C9orf72* hexanucleotide expansions forms toxic amyloids possessing cell-to-cell transmission properties. *J Biol Chem.* 2016; 291:4903–4911. [PubMed: 26769963]
 191. Ohki Y, et al. Glycine–alanine dipeptide repeat protein contributes to toxicity in a zebrafish model of *C9orf72* associated neurodegeneration. *Mol Neurodegener.* 2017; 12:6. [PubMed: 28088213]
 192. Flores BN, et al. Distinct *C9orf72*-associated dipeptide repeat structures correlate with neuronal toxicity. *PLoS ONE.* 2016; 11:e0165084. [PubMed: 27776165]
 193. Prpar Mihevc S, et al. Nuclear trafficking in amyotrophic lateral sclerosis and frontotemporal lobar degeneration. *Brain.* 2017; 140:13–26. [PubMed: 27497493]
 194. Lin Y, et al. Toxic PR poly-dipeptides encoded by the *C9orf72* repeat expansion target LC domain polymers. *Cell.* 2016; 167:789–802.e12. [PubMed: 27768897]

195. Lagier-Tourenne C, Polymenidou M, Cleveland DW. TDP-43 and FUS/TLS: emerging roles in RNA processing and neurodegeneration. *Hum Mol Genet.* 2010; 19:R46–R64. [PubMed: 20400460]
196. Molliex A, et al. Phase separation by low complexity domains promotes stress granule assembly and drives pathological fibrillization. *Cell.* 2015; 163:123–133. [PubMed: 26406374]
197. Patel A, et al. A liquid-to-solid phase transition of the ALS protein FUS accelerated by disease mutation. *Cell.* 2015; 162:1066–1077. [PubMed: 26317470]
198. Murakami T, et al. ALS/FTD mutation-induced phase transition of FUS liquid droplets and reversible hydrogels into irreversible hydrogels impairs RNP granule function. *Neuron.* 2015; 88:678–690. [PubMed: 26526393]
199. Jain A, Vale RD. RNA phase transitions in repeat expansion disorders. *Nature.* 2017; 546:243–247. [PubMed: 28562589]
200. Halliday M, Mallucci GR. Review: Modulating the unfolded protein response to prevent neurodegeneration and enhance memory. *Neuropathol Appl Neurobiol.* 2015; 41:414–427. [PubMed: 25556298]
201. Bowden HA, Dormann D. Altered mRNP granule dynamics in FTL D pathogenesis. *J Neurochem.* 2016; 138(Suppl 1):112–133. [PubMed: 26938019]
202. Tabet R, et al. CUG initiation and frameshifting enable production of dipeptide repeat proteins from ALS/FTD *C9orf72* transcripts. *Nat Commun.* 2018; 9:152. [PubMed: 29323119]
203. Miller TM, et al. An antisense oligonucleotide against *SOD1* delivered intrathecally for patients with *SOD1* familial amyotrophic lateral sclerosis: a phase 1, randomised, first-in-man study. *Lancet Neurol.* 2013; 12:435–442. [PubMed: 23541756]
204. Zamiri B, Reddy K, Macgregor RB Jr, Pearson CE. TMPyP4 porphyrin distorts RNA G-quadruplex structures of the disease-associated r(GGGGCC)_n repeat of the *C9orf72* gene and blocks interaction of RNA-binding proteins. *J Biol Chem.* 2014; 289:4653–4659. [PubMed: 24371143]
205. Simone R, et al. G-quadruplex-binding small molecules ameliorate *C9orf72* FTD/ALS pathology in vitro and in vivo. *EMBO Mol Med.* 2018; 10:22–31. [PubMed: 29113975]
206. Alniss H, Zamiri B, Khalaj M, Pearson CE, Macgregor RB Jr. Thermodynamic and spectroscopic investigations of TMPyP4 association with guanine- and cytosine-rich DNA and RNA repeats of *C9orf72*. *Biochem Biophys Res Commun.* 2018; 495:2410–2417. [PubMed: 29274339]
207. Kramer NJ, et al. Spt4 selectively regulates the expression of *C9orf72* sense and antisense mutant transcripts. *Science.* 2016; 353:708–712. [PubMed: 27516603]
208. Hu J, Rigo F, Prakash TP, Corey DR. Recognition of *c9orf72* mutant RNA by single-stranded silencing RNAs. *Nucleic Acid Ther.* 2017; 27:87–94. [PubMed: 28005462]
209. Pinto BS, et al. Impeding transcription of expanded microsatellite repeats by deactivated Cas9. *Mol Cell.* 2017; 68:479–490.e5. [PubMed: 29056323]
210. Batra R, et al. Elimination of toxic microsatellite repeat expansion RNA by RNA-targeting Cas9. *Cell.* 2017; 170:899–912.e10. [PubMed: 28803727]
211. Barry AE, et al. Alzheimer's disease brain-derived amyloid- β -mediated inhibition of LTP in vivo is prevented by immunotargeting cellular prion protein. *J Neurosci.* 2011; 31:7259–7263. [PubMed: 21593310]
212. Asuni AA, Boutajangout A, Quartermain D, Sigurdsson EM. Immunotherapy targeting pathological tau conformers in a tangle mouse model reduces brain pathology with associated functional improvements. *J Neurosci.* 2007; 27:9115–9129. [PubMed: 17715348]
213. Boutajangout A, Ingadottir J, Davies P, Sigurdsson EM. Passive immunization targeting pathological phospho-tau protein in a mouse model reduces functional decline and clears tau aggregates from the brain. *J Neurochem.* 2011; 118:658–667. [PubMed: 21644996]
214. Masliah E, et al. Effects of α -synuclein immunization in a mouse model of Parkinson's disease. *Neuron.* 2005; 46:857–868. [PubMed: 15953415]
215. Masliah E, et al. Passive immunization reduces behavioral and neuropathological deficits in an alpha-synuclein transgenic model of Lewy body disease. *PLoS ONE.* 2011; 6:e19338. [PubMed: 21559417]

216. Zhou Q, et al. Antibodies inhibit transmission and aggregation of *C9orf72* poly-GA dipeptide repeat proteins. *EMBO Mol Med.* 2017; 9:687–702. [PubMed: 28351931]
217. Yang YM, et al. A small molecule screen in stem-cell-derived motor neurons identifies a kinase inhibitor as a candidate therapeutic for ALS. *Cell Stem Cell.* 2013; 12:713–726. [PubMed: 23602540]

Box 1**Epigenetic modification of *C9orf72***

Analysis of blood, frontal cortex and cervical spinal cord DNA from patients with *C9orf72*-associated frontotemporal dementia and/or amyotrophic lateral sclerosis (C9FTD/ALS) has revealed that in 20–40% of cases, the CpG island in the *C9orf72* promoter region upstream of the pathogenic repeats is hypermethylated^{22,140,147,148}. This hypermethylation is associated with increased repeat length and reduced transcription of *C9orf72*. Moreover, analysis of blood DNA from patients with C9FTD/ALS showed that in 97% of cases, the expanded hexanucleotide repeat itself was methylated¹⁴⁸. A subset of *C9orf72* bacterial artificial chromosome transgenic mice also demonstrate hypermethylation of the *C9orf72* promoter and increasing hexanucleotide repeat methylation with age¹⁴⁹. Treatment of C9FTD/ALS patient fibroblasts with a DNA and histone demethylating agent led to an increase in *C9orf72* transcript levels¹⁴³. Bromodomain-containing proteins are involved in epigenetic regulation, and bromodomain inhibitors increase *C9orf72* expression in cells from patients with *C9orf72*-associated ALS and control individuals¹⁵⁰.

Histone trimethylation is another epigenetic modification that can reduce gene expression. Chromatin immunoprecipitation experiments in frontal cortex and cerebellar tissue showed that the *C9orf72* promoter region is bound to trimethylated histones in *C9orf72* repeat expansion carriers¹⁴³.

Considerable evidence points to amelioration of disease phenotypes by *C9orf72* hypermethylation. Hypermethylation is associated with reductions in RNA foci and dipeptide repeat proteins (DPRs) in patients with C9FTD/ALS⁴¹ and *C9orf72* cell models¹⁵¹, as well as reduced neuronal and grey matter loss¹⁵². In addition, hypermethylation has been linked to longer survival in patients with *C9orf72*-associated FTD¹⁵³ and a later age of onset in ALS and FTD²². However hypermethylation also correlates with reduced disease duration before death in patients with *C9orf72*-associated ALS¹⁴⁰. One possibility is that some phenotypes are dependent on gain of function and others on loss of function, so hypermethylation could have pleiotropic effects.

One might predict that reducing the levels of transcript variants 1 and 3 (Fig. 1) which are responsible for producing RNA foci and DPRs, would be protective. However, it is unclear whether reduction of variant 2 would be beneficial — this is the most highly expressed variant so is likely to be the main contributor to the functional pool of *C9orf72* protein.

Box 2**RNA-binding protein sequestration**

RNA-binding proteins (RBPs) have diverse roles in splicing, translational regulation and RNA transport and degradation. In repeat-mediated diseases, possibly including *C9orf72*-associated disease, RBPs can become sequestered by RNA foci, leading to downstream consequences.

A number of RBPs have been shown to interact with *C9orf72* repeat RNA in human tissue^{12,66,84,169–174}, *in vitro* assays^{66,84,162,170,172,174}, cell models^{66,70,79,169,170,172,174,175}, induced pluripotent stem cell-derived neurons^{12,66,126} and *in vivo* models^{79,90,169,172}. The most frequently identified RBPs that interact with *C9orf72* repeat RNA are the heterogeneous nuclear ribonucleoprotein (hnRNPs), in particular, hnRNP H, although hnRNP A1 and hnRNP A3 are also detected. Other RBPs include ALYREF, ASF/SF2, ADARB2, nucleolin, Pur- α and SRSF2. These RBPs are not found consistently across different studies, possibly reflecting the diversity of models and methodologies. However, several of these proteins have been identified in multiple independent studies, indicating that specific RBPs can be sequestered by GGGGCC repeat RNA. Only a small subset of RNA foci seem to colocalize with RBPs in human tissue¹⁷⁰ and cells⁷⁹, indicating that sequestration of these proteins is a dynamic process, or that diffuse *C9orf72* repeat RNA that is not contained within foci can sequester RBPs.

Overexpression of Pur- α , an RBP that is involved in transcription regulation^{176,177}, mRNA localization^{178,179} and stress granule formation⁷⁷ and was shown to interact with *C9orf72* repeat RNA^{79,172}, can ameliorate neurodegeneration in *C9orf72* repeat-expressing *Drosophila* and neuronal cell lines¹⁷². Overexpression of Zfp106 also suppresses toxicity in *C9orf72* repeat *Drosophila*¹⁸⁰. To date, direct evidence regarding the effects of RBP sequestration in tissue from patients with C9FTD/ALS is limited, although some hnRNP H targets are altered^{86,174}. A crucial next step is to establish how the implicated RBPs and their targets are mechanistically linked to pathogenesis.

Key points

- Rapid progress has been made in the understanding how repeat expansions in C9orf72 cause C9FTD/ALS
- Both loss of function of C9orf72 and gain of toxic function of the repeats are implicated
- A range of new models including mice, *Drosophila* and patient neurons have provided new insights
- Several cellular pathways are affected and could provide new options for treatment
- Targeted therapeutic strategies against the repeats themselves are most advanced and progressing towards clinical trials

Glossary

Repeat-associated non-ATG (RAN) translation

Translation is canonically dependent on an ATG start codon for initiation. RAN translation is a non-canonical form of translation that in the presence of repetitive sequences can start without the need for an ATG codon.

Hairpins

A secondary structure in which an RNA or DNA molecule folds back onto itself to resemble a hairpin.

G-quadruplexes

A secondary structure formed by guanine rich RNA or DNA molecules consisting of a stack of G-quartets (four guanine residues aligned in a square planar configuration).

i-motifs

A four-stranded secondary structure formed by cytosine-rich DNA or RNA molecules.

Frontotemporal lobar degeneration

Frontotemporal lobar degeneration (FTLD) describes the pathological findings observed in patients with frontotemporal dementia (FTD), however FTLD and FTD are also often used interchangeably to describe the clinical syndrome.

Bacterial artificial chromosome (BAC)

A vector for maintaining large pieces of DNA, often 50-200 kilobases in size.

Cryo-electron tomography

A high-resolution technique that involves collecting a series of tilted images of frozen hydrated samples using an electron microscope to produce a 3D reconstruction of the sample

P-bodies

Processing bodies – membraneless organelles within the cytoplasm that are involved in translational repression of mRNAs and mRNA silencing and degradation




Figure 1. *C9orf72* structure, transcript variants and protein isoforms.

The *C9orf72* gene consists of 11 exons, has three main alternatively spliced transcript variants and produces two protein isoforms. In the figure, coding exons are indicated in yellow and noncoding exons in blue (not to scale). The GGGGCC hexanucleotide repeat expansion mutation is located in the first intron of variants 1 and 3 and within the promoter region of variant 2. Variant 1 encodes C9orf72-S (short), a 222-amino-acid protein of 24 kDa, and variants 2 and 3 encode C9orf72-L (long), a 481-amino-acid protein of 54 kDa




Figure 2. Dipeptide repeat proteins.

The figure shows the dipeptide repeat proteins that are generated by GGGGCC repeat-associated non-ATG (RAN) translation. The sense strand generates poly-GA, poly-GP and poly-GR and the antisense strand generates poly-GP, poly-PA and poly-PR.




Figure 3. C9FTD/ALS neuropathology.

Sense and antisense RNA foci are a common feature in the brains of patients with *C9orf72*-associated frontotemporal dementia and/or amyotrophic lateral sclerosis (C9FTD/ALS). **a,b** | Representative images show neurons from the frontal cortex of a patient with C9FTD/ALS, containing multiple sense (red; part a) and antisense (green; part b) foci in nuclei (stained blue with DAPI). Scale bar: 2.5 μ m. **c** | TAR DNA-binding protein 43 (TDP-43) pathology in a patient with C9FTD/ALS. Arrow indicates a neuronal cytoplasmic TDP-43 inclusion in the frontal cortex, with concomitant depletion of nuclear TDP-43. Scale bar: 50 μ m. **d** |

Dipeptide repeat protein (DPR) pathology in a patient with C9FTD/ALS. Inclusions consisting of sense and antisense DPRs are produced by repeat-associated non-ATG (RAN) translation. Arrows indicate neuronal cytoplasmic inclusions of poly-GA protein. Scale bar: 50 μm .




Figure 4. Cellular processes implicated in *C9orf72*-associated FTD and ALS.

A wide range of cellular pathways have been implicated in *C9orf72*-mediated disease, several of which have previously been linked to amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD). *C9orf72* loss-of-function and toxic gain-of-function mechanisms can both alter RNA processing and metabolism pathways, with alterations in stress granules and P-bodies, and *C9orf72* gain-of-function mechanisms can lead to nucleolar dysfunction, affect RNA splicing and transcription and cause DNA damage. Proteostasis pathways have also been implicated, with impairments in autophagy and lysosomal function, the unfolded protein response and the endoplasmic reticulum, and the ubiquitin–proteasome system. Other cellular processes including nucleocytoplasmic transport, vesicular trafficking and transport granule function, and mitochondrial function, can also be impaired. In addition, neuron-specific processes, including hyperexcitability and hypoexcitability, glutamate excitotoxicity, axonal transport and neuronal branching defects, have been implicated in C9FTD/ALS. Finally, loss of *C9orf72* function alters immune system and microglial function (Table 3).

Mouse models of *C9orf72* loss of function**Table 1**

Study	Method(s)	Motor phenotypes in homozygotes	Cognitive and behavioural phenotypes in homozygotes	Other phenotypes in homozygotes	Survival in homozygotes
Clotilde Lagier-Tourenne et al. (2013)13	Somatic brain transgenesis with antisense oligonucleotide	Normal function	Normal function	None reported	Not reported
Panda et al. (2013)49	Non-conditional TALEN-mediated knockout	Not reported	Not reported	None reported	Not reported
Suzuki et al. (2013)50	Non-conditional knockout of exons 2–6	Not reported	Not reported	None reported	Not reported
Koppers et al. (2015)51	Conditional Cre-loxP-mediated knockout in neurons and glia	Normal function	Not reported	6% reduction in body weight in homozygotes compared with controls	Normal: oldest mice lived >24 months
Jiang et al. (2016)52	Non-conditional knockout of exons 2–6	Mild motor deficits on rotarod test	Mild social interaction and social recognition abnormalities	Reduced body weight, splenomegaly and cervical lymphadenopathy	Normal until 11 months, 7% of mice survived to 20 months
Atanasio et al. (2016)53	Non-conditional knockout of the full gene	Mild motor deficits at 40 weeks	Not reported	Lymphadenopathy at 12 months	9 of 17 mice survived to the end of the neurological assay period
O'Rourke et al. (2016)54	1. Non-conditional knockout of exons 2–6 2. Non-conditional knockout with zinc finger deletion	Normal function	Not reported	Cervical lymphadenopathy and splenomegaly	Normal lifespan
Sudria-Lopez et al. (2016)55	Not reported — full knockout in all tissues	Normal function	Not reported	Reduced body weight, lymphadenopathy and splenomegaly	Reduced survival: median lifespan ~500 days
Sullivan et al. (2016)56	CRISPR-Cas9-mediated non-conditional knockout	Not reported	Not reported	Lymphadenopathy and splenomegaly	Not reported
Burberry et al. (2016)57	1. Non-conditional knockout of exons 2–6 in a C57BL/6 inbred background (model 1) 2. Same as model 1, but on an outbred background (model 2) 3. CRISPR-Cas9-mediated non-conditional knockout (model 3)	Not reported	Not reported	Reduced body weight, splenomegaly, cervical lymphadenopathy and hepatomegaly	Model 1: 7% alive by 400 days Model 2: 64% alive by 300 days Model 3: reduced survival
Ugolino et al. (2017)58	Non-conditional knockout of exons 2–6	Not reported	Not reported	Splenomegaly and lethargy	Homozygotes: >50% dead in 600 days Heterozygotes: 20% dead in 600 days

A range of techniques to reduce expression of or knock out the mouse *C9orf72* orthologue have been used to investigate the normal function of *C9orf72* protein. Homozygous *C9orf72* knockouts suggest roles for the protein in immune system function, autophagy and endosomal processes. TALEN, transcription activator-like effector nuclease.

Table 2
Mouse models of *C9orf72* gain of function

Study	Mouse strain and methodology	RNA foci	DPRs detected by immunocytochemistry	DPRs detected by Immunoassay	TDP-43 pathology	Motor and cognitive phenotypes	Survival
Chew et al. (2015)59	C57BL/6J mice Somatic brain transgenesis: AAV-mediated expression of (GGGGCC) ₆₆ repeats and 119 bp upstream and 100 bp downstream <i>C9orf72</i> sequence	Sense foci throughout CNS	Poly-GA and poly-GP inclusions	Poly-GP expression in (GGGGCC) ₆₆ mice	Nuclear and occasionally cytoplasmic pTDP-43 inclusions	Rotarod impairments from day 2 of testing onwards, plus anxiety-like behaviour and hyperactivity	Not reported
Peters et al. (2015)60	SJL/B6 mice BAC transgenic, 140.5 kb upstream <i>C9orf72</i> sequence, exons 1–5, and 300 or 500 repeats	Abundant sense foci throughout CNS, antisense foci more sparse	Poly-GP inclusions, increase with age	Poly-GP throughout brain, lower than in patients with C9FTD/ALS	No TDP-43 pathology	Normal rotarod and grip strength testing, normal social behaviour	Normal survival
O'Rourke et al. (2015)61	C57BL/6J mice BAC transgenic, 110 kb upstream <i>C9orf72</i> sequence, full gene and 100–1000 repeats, 20 kb downstream sequence	Sense and antisense RNA foci throughout CNS	Poly-GP inclusions, increase with age	Soluble and insoluble poly-GP, similar levels to patients with <i>C9orf72</i> -associatedFTLD	No TDP-43 pathology	Normal grip strength, rotarod and open field testing; normal behaviour	Not reported
Liu et al. (2016)62	FVB/NJ mice BAC transgenic, 52 kb upstream <i>C9orf72</i> sequence, full gene and different lines with repeat lengths and copy numbers up to 500 repeats, 19.4 kb downstream sequence	Sense and antisense RNA foci	Poly-GA aggregates throughout CNS, increase with age	Not reported	Nuclear and cytoplasmic TDP-43 aggregates	Acute rapidly progressive disease with motor phenotype; normal open field behaviour test	Mice with acute rapidly progressive disease have decreased survival
Jiang et al. (2016)52	C57BL/6 mice BAC transgenic, 140 kb upstream <i>C9orf72</i> sequence, exons 1–5 and ~110	Sense and antisense foci	Poly-GA, poly-GP and poly-GR inclusions	2% SDS-soluble poly-GP detected	Increased levels of pTDP-43, no mislocalization or aggregates	No motor deficits; spatial and working memory deficits and anxiety found	Not reported

Study	Mouse strain and methodology	RNA foci	DPRs detected by immunocytochemistry	DPRs detected by Immunoassay	TDP-43 pathology	Motor and cognitive phenotypes	Survival
	repeats or ~450 repeats						
Zhang et al. (2016) ⁶³	C57BL/6J mice Somatic brain transgenesis: AAV-mediated, GFP-(GA) ₅₀ or GFP-(GA) ₅₀ -mutated	Not applicable	GFP-(GA) ₅₀ inclusions	Poly-GA levels twofold higher than in (GGGGCC) ₆₆ mice from the Chew et al. study ⁵⁹	Rare pTDP-43 inclusions	Motor phenotype, hyperactivity and anxiety	Not reported
Schludi et al. (2017) ⁶⁴	C57BL/6 mice Germline transgenesis: neuronal expression, (GA) ₁₄₉ -CFP + 31 carboxy-terminal amino acids from endogenous human locus	Not applicable	(GA) ₁₄₉ -CFP inclusions in brainstem, cerebellar nuclei and spinal cord, increase with age	Similar poly-GA levels in mouse spinal cord and motor neurons from patients with C9FTD/ALS	pTDP-43 levels higher in the urea-soluble fraction, as determined by ELISA; no TDP-43 inclusions or mislocalization.	Progressive gait and balance deficits; muscle strength and spatial memory normal	Not reported
Herranz-Martin et al. (2017) ⁶⁵	C57BL/6J mice Somatic brain transgenesis: AAV-mediated expression of 10 pure or 102 interrupted (by TCGAG linker) GGGGCC repeats No <i>C9orf72</i> flanking sequence	Sense foci throughout CNS, less frequent in spinal cord	Poly-GA, primarily in cerebellum and brainstem	Not reported	Infrequent TDP-43 aggregates found equally in 10 pure or 102 interrupted repeats	Progressive gait and behavioural deficits on open field and novel object recognition	Not reported

Several mouse gain-of-function models are described. The approaches used have included somatic brain or germline transgenesis and expression of BAC constructs containing the full-length or partial human *C9orf72* gene and repeat expansion. AAV, adeno-associated virus; BAC, bacterial artificial chromosome; C9FTD/ALS, *C9orf72*-associated frontotemporal dementia and/or amyotrophic lateral sclerosis; CFP, cyan fluorescent protein; DPRs, dipeptide repeat proteins; ELISA, enzyme-linked immunosorbent assay; FTL, frontotemporal lobar degeneration; GFP, green fluorescent protein; pTDP-43, phospho-TDP-43; SDS, sodium dodecyl sulphate; TDP-43, TAR DNA-binding protein 43.

Table 3

Downstream mechanisms implicated in C9FTD/ALS

	C9orf72 human tissue/model				Other C9orf72 in vitro models				C9orf72 in vivo models			
	Patient-derived neurons/glia	Post-mortem tissue	Other system/tissue/in patients	Cell lines	Primary neurons	Yeast	Worm and zebrafish	Fly	Mouse			
RNA metabolism												
Nucleolar function/LLPS	Yes66	Yes42,66,67	Yes42,66	Yes68-73	Yes36	-	-	Yes67,72	-			
Processing bodies	Yes66	-	-	Yes74	Yes42,74	-	-	-	-			
Stress granules/LLPS	Yes75	Yes76	Yes77	Yes69,72-74,78-83	Yes42,74	-	-	Yes72	-			
RNA processing, splicing, transcription and transport	Yes84,85	Yes84-86	-	Yes68,84,85,87,88	Yes84	Yes89	-	Yes84	-			
Nuclear speckles/LLPS	-	-	-	Yes72	-	-	-	Yes72	-			
Cajal bodies/LLPS	-	-	-	Yes72	-	-	-	Yes72	-			
Proteostasis												
Autophagy and lysosomal function	Yes75,90-94	Yes54	Yes93,95	Yes54,58,78,88,91,96-101	Yes91,96,102	-	-	-	Yes54,56,58			
Endoplasmic reticulum/UPR	Yes66,75	Yes86,103,104	-	Yes80,83,88	Yes80,104	-	-	-	-			
Translational inhibition	-	-	-	Yes79,80,87	-	-	-	-	-			
Ubiquitin-proteasome system	-	-	-	Yes88,99,105,106	Yes63,102,104,107	-	-	-	Yes63			
Other cellular processes												
Nucleocytoplasmic transport	Yes82,84,89,108,109	Yes86,109-111	-	Yes68,79,82,84,88,108,110-114	Yes84,89	Yes89	-	Yes82,84,108,109,115	Yes63			
Vesicle trafficking	Yes93	Yes86	Yes93	Yes78,96	-	-	-	Yes116	-			
DNA damage	Yes117	Yes118,119	-	Yes118,119	Yes118	-	-	Yes117	-			
Transport granule function	Yes120	Yes120	-	Yes121	Yes120	-	-	Yes120	-			
Mitochondrial function	Yes75,117	-	Yes95	Yes122	-	-	-	-	-			
Notch signalling	Yes123	Yes123	-	-	-	-	-	Yes123	-			
Arginine methylation	-	-	-	Yes115	-	-	-	Yes115	-			
Nervous system-specific processes												
Neuronal excitability and glutamate toxicity	Yes12,94,124-127	Yes12,124-126	Yes128-131	Yes71	-	-	-	-	Yes94			
Neuronal branching and growth	-	Yes105	-	-	Yes104,105,120	-	Yes132-134	Yes120,123	-			
Axonal transport	-	-	-	-	-	-	-	Yes135	-			

	<i>C9orf72</i> human tissue/model			Other <i>C9orf72</i> <i>in vitro</i> models			<i>C9orf72</i> <i>in vivo</i> models			
	Patient-derived neurons/glia	Post-mortem tissue	Other system/tissue/in patients	Cell lines	Primary neurons	Yeast	Worm and zebrafish	Fly	Mouse	
Actin/growth cone dynamics	Yes136	-	-	-	Yes136	-	-	-	-	
Glial dysfunction	Yes137	-	-	-	-	-	-	-	-	
Systemic functions										
Immune system function	-	Yes54	-	Yes54	-	-	-	-	Yes (Table 1)	

Several studies (referenced within the relevant cell of the table) have identified a spectrum of downstream processes involved in *C9orf72*-mediated disease, using patient-derived tissue, cells or induced pluripotent stem cell-derived neurons, and *in vitro* or *in vivo* disease models. Globally, these processes can be classified into those involved in RNA metabolism and processing, proteostasis, other cellular processes, nervous system-specific functions or systemic functions. LLPS, liquid-liquid phase separation; UPR, unfolded protein response.