



Published in final edited form as:

*Clin Cancer Res.* 2012 April 01; 18(7): 2099–2107. doi:10.1158/1078-0432.CCR-11-2484.

## Sorafenib is an Inhibitor of UGT1A1 but is Metabolized by UGT1A9: Implications of Genetic Variants on Pharmacokinetics and Hyperbilirubinemia

Cody J. Peer<sup>1,\*</sup>, Tristan M. Sissung<sup>1,\*</sup>, AeRang Kim<sup>2</sup>, Lokesh Jain<sup>1</sup>, Sukyung Woo<sup>1</sup>, Erin R. Gardner<sup>1</sup>, C. Tyler Kirkland<sup>3</sup>, Sarah M. Troutman<sup>3</sup>, Bevin C. English<sup>3</sup>, Emily D. Richardson<sup>3</sup>, Joel Federspiel<sup>1</sup>, David Venzon<sup>4</sup>, William Dahut<sup>5</sup>, Elise Kohn<sup>5</sup>, Shivaani Kummar<sup>5</sup>, Robert Yarchoan<sup>6</sup>, Giuseppe Giaccone<sup>5</sup>, Brigitte Widemann<sup>2</sup>, and William D. Figg<sup>1,3</sup>

<sup>1</sup>Clinical Pharmacology Program, National Cancer Institute, Bethesda, MD;

<sup>2</sup>Pharmacology and Experimental Therapeutics Section, National Cancer Institute, Bethesda, MD;

<sup>3</sup>Molecular Pharmacology Section, National Cancer Institute, Bethesda, MD;

<sup>4</sup>Biostatistics and Data Management Branch, National Cancer Institute, Bethesda, MD;

<sup>5</sup>Medical Oncology Branch, National Cancer Institute, Bethesda, MD;

<sup>6</sup>HIV/AIDS Malignancy Branch, National Cancer Institute, Bethesda, MD;

### Abstract

**Purpose:** Several case reports suggest sorafenib exposure and sorafenib-induced hyperbilirubinemia may be related to a (TA)<sub>5/6/7</sub> repeat polymorphism in *UGT1A1\*28*. We hypothesized that sorafenib inhibits UGT1A1 and individuals carrying *UGT1A1\*28* and/or UGT1A9 variants experience greater sorafenib exposure and greater increase in sorafenib-induced plasma bilirubin concentration.

**Experimental Design:** Inhibition of UGT1A1-mediated bilirubin glucuronidation by sorafenib was assessed *in vitro*. *UGT1A1\*28* and *UGT1A9\*3* genotypes were ascertained using fragment analysis or direct sequencing in 120 cancer patients receiving sorafenib on five different clinical trials. Total bilirubin measurements were collected in prostate cancer patients prior to receiving sorafenib ( $n=41$ ) and 19–30 days following treatment and were compared to *UGT1A1\*28* genotype.

---

Corresponding author: William D. Figg, Medical Oncology Branch, CCR, NCI/NIH, 9000 Rockville Pike, Building 10, Room 5A01, Bethesda, Maryland, 20892 Tel: +1-301-402-3622 Fax: +1 301-402-8606, wf13e@nih.gov.

\*Authors had an equal contribution.

Authorship Contributions

*Participated in research design:* Peer, Sissung, Jain, Gardner, Dahut, Giaccone, Kummar, Kohn, Yarchoan

*Conducted experiments:* Peer, Kirkland, Troutman, English, Richardson, Federspiel

*Performed data analysis:* Peer, Sissung, Venzon, Jain, Woo

*Wrote or contributed to the writing of the manuscript:* Peer, Sissung, Kim, Woo, Widemann, Kirkland, Troutman, Richardson, Venzon, Yarchoan, Figg

Disclaimer

The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organization imply endorsement by the U.S. Government.

**Results:** Sorafenib exhibited mixed-mode inhibition of UGT1A1-mediated bilirubin glucuronidation ( $IC_{50}=18\mu\text{M}$ ;  $K_i=11.7\mu\text{M}$ ) *in vitro*. Five patients carrying *UGT1A1\*28/\*28* ( $n=4$ ) or *UGT1A9\*3/\*3* ( $n=1$ ) genotypes had first-dose, dose-normalized sorafenib AUCs that were in the 93<sup>rd</sup> percentile, while three patients carrying *UGT1A1\*28/\*28* had AUCs in the bottom quartile of all genotyped patients. DMET genotyping on six patients revealed the *ABCC2-24C>T* genotype cosegregated with sorafenib AUC phenotype. Sorafenib exposure was related to plasma bilirubin increases in patients carrying 1 or 2 copies of *UGT1A1\*28* alleles ( $n=12$  and  $n=5$ ;  $R^2=0.38$  and  $R^2=0.77$ ;  $P=0.032$  and  $P=0.051$ , respectively). *UGT1A1\*28* carriers demonstrated two distinct phenotypes that could be explained by *ABCC2-24C>T* genotype and are more likely to experience plasma bilirubin increases following sorafenib if they had high sorafenib exposure.

**Conclusions:** This pilot study indicates that genotype status of *UGT1A1*, *UGT1A9*, and *ABCC2* and serum bilirubin concentration increases reflect abnormally high AUC in patients treated with sorafenib.

### Keywords

Sorafenib; glucuronidation; hyperbilirubinemia; pharmacokinetics; pharmacogenetics

## INTRODUCTION

Sorafenib tosylate inhibits VEGF receptors and several tyrosine kinases and thus is considered an antiangiogenic agent with dual multikinase activity (1). Although sorafenib is currently approved for the treatment of renal cell and unresectable hepatocellular carcinomas, it may also have activity in other solid tumors, which led to the initiation of several sorafenib-based clinical trials (2–5). There is wide variation in the response and toxicity between patients following sorafenib treatment that appears to be, at least in part, related to cumulative drug exposure (3). For this reason, studies investigating the sources of inter-individual variation in sorafenib exposure are needed.

In humans, the majority (77%) of the sorafenib dose is either not absorbed or is eliminated through the hepatobiliary route (50% unchanged), while 19% of the dose (mostly glucuronides) is excreted in urine (6). Both routes of elimination require glucuronidation catalyzed by UGT1A9 (7, 8), although it remains unclear if other UGTs are responsible for glucuronidation of oxidized sorafenib metabolites formed through CYP3A4/5 metabolism. Moreover, these enzymes demonstrate phenotypic variability based on multiple polymorphisms, (i.e., *UGT1A9\*3*, *UGT1A9-118<sub>d</sub>T<sub>9/10</sub>*, *UGT1A9IVS+1 399 C>T* (9); *CYP3A4\*1b* and *CYP3A5\*3C* (10)). Once in the gut, intestinal microflora deglucuronidate and reduce sorafenib resulting in enterohepatic circulation allowing systemic re-exposure (10, 11). However, renal elimination appears to be irreversible and individuals with low creatinine clearance ( $\text{CrCl}<60\text{mL}/\text{min}$ ) require more sorafenib dose reductions than patients with normal renal function (12). Thus, sorafenib glucuronidation is a significant route of sorafenib metabolism (6, 8) and can potentially alter sorafenib exposure.

It is known that seven TA nucleotide repeats in the  $(\text{TA})_n\text{TAA}$  promoter region of *UGT1A1* (*UGT1A1\*28*) leads to decreased expression of UGT1A1, resulting in high plasma bilirubin levels and is often diagnosed as Gilbert's syndrome (13). Previous reports suggested that

bilirubin concentrations were elevated by sorafenib (14, 15). Interestingly, one report suggested that sorafenib induced jaundice in individuals carrying *UGT1A1\*28* alleles due to a proposed UGT1A1 inhibition (14). This is consistent with three additional reports that also suggested sorafenib might inhibit UGT1A1-mediated bilirubin glucuronidation resulting in elevated bilirubin concentration (14–17). Another study profiled a patient receiving sorafenib who had yellow skin coloration despite a normal serum concentration of bilirubin and determined that the outcome was likely, if not definitely, attributable to sorafenib treatment (18). Furthermore, a phase I dose-escalation trial ( $n=34$ ) of sorafenib with irinotecan, a UGT1A1 and UGT1A9 substrate, resulted in elevated irinotecan and SN-38 exposure with the highest sorafenib dose (400 mg BID) (17). In that study, sorafenib was reported to have an *in vitro* inhibitor constant ( $K_i$ ) of 2.7  $\mu\text{M}$  in human liver microsomes. This suggested that the increased SN-38 exposure was due to sorafenib-induced inhibition of UGT1A1- and/or UGT1A9-mediated SN-38 glucuronidation. However, none of the above case reports evaluated sorafenib plasma concentration in these patients, thus further confirmation of these results in larger patient cohorts undergoing sorafenib treatment is needed.

Herein, we present a case report of a child with Gilbert's syndrome who underwent sorafenib treatment and experienced abnormally high sorafenib exposure. Based on this observation and the aforementioned case studies, we hypothesized that sorafenib and/or CYP3A4/5-mediated sorafenib oxide may be glucuronidated by UGT1A1 and/or UGT1A9, and that sorafenib acts as an inhibitor of UGT1A1-mediated bilirubin glucuronidation. Additionally, sorafenib AUC and sorafenib-induced hyperbilirubinemia might be related to the *UGT1A1\*28* allele that is responsible for most cases of Gilbert's syndrome. Since UGT1A9 and CYP3A4/5 are known to metabolize sorafenib (6, 8), we hypothesized that sorafenib exposure would also be related to allelic variation in the genes encoding these enzymes (19). To this end, we compared sorafenib AUC with genetic variation in *CYP3A4/5* and *UGT1A1/9* in patients with various solid tumors undergoing sorafenib therapy, as well as with sorafenib-related toxicities (hand-foot skin reaction; HFSR) and clinical outcome.

## METHODS

### Materials

The following chemicals were purchased from their respective suppliers: Sorafenib tosylate (CTEP, c/o Bayer Schering Pharma, Berlin, Germany), Human CYP3A4 supersomes containing CYP450 oxidoreductase and NADPH generating system, 0.5 M potassium phosphate, pH 7.4, 100 mM Tris buffer, pH 7.4, Human UGT1A1 and UGT1A9 Supersomes and UGT Reaction Mix (BD Biosciences, San Jose, CA), methanol and acetonitrile (Optima grade, Fisher Scientific, Pittsburgh, PA), formic acid and acetic acid (Sigma-Aldrich, St. Louis, MO).  $\beta$ -Glucuronidase was purchased from Roche (Roche Diagnostics, Indianapolis, IN). All water used was deionized and purified using a Millipore system.

### In vitro studies

Sorafenib and cytochrome P450 (CYP)-mediated oxidized sorafenib (sorafenib-N-oxide; M-2) were subjected to uridine glucuronosyl transferase (UGT)-catalyzed glucuronidation

by members of the UGT1A family, UGT1A1 and UGT1A9. Furthermore, the role of sorafenib as an inhibitor of UGT1A1-mediated bilirubin glucuronidation was also studied. Details are discussed in Supplemental Methods.

### Patients and Treatment

Patients ( $n=120$ ) from five clinical trials involving sorafenib treatment were used for subsequent pharmacogenetic analysis, consisting of two phase I trials and three phase II trials. The phase I trials were BAY-BEV (200 mg bid sorafenib with bevacizumab;  $n=27$ ) (3), and BAY-KS (200 mg qd or 200–400 mg bid sorafenib with ritonavir in Kaposi's sarcoma;  $n=8$ ; data not yet published), AUC data from patients on the BAY-KS trial who received ritonavir were not available; thus potential AUC-influencing drug-drug interactions between ritonavir and sorafenib were not accounted for in future analyses. The phase II trials were BAY-CRPC (400 mg bid sorafenib in castration resistant prostate cancer;  $n=46$ ) (2, 4), BAY-NSCLC (400 mg bid sorafenib in non small-cell lung cancer;  $n=22$ ) (20, 21), and BAY-CRC (400 mg bid sorafenib with cetuximab in colorectal cancer;  $n=17$ ; data not yet published). Written informed consent was obtained from all patients before enrollment on the trials and genotyping was approved by the Institutional Review Board of the National Cancer Institute. All inclusion/exclusion criteria and genotyping methods are detailed in the Supplemental Methods.

### Sorafenib Exposure

Exposure (AUC) data were represented as day 1 dose-normalized  $AUC_{0-12h}$  (ng-h/mL/mg), as previously reported (22). Steady-state exposures were not available for all 120 patients. Furthermore, exposure values were dose-normalized to compare AUC from the five different trials with patients administered different doses. Linear pharmacokinetics were not assumed from first-dose AUC values, rather individual patient exposures were correlated to physiological changes (i.e. HFSR, PFS, and plasma bilirubin concentration).

### Statistical considerations

Hardy-Weinberg equilibrium was tested using the chi-squared test. Genetic linkage statistics were obtained using Haploview (Broad Institute, Cambridge, MA). Comparisons of genotype versus demographics, preclinical measures, and pharmacokinetics were conducted with nonparametric statistical tests including the Wilcoxon Rank-Sum test or Kruskal Wallis ANOVA. Kruskal Wallis ANOVA was also used to compare sorafenib AUC to the clinical grade of hand-foot skin reactions (HFSR). Fisher's Exact Test was used to compare between different AUC percentiles and genotype. Linear regression was conducted to compare the creatinine clearance (CrCl) and serum glutamic oxaloacetic transaminase (SGOT) to sorafenib AUC and was also utilized in comparisons between AUC and the bilirubin change from baseline in the different genotype groupings. Cox model analysis was conducted to compare genotypes versus progression-free survival (PFS). Statistical significance was assigned if  $P<0.05$  as this study was conducted in an exploratory mode for potential confirmation in independent data.

## RESULTS

### Case Report

The patient was a 12 year-old boy who enrolled on a phase I trial of sorafenib for children with neurofibromatosis type 1 and inoperable plexiform neurofibromas (23). Prior to treatment, he had a total elevated bilirubin of 2.0 mg/dL with a direct of 0.3 mg/dL, normal serum ALT and AST, and was clinically diagnosed with Gilbert's syndrome. This was later confirmed with genetic testing as he was found to be homozygous for the A(TA)<sub>7</sub>TAA allele of the *UGT1A1* gene (i.e., *UGT1A1*\*28/\*28). The protocol required a bilirubin concentration within normal limits for study entry, except for patients with Gilbert's syndrome. He was treated with a dose of 115 mg/m<sup>2</sup> twice daily (approximately 50% of the adult maximum tolerated dose (MTD) based on an average adult body surface area of 1.8 m<sup>2</sup>).

The patient had day 1 pharmacokinetics performed, and his AUC<sub>0-24h</sub> was noted to be 81 µg·h/mL, which is greater than the average AUC<sub>0-24h</sub> of 28 ± 17 µg·h/mL observed in children treated at the MTD of 200 mg/m<sup>2</sup> on the refractory solid tumor phase I trial (24). The patient came off treatment after 9 days due to dose limiting grade 3 tumor pain. The protocol was subsequently amended to exclude patients with known Gilbert's syndrome from trial participation.

Based on this case observation and the published case reports that sorafenib induced hyperbilirubinemia in a small number of patients who carry a *UGT1A1*\*28 allele (14, 15), we hypothesized that genetic variation in *UGT1A1* was a potential source of alterations in sorafenib exposure and sorafenib-induced hyperbilirubinemia. Moreover, since *UGT1A9* is known to primarily glucuronidate sorafenib (8), we hypothesized that *UGT1A9* alleles might also contribute to both endpoints; thus, we studied sorafenib and sorafenib-N-oxide glucuronidation by *UGT1A1* and *UGT1A9* *in vitro* (Supplemental Methods and Results), and ascertained *UGT1A1* and *UGT1A9* genotypes in patients treated with sorafenib for comparison with pharmacokinetics and sorafenib-induced hyperbilirubinemia endpoints. Patient characteristics are reported in Table 1.

### *In Vitro* Sorafenib Glucuronidation

Sorafenib glucuronidation by *UGT1A9* was confirmed via *in vitro* metabolism experiments with recombinant *UGT1A9* via liquid chromatography-mass spectrometry (LC-MS), whereas a similar experiment using *UGT1A1* did not metabolize sorafenib (see Supplemental Results). This is in agreement with literature (7, 8). Based on exploratory studies, neither *UGT1A1* nor *UGT1A9* glucuronidated the CYP3A4-mediated sorafenib-N-oxide (M-2) (data not shown).

Based on previous reports (14, 15, 17), we hypothesized that *UGT1A1* could bind sorafenib and that this binding event could inhibit bilirubin glucuronidation. Increasing amounts of sorafenib were added to an *in vitro* *UGT1A1*-catalyzed bilirubin glucuronidation enzyme activity assay to determine the extent and mechanism of inhibition by sorafenib. Bilirubin concentrations used were based on literature reports of its K<sub>m</sub> for *UGT1A1*-mediated glucuronidation, ranging between 0.2–26 µM (25–28). Graphical modeling suggested

sorafenib best fits a mixed-mode (mixed-type) inhibitor of UGT1A1 (model correlation  $r=0.96$ ), which demonstrates properties of both a competitive and noncompetitive inhibitor (Supplemental Results). Based on model correlations (noncompetitive  $r=0.93$ ; competitive  $r=0.88$ ), it was suggested that while sorafenib is a mixed-type inhibitor, it exhibits more noncompetitive-type inhibitor characteristics than competitive.

A bilirubin  $K_m$  of  $5.9 \mu\text{M}$  and an inhibitor constant ( $K_i$ ) of  $11.8 \mu\text{M}$  were determined empirically through the mixed-type model. The  $\text{IC}_{50}$  of sorafenib for UGT1A1-mediated bilirubin glucuronidation was determined to be  $18 \mu\text{M}$  following a separate experiment (see Supplemental Methods). These values were slightly higher than the literature values obtained by studying the mixed-type sorafenib-mediated inhibition of SN38 glucuronidation by UGT1A1, where  $K_i$  was found to be  $2.7 \mu\text{M}$  (17). The most likely reason for this discrepancy from literature is due to the different substrates used in the experiment (SN38 vs bilirubin). Since sorafenib demonstrates a competitive inhibition factor (model correlation 0.88), the substrate likely has an effect on inhibitor  $K_i$ .

### Genotype versus sorafenib exposure

Based on the case report presented above, data indicating that sorafenib is glucuronidated by UGT1A9 (Supplemental Results), and the sorafenib-mediated inhibition of bilirubin glucuronidation through UGT1A1 (Supplemental Results), we next hypothesized that genetic variants in *UGT1A1* and *UGT1A9* would affect sorafenib pharmacokinetics and bilirubin metabolism in patients with solid tumors who received sorafenib.

We excluded patients with low CrCl ( $<60\text{mL}/\text{min}$ ) and a single patient with abnormally high SGOT ( $90\text{U}/\text{L}$ ) given the importance of hepatic and renal function in sorafenib PK (Supplemental Results). Carriers of *UGT1A1*\*28/\*28 tended to be younger and have higher median total bilirubin. Patients with wild-type *CYP3A4* and *CYP3A5* alleles had lower median SGOT and median CrCl than variant allele carriers, respectively. A more detailed summary can be found in the Supplemental Results section. None of the above associations are likely to have altered the results presented in later sections.

Initial analysis of *UGT1A1* A(TA)<sub>n</sub>TAA and *UGT1A9*\*3 genotypes versus day 1 dose-normalized sorafenib  $\text{AUC}_{0-12\text{h}}$  revealed five patients carrying *UGT1A1*\*28/\*28 ( $n=4$ ) and *UGT1A9*\*3/\*3 ( $n=1$ ) have higher AUCs than those patients corresponding to any other genotypes with normal creatinine clearance (CrCl) and SGOT (i.e., all  $> 93^{\text{rd}}$  percentile; range =  $109.9 - 198.6 \text{ ng}\cdot\text{h}/\text{mL}/\text{mg}$ ), and one patient carrying *UGT1A1*\*28/\*28 had an AUC in the  $76^{\text{th}}$  percentile (AUC =  $72.5 \text{ ng}\cdot\text{h}/\text{mL}/\text{mg}$ ). Interestingly, of the three remaining patients carrying *UGT1A1*\*28/\*28, two had the lowest AUCs (i.e.  $< 3^{\text{rd}}$  percentile; range =  $2.7 - 6.3 \text{ ng}\cdot\text{h}/\text{mL}/\text{mg}$ ) while one had an AUC in the  $21^{\text{st}}$  percentile (AUC =  $18.0 \text{ ng}\cdot\text{h}/\text{mL}/\text{mg}$ ). For this reason, *UGT1A1*\*28 status was considered to confer different phenotypes: those having an abnormally high exposure, those having exposures matching the rest of the cohort, and those having low exposure (Figure 1). Analysis of *UGT1A1* and *UGT1A9* genotypes versus sorafenib AUC in the different genotype groupings did not lead to a statistically significant result due to the wide variability in phenotype in the *UGT1A1*\*28/\*28 genotype grouping ( $P=0.32$ ; Kruskal-Wallis ANOVA; Figure 1); however, there were strongly significant differences in the odds of having AUCs  $>93^{\text{rd}}$



percentile (i.e., 107 ng·h/mL/mg) and also carrying *UGT1A1*\*28/\*28 ( $n=4$ ) or *UGT1A1*\*3/\*3 ( $n=1$ ) (OR (95% CI) = 179.7 (8.5 – 3787);  $P<0.0001$ ; Fisher's Exact Test). In addition, there was a strongly significant difference in the odds of carrying *UGT1A1*\*28/\*28 and also having AUCs  $\leq$  3<sup>rd</sup> percentile (OR (95% CI) = 57.3 (2.5 – 1326);  $P=0.0084$ ; Fisher's Exact Test), further justifying consideration of *UGT1A1*\*28 alleles as conferring different phenotypes. Although *UGT1A1* IVS+1 (399C>T) and -118dT<sub>9/10</sub> were in linkage disequilibrium with *UGT1A1* A(TA)<sub>n</sub>TAA (Supplementary Results), these polymorphisms were not associated with alterations in AUC ( $P=0.39$ ; data not shown). Neither *CYP3A4*\*1B ( $P=0.42$ ) nor *CYP3A5*\*3C ( $P=0.52$ ) status was associated with increased sorafenib exposure. Therefore, *UGT1A1*\*28 and possibly the *UGT1A1*\*3 SNP appeared to be the major predictive alleles associated with phenotype.

To further study the apparently different phenotypes for patients carrying only *UGT1A1*\*28 alleles ( $n=8$ ), we genotyped 1,936 polymorphisms in 225 genes involved in clinical pharmacology using the DMET Plus panel (Coriell Institute, N.J., USA). DMET genotyping was only successful (90% call rate) in a total of 6 patients with the following AUCs (ng·h/mL/mg): 2.7, 6.3, 18.0, 109.9, 116.2, 139.6. DMET analysis revealed that only the *ABCC2*-24C>T SNP cosegregated with sorafenib metabolism phenotype. After sequencing the *ABCC2*-24C>T SNP in all patients carrying *UGT1A1*\*28/\*28, it was determined that the patient with the lowest observed AUC (i.e. 2.7 ng·h/mL/mg) was double variant, whereas patients with the next lowest AUCs (6.3, 18.0, and 72.5 ng·h/mL/mg) were heterozygous followed by those with the highest AUCs (109.9, 116.2, and 139.6) who carried homozygous wild-type alleles. A single patient with AUC = 198.6 ng·h/mL/mg was not ascertainable as *ABCC2* genotyping by direct sequencing was not successful. Upon sequencing the whole population for *ABCC2*-24C>T, it was determined that individuals carrying only variant alleles in this SNP tended to have lower median AUC (29.8 versus 40.5 ng·h/mL/mg) compared to individuals carrying one or two copies of wild-type allele, but this was not statistically significant ( $P=0.21$ ). Therefore, the *ABCC2*-24C>T SNP only appears to modify AUC phenotype in those carrying only *UGT1A1*\*28 alleles.

### Genotype versus bilirubin change following sorafenib

Since previous case-report data indicated that sorafenib might induce bilirubin changes in patients based on *UGT1A1* allele status (14, 15), we hypothesized that sorafenib exposure would correspond to greater increases in post-sorafenib bilirubin concentration in those patients with low-functioning *UGT1A1* alleles (i.e., *UGT1A1*\*28). Analysis of bilirubin versus genotype was only conducted in men with prostate cancer receiving sorafenib as comprehensive bilirubin data were not obtained in other trials. The median change in bilirubin plasma concentration was 0 mg/dL (range = -0.3 to 0.5 mg/dL;  $n=45$ ). A total of 3 patients with normal CrCL developed hyperbilirubinemia (i.e., bilirubin concentration  $\geq$  1.0mg/dL) following sorafenib (*UGT1A1* (TA)<sub>5</sub>/(TA)<sub>6</sub>  $n=1$ , (TA)<sub>6</sub>/(TA)<sub>7</sub>  $n=2$ ), and 2 patients that presented with hyperbilirubinemia prior to the sorafenib dose had a further rise in bilirubin concentration following sorafenib (*UGT1A1* (TA)<sub>6</sub>/(TA)<sub>7</sub>  $n=1$ , (TA)<sub>7</sub>/(TA)<sub>7</sub>  $n=1$ ; the latter patient had a 0.4mg/dL increase).

*UGT1A1* A(TA)<sub>n</sub>TAA status was not related to change in bilirubin from baseline ( $P=0.39$ ; Figure 2A). However, regression analysis indicated that sorafenib exposure was related to bilirubin serum concentration in patients with normal CrCl ( $R^2=0.29$ ;  $P=0.0005$ ; Figure 2B). When regression analyses were stratified on the basis of *UGT1A1* A(TA)<sub>n</sub>TAA genotype status, this analysis revealed that sorafenib exposure was not related to bilirubin increases in patients carrying either *UGT1A1* (TA)<sub>5/5</sub> or (TA)<sub>5/6</sub>, or *UGT1A1* (TA)<sub>6/6</sub> genotypes ( $R^2=0.43$  and  $R^2=0.030$  respectively;  $P=0.35$  and  $P=0.51$ , respectively; Figure 2C–D). However, only 4 individuals carried a copy of *UGT1A1* (TA)<sub>5</sub>, and there is an apparent (albeit non-significant) proportional increase in both AUC and sorafenib-induced bilirubin changes consistent with the rather high  $R^2$  for this genotype grouping. When the data were stratified by *UGT1A1* (TA)<sub>6/7</sub> and *UGT1A1* (TA)<sub>7/7</sub> genotypes, a significant (or marginally non-significant) relationship was observed in both cases with a relatively high correlation ( $R^2=0.38$  and  $R^2=0.77$  respectively;  $P=0.032$  and  $P=0.051$ , respectively; Figure 2E–F). For patients with the *UGT1A1* (TA)<sub>6/7</sub> genotype, the data indicate that bilirubin increased by 0.1 mg/dL for every 25.7 (ng/mL\*hr/mg) unit increase in sorafenib AUC. Carriers of *UGT1A1* (TA)<sub>7/7</sub> had a similar relationship between bilirubin and AUC (i.e., a 30.9 ng/mL\*hr/mg unit increase in AUC corresponded to a 0.1 mg/dL increase in bilirubin). These data are consistent with previous case reports where *UGT1A1* (TA)<sub>6/7</sub> carriers developed jaundice following sorafenib treatment and are also consistent with our results that sorafenib is a mixed inhibitor of *UGT1A1*.

### Genotype versus progression-free survival and toxicity

The *CYP3A4\*1B* allele was weakly associated with PFS according to a Cox model analysis accounting for the multiple clinical trials where the present patient cohort was ascertained (data not shown). Those patients carrying variant alleles at *CYP3A4\*1B* ( $n=19$ ) tended to have shorter PFS than those patients carrying homozygous wild-type alleles ( $n=99$ ;  $P=0.034$ ). None of the other alleles studied herein were related to PFS ( $P>0.05$ ); however, the small numbers of variants within each trial led to wide confidence intervals for the individual hazard ratio estimates and the present results with respect to PFS should be interpreted with caution.

Consistent with previous literature (22), the incidence of HFSR was associated with increases in sorafenib AUC ( $P=0.0054$ ; Figure 3). However, all patients who had AUC>100 ng-h/mL/mg ( $n=7$ ) developed HFSR regardless of genotype ( $P=0.0085$ ); thus, while *UGT1A* genotypes were not associated with HFSR in the present study (due to heterogeneity of phenotype and low allele frequency), it is likely that *UGT1A1\*28\*28* and *UGT1A9\*3* carriers are subject to increased incidence of HFSR as they likely have higher exposure to sorafenib, and this should be confirmed by future studies.

## DISCUSSION

The case report presented here involved a child with Gilbert's syndrome (*UGT1A1\*28\*28*) who upon receiving sorafenib for treatment of a neurofibroma had higher than expected sorafenib exposure. Based on this and previously published observations (14–17), we hypothesized and assessed if *UGT1A1* genotype was associated with sorafenib systemic



exposure and serum bilirubin concentrations in patients treated with sorafenib. We confirmed previous hypotheses and findings (14–17) that sorafenib is an inhibitor of, but is not metabolized by, UGT1A1; rather, UGT1A9 is involved in sorafenib glucuronidation, which also confirms another report (8). It is unclear as to the exact reason for the approximately 4–10-fold difference in  $K_i$  values, however there is one plausible reason. Both this study and literature described sorafenib as a mixed-type inhibitor of UGT1A1, and although sorafenib demonstrates more noncompetitive characteristics (based on higher model correlation), there is a competitive inhibitor portion to the  $K_i$  calculation that is potentially altered based on the substrate used. The UGT1A1 substrate used in the literature was SN38, whereas bilirubin was used in this study, and the affinity of that substrate may affect the  $K_i$  of sorafenib. SN38 has a  $K_m$  of 11  $\mu\text{M}$  for UGT1A1 (29), whereas bilirubin has a reported  $K_m$  range of 0.2–26  $\mu\text{M}$  (25–28) due to the numerous stereoisomers present. Furthermore, the  $K_i$  of 11  $\mu\text{M}$  and  $\text{IC}_{50}$  of 18  $\mu\text{M}$  are clinically-relevant plasma concentrations following 400 mg BID dosing, which typically result in maximum plasma concentrations between 11–21  $\mu\text{M}$  after a cycle lasting either 7 or 28 days (30). Although sorafenib is >99% plasma protein bound, there is a high enough intra-hepatic sorafenib concentration in patients to inhibit the UGT1A1-mediated bilirubin glucuronidation. From our data, patients carrying at least one allele of *UGT1A1\*28* and having sorafenib AUC >60 ng\*h/mL, or 130 nM\*h, were related to increases in serum bilirubin following sorafenib treatment (Figure 2E–F). This provides clinical evidence that intra-hepatic sorafenib concentrations can be achieved in high enough levels to inhibit UGT1A1.

The present results suggest that patients carrying only *UGT1A1\*28*, and possibly *UGT1A9\*3*, alleles are at an increased risk of elevated sorafenib concentrations, as well as a greater incidence of HFSR. However, there remained a group of patients carrying *UGT1A1\*28/\*28* and *ABCC2-24C>T* that had abnormally low sorafenib AUC, thereby complicating the present analysis. Nonetheless, sorafenib AUC was well correlated with change in bilirubin concentrations in patients carrying at least a single *UGT1A1\*28* allele; conversely, it was not correlated with AUC in patients who carried only wild-type genotypes at this site.

The mechanism underlying the observation that individuals carrying *UGT1A1\*28/\*28* also have high AUC, a phenotype that appears to be modified by *ABCC2-24C>T*, is currently unclear. We tested the hypothesis that *UGT1A1\*28* alleles were merely in linkage disequilibrium with *UGT1A9* alleles (i.e., *UGT1A9\*3*, *UGT1A9IVS+1* (399C>T) and –118dT<sub>9/10</sub>) that were truly responsible for the observed differences in AUC. While linkage was observed, the current *UGT1A9* alleles did not explain the association between *UGT1A1\*28/\*28* and AUC. We also tested whether or not *ABCC2-24C>T* was itself related to AUC regardless of *UGT1A1* genotype; this also revealed no apparent association outside of those patients carrying *UGT1A1\*28/\*28*. Only parent sorafenib pharmacokinetics were analyzed, however no bias was expected through analysis of metabolites, since it has previously been demonstrated that altering one metabolic pathway of sorafenib does not alter parent drug pharmacokinetics (6).

The present results demonstrate that UGT1A1 binds sorafenib; others demonstrated that sorafenib binds to *ABCC2* and inhibits transport of other substrates but is not transported by

ABCC2 (31). Elimination of sorafenib glucuronides through ABCC2 may also be involved, however it is unclear if sorafenib glucuronides are actually transported by ABCC2. One previous report has also shown that the liver contains binding sites for drugs that act as “sinks” to slowly dissociate bound drug that is subsequently metabolized by liver enzymes (32). It is therefore possible that given the extensive enterohepatic circulation of sorafenib (10, 11), individuals expressing relatively high levels of liver UGT1A1 (i.e., those not carrying *UGT1A1\*28*) and possibly other proteins that bind sorafenib have a greater propensity to extrude sorafenib from the serum and hold sorafenib in the liver where it can be metabolized more extensively prior to hepatobiliary elimination. This may explain the significantly higher sorafenib AUCs of individuals who have reduced expression of UGT1A1 due to genetic polymorphisms. Still, our hypothesis does not explain why individuals carrying both *UGT1A1\*28* and *ABCC2-24C>T* variants have some of the lowest AUCs. As both proteins are involved in bilirubin elimination (19), we expected that patients carrying both *UGT1A1\*28* and *ABCC2-24C>T* would have less ability to glucuronidate and eliminate bilirubin due to decreased sorafenib-mediated UGT1A1 inhibition; however, this was not the case in the present patient cohort (data not shown). Therefore, the fact that both proteins regulate bilirubin concentrations that could in turn influence binding of sorafenib to UGT1A1 in the liver is not likely to be the cause of *ABCC2-24C>T* modifying the sorafenib exposure phenotype of patients carrying *UGT1A1\*28/\*28*.

Although our data did not point to a specific mechanism underlying associations between *UGT1A1\*28/\*28* and AUC, we observed a clear relationship between sorafenib exposure and bilirubin concentration. We demonstrated that sorafenib inhibits bilirubin glucuronidation as a mixed inhibitor *in vitro*. Based on this observation and the aforementioned case reports, we assessed whether or not sorafenib inhibited bilirubin glucuronidation in patients and whether or not this depended on *UGT1A1 A(TA)<sub>n</sub>TAA* status. The results indicated that sorafenib exposure was not correlated with total bilirubin concentration in patients that do not carry *UGT1A1\*28*; however, there was an increasingly strong correlation between sorafenib AUC and total bilirubin in patients carrying a single copy or two copies of *UGT1A1\*28*. Therefore, hyperbilirubinemia appears to be a marker of high sorafenib exposure in patients expressing low levels of UGT1A1 and care should be taken in monitoring patients carrying *UGT1A1\*28* that are known to have high sorafenib exposure. Nonetheless, none of the genes studied in this investigation were related to the PFS of the various sorafenib studies with the exception of the weak association with the *CYP3A4\*1B* allele. The results of the present Cox model should be interpreted with caution.

To our knowledge, the present pilot study represents the first exploration of sorafenib AUC in patients based on *UGT1A1* and *UGT1A9* genotype status and suggests that future studies should focus on the *UGT1A1 A(TA)<sub>n</sub>TAA* and *UGT1A9\*3* alleles and bilirubin increases in relation to sorafenib exposure, and HFSR.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgements

The authors thank Kathleen M. Wyvill and Thomas S. Uldrick for their contributions to the BAY-KS trial. The authors also thank all participating patients in each of the trials.

## Abbreviations:

<b>UGT</b>	UDP-glucuronyltransferase
<b>AUC</b>	area under the plasma-concentration time curve
<b>VEGF</b>	vascular endothelial growth factor
<b>CYP</b>	cytochrome P450
<b>MTD</b>	maximum tolerated dose
<b>CrCL</b>	creatinine clearance
<b>SGOT</b>	serum glutamic oxaloacetic transaminase
<b>CTEP</b>	Cancer Therapy Evaluation Program
<b>NADPH</b>	nicotinamide adenine dinucleotide phosphate
<b>DMSO</b>	dimethyl sulfoxide
<b>LC/MS</b>	liquid chromatography/mass spectrometry

## REFERENCES

- Jain L, Sissung TM, Danesi R, Kohn EC, Dahut WL, Kummar S, et al. Hypertension and hand-foot skin reactions related to VEGFR2 genotype and improved clinical outcome following bevacizumab and sorafenib. *J Exp Clin Cancer Res* 2010; 29:95. [PubMed: 20630084]
- Aragon-Ching JB, Jain L, Gulley JL, Arlen PM, Wright JJ, Steinberg SM, et al. Final analysis of a phase II trial using sorafenib for metastatic castration-resistant prostate cancer. *BJU Int* 2009; 103:1636–40. [PubMed: 19154507]
- Azad NS, Posadas EM, Kwitkowski VE, Steinberg SM, Jain L, Annunziata CM, et al. Combination targeted therapy with sorafenib and bevacizumab results in enhanced toxicity and antitumor activity. *J Clin Oncol* 2008; 26:3709–14. [PubMed: 18669456]
- Dahut WL, Scripture C, Posadas E, Jain L, Gulley JL, Arlen PM, et al. A phase II clinical trial of sorafenib in androgen-independent prostate cancer. *Clin Cancer Res* 2008; 14:209–14. [PubMed: 18172272]
- Jain L, Venitz J, Figg WD. Randomized discontinuation trial of sorafenib (BAY 43–9006). *Cancer Biol Ther* 2006; 5:1270–2. [PubMed: 16969070]
- Lathia C, Lettieri J, Cihon F, Gallentine M, Radtke M, Sundaresan P. Lack of effect of ketoconazole-mediated CYP3A inhibition on sorafenib clinical pharmacokinetics. *Cancer Chemother Pharmacol* 2006; 57:685–92. [PubMed: 16133532]
- Keating GM, Santoro A. Sorafenib: a review of its use in advanced hepatocellular carcinoma. *Drugs* 2009; 69:223–40. [PubMed: 19228077]
- Nexavar (Sorafenib) Prescribing Information, Bayer Pharmaceuticals. 2009 [cited; Available from: [http://www.nexavar.com/html/download/Nexavar\\_PL.pdf](http://www.nexavar.com/html/download/Nexavar_PL.pdf)]
- Girard H, Villeneuve L, Court MH, Fortier LC, Caron P, Hao Q, et al. The novel UGT1A9 intronic I399 polymorphism appears as a predictor of 7-ethyl-10-hydroxycamptothecin glucuronidation levels in the liver. *Drug Metab Dispos* 2006; 34:1220–8. [PubMed: 16595709]


10. Jain L, Woo S, Gardner ER, Dahut WL, Kohn EC, Kummar S, et al. Population Pharmacokinetic Analysis of Sorafenib in Patients with Solid Tumors. *Br J Clin Pharmacol* 2011.
11. European Public Assessment Reports (EPAR)-Scientific Discussion. 2009 [cited; Available from: [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/EPAR\\_-\\_Scientific\\_Discussion/human/000690/WC500027707.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Scientific_Discussion/human/000690/WC500027707.pdf)]
12. Parsa V, Heilbrun L, Smith D, Sethi A, Vaishampayan U. Safety and efficacy of sorafenib therapy in patients with metastatic kidney cancer with impaired renal function. *Clin Genitourin Cancer* 2009; 7:E10–5. [PubMed: 19692316]
13. Strassburg CP. Hyperbilirubinemia syndromes (Gilbert-Meulengracht, Crigler-Najjar, Dubin-Johnson, and Rotor syndrome). *Best Pract Res Clin Gastroenterol* 2010; 24:555–71. [PubMed: 20955959]
14. Meza-Junco J, Chu QS, Christensen O, Rajagopalan P, Das S, Stefanyshyn R, et al. UGT1A1 polymorphism and hyperbilirubinemia in a patient who received sorafenib. *Cancer Chemother Pharmacol* 2009; 65:1–4. [PubMed: 19672597]
15. Miller AA, Murry DJ, Owzar K, Hollis DR, Kennedy EB, Abou-Alfa G, et al. Phase I and pharmacokinetic study of sorafenib in patients with hepatic or renal dysfunction: CALGB 60301. *J Clin Oncol* 2009; 27:1800–5. [PubMed: 19255312]
16. Abou-Alfa GK, Amadori D, Santoro A, Figer A, De Greve J, Lathia C, et al. Safety and Efficacy of Sorafenib in Patients with Hepatocellular Carcinoma (HCC) and Child-Pugh A versus B Cirrhosis. *Gastrointest Cancer Res* 2011; 4:40–4. [PubMed: 21673874]
17. Mross K, Steinbild S, Baas F, Gmehling D, Radtke M, Voliotis D, et al. Results from an in vitro and a clinical/pharmacological phase I study with the combination irinotecan and sorafenib. *Eur J Cancer* 2007; 43:55–63. [PubMed: 17095207]
18. Dasanu CA, Alexandrescu DT, Dutcher J. Yellow skin discoloration associated with sorafenib use for treatment of metastatic renal cell carcinoma. *South Med J* 2007; 100:328–30. [PubMed: 17396743]
19. Jedlitschky G, Hoffmann U, Kroemer HK. Structure and function of the MRP2 (ABCC2) protein and its role in drug disposition. *Expert Opin Drug Metab Toxicol* 2006; 2:351–66. [PubMed: 16863439]
20. Kelly RJ, Rajan A, Force J, Keen C, Cao L, Yu Y, et al. Evaluation of KRAS mutations, angiogenic biomarkers and DCE-MRI in patients with advanced non-small cell lung cancer receiving sorafenib. *Clin Cancer Res* 2011.
21. M Gutierrez SK AD, Turkbey B, Choyke P, Wright JJ, Kurkjian C, Giaccone G, Doroshow JH, Murgo AJ. A phase II study of multikinase inhibitor sorafenib in patients with relapsed non-small cell lung cancer (NSCLC). *American Society of Clinical Oncology (ASCO)* 2008.
22. Azad NS, Aragon-Ching JB, Dahut WL, Gutierrez M, Figg WD, Jain L, et al. Hand-foot skin reaction increases with cumulative sorafenib dose and with combination anti-vascular endothelial growth factor therapy. *Clin Cancer Res* 2009; 15:1411–6. [PubMed: 19228742]
23. Kim A DE, Tepas K, Fox E, Balis FM, Korf B, and Widemann BC. Phase I trial of sorafenib in children with neurofibromatosis Type I and inoperable plexiform neurofibromas. *Neuro-Oncology*; 2010.
24. Widemann BC, Fox E, Adamson PC, Baruchel S, Kim A, Ingle AM, et al. Phase I study of sorafenib in children with refractory solid tumors: A Children's Oncology Group Phase I Consortium Trial. *Proc Am Soc Clin Oncology* 2009.
25. Senafi SB, Clarke DJ, Burchell B. Investigation of the substrate specificity of a cloned expressed human bilirubin UDP-glucuronosyltransferase: UDP-sugar specificity and involvement in steroid and xenobiotic glucuronidation. *Biochem J* 1994; 303 ( Pt 1):233–40. [PubMed: 7945246]
26. Seppen J, Bosma PJ, Goldhoorn BG, Bakker CT, Chowdhury JR, Chowdhury NR, et al. Discrimination between Crigler-Najjar type I and II by expression of mutant bilirubin uridine diphosphate-glucuronosyltransferase. *J Clin Invest* 1994; 94:2385–91. [PubMed: 7989595]
27. Zhang D, Chando TJ, Everett DW, Patten CJ, Dehal SS, Humphreys WG. In vitro inhibition of UDP glucuronosyltransferases by atazanavir and other HIV protease inhibitors and the relationship of this property to in vivo bilirubin glucuronidation. *Drug Metab Dispos* 2005; 33:1729–39. [PubMed: 16118329]

28. Udomuksorn W, Elliot DJ, Lewis BC, Mackenzie PI, Yoovathaworn K, Miners JO. Influence of mutations associated with Gilbert and Crigler-Najjar type II syndromes on the glucuronidation kinetics of bilirubin and other UDP-glucuronosyltransferase 1A substrates. *Pharmacogenet Genomics* 2007; 17:1017–29. [PubMed: 18004206]
29. Jinno H, Tanaka-Kagawa T, Hanioka N, Saeki M, Ishida S, Nishimura T, et al. Glucuronidation of 7-ethyl-10-hydroxycamptothecin (SN-38), an active metabolite of irinotecan (CPT-11), by human UGT1A1 variants, G71R, P229Q, and Y486D. *Drug Metab Dispos* 2003; 31:108–13. [PubMed: 12485959]
30. Strumberg D, Clark JW, Awada A, Moore MJ, Richly H, Hendlitz A, et al. Safety, pharmacokinetics, and preliminary antitumor activity of sorafenib: a review of four phase I trials in patients with advanced refractory solid tumors. *Oncologist* 2007; 12:426–37. [PubMed: 17470685]
31. Hu S, Chen Z, Franke R, Orwick S, Zhao M, Rudek MA, et al. Interaction of the multikinase inhibitors sorafenib and sunitinib with solute carriers and ATP-binding cassette transporters. *Clin Cancer Res* 2009; 15:6062–9. [PubMed: 19773380]
32. Rubin GM, Tozer TN. Hepatic binding and Michaelis-Menten metabolism of drugs. *J Pharm Sci* 1986; 75:660–3. [PubMed: 3761165]

### Translational Relevance


We investigated UGT1A-mediated sorafenib glucuronidation *in vitro*, inhibition of UGT1A1 bilirubin conjugation by sorafenib, and ascertained whether patients carrying *UGT1A1*\*28 treated with sorafenib had increased sorafenib exposure or increased risk of developing hyperbilirubinemia. We also investigated genetic variation in genes encoding sorafenib metabolizing enzymes (*UGT1A9*, *CYP3A4/5*) and the glucuronide transporter, *ABCC2*. *In vitro* data demonstrate that sorafenib inhibits UGT1A1. Patients carrying *UGT1A1*\*28/\*28 had abnormally low exposure to sorafenib if they also carried the -24C>T variant in *ABCC2*, but had abnormally high exposure to sorafenib if they were wild-type for *ABCC2*. Patients carrying *UGT1A1*\*28/\*28 also had greater increases in total bilirubin if sorafenib exposure was high. The clinical data suggest that sorafenib can cause hyperbilirubinemia in patients with Gilbert's syndrome, which can lead to abnormally high or low sorafenib exposure. Therefore, patients with Gilbert's syndrome must be carefully monitored when treated with sorafenib or alternative treatment options should be considered.






**Figure 1.**

Dose-normalized sorafenib AUC versus *UGT1A* genotype. *UGT1A9*\*1/\*3 carriers each also carried *UGT1A1* (*TA*)<sub>6</sub>/*(TA)*<sub>7</sub> (AUC = 17.3 and 20.2 ng/mL\*hr/mg) while the single *UGT1A9*\*3/\*3 carrier also carried *UGT1A1* (*TA*)<sub>6</sub>/*(TA)*<sub>6</sub> (AUC = 153.3 ng/mL\*hr/mg, as indicated by \*). *UGT1A-impaired* implies patients with deficient *UGT1A1* or *UGT1A9* metabolism based on genetics. Excluded patients ( $n=38$ ) are described in the supplementary results section and  $n=82$  individuals were included in the present analysis. Of these, AUC data for  $n=8$  patients participating on BAY-KS were not available; thus potential drug-drug interactions between ritonavir and sorafenib were not accounted for. There was no association between *UGT1A1* and *UGT1A9* genotype status when compared to sorafenib AUC ( $P=0.20$ ; Kruskal Wallis ANOVA).



**Figure 2.**

*UGT1A1* A(TA)<sub>n</sub>TAA status versus bilirubin change from baseline following sorafenib (mg/dL) (A); and sorafenib AUC (ng/mL\*hr/mg) versus bilirubin change from baseline (mg/dL) (B) in patients with *UGT1A1*(TA)<sub>5/5</sub> or *UGT1A1*(TA)<sub>5/6</sub> (C), *UGT1A1*(TA)<sub>6/6</sub> (D), *UGT1A1*(TA)<sub>6/7</sub> (E), or *UGT1A1*(TA)<sub>7/7</sub> (F). \* Indicates a patient who carried *UGT1A1*(TA)<sub>6/6</sub> that also carried two variants at *UGT1A9*\*3 and was thus excluded from analysis in the *UGT1A1*(TA)<sub>6/6</sub> cohort.



**Figure 3.** Sorafenib exposure is related to hand-foot skin reaction. HFSR was associated with increases in sorafenib AUC ( $P=0.0054$ ).

**Table 1.**

## Patient demographics and baseline characteristics

Characteristic	Values <i>n</i> (%) or median (95%CI)
Total	120 (100.0)
Male	86(71.7)
Female	34 (28.3)
Age	63.2(60.9–65.4)
<i>Race</i>	
Caucasian	99 (82.5)
African American	11 (9.2)
Hispanic	4 (3.3)
Asian	5 (4.2)
Unknown	1 (0.8)
BSA (m <sup>2</sup> )	2.0(1.9–2.0)
<i>Study/Disease</i>	
BAY-BEV / Solid tumors	27 (22.5)
BAY-KS / Kaposi's sarcoma	8 (6.7)
BAY-CRPC / Prostate	46 (38.3)
BAY-NSCLC / Lung	22(18.3)
BAY-CRC / Colorectal	17(14.2)
Albumin (g/dL)	3.6(3.5–3.7)
Alkaline Phosphatase (U/L)	81.0(74.0–89.0)
Total Bilirubin (mg/dL)	0.6 (0.6–0.7)
SGOT (U/L)	26 (24–28)
CrCl (mL/min)	90.9 (84.8–97.7)
CrCl <60mL/min	17(14.2)