

Original Article
Preventive &
Social Medicine



Projection of the Years of Life Lost, Years Lived with Disability, and Disability-Adjusted Life Years in Korea for 2030

Bomi Park , Bohyun Park , Hyejin Han , Eun Jung Choi , Nam-eun Kim , Yoonhee Shin , and Hyesook Park

Department of Preventive Medicine, Ewha Womans University School of Medicine, Seoul, Korea

OPEN ACCESS

Received: Jul 8, 2018

Accepted: Oct 10, 2018

Address for Correspondence:

Hyesook Park, MD, PhD

Department of Preventive Medicine, Ewha Womans University School of Medicine, 1071 Anyangcheon-ro, Yangcheon-gu, Seoul 07985, Korea.
E-mail: hpark@ewha.ac.kr

© 2019 The Korean Academy of Medical Sciences.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Bomi Park <https://orcid.org/0000-0001-5834-9975>
Bohyun Park <https://orcid.org/0000-0002-8538-9662>
Hyejin Han <https://orcid.org/0000-0002-1669-3177>
Eun Jung Choi <https://orcid.org/0000-0002-7698-3424>
Nam-eun Kim <https://orcid.org/0000-0002-8826-7644>
Yoonhee Shin <https://orcid.org/0000-0002-9950-9362>
Hyesook Park <https://orcid.org/0000-0002-9359-6522>

ABSTRACT

Background: Projection of future trends in disease burden can facilitate setting of priorities for health policies and resource allocation. We report here projections of disease-specific mortality and the burdens of various diseases in Korea from 2016 to 2030.

Methods: Separate age- and sex-specific projection models for 21 major cause clusters from 2016 to 2030 were developed by applying coherent functional data models based on historical trends from 2002 to 2015. The age- and sex-specific years of life lost (YLL) for each cause cluster were projected based on the projected number of deaths. Years lived with disability (YLD) projections were derived using the 2015 age- and sex-specific YLD to YLL ratio. The disability-adjusted life years (DALYs) was the sum of YLL and YLD.

Results: The total number of deaths is projected to increase from 275,777 in 2015 to 421,700 in 2030, while the age-standardized death rate is projected to decrease from 586.9 in 2015 to 447.3 in 2030. The largest number of deaths is projected to be a result of neoplasms (75,758 deaths for males; 44,660 deaths for females), followed by cardiovascular and circulatory diseases (34,795 deaths for males; 48,553 deaths for females). The three leading causes of DALYs for both sexes are projected to be chronic respiratory diseases, musculoskeletal disorders, and other non-communicable diseases (NCDs).

Conclusion: We demonstrate that NCDs will continue to account for the majority of the disease burden in Korea in the future.

Keywords: Burden of Disease; Republic of Korea; Projection; Coherent Functional Data Model; Time Series

INTRODUCTION

Assessment and quantification of the population health scenarios expected to occur in the future provide information on probable trends in health and enable setting of priorities for healthcare and public health policies. This is particularly important for Korea, which is currently undergoing a major epidemiologic and demographic transition; the Korean population has aged more rapidly than those of other developed countries, and Korea is expected to be a super-aged society by 2026.¹ Comprehensive assessments of current burden of disease in Korea have been conducted.^{2,3} However, little effort has focused on projections

Funding

This study was supported by a grant from the Korean Health Technology R & D Project, Ministry of Health & Welfare, Republic of Korea (No. HI13C0729).

Disclosure

The authors have no potential conflicts of interest to disclose.

Author Contributions

Conceptualization: Park B,¹ Park H. Data curation: Han H, Choi EJ, Kim N, Shin Y. Formal analysis: Park B,¹ Park B,² Park H. Investigation: Park B.² Methodology: Park B.¹ Writing - original draft: Park B,¹ Park B,² Park H. Writing - review & editing: Park B,¹ Park B,² Han H, Choi EJ, Kim N, Shin Y, Park H.

Park B,¹ Park Bomi; Park B,² Park Bohyun.

of disease-specific mortality and the burden of disease for the entire disease spectrum at the national level.

At the global level, projections for a complete set of causes were conducted as part of the Global Burden of Disease (GBD) study.^{4,5} The GBD projections used a structural model based on relationships between mortality and projected changes in key system components. However, the structural model requires considerable knowledge of the important system components and the relationships among them. In addition, when the system is complex and highly sensitive to its components, the structural model provides less reliable estimates. In contrast, aggregate models are based on the assumption that previous trends will continue into the future. This method requires good mortality and morbidity registration data.

We prepared projections of mortality and burden of disease from 2016 to 2030 by developing separate age-, sex-, and cause-specific projection models using coherent functional data models and time-series methods based on the trends from 2002 to 2015.

METHODS**Data source**

To estimate the mortality rates from 2002 to 2015 and to project future mortality rates from 2016 to 2030 of diseases and injuries in the entire Korean population, the 2002–2015 Causes of Death Statistics and database of residents registered with the Ministry of Interior, the 2016–2030 projected population, and the 2002–2030 Life Expectancy Table from Statistics Korea were used.⁶ To estimate the prevalence and incidence rates of the selected causes in 2015, we used National Health Insurance Service claims data for 2015, the 2015 National Cancer Registration and Statistics from the Korea Central Cancer Registry, and the 2015 Korea National Hospital Discharge Injury Surveillance data from the Korea Centers for Disease Control and Prevention.

Case definitions

In total, 260 mutually exclusive diseases and injuries were classified into the following 21 cause clusters: 1) human immunodeficiency virus/acquired immune deficiency syndrome (HIV/AIDS) and tuberculosis; 2) diarrhea, lower respiratory infections, meningitis, and other common infectious diseases; 3) neglected tropical diseases and malaria; 4) maternal disorders; 5) neonatal disorders; 6) nutritional deficiencies; 7) other communicable, maternal, neonatal, and nutritional disorders; 8) neoplasms; 9) cardiovascular and circulatory diseases; 10) chronic respiratory diseases; 11) cirrhosis of the liver; 12) digestive diseases except cirrhosis; 13) neurological disorders; 14) mental and behavioral disorders; 15) diabetes, urogenital, blood, and endocrine diseases; 16) musculoskeletal disorders; 17) other non-communicable diseases (NCDs); 18) transport injuries; 19) unintentional injuries other than transport injuries; 20) self-harm and interpersonal violence; and 21) forces of nature, war, and legal intervention. These 21 cause clusters were further grouped into three broad cause groups: 1) communicable, maternal, neonatal, and nutritional conditions (CMNNs); 2) NCDs; and 3) injuries. Detailed information on the definition of cause of death and epidemiologic parameters is available.³

Mortality projection methods

We used these 21 cause clusters to estimate mortality rates from 2002 to 2015 for both sexes and eighteen 5-year age groups (0–4, 5–9, 10–14, 15–19, 20–24, 25–29, 30–34, 35–39,

40–44, 45–49, 50–54, 55–59, 60–64, 65–69, 70–74, 75–79, 80–84, and ≥ 85 years). For analysis of maternal, neonatal, and childhood behavioral disorders, age restriction was applied depending on the characteristics of the diseases. Only females 10–59 years of age were included for maternal disorders, children 0–9 years of age for infant disorders, newborns 0–1 years of age for sepsis and other infectious disorders of newborns, and children under 20 years of age for childhood behavioral disorders.

To forecast mortality rates from 2016 to 2030, we developed coherent functional time-series models for each age-sex group for the 21 cause-of-death clusters using the demography package in R software (version 3.4.4; R Foundation, Vienna, Austria). This model uses smoothed historical mortality data, and we used the mortality rates for the years 2002–2015 as the basis for the projection of mortality rates. The past mortality rates were calculated by the ratio between the number of deaths and mid-year population. The square roots of the products and ratios of the smoothed rates for each sex were modeled using functional time series models.⁷ The projected number of deaths was generated by applying the projected population to the projected mortality rate.

Calculation of years of life lost (YLLs), years lived with disability (YLDs), and disability-adjusted life years (DALYs)

Details of the methods for estimating YLLs, YLDs, and DALYs are provided elsewhere.^{8,9} The cause-specific YLLs for 2015 were computed by multiplying the number of deaths for each cause cluster by the residual expected life expectancy at the age of death. We used a 4% age weighting and 3% time discount rate. The YLDs for 2015 were calculated by multiplying the number of incident cases by the average duration of a case and the disability weight of the disease. To calculate the age at onset and disease duration, we used DisMod-II software (EpiGear, Noosa, Australia) with appropriate input parameters, including the incidence rate, prevalence rate, mortality rate, and fatality rate. Detailed information on the disability weights used in this study is available.¹⁰ The DALY was the sum of YLL and YLD.

YLLs from 2016 to 2030 were calculated based on the projected mortality by cause, age, and sex. YLD projections were derived from the YLL projections by applying the ratio of YLDs to YLLs for 2015 with the assumption that these ratios would be constant from 2015 to 2030, as estimated in the original GBD projections.⁴ For causes for which there was no mortality, the 2015 YLD rates per capita were assumed to remain constant until 2030. We used the projected population to calculate age-standardized YLL, YLD, and DALY rates per capita. All the analysis was done by age-, and sex-specific models.

Ethics statement

This study was approved by the Korea University Institutional Review Board (KU-IRB-18-EX-51-A-1). The informed consent was waived by the board.

RESULTS

Overall, the number of deaths is projected to increase from 275,777 in 2015 to 421,700 in 2030; and the age-standardized death rate (ASDR) is projected to decrease from 586.9 in 2015 to 447.3 in 2030. NCDs are projected to account for approximately 80% of all deaths in 2030. In contrast, injuries and CMNNs will account for 13.7% and 5.9%, respectively, of the total deaths in 2030.

Table 1 summarizes the projected number of deaths and ASDRs for the 21 cause clusters in 2030 by sex in comparison with those in 2015, as well as the percentage changes in ASDRs for 2015–2030. The five leading causes of death are projected to be NCDs. The largest number of deaths is projected to be caused by neoplasms (75,758 deaths for males; 44,660 deaths for females), followed by cardiovascular and circulatory diseases (34,795 deaths for males; 48,553 deaths for females); these were also the leading causes of death in 2015. Although the ASDRs for both are projected to decline from 2015 to 2030, the number of deaths from those causes is projected to increase, likely due to population growth and/or aging. For injuries, the rank orders of the causes drop or are unchanged from those in 2015. Unintentional injuries and self-harm and interpersonal violence are projected to account for almost 90% of the injury-related deaths in 2030. Generally, CMNNs had the lowest rank orders in both 2015 and 2030, with the exceptions of diarrhea, lower respiratory infections, meningitis, and other common infectious diseases (ninth for males and seventh for females in 2030) and HIV/AIDS (11th for both males and females in 2030), each of which is projected to move up one place in the rankings.

The estimates of YLL per 100,000 population for 2015 and 2030 and the changes in rank order by sex are shown in **Table 2**. Neoplasms were the leading cause of YLL for both sexes in 2015 and is projected to remain so in 2030. For males, self-harm and interpersonal violence is projected to move up one place in the rankings to be the second leading cause of YLL. Cardiovascular and circulatory diseases is projected to move down one place in the rankings and be the third leading cause of YLL. For females, the ranking of cardiovascular and

Table 1. Projected mortality for selected 21 causes in 2030 in comparison to mortality in 2015

Diseases	Male					Female				
	2015		2030		% change in ASDR	2015		2030		% change in ASDR
	Deaths	ASDR	Deaths	ASDR		Deaths	ASDR	Deaths	ASDR	
CMNNs										
HIV/AIDS and tuberculosis	2,181	4.5	4,669	4.8	7.0	1,395	2.7	2,628	2.7	0.4
Other infectious ds ^a	3,031	5.8	6,705	5.2	-9.6	3,436	6.4	8,046	7.1	9.8
Neglected tropical ds and malaria	131	0.3	266	0.3	1.7	129	0.3	227	0.3	-5.1
Maternal disorder	N/A					64	0.3	65	0.4	11.6
Neonatal disorder	418	2.6	563	3.8	49.5	330	1.9	378	2.5	27.2
Nutritional deficiencies	312	0.6	664	0.6	-3.6	374	0.7	857	0.7	5.4
Other CMNN	309	0.7	568	0.7	-6.1	289	0.7	491	0.7	-3.5
NCDs										
Neoplasms	51,049	102.9	75,758	71.0	-31.0	32,848	75.6	44,660	53.7	-28.9
Cardiovascular and circulatory ds	32,964	63.6	34,795	26.2	-58.9	37,848	72.2	48,553	40.5	-43.9
Chronic respiratory ds	9,114	15.2	23,313	16.1	6.0	6,855	12.5	14,777	12.9	3.5
Cirrhosis of the liver	5,882	14.2	8,494	14.7	3.9	2,332	5.7	3,568	5.6	-2.0
Digestive ds except cirrhosis	2,672	5.3	5,864	5.5	3.8	3,177	6.0	6,390	6.2	3.0
Neurological disorders	7,292	13.4	16,676	12.8	-4.3	11,000	20.4	24,441	21.6	6.1
Mental and behavioral disorders	924	2.3	1,228	2.4	2.6	197	0.5	247	0.5	-5.2
DM, urogenital, blood, and endocrine ds	9,278	18.0	14,344	10.2	-43.4	10,357	20.5	18,353	16.0	-21.8
Musculoskeletal disorders	510	1.0	1,032	0.9	-5.9	970	2.0	1,864	2.0	1.0
Other NCDs	691	2.9	1,067	2.9	0.5	805	3.0	1,222	3.0	1.8
Injuries										
Transport injuries	4,566	13.3	3,710	6.8	-48.9	2,154	6.0	1,809	3.2	-47.7
Unintentional injuries ^b	7,497	18.6	12,212	19.2	3.1	5,576	13.0	9,663	12.4	-4.5
Self-harm and interpersonal violence	11,393	33.8	14,760	33.1	-1.9	5,136	16.9	6,250	17.3	1.9
Other injuries ^c	180	0.4	348	0.6	54.8	111	0.2	174	0.2	-3.3

ASDR = age-standardized death rate, ds = diseases, N/A = not applicable, CMNN = communicable, maternal, neonatal, and nutritional condition, HIV/AIDS = human immunodeficiency virus/acquired immune deficiency syndrome, NCD = non-communicable disease, DM = diabetes mellitus.

^aDiarrhea, lower respiratory infections, meningitis, and other common infectious diseases; ^bUnintentional injuries other than transport injuries; ^cForces of natures, war, and legal intervention.

Table 2. Projected YLLs and changes in rankings for selected 21 causes for 2015 and 2030

Diseases	Male					Female				
	2015		2030		Change in rank	2015		2030		Change in rank
	YLL ^a	Rank	YLL ^a	Rank		YLL ^a	Rank	YLL ^a	Rank	
CMNNs										
HIV/AIDS and tuberculosis	72	12	125	12	-	31	14	56	13	↑1
Other infectious ds ^b	76	11	135	11	-	58	10	136	8	↑2
Neglected tropical ds and malaria	4	20	8	20	-	3	21	5	20	↑1
Maternal disorder	N/A					7	18	7	19	↓1
Neonatal disorder	54	14	70	13	↑1	43	13	48	14	↓1
Nutritional deficiencies	10	18	16	19	↓1	6	19	14	17	↑2
Other CMNNs	12	17	18	17	-	10	17	14	16	↑1
NCDs										
Neoplasms	1,642	1	1,940	1	-	1,108	1	1,146	1	-
Cardiovascular and circulatory ds	941	2	680	3	↓1	705	2	772	2	-
Chronic respiratory ds	164	9	434	5	↑4	104	8	252	7	↑1
Cirrhosis of the liver	296	5	390	6	↓1	95	9	125	9	-
Digestive ds except cirrhosis	80	10	147	10	-	56	12	122	10	↑2
Neurological disorders	172	8	332	7	↑1	179	6	415	3	↑3
Mental and behavioral disorders	49	15	61	14	↑1	11	16	12	18	↓2
DM, urogenital, blood, and endocrine ds	265	7	268	8	↓1	214	4	306	5	↓1
Musculoskeletal disorders	15	16	25	16	-	23	15	41	15	-
Other NCDs	58	13	59	15	↓2	56	11	60	12	↓1
Injuries										
Transport injuries	281	6	164	9	↓3	109	7	69	11	↓4
Unintentional injuries ^c	357	4	479	4	-	182	5	255	6	↓1
Self-harm and interpersonal violence	728	3	785	2	↑1	356	3	363	4	↓1
Other injuries ^d	8	19	17	18	↑1	3	20	5	21	↓1

YLL = years of life lost, CMNN = communicable, maternal, neonatal, and nutritional condition, HIV/AIDS = human immunodeficiency virus/acquired immune deficiency syndrome, ds = diseases, N/A = not applicable, NCD = non-communicable disease, DM = diabetes mellitus.

^aYLLs per 100,000 population; ^bDiarrhea, lower respiratory infections, meningitis, and other common infectious diseases; ^cUnintentional injuries other than transport injuries; ^dForces of natures, war, and legal intervention.

circulatory diseases is projected to remain unchanged as the second leading cause of YLL, and neurological disorders is projected to move up three places to be the third leading cause of YLL. In general, there is projected to be little or no change in the YLL rankings between 2015 and 2030, while chronic respiratory diseases is projected to move up four places for males, neurological disorders is projected to move up three places for females, and transport injuries is projected to move down three places for males and four places for females.

Table 3 summarizes the changes of YLDs and DALYs per 100,000 population for the 21 cause clusters from 2015 to 2030 by sex. For males, the leading cause of YLD in 2015 was digestive diseases except cirrhosis, followed by chronic respiratory diseases and other NCDs, while the leading cause of YLD in 2030 was chronic respiratory diseases followed by other NCDs and digestive diseases except cirrhosis. For females, the largest number of YLD per 100,000 in 2015 was caused by musculoskeletal disorders, followed by diabetes mellitus (DM), urogenital, blood, and endocrine diseases, and other NCDs. The top leading causes of YLD were unchanged in 2030. Other NCDs and digestive diseases except cirrhosis were projected to be second and third in the rankings, respectively, while DM, urogenital, blood, and endocrine diseases is projected to move down five places to be seventh in the rankings.

DALYs are projected to increase from 26,799,184 in 2015 to 30,685,593 in 2030, an increase of 14.5%. During the same period, the population is projected to increase by 3.9%. NCDs are projected to account for 90% of the total DALYs and to represent a leading burden of disease, while CMNNs and injuries are projected to account for 5% each. The three leading causes

Table 3. Projected YLDs and DALYs for selected 21 causes for 2015 and 2030

Diseases	Male				Female			
	2015		2030		2015		2030	
	YLD ^a	DALY ^b	YLD ^a	DALY ^b	YLD ^a	DALY ^b	YLD ^a	DALY ^b
CMNNs								
HIV/AIDS and tuberculosis	107	179	140	265	54	85	90	146
Other infectious ds ^c	60	136	74	209	77	135	80	216
Neglected tropical ds and malaria	0	5	0	8	0	3	0	6
Maternal disorder	N/A				3,001	3,008	4,417	4,424
Neonatal disorder	14	68	18	88	10	54	12	59
Nutritional deficiencies	19	29	32	48	96	102	165	179
Other CMNNs	2	14	2	20	19	28	15	29
NCDs								
Neoplasms	1,181	2,805	1,224	3,164	1,218	2,326	894	2,040
Cardiovascular and circulatory ds	3,249	4,190	1,676	2,356	2,754	3,459	1,629	2,401
Chronic respiratory ds	6,242	6,406	10,919	11,352	7,108	7,212	7,429	7,681
Cirrhosis of the liver	975	1,271	1,105	1,495	501	596	548	673
Digestive ds except cirrhosis	6,924	7,004	7,671	7,818	7,116	7,172	8,964	9,086
Neurological disorders	1,381	1,553	2,180	2,512	2,223	2,401	3,965	4,380
Mental and behavioral disorders	2,081	2,131	2,459	2,520	2,572	2,583	3,100	3,112
DM, urogenital, blood, and endocrine ds	5,493	5,758	2,846	3,114	7,971	8,185	3,910	4,216
Musculoskeletal disorders	6,095	6,109	7,245	7,270	10,223	10,246	11,675	11,716
Other NCDs	6,120	6,178	7,682	7,741	7,734	7,790	9,728	9,788
Injuries								
Transport injuries	923	1,205	471	635	633	742	303	372
Unintentional injuries ^d	1,310	1,667	1,566	2,045	895	1,078	1,161	1,416
Self-harm and interpersonal violence	111	840	104	888	65	421	63	426
Other injuries ^e	3	11	5	21	1	4	1	6

YLD = years lived with disability, DALY = disability-adjusted life year, CMNN = communicable, maternal, neonatal, and nutritional condition, HIV/AIDS = human immunodeficiency virus/acquired immune deficiency syndrome, ds = diseases, N/A = not applicable, NCD = non-communicable disease, DM = diabetes mellitus. ^aYLDs per 100,000 population; ^bDALYs per 100,000 population; ^cDiarrhea, lower respiratory infections, meningitis, and other common infectious diseases; ^dUnintentional injuries other than transport injuries; ^eForces of natures, war, and legal intervention.

of DALYs for both sexes are projected to be chronic respiratory diseases, musculoskeletal disorders, and other NCDs.

Figs. 1 and 2 show the rank orders of DALYs in 2015 and 2030 by sex. The three leading causes of DALY are projected to be chronic respiratory diseases, digestive diseases except cirrhosis, and other NCDs for males; and musculoskeletal disorders, other NCDs, and digestive diseases except cirrhosis for females. Overall, NCDs are ranked highly and CMNNs are ranked lowly. Generally, there was little or no change in the DALY rankings between 2015 and 2030, while cardiovascular and circulatory diseases is projected to move down three places for males, neurological disorders is projected to move up three places for females, and both DM, urogenital, blood, and endocrine disorders and neonatal disorders are projected to move down five places for females.

Table 4 lists the five leading causes of DALYs in 2030 by age group. Chronic respiratory disease is projected to be an important cause of DALYs in all age groups. Mental and behavioral disorders as well as unintentional injuries other than transport injuries are projected to be the major causes of DALYs for males in the 10–19 years age group, and maternal disorders is projected to be the major cause of DALYs for females of childbearing age. Musculoskeletal disorders are projected to be ranked highly in the 20–64 years age group, and neoplasms for males and neurological disorders for females are the leading causes of DALY in the > 65 years age group.

Rank	2015	2030	Rank
1	Digestive diseases except cirrhosis	Chronic respiratory diseases	1
2	Chronic respiratory diseases	Digestive diseases except cirrhosis	2
3	Other non-communicable diseases	Other non-communicable diseases	3
4	Musculoskeletal disorders	Musculoskeletal disorders	4
5	Diabetes, urogenital, blood, and endocrine diseases	Neoplasms	5
6	Cardiovascular and circulatory diseases	Diabetes, urogenital, blood, and endocrine diseases	6
7	Neoplasms	Mental and behavioral disorders	7
8	Mental and behavioral disorders	Neurological disorders	8
9	Unintentional injuries other than transport injuries	Cardiovascular and circulatory diseases	9
10	Neurological disorders	Unintentional injuries other than transport injuries	10
11	Cirrhosis of the liver	Cirrhosis of the liver	11
12	Transport injuries	Self-harm and interpersonal violence	12
13	Self-harm and interpersonal violence	Transport injuries	13
14	HIV/AIDS and Tuberculosis	HIV/AIDS and Tuberculosis	14
15	Diarrhea, lower respiratory infections, meningitis, and other common infectious diseases	Diarrhea, lower respiratory infections, meningitis, and other common infectious diseases	15
16	Neonatal disorder	Neonatal disorder	16
17	Nutritional deficiencies	Nutritional deficiencies	17
18	Other communicable, maternal, neonatal, and nutritional disorders	Forces of natures, war, and legal intervention	18
19	Forces of natures, war, and legal intervention	Other communicable, maternal, neonatal, and nutritional disorders	19
20	Neglected tropical diseases and malaria	Neglected tropical diseases and malaria	20

Fig. 1. Change in rank order of DALYs between 2015 and 2030 for the 21 selected causes in male. DALY = disability-adjusted life year, HIV/AIDS = human immunodeficiency virus/acquired immune deficiency syndrome.

DISCUSSION

We performed projections of the number of deaths and burden of disease from 2016 to 2030 by 21 cause clusters, age, and sex. The number of deaths is projected to increase, and the ASDRs are projected to decrease, from 2015 and 2030. By 2030, the largest number of deaths is projected to be caused by neoplasms and cardiovascular and circulatory diseases. There is no substantial change in the patterns of DALYs between 2015 and 2030. In 2030, the three leading causes of DALY will be chronic respiratory diseases, musculoskeletal disorders, and other NCDs. The proportions of deaths and DALYs due to NCDs were projected to be 82% and 90%, respectively.

The projected trends of deaths and DALYs can be attributed in part to population growth and demographic transition in addition to improvements in medical care and health policy. According to the population projections reported by Statistics Korea, the proportion of the population aged 0–14 years is projected to decrease from 13.8% in 2015 to 11.5% in 2030, the proportion of the population of working age is projected to decrease from 73.4% in 2015 to

Rank	2015	2030	Rank
1	Musculoskeletal disorders	Musculoskeletal disorders	1
2	Diabetes, urogenital, blood, and endocrine diseases	Other non-communicable diseases	2
3	Other non-communicable diseases	Digestive diseases except cirrhosis	3
4	Chronic respiratory diseases	Chronic respiratory diseases	4
5	Digestive diseases except cirrhosis	Maternal disorder	5
6	Maternal disorder	Neurological disorders	6
7	Cardiovascular and circulatory diseases	Diabetes, urogenital, blood, and endocrine diseases	7
8	Mental and behavioral disorders	Mental and behavioral disorders	8
9	Neurological disorders	Cardiovascular and circulatory diseases	9
10	Neoplasms	Neoplasms	10
11	Unintentional injuries other than transport injuries	Unintentional injuries other than transport injuries	11
12	Transport injuries	Cirrhosis of the liver	12
13	Neonatal disorder	Self-harm and interpersonal violence	13
14	Cirrhosis of the liver	Transport injuries	14
15	Self-harm and interpersonal violence	Diarrhea, lower respiratory infections, meningitis, and other common infectious diseases	15
16	Diarrhea, lower respiratory infections, meningitis, and other common infectious diseases	Nutritional deficiencies	16
17	Nutritional deficiencies	HIV/AIDS and Tuberculosis	17
18	HIV/AIDS and Tuberculosis	Neonatal disorder	18
19	Other communicable, maternal, neonatal, and nutritional disorders	Other communicable, maternal, neonatal, and nutritional disorders	19
20	Forces of natures, war, and legal intervention	Forces of natures, war, and legal intervention	20
21	Neglected tropical diseases and malaria	Neglected tropical diseases and malaria	21

Fig. 2. Change in rank order of DALYs between 2015 and 2030 for the 21 selected causes in female. DALY = disability-adjusted life year, HIV/AIDS = human immunodeficiency virus/acquired immune deficiency syndrome.

64.0% in 2030, and the proportion of the population ≥ 65 years is projected to increase from 12.8% in 2015 to 24.5% in 2030.¹

According to the GBD projection, the three leading causes of burden of disease worldwide in 2030 will be HIV/AIDS, depression, and ischemic heart disease, and those in high-income countries will be depression, ischemic heart disease, and dementias. Comparing the GBD projection with our results is difficult because the GBD study performed projections for 132 detailed causes, whereas we conducted projections for 21 cause-clusters. However, the future disease burden in Korea is in accordance with that in high-income countries in the GBD projection, in that NCDs will account for the majority.

To the best of our knowledge, this is the first study to predict cause-specific mortality and DALYs for a comprehensive set of causes in Korea; therefore, comparisons with previous studies are impossible. Our projections for all-cause mortality are similar to those by Statistics Korea, despite our use of the sum of the projections for 21 cause clusters compared

Table 4. Ranking for 5 leading disease groups of DALYs^a by sex and age group in 2030

Age	Rank	Male (%)	Female (%)
Total	1	Chronic respiratory diseases (21)	Musculoskeletal disorders (19)
	2	Digestive diseases except cirrhosis (15)	Other NCDs (16)
	3	Other NCDs (14)	Digestive diseases except cirrhosis (15)
	4	Musculoskeletal disorders (14)	Chronic respiratory diseases (12)
	5	Neoplasms (6)	Maternal disorders (7)
0–9 yr	1	Chronic respiratory disease (78)	Chronic respiratory diseases (47)
	2	Other NCDs (10)	Other NCDs (25)
	3	Digestive diseases except cirrhosis (5)	Digestive diseases except cirrhosis (17)
	4	Neonatal disorder (1)	Mental and behavioral disorders (4)
	5	Mental and behavioral disorders (1)	Diabetes, urogenital, blood, and endocrine diseases (2)
10–19 yr	1	Other NCDs (23)	Other NCDs (51)
	2	Chronic respiratory diseases (17)	Digestive diseases except cirrhosis (11)
	3	Mental and behavioral disorders (17)	Chronic respiratory diseases (11)
	4	Unintentional injuries other than transport injuries (10)	Maternal disorder (7)
	5	Musculoskeletal disorders (9)	Mental and behavioral disorders (6)
20–64 yr	1	Digestive diseases except cirrhosis (18)	Musculoskeletal disorders (24)
	2	Musculoskeletal disorders (17)	Digestive diseases except cirrhosis (16)
	3	Other NCDs (16)	Other NCDs (13)
	4	Chronic respiratory diseases (14)	Maternal disorder (13)
	5	Diabetes, urogenital, blood, and endocrine diseases (6)	Chronic respiratory diseases (11)
65 yr and up	1	Other NCDs (13)	Neurological disorders (18)
	2	Digestive diseases except cirrhosis (13)	Other NCDs (13)
	3	Chronic respiratory diseases (12)	Musculoskeletal disorders (13)
	4	Neoplasms (12)	Digestive diseases except cirrhosis (11)
	5	Musculoskeletal disorders (12)	Cardiovascular and circulatory diseases (9)

The value in parentheses are percentages of the DALY of the disease among the total DALY by sex and age group.

DALY = disability-adjusted life year, NCD = non-communicable disease.

^aDALYs per 100,000 population.

to past trends in all-cause mortality by Statistics Korea. The projected total number of deaths in 2030 is 421,700 (227,036 for males; 194,664 for females) and that by Statistics Korea is 423,000 (222,000 for males; 201,000 for females).

The GBD projection used a methodology based on the assumption of a relationship between cause-specific mortality rates and a set of socioeconomic variables. The GBD study considered a limited number of socioeconomic variables — including average income per capita, the average number of years of schooling in adults, and time — to capture the impact of technological change on health status. In addition, tobacco use was included in the projections for cancers, cardiovascular diseases, and chronic respiratory diseases.⁵

We applied coherent functional data models to forecast cause-, age-, and sex-specific mortality in Korea from 2016 to 2030. The coherent functional data model is a variant of the Lee–Carter model, which extrapolates historical trends and forecasts the probability distributions of age-specific death rates using a statistical time-series method.¹¹ The time-series method presupposes that the historical trend will continue into the future, and is regarded to be superior to structural modeling using exogenous variables for forecasting.¹² The coherent functional data model is a product-ratio method based on forecasting of the simple product function (the geometric mean of subpopulation death rates) and the ratio function (ratios of the subpopulation death rates to the geometric mean rates).⁷ In this

paradigm, coherence is imposed and the forecasts of cause-specific mortality for the two sexes maintain certain structural relationships without continuing to diverge.^{7,13-15}

This study had limitations that should be considered. The forecasting is based on the assumption that past trends would be reflected in future trends, and unpredictable future occurrences are not considered.¹⁵ In addition, we assumed that the YLD to YLL ratio in 2015 would remain unchanged in the future. However, the YLD to YLL ratio can increase if the mortality rate decreases or decrease if the disease burden decreases due to improvements in medical technologies and health interventions.

To the best of our knowledge, this is the first projection of the future burden of a comprehensive set of diseases in Korea. Our results will facilitate establishment of appropriate health interventions to reduce the burdens imposed by various diseases.

REFERENCES

1. Korean Statistical Information Service. Population projections and summary indicators by scenarios annual 2015–2065. http://kosis.kr/eng/statisticsList/statisticsList_01List.jsp?vwcd=MT_ETITLE&parmTabId=M_01_01#SubCont. Updated 2017. Accessed October 24, 2017.
2. Yoon J, Oh IH, Seo H, Kim EJ, Gong YH, Ock M, et al. Disability-adjusted life years for 313 diseases and injuries: the 2012 Korean Burden of Disease Study. *J Korean Med Sci* 2016;31 Suppl 2:S146-57.
[PUBMED](#) | [CROSSREF](#)
3. Kim YE, Park H, Jo MW, Oh IH, Go DS, Jung J, et al. Trends and patterns of burden of disease in Korea using disability-adjusted life years. *J Korean Med Sci* 2019;34 Suppl 1:e75.
[CROSSREF](#)
4. Murray CJ, Lopez AD. Alternative projections of mortality and disability by cause 1990–2020: Global Burden of Disease Study. *Lancet* 1997;349(9064):1498-504.
[PUBMED](#) | [CROSSREF](#)
5. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* 2006;3(11):e442.
[PUBMED](#) | [CROSSREF](#)
6. Korean Statistical Information Service. Life expectancy at birth. http://kosis.kr/eng/statisticsList/statisticsList_01List.jsp?vwcd=MT_ETITLE&parmTabId=M_01_01#SubCont. Updated 2017. Accessed October 24, 2017.
7. Hyndman RJ, Booth H, Yasmeeen F. Coherent mortality forecasting: the product-ratio method with functional time series models. *Demography* 2013;50(1):261-83.
[PUBMED](#) | [CROSSREF](#)
8. GBD 2015 Mortality and Causes of Death Collaborators. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016;388(10053):1459-544.
[PUBMED](#) | [CROSSREF](#)
9. GBD 2015 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries during 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016;388(10053):1545-602.
[PUBMED](#) | [CROSSREF](#)
10. Ock M, Lee JY, Oh IH, Park H, Yoon SJ, Jo MW. Disability weights measurement for 228 causes of disease in the Korean Burden of Disease Study 2012. *J Korean Med Sci* 2016;31 Suppl 2:S129-38.
[PUBMED](#) | [CROSSREF](#)
11. Lee RD, Carter LR. Modeling and forecasting U.S. mortality. *J Am Stat Assoc* 1992;87(419):659-71.
[CROSSREF](#)
12. Booth H. Demographic forecasting: 1980 to 2005 in review. *Int J Forecast* 2006;22(3):547-81.
[CROSSREF](#)

13. Li N, Lee R. Coherent mortality forecasts for a group of populations: an extension of the Lee–Carter method. *Demography* 2005;42(3):575-94.
[PUBMED](#) | [CROSSREF](#)
14. Hyndman RJ, Ullah MS. Robust forecasting of mortality and fertility rates: a functional data approach. *Comput Stat Data Anal* 2007;51(10):4942-56.
[CROSSREF](#)
15. Hyndman RJ, Booth H. Stochastic population forecasts using Functional data models for mortality, fertility and migration. *Int J Forecast* 2008;24(3):323-42.
[CROSSREF](#)