

Research Article

# Expression of human miR-200b-3p and -200c-3p in cytomegalovirus-infected tissues

Kyoung Hwa Lee<sup>1,\*</sup>, Beom Jin Lim<sup>2,\*</sup>, Victor H. Ferreira<sup>3</sup>, Seo Yeon Min<sup>1</sup>, Yeon-Mi Hong<sup>1</sup>, Jeong-Hyeon Jo<sup>2</sup> and Sang Hoon Han<sup>1</sup>

<sup>1</sup>Division of Infectious Disease, Department of Internal Medicine, Yonsei University College of Medicine, Seoul, Republic of Korea; <sup>2</sup>Department of Pathology, Yonsei University College of Medicine, Seoul, Republic of Korea; <sup>3</sup>Multi-Organ Transplant Program, University Health Network, Toronto, Ontario, Canada

Correspondence: Sang Hoon Han (shhan74@yuhs.ac)



Human cytomegalovirus (HCMV) infection can cause inflammatory tissue-invasive end-organ diseases upon lytic replication. In humans, mature miR-200b-3p and -200c-3p suppress the synthesis of HCMV immediate early 2 (IE2) protein by binding to the 3'-UTR of the mRNA encoded by the unique long (UL) 122-123 region in human foreskin fibroblasts and pre-transplant peripheral blood mononuclear cells stimulated with HCMV. The present study aimed to quantitate the expression of *Homo sapiens* (hsa)-miR-200b-3p and 200c-3p in HCMV-infected tissues. We collected 240 HCMV-infected and 154 HCMV-non-infected, formalin-fixed, paraffin-embedded tissue samples of the gastrointestinal (GI) tract and bronchi/lungs. MiRNAs, HCMV, and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) were quantitated by quantitative reverse transcription-PCR (qRT-PCR) and quantitative PCR (qPCR) on the basis of standard curves generated using miRNA mimics, the HCMV strain from National Institute for Biological Standards and Control (NIBSC) 09/162, and GAPDH control. To avoid the effect of cell counts on the qRT-PCR and qPCR results, the data were normalized to GAPDH levels. HCMV-infected tissues had significantly lower levels of 200b-3p/GAPDH ( $3.03 \pm 1.50$  compared with  $3.98 \pm 1.08 \log_{10}$  copies/ $\mu\text{l}$ ,  $P < 0.001$ ) and 200c-3p/GAPDH ( $4.67 \pm 1.84$  compared with  $6.35 \pm 1.47 \log_{10}$  copies/ $\mu\text{l}$ ,  $P < 0.001$ ) than normal tissues. The values for 200b-3p/GAPDH ( $r = -0.51$ ,  $P < 0.001$ ) and 200c-3p/GAPDH ( $r = -0.54$ ,  $P < 0.001$ ) were significantly inversely correlated with HCMV load. Low tissue levels of 200b-3p and 200c-3p in humans are associated with cytopathic inflammation due to HCMV infection.

## Introduction

Human cytomegalovirus (HCMV) replication can cause serious harm in solid organ transplantation (SOT) and hematopoietic stem cell transplantation (HSCT) recipients [1–3]. In particular, post-transplant HCMV viremia has various indirect immunomodulatory effects, such as chronic allograft dysfunction and failure, in SOT recipients [1,2]. HCMV reactivation has been associated with increased mortality and prolonged treatment of mechanical ventilation in critically ill intensive care unit patients without prior objective immunocompromised conditions [4–7]. This worsening of outcomes by HCMV infection has led to the implementation of regular HCMV monitoring as well as well-organized and patient-tailored preventive strategies with risk stratification in routine clinical practice of SOT and HSCT recipients [1–3]. In addition, clinical trials have been conducted for evaluating the effects of HCMV prevention on outcomes in non-immunocompromised critically ill patients with acute respiratory distress syndrome [8].

Numerous basic and clinical studies fully support the need for continuous suppression and elimination of HCMV replication, a need that is currently unmet [9]. *In vitro* experiments on cultured cells have shown that epigenetic regulation through chromatin structural changes in the major immediate-early promoter (MIEP) and miRNA originating from the human host and/or HCMV

\* These authors contributed equally to this work.

Received: 14 June 2018  
Revised: 24 October 2018  
Accepted: 25 October 2018

Accepted Manuscript Online:  
26 October 2018  
Version of Record published:  
07 December 2018

play important roles in balancing latency and cytopathic replication [10–14]. Therefore, complete blockage of the production of immediate-early (IE) protein 2 (pp86), which is essential for the lytic virion structure and is encoded by the HCMV UL122-123 region, could be an effective approach to maintaining latency [12–15].

Seed sequences of human (*Homo sapiens* (Hsa)) miRNAs bind to the 3'-UTR of certain mRNAs and generally inhibit protein synthesis [16]. *In vitro* studies revealed that hsa-mature miRNA (miR)-200b-3p, -200c-3p, and -429, belonging to the miRNA-200 family, bind the 3'-UTR of HCMV UL122-123, in accordance with *in silico* predictions, and decrease IE2 synthesis in HCMV-infected fibroblasts [12,17]. Moreover, higher levels of hsp-miR-200b-3p and -200c-3p in pre-transplant blood with *in vitro* HCMV stimulation were associated with lower post-transplant HCMV replication rates in SOT recipients [17]. These results suggest that hsa-miR-200b-3p, -200c-3p, and -429 might serve as biomarkers to predict HCMV infection and/or disease and as therapeutic targets to control HCMV replication. The present study aimed to confirm the association between the levels of hsa-miR-200b-3p or -200c-3p or -429 and HCMV load in a clinical setting, with a focus on tissue-invasive disease, using formalin-fixed paraffin-embedded (FFPE) tissues.

## Methods

### FFPE tissue collection

We collected 240 whole-block HCMV-infected FFPE tissues between January 2007 and August 2016 in Shinchon and Gangnam Severance Hospital (Seoul, Korea) on the basis of information retrospectively extracted from electronic medical records. HCMV-infected tissues were strictly defined as those distinctly positive in immunohistochemical staining (IHS) using mouse anti-HCMV monoclonal antibody (mAb) (Clone DDG9 and CCH2; Agilent Dako Technologies, Seoul, South Korea) along with cytopathic histopathological findings of inflammation with/without necrosis as well as ulcerative/erosive lesions. After the exhaustive review of pathologic reports, we excluded FFPE tissues with following features: (i) suspected viral infection as indicated by the presence of giant cells with macronuclei/smudged chromatin/inclusion bodies, but IHS for HCMV was negative or not performed; (ii) co-infections with other organisms; (iii) patient was  $\leq 18$  years of age; and (iv) pathologic findings such as malignancies and acute/chronic inflammation caused by factors other than HCMV. As controls, 154 blocks of HCMV-non-infected age- and sex-matched FFPE tissues were obtained, between 2011 and 2016, from the resection margin of solid cancers and without any abnormal histological findings were included. The present study was approved by the institutional review board with waiver of written informed consent.

### DNA and RNA preparation

Total RNA was isolated from 20- $\mu\text{m}$ -thick FFPE tissue sections using a miRNeasy<sup>®</sup> FFPE Kit per the manufacturer's instructions (Qiagen, Seoul, Korea). Genomic DNA (gDNA) was extracted with a QIAamp<sup>®</sup> DNA FFPE Tissue kit per the manufacturer's instructions (Qiagen). All RNA and gDNA samples had  $A_{260}/A_{280}$  values of 1.8–2.0 as measured in a NanoDrop<sup>™</sup> 2000 spectrophotometer (Thermo Fisher Scientific, Seoul, Korea). All samples were stored at  $-80^{\circ}\text{C}$  until use.

### Quantitative real-time PCR and reverse transcription PCR

HCMV was quantitated by quantitative PCR (qPCR) using the National Institute for Biological Standards and Control (NIBSC) code 09/162 (NIBSC, Hertfordshire, U.K.) to obtain standard curves according to the World Health Organization (WHO) international standard guidelines for HCMV nucleic acid amplification techniques [18,19]. One vial of NIBSC code 09/162 was reconstituted in 1 ml, yielding  $5 \times 10^6$  international unit (IU)/ml [18,19]. The primers and probe for amplification of HCMV UL83 region were: forward, 5'-GCAGCCACGGGATCGTACT-3'; reverse, 5'-GGCTTTTACCTCACACGAGCATT-3'; probe, 5'-6(FAM)-CGCGAGACCGTGGAACTGCG-(TAMRA)-3' (Applied Biosystems/Thermo Fisher Scientific, Foster City, CA, U.S.A.) [20,21]. qPCRs were conducted using 50 ng/ $\mu\text{l}$  of gDNA, and for the standard curve,  $5 \times 10^1$  to  $5 \times 10^6$  IU/ml, and TaqMan<sup>™</sup> universal PCR master mixture under the following conditions: 2 min at  $50^{\circ}\text{C}$ , 10 min at  $95^{\circ}\text{C}$ , 45 cycles of 15 s at  $95^{\circ}\text{C}$ , and 1 min at  $60^{\circ}\text{C}$  [20,21].

Quantitative reverse transcription-PCR (qRT-PCR) of hsa-miR-200b-3p, -200c-3p, and -429 was performed with stem-loop primers and probes from TaqMan<sup>™</sup> Small RNA Assays (Assay IDs: 002251, 002300, and 001024, respectively) as well as TaqMan<sup>™</sup> universal PCR master mixture (Applied Biosystems/Thermo Fisher Scientific). Using 6 ng/ $\mu\text{l}$  of total RNA, RT was performed in a C1000 Touch<sup>™</sup> thermal cycler (Bio-Rad, Seoul, Korea) under the following conditions: 30 min at  $16^{\circ}\text{C}$ , 30 min at  $42^{\circ}\text{C}$ , and 5 min at  $85^{\circ}\text{C}$ . qRT-PCRs were run using 1.33  $\mu\text{l}$  of cDNA,

**Table 1 Characteristics of HCMV-infected and -non-infected, normal FFPE tissues**

| Characteristics   | Total (n=394) | HCMV-infected (n=240) | Normal (n=154) | P-value              |
|-------------------|---------------|-----------------------|----------------|----------------------|
| Age, years        | 57.7 ± 17.2   | 56.8 ± 16.9           | 59.2 ± 17.7    | 0.188 <sup>1</sup>   |
| Sex, male (%)     | 256 (65.0)    | 154 (64.2)            | 102 (66.2)     | 0.872 <sup>2</sup>   |
| Tissue type       |               |                       |                | 0.918 <sup>2,3</sup> |
| GI tract          | 370 (93.9)    | 235 (97.9)            | 135 (87.7)     |                      |
| Upper             | 122 (31.0)    | 67 (27.9)             | 56 (36.4)      |                      |
| Esophagus         | 58 (14.7)     | 31 (12.9)             | 27 (17.5)      |                      |
| Stomach           | 64 (16.2)     | 36 (15.0)             | 29 (18.9)      |                      |
| Lower             | 248 (62.9)    | 168 (65.8)            | 79 (51.3)      |                      |
| Small intestine   | 13 (3.3)      | 13 (5.4)              | 0 (0.0)        |                      |
| Large intestine   | 235 (59.6)    | 155 (64.6)            | 79 (51.3)      |                      |
| Colon             | 164 (41.6)    | 104 (43.3)            | 54 (35.1)      |                      |
| Rectum            | 71 (18.0)     | 51 (21.3)             | 25 (16.2)      |                      |
| Lung and bronchus | 24 (6.1)      | 5 (2.1)               | 19 (12.3)      |                      |

Data are expressed as mean ± S.D. or number (percent).

<sup>1</sup>Independent *t* test.

<sup>2</sup>Chi-square test.

<sup>3</sup>Comparison between three variables of upper GI tract, lower GI tract, and lung/bronchus.

with the following thermal cycles: 10 min at 95°C, 45 cycles of 15 s at 95°C, and 60 s at 60°C. Standard curves for quantitation were generated using mirVana™ miRNA mimics [22] at 10<sup>1</sup>–10<sup>9</sup> copies/μl.

To exclude the effects of severity of necrosis on HCMV quantity [14,23], we measured the expression of the house-keeping protein glyceraldehyde-3-phosphate dehydrogenase (GAPDH) in 100 ng/μl of gDNA through qPCR using TaqMan™ human GAPDH endogenous control (6FAM™/MGB probe, non-primer limited) according to the manufacturer's instructions (Applied Biosystems/Thermo Fisher Scientific). qPCRs were run in a LightCycler® 480 platform (Roche Diagnostics, Seoul, Korea). Each sample was tested in triplicate. Reliability analysis using inter-class correlation of C<sub>t</sub> values for each qRT-PCR or qPCR showed nearly perfect reproducibility as 0.996 for hsa-miR-200b-3p, 0.994 for -200c-3p, 0.987 for HCMV, and 0.959 for GAPDH.

The quantity of miRNAs or GAPDH was expressed as log<sub>10</sub> copies/μl of input RNA or log<sub>10</sub> copies/μl of input DNA. The HCMV load was expressed as log<sub>10</sub> IU/μl of input DNA. The limit of detection of all qRT-PCRs was defined as 2 log<sub>10</sub> copies/μl. All data under this threshold were considered as undetectable for categorical and zero for continuous variable analysis.


## Statistical analyses

Data were expressed as mean ± S.D. or median (interquartile range) or number (percent). Statistical analyses were performed using SPSS V23.0 (SPSS, Chicago, IL, U.S.A.) and GraphPad Prism V6 (GraphPad Software, La Jolla, CA). Differences in categorical and continuous variables amongst two groups were analyzed by chi-square and independent *t* test or Mann–Whitney U test, respectively. We used one-way ANOVA test and post-hoc Bonferroni's multiple comparison tests to compare three groups. Correlations between two continuous variables were expressed as Pearson's *r* value and 95% confidence interval (CI). To adjust the type I error by multiple independent comparisons, we considered the significant two-tailed *P*-value as less than 0.002 (0.05/25 of independent comparisons in the present study).

## Results

### Clinical features of patients and FFPE tissues

Table 1 summarizes the clinical characteristics of patients and FFPE tissues. All tissues originated from HCMV seropositive subjects. Mean age (*P*=0.188) and sex (*P*=0.872) were not different between HCMV-infected and non-infected tissues. The majority of tissues originated from the gastrointestinal (GI) tract, including the esophagus, stomach, colon, and rectum, and there was no significant difference in organ distribution between the two groups (*P*=0.918).



**Figure 1. Differences and correlation between levels of hsa-miR-200b-3p, -200c-3p, and -429 in 394 analyzed tissues samples**

(A) Differences in hsa-miR-200b-3p, -200c-3p, and -429 levels after normalization to GAPDH. Bars represent means and S.D. Dotted line indicates limit of detection (100 copies/ $\mu\text{L}$  of input nucleic acid). \* $P$ -value between three miRNAs by ANOVA test, \*\* $P < 0.001$  between two miRNAs by Bonferroni's multiple comparisons tests. (B) Correlation between 200b-3p and 200c-3p after normalization to GAPDH. The levels of hsa-miR-200b-3p/GAPDH and -200c-3p/GAPDH are strongly correlated, with  $r = 0.87$  (95% CI, 0.86–0.87) and  $P < 0.001$ . 'Nucleic acid' stands for DNA for GAPDH and RNA for miRNAs. Each dot corresponds to the miRNA/GAPDH level expressed as  $\log_{10}$  copies/ $\mu\text{L}$  of input nucleic acid in all figures.


### miRNA and HCMV levels in HCMV-infected and normal FFPE tissues

The levels of hsa-miR-200c-3p ( $7.19 \pm 1.71 \log_{10}$  copies/ $\mu\text{L}$ ) in all 394 tissues were significantly ( $P < 0.001$ ) higher than those of 200b-3p ( $4.76 \pm 1.47 \log_{10}$  copies/ $\mu\text{L}$ ). The 200c-3p/GAPDH had significantly higher values compared with 200b-3p/GAPDH ( $6.15 \pm 2.26$  compared with  $3.83 \pm 2.15 \log_{10}$  copies/ $\mu\text{L}$ ) with a 211-fold difference. The levels of miR-429/GAPDH ( $0.95 \pm 0.85 \log_{10}$  copies/ $\mu\text{L}$ ) was lowest amongst three miRNAs ( $P < 0.001$ ) (Figure 1A). Because few samples (34 of 394, 8.6%) had expression levels above limit of detection for miR-429, we performed additional analyses with 200b-3p and -200c-3p. The expression of 200b-3p and 200c-3p showed a strong positive correlation ( $r = 0.85$ , 95% CI, 0.82–0.88,  $P < 0.001$ ), even when the level of each miRNA was normalized to that of GAPDH mRNA in all 394 tissues ( $r = 0.87$ , 95% CI, 0.86–0.87,  $P < 0.001$ ) (Figure 1B). The HCMV load in the 240 HCMV-infected FFPE tissues was  $6.15 \pm 1.19$  (range: 4.11–9.69)  $\log_{10}$  IU/ml, whereas that in all normal tissues was less than 0.7  $\log_{10}$  IU/ml of the lower limit of quantitation.

HCMV-infected tissues had significantly lower levels of both hsa-miR-200b-3p ( $4.26 \pm 1.44$  compared with  $5.55 \pm 1.13 \log_{10}$  copies/ $\mu\text{L}$ ,  $P < 0.001$ ) and 200c-3p ( $6.50 \pm 1.60$  compared with  $8.26 \pm 1.27 \log_{10}$  copies/ $\mu\text{L}$ ,  $P < 0.001$ ) than normal tissues, with 19.5- and 57.5-fold differences, respectively. After normalization to GAPDH, fold-differences in both 200b-3p and 200c-3p were decreased by 8.9 and 47.9, respectively (200b-3p/GAPDH:  $3.03 \pm 1.50$  compared with  $3.98 \pm 1.08 \log_{10}$  copies/ $\mu\text{L}$ ,  $P < 0.001$  and 200c-3p;  $4.67 \pm 1.84$  compared with  $6.35 \pm 1.47 \log_{10}$  copies/ $\mu\text{L}$ ,  $P < 0.001$ ) (Figure 2A,B). The number of samples with levels below the detection limit for hsa-miR-200b-3p (9.2 compared with 0.0%,  $P < 0.001$ ) and 200b-3p/GAPDH (23.3 compared with 1.3%,  $P < 0.001$ ) as well as 200c-3p (3.3 compared with 0.0%,  $P = 0.025$ ) and 200c-3p/GAPDH (10.0 compared with 0.0%,  $P < 0.001$ ) was significantly higher in the HCMV-infected tissues.


### Hsa-miR-200b-3p and -200c-3p levels according to tissue type

HCMV-infected tissues from the upper GI tract ( $n = 70$ ) had significantly lower levels of hsa-miR-200b-3p/GAPDH ( $2.93 \pm 1.47$  compared with  $4.02 \pm 1.03 \log_{10}$  copies/ $\mu\text{L}$ ,  $P < 0.001$ ) than normal upper GI tissues ( $n = 56$ ). The levels of 200c-3p/GAPDH ( $5.38 \pm 1.49$  compared with  $6.44 \pm 1.59 \log_{10}$  copies/ $\mu\text{L}$ ,  $P < 0.001$ ) were significantly lower in HCMV-infected than in normal upper GI tissues (Figure 3A). 200b-3p/GAPDH ( $3.11 \pm 1.49$  compared with  $4.00 \pm 0.93 \log_{10}$  copies/ $\mu\text{L}$ ,  $P < 0.001$ ) and 200c-3p/GAPDH ( $4.39 \pm 1.88$  compared with  $6.35 \pm 1.43 \log_{10}$  copies/ $\mu\text{L}$ ,  $P < 0.001$ ) were significantly lower in HCMV-infected lower GI tissues ( $n = 165$ ) than in normal lower GI tissues ( $n = 79$ ) (Figure 3B). Amongst the lung tissue samples, 200b-3p/GAPDH ( $P = 0.297$ ) and 200c-3p/GAPDH ( $P = 0.120$ ), levels were not different between HCMV-infected ( $n = 5$ ) and normal ( $n = 19$ ) tissues (Figure 3C).



**Figure 2.** Difference of hsa-miR-200b-3p and -200c-3p levels between HCMV-infected ( $n=240$ ) and non-infected ( $n=154$ ) FFPE tissues

(A), (B) Bars indicate means and S.D. Dotted lines represent limit of detection (100 copies/ $\mu\text{L}$  of input nucleic acid). ‘Nucleic acid’ stands for DNA for GAPDH and RNA for miRNAs. Each dot corresponds to the miRNA/GAPDH level expressed  $\log_{10}$  copies/ $\mu\text{L}$  of input nucleic acid in all figures.




**Figure 3.** Difference in hsa-miR-200b-3p and -200c-3p levels according to tissue type

(A) Upper-GI tract samples originating from esophagus and stomach, including 67 HCMV-infected and 56 non-infected FFPE tissues. The percentage of samples with undetectable level of hsa-miR-200b-3p/GAPDH (21.7 compared with 0.0%) was significantly higher in HCMV-infected than non-infected upper-GI tract tissues ( $P<0.001$ ). However, the percentage of undetectable 200c-3p/GAPDH (4.3 compared with 0.0%) was similar between the two groups ( $P=0.121$ ). (B) Lower-GI tract samples from duodenum, cecum, terminal ileum, colon, and rectum, including 168 HCMV-infected and 79 normal FFPE tissues. The percentage of samples with undetectable level of hsa-miR-200b-3p/GAPDH (23.0 compared with 0.0%) was significantly higher in HCMV-infected than non-infected lower-GI-tract tissues ( $P<0.001$ ). For 200c-3p, values normalized to GAPDH had a significantly high undetectable percentage (12.1 compared with 0.0%,  $P<0.001$ ). (C) Lung samples, including 5 HCMV-infected and 19 normal FFPE tissues. In lung, the 200b-3p/GAPDH and 200c-3p/GAPDH levels did not show significant difference between two groups ( $P=0.297$  and  $0.120$ , respectively). \* $P<0.001$ .  $P$ -values for GI tract and lung samples were obtained through independent  $t$  test and Mann-Whitney U test, respectively. In (A,B), height of boxes and upper/lower lines indicate mean and S.D. In (C), bars represent median and 25 or 75 percentile values. Each dot corresponds to the miRNA/GAPDH levels expressed as  $\log_{10}$  copies/ $\mu\text{L}$  of input nucleic acid in all figures.

## Association of hsa-miR-200b-3p and -200c-3p expression with HCMV levels

Both hsa-miR-200b-3p and -200c-3p were negatively correlated with HCMV levels (200b-3p;  $r = -0.69$ , 95% CI =  $-0.76$  to  $-0.60$ ,  $P<0.001$ , 200c-3p;  $r = -0.68$ , 95% CI =  $-0.74$  to  $-0.58$ ,  $P<0.001$ ) in HCMV-infected tissues. The



**Figure 4. Correlation between HCMV and hsa-miR-200b-3p or -200c-3p levels in HCMV-infected FFPE tissues**

Straight and dotted lines represent each equation according to  $r$  coefficient and 95% CI values, respectively. Each dot corresponds to the miRNA/GAPDH or HCMV level expressed as  $\log_{10}$  copies/ $\mu\text{L}$  of input nucleic acid or DNA in all figures. ‘Nucleic acid’ stands for DNA for GAPDH and RNA for miRNAs.

correlation coefficients after normalization to GAPDH maintained the statistical significance (200b-3p/GAPDH;  $r = -0.51$ , 95% CI =  $-0.61$  to  $-0.39$ ,  $P < 0.001$ , 200c-3p;  $r = -0.54$ , 95% CI =  $-0.64$  to  $-0.042$ ,  $P < 0.001$ ) (Figure 4A,B).

## Discussion

Our study revealed that the expression of hsa-miR-200b-3p and -200c-3p was decreased in HCMV-infected tissues. This finding is in-line with those in two previous studies that evaluated the levels of two miRNAs in pre-transplant peripheral blood mononuclear cells (PBMCs) from SOT recipients with *in vitro* HCMV stimulation, and performed co-transfection assays using a laboratory HCMV strain and two miRNA mimics in fibroblasts [12,17]. This is the first study using HCMV-infected FFPE tissues, which are more representative of the clinical situation because these tissues definitely indicate HCMV end-organ disease.

The suppressive role of hsa-miR-200b-3p and -200c-3p on HCMV lytic replication was supported by two stepwise objective bases: (i) accurate prediction through *in silico* analysis of miRNAs binding to the 3'-UTR in the HCMV UL122-123 rather than high-throughput screening methods, such as oligonucleotide microarrays, in which expression profiles could be biased due to sample characteristics, platforms, or statistical analyses [24,25], (ii) verification of the binding of identical seed sequences in hsa-miR-200b-3p and -200c-3p to the HCMV UL122-123 by luciferase reporter assay using wild-type and mutant recombinant vectors, the results of which precisely corresponded with *in silico* prediction [12,17].

The results of the different experiments in the present study were consistent, allowing us to derive a reasonable conclusion. In spite of the relatively small differences in hsa-miR-200b-3p or -200c-3p levels between HCMV-infected and non-infected upper GI tissues, the levels of both miRNAs were significantly decreased in HCMV-infected upper/lower GI tissues. These findings were consistent between the quantitative and categorical analyses. We identified a strong negative correlation between the two miRNAs and HCMV levels; however, we were not able to establish that this correlation is directly caused by a reduction in IE2 production. This inability could be attributed to the fact that the HCMV-infected tissues do not have measurable levels of IE2 owing to the presence of large numbers of complete virions. In fact, we could not detect the mRNA encoding IE2 by qRT-PCR (data not shown). Finally, normalization of the expression data to that of GAPDH did not significantly alter the findings [14,23]. Our findings suggest that hsa-miR-200b-3p and -200c-3p could play relevant roles in regulating the initiation of HCMV active replication and/or latent-to-lytic switch by blocking the synthesis of major IE2 protein. This IE2 suppression could result in a serial reduction in early and late proteins, including HCMV DNA polymerase or glycoproteins [26]. Finally, this process could inhibit the extracellular budding of complete HCMV virion with envelope. However, the cross-reaction between these hsa-miRNAs and miRNAs encoded by HCMV, as well as the effect of MIEP, a strong transcription factor for major IE proteins, should be explored [27–30].

The major limitation of the present study would be the quality of RNA extracted from the FFPE tissues. MiRNAs can be degraded in FFPE tissues, and this damage would affect qRT-PCR results [31,32]. We did not check RNA integrity number by capillary electrophoresis. However, in addition to good  $A_{260}/A_{280}$  values, the following features

were considered while concluding that our RNA samples were of sufficient quality to yield satisfactory results: (i) triplicate qRT-PCRs and qPCRs for all samples showed nearly perfect inter-class correlation of  $C_t$  values; (ii) very strong correlation between the levels of hsa-miR-200b-3p and -200c-3p, with  $r = 0.85$ ; and (iii) 200c-3p levels were higher than 200b-3p levels, similar to the findings in human PBMCs [17]. Most tissue samples were of the GI tract and a small number of the lungs, because clinical specimens of HCMV-infected tissues such as liver, lung, and central nervous system, are hard to obtain. Despite these limitations, this first evaluation of hsa-miR-200b-3p and -200c-3p in end-organ FFPE tissues gives insights in HCMV infection in association with these miRNAs in the real clinical setting.

In conclusion, the levels of hsa-miR-200b-3p and -200c-3p are reduced in HCMV-infected tissues. These miRNAs likely play a biologically relevant role in the control of HCMV after transplantation.

## Clinical perspectives

- HCMV infection can result in cytopathic tissue-invasive inflammatory diseases. In addition, active replication of HCMV is closely associated with mortality and several morbidities such as graft rejection/dysfunction/failure, graft-versus-host disease, increase in other opportunistic infections in recipients of SOT and HSCT.
- hsa-miR-200b-3p and -200c-3p suppressed the synthesis of IE2 by binding the 3'-UTR of mRNA encoded by the HCMV UL122-123 region in human foreskin fibroblasts (*in vitro*) and in pre-transplant PBMCs stimulated by HCMV (in SOT recipients).
- MiR-200b-3p and -200c-3p levels were lower in HCMV-infected tissues than in normal tissues. The levels of miR-200b-3p and -200c-3p were inversely correlated with HCMV load. These miRNAs likely play a biologically relevant role in the control of cytopathic inflammation or HCMV infection after transplantation.

## Acknowledgements

We thank Atul Humar and Deepali Kumar at the Multi-Organ Transplant Program, University Health Network, Toronto, Ontario, Canada for review and comment on manuscript.

## Funding

This work was supported by the Faculty Research Grant from the Department of Internal Medicine, Yonsei University College of Medicine for 2016.

## Competing interests

The authors declare that there are no competing interests associated with the manuscript.

## Author contribution

All authors revised the manuscript critically and approved the final version of the submitted manuscript. K.H.L. and B.J.L. contributed equally to the design of the study, co-ordination of the research, and analysis and interpretation of the data. S.Y.M., Y.-M.H., and S.H.H. performed the experiments. K.H.L., B.J.L., V.H.F, and S.H.H. wrote the manuscript. B.J.L. and J.-H.J. selected and prepared all tissue specimens.

## Abbreviations

CI, confidence interval; FFPE, formalin-fixed paraffin-embedded; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; gDNA, genomic DNA; GI, gastrointestinal; HCMV, human cytomegalovirus; hsa, *Homo sapiens*; HSCT, hematopoietic stem cell transplantation; IE, immediate early protein; IHS, immunohistochemical staining; miR, mature microRNA; NIBSC, National Institute for Biological Standards and Control; PBMC, peripheral blood mononuclear cell; qPCR, quantitative PCR; qRT-PCR, quantitative reverse transcription-PCR; SOT, solid organ transplantation; UL, unique long.

## References

- 1 Razonable, R.R. and Humar, A. (2013) Cytomegalovirus in solid organ transplantation. *Am. J. Transplant.* **13**, 93–106, <https://doi.org/10.1111/ajt.12103>
- 2 Kotton, C.N. et al. (2013) Updated international consensus guidelines on the management of cytomegalovirus in solid-organ transplantation. *Transplantation* **96**, 333–360, <https://doi.org/10.1097/TP.0b013e31829df29d>
- 3 Fuji, S., Einsele, H. and Kapp, M. (2017) Cytomegalovirus disease in hematopoietic stem cell transplant patients: current and future therapeutic options. *Curr. Opin. Infect. Dis.* **30**, 372–376, <https://doi.org/10.1097/QCO.0000000000000375>
- 4 Kalil, A.C. and Florescu, D.F. (2009) Prevalence and mortality associated with cytomegalovirus infection in nonimmunosuppressed patients in the intensive care unit. *Crit. Care Med.* **37**, 2350–2358, <https://doi.org/10.1097/CCM.0b013e3181a3aa43>
- 5 Limaye, A.P. et al. (2008) Cytomegalovirus reactivation in critically ill immunocompetent patients. *JAMA* **300**, 413–422, <https://doi.org/10.1001/jama.2008.697>
- 6 Papazian, L. et al. (2016) Cytomegalovirus reactivation in ICU patients. *Intensive Care Med.* **42**, 28–37, <https://doi.org/10.1007/s00134-015-4066-9>
- 7 Ong, D.S. et al. (2016) Cytomegalovirus reactivation and mortality in patients with acute respiratory distress syndrome. *Intensive Care Med.* **42**, 333–341, <https://doi.org/10.1007/s00134-015-4071-z>
- 8 Limaye, A.P. et al. (2017) Effect of ganciclovir on IL-6 levels among cytomegalovirus-seropositive adults with critical illness: a randomized clinical trial. *JAMA* **318**, 731–740, <https://doi.org/10.1001/jama.2017.10569>
- 9 Griffiths, P.D. (2012) Burden of disease associated with human cytomegalovirus and prospects for elimination by universal immunisation. *Lancet Infect. Dis.* **12**, 790–798, [https://doi.org/10.1016/S1473-3099\(12\)70197-4](https://doi.org/10.1016/S1473-3099(12)70197-4)
- 10 Reeves, M.B. (2011) Chromatin-mediated regulation of cytomegalovirus gene expression. *Virus Res.* **157**, 134–143, <https://doi.org/10.1016/j.virusres.2010.09.019>
- 11 Kumar, A. and Herbein, G. (2014) Epigenetic regulation of human cytomegalovirus latency: an update. *Epigenomics* **6**, 533–546, <https://doi.org/10.2217/epi.14.41>
- 12 O'Connor, C.M., Vanicek, J. and Murphy, E.A. (2014) Host microRNA regulation of human cytomegalovirus immediate early protein translation promotes viral latency. *J. Virol.* **88**, 5524–5532, <https://doi.org/10.1128/JVI.00481-14>
- 13 Moller, R. et al. (2018) miRNA-mediated targeting of human cytomegalovirus reveals biological host and viral targets of IE2. *Proc. Natl. Acad. Sci. U.S.A.* **115**, 1069–1074, <https://doi.org/10.1073/pnas.1719036115>
- 14 Lau, B. et al. (2016) Human cytomegalovirus miR-UL112-1 promotes the down-regulation of viral immediate early-gene expression during latency to prevent T-cell recognition of latently infected cells. *J. Gen. Virol.* **97**, 2387–2398, <https://doi.org/10.1099/jgv.0.000546>
- 15 Paulus, C. and Nevels, M. (2009) The human cytomegalovirus major immediate-early proteins as antagonists of intrinsic and innate antiviral host responses. *Viruses* **1**, 760–779, <https://doi.org/10.3390/v1030760>
- 16 Bartel, D.P. (2004) MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* **116**, 281–297, [https://doi.org/10.1016/S0092-8674\(04\)00045-5](https://doi.org/10.1016/S0092-8674(04)00045-5)
- 17 Han, S.H. et al. (2017) Human microRNA responses predict cytomegalovirus replication following solid organ transplantation. *J. Infect. Dis.* **215**, 537–546
- 18 Fryer, J.F., Heath, A.B. and Minor, P.D. (2016) A collaborative study to establish the 1st WHO International Standard for human cytomegalovirus for nucleic acid amplification technology. *Biologicals* **44**, 242–251, <https://doi.org/10.1016/j.biologicals.2016.04.005>
- 19 Fryer, J.F., Heath, A.B., Anderson, R., Minor, P.D. and The Collaborative Study Group (2010) Collaborative study to evaluate the proposed 1st WHO International Standard for human cytomegalovirus (HCMV) for nucleic acid amplification (NAT)-based assays. *WHO ECBS Report*, WHO/BS/10.2138
- 20 Smithers-Sheedy, H. et al. (2017) Congenital cytomegalovirus among children with cerebral palsy. *J. Pediatr.* **181**, 267–271, <https://doi.org/10.1016/j.jpeds.2016.10.024>
- 21 Griscelli, F. et al. (2001) Quantification of human cytomegalovirus DNA in bone marrow transplant recipients by real-time PCR. *J. Clin. Microbiol.* **39**, 4362–4369, <https://doi.org/10.1128/JCM.39.12.4362-4369.2001>
- 22 Di Martino, M.T. et al. (2016) Functional analysis of microRNA in multiple myeloma. *Methods Mol. Biol.* **1375**, 181–194, [https://doi.org/10.1007/7651\\_2015\\_250](https://doi.org/10.1007/7651_2015_250)
- 23 Zhang, J. et al. (2013) Allitridin inhibits human cytomegalovirus replication *in vitro*. *Mol. Med. Rep.* **7**, 1343–1349, <https://doi.org/10.3892/mmr.2013.1328>
- 24 Balazsi, G. and Oltvai, Z.N. (2007) A pitfall in series of microarrays: the position of probes affects the cross-correlation of gene expression profiles. *Methods Mol. Biol.* **377**, 153–162, [https://doi.org/10.1007/978-1-59745-390-5\\_9](https://doi.org/10.1007/978-1-59745-390-5_9)
- 25 Jaksik, R. et al. (2015) Microarray experiments and factors which affect their reliability. *Biol. Direct* **10**, 46, <https://doi.org/10.1186/s13062-015-0077-2>
- 26 Landais, I. and Nelson, J.A. (2013) Functional genomics approaches to understand cytomegalovirus replication, latency and pathogenesis. *Curr. Opin. Virol.* **3**, 408–415, <https://doi.org/10.1016/j.coviro.2013.06.002>
- 27 Martinez, F.P. et al. (2014) CTCF binding to the first intron of the major immediate early (MIE) gene of human cytomegalovirus (HCMV) negatively regulates MIE gene expression and HCMV replication. *J. Virol.* **88**, 7389–7401, <https://doi.org/10.1128/JVI.00845-14>
- 28 Ng, K.R., Li, J.Y. and Gleadle, J.M. (2015) Human cytomegalovirus encoded microRNAs: hitting targets. *Expert Rev. Anti Infect. Ther.* **13**, 1469–1479, <https://doi.org/10.1586/14787210.2015.1106939>
- 29 Shen, K. et al. (2018) Human cytomegalovirus-encoded miR-UL112 contributes to HCMV-mediated vascular diseases by inducing vascular endothelial cell dysfunction. *Virus Genes* **54**, 172–181, <https://doi.org/10.1007/s11262-018-1532-9>
- 30 Stinski, M.F. and Isomura, H. (2008) Role of the cytomegalovirus major immediate early enhancer in acute infection and reactivation from latency. *Med. Microbiol. Immunol.* **197**, 223–231, <https://doi.org/10.1007/s00430-007-0069-7>



- 31 Evers, D.L. et al. (2011) Paraffin embedding contributes to RNA aggregation, reduced RNA yield, and low RNA quality. *J. Mol. Diagn.* **13**, 687–694, <https://doi.org/10.1016/j.jmoldx.2011.06.007>
- 32 Howe, K. (2017) Extraction of miRNAs from Formalin-Fixed Paraffin-Embedded (FFPE) tissues. *Methods Mol. Biol.* **1509**, 17–24, [https://doi.org/10.1007/978-1-4939-6524-3\\_3](https://doi.org/10.1007/978-1-4939-6524-3_3)