



Published in final edited form as:

Maturitas. 2019 April ; 122: 66–72. doi:10.1016/j.maturitas.2019.01.010.

Childhood experiences of parenting and age at menarche, age at menopause and duration of reproductive lifespan: Evidence from the English Longitudinal Study of Ageing

Panayotes Demakakos¹, Nora Pashayan², Georgios Chrousos³, Eleni Linara-Demakakou⁴, and Gita D. Mishra⁵

¹Department of Epidemiology and Public Health, University College London, London, United Kingdom

²Department of Applied Health Research, University College London, London, United Kingdom

³First Department of Pediatrics, National and Kapodistrian University of Athens Medical School, Aghia Sophia Children's Hospital, Athens, Greece

⁴The London Women's Clinic, London, United Kingdom

⁵School of Public Health, The University of Queensland, Brisbane, Australia.

Abstract

Objectives: The parent-child relationship is critical for human development, yet little is known about its association with offsprings' reproductive health outside the context of abuse and neglect. We investigated whether childhood experiences of poor-quality parenting (characterized as decreased parental care and increased parental overprotection) are associated with women's reproductive timing and lifespan.

Study design: Observational study of 2,383 women aged 55–89 years in 2007 from the English Longitudinal Study of Ageing (ELSA). Multinomial logistic regression models were estimated.

Corresponding author: Panayotes Demakakos, Department of Epidemiology and Public Health, University College London, 1-19 Torrington Place, London, WC1E 6BT, UK. p.demakakos@ucl.ac.uk.

Contributors

Panayotes Demakakos conceived and designed the study, analyzed data and drafted the manuscript.

Nora Pashayan, Georgios Chrousos, Eleni Linara-Demakakou and Gita D. Mishra contributed to the design of the study, and critically revised the manuscript for important intellectual content and approved its submission.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of interest

The authors declare that they have no conflict of interest.

Ethical approval

This study has been conducted in accordance with all relevant ethical regulations. It involves the analysis of publicly available secondary data from the ELSA study (www.elsa-project.ac.uk). ELSA has been approved by the London Multi-Centre Research Ethics Committee (MREC/01/2/91) and informed consent has been obtained by all ELSA participants.

Provenance and peer review

This article has undergone peer review.

Research data (data sharing and collaboration)

The ELSA data can be downloaded from the UK Data Service: <https://beta.ukdataservice.ac.uk/datacatalogue/studies/study?id=5050>

Main outcome measures: Self-reported ages at menarche and menopause and duration of reproductive lifespan.

Results: Increasing maternal and paternal overprotection were associated with later menarche (16 years) after adjustment for age and childhood socioeconomic position (relative risk ratio (RRR) 1.11, 95% CI 1.02–1.21 and 1.11, 95% CI 1.01–1.21, respectively, per unit increase in the predictor). Increasing parental overprotection and decreasing paternal care were associated with earlier menarche (10 years). However, these associations were marginally non-significant. Maternal and paternal overprotection were also inversely associated with age at natural menopause after adjustment for age, childhood socioeconomic position and age at menarche (p value for linear trend=0.041 and 0.004, respectively). Further, increasing paternal overprotection was associated with a shorter reproductive lifespan (33 years) (RRR 1.09 (1.01–1.18), per unit increase in the predictor) after adjustment for age and childhood socioeconomic position. Adjustment for additional childhood and adult factors did not explain these associations.

Conclusions: Women who experienced poor-quality parenting in childhood, especially increased levels of parental overprotection, might be at increased risk of an unfavourable reproductive health profile that is characterized by late or early menarche, premature menopause and a shorter reproductive lifespan.

Keywords

ageing; childhood; cohort; life course; menarche; menopause; parental overprotection; parental care; parenting; reproductive lifespan

1. Introduction

Menarche and menopause are two landmarks in women's reproductive history that define the duration of reproductive lifespan. They are also major determinants of women's health. Early menarche is associated with a number of health problems, including an unfavourable cardiovascular risk profile, and increased risk of breast, endometrial and ovarian cancer, and mortality [1–5]. Late menarche has been associated with health symptoms and conditions such as asthma [2]. Premature and early menopause are associated with an increased risk of chronic conditions including cardiovascular disease and mortality [6,7], while late menopause has been linked to an increased risk of breast, endometrial and ovarian cancer [1,5,8]. The duration of reproductive lifespan has also been associated with health problems, such as cardiovascular disease [9] and hormone-sensitive cancers, such as breast cancer [1].

Evidence suggests that childhood family environment can affect the timing of both menarche and menopause [10]. There is an extensive literature on the importance of abuse, neglect and an unfavourable family environment in the determination of age at menarche (AAM) [11–13], while familial and parental factors are also associated with earlier menopause [10]. However, most of this evidence stems from studies of smaller selective samples with only few studies having used large or nationally representative samples to examine the associations between the childhood experiences of parenting and AAM [14–16], age at natural menopause (AANM) and duration of reproductive lifespan in the offspring [14]. For this reason, and because the parent-child relationship is critical for human development and

childhood experiences of poor quality parenting are associated with increased risk of mortality [17] and cancer [18], we studied whether childhood experiences of poor quality parenting were also associated with AAM, AANM and the duration of reproductive lifespan in a national sample of older women. Drawing on earlier research [19], we defined poor quality parenting as low levels of paternal and maternal care and affection and high levels of paternal and maternal overprotection. Our hypothesis is that poor quality parenting is a potent childhood stressor and as such it could influence women's reproductive timing and health over the life course in multiple ways.

2. Methods

2.1 Study population

Our sample was drawn from the English Longitudinal Study of Aging (ELSA) (www.elsa-project.ac.uk). ELSA is an ongoing nationally representative observational study that begun in 2002–03 (ELSA wave 1) with a sample of 11,391 individuals (6,205 women) aged 50 years. For the needs of our study, we used data from the second follow-up interview (ELSA wave 3), which took place in 2006–07, and the 2007 ELSA Life History Interview, which was an one-off survey that collected retrospective information about the material circumstances, experiences and health of the ELSA participants before joining ELSA.

4,181 women participated in ELSA wave 3 of whom 3,442 participated in the ELSA Life History Interview. The analytical sample comprised 2,383 women aged 55 years in 2007 after the exclusion of 59 women due to very old age (>90 years), 491 women who did not complete the childhood experiences questionnaire, 298 women with missing values in the parenting measures, 180 women who were not reared by both natural parents and 31 with missing information on AAM (including 2 with AAM >20 years). For the needs of the AANM and duration of reproductive lifespan analyses, we used an analytical sample of 1674 women, after further excluding 561 women who experienced non-natural menopause (including 11 with missing information on age at menopause), 84 who had their natural menopause at unusually old >60 or young age <30 years, and 64 with missing values in covariates. The sample selection flowchart can be found in the Online Supplement (eFigure 1). ELSA has been approved by the London Multi-Centre Research Ethics Committee (MREC/01/2/91) and informed consent has been obtained by the participants.

2.2 Measures of childhood experiences of parenting

Parenting was measured as part of the ELSA Life History interview using the seven-item Parental Bonding Instrument (PBI). PBI is designed to collect retrospective information about the childhood experiences of parenting (at age 15 years) in adult samples and focuses on two fundamental dimensions of parenting, care and overprotection. Parental care refers to parental emotional warmth, affection, empathy, closeness and care for one's child as opposed to emotional coldness, indifference and neglect [19]. Parental overprotection refers to parental control, overprotection, intrusion, excessive contact and prevention of independent behaviour as opposed to allowance of independence and autonomy [19]. The seven-item PBI includes three care and four overprotection items and can be found here: <https://bit.ly/2LqwFMx> (see question 1). We generated care and overprotection summary

scores for both natural parents. To avoid the unnecessary exclusion of participants with few missing values in any of the parenting scales, we imputed up to one missing value per scale with the mean score of that scale (maternal overprotection was the scale with the largest number of such imputations, $n=69$). For comparison reasons, the analyses of the non-imputed data are presented in eTables 1–3.

2.2 Reproductive health outcomes

Information on women's health and reproductive history was self-reported and retrospectively collected. AAM, the age at first menstrual period, was measured as an ordinal variable with the following categories: 10, 11, 12, 13, 14, 15 and 16 years. AANM, was calculated by subtracting the year of birth from the year of last menstrual period for women who had natural menopause. We categorized the continuous AANM variable as follows: 30–39 years (premature menopause), 40–44 years (early menopause), 45–52 years and 53–60 years (late menopause). The duration of reproductive lifespan was calculated by subtracting AAM from AANM and categorized into groups of 3-year incremental differences [9] as follows: 33 years, 34–36 years, 37–39 years, 40 years.

2.3 Statistical analyses

We estimated multinomial logistic regression models. The predictor measures were used as continuous variables; risk estimates denote change in the outcome measure per unit decrease in maternal and paternal care scores or per unit increase in maternal and paternal overprotection scores. When modelling AAM, first, we estimated the unadjusted associations, which we then adjusted for age and childhood socioeconomic position (father's or main carer's occupation when respondent aged 14 years and number of books in the household when respondent aged 10 years). We followed a different modelling approach when analysing AANM and duration of the reproductive lifespan. We first estimated the unadjusted associations, which we then initially adjusted for age, and childhood socioeconomic position (in the AANM analyses we also included AAM in this model), and then adult socioeconomic position (education and total net non-pension household wealth including property, savings, and other assets), marital status, adult obesity (body mass index and waist circumference), lifetime smoking, and parity. In supplementary analyses, we adjusted our models for a number of additional childhood and adult factors that could have confounded the associations (see eTables 1–3).

3. Results

The mean age of the sample was 67.9 years (Table 1). The mean AANM was 50.3 years, the mean AAM was 13 years, and mean duration of reproductive lifespan was 37.2 years (Table 1). Childhood experiences of poor parenting were related with AAM (Table 2). Increasing paternal and maternal overprotection were significantly associated with a later menarche (16 years) (age- and childhood SEP-adjusted relative risk ratio (RRR): 1.11, 95% CI, 1.01, 1.21 and 1.11, 95% CI, 1.02, 1.21, respectively, per unit increase in the predictor). Along with decreasing paternal care, they were also associated with early menarche (10 years), but these associations were marginally non-significant. Further, we observed inverse associations between paternal and maternal overprotection and AANM (*P value* for linear

trend: 0.004 and 0.041, respectively, after adjustment for age, childhood socioeconomic position and AAM) (Table 3). Finally, we found that paternal overprotection was associated with a shorter reproductive lifespan (\approx 33 years) (RRR: 1.09, 95% CI, 1.01, 1.18, per unit increase in the predictor, after adjustment for age, childhood socioeconomic position and AAM) (Table 4). Additional adjustments for childhood and adult covariates did not explain these associations.

4. Discussion

In a national sample of older women, we found childhood experiences of poor parenting to be associated with an unfavourable reproductive health profile characterized by late or early menarche, premature natural menopause and a shorter reproductive lifespan. Maternal care, which is the most extensively studied parental factor in both animals and humans, appears to be less important for women's reproductive timing than parental overprotection, which was associated with both age at menarche and age at natural menopause. The preponderance of parental overprotection as a childhood determinant of reproductive development and lifespan over parental care is not surprising and concurs with literature highlighting parental overprotection as a risk factor for psychosocial development [20], and meta-analytic evidence suggesting that autonomy restriction, which is a hallmark of overprotective parenting, is the parental factor most strongly associated with an increased risk of depression in adolescence [21].

Our findings highlight the importance of the role of father for daughters' reproductive lifespan. Paternal overprotection was more strongly associated with a shorter reproductive lifespan than maternal overprotection in our data. There is extensive literature on the role of the father in the determination of AAM in the female offspring [12,13,22]. From an evolutionary perspective, fathers, unlike mothers, are expected to grant more autonomy, encourage independence, and prepare the offspring for the challenges of the life outside the family environment [23]. Based on this evidence, we can speculate that having an autonomy-restricting overprotective father can be more stressful and because of that potentially more harmful and more strongly associated with a shorter female offspring reproductive lifespan than having an overprotective mother.

4.1 Previous evidence

Our findings are partially discordant with those of a recent study that did not find an association between maternal overprotection and AAM [14]. Evidence suggests that a stressful family environment that is characterized by family conflict and disruption and father's absence is associated with earlier menarche [12], but studies that specifically examined factors such as a parental control over the child reported that harsh maternal and paternal control were associated with older age at menarche [11]. Our findings largely concur with this evidence. We found associations between decreased parental care and increased parental overprotection and both early menarche (\approx 10 years) (these associations were borderline non-significant though) and late menarche (\approx 16 years). Our findings are also concordant with evidence from national birth cohort studies suggesting that parental abuse is strongly associated with late menarche and more weakly with early menarche [16], and that

parental neglect, that is lack of interest in the offspring at age 7 years, is strongly associated with later menarche [15].

Fewer studies have examined the association between familial factors in childhood and menopause. Our findings are consistent with evidence suggesting an association between an unfavourable family environment in childhood that is characterized by conflict and parental divorce and an earlier age at menopause [25], but are at odds with findings suggesting that maternal overprotection is not associated with AANM and reproductive lifespan [14].

4.2 Strengths and weaknesses

Evidence on the association between childhood experiences of parenting and women's reproductive lifespan from large well-characterized studies is scarce. Our findings substantially add to the literature and improve our understanding of this relationship. The use of data from a nationally representative study such as ELSA also makes our findings more generalizable to community-dwelling women aged 55 years. Further, we were able to ascertain that adjustment for known childhood risk factors, such as childhood experiences of abuse and parental mental health and addiction problems, and adult risk factors, such as history of cancer, did not explain the observed associations. Finally, the use of PBI, which is a validated widely used instrument of parenting experiences, makes the replication of our work by future research easier.

Our study has weaknesses that should be considered. Its observational design makes it impossible to account for all potential confounders and eliminate the possibility of spurious associations. Further, our study adopted a simple "traditional" mediation approach, which allows neither a fuller exploration of the interrelationships between the study variables nor the estimation of direct and indirect effects. However, the diversity of our findings, that is different parenting measures being associated with three different outcome measures, and their consistency with earlier findings [17,18], makes it unlikely that they are a statistical artefact caused by unaccounted confounding. Further, in complementary analyses, we also found that potentially confounding factors that might introduce recall bias, such as mood and memory impairment, did not alter our findings.

The use of retrospectively collected childhood data makes our findings susceptible to measurement bias. Nevertheless, our parenting and childhood socioeconomic position measures have been used before and found to have good predictive validity, while a comparison of our retrospective menarche and menopause data with those of previous reports [26] provides good evidence for their validity, including capturing the well-documented downward secular trend in age at menarche (eTable 4 and eFigure 2). The same applies to reproductive lifespan duration; our estimate of mean lifespan duration of 37.2 years is almost identical with estimates reported by large US studies [9,27]. Further, the concordance of our findings with those from national birth cohort studies is reassuring and likely indicates that the observed associations represent real phenomena.

Non-response is another source of bias in our data. The overall individual response rate in ELSA wave 3 (after excluding people who died, became institutionalized or migrated) was 73%, with no noticeable gender differences. 84.4% of responders in wave 3 participated in

the ELSA Life History in 2007 [28], but again not of all of these people completed the self-completion questionnaire on childhood experiences that contained the parenting questions. Analyses of non-response in the ELSA Life History survey found significant differences in key characteristics such as socioeconomic position and health between responders and non-responders [17,29]. Based on these earlier findings, we can speculate that to an extent our findings are likely biased towards the null. Finally, statistical power is an issue as some analytical categories contained a relatively small number of participants and this led to wider 95% CI and increased uncertainty.

4.3 Pathways – poor quality parenting and age at menarche

Childhood experiences of poor parenting appear to be associated with AAM independently of low childhood socioeconomic position, adverse childhood experiences, such as abuse and parental mental health and addiction problems, and childhood health problems known to affect parenting. Notwithstanding our inability to account for other risk factors, such as maternal M, and childhood nutrition and obesity, these key findings point to the direction of a direct biological effect that can at least partially explain the association. quality parenting can be a chronic childhood stressor that may induce chronic alterations and dysregulations in the function of the neuroendocrine and immune systems and affect the developing brain, which in turn, could affect AAM.

We found that childhood experiences of poor parenting were associated with late menarche. We also found marginally non-significant associations between childhood experiences of poor parenting and early menarche. Considered together, these findings indicate that the effect of stress stemming from poor parenting experiences in childhood on AAM is not unidirectional and possibly there are important modifiers that determine the direction of this association. A recent review suggested that one such modifier might be the timing of the action of stressors, with early life stress leading to an earlier onset of puberty and juvenile or peripubertal stress delaying the onset of puberty [30]. Another such modifier can be genes. Evidence supports a gene-environment interaction hypothesis as the quality of the family environment has been found to be positively associated with AAM in participants homozygous for minor alleles of the estrogen receptor alpha gene (*ESR1*), but not in participants with other *ESR1* genotypes [31].

For any childhood exposure to delay or accelerate puberty and menarche, it should ultimately influence the activation of the hypothalamic–pituitary–gonadal (HPG) axis, whose core component is the pulsatile secretion of the Gonadotropin-releasing Hormone (GnRH) by hypothalamic GnRH neurons. GnRH is necessary for the secretion of gonadotropins, that is the follicle-stimulating hormone (FSH) and luteinizing hormone (LH), which are master regulators of the menstrual cycle and necessary for ovulation. Stress stemming from poor parenting experiences in childhood could affect multiple pathways involved in the activation of GnRH pulse generator. It may inhibit kisspeptin-mediated GnRH release. Kisspeptin (Kiss1) is a protein that plays a key stimulatory role in the activation of the GnRH pulse generator and the initiation of menarche [32]. It may also delay the onset of puberty via gamma-amino butyric acid- (GABA) and glutamate-mediated pathways [30], which play a critical role in the pubertal release of GnRH [33]. Further,

chronic stress in childhood stemming from experiences of poor quality parenting may also affect AAM by inducing epigenetic alterations [34].

4.4 Pathways – poor quality parenting and age at natural menopause

Low socioeconomic position, lifetime smoking, obesity, history of cancer, ages at menarche and first natural birth, and parity did not explain the association between poor quality parenting and AANM. Based on these findings, we hypothesize that childhood experiences of poor quality parenting could be directly associated with a younger AANM via biological mediating pathways. Multiple stress-related pathways might be implicated in this association, however all these pathways should influence a single biological parameter of crucial importance, the ovarian reserve, the number of non-growing primordial follicles in the ovaries.

A dysregulated stress system and prolonged activation of the HPA axis are expected to suppress the function of the HPG axis and the secretion of FSH and LH [35] and increase follicular atresia and degeneration [36]. Chronic stress could also affect the function of sympathetic nervous system, which releases norepinephrine in peripheral tissues. In the ovaries, norepinephrine is critical in the regulation of follicular development, ovulation and ovarian steroidogenesis [37]. Of importance in explaining our findings might also be stress-related pathways implicated in the decrease of the ovarian reserve before puberty, when the HPG axis is inactive. Such pathways may involve growth factors such as members of the transforming growth factor- β (TGF- β) superfamily [38], whose overactivation due to suppression of their regulators resulted in a considerable decrease of the ovarian reserve in prepubertal mice [39]. Also very important for premature menopause and regulated by growth factors, such as the insulin-like growth factor 1 (IGF1), is the intracellular phosphatidylinositol 3-kinase/Akt/mammalian target of rapamycin (PI3K/Akt/mTOR) signaling pathway, which is the master regulator of follicular activation and proliferation [40]. Increased activity of PI3K and mTOR may lead to increased activation of primordial follicles and premature “exhaustion” of the ovarian reserve. PI3K and mTOR pathways are also downregulated by different factors including oxytocin, a hypothalamic hormone that is related to maternal bonding with the newborn baby and parental behaviour, and its levels are lower in people who have experienced childhood adversity [41].

4.5 Conclusions

Using retrospectively collected childhood data, we found that childhood experiences of parenting might be a lifelong determinant of women’s reproductive timing and lifespan independently of other childhood and adult risk factors. On the understanding that these findings cannot simply be an artefact of measurement error and selection bias, our study adds to the current understanding of the role of childhood factors in women’s reproductive health. The importance of AAM and AANM for many health conditions, including cardiovascular disease, cancer and mortality, and the relevance of parenting to the vast majority of the population add to the scientific and societal value of our findings. Based on the assumption that poor quality parenting is a modifiable trait, our findings can inform prevention strategies and health policies. Future research should try to replicate our findings

and add to the exploration of the association between childhood experiences of poor quality parenting and reproductive lifespan in women.[24]

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding

The English Longitudinal Study of Ageing is supported by the National Institute on Aging (Grants 2RO1AG7644 and 2RO1AG017644–01A1) and a consortium of the UK government departments coordinated by the Economic and Social Research Council (ESRC). The National Institute on Aging and the consortium of the UK government departments had no role in the design and conduct of this study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript.

REFERENCES

- [1]. Collaborative Group on Hormonal Factors in Breast Cancer, Menarche, menopause, and breast cancer risk: individual participant meta-analysis, including 118 964 women with breast cancer from 117 epidemiological studies., *Lancet. Oncol* 13 (2012) 1141–51. doi:10.1016/S1470-2045(12)70425-4. [PubMed: 23084519]
- [2]. Day FR, Elks CE, Murray A, Ong KK, Perry JB, Puberty timing associated with diabetes, cardiovascular disease and also diverse health outcomes in men and women: the UK Biobank study, *Sci. Rep* 5 (2015) 11208. doi:10.1038/srep11208. [PubMed: 26084728]
- [3]. Charalampopoulos D, McLoughlin A, Elks CE, Ong KK, Age at menarche and risks of all-cause and cardiovascular death: a systematic review and meta-analysis., *Am. J. Epidemiol* 180 (2014) 29–40. doi:10.1093/aje/kwu113. [PubMed: 24920784]
- [4]. Gong T-T, Wu Q-J, Vogtmann E, Lin B, Wang Y-L, Age at menarche and risk of ovarian cancer: a meta-analysis of epidemiological studies., *Int. J. Cancer* 132 (2013) 2894–900. doi:10.1002/ijc.27952. [PubMed: 23175139]
- [5]. Brett M R., Jennifer B P., Thomas A S., Brett M R., Jennifer B P., Thomas A S., Epidemiology of ovarian cancer: a review, *Cancer Biol. Med* 14 (2017) 9–32. doi:10.20892/j.issn.2095-3941.2016.0084. [PubMed: 28443200]
- [6]. Fehring G, Kraft P, Pharoah PD, Eeles RA, Chatterjee N, Schumacher FR, Schildkraut JM, Lindström S, Brennan P, Bickeböller H, Houlston RS, Landi MT, Caporaso N, Risch A, Amin Al Olama A, Berndt SI, Giovannucci EL, Grönberg H, Kote-Jarai Z, Ma J, Muir K, Stampfer MJ, Stevens VL, Wiklund F, Willett WC, Goode EL, Permuth JB, Risch HA, Reid BM, Bezieau S, Brenner H, Chan AT, Chang-Claude J, Hudson TJ, Kocarnik JK, Newcomb PA, Schoen RE, Slattery ML, White E, Adank MA, Ahsan H, Aittomäki K, Baglietto L, Blomquist C, Canzian F, Czene K, Dos-Santos-Silva I, Eliassen AH, Figueroa JD, Flesch-Janys D, Fletcher O, Garcia-Closas M, Gaudet MM, Johnson N, Hall P, Hazra A, Hein R, Hofman A, Hopper JL, Irwanto A, Johansson M, Kaaks R, Kibriya MG, Lichtner P, Liu J, Lund E, Makalic E, Meindl A, Müller-Miyhsok B, Muranen TA, Nevanlinna H, Peeters PH, Peto J, Prentice RL, Rahman N, Sanchez MJ, Schmidt DF, Schmutzler RK, Southey MC, Tamimi R, Travis RC, Turnbull C, Uitterlinden AG, Wang Z, Whittemore AS, Yang XR, Zheng W, Buchanan DD, Casey G, V Conti D, Edlund CK, Gallinger S, Haile RW, Jenkins M, Le Marchand L, Li L, Lindor NM, Schmit SL, Thibodeau SN, Woods MO, Rafnar T, Gudmundsson J, Stacey SN, Stefansson K, Sulem P, Chen YA, Tyrer JP, Christiani DC, Wei Y, Shen H, Hu Z, Shu X-O, Shiraishi K, Takahashi A, Bossé Y, Obeidat M, Nickle D, Timens W, Freedman ML, Li Q, Seminara D, Chanock SJ, Gong J, Peters U, Gruber SB, Amos CI, Sellers TA, Easton DF, Hunter DJ, Haiman CA, Henderson BE, Hung RJ, Ovarian Cancer Association Consortium (OCAC), PRACTICAL Consortium, Hereditary Breast and Ovarian Cancer Research Group Netherlands (HEBON), Colorectal Transdisciplinary (CORECT) Study, African American Breast Cancer Consortium (AABC) and African Ancestry

- Prostate Cancer Consortium (AAPC), Cross-Cancer Genome-Wide Analysis of Lung, Ovary, Breast, Prostate, and Colorectal Cancer Reveals Novel Pleiotropic Associations., *Cancer Res* 76 (2016) 5103–14. doi:10.1158/0008-5472.CAN-15-2980. [PubMed: 27197191]
- [7]. Roeters van Lennep JE, Heida KY, Bots ML, Hoek A, on behalf of the collaborators of the D.M.G.D.G. on C.R.M. after R. Disorders, Cardiovascular disease risk in women with premature ovarian insufficiency: A systematic review and meta-analysis, *Eur. J. Prev. Cardiol* 23 (2016) 178–186. doi:10.1177/2047487314556004. [PubMed: 25331207]
- [8]. Karageorgi S, Hankinson SE, Kraft P, De Vivo I, Reproductive factors and postmenopausal hormone use in relation to endometrial cancer risk in the Nurses' Health Study cohort 1976–2004, *Int. J. Cancer* 126 (2010) 208–216. doi:10.1002/ijc.24672. [PubMed: 19551854]
- [9]. Ley SH, Li Y, Obias DK, Manson JE, Rosner B, Hu FB, Rexrode KM, Duration of Reproductive Life Span, Age at Menarche, and Age at Menopause Are Associated With Risk of Cardiovascular Disease in Women, *J. Am. Heart Assoc* 6 (2017) e006713. doi:10.1161/JAHA.117.006713. [PubMed: 29097389]
- [10]. Mishra GD, Cooper R, Tom SE, Kuh D, Early life circumstances and their impact on menarche and menopause., *Women's Heal* 5 (2009) 175–90. doi:10.2217/17455057.5.2.175.
- [11]. Belsky J, Steinberg LD, Houts RM, Friedman SL, DeHart G, Cauffman E, Roisman GI, Halpern-Felsher BL, Susman E, Family Rearing Antecedents of Pubertal Timing, *Child Dev* 78 (2007) 1302–1321. doi:10.1111/j.1467-8624.2007.01067.x. [PubMed: 17650140]
- [12]. Yermachenko A, Dvornyk V, Nongenetic determinants of age at menarche: a systematic review., *Biomed Res. Int* 2014 (2014) 371583. doi:10.1155/2014/371583. [PubMed: 25050345]
- [13]. Webster GD, Graber JA, Gesselman AN, Crosier BS, Schember TO, A Life History Theory of Father Absence and Menarche: A Meta-Analysis, *Evol. Psychol* 12 (2014). doi: 10.1177/147470491401200202.
- [14]. Magnus MC, Anderson EL, Howe LD, Joinson J, Penton-Voak IS, Fraser A, Childhood psychosocial adversity and female reproductive timing: a cohort study of the ALSPAC mothers, *J. Epidemiol. Community Health* 72 (2018) 34–40. doi:10.1136/jech-2017-209488. [PubMed: 29122994]
- [15]. Li L, Denholm R, Power C, Child maltreatment and household dysfunction: associations with pubertal development in a British birth cohort., *Int. J. Epidemiol* 43 (2014) 1163–73. doi: 10.1093/ije/dyu071. [PubMed: 24706731]
- [16]. Boynton-Jarrett R, Wright RJ, Putnam FW, Lividoti Hibert E, Michels KB, Forman MR, Rich-Edwards J, Childhood abuse and age at menarche., *J. Adolesc. Health* 52 (2013) 241–7. doi: 10.1016/j.jadohealth.2012.06.006. [PubMed: 23332491]
- [17]. Demakakos P, Pillas D, Marmot M, Steptoe A, Parenting style in childhood and mortality risk at older ages: a longitudinal cohort study, *Br. J. Psychiatry* 209 (2016) 135–141. doi:10.1192/bjp.bp.115.163543. [PubMed: 26941265]
- [18]. Demakakos P, Chrousos GP, Biddulph JP, Childhood experiences of parenting and cancer risk at older ages: Findings from the English Longitudinal Study of Ageing, *Int. J. Public Health* 63 (2018) 823–832. doi:10.1007/s00038-018-1117-3. [PubMed: 29947829]
- [19]. Parker G, Tupling H, Brown LB, A Parental Bonding Instrument, *Br. J. Med. Psychol* 52 (1979) 1–10. doi:10.1111/j.2044-8341.1979.tb02487.x.
- [20]. Parker G, Parental overprotection: A risk factor in psychosocial development., *Grune & Stratton*, 1983.
- [21]. Yap MBH, Pilkington PD, Ryan SM, Jorm F, Parental factors associated with depression and anxiety in young people: a systematic review and meta-analysis., *J. Affect. Disord* 156 (2014) 8–23. doi:10.1016/j.jad.2013.11.007. [PubMed: 24308895]
- [22]. Ellis BJ, Timing of Pubertal Maturation in Girls: An Integrated Life History Approach., *Psychol. Bull* 130 (2004) 920–958. doi:10.1037/0033-2909.130.6.920. [PubMed: 15535743]
- [23]. Paquette D, Theorizing the Father-Child Relationship: Mechanisms and Developmental Outcomes, *Hum. Dev* 47 (2004) 193–219. doi:10.1159/000078723.
- [24]. Tither JM, Ellis BJ, Impact of fathers on daughters' age at menarche: A genetically and environmentally controlled sibling study., *Dev. Psychol* 44 (2008) 1409–1420. doi:10.1037/a0013065. [PubMed: 18793072]

- [25]. Mishra G, Hardy R, Kuh D, Are the effects of risk factors for timing of menopause modified by age? Results from a British birth cohort study, *Menopause*. PAP (2007) 717–24. doi:10.1097/GME.0b013e31802f3156.
- [26]. Gentry-Maharaj A, Glazer C, Burnell M, Ryan A, Berry H, Kalsi J, Woolas R, Skates SJ, Campbell S, Parmar M, Jacobs I, Menon U, Changing trends in reproductive/lifestyle factors in UK women: descriptive study within the UK Collaborative Trial of Ovarian Cancer Screening (UKCTOCS)., *BMJ Open* 7 (2017) e011822. doi:10.1136/bmjopen-2016-011822.
- [27]. Nichols HB, Trentham-Dietz A, Hampton JM, Titus-Ernstoff L, Egan KM, Willett WC, Newcomb PA, From Menarche to Menopause: Trends among US Women Born from 1912 to 1969, *Am. J. Epidemiol* 164 (2006) 1003–1011. doi:10.1093/aje/kwj282. [PubMed: 16928728]
- [28]. Scholes S, Medina J, Cheshire H, Cox K, Hacker E, Carli L, Living in the 21st century: older people in England The 2006 English Longitudinal Study of Ageing - Technical Report, London, 2009 <http://doc.ukdataservice.ac.uk/doc/5050/mrdoc/pdf/5050Wave3TechnicalReport.pdf>.
- [29]. Ward K, Medina J, Mo M, Cox K, ELSA Wave Three: Life History Interview user guide to the data, London, 2009 http://www.esds.ac.uk/doc/5050/mrdoc/pdf/5050_Wave_3_Life_History_Documentation.pdf.
- [30]. Camille Melón L, Maguire J, GABAergic regulation of the HPA and HPG axes and the impact of stress on reproductive function, *J. Steroid Biochem. Mol. Biol* 160 (2016) 196–203. doi:10.1016/J.JSBMB.2015.11.019. [PubMed: 26690789]
- [31]. Manuck SB, Craig AE, Flory JD, Halder I, Ferrell RE, Reported early family environment covaries with menarcheal age as a function of polymorphic variation in estrogen receptor- α , *Dev. Psychopathol* 23 (2011) 69–83. doi:10.1017/S0954579410000659. [PubMed: 21262040]
- [32]. Livadas S, Chrousos GP, Control of the onset of puberty, *Curr. Opin. Pediatr* 28 (2016) 551–558. doi:10.1097/MOP.0000000000000386. [PubMed: 27386974]
- [33]. Herbison AE, Control of puberty onset and fertility by gonadotropin-releasing hormone neurons, *Nat. Rev. Endocrinol* 12 (2016) 452–466. doi:10.1038/nrendo.2016.70. [PubMed: 27199290]
- [34]. Cecil CAM, Smith RG, Walton E, Mill J, McCrory EJ, Viding E, Epigenetic signatures of childhood abuse and neglect: Implications for psychiatric vulnerability, *J. Psychiatr. Res* 83 (2016) 184–194. doi:10.1016/J.JPSYCHIRES.2016.09.010. [PubMed: 27643477]
- [35]. Chrousos GP, Torpy DJ, Gold PW, Interactions between the Hypothalamic-Pituitary-Adrenal Axis and the Female Reproductive System: Clinical Implications, *Ann. Intern. ed* 129 (1998) 229. doi:10.7326/0003-4819-129-3-199808010-00012.
- [36]. Yuan H-J, Han X, He N, Wang G-L, Gong S, Lin J, Gao M, Tan J-H, Glucocorticoids impair oocyte developmental potential by triggering apoptosis of ovarian cells via activating the Fas system, *Sci. Rep* 6 (2016) 24036. doi:10.1038/srep24036. [PubMed: 27040909]
- [37]. Roa J, Garcia-Galiano D, Varela L, Sánchez-Garrido MA, Pineda R, Castellano JM, Ruiz-Pino F, Romero M, Aguilar E, López M, Gaytan F, Diéguez C, Pinilla L, Tena-Sempere M, The Mammalian Target of Rapamycin as Novel Central Regulator of Puberty Onset via Modulation of Hypothalamic Kiss1 System, *Endocrinology* 150 (2009) 5016–5026. doi:10.1210/en.2009-0096. [PubMed: 19734277]
- [38]. Knight PG, Glistler C, TGF-beta superfamily members and ovarian follicle development., *Reproduction* 132 (2006) 191–206. doi:10.1530/rep.1.01074. [PubMed: 16885529]
- [39]. Rimon-Dahari N, Heinemann-Yerushalmi L, Hadas R, Kalich-Philosoph L, Ketter D, Nevo N, Galiani D, Dekel N, Vasorin: a newly identified regulator of ovarian folliculogenesis, *FASEB J* 32 (2018) 2124–2136. doi:10.1096/fj.201700057RRR. [PubMed: 29259033]
- [40]. Sengupta S, Peterson TR, Sabatini DM, Regulation of the mTOR Complex 1 Pathway by Nutrients, Growth Factors, and Stress, *Mol. Cell* 40 (2010) 310–322. doi:10.1016/J.MOLCEL.2010.09.026. [PubMed: 20965424]
- [41]. Johnson JL, Buisman-Pijlman FTA, Adversity impacting on oxytocin and behaviour, *Behav. Pharmacol* 27 (2016) 659–671. doi:10.1097/FBP.0000000000000269. [PubMed: 27755016]

Highlights

- Childhood experiences of poor-quality parenting, especially increased paternal overprotection, were associated with late menarche (16 years).
- Parental overprotection and decreased paternal care were associated with early menarche (10 years), but these associations were marginally non-significant.
- Maternal and paternal overprotection were also inversely associated with age at natural menopause.
- Increased paternal overprotection was associated with a shorter reproductive lifespan (33 years).
- Adjustment for several childhood and adult risk factors, including childhood experiences of abuse and low socioeconomic position, did not explain these associations.
- On the understanding that poor parenting experiences are a major childhood stressor with lifelong implications, and that known childhood and adult risk factors did not explain the associations, we hypothesize that our findings can partially be explained by stress-related biological pathways.

Table 1.

The baseline characteristics of the sample, English Longitudinal Study of Ageing, 2007 (n=2,383)^a

	N ^a (%)
Mean age (SD)	67.9 (8.8)
Paternal or main carer's occupation when respondent aged 14 years	
Manager/professional/administrator/own business	837 (35.1)
Trade/care/sales/services	724 (30.4)
Manual or casual jobs/unemployed	722 (30.3)
Other (including retired)	100 (4.2)
Number of books in the household when respondent aged 10 years	
Enough to fill two bookcases or more (>100 books)	474 (19.9)
Enough to fill one bookcase (26 to 100 books)	705 (29.6)
Enough to fill one shelf (11 to 25 books)	585 (24.5)
None or very few (0 to 10 books)	531 (22.3)
Missing	88 (3.7)
Current marital status	
Married	1514 (63.5)
Non-married	869 (36.5)
Education	
A-level or higher	765 (32.1)
Secondary or equivalent	830 (34.8)
No educational qualifications	788 (33.1)
Total household wealth (N=2335)	
Wealthiest tertile (£304,000)	787 (33.7)
Intermediate tertile (<£304,000 & £157,500)	782 (33.5)
Lowest tertile (<£157,500)	766 (32.8)
Smoking	
Never	1098 (46.1)
Ex-smoker	1005 (42.2)
Current smoker	280 (11.7)
Body mass index (kg/m²) (categories)	
<25	635 (26.7)
25 to <30	794 (33.3)
30	637 (26.7)
Missing	317 (13.3)
Waist circumference (categories)	
<94 cm in men / <80 cm in women	435 (18.3)
94 to 101 cm in men / 80 to 87 cm in women	490 (20.6)
102 cm in men / 88 cm in women	1183 (49.6)

	N^a (%)
Mean age (SD)	67.9 (8.8)
Missing	275 (11.5)
N of natural children (parity)	
None	336 (14.1)
1 child	441 (18.5)
2 children	935 (39.2)
3 children	671 (28.2)
Mean age at natural menopause (SD) (n=1674)	50.3 (4.6)
Age at natural menopause (categories) (n=1674)	
<40 years (premature menopause)	34 (2.1)
40 to 44 years (early menopause)	136 (8.1)
45 to 52 years	958 (57.2)
53 years (late menopause)	546 (32.6)
Mean age at menarche (SD)	13.0 (1.7)
Age at menarche (categories)	
10 years	120 (5.0)
11 years	394 (16.5)
12 years	366 (15.4)
13 years	543 (22.8)
14 years	504 (21.2)
15 years	291 (12.2)
16 years	165 (6.9)
Mean duration of reproductive lifespan (SD) (n=1674)	37.2 (4.9)
Duration of reproductive lifespan (categories) (n=1674)	
33 years	309 (18.5)
34 to 36 years	334 (19.9)
37 to 39 years	480 (28.7)
40 years	551 (32.9)

^a unless otherwise stated

Table 2.

The associations between parenting measures and age at menarche^a (N=2,383)

	10 years (n=120)	11 years (n=394)	12 years (n=366)	13 years (reference category) (n=543)	14 years (n=504)	15 years (n=291)	16 years (n=165)
Maternal Care Score (range: 0-highest levels of care to 9-lowest levels of care)							
Model 1 ^b	1.04 (0.94 to 1.14)	1.02 (0.96 to 1.09)	1.02 (0.95 to 1.09)	1.00	1.01 (0.95 to 1.08)	0.99 (0.92 to 1.07)	1.04 (0.95 to 1.13)
Model 2 ^c	1.03 (0.94 to 1.14)	1.02 (0.95 to 1.09)	1.02 (0.95 to 1.09)	1.00	1.02 (0.95 to 1.08)	1.00 (0.93 to 1.08)	1.04 (0.95 to 1.14)
Maternal Overprotection Score (range: 0-lowest levels of overprotection to 12-highest levels of overprotection)							
Model 1 ^b	1.08 (0.98 to 1.18)	1.05 (0.99 to 1.12)	1.02 (0.96 to 1.09)	1.00	1.04 (0.98 to 1.10)	1.00 (0.93 to 1.07)	1.11 (1.02 to 1.21) ^d
Model 2 ^c	1.08 (0.98 to 1.18)	1.05 (0.99 to 1.12)	1.02 (0.96 to 1.09)	1.00	1.04 (0.98 to 1.11)	1.00 (0.93 to 1.07)	1.11 (1.02 to 1.21) ^d
Paternal Care Score (range: 0- highest levels of care to 9-lowest levels of care)							
Model 1 ^b	1.09 (0.99 to 1.21)	1.03 (0.96 to 1.11)	1.00 (0.93 to 1.07)	1.00	0.99 (0.93 to 1.06)	0.97 (0.90 to 1.05)	1.03 (0.94 to 1.14)
Model 2 ^c	1.09 (0.99 to 1.21)	1.02 (0.95 to 1.10)	0.99 (0.92 to 1.07)	1.00	1.00 (0.93 to 1.07)	0.98 (0.90 to 1.06)	1.04 (0.95 to 1.15)
Paternal Overprotection Score (range: 0-lowest levels of overprotection to 12-highest levels of overprotection)							
Model 1 ^b	1.07 (0.97 to 1.18)	1.04 (0.97 to 1.11)	0.99 (0.93 to 1.06)	1.00	1.02 (0.96 to 1.08)	1.01 (0.94 to 1.09)	1.10 (1.00 to 1.20) ^d
Model 2 ^c	1.07 (0.97 to 1.18)	1.03 (0.97 to 1.11)	0.99 (0.92 to 1.06)	1.00	1.02 (0.96 to 1.09)	1.01 (0.94 to 1.09)	1.11 (1.01 to 1.21) ^d

^aThe estimates are relative risk ratios and denote change in the risk of experiencing younger or older age at menarche compared with the reference category per unit change in the predictor variable

^bThis is the unadjusted association

^cModel 2 is adjusted for age and childhood socioeconomic position (i.e. number of books in the household at age 10 years, and father's or main carer's occupational class at age 14 years)

^d*p* < 0.05

Table 3.

The association between parenting measures and age at natural menopause^a (N=1,674)

	30 to 39 years (premature menopause) (n=34)	40 to 44 years (early menopause) (n=136)	45 to 52 years (reference category) (n=958)	53 to 60 years (n=546)	P value for linear trend
Maternal Care Score					
Model 1 ^b	1.08 (0.92 to 1.28)	0.97 (0.88 to 1.07)	1.00	1.00 (0.94 to 1.05)	0.74
Model 2 ^c	1.15 (0.96 to 1.37)	0.96 (0.87 to 1.06)	1.00	0.99 (0.93 to 1.05)	0.55
Model 3 ^d	1.15 (0.96 to 1.37)	0.96 (0.87 to 1.06)	1.00	0.99 (0.93 to 1.05)	0.54
Maternal Overprotection Score					
Model 1 ^b	1.09 (0.92 to 1.28)	0.97 (0.89 to 1.07)	1.00	0.94 (0.89 to 0.99) ^e	0.040
Model 2 ^c	1.12 (0.94 to 1.34)	0.97 (0.88 to 1.06)	1.00	0.94 (0.89 to 0.99) ^e	0.041
Model 3 ^d	1.13 (0.93 to 1.36)	0.96 (0.87 to 1.05)	1.00	0.94 (0.89 to 0.99) ^e	0.035
Paternal Care Score					
Model 1 ^b	1.08 (0.91 to 1.30)	1.03 (0.93 to 1.14)	1.00	1.01 (0.95 to 1.07)	0.57
Model 2 ^c	1.14 (0.95 to 1.38)	1.03 (0.93 to 1.14)	1.00	1.00 (0.94 to 1.06)	0.40
Model 3 ^d	1.13 (0.93 to 1.37)	1.04 (0.94 to 1.15)	1.00	1.00 (0.94 to 1.07)	0.35
Paternal Overprotection Score					
Model 1 ^b	1.14 (0.97 to 1.34)	1.02 (0.93 to 1.11)	1.00	0.94 (0.89 to 1.00) ^e	0.007
Model 2 ^c	1.20 (1.01 to 1.43) ^e	1.01 (0.92 to 1.11)	1.00	0.94 (0.89 to 0.99) ^e	0.004
Model 3 ^d	1.18 (0.98 to 1.40)	1.01 (0.92 to 1.11)	1.00	0.94 (0.89 to 0.99) ^e	0.005

^aThe estimates are relative risk ratios and denote change in the risk of experiencing premature, early or later menopause compared with the reference category per unit change in the predictor variable

^bThis is the unadjusted association

^cModel 2 is adjusted for age, childhood socioeconomic position (i.e. number of books in the household at age 10 years and father's or main carer's occupational class at age 14 years), and age at menarche

^dModel 3 is adjusted for age, childhood socioeconomic position (i.e. number of books in the household at age 10 years and father's or main carer's occupational class at age 14 years), age at menarche, adult socioeconomic position (i.e. education and total net household wealth), marital status, smoking, body mass index, waist circumference, and parity

^e p < 0.05

Table 4. The association between parenting measures and duration of the reproductive lifespan^a (N=1,674)

	33 years (n=309)	34 to 36 years (n=334)	37 to 39 years (reference category) (n=480)	40 years (n=551)
Maternal Care Score				
Model 1 ^b	0.99 (0.92 to 1.06)	0.99 (0.92 to 1.07)	1.00	1.00 (0.93 to 1.06)
Model 2 ^c	1.00 (0.92 to 1.08)	1.00 (0.93 to 1.08)	1.00	1.00 (0.93 to 1.06)
Model 3 ^d	0.99 (0.92 to 1.07)	1.00 (0.93 to 1.08)	1.00	1.00 (0.93 to 1.06)
Maternal Overprotection Score				
Model 1 ^b	1.03 (0.96 to 1.11)	1.07 (1.00 to 1.15) ^e	1.00	1.00 (0.94 to 1.06)
Model 2 ^c	1.03 (0.96 to 1.11)	1.08 (1.00 to 1.16) ^e	1.00	1.00 (0.94 to 1.07)
Model 3 ^d	1.03 (0.96 to 1.11)	1.08 (1.00 to 1.16) ^e	1.00	1.00 (0.94 to 1.07)
Paternal Care Score				
Model 1 ^b	1.03 (0.95 to 1.12)	0.99 (0.92 to 1.07)	1.00	1.03 (0.96 to 1.10)
Model 2 ^c	1.05 (0.97 to 1.14)	1.00 (0.92 to 1.08)	1.00	1.03 (0.96 to 1.10)
Model 3 ^d	1.06 (0.97 to 1.15)	1.00 (0.92 to 1.08)	1.00	1.03 (0.96 to 1.11)
Paternal Overprotection Score				
Model 1 ^b	1.08 (1.01 to 1.17) ^e	1.09 (1.01 to 1.17) ^e	1.00	1.02 (0.96 to 1.09)
Model 2 ^c	1.09 (1.01 to 1.18) ^e	1.10 (1.02 to 1.18) ^e	1.00	1.02 (0.96 to 1.09)
Model 3 ^d	1.10 (1.02 to 1.19) ^e	1.10 (1.02 to 1.19) ^e	1.00	1.03 (0.97 to 1.10)

^aThe estimates are relative risk ratios and denote change in the risk of having a shorter or longer reproductive lifespan compared with the reference category per unit change in the predictor variable

^bThis is the unadjusted association

^cModel 2 is adjusted for age and childhood socioeconomic position (i.e. number of books in the household at age 10 years and father's or main carer's occupational class at age 14 years)

^dModel 3 is adjusted for age, childhood (i.e. number of books in the household at age 10 years and father's or main carer's occupational class at age 14 years), and adult socioeconomic position (i.e. education and total net household wealth), marital status, smoking, body mass index, waist circumference, and parity

^eP < 0.05