



Published in final edited form as:

J Magn Reson. 2010 April ; 203(2): 213–219. doi:10.1016/j.jmr.2009.12.017.

EPR characterization of ascorbyl and sulfur dioxide anion radicals trapped during the reaction of bovine Cytochrome *c* Oxidase with molecular oxygen

Michelle A. Yu, Tsuyoshi Egawa, Syun-Ru Yeh, Denis L. Rousseau*, and Gary J. Gerfen*

Department of Physiology and Biophysics, Albert Einstein College of Medicine, 1300 Morris Park Ave., Bronx, NY 10461

Abstract

The reaction intermediates of reduced bovine Cytochrome *c* Oxidase (CcO) were trapped following its reaction with oxygen at 50 μ s to 6 msec by innovative freeze-quenching methods and studied by EPR. When the enzyme was reduced with either ascorbate or dithionite, distinct radicals were generated; X-band (9 GHz) and D-band (130 GHz) CW-EPR measurements support the assignments of these radicals to ascorbyl and sulfur dioxide anion radical (SO₂^{-•}) respectively. The X-band spectra show a linewidth of 12 G for the ascorbyl radical and 11 G for the SO₂^{-•} radical and an isotropic *g*-value of 2.005 for both species. The D-band spectra reveal clear distinctions in the *g*-tensors and powder patterns of the two species. The ascorbyl radical spectrum displays approximate axial symmetry with *g*-values of $g_x = 2.0068$, $g_y = 2.0066$, and $g_z = 2.0023$. The SO₂^{-•} radical has rhombic symmetry with *g*-values of $g_x = 2.0089$, $g_y = 2.0052$, and $g_z = 2.0017$. When the contributions from the ascorbyl and SO₂^{-•} radicals were removed, no protein-based radical on CcO could be identified in the EPR spectra.

1. INTRODUCTION

Radical-based catalysis is integral to biological processes in organisms ranging from archaeobacteria [1] to mammals [2]. Protein-based radicals are essential to a variety of enzymatic systems, [3], such as ribonucleotide reductases [4, 5] and heme peroxidases [6–12]. Cytochrome *c* Oxidase (CcO), which catalyzes the reduction of oxygen to water, has been proposed to involve a protein-based radical [13–15] in its catalytic chemistry. Such a radical would explain how CcO sustains turnover without releasing reactive oxygen species even under conditions of limited electron availability. Despite impressive efforts and a variety of experimental approaches, the identity and role of radical species formed during the reaction of oxygen with CcO remain elusive.

Mammalian CcO is a 204 kDa membrane protein, which functions as Complex IV in the electron transport chain. Electrons derived from catabolism of fats, amino acids, and sugars

*To Whom Correspondence should be addressed: Denis.Rousseau@einstein.yu.edu, Gary.Gerfen@einstein.yu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

enter the chain at Complex I and arrive via Cytochrome *c* (Cyt *c*) at CcO, where a bound oxygen molecule is the terminal electron acceptor. CcO catalyzes this exergonic reduction of oxygen to water, harnessing the released energy to pump four net protons across the inner mitochondrial membrane. The reaction of CcO with oxygen has been heavily studied using small molecule reductants such as ascorbate or dithionite as the external electron source for Cyt *c* and ultimately CcO. From these studies, several intermediates along the catalytic pathway have been identified. One such intermediate, termed “**P**,” is believed to contain a protein-based radical, although its identity is highly debated [14, 16–18].

Wilson *et al.* detected a radical during the reaction of oxygen with fully reduced bovine CcO; the authors concluded that it was an artifact from the reductant, although they could not rule out a protein-based radical [18]. In a radioactive iodide study, Babcock and coworkers reported a radical on the modified tyrosine, Y244, which is believed to donate a proton and an electron during catalysis [13]. In the reaction of oxygen with the fully reduced forms of two bacterial oxidases, *P. denitrificans aa₃* and *E. coli bo₃*, de Vries and coworkers suggested that as many as three radicals, ascribed to tryptophanyl and an unknown “6 msec” species, form during turnover [14, 17]. However, neither the locations of the radicals nor their role in catalysis could be determined; furthermore, their formation could not be correlated with the expected steps in the catalytic cycle. In addition, L-ascorbic acid was used as the primary reductant, but its effect on the EPR spectra was not addressed.

L-ascorbic acid (AH₂) is an endogenous water-soluble antioxidant present in up to millimolar concentrations in the human body [19]. It has pK_a values of 4.2 and 11.6 [20] and exists predominantly as the anion, AH⁻, at neutral pH. Ascorbate oxidation involves the loss of a proton and an electron, which may occur via several possible mechanisms: (1) AH⁻ may lose an electron forming the neutral free radical, AH[•], which deprotonates to form the semidehydroascorbate radical anion, A^{-•}; (2) AH⁻ deprotonates to form the dianion, A²⁻, which then releases an electron to form A^{-•}; (3) AH⁻ forms A^{-•} directly by a concerted proton-electron transfer. The first mechanism can be ruled out by its unfavorable midpoint potential ($E^0 = +766$ mV) [21]. The second mechanism is suggested to occur in the reduction of Cyt *c* [22]. The third mechanism has been demonstrated in cytochrome *b561* [23]. In addition, recent energy calculations propose that a di-radical forms, although the mechanism is unclear [24, 25].

Although the mechanism may vary, the end product is conventionally accepted to be A^{-•}, which has been used as a spectroscopic indicator of oxidative stress due to its half-life of ~50 sec [26, 27]. A^{-•} has an unpaired electron in a highly delocalized π -system, conferring stability as the “terminal small-molecule antioxidant” [28]. A second one-electron oxidation produces dehydroascorbate, which is reduced and recycled in mitochondria to ascorbate [29–31]. Thus, the redox chemistry of ascorbate has immense physiological relevance as well as value as a laboratory reagent.

Sodium dithionite, which is another common reductant, has a midpoint potential of $E^0 = -420$ mV at pH 7 [32]. The dithionite ion, S₂O₄²⁻ can be considered a dimer of SO₂^{-•} radicals, as the S-S bond is exceptionally long [33]; the reduction reactions of dithionite were suggested to occur via the SO₂^{-•} radical, whose EPR spectra have been studied [34,

35]. $\text{SO}_2^{\bullet-}$ may arise from thermal decomposition prior to reduction or in the commercial preparation of sodium dithionite [35].

In biological systems, multiple radical species often occur in mechanisms involving radical migration between tryptophan, tyrosine, glycine, and cysteine residues as well as from a porphyrin pi-cation radical in heme systems [3]. These systems give rise to complex EPR signatures. In addition, radicals originating from reductants used for reducing biological systems may further complicate EPR spectra. Thus, it is essential to identify these reductant-based radical components in order to make assignments of biologically relevant protein-based radicals.

To determine how the radicals formed from dithionite and ascorbate can affect EPR spectra obtained during the oxygen chemistry of CcO, we systematically investigated the reactions of these reductants with CcO. We found that when ascorbate/Cyt *c* or dithionite are used to reduce bovine CcO prior to its reaction with oxygen, ascorbyl or $\text{SO}_2^{\bullet-}$ radicals are generated and may be trapped under conditions typically used to detect reaction intermediates. We report the first multi-frequency EPR characterization of these radicals.

2. MATERIALS AND METHODS

CcO was purified from bovine heart tissue via the method described by Yoshikawa [36]. Enzyme concentrations were determined by taking the optical absorption difference between fully-reduced CcO at 604 nm and the oxidized enzyme at 630 nm, using an extinction coefficient of $23.3 \text{ mM}^{-1} \text{ cm}^{-1}$ [37].

Natural abundance L-ascorbic acid was obtained from Fisher Scientific. Isotopically labeled L-ascorbic acid was obtained from Omicron Biochemicals. A 1 M stock solution was prepared in degassed 0.2 M sodium phosphate buffer with the pH adjusted to 7.4 with NaOH. A final concentration of 10 mM ascorbate was added to degassed 160 μM CcO and incubated with 0.5 μM Cyt *c* as the electron carrier.

Sodium dithionite was obtained from Sigma. A 1 M stock solution was prepared in degassed 0.2 M sodium phosphate buffer at pH 7.4. A final concentration of 6 mM dithionite was added to degassed 600 μM CcO without Cyt *c*.

Rapid freeze-quench (RFQ) is a novel technology used to trap transient radicals formed at room temperature for spectroscopic characterization at low temperature. RFQ samples were prepared using a custom-built device described by Lin *et al.* [38] and Egawa *et al.* [39]. Resting CcO, reduced with either ascorbate or dithionite in a gas-tight syringe, was mixed with oxygenated buffer in the RFQ device at room temperature. Samples were freeze-quenched on a timescale of 50 μsec to 6 msec. The frozen powder was packed into 4-mm O.D. (X-band) and 0.55 mm O.D. (D-band) precision-bore quartz EPR tubes immersed in liquid nitrogen. Hand-quenched samples were prepared in similar EPR tubes and frozen in liquid nitrogen on the timescale of minutes.

X-band (9 GHz) measurements were made on a Varian E-line spectrometer. A finger dewar was filled with liquid nitrogen and inserted into the EPR cavity preserved the sample at 77

K. Experimental conditions were: modulation amplitude, 3.2 G; microwave power, 1 mW; receiver gain, 2.5×10^4 ; microwave frequency, 9.107 GHz.

D-band (130 GHz) two pulse echo-detected spectra were obtained on a spectrometer described elsewhere [40, 41] using the following parameters: temperature, 7 K; repetition rate, 30 Hz; 30 averages per point; 90 degree pulse, 50 ns; time τ between pulses, 130 ns. For both X- and D-band spectra, the field was calibrated using Mn^{2+} doped in MgO [42].

Low temperature optical absorption was measured at 77 K on a modified Linkam THMS600 microscope stage system with a halogen light source.

3. RESULTS AND DISCUSSION

3.1 Single-turnover reaction of O_2 with ascorbate-reduced CcO

Fig. 1A shows optical absorption measurements of the reaction intermediates freeze-quenched at 50 to 6000 μs . The visible band exhibits a red-shift from 603 nm (50 μs) to 604 nm (150 μs), where it remains for ~ 1000 μs until shifting to 602 nm and finally to 599 nm (6000 μs). These shifts suggest progression from the fully reduced form, **R**, to the pulsed oxidized form of the enzyme, **O_H**. We find that the visible α -band of the pulsed oxidized form at 599 nm is red-shifted relative to the resting oxidized form at 597 nm, consistent with that reported in the literature [43–45]. The Soret band is saturated at the high enzyme concentration needed for EPR measurements and hence, is not shown.

In the corresponding X-band CW-EPR spectra for the RFQ samples, a narrow radical with a linewidth of 12 G overlies features of cupric Cu_A in all samples containing oxygen-containing intermediates (Fig. 1B). The isolated Cu_A^{2+} signal is seen clearly in the RFQ resting CcO sample and can be simulated with the parameters, $g_x = 1.99$, $g_y = 2.02$, and $g_z = 2.18$, consistent with reported values [46–48]. To demonstrate that the narrow radical was not formed due to passing a non-specific protein solution through the RFQ device, resting CcO and BSA were mixed with buffer and freeze-quenched in a similar fashion. As shown in Fig. 1B, the narrow radical was not generated in these controls.

High frequency EPR was used to determine the g -tensor of the radical. The radical has a powder pattern consistent with near axial symmetry and can be simulated with $g_x = 2.0068$, $g_y = 2.0066$, and $g_z = 2.0023$ (Fig. 2a). The isotropic g -value calculated from the trace of the g -tensor matches values reported for the ascorbyl radical at room temperature, $g = 2.0052$ [49].

Annealing experiments were performed, in which the 150 μs sample was warmed to either 180 K or 298 K then cooled back to 7 K for EPR measurements. The paramagnetic species diminishes in the D-band spectra after annealing (Fig. 2b and c). The stability of this radical species is consistent with its assignment as the ascorbyl radical. The same radical species could be observed by hand-quenching the CcO reaction during multiple-turnover conditions as shown in Fig. 3.

3.2 Multiple-turnover reaction of O₂ with ascorbate-reduced CcO

The stability of the radical observed in the single-turnover reaction suggested that large amounts of radical could be trapped in the multiple-turnover reaction. In particular, an increase in radical would be seen if its production depended on the CcO reaction—either in the reduction of CcO to produce **R** or in the oxygen chemistry to produce **P**, **F**, or **O_H**. To test this, we prepared hand-quenched samples in which the ascorbate-reduced CcO was mixed with O₂-saturated buffer and allowed to turn over for several minutes. An accumulation of the relatively stable radical species was detected by X-band CW-EPR. Fig. 3 shows the amount of unpaired spin in the multiple-turnover compared to the single-turnover samples. The signal-to-noise is higher in the multiple-turnover samples, which are frozen liquid, compared to the RFQ samples, which are frozen powders. The origin of the feature at ~3360 G in the RFQ sample spectrum is unknown at present but it does not interfere with our analysis of the signal at $g \sim 2.005$.

The principal g -values and stability of the radical species at room temperature were consistent with the ascorbyl radical rather than a protein radical arising from CcO. To confirm the assignment, experiments were carried out with isotopically-labeled ascorbate. L-[1-¹³C]-ascorbic acid was used to repeat the hand-quenched and RFQ experiments. A characteristic splitting of the radical signal due to its coupling to the ¹³C nucleus was observed in both samples (Fig. 4). Lowering the power to 0.3 mW and narrowing the field sweep to 50 G allowed detection of the radical species with minimal contributions from Cu_A²⁺. The linewidth of the natural abundance ascorbyl radical is remarkably narrow, measuring 9 G, yet subtle hyperfine coupling broaden the edges of the signal (arrows). The hyperfine structure can be fit by adding 4 protons, expected from the hydrogen-atoms attached at the C₄, C₅, and C₆ positions [49]. Adding the ¹³C nucleus with an isotropic hyperfine coupling of 6.54 in the C₁ position simulates the splitting seen in the isotopic data, similar to that reported for the ascorbyl radical (5.74 G) in the literature [49, 50]. The hyperfine structure was well-fit using isotropic values in both the frozen liquid and RFQ samples, suggesting that the hyperfine anisotropy is too small to significantly influence the quality of the simulations. The simulation parameters are shown in Table 1.

To determine if the ascorbyl radical originated from the one-electron reduction reaction of Cyt *c*, Cyt *c* and ascorbate were mixed in various ratios and frozen within 5 minutes. One study on the reduction of single-site metalloenzymes suggested that the dianion acts as the major reductant, forming the anionic ascorbate radical, which decays by a slow disproportionation reaction [22]. However, unlike the situation in the presence of CcO, we found that no radicals accumulate at 5 minutes in preparations of 1:1 or 20:1 Cyt *c* to reductant (Fig. 5a and b). Thus, upon the reduction of Cyt *c* alone, the ascorbyl radical initially formed must undergo further reactions which render it EPR silent. One possibility is that the disproportionation reaction is somehow accelerated under our experimental conditions. Another possibility is that the ascorbyl radical also acts as a reductant for Cyt *c*, producing the diamagnetic product dehydroascorbate in a reaction that is somehow prevented in the presence of CcO. Although the chemistry of the process is unclear at this stage, the end result is that ascorbate reduces Cyt *c* alone without the generation of an EPR-

detectable concentration of radicals, while reduction in the presence of CcO produces ascorbyl radical signals as early as 50 μ s which are stable on the timescale of minutes.

A major physiological function of cellular ascorbate is to quench cytotoxic oxoferryl species and protein-based radicals produced on hemoglobin during oxidative stress [51]. Similar reactions have been demonstrated in leghaemoglobin [52] and Mb [53]. Thus, a possible mechanism in CcO is that ascorbate reacts with an oxoferryl heme or a protein-based radical to generate the observed radical species. We stress that detection of the ascorbyl free radical neither proves nor disproves that CcO produces a protein-based radical in the oxygen reaction.

3.3 Previously reported radicals in the reaction of O₂ with ascorbate-reduced CcO

Wilson *et al.* studied bovine CcO using ascorbate as the reductant to create multiple turnover conditions and reported a $g \sim 2$ radical with a linewidth of 8 G that amounted to as much as 10% of the enzyme concentration [18]. They noted that when the enzyme is reduced with only ferrocycytochrome *c*, which was passed on a column to remove excess ascorbate, the magnitude of the radical signal decreased to $\sim 1\%$ of the enzyme concentration; however, they were unable to determine if the signal resulted from a residual ascorbyl radical or a protein-based radical.

In 2007, Wiertz *et al.* reported the presence of two radicals in the reaction of ascorbate-reduced *Paracoccus denitrificans aa₃* with oxygen [14]. The first species, present from 83 μ s to ~ 1 ms, was assigned as a tryptophanyl radical derivative of W272 (*P. denitrificans* CcO numbering, W236 in bovine) based on simulations of Q-band data. A radical with similar features in X-band spectra was reported in the *bo₃* oxidase from *E. coli* [17]. A second unidentified, narrow radical existed from 83 μ s through 6 msec. The authors suggested the source to be a “main-chain radical” but concluded that its identity and functionality could not be determined. This second “6 msec” species accounted for $\sim 0.5\%$ of the enzyme concentration and had g -values reported as 2.0022, 1.9965, 1.9994, which differ from those reported here for the ascorbyl radical.

For the bovine enzyme, when we subtract the ascorbyl radical and oxidized Cu_A from our data, we are unable to identify a protein-based radical. In addition, no protein-based radical was detected when CcO was reduced with dithionite, which was subsequently removed by a gel-filtration column. The data indicate that protein-based are not observed in bovine CcO when the fully-reduced enzyme is reacted with oxygen; in addition, ascorbate produces an ascorbyl radical in the oxygen reaction of bovine CcO. These data contrast the bacterial system, in which a tryptophanyl radical arises and no ascorbyl radical is reported even when the same concentration (10 mM) is used as a reductant [14].

Either the mechanism of the bovine enzyme or the reagents—specifically, the reductants and mediators—may cause these differences between the bovine and the bacterial systems. The observations of Wilson *et al.* along with ours are most consistent with differing mechanisms between mammalian and bacterial enzymes. This would suggest that electron donation from an amino acid is unnecessary in the robust mammalian chemistry but occurs in bacterial enzymes. This difference is interesting because in both systems, the metal centers of the

fully-reduced enzyme are capable of donating the four electrons necessary to complete the reduction. If radicals were to form in the reaction of bovine CcO, they could be quenched by ascorbate, obscuring the detection of any protein-based radicals. However, it should be noted that in the studies of the bacterial enzymes by de Vries and coworkers, phenazine ethosulfate (PES) was used as the redox mediator in lieu of the physiological reductant, Cyt *c*, used by us and Wilson *et al.*

3.4 Reaction of O₂ with dithionite-reduced CcO

To avoid the complication from the ascorbyl radical, we examined the CcO reaction by using dithionite as a reductant instead of ascorbate/Cyt *c*. In this series of experiments, the enzyme was exposed to a ten-fold excess of sodium dithionite prior to the reaction with O₂. RFQ samples were prepared as in the ascorbate experiments. X-band CW-EPR measurements show a narrow radical signal with a linewidth of 11 G in intermediates trapped from 50 μs through 6000 μs (Fig. 6a–e). When the narrow radical signal was subtracted from the data, no additional protein-based radicals were detected. The narrow radical disappeared upon annealing for 1 minute, leaving only the Cu_A²⁺ signal (Fig. 6f). In addition, hand-quench samples, prepared by mixing CcO reduced by dithionite and oxygen and quenched at one minute, showed only cupric Cu_A with no evidence of the narrow radical (data not shown).

Because previous studies suggested that dithionite spontaneously dissociates into breakdown products such as SO₂^{•-} [34, 35], the reductant was tested with both redox and non-redox active proteins. 115 μM CcO, BSA, or Mb was mixed with 50 mM dithionite and freeze-quenched at 50 μs. An intense signal with a linewidth of 11 G was seen in all three enzyme samples (Fig. 7a–c). These observations suggest that the radical arises spontaneously from dithionite and not from interactions with proteins.

To further characterize the radical species, 50 mM dithionite solutions were freeze-quenched with oxygenated or deoxygenated buffer and the samples were measured by D-band and X-band EPR. High-frequency spectra show perturbations of the *g_x* and *g_z* features in the oxygenated sample (Fig. 8b) compared to the deoxygenated sample (Fig. 8a), the origin of which are unknown at this time. However, both of these high-frequency oxygenated and deoxygenated spectra are well-fit by simulations with *g*-tensor values at 2.0089, 2.0052, and 2.0017 (Fig. 8–b, dotted lines). The isotropic *g*-value is 2.0053, which is in good agreement with the literature value of SO₂^{•-} [54–56]. This study is the first report of the *g*-tensor components of SO₂^{•-}. The radical in the samples of 50 mM dithionite with (Fig. 8e) and without oxygen (Fig. 8d) is identical to the radical produced when reacting CcO with oxygen (Fig. 8f). Using the well-defined *g*-values, the X-band data can be simulated using parameters in Table 2. The simulations, which are optimized to fit the slight asymmetry of the lineshape (Fig. 7, arrows), include two protons. These protons are likely H-bonding water molecules in the vicinity of SO₂^{•-}. In addition, the radical nearly disappears when annealed for 1 min. at 180 K (Fig. 8c), which is consistent with our annealing experiments performed at X-band. Unlike the ascorbyl radical, the SO₂^{•-} radical is unstable at room temperature under these experimental conditions.

4. CONCLUSIONS

This study reports *g*-tensors for the ascorbyl and $\text{SO}_2^{\cdot-}$ radicals and establishes that these radicals are formed and can be trapped in the course of CcO reacting with oxygen when ascorbate/Cyt *c* or dithionite is used as the reductant. In the frozen solution X-band spectra, the ascorbyl and $\text{SO}_2^{\cdot-}$ radicals both appear as relatively featureless single-line spectra with zero crossings at $g \sim 2.005$. The high-frequency spectra provide the resolution of peaks associated with the principal *g*-values and reveal that the ascorbyl radical has near-axial symmetry while the $\text{SO}_2^{\cdot-}$ is much more rhombic. This study demonstrates the utility of high-frequency EPR to distinguish between radicals that appear similar at X-band, thereby allowing for unequivocal identification.

Protein-based free radicals play an integral role in the chemistry of life processes and are often studied by EPR spectroscopy. In assessing protein-based radical mechanisms, byproducts from reductants must not be neglected as they may significantly impact the analyses. In bacterial CcO, catalytically relevant radicals are formed, but no such radical has yet been unequivocally identified in the mammalian enzyme. Identifying the origin of reducing equivalents used in catalysis is crucial to understanding the link between redox chemistry and proton translocation. However, the question of whether a radical arises in the catalytic reaction of mammalian CcO with O_2 remains unclear. Our studies show that ascorbate and dithionite—two universally used reductants in studies of redox proteins—can contribute to EPR spectra. Hence, great care must be taken in the interpretation of data to understand their roles in the reactions studied.

ACKNOWLEDGEMENTS

This work was funded by the National Institutes of Health (NIH) grants GM074982 to D.L.R. and GM075920 to G.J.G.

REFERENCES

- [1]. Buckel W and Golding BT, Radical enzymes in anaerobes, *Annu Rev Microbiol* 60 (2006) 27–49. [PubMed: 16704345]
- [2]. Marsh EN, A radical approach to enzyme catalysis, *Bioessays* 17 (1995) 431–41. [PubMed: 7786289]
- [3]. Stubbe J and van Der Donk WA, Protein Radicals in Enzyme Catalysis, *Chem Rev* 98 (1998) 705–762. [PubMed: 11848913]
- [4]. Sjoberg BM, Reichard P, Graslund A and Ehrenberg A, The tyrosine free radical in ribonucleotide reductase from *Escherichia coli*, *J Biol Chem* 253 (1978) 6863–5. [PubMed: 211133]
- [5]. Licht S, Gerfen GJ and Stubbe J, Thiyl radicals in ribonucleotide reductases, *Science* 271 (1996) 477–81. [PubMed: 8560260]
- [6]. Yonetani T, Schleyer H and Ehrenberg A, Studies on Cytochrome *c* Peroxidase: Electron Paramagnetic Resonance Absorptions of the Enzyme and Complex ES in Dissolved and Crystalline Forms, *J. Biol. Chem* 241 (1966) 3240–43. [PubMed: 4287913]
- [7]. Wittenberg BA, Kampa L, Wittenberg JB, Blumberg WE and Peisach J, The electronic structure of protoheme proteins. II. An electron paramagnetic resonance and optical study of cytochrome *c* peroxidase and its derivatives, *J Biol Chem* 243 (1968) 1863–70. [PubMed: 5646480]
- [8]. Goodin DB, Mauk AG and Smith M, Studies of the radical species in compound ES of cytochrome *c* peroxidase altered by site-directed mutagenesis, *Proc Natl Acad Sci U S A* 83 (1986) 1295–9. [PubMed: 3006043]

- [9]. Hiner AN, Martinez JI, Arnao MB, Acosta M, Turner DD, Lloyd Raven E and Rodriguez-Lopez JN, Detection of a tryptophan radical in the reaction of ascorbate peroxidase with hydrogen peroxide, *Eur J Biochem* 268 (2001) 3091–8. [PubMed: 11358529]
- [10]. Choinowski T, Blodig W, Winterhalter KH and Piontek K, The crystal structure of lignin peroxidase at 1.70 Å resolution reveals a hydroxy group on the Cβ of tryptophan 171: a novel radical site formed during the redox cycle, *J Mol Biol* 286 (1999) 809–27. [PubMed: 10024453]
- [11]. Pogni R, Baratto MC, Giansanti S, Teutloff C, Verdin J, Valderrama B, Lenzian F, Lubitz W, Vazquez-Duhalt R and Basosi R, Tryptophan-based radical in the catalytic mechanism of versatile peroxidase from *Bjerkandera adusta*, *Biochemistry* 44 (2005) 4267–74. [PubMed: 15766255]
- [12]. Pogni R, Baratto MC, Teutloff C, Giansanti S, Ruiz-Duenas FJ, Choinowski T, Piontek K, Martinez AT, Lenzian F and Basosi R, A tryptophan neutral radical in the oxidized state of versatile peroxidase from *Pleurotus eryngii*: a combined multifrequency EPR and density functional theory study, *J Biol Chem* 281 (2006) 9517–26. [PubMed: 16443605]
- [13]. Proshlyakov DA, Pressler MA, DeMaso C, Leykam JF, DeWitt DL, Babcock GT, Oxygen activation and reduction in respiration: involvement of redox-active Tyrosine 244, *Science* 290 (2000) 1588–90. [PubMed: 11090359]
- [14]. Wiertz FG, Richter OM, Ludwig B and de Vries S, Kinetic resolution of a tryptophan-radical intermediate in the reaction cycle of *Paracoccus denitrificans* cytochrome c oxidase, *J Biol Chem* 282 (2007) 31580–91. [PubMed: 17761680]
- [15]. Michel H, The mechanism of proton pumping by cytochrome c oxidase, *Proc Natl Acad Sci U S A* 95 (1998) 12819–24. [PubMed: 9788998]
- [16]. Svistunenko DA, Wilson MT and Cooper CE, Tryptophan or tyrosine? On the nature of the amino acid radical formed following hydrogen peroxide treatment of cytochrome c oxidase, *Biochim Biophys Acta* 1655 (2004) 372–80. [PubMed: 15100053]
- [17]. Wiertz FG, Richter OM, Cherepanov AV, MacMillan F, Ludwig B and de Vries S, An oxo-ferryl tryptophan radical catalytic intermediate in cytochrome c and quinol oxidases trapped by microsecond freeze-hyperquenching (MHQ). *FEBS Lett.* 575 (2004) 127–30. [PubMed: 15388346]
- [18]. Wilson MT, Jensen P, Aasa R, Malmstrom BG and Vanngard T, An investigation by e.p.r. and optical spectroscopy of cytochrome oxidase during turnover, *Biochem J* 203 (1982) 483–92. [PubMed: 6288002]
- [19]. Bergsten P, Amitai G, Kehrl J, Dhariwal KR, Klein HG and Levine M, Millimolar concentrations of ascorbic acid in purified human mononuclear leukocytes. Depletion and reaccumulation, *J Biol Chem* 265 (1990) 2584–7. [PubMed: 2303417]
- [20]. Shaskus J and Haake P, AScorbic Acid. 2. Nucleophilic Reactivity of Ascorbate Anion towards Acyl Carbon and Phosphorus, *J. Org. Chem* 49 (1984) 197–199.
- [21]. Pelizzetti E, Metnasti E and Pramauro E, Kinetics and mechanism of the oxidation of ascorbic acid by Tris(1,10-phenanthroline)iron(III) and its derivatives in aqueous acidic perchlorate media, *Inorganic Chemistry* 15 (1976) 2898–900.
- [22]. Al-Ayash AI and Wilson MT, The mechanism of reduction of single-site redox proteins by ascorbic acid, *Biochem J* 177 (1979) 641–8. [PubMed: 35158]
- [23]. Njus D, Jalukar V, Zu JA and Kelley PM, Concerted proton-electron transfer between ascorbic acid and cytochrome b561, *Am J Clin Nutr* 54 (1991) 1179S–1183S. [PubMed: 1660216]
- [24]. Bailey DM, George WO and Gutowski M, Theoretical studies of L-ascorbic acid (vitamin C) and selected oxidized, anionic and free-radical forms, *J. of Molecular Structure: THEOCHEM* 910 (2009) 61–68.
- [25]. Allen RN, Shukla MK, Reed D and Leszczynski J, Ab Initio Study of the Structural Properties of Ascorbic Acid (Vitamin C), *International Journal of Quantum Chemistry* 106 (2006) 2934–2943.
- [26]. Buettner GR and Chamulitrat W, The catalytic activity of iron in synovial fluid as monitored by the ascorbate free radical, *Free Radic Biol Med* 8 (1990) 55–6. [PubMed: 2157634]
- [27]. Buettner GR and Jurkiewicz BA, Ascorbate free radical as a marker of oxidative stress: an EPR study, *Free Radic Biol Med* 14 (1993) 49–55. [PubMed: 8384150]

- [28]. Buettner GR, The pecking order of free radicals and antioxidants: lipid peroxidation, alpha-tocopherol, and ascorbate, *Arch Biochem Biophys* 300 (1993) 535–43. [PubMed: 8434935]
- [29]. May JM, Li L, Qu ZC and Cobb CE, Mitochondrial recycling of ascorbic acid as a mechanism for regenerating cellular ascorbate, *Biofactors* 30 (2007) 35–48. [PubMed: 18198400]
- [30]. Li X, Cobb CE and May JM, Mitochondrial recycling of ascorbic acid from dehydroascorbic acid: dependence on the electron transport chain, *Arch Biochem Biophys* 403 (2002) 103–10. [PubMed: 12061807]
- [31]. Li X, Cobb CE, Hill KE, Burk RF and May JM, Mitochondrial uptake and recycling of ascorbic acid, *Arch Biochem Biophys* 387 (2001) 143–53. [PubMed: 11368176]
- [32]. Mayhew SG, The redox potential of dithionite and SO₂ from equilibrium reactions with flavodoxins, methyl viologen and hydrogen plus hydrogenase, *Eur J Biochem* 85 (1978) 535–47. [PubMed: 648533]
- [33]. Dunitz JD, Structure of Dithionite, *J. Amer. Chem. Soc* 78 (1956) 878–9.
- [34]. Hodgson WG, Neaves A and Parker CA, Detection of Free Radicals in Sodium Dithionite by Paramagnetic Resonance, *Nature* 178 (1956) 489.
- [35]. Janzen EG, Electron spin resonance study of the SO₂⁻ formation in the thermal decomposition of sodium dithionite, sodium and potassium metabisulfite, and sodium hydrogen sulfite., *J. of Physical Chemistry* 76 (1972) 157–162.
- [36]. Yoshikawa S, Choc MG, O'Toole MC and Caughey WS, An infrared study of CO binding to heart cytochrome c oxidase and hemoglobin A. Implications re O₂ reactions, *J Biol Chem* 252 (1977) 5498–508. [PubMed: 195952]
- [37]. Tsubaki M, Shinzawa K and Yoshikawa S, Effects of crystallization on the heme-carbon monoxide moiety of bovine heart cytochrome c oxidase carbonyl, *Biophys J* 63 (1992) 1564–71. [PubMed: 1336988]
- [38]. Lin Y, Gerfen GJ, Rousseau DL and Yeh SR, Ultrafast microfluidic mixer and freeze-quenching device, *Anal Chem* 75 (2003) 5381–6. [PubMed: 14710815]
- [39]. Egawa T, Durand JL, Hayden EY, Rousseau DL and Yeh SR, Design and evaluation of a passive alcove-based microfluidic mixer, *Anal Chem* 81 (2009) 1622–7. [PubMed: 19140669]
- [40]. Krymov V and Gerfen GJ, Analysis of the tuning and operation of reflection resonator EPR spectrometers, *J Magn Reson* 162 (2003) 466–78. [PubMed: 12810032]
- [41]. Rangelova K, Girotto S, Gerfen GJ, Yu S, Suarez J, Metlitsky L and Magliozzo RS, Radical sites in *Mycobacterium tuberculosis* KatG identified using electron paramagnetic resonance spectroscopy, the three-dimensional crystal structure, and electron transfer couplings, *J Biol Chem* 282 (2007) 6255–64. [PubMed: 17204474]
- [42]. Burghaus O, Rohrer M, Plato M and Mobius K, A novel high-field/high-frequency EPR and ENDOR spectrometer operating at 3 mm wavelength, *Meas Sci Tech* 3 (1992) 765–774.
- [43]. Antonini E, Brunori M, Greenwood C, Colosimo A, Wilson MT, Oxygen “pulsed” cytochrome c oxidase: functional properties and catalytic relevance, *PNAS* 74 (1977) 3128–32. [PubMed: 198771]
- [44]. Young LJ and Palmer G, Redox-cycled oxidase. One of the reaction products of reduced cytochrome c, cytochrome c oxidase, and dioxygen, *J Biol Chem* 261 (1986) 13031–3. [PubMed: 3020020]
- [45]. Okunuki K, Hagihara B, Sekuzu I and Horio T, Studies on the cytochrome system, *Proceedings of the International Symposium on Enzyme Chemistry* (1958) 264–272.
- [46]. Hoffman BM, Roberts JE, Swanson M, Speck SH and Margoliash E, Copper electron-nuclear double resonance of cytochrome c oxidase, *Proc Natl Acad Sci U S A* 77 (1980) 1452–6. [PubMed: 6246493]
- [47]. Beinert H, Griffiths DE, Wharton DC, Sands RH, Properties of the copper associated with cytochrome oxidase as studied by paramagnetic resonance spectroscopy, *J. Biol. Chem* 237 (1962) 2337–46. [PubMed: 13866633]
- [48]. Zhen Y, Schmidt B, Kang UG, Antholine W and Ferguson-Miller S, Mutants of the CuA site in cytochrome c oxidase of *Rhodobacter sphaeroides*: I. Spectral and functional properties, *Biochemistry* 41 (2002) 2288–97. [PubMed: 11841221]

- [49]. Laroff GP, Fessenden RW and Schuler RH, The electron spin resonance spectra of radical intermediates in the oxidation of ascorbic acid and related substances, *J Am Chem Soc* 94 (1972) 9062–73. [PubMed: 4345160]
- [50]. O'Malley PJ, Density Functional Calculations Modelling the Spin Density Distribution, Hyperfine Couplings, and Hydrogen Bonding Environment of the Ascorbate (Vitamin C) Free Radical, *J. Phys. Chem. B* 105 (2001) 11290–11293.
- [51]. Dunne J, Caron A, Menu P, Alayash AI, Buehler PW, Wilson MT, Silaghi-Dumitrescu R, Faivre B and Cooper CE, Ascorbate removes key precursors to oxidative damage by cell-free haemoglobin in vitro and in vivo, *Biochem J* 399 (2006) 513–24. [PubMed: 16848758]
- [52]. Moreau S, Puppo A and Davies MJ, The reactivity of ascorbate with different redox states of leghaemoglobin, *Phytochemistry* 39 (1995) 1281–6. [PubMed: 7669277]
- [53]. Giulivi C and Cadenas E, The reaction of ascorbic acid with different heme iron redox states of myoglobin. Antioxidant and prooxidant aspects, *FEBS Lett* 332 (1993) 287–90. [PubMed: 8405472]
- [54]. Streeter I, Wain AJ, Davis J and Compton RG, Cathodic reduction of bisulfite and sulfur dioxide in aqueous solutions on copper electrodes: an electrochemical ESR study, *J Phys Chem B* 109 (2005) 18500–6. [PubMed: 16853382]
- [55]. Zhu J, Petit K, Colson O, DeBolt S and Sevilla MD, Reactions of HS and S* with Molecular Oxygen, H₂S, HS⁻, and S²⁻: Formation of SO₂^{•-}, HSSH[•], HSS^{2•-}, and HSS^{•-}, *Journal of Physical Chemistry* 95 (1991) 3676–81.
- [56]. Ozawa T, Setaka M and Kwan T, ESR studies of the Sulfite Radical Anion, *Bull. Chem. Soc. Japan* 44 (1971) 3473–3474.




Fig. 1. Low-temperature optical absorption and X-band CW-EPR measurements of bovine CcO. A. 80 μM CcO completely reduced by 10 mM ascorbate and catalytic amounts of Cyt *c* under Ar then mixed with O_2 -saturated buffer and rapid freeze-quenched at time points (a) 50 μs , (b) 150 μs , (c) 300 μs , (d) 400 μs , (e) 540 μs , (f) 1 ms, (g) 6 ms. (h) is a sample of the resting enzyme. Optical absorption measurements were made at 83 K. Solutions were prepared in 200 mM NaPi buffer, pH 7.4, containing 0.2% w/v n-decyl- β -D-maltoside (DM). B. X-band EPR under the same conditions with quench times of (a) 50 μs , (b) 150 μs , (c) 300 μs , (d) 540 μs , (e) 1 ms, (f) 6 ms. (h) is a sample made by running resting enzyme through the RFQ device. The bottom two spectra are control samples of BSA run through the RFQ device (upper) and simply frozen in solution (lower). The conditions of EPR spectroscopy were: microwave power, 1 mW; microwave frequency, 9.1 GHz; modulation amplitude, 3.2 G; temperature 77 K.




Fig. 2. D-band EPR spectra of RFQ CcO prepared with ascorbate. (a) D-band EPR measurements of 80 μM CcO, completely reduced by 10 mM ascorbate and catalytic amounts of Cyt *c* under Ar, then mixed with O₂-saturated buffer and rapid freeze-quenched at 150 μs . A simulation with $g_x = 2.0068$, $g_y = 2.0066$, and $g_z = 2.0023$ in the dotted line. Sample from (a) annealed at the conditions: (b) 180 K for 1 min and (c) 298 K for 5 min. The conditions for the D-band Hahn echo-detected spectra were: microwave frequency, 129.998 GHz; repetition rate, 50 Hz; averages per point, 100; 90 degree pulse, 50 ns; time between pulses, 130 ns; temperature, 7K.




Fig. 3.

X-band EPR spectra of CcO under multiple-turnover and single-turnover. (a) 150 μ M CcO completely reduced by 10 mM ascorbate and catalytic amounts of Cyt *c* under Ar then mixed with O₂-saturated buffer and hand-quenched at 5 min. (b) 150 μ M CcO completely reduced by 10 mM. ascorbate and catalytic amounts of Cyt *c* under Ar then mixed with O₂-saturated buffer and rapid freeze-quenched at 150 μ s. EPR conditions as in Fig. 1.




Fig. 4.

X-band EPR spectra of CcO prepared with isotopically-labeled ascorbate. 150 μM CcO completely reduced by (a) 10 mM natural abundance ascorbate or (b) 10 mM L-[1-¹³C] ascorbate and catalytic amounts of Cyt *c* under Ar then mixed with O₂-saturated buffer and hand-quenched at 5 min. 150 μM CcO completely reduced by (c) 10 mM natural abundance ascorbate or (d) 10 mM L-[1-¹³C] ascorbate and catalytic amounts of Cyt *c* under Ar then mixed with O₂-saturated buffer and rapid freeze-quenched at 150 μs. Simulations are shown in the dotted lines with parameters shown in Table 1. EPR conditions were: frequency, 9.1 GHz; power, 0.3 mW; temperature, 77 K.




Fig. 5. X-band EPR spectra of Cyt *c* and ascorbate. (a) 5 mM Cyt *c* is reduced with 5 mM ascorbate and frozen at 77 K at 5 min. (b) 0.5 mM Cyt *c* is reduced with 10 mM ascorbate and frozen at 77 K at 5 min. (c) 150 μ M CcO and 0.5 μ M Cyt *c* are reduced with 10 mM ascorbate and frozen at 77 K at 5 min. The pH of all samples is 7.4. EPR conditions as in Fig. 1.




Fig. 6. X-band EPR spectra of CcO prepared with dithionite. 150 μ M CcO completely reduced by 1.5 mM dithionite under Ar then mixed with O₂-saturated buffer and rapid freeze-quenched at (a) 50, (b) 150, (c) 350, (d) 450, and (e) 6000 μ s. (f) 350 μ s sample annealed for 1 min. at 298 K. EPR conditions as in Fig. 1.




Fig. 7. X-band EPR spectra of various proteins prepared with dithionite. (a) 115 μM CcO and 50 mM $\text{Na}_2\text{S}_2\text{O}_4$ rapid freeze-quenched at 50 μs . (b) 115 μM bovine serum albumin and 50 mM $\text{Na}_2\text{S}_2\text{O}_4$ rapid freeze-quenched at 50 μs . X-band EPR conditions as in Fig. 1.




Fig. 8. D-band (top) and X-band (bottom) EPR spectra of dithionite with and without oxygen. (a) and (d) 50 mM $\text{Na}_2\text{S}_2\text{O}_4$ in degassed buffer rapid freeze-quenched at 50 μs . (b) and (e) 50 mM $\text{Na}_2\text{S}_2\text{O}_4$ with oxygenated buffer and 115 μM CcO rapid freeze-quenched at 50 μs . (c) Sample from (b) annealed at 180 K for 1 min. (f) 50 mM $\text{Na}_2\text{S}_2\text{O}_4$ in oxygenated buffer rapid freeze-quenched at 50 μs . D-band and X-band EPR conditions as in Figs. 2 and 1, respectively.

Table 1

Simulation parameters for the ascorbyl radical.

	L-[¹² C] Ascorbic Acid	L-[1- ¹³ C] Ascorbic Acid
g_x	2.0068	2.0068
g_y	2.0066	2.0066
g_z	2.0023	2.0023
a_{H4}	1.76 ^a	1.76 ^a
a_{H5}	0.07 ^a	0.07 ^a
$a_{H6,1}$	0.19 ^a	0.19 ^a
$a_{H6,2}$	0.19 ^a	0.19 ^a
a_{c1}	-	6.54 ^a

^aHyperfine couplings are given in units of Gauss.

Table 2

Simulation parameters for the sulfur dioxide anion radical.

<hr/> $\text{SO}_2^{\cdot-}$ <hr/>	
g_x	2.0089
g_y	2.0052
g_z	2.0017
A_{H2O}	0.5 ^a
	2.1 ^a
	1.1 ^a
A_{H2O}	0.5 ^a
	2.1 ^a
	1.1 ^a

^aHyperfine couplings are given in units of Gauss.