

Trends in VA Telerehabilitation Patients and Encounters Over Time and by Rurality

Diane C. Cowper-Ripley, PhD; Huanguang Jia, PhD; Xinping Wang, PhD; I. Maggie Freytes, PhD; Jennifer Hale-Gallardo, PhD; Gail Castaneda, PhD; Kimberly Findley, RN; and Sergio Romero, PhD

Telerehabilitation fills a need and helps ensure treatment adherence for rural and other veterans who find it difficult to access health care.

Diane Cowper-Ripley, Huanguang Jia, Maggie Freytes, and Sergio Romero are Research Health Scientists, and **Xinping Wang, Jennifer Hale-Gallardo, and Kimberly Findley** are Health Science Specialists, all at the Center of Innovation on Disability and Rehabilitation Research in Gainesville, Florida.
Correspondence: Huanguang Jia (huanguang.jia@va.gov)

Historically, the Veterans Health Administration (VHA) has excelled at improving veterans' access to health care and enhancing foundational services, such as prosthetics and other veteran-centric services, and this continues to be the VHA's top priority.¹ Travel distance and time are often barriers to accessing health care for many veterans.²⁻¹¹ For veterans with disabilities who must overcome additional physical, cognitive, and emotional obstacles to access vital rehabilitation services, these geographic obstacles are magnified. Further compounding the challenge is that rehabilitation therapies frequently require multiple encounters. Telerehabilitation is a promising solution for veterans in need of rehabilitation to regain optimal functioning. This alternative mode of service delivery can help veterans overcome geographic access barriers by delivering health care directly to veterans in their homes or nearby community-based outpatient clinics.^{12,13}

A growing body of evidence supports telerehabilitation. In a 2017 systematic review and meta-analysis, Cottrell and colleagues reviewed and analyzed data from 13 studies that met their inclusion criteria; specifically, their meta-analytic sample comprised adults aged ≥ 18 years presenting with any diagnosed primary musculoskeletal condition; treatment interventions via a real-time telerehabilitation medium, trials that had a comparison group with the same condition; provided clinical outcomes data, and included published randomized and nonrandomized controlled trials.¹⁴ Based on their aggregated results,

they concluded that real-time telerehabilitation was effective in improving physical function (standardized mean difference [SMD], 0.63; 95% CI, 0.92-2.33; I^2 , 93%), and reducing pain (SMD, 0.66; 95% CI, -0.27- .60; I^2 , 96%) in patients with any diagnosed primary musculoskeletal condition.¹⁴

Two other systematic reviews conducted by Pietrzak and colleagues and Agostini and colleagues also demonstrated the clinical effectiveness of telerehabilitation.^{15,16} Clinical effectiveness was defined as changes in health, functional status, and satisfaction with the telerehabilitation services delivered. The studies examined in the review included those that provided online self-management and education in addition to exercise via teleconferencing in real time.

Pietrzak and colleagues found that Internet-based osteoarthritis self-management interventions significantly improved 4 of 6 health status measures reviewed (ie, pain, fatigue, activity limitation, health distress, disability, and self-reported global health).¹⁵ User acceptance and satisfaction were high ($\geq 70\%$ satisfied) in all studies meeting the inclusion criteria.

Agostini and colleagues found that telerehabilitation was more effective than other modes of delivering rehabilitation to regain motor function in cardiac (SMD, 0.24; 95% CI, 0.04-0.43) and total knee arthroplasty (Timed Up and Go test: SMD, -5.17; 95% CI, -9.79- -0.55) patients.¹⁶ Some evidence from VHA and non-VHA studies also support the use of telerehabilitation to reduce health care costs,¹⁷⁻¹⁹ improve treatment adherence,^{12,20} and enhance patient physical, cognitive and

TABLE 1 Number of Unique Patients Seen and Total Number of Encounters by Clinic

Clinics	FY 2012		FY 2013		FY 2014		FY 2015		FY 2016		FY 2017	
	P	E	P	E	P	E	P	E	P	E	P	E
Traumatic brain injury-individual	700	996	1,036	1,569	1,562	2,196	1,700	2,424	1,686	2,926	1,635	2,465
PM&R Service	918	1,109	655	903	709	983	766	1,141	591	807	503	685
Physical therapy	1,676	3,016	3,446	5,615	6,503	9,293	8,218	11,969	8,106	11,063	9,136	11,834
Occupational therapy	147	292	966	1,810	1,262	2,077	1,775	2,638	2,457	3,584	2,495	3,538
PM&R amputation	492	664	668	965	821	1,210	973	1,364	1,109	1,673	1,116	1,669
Kinesiotherapy	23	23	183	259	267	353	357	443	496	551	563	624
Amputation	439	540	598	891	751	1,132	986	1,518	1,005	1,447	851	1,339
PM&R Service AT	2	3	4	6	3	5	10	16	13	18	20	25
Total	4,397	6,643	7,556	12,018	11,878	17,249	14,785	21,513	15,463	22,069	16,319	22,179

Abbreviations: AT, assistive technology; E, encounter; FY, fiscal year; P, patient; PM&R, physical medicine and rehabilitation.

mobility function, as well as patient satisfaction and health-related quality of life.^{13,21-24}

Since the first recorded use of telehealth in 1959, the application of technology to deliver health care, including rehabilitation services, has increased exponentially.¹⁴ In fiscal year (FY) 2017 alone, the VA provided > 2 million episodes of care for > 700,000 veterans using telehealth services.²⁵

Although the process for accessing telerehabilitation may vary throughout the VA, typically a few common factors make a veteran eligible for this mode of rehabilitation care delivery: Veterans must meet criteria for a specific program (eg, amputation, occupational therapy, and physical therapy) and receive VA care from a VA medical facility or clinic that offers telehealth services. Care providers must believe that the veteran would benefit from telerehabilitation (eg, limited mobility and long-distance travel to the facility) and that they would be able to receive an appropriate consult. The veteran must meet the following requirements: (1) willingness to consent to a visit via telehealth; (2) access to required equipment/e-mail; and (3) a caregiver to assist if they are unable to complete a visit independently.

In this article, we provide an overview of the growth of telerehabilitation in the VHA. Data are presented for specific telerehabilitation programs over time and by rurality.

METHODS

The VHA Support Service Center works with VHA program offices and field users to provide field-focused business, clinical, and special topic reports. An online portal provides access to these customizable reports organized as data cubes, which represent data dimensions (ie, clinic type) and measures (ie, number of unique patients). For this study, we used the Connected Care, Telehealth, Call Centers Clinical Video Telehealth/Store and Forward Telehealth data cube clinical stop codes to identify the numbers of telerehabilitation veteran users and encounters across time. The following telerehabilitation clinic-stop codes were selected: 197 (polytrauma/traumatic brain injury [TBI]—individuals), 201 (Physical Medicine and Rehabilitation [PM&R] Service), 205 (physical therapy), 206 (occupational therapy), 211 (PM&R amputation clinic), 418 (amputation clinic), 214 (kinesiotherapy), and 240 (PM&R assistive technology clinic). Data for total unique patients served and the total number of encounters were extracted at the national level and by rurality from FY 2012 to FY 2017, providing the past 5 years of VHA telerehabilitation data.

It is important to note that in FY 2015, the VHA changed its definition of rurality to a rural-urban commuting areas (RUCA)-based system (www.ruralhealth.va.gov/rural

TABLE 2 Number of Unique Patients Treated in Telerehabilitation Clinics by Rurality

Clinics	FY 2012		FY 2013		FY 2014		FY 2015		FY 2016		FY 2017	
	Rural	Urban	Rural	Urban	Rural	Urban	Rural	Urban	Rural	Urban	Rural	Urban
Traumatic brain injury-individual	235	434	344	671	512	995	653	1,011	584	1,072	682	936
PM&R Service	138	770	155	496	100	603	162	604	92	499	94	409
Physical therapy	828	838	2,062	1,364	3,644	2,835	4,812	3,406	4,652	3,453	5,280	3,856
Occupational therapy	73	73	454	509	604	649	1,048	727	1,418	1,038	1,397	1,098
PM&R amputation	283	207	382	276	435	385	475	498	573	536	623	492
Kinesiotherapy	14	9	87	87	109	156	187	170	260	236	303	260
Amputation	176	260	226	364	285	443	380	606	433	571	369	480
PM&R Service AT	0	2	1	3	0	3	4	6	3	10	10	10
Total	1,747	2,593	3,711	3,770	5,689	6,069	7,721	7,028	8,015	7,415	8,758	7,541

Abbreviations: AT, assistive technology; E, encounter; FY, fiscal year; P, patient; PM&R, physical medicine and rehabilitation.

-definition.asp). Prior to FY 2015, the VHA used the US Census Bureau (CB) urbanized area definitions. According to CB, an urbanized area contains a central city and surrounding area that totals > 50,000 in population. It also includes places outside of urbanized areas with populations > 2,500. Rural areas are defined as all other areas. VHA added a third category, highly rural, which is defined as areas that had < 7 people per square mile. In the RUCA system, each census tract defined by the CB is given a score. The VHA definitions are as follows:

- Urban (U)—census tracts with RUCA scores of 1.0 or 1.1. These tracts are determined by the CB as being in an urban core and having the majority of their workers commute within that same core (1.0). If 30% to 49% commute to an even larger urban core, then the code is 1.1;
- Rural (R)—all tracts not receiving scores in the urban or highly rural tiers; and
- Highly rural (H)—tracts with a RUCA score of 10.0. These are the most remote occupied land areas. Less than 10% of workers travel to CB-defined urbanized areas or urban clusters.

In addition, VHA recently added an “I” category to complement “U,” “R,” and “H.” The “I” value is assigned to veterans living on the US insular islands (ie, territories): Guam, American Samoa, Northern Marianas, and

US Virgin Islands. For the analysis by rurality in this study, we excluded veterans living in the insular islands and those of unknown rurality (< 1.0% of patients and encounters). Further, because the numbers of highly rural veterans were relatively small (< 2% of patients and encounters), the rural and highly rural categories were combined and compared with urban-dwelling veterans.

RESULTS

Overall, the workload for telerehabilitation nearly quadrupled over the 5-year period (Table 1 and Figure 1). In FY 2012, there were 4,397 unique individuals receiving telerehabilitation in the selected telerehabilitation clinics. By FY 2017, this number had grown to 16,319 veterans. Similar increases were seen for total encounters, growing from 6,643 in FY 2012 to 22,179 in FY 2017 (Figure 2). The rate of the increase for the number of unique patients seen and telerehabilitation encounter totals across years were higher from FY 2012 to FY 2015 than from FY 2015 to FY 2017.

Interesting trends were seen by clinic type. Some clinics increased substantially, whereas others showed only moderate increases, and in 1 case (PM&R Service), a decrease. For example, there is significant growth in the number of patients and encounters involving physical therapy

through telerehabilitation. This telerehabilitation clinic increased its workload from 1,676 patients with 3,016 encounters in FY 2012 to 9,136 patients with 11,834 encounters in FY 2017, accounting for 62.6% of total growth in patients and 56.8% of total growth in encounters.

Other clinics showing substantial growth over time included occupational therapy and polytrauma/TBI-individual secondary evaluation. Kinesiotherapy telerehabilitation was almost nonexistent in the VHA during FY 2012, with only 23 patients having 23 encounters. By FY 2017, there were 563 patients with 624 kinesiotherapy telerehabilitation encounters, equating to staggering increases in 5 years: 2,348% for patients and 2,613% for encounters. Similarly, the Physical Medicine and Rehabilitation Assistive Technology clinics had very low numbers in FY 2012 (patients, 2; encounters, 3) and increased over time; albeit, at a slow rate.

Trends by Rurality


Trends by rural location of patients and encounters must be interpreted with caution because of the changing rural definition between FY 2014 and FY 2015 (Tables 2 and 3; Figures 3 and 4). Nevertheless, the number of veterans seen and encounters performed via telerehabilitation increased in both urban and rural settings during the time under investigation. Under both the legacy and RUCA definitions of rural, the percentage increase was greater for rural veterans than that for urban veterans.

The increased total number of patients seen between FY 2012 and FY 2014 (old definition) was 225% for rural veterans vs 134% for urban veterans. Between FY 2015 and FY 2017 (new definition), the increase was lower for both groups (rural, 13.4%; urban, 7.3%), but rural veterans still increased at a higher rate than did urban dwellers.

DISCUSSION


Our primary aim was to provide data on the growth of telerehabilitation in the VHA over the past 5 years. Our secondary aim was to examine growth in the use of telerehabilitation by rurality. Specifically, we provided an overview of telerehabilitation growth in terms of unique patients and overall encounters in the VHA by rural-

FIGURE 1 Unique Patients Seen Via Telerehabilitation by Clinic



Abbreviations: AT, assistive technology; FY, fiscal year; P, patients; PM&R, physical medicine and rehabilitation; TBI, traumatic brain injury.

FIGURE 2 Number of Telerehabilitation Encounters by Clinic



Abbreviations: AT, assistive technology; E, encounters; FY, fiscal year; PM&R, physical medicine and rehabilitation; TBI, traumatic brain injury.

ity from FY 2012 to FY 2014 and FY 2015 to FY 2017 using the following programs: Polytrauma/TBI, PM&R Service, physical therapy, occupational therapy, PM&R amputation clinic, amputation clinic, kinesiotherapy, and PM&R assistive technology clinic. Our findings demonstrated a noteworthy increase in telerehabilitation encounters and unique patients over time for these programs. These findings were consistent with

TABLE 3 Number of Encounters in Telerehabilitation Clinics by Rurality

Clinics	FY 2012		FY 2013		FY 2014		FY 2015		FY 2016		FY 2017	
	Rural	Urban	Rural	Urban	Rural	Urban	Rural	Urban	Rural	Urban	Rural	Urban
Traumatic brain injury-individual	348	601	570	977	727	1385	915	1431	934	1,930	954	1,481
PM&R Service	234	835	251	627	131	842	257	884	160	647	137	548
Physical therapy	1,836	1,129	3,717	1,861	5,507	3,754	7,191	4,778	6,564	4,498	7,090	4,744
Occupational therapy	214	77	756	1,051	968	1,092	1,481	1,157	2,024	1,559	1,905	1,633
PM&R amputation	365	297	541	410	608	601	667	697	895	778	909	756
Kinesiotherapy	14	9	104	155	123	226	227	216	281	270	332	292
Amputation	201	336	322	558	416	668	581	937	588	857	591	744
PM&R Service AT	0	3	1	5	0	5	4	12	3	15	14	11
Total	3,212	3,287	6,262	5,644	8,480	8,573	11,323	10,112	11,449	10,554	11,932	10,209

Abbreviations: AT, assistive technology; E, encounter; FY, fiscal year; P, patient; PM&R, Physical Medicine and Rehabilitation.

FIGURE 3 Number of Unique Patients by Rurality





FIGURE 4 Number of Encounters by Rurality



the overall trend of continued growth and expansion of telehealth within the VHA.

Our findings reveal an upward trend in the total number of rural encounters and rural unique patients despite the change in the VA's definition of rurality in FY 2015. To our knowledge, urban and rural use of telerehabilitation has not been examined previously. Under both definitions of rural-

ity, encounters and unique patients show an important increase over time, and by year-end 2017, more than half of all patients and encounters were attributed to rural patients (53.7% and 53.9%, respectively). Indeed, the upward trend may have been more pronounced if the rural definition had not changed in FY 2015. Our early VHA stroke patients study on the difference between

rural-urban patients and taxonomies showed that the RUCA definition was more likely to reduce the number of rural patients by 8.5% than the early definition used by the VHA.²⁶

It is notable that although the use of tele-delivery of rehabilitation has continually increased, the rate of this increase was steeper from FY 2012 to FY 2014 than FY 2015 to FY 2017. For the programs under consideration in this study, the total number of rural patients/encounters increased throughout the observed periods. However, urban patients and encounters increased through FY 2016 and experienced a slight decrease in FY 2017.

The appearance of a slower rate of increase may be due to a rapid initial rate of increase through early adopters and “crossing the diffusion chasm,” a well-documented process of slower diffusion between the time of invention to penetration that often characterizes the spread of successful telehealth innovations.²⁷ Integrating technology into care delivery innovation requires the integration of technical, clinical, and administrative processes and can take time to scale successfully.²⁸

With an emphasis on increasing access to rehabilitation services, the VHA can expect to see a continuing increase in both the number and the percentage of telerehabilitation rural patients and encounters. The VHA has several telerehabilitation initiatives underway through the VHA's Physical Medicine and Rehabilitation Telerehabilitation Enterprise Wide Initiative (TREWI) and Rural Veterans Telerehabilitation Initiative. These projects demonstrate the feasibility of this delivery approach and facilitate integration of this modality in clinical workflows. However, to sustain these efforts, facilities will need more infrastructure and personnel resources dedicated to the delivery of services.

In an ongoing evaluation of the TREWI, several factors seem to influence the uptake of the VHA Office of Rural Health TREWI programs. These factors are the presence or absence of a local site champion; the quality of hospital leadership support; the quality of past relationships between telerehabilitation sending sites and receiving sites; barriers to getting a telehealth service agreement in place; the availability of space; administrative know-how on setting up clinics appropriately; time involved to bring on staff;

contracting issues; equipment availability and installation; cultural issues in embracing technologic innovation; training burden; hassle factors; and limited funds. Although early adopters may be able to negotiate and push through many of the barriers associated with the diffusion of telerehabilitation, the numerous barriers may slow its larger systemwide diffusion.

Telerehabilitation is a promising mode to deliver care to rural veterans who otherwise may not have access to this type of specialty care. Therefore, the identification of elements that foster telerehabilitation growth in future investigations can assist policy makers and key stakeholders in optimally leveraging program resources for maximal productivity. Future studies investigating the drivers of increases in telerehabilitation growth by rurality are warranted. Furthermore, more research is needed to examine telerehabilitation growth quality of care outcomes (eg, patient and provider satisfaction) to ensure that care is not only timely and accessible, but of high quality.

CONCLUSION

Disparities between rural and urban veterans compel a mode of expanding delivery of care. The VHA has embraced the use of telehealth modalities to extend its reach of rehabilitation services to veterans with disability and rehabilitation needs. Growth in telerehabilitation rural patient encounters increases access to rehabilitative care, reduces patient and caregiver travel burden, and helps ensure treatment adherence. Telerehabilitation utilization (unique patients and total encounters) is growing more rapidly for rural veterans than for their urban counterparts. Overall, telerehabilitation is filling a gap for rural veterans, as well as veterans in general with challenges in accessibility to health care. In order to make full use of the telerehabilitation services across its health care system, VA health care facilities may need to expand their effort in telerehabilitation dissemination and education among providers and veterans, particularly among providers who are less familiar with telerehabilitation services and among veterans who live in rural or highly rural areas and need special rehabilitation care.

Author disclosures

The authors report no actual or potential conflicts of interest with regard to this article.

Disclaimer

The opinions expressed herein are those of the authors and do not necessarily reflect those of *Federal Practitioner*, Frontline Medical Communications Inc., the US Government, or any of its agencies.

References

- Shane L. What's in the VA secretary's 10-point plan to reform his department? <https://rebootcamp.militarytimes.com/news/pentagon-congress/2017/02/28/what-s-in-the-va-secretary-s-10-point-plan-to-reform-his-department>. Published February 28, 2017. Accessed November 21, 2018.
- Burgess JF, DeFiore DA. The effect of distance to a VA facility on the choice and level of utilization of VA outpatient services. *Soc Science Med*. 1994;39(1):95-104.
- LaVela SL, Smith B, Weaver FM, Miskevics SA. Geographical proximity and health care utilization in veterans with SCI&D in the USA. *Soc Science Med*. 2004;59:2387-2399.
- Piette JD, Moos RH. The influence of distance on ambulatory care use, death, and readmission following a myocardial infarction. *Health Serv Res*. 1996;31(5):573-591.
- Schmitt SK, Phibbs CS, Piette JD. The influence of distance on utilization of outpatient mental health aftercare following inpatient substance abuse treatment. *Addictive Behav*. 2003;28(6):1183-1192.
- Fortney JC, Booth BM, Blow FC, Bunn JY. The effects of travel barriers and age on the utilization of alcoholism treatment aftercare. *Am J Drug Alcohol Abuse*. 1995;21(3):391-406.
- McCarthy JF, Blow FC, Valenstein M, et al. Veterans Affairs Health System and mental health treatment retention among patients with serious mental illness: evaluating accessibility and availability barriers. *Health Serv Res*. 2007;42(3):1042-1060.
- Mooney C, Zwanziger J, Phibbs CS, Schmitt S. Is travel distance a barrier to veterans' use of VA hospitals for medical surgical care? *Soc Sci Med*. 2000;50(12):1743-1755.
- Friedman SA, Frayne SM, Berg E, et al. Travel time and attrition from VHA care among women veterans: how far is too far? *Med Care*. 2015;53(4)(suppl 1):S15-S22.
- Buzza C, Ono SS, Turvey C, et al. Distance is relative: unpacking a principal barrier in rural healthcare. *J Gen Intern Med*. 2011;26(suppl 2):648-654.
- Goins RT, Williams KA, Carter MW, Spencer SM, Solovieva T. Perceived barriers to health care access among rural older adults: a qualitative study. *J Rural Health*. 2005;21(3):206-213.
- Kairy D, Lehoux P, Vincent C, Visintin M. A systematic review of clinical outcomes, clinical process, healthcare utilization and costs associated with telerehabilitation. *Disabil Rehabil*. 2009;31(6):427-447.
- McCue M, Fairman A, Pramuka M. Enhancing quality of life through telerehabilitation. *Phys Med Rehabil Clin N Am*. 2010;21(1):195-205.
- Cottrell MA, Galea OA, O'Leary SP, Hill AJ, Russell TG. Real-time telerehabilitation for the treatment of musculoskeletal conditions is effective and comparable to standard practice: a systematic review and meta-analysis. *Clin Rehabil*. 2017;31(5):625-638.
- Pietrzak E, Cotea C, Pullman S, Nasveld P. Self-management and rehabilitation in osteoarthritis: is there a place for internet-based interventions? *Telemed J E Health*. 2013;19(10):800-805.
- Agostini M, Moja L, Banzi R, et al. Telerehabilitation and recovery of motor function: a systematic review and meta-analysis. *J Telemed Telecare*. 2015;21(4):202-213.
- Kortke H, Stromeyer H, Zittermann A, et al. New East-Westfalian Postoperative Therapy Concept: A telemedicine guide for the study of ambulatory rehabilitation of patients after cardiac surgery. *Telemed J E-Health*. 2006;12(4):475-483.
- Tousignant M, Boissy P, Corriveau H, Moffet H. In home telerehabilitation for older adults after discharge from an acute hospital or rehabilitation unit: A proof-of- concept study and costs estimation. *Disabil Rehabil Assist Technol*. 2006;1(4):209-216.
- Sanford JA, Griffiths PC, Richardson P, et al. The effects of in-home rehabilitation on task self-efficacy in mobility-impaired adults: a randomized clinical trial. *J Am Geriatr Soc*. 2006;54(11):1641-1648.
- Nakamura K, Takano T, Akao C. The effectiveness of videophones in home healthcare for the elderly. *Med Care*. 1999;37(2):117-125.
- Levy CE, Silverman E, Jia H, Geiss M, Omura D. Effects of physical therapy delivery via home video telerehabilitation on functional and health-related quality of life outcomes. *J Rehabil Res Dev*. 2015;52(3):361-370.
- Guilfoyle C, Wootton R, Hassall S, et al. User satisfaction with allied health services delivered to residential facilities via videoconferencing. *J Telemed Telecare*. 2003;9(1):S52-S54.
- Mair F, Whitten P. Systematic review of studies of patient satisfaction with telemedicine. *BMJ*. 2000;320(7248):1517-1520.
- Williams T L, May C R, Esmail A. Limitations of patient satisfaction studies in telehealthcare: a systematic review of the literature. *Telemed J E-Health*. 2001;7(4):293-316.
- US Department of Veterans Affairs, Office of Telehealth Services. <http://vaww.telehealth.va.gov/quality/data/index.asp>. Accessed June 1, 2018. [Nonpublic document; source not verified.]
- Jia H, Cowper D, Tang Y, et al. Post-acute stroke rehabilitation utilization: Are there difference between rural-urban patients and taxonomies? *J Rural Health*. 2012;28(3):242-247.
- Cho S, Mathiassen L, Gallivan M. Crossing the chasm: from adoption to diffusion of a telehealth innovation. In: León G, Bernardos AM, Casar JR, Kautz K, De Gross JI, eds. *Open IT-Based Innovation: Moving Towards Cooperative IT Transfer and Knowledge Diffusion*. Boston, MA: Springer; 2008.
- Broderick A, Lindeman D. Scaling telehealth programs: lessons from early adopters. <https://www.commonwealthfund.org/publications/case-study/2013/jan/scaling-telehealth-programs-lessons-early-adopters>. Published January 2013. Accessed June 1, 2018.