



Cochrane
Library

Cochrane Database of Systematic Reviews

Prophylactic chemotherapy for hydatidiform mole to prevent gestational trophoblastic neoplasia (Review)

Fu J, Fang F, Xie L, Chen H, He F, Wu T, Hu L, Lawrie TA

Fu J, Fang F, Xie L, Chen H, He F, Wu T, Hu L, Lawrie TA.

Prophylactic chemotherapy for hydatidiform mole to prevent gestational trophoblastic neoplasia.

Cochrane Database of Systematic Reviews 2012, Issue 10. Art. No.: CD007289.

DOI: 10.1002/14651858.CD007289.pub2.

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON	4
BACKGROUND	7
OBJECTIVES	8
METHODS	8
RESULTS	10
Figure 1.	11
Figure 2.	13
DISCUSSION	15
AUTHORS' CONCLUSIONS	18
ACKNOWLEDGEMENTS	18
REFERENCES	19
CHARACTERISTICS OF STUDIES	22
DATA AND ANALYSES	29
Analysis 1.1. Comparison 1 Prophylactic chemotherapy versus no prophylactic chemotherapy, Outcome 1 Incidence of GTN (overall).	30
Analysis 1.2. Comparison 1 Prophylactic chemotherapy versus no prophylactic chemotherapy, Outcome 2 Incidence of GTN (high-risk HM only).	31
Analysis 1.3. Comparison 1 Prophylactic chemotherapy versus no prophylactic chemotherapy, Outcome 3 Time to GTN diagnosis.	32
Analysis 1.4. Comparison 1 Prophylactic chemotherapy versus no prophylactic chemotherapy, Outcome 4 Number of courses of chemotherapy to cure.	33
Analysis 1.5. Comparison 1 Prophylactic chemotherapy versus no prophylactic chemotherapy, Outcome 5 Mortality rate.	33
ADDITIONAL TABLES	33
APPENDICES	39
WHAT'S NEW	40
HISTORY	40
CONTRIBUTIONS OF AUTHORS	41
DECLARATIONS OF INTEREST	41
SOURCES OF SUPPORT	41
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	41
INDEX TERMS	42

[Intervention Review]

Prophylactic chemotherapy for hydatidiform mole to prevent gestational trophoblastic neoplasia

Jing Fu¹, Fang Fang², Lingxia Xie², Hengxi Chen¹, Fan He³, Taixiang Wu⁴, Lina Hu⁵, Theresa A Lawrie⁶

¹Department of Obstetrics and Gynecology, West China Second University Hospital, Sichuan University, Chengdu, China. ²Department of Obstetrics and Gynecology, West China Second University Hospital, West China Women's and Children's Hospital, Chengdu, China. ³West China Hospital, Sichuan University, Chengdu, China. ⁴Chinese Clinical Trial Registry, Chinese Ethics Committee of Registering Clinical Trials, West China Hospital, Sichuan University, Chengdu, China. ⁵The Obstetrics and Gynecology Department, The Second Affiliated Hospital of Chongqing Medical University, Chongqing, China. ⁶Cochrane Gynaecological, Neuro-oncology and Orphan Cancer Group, Royal United Hospital, Bath, UK

Contact address: Lina Hu, The Obstetrics and Gynecology Department, The Second Affiliated Hospital of Chongqing Medical University, 76 Linjiang Rd, Chongqing, 400010, China. cqhulina@126.com.

Editorial group: Cochrane Gynaecological, Neuro-oncology and Orphan Cancer Group.

Publication status and date: Edited (no change to conclusions), published in Issue 9, 2016.

Review content assessed as up-to-date: 10 August 2012.

Citation: Fu J, Fang F, Xie L, Chen H, He F, Wu T, Hu L, Lawrie TA. Prophylactic chemotherapy for hydatidiform mole to prevent gestational trophoblastic neoplasia. *Cochrane Database of Systematic Reviews* 2012, Issue 10. Art. No.: CD007289. DOI: 10.1002/14651858.CD007289.pub2.

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Hydatidiform mole (HM), also called a molar pregnancy, is characterised by an overgrowth of foetal chorionic tissue within the uterus. HMs may be partial (PM) or complete (CM) depending on their gross appearance, histopathology and karyotype. PMs usually have a triploid karyotype, derived from maternal and paternal origins, whereas CMs are diploid and have paternal origins only. Most women with HM can be cured by evacuation of retained products of conception (ERPC) and their fertility preserved. However, in some women the growth persists and develops into gestational trophoblastic neoplasia (GTN), a malignant form of the disease that requires treatment with chemotherapy. CMs have a higher rate of malignant transformation than PMs. It may be possible to reduce the risk of GTN in women with HM by administering prophylactic chemotherapy (P-Chem). However, P-Chem given before or after evacuation of HM to prevent malignant sequelae remains controversial, as the risks and benefits of this practice are unclear.

Objectives

To systematically review the evidence for the effectiveness and safety of P-Chem to prevent GTN in women with a molar pregnancy.

Search methods

We performed electronic searches in the Cochrane Gynaecological Cancer Specialised Register, the Cochrane Central Register of Controlled Trials (CENTRAL, Issue 2, 2012), MEDLINE (1946 to February week 4, 2012) and EMBASE (1980 to week 9, 2012). The search strategy was developed using free text and medical subject headings (MESH). We handsearched reference lists of relevant literature to identify additional studies.

Selection criteria

We included randomised controlled trials (RCTs) of P-Chem for HM.

Data collection and analysis

Two review authors independently assessed studies for inclusion in the review and extracted data using a specifically designed data collection form. Meta-analyses were performed by pooling data from individual trials using RevMan 5.1 software.

Main results

We included three RCTs with a combined total of 613 participants. One study compared prophylactic dactinomycin to no prophylaxis (60 participants); the other two studies compared prophylactic methotrexate to no prophylaxis (420 and 133 participants). All participants were diagnosed with CMs. We considered the latter two studies to be of poor methodological quality.

P-Chem reduced the risk of GTN occurring in women following a CM (3 studies, 550 participants; RR 0.37; 95% confidence interval (CI) 0.24 to 0.57; $I^2 = 0\%$; $P < 0.00001$). However, owing to the poor quality of two of the included studies, we performed sensitivity analyses excluding these two studies. This left only one small study of high-risk women to contribute data for this primary outcome (59 participants; RR 0.28; 95% CI 0.10 to 0.73; $P = 0.01$), therefore we consider this evidence to be of a low quality.

The time to diagnosis was longer in the P-Chem group than the control group (2 studies, 33 participants; mean difference (MD) 28.72; 95% CI 13.19 to 44.24; $P = 0.0003$) and the P-Chem group required more courses to cure subsequent GTN (1 poor-quality study, 14 participants; MD 1.10; 95% CI 0.52 to 1.68; $P = 0.0002$). We consider this evidence to be of a low to very low quality for similar reasons to those listed above.

There were insufficient data to perform meta-analyses for toxicity, overall survival, drug resistance and reproductive outcomes.

Authors' conclusions

P-Chem may reduce the risk of progression to GTN in women with CMs who are at a high risk of malignant transformation; however, current evidence in favour of P-Chem is limited by the poor methodological quality and small size of the included studies. As P-Chem may increase drug resistance, delay treatment of GTN and expose women unnecessarily to toxic side effects, this practice cannot currently be recommended.

PLAIN LANGUAGE SUMMARY

Prophylactic (preventive) chemotherapy for hydatidiform mole (molar pregnancy) to prevent cancerous growth later

A molar pregnancy (hydatidiform mole) develops following an abnormal process of conception, whereby placental tissue overgrows inside the womb (uterus). Molar pregnancies are classified as complete (CM) or partial (PM) based on their appearance (gross and microscopic), and their chromosome pattern. Moles are usually suspected at the early pregnancy scan and women often present with bleeding, similar to a miscarriage. The molar tissue is removed by evacuation of retained products of conception (ERPC), also known as dilatation and curettage (D&C) and women generally make a full recovery. However, some women go on to develop a cancer in the womb (about 1 in every 5 women with a CM and 1 in 200 with a PM). Women are generally at a higher risk of getting this cancer, which is known as gestational trophoblastic neoplasia (GTN), if they are over 40 years old, have a large increase in the size of the womb, have large cysts in the ovaries or have high initial levels of β -human chorionic gonadotrophin (hCG) (the pregnancy hormone) in their blood. Although treatment of the cancer with chemotherapy (anti-cancer drugs) is almost always effective, it has been suggested that routinely giving women anti-cancer drugs (P-Chem) before or after the removal the molar tissue may reduce the risk of the cancerous tissue developing.

By doing this review, we tried to assess the benefits and risks of giving P-Chem to women with molar pregnancies, before or after ERPC. We found three randomised studies involving a total of 613 women. Two studies tested methotrexate in all women with a CM and one study tested dactinomycin in women with a CM who were at a high risk of getting GTN. The two methotrexate studies are older studies that used relatively poor research methods, therefore their findings cannot be relied upon. Overall the review findings suggest that P-Chem reduces the number of women developing cancer after molar pregnancy; however, this is probably only be true for women with high-risk moles. In addition, P-Chem might make the time to diagnosing the cancer longer and might increase the number of anti-cancer treatments needed to cure the cancer if it develops. We were unable to assess the short- and long-term side-effects of P-Chem in this review because there was not enough available data; however, we are concerned that the five- and eight-day courses of P-Chem used by researchers in these studies are too toxic to be given to women routinely.

Currently there is insufficient evidence to support giving anti-cancer drugs to women with molar pregnancies. However, GTN is almost always cured with modern care and P-Chem for molar pregnancy would only reduce the risk of needing full-scale chemotherapy,

but would not remove that risk. In addition, it would not change the need for careful monitoring and follow-up of women with hydatidiform moles.

SUMMARY OF FINDINGS FOR THE MAIN COMPARISON *[Explanation]*

Prophylactic chemotherapy compared with no prophylactic chemotherapy for hydatidiform mole						
Patient or population: women with a molar pregnancy Settings: inpatient Intervention: methotrexate or dactinomycin Comparison: placebo or no prophylaxis						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No prophylaxis	P-Chem				
Incidence of GTN (including low-quality studies)	Mixed-risk population		RR 0.37 (0.24 to 0.57)	550 women (3 studies)	⊕⊕○○ low	The NNTB to prevent 1 woman developing GTN after evacuation of HM was 6 (95% CI 5 to 10) . We downgraded this
	254 per 1000	94 per 1000 (61 to 145)				
	High-risk population		RR 0.29 (0.14 to 0.60)	99 women (2 studies)	⊕⊕○○ low	
490 per 1000	142 per 1000 (69 to 294)					
Incidence of GTN (excluding low-quality studies)	High-risk population		RR 0.28 (0.10 to 0.73)	59 women (1 study)	⊕⊕○○ low	The NNTB to prevent 1 woman developing GTN after evacuation of high-risk HM was 3 (95% CI 2 to

						20). We downgraded this evidence because only 1 small study (Limpongsanurak 2001) contributed data, giving an imprecise result
	500 per 1000	140 per 1000 (50 to 365)				
Time to GTN diagnosis (days)	The mean time to GTN diagnosis ranged from 35.7 days to 59.5 days	The mean time to GTN diagnosis in the intervention groups was 65.5 days to 81.8 days (higher)	MD 28.72 (13.19 to 44.24)	33 women (2 studies)	⊕⊕○○ low	We downgraded this evidence because the meta-analysis included 1 study of poor methodological quality (Kim 1986). When this study was excluded, the results of the remaining study (Limpongsanurak 2001 ; 19 women) were: MD 22.30; 95% CI -9.05 to 53.65
Number of courses of chemotherapy to cure	The mean number of courses of chemotherapy required to cure subsequent GTN was 1.4 courses (10 women)	The mean number of courses of chemotherapy required to cure subsequent GTN was 2.5 courses (4 women)	MD 1.10 (0.52 to 1.68)	14 women (1 study)	⊕○○○ very low	This analysis only included 1 study (Kim 1986) that we considered to be of a poor methodological quality

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).
CI: confidence interval; HM: hydatidiform mole; NNTB: number needed to treat for an additional beneficial outcome; RR: risk ratio; MD: mean difference; GTN: gestational trophoblastic neoplasia

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

The assumed risk for the mixed-risk population was calculated by using the weighted mean risk across the control group for this outcome. The assumed risk for the high-risk population was based on the control group of [Limpongsanurak 2001](#), which was the only study to evaluate a high-risk population only.

BACKGROUND

Description of the condition

Gestational trophoblastic disease (GTD) is a spectrum of disease characterised by an autonomous overgrowth of foetal chorionic tissue or trophoblast. Hydatidiform mole (HM) is the most common and benign form of the disease. The prevalence of HM is the highest in Asia, with rates ranging from 1 to 2 per 1000 pregnancies in Japan and China (Palmer 1994; Song 1987; Takeuchi 1987) to 12 per 1000 pregnancies in Indonesia, India and Turkey (Aziz 1984; Gul 1997; Steigrad 2003). In North America and Europe, the incidence is reported to be lower, at 0.5 to 1 per 1000 pregnancies (Lee 2009; Steigrad 2003). The incidence has been reported to vary with race (Tham 2003), maternal age (Parazzini 1986), parity (Bagshawe 1986) and diet (Berkowitz 1985; Parazzini 1988). Variations in prevalence may be because of differences in reporting between hospital-based and population-based data or in the availability of central pathology review (Seckl 2010; Smith 2003). HMs are categorised as partial (PM) or complete moles (CM) based on their gross morphology, histopathology and karyotype. CMs usually occur when a duplicated haploid sperm fertilises an anucleate or 'empty' ovum; the ensuing diploid product (usually 46 XX) is therefore paternally derived (Fisher 2009). PMs are usually triploid (69XXX, 69XXY or 69XYY), with two sets of paternal haploid genes and one set of maternal haploid genes, and occur when two sperm fertilise one ovum (Fisher 2009). With CMs there is no evidence of foetal tissue; however, with PM an embryo or foetus frequently dies in early pregnancy and foetal tissue and blood cells may be identified in 20% and 50% of PM specimens, respectively (Sebire 2009).

HMs usually present with vaginal bleeding. Associated features (including excessive uterine enlargement, theca lutein ovarian cysts, hyperemesis, pre-eclampsia and hyperthyroidism) are more common in CMs; however, they occur less frequently as the routine use of ultrasound has led to earlier diagnosis (Seckl 2010). The management of PMs and CMs is similar (Berkowitz 2009a; Berkowitz 2009b). For women who want to preserve their fertility, an evacuation of the retained products of conception (ERPC) is performed, ideally by suction curettage, to remove all trophoblastic tissue completely (Seckl 2010). Most women are cured in this way; however, in some women HM persists and becomes malignant (gestational trophoblastic neoplasia (GTN)), requiring treatment with chemotherapy. In Japan, second ERPCs are performed routinely for HMs within one week of the initial ERPC, to ensure that there is no residual molar tissue in the uterus (Sasaki 2009). Second ERPCs may reduce the risk of GTN; however, to our knowledge, there is currently insufficient evidence to support this routine practice. Hysterectomy may reduce the risk of GTN by up to 10% (Bahar 1989; Curry 1975) and is an option for women who do not wish to retain their fertility or who experience life-threatening bleeding

at the time of evacuation; however, it does not avoid the need for subsequent monitoring or chemotherapy (Tidy 2009).

Transformation to GTN is considered to have occurred when trophoblastic activity remains following evacuation, as shown by a plateau or rise in serial β -human chorionic gonadotrophin (hCG) levels, raised hCG levels six months after evacuation or if the histopathological examination indicates choriocarcinoma (Kohorn 2009). The risk of developing GTN is reported to be 16% to 20% in women with CM (Berkowitz 1995; Curry 1975; Felemban 1998; Seckl 2009) and 0.5% to 1% in women with PM (Bagshawe 1990; Seckl 2009). In the UK, this translates to a GTN transformation rate of approximately 8% of all molar pregnancies (Seckl 2009). Thresholds for treating persistent GTD differ by region with, for example, more than twice as many women in the US (20%) receiving chemotherapy for persistent GTD than in the UK (Hancock 2009).

HMs may be categorised as being at a low or high risk of malignant transformation based on criteria first introduced by Bagshawe 1976 (Table 1; Berkowitz 1987). Women with high-risk HMs have more than one of the following characteristics: an initial serum β -hCG more than 100,000 mIU/mL, uterine size larger than gestational age, theca lutein cysts more than 6 cm in diameter, maternal age over 40 years, and other associated medical and epidemiological factors, including previous GTD, hyperthyroidism and trophoblastic embolisation (Berkowitz 1995). Approximately 30% to 50% of high-risk HMs will progress to GTN (Goldstein 1981; Limpongsanurak 2001; Uberti 2009).

GTN, which may also follow a 'normal' pregnancy, ectopic or miscarriage, is classified as low or high risk using a modified World Health Organization (WHO) scoring system adapted by the International Federation of Gynaecology and Obstetrics (FIGO 2009). Low-risk GTN accounts for 95% of cases in the UK and has a cure rate of almost 100% (Seckl 2010). High-risk GTN has a cure rate of between 80% and 90%; these lesions require combination chemotherapy regimens and frequently develop drug resistance (Goldstein 2012).

Description of the intervention

Methotrexate was first reported to be active against trophoblastic tissue in the mid-1950s (Hertz 1956). Since then, GTN has been shown to be a highly chemosensitive disease, with various chemotherapeutic agents achieving good rates of cure. All women with 'low-risk' GTN and approximately 80% to 90% of women with 'high-risk' GTN will be cured following treatment with one or more chemotherapy regimens (Seckl 2010; Goldstein 2012). Since chemotherapy drugs are associated with various toxic effects, most commonly myelotoxicity, gastrointestinal toxicity, stomatitis and alopecia, the chemotherapeutic aim when treating GTN is to provide the most effective treatment with the least toxicity. Methotrexate and dactinomycin are considered to be relatively safe agents that are commonly administered as first-line chemotherapy for

GTN, alone or in combination with other agents (Alazzam 2012). They have not been shown to be associated with adverse reproductive outcomes, ovarian failure or second tumours (Goldstein 1995).

The use of prophylactic chemotherapy (P-Chem) in women with molar pregnancy was first described in 1966 (Lewis 1966). Since then, studies of dactinomycin and methotrexate administered before, during or after evacuation of a molar pregnancy have reported encouraging results (see Table 2). Several studies have found a significant reduction in GTN for high-risk HMs only (Kim 1986; Fasoli 1982; Park 1996). Various dosing schedules have been described including five-day dactinomycin (Goldstein 1974; Goldstein 1981; Limpongsanurak 2001; Park 1996), eight-day methotrexate-folinic acid (Goldstein 1971; Kim 1986; Park 1996) and single-dose dactinomycin (Uberti 2006; Uberti 2009).

How the intervention might work

As GTN is a highly chemosensitive disease, prophylaxis with chemotherapy agents that have known activity against trophoblast tumour cells may prevent progression to GTN. The use of P-Chem has been based on an assumption that the development of GTN is pre-determined, that metastatic GTN spreads via the bloodstream and that high serum levels of cytotoxic agents around the time of evacuation should reduce the ability of the trophoblast cells to invade or metastasise (Goldstein 1995).

P-Chem may be particularly useful in women with high-risk CMs, who have poor access to health care, for whom hormonal follow-up is not available, or where poor compliance may be an issue (Limpongsanurak 2001; Uberti 2006; Berkowitz 2009a). In Latin America, loss to follow-up may be as high as 44% in some areas; hence numerous referral centres in this region are reported to use P-Chem (Charry 2009). However, the use of P-Chem exposes women unnecessarily to toxic side effects (Kaye 2002 and Ratnam 1971 have reported toxicity-related deaths with methotrexate prophylaxis), may lead to inadequate follow-up, and incompletely protects women against persistent tumour (Goldstein 1995; Hancock 2009). Furthermore, P-Chem may favour the development of drug resistance (Kim 1986) and delay the time to effective treatment, thereby having adverse effects on survival.

Why it is important to do this review

P-Chem for high-risk HM appears to be routine clinical practice in some regions of the world (Charry 2009). Although several studies have been reported, it remains unclear whether P-Chem, which may be associated with substantial toxicity, will prevent malignant transformation of HM. Furthermore, if benefits to P-Chem exist, it is not clear which drug regimen might have the best effectiveness-to-toxicity ratio. We undertook this review in an

attempt to clarify the benefits and risks associated with P-Chem for HM.

OBJECTIVES

To evaluate the effectiveness and safety of P-Chem for the prevention of GTN in women with molar pregnancy.

To investigate whether any subgroup of women with HM may benefit more from P-Chem than others.

METHODS

Criteria for considering studies for this review

Types of studies

Randomised controlled trials (RCTs).

Types of participants

Inclusions

All women diagnosed with HM.

Exclusions

Women who were diagnosed with other types of GTD such as invasive moles, choriocarcinoma and placental site trophoblastic tumour (PSTT).

Types of interventions

P-Chem compared with no or other treatments (e.g. placebo or analgesic drugs).

Types of outcome measures

Primary outcomes

- Incidence of GTN (invasive mole, choriocarcinoma, PSTT and epithelioid trophoblastic tumour (ETT)).

Secondary outcomes

- Drug toxicity, including myelotoxicity, gastrointestinal toxicity, stomatitis and alopecia.
- Overall survival (more than five years).
- Time to negative conversion of serum or urine β -hCG.
- Time to GTN diagnosis.
- Incidence and nature of subsequent pregnancies.
- Quality of life (QoL).

Search methods for identification of studies

Electronic searches

We searched the MEDLINE (1946 to February 2012) (Appendix 1), EMBASE (1980 to September 2012) (Appendix 2) and the Cochrane Central Register of Controlled Trials (CENTRAL, Issue 2, 2012) (Appendix 3). No language restriction was applied.

Searching other resources

All relevant articles were identified on PubMed, and, using the 'related articles' feature, a further search was carried out for newly published articles. The reference lists from identified published trials were handsearched for further clinical trials. Papers in all languages were sought and translated as necessary. We searched the *meta*Register of Controlled Trials (*mRCT*) and the National Research Register (NRR) archive for ongoing trials.

Data collection and analysis

Selection of studies

We downloaded all titles and abstracts retrieved by electronic searching to a reference management database, duplicates were removed and the remaining references were examined by three review authors (JF, FH) independently. Those studies that clearly did not meet the inclusion criteria were excluded and copies of the full text of potentially relevant references were obtained. The eligibility of retrieved papers was assessed independently by three review authors (JF, LX, TL). We documented the reasons for exclusion.

Data extraction and management

For included studies, data on patient characteristics; number recruited to each arm; number excluded from analysis; type of intervention; proportion of participants who received all, part or none of the intended treatment; methods of randomisation, blinding and allocation concealment; length of follow-up and data on outcome were extracted independently by three review authors (JF,

HC, TL). Differences between review authors were resolved by discussion or by appeal to a fourth review author (LH, FF or TW) if necessary.

Assessment of risk of bias in included studies

We used the Cochrane Collaboration's tool for assessing risk of bias (Higgins 2011). Three review authors (JF, LX, TL) independently assessed the risk of bias within each included study based on the following six domains, with review authors' judgements presented as answers of 'Yes' (low risk of bias); 'No' (high risk of bias), and 'Unclear' (uncertain risk of bias).

- Selection bias: random sequence generation and allocation concealment.
- Performance bias: blinding of participants and personnel (women and treatment providers).
- Detection bias: blinding of outcome assessment.
- Attrition bias: incomplete outcome data. We considered studies to be at a high risk of bias if more than 20% of women were lost to follow-up or reasons for loss to follow-up differed between treatment arms.
- Reporting bias: selective reporting of outcomes.
- Other possible sources of bias.

Assessment of heterogeneity

Random-effects models were used for all meta-analyses (DerSimonian 1986). Heterogeneity between studies was assessed by visual inspection of forest plots, by estimation of the I^2 statistic, which summarises the percentage heterogeneity between trials that cannot be ascribed to sampling variation, and by a formal statistical test of the significance of the heterogeneity (Deeks 2001). A level of I^2 of less than 25% was considered as low level heterogeneity, 25% to 50% as moderate level, and higher than 50% as substantial heterogeneity (Higgins 2003). If there was evidence of substantial heterogeneity, the possible reasons for this were investigated and reported.

Data synthesis

Meta-analysis was carried out using the Review Manager 5.1 software (RevMan 2011). We used random-effects models for all meta-analyses (DerSimonian 1986). For dichotomous outcomes, we calculated risk ratios (RR) and associated 95% confidence intervals (CIs). For continuous outcomes we pooled the mean differences (MD) between the treatment arms where trials measured the outcome on the same scale.

Subgroup analysis and investigation of heterogeneity

We subgrouped women by the type of chemotherapy agent (i.e. methotrexate and dactinomycin). We had planned to perform other subgroup analyses, including subgroups of women at a low and high risk of GTN, and according to drug regimens; however,

this was not possible since these data were not reported in the included trials.

Sensitivity analysis

We performed sensitivity analysis to assess the robustness of the meta-analyses by comparing the results using all trials and then excluding trials of lower methodological quality or those considered to be at a higher risk of bias.


RESULTS

Description of studies

Results of the search

From the search, we identified and screened 133 citations, retrieving the full text of 86 citations that we considered potentially eligible for inclusion in this review. We excluded 79 out of these 86 papers. Of the remaining seven reports, we excluded four on the basis that they were not RCTs ([Geng 2011](#); [Uberti 2006](#); [Uberti 2009](#) - two papers), and included the remaining three studies ([Kashimura 1986](#); [Kim 1986](#); [Limpongsanurak 2001](#)) ([Figure 1](#)).

Figure 1. Study flow diagram.



Included studies

We included three studies with a total of 613 participants ([Kashimura 1986](#); [Kim 1986](#); [Limpongsanurak 2001](#)). Two studies used prophylactic methotrexate and one trial used prophylactic dactinomycin. The number of evaluable participants in the three studies was 550. See [Characteristics of included studies](#).

[Kashimura 1986](#) 'randomly selected' 420 women with low- or high-risk CM to receive one course of prophylactic methotrexate (10 mg daily for seven days) within three weeks of ERPC, or no prophylaxis, and evaluated subsequent rates of GTN in the two groups.

[Kim 1986](#) randomised 133 women with low- or high-risk CM to prophylactic methotrexate or no prophylaxis. Only 71 out of 133 women completed this trial and were included in the analyses (39 in the treatment group and 32 in the untreated group). The intervention group (18 out of 31 low-risk women and 21 out of 40 high-risk women) received one course of methotrexate with citrovorum rescue factor (methotrexate 1.0 mg/kg/day intramuscular (IM) on days 1, 3, 5 and 7; citrovorum rescue factor 0.1 mg/kg/day IM on days 2, 4, 6 and 8); the control group received no treatment. The ERPC in the intervention group was done on the

third or fourth day of P-Chem.

[Limpongsanurak 2001](#) randomised 60 women with high-risk CM to dactinomycin prophylaxis (10 µg/kg body weight daily for 5 days; 30 women) or no prophylaxis (30 women) within one week after ERPC. One woman was lost to follow-up.

Excluded studies

We excluded three retrospective studies ([Geng 2011](#); [Uberti 2006](#); [Uberti 2009](#) - two reports). See [Characteristics of excluded studies](#).

Risk of bias in included studies

We assessed the risk of bias of the three included RCTs using the Cochrane Collaboration's 'Risk of bias' tool, see [Figure 2](#). Overall, we consider the older studies of [Kashimura 1986](#) and [Kim 1986](#) to be at a high risk of bias and [Limpongsanurak 2001](#) to be at a low risk of bias. Dr. Limpongsanurak provided us with additional methodological details for [Limpongsanurak 2001](#) via e-mail. Although we attempted to contact the other authors by e-mail for more details, we were unsuccessful as we had no contact details for the authors of [Kim 1986](#), and received no reply to our queries from Dr. Kashimura ([Kashimura 1986](#)).

Figure 2. 'Risk of bias' summary: review authors' judgements about each 'Risk of bias' item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Kashimura 1986	-	?	?	?	?	-	-
Kim 1986	?	?	?	?	-	-	-
Limpongsanurak 2001	+	+	+	?	+	+	+

Allocation

[Limpongsanurak 2001](#) used lot-drawing to randomise women and sealed opaque envelopes to conceal random group allocations; we assessed these methods to be at a low risk of bias. We assessed the randomisation and allocation methods used in [Kim 1986](#) to be of unclear risk of bias and assessed the random sequence generation in [Kashimura 1986](#) to be at a high risk of bias. In the latter report, participants were 'selected at random', which suggests that participant selection in this study may not have been a truly random process. Furthermore, no randomisation ratio was described and yet the intervention and control groups were very different sizes (293 women versus 127 women).

Blinding

The only trial that reported blinding was [Limpongsanurak 2001](#). Although precise details were not reported, this trial was described as 'double-blind' as control participants received a similar-looking intravenous (IV) solution with analgesic drugs for five days, and neither the participant nor the attending doctor knew to which group the participant had been allocated. It is unclear whether outcome assessment was also blind.

Incomplete outcome data

We assessed [Kim 1986](#) as being at a high risk of attrition bias as 62 out of 133 women were excluded from the analyses owing to loss to follow-up (36 women), insufficient length of follow-up (7 women) and hysterectomy (19 women). Of the 60 women in [Limpongsanurak 2001](#), one woman in the P-Chem group was lost to follow-up one month after treatment and was not included in the main analyses. [Kashimura 1986](#) reported complete data sets for the main outcomes.

Selective reporting

All pre-specified outcomes were reported for [Kim 1986](#) and [Limpongsanurak 2001](#). [Kashimura 1986](#) failed to report baseline characteristics of the two study groups that could represent reporting bias, especially since there were proportionally more women aged 40 years and over in the control group.

Other potential sources of bias

In [Kashimura 1986](#), more women in the control group were 40 years old and over (22% in control group versus 11% in P-Chem group). The older women were more likely to progress to GTN (39% versus 12%). This may have biased the results in favour of the P-Chem group.

[Limpongsanurak 2001](#) included women with high-risk CM only, whereas [Kim 1986](#) and [Kashimura 1986](#) included women with

low- and high-risk CM. Since [Kim 1986](#) showed that P-Chem was not beneficial to women with low-risk CM, by including these women the meta-analysis results may be biased in the direction of the control arm.

[Limpongsanurak 2001](#) and [Kashimura 1986](#) gave P-Chem after ERPC whereas [Kim 1986](#) started P-Chem before the ERPC. If treatment was commenced before ERPC, this trial may have included women with PM and hydropic abortion that would otherwise have been excluded following histological diagnosis of evacuation products. It is unclear whether women with PM or hydropic abortion were present in the same numbers within the allocated groups.

[Kashimura 1986](#) and [Kim 1986](#) were older studies that took place before rigorous RCT guidelines were in place and therefore are lacking in methodological quality. It is not possible to determine whether these were true RCTs so we have assumed that they were but considered them to be at a high risk of bias overall; therefore we have performed sensitivity analysis and downgraded the results accordingly.

Effects of interventions

See: [Summary of findings for the main comparison](#)

Incidence of GTN

Incidence of GTN (overall)

P-Chem was associated with a significant reduction in the incidence of GTN compared with the control group (3 studies; 550 participants; 30 out of 361 versus 48 out of 189; RR 0.37; 95% CI 0.24 to 0.57; $I^2 = 0\%$; $P < 0.00001$; [Analysis 1.1](#)) with no significant difference between methotrexate and dactinomycin prophylaxis subgroups.

We performed sensitivity analysis for this outcome as two of the included studies were at a high risk of bias. When these two studies (both in the methotrexate subgroup) were excluded, only one trial remained (59 participants; 4 out of 29 versus 15 out of 30; RR 0.28; 95% CI 0.10 to 0.73; $P = 0.01$); therefore we consider this evidence to be of a low quality.

Incidence of GTN (high-risk HM only)

When only high-risk women were included, P-Chem was similarly associated with a significant reduction in the incidence of GTN compared with the control group (2 studies; 99 participants; 7 out of 50 versus 24 out of 49; RR 0.29; 95% CI 0.14 to 0.60; $I^2 = 0\%$; $P = 0.001$; [Analysis 1.2](#)). Sensitivity analysis gave the same results as above when the only well-conducted trial was included

(59 participants; 4 out of 29 versus 15 out of 30; RR 0.28; 95% CI 0.10 to 0.73; P = 0.01).

Invasive mole and choriocarcinoma

There were insufficient data from these trials to analyse the rates of invasive mole and choriocarcinoma in the study groups. [Limpongsanurak 2001](#) utilised a prognostic scoring system to diagnose GTN (not histology) and secondary histology was known in only three participants, two of whom underwent hysterectomy for excessive bleeding. [Kashimura 1986](#) diagnosed 27 out of 45 cases of GTN histologically and 18 out of 45 by a Japanese prognostic scoring system. Four of these participants had choriocarcinoma (two in each study group) and 41 were considered to have invasive mole. [Kim 1986](#) diagnosed most cases of GTN based on persistent or rising hCG levels, persistent or recurrent uterine haemorrhage, or clinical/histological evidence of metastases, and did not distinguish between invasive mole and choriocarcinoma.

Time to GTN diagnosis

Two studies reported this outcome for 33 participants who developed GTN ([Kim 1986](#); [Limpongsanurak 2001](#)). The time to GTN diagnosis was significantly longer in the P-Chem group compared with the control group (MD 28.72 days; 95% CI 13.9 to 44.24; $I^2 = 0\%$; P = 0.0003; [Analysis 1.3](#)). When we excluded [Kim 1986](#) from the sensitivity analysis, results for the one remaining study were similar but not significant (19 participants; MD 22.30 days; 95% CI to 9.05 to 53.65; P = 0.16).

Toxicity

None of the studies reported adverse effects in the control groups and we have not imputed these data. [Limpongsanurak 2001](#) (five-day dactinomycin) only reported side effects as percentages, including stomatitis (10%), nausea or vomiting (10%), oral ulcers (3.3%) and hair loss (13.3%). All adverse effects in this study were grade 1 except for two women with patchy alopecia (grade 2). The other two studies reported the following:

- [Kashimura 1986](#) (239 women): stomatitis (10.3%), nausea or vomiting (6.8%) and leukopenia (4.4%);
- [Kim 1986](#) (39 women): epithelial (5.1%), hepatic (7.7%), haematological (7.7%) and neuromuscular (2.6%) toxicity.

In these latter two studies, grades of toxicity were not reported. Both reports state that there were no severe complications or drug-related deaths.

Courses of chemotherapy

Only one poor-quality study reported this outcome. Women in the methotrexate prophylaxis group needed more subsequent chemotherapy courses for GTN treatment than the control group (14 women; MD 1.10; 95% CI 0.52 to 1.68; P = 0.0002).

Survival

There were no deaths during the follow-up periods in [Kim 1986](#) and [Limpongsanurak 2001](#); however, [Kashimura 1986](#) reported three deaths from metastatic disease that occurred two to 12 years after ERPC, including two women in P-Chem group and one in the control group.

Subsequent pregnancies

[Limpongsanurak 2001](#) did not report subsequent pregnancy rates. Women in [Kim 1986](#) were given contraception for one year after ERPC and subsequent pregnancy rates were assessed. Out of 51 women who attempted to become pregnant, 40 had 44 pregnancies including 24 out of 29 in the P-Chem group and 16 out of 22 in the control group. The rates of full-term pregnancies were reported as being similar in the two groups (25 out of 27 (93%) in the P-Chem group versus 16 out of 17 (94%) in the control group).

[Kashimura 1986](#) obtained these data for 101 out of 420 participants only (24%): 74 out of 112 (67%) subsequent pregnancies were normal full-term pregnancies in the P-Chem group compared with 19 out of 31 (61%) in the control group. The induced abortion rate was 22% of pregnancies in the P-Chem group and 13% in the control group.

It was not possible to perform a meta-analysis of these very limited data.

DISCUSSION

Summary of main results

In a limited meta-analysis of three studies that included low- and high-risk molar pregnancies, P-Chem reduced the incidence of subsequent GTN by approximately two-thirds overall. On sensitivity analysis, where the only satisfactory study (high-risk women only) contributed data, the results showed a similar effect but with a wider CI. (See [Summary of findings for the main comparison](#)). The interpretation of these results was influenced by the poor methodological quality of the two older included studies, and the small number of participants in the only other study. We therefore consider this evidence to be of low quality.

The diagnosis of GTN occurred approximately one month later in the P-Chem group compared with the control group in the meta-analysis of two studies that reported this outcome. The P-Chem group needed approximately one extra course of chemotherapy than the control group to achieve a cure in the participants whose disease progressed to GTN. We consider this evidence to be of a very low quality for similar reasons as those given above.

Data on toxicity were insufficient for meta-analysis; however, it was reported that no participants in any of the studies experienced severe drug-related complications or deaths.

Overall completeness and applicability of evidence

We consider this evidence to be incomplete and not widely applicable:

Incidence of GTN

This limited evidence in favour of P-Chem may apply to women with high-risk CM only. [Kim 1986](#) found that P-Chem did not benefit women with low-risk HM and some non-randomised studies have drawn similar conclusions ([Fasoli 1982](#); [Park 1996](#)). This is probably why the more recent studies have excluded low-risk women (e.g. [Limpongsanurak 2001](#); [Uberti 2009](#)). Furthermore, the incidence of GTN in the high-risk control groups of included studies was high, at 47% and 50% for [Kim 1986](#) and [Limpongsanurak 2001](#), respectively. This may reflect regional differences in the rates of CM transformation or the selection/diagnostic criteria applied, and needs further clarification.

Only two chemotherapeutic agents, namely, methotrexate and actinomycin D, were investigated. None of the included studies evaluated the less toxic single-dose regimens of these agents that have been shown to be useful to treat low-risk GTN ([Alazzam 2012](#)). One RCT of single-dose dactinomycin for P-Chem was proposed by [Goldstein 1995](#), but has never been conducted. [Uberti 2006](#) and [Uberti 2009](#) report the results of two retrospective studies of single-dose dactinomycin administered before evacuation of high-risk molar pregnancies, showing a reduction in post-molar GTN of 76% and 46%, respectively, with minimal adverse effects. Since this low-dose dactinomycin regimen appears to be in practice (personal communication with Dr Uberti) it should be evaluated in an RCT.

Toxicity

Toxicity was not rigorously reported in the included studies and meta-analyses of these data were not possible. When used to treat low-risk GTN, five-day dactinomycin and five- and eight-day methotrexate regimens have been associated with severe adverse effects including myelosuppression, hepatotoxicity and alopecia (dactinomycin); hence there is a move towards less toxic chemotherapy regimens for the treatment of low-risk GTN ([Alazzam 2012](#)). For this reason, the more toxic five- and eight-day regimens that have historically been used in studies of P-Chem ([Table 2](#)) are unlikely to be favoured as prophylaxis for HM.

Survival

Only one study ([Kashimura 1986](#)) reported long-term follow-up, with three deaths occurring from metastatic disease between two and 12 years after ERPC (two in the P-Chem group and one in the control group); neither [Kim 1986](#) nor [Limpongsanurak 2001](#) assessed the impact of P-Chem on long-term overall survival, hence the evidence for this outcome is incomplete. Longer follow-up of participants is needed in any future studies of P-Chem interventions for HM.

Subsequent pregnancy and QoL

Although subsequent pregnancies were reported by [Kashimura 1986](#) and [Kim 1986](#), these data were incomplete (owing to high attrition) and it was not possible to draw any conclusions. [Kim 1986](#) reported that 78.4% of their participants experienced at least one pregnancy after one year of contraception during the follow-up period and the frequency of full-term delivery was 92.6% and 94.1%, for the treatment and control groups, respectively. In [Kashimura 1986](#), comparable rates of secondary infertility and a similar time to first menstruation after ERPC were reported in the P-Chem and control groups. Neither [Kim 1986](#) nor [Limpongsanurak 2001](#) assessed the impact of P-Chem on long-term ovarian function or QoL. However, these chemotherapy agents have been used extensively to treat GTN over several decades and have not been shown to adversely affect reproductive outcomes, ovarian function or to be associated with second tumours ([Garner 2002](#); [Garrett 2008](#); [Goldstein 1995](#); [Uberti 2006](#); [Uberti 2009](#)).

[Uberti 2009](#) has suggested that P-Chem reduces the emotional complications associated with HM but we were unable to corroborate this owing to a lack of QoL data.

Drug resistance

Drug resistance may occur following P-Chem, as the agents used for P-Chem are also used as first-line treatment for GTN. Only one included study compared the number of courses of chemotherapy required to treat subsequent GTN ([Kim 1986](#)). These investigators found that women in the P-Chem group (4 women) required more courses than women in the control group (10 women), suggesting that P-Chem may increase resistance to subsequent chemotherapy. Owing to the poor methodological quality of the [Kim 1986](#) study and the small number of participants concerned, we are very uncertain of this estimate of effect ([Summary of findings for the main comparison](#)). To prevent resistance to treatment it has been suggested that an alternative agent be used for the treatment of persistent disease ([Goldstein 1995](#)). Drug resistance could not be adequately evaluated in this review and warrants further investigation.

Time to GTN diagnosis

The time interval from the index pregnancy to the diagnosis of GTN is considered to be a risk factor for the development of GTN and is included in the Modified WHO Prognostic Scoring System (FIGO 2009). Again, extremely limited evidence from Kim 1986 suggests that P-Chem may delay the time to diagnosis of GTN. It is unclear whether such a delay might potentially lead to up-scoring of GTN lesions from low to high risk. Investigators of some retrospective studies of single-dose dactinomycin found no difference in the time to GTN diagnosis among their participants (Uberti 2006; Uberti 2009). Thus further research is needed to clarify the impact of low-dose P-Chem on the time to diagnosis and subsequent GTN risk scores.

Timing of ERPC

Most studies of P-Chem to date have administered P-Chem before performing the ERPC with the exception of Limpongsanurak 2001, Kashimura 1986 and Koga 1968, where participants received their P-Chem within one week and three weeks of the ERPC. Only Limpongsanurak 2001 included participants based on a histological diagnosis. Goldstein 1971 was the first to describe performing the ERPC on P-Chem day three (see Table 2) and this practice was included in several subsequent studies, including Kim 1986. Administering P-Chem before ERPC may theoretically be more effective in preventing haematogenous spread of the molar tissue during the procedure. Since the incidence of GTN in the high-risk groups of Kim 1986 and Limpongsanurak 2001 were similar, it would appear that administering P-Chem before ERPC may not confer any additional benefit. Treating women with P-Chem before evacuation carries the inherent risk of over-treating women, as 10% of cases that are thought to be molar on ultrasound may turn out to be non-molar hydropic abortions (Fowler 2006). Thus we propose that, in any future studies of P-Chem, the intervention is administered after ERPC, and following a histological diagnosis of CM.

Quality of the evidence

We consider this evidence to be of a low to very low quality. This conclusion is based on our assessment that two of the included studies were of poor methodological quality and at a high risk of bias (See Risk of bias in included studies); the third study was of a good quality but consisted of only 60 participants. Kashimura 1986 and Kim 1986 are older studies that took place before rigorous RCT guidelines were in place.

With reference to the evidence for high-risk CMs only, we calculated that the number needed to treat for an additional beneficial outcome (NNTB) to prevent one woman with high-risk CM developing GTN was 3, with a 95% CI of 2 to 20 women; this wide CI illustrates the uncertainty concerning this evidence. Therefore, we believe that further research is very likely to have an important

impact on our confidence in the estimate of effects, and is likely to change the estimates.

With regard to the number of courses of chemotherapy required to cure post-molar GTN, we are very uncertain about this estimate of effect (see Summary of findings for the main comparison).

Potential biases in the review process

We included all identified RCTs in this review, including two older studies of poor methodological quality. We rigorously debated the merits of including these weaker studies as, in so doing, we might have biased the results in favour of P-Chem. In particular, Kashimura 1986 describes its study design as 'prospective', with participants 'selected at random'. While some authors have interpreted this study design as an RCT (Limpongsanurak 2001), it has also been referred to as a retrospective study (Goldstein 1995). We were unsuccessful in making contact with the investigators of either Kim 1986 or Kashimura 1986 and therefore it was not possible to determine whether these were true RCTs. We decided to include them in our meta-analyses. This may seem controversial; however, we performed sensitivity analyses and downgraded the meta-analyses results accordingly. Sensitivity analysis of the main outcome produced similar findings when these studies were excluded. Furthermore, these weaker studies included low- and high-risk women in their sample, which may have biased the results in the direction of the control group, as P-Chem was found in these studies to have little benefit for women with a low risk of developing GTN.

Agreements and disagreements with other studies or reviews

Most recent studies of P-Chem have been conducted in Asia and South America. This may be indicative of higher prevalences of GTD, limited health resources or lower rates of follow-up experienced by in these regions. However, concerns regarding the exposure of women to unnecessary side effects (Goldstein 1995) may have played a role in the lack of contributing data from centres in North America and Europe.

Most recently, investigators in Brazil conducted two retrospective case-control studies and reported a significant reduction in the rate of GTN transformation with the use of a single bolus dose of dactinomycin before evacuation for high-risk HM, with minimal side effects (see Table 2; Uberti 2006; Uberti 2009). Baseline risk scores and hCG levels were significantly greater in the P-Chem arm of the latter study, yet the incidence of GTN was significantly lower in the P-Chem group compared with the control group. These studies suggest that low-dose dactinomycin may have an improved effectiveness-to-toxicity ratio and, hence, greater general acceptability as a P-Chem regimen.

It has been argued that because of the excellent primary cure rates among women with GTN, most doctors prefer to monitor hCG levels in women following HM, rather than administer P-Chem (Hurteau 2003). In addition, even if prophylaxis reduces the risk of GTN, women who are given prophylaxis would still require the same monitoring and follow-up as those who are not. However, P-Chem this may reduce emotional costs for affected women, as well as operating costs for institutions (Uberti 2009). In Uberti 2009, the number of women with high-risk HM who needed to be treated (NNTB) to prevent one case of GTN was seven; at this rate, P-Chem would apparently result in substantial cost savings to their GTD centre. Following a personal communication with Dr Uberti, we understand that this regimen is already in clinical practice in this GTD Centre in Brazil. The emotional (QoL) and cost implications of P-Chem versus no P-Chem could not be evaluated in this review,

P-Chem may not be the only method of reducing the incidence of GTN. In Japan, second ERPCs are performed within one week of the first ERPC following histological confirmation of HM (Sasaki 2009), and, in Indonesia, women with HM are given vitamin A supplementation (personal communication; Andrijono 2010). In one double-blind RCT conducted in Indonesia, vitamin A prophylaxis was compared with an identical placebo in women with CMs (Andrijono 2010). The theoretical basis for this intervention was that vitamin A has been shown to cause trophoblastic cells to undergo apoptosis (Andrijono 2010). Investigators reported that only 2 out of 30 women (6.7%) with CM who received 200,000 IU of vitamin A per day progressed to GTN compared with 10 out of 35 in the placebo group (28.6%) (P = 0.029). Side effects appeared to have been minimal. The proportion of high-risk CMs and the median duration of prophylaxis were not reported; however, these results are encouraging and warrant further research into vitamin A supplementation in women with HM.

AUTHORS' CONCLUSIONS

Implications for practice

P-Chem may reduce the risk of progression to GTN in women with CMs who are at a high risk of malignant transformation. However, the five and eight-day methotrexate and dactinomycin regimens studied in this review were too toxic for routine use, may delay the time to GTN diagnosis and may lead to subsequent drug resistance. The current evidence in favour of P-Chem is limited by the small numbers and poor methodological quality of available

RCTs in this field. Hence there is currently insufficient evidence to support the use of P-Chem in clinical practice.

Implications for research

One well-conducted RCT of single-dose dactinomycin compared with placebo for high-risk CM may be appropriate in countries with limited healthcare resources and where follow-up and hCG surveillance are more difficult. Baseline histology should be reported and outcomes should include time to GTN diagnosis, HM and GTN risk scores at diagnosis, the number of courses to cure, long-term survival, QoL, subsequent pregnancies and an economic evaluation of the intervention groups over time.

Other RCTs that may be helpful include:

- vitamin A studies (e.g. a 200,000 IU bolus or monthly) versus placebo, in women with a histological diagnosis of HM;
- second ERPC versus no additional (routine) ERPC in women with a histological diagnosis of HM. (There is currently a pilot Phase II study [GOG-0242] underway to investigate the effect of a second ERPC in women with low-risk GTN on the frequency of surgical cure and the rate of persistent GTN (GOG 0242)).

ACKNOWLEDGEMENTS

We would like to thank:

- Clare Jess, Gail Quinn, Jo Morrison and the staff of the Cochrane Gynaecological Cancer Review Group for their advice and administrative support throughout the review process;
- Library staff at the Royal United Hospital, Bath, UK, for their assistance with the sourcing of articles;
- Donald Goldstein for responding to reprint requests;
- Elza Uberti for providing copies of Uberti 2006 and Uberti 2009 and responding to e-mailed queries;
- Andri Andrijono for responding to e-mailed queries.

This review received methodological and statistical support as part of the 10/4001/12 NIHR Cochrane Programme Grant Scheme - *Optimising Care, Diagnosis and Treatment Pathways to Ensure Cost Effectiveness and Best Practice in Gynaecological Cancer: Improving Evidence for the NHS*

REFERENCES

References to studies included in this review

Kashimura 1986 *{published data only}*

Kashimura Y, Kashimura M, Sugimori H, Tsukamoto N, Matsuyama T, Matsukuma K, et al. Prophylactic chemotherapy for hydatidiform mole: five to 15 years follow up. *Cancer* 1986;**58**:624–9.

Kim 1986 *{published data only}*

Kim DS, Moon H, Kim KT, Moon YJ, Hwang YY. Effects of prophylactic chemotherapy for persistent trophoblastic disease in patients with complete hydatidiform mole. *Obstetrics and Gynecology* 1986;**67**:690–4.

Limpongsanurak 2001 *{published data only}*

Limpongsanurak S. Prophylactic actinomycin D for high-risk complete hydatidiform mole. *The Journal of Reproductive Medicine* 2001;**46**:110–6.

References to studies excluded from this review

Geng 2011 *{published data only}*

Geng S, Feng FZ, Xiang Y, Wan XR, Zhou Y, Zhonghua Fu, et al. Analysis of prophylactic chemotherapy outcome and clinical characteristics in patients of high-risk hydatidiform mole. *Zhonghua Fu Chan Ke Za Zhi* 2011;**46**(1):24–7. [Chinese]

Uberti 2006 *{published data only}*

Uberti EMH, Diestel MCF, Guimaraes FE, De Napoli G, Schmid H. Single-dose actinomycin D: efficacy in the prophylaxis of post molar gestational trophoblastic neoplasia in adolescents with high-risk hydatidiform mole. *Gynecologic Oncology* 2006;**102**:325–32.

Uberti 2009 *{published data only}*

Uberti EMH, Fajardo MC, Ferreira SVVR, Pereira MV, Seger RC, Moreira MAR, et al. Reproductive outcome after discharge of patients with high-risk hydatidiform mole with or without use of one bolus dose of actinomycin D, as prophylactic chemotherapy, during the uterine evacuation of molar pregnancy. *Gynecologic Oncology* 2009;**115**:476–81.

* Uberti EMH, Fajardo MC, da Cunha AGV, Rosa MW, Ayub ACK, Graudenz MS, et al. Prevention of post molar gestational trophoblastic neoplasia using prophylactic single bolus dose of actinomycin D in high-risk hydatidiform mole: a simple, effective, secure and low-cost approach without adverse effects on compliance to general follow-up or subsequent treatment. *Gynecologic Oncology* 2009;**114**:299–305.

References to ongoing studies

GOG 0242 *{published data only}*

Gynecologic Oncology Group. Second Curettage in Treating Patients With Persistent Non-Metastatic Gestational Trophoblastic Tumor. NCT00521118. <http://clinicaltrials.gov/ct2/show/NCT00521118?term=GOG-0242&rank=1>.

Additional references

Alazzam 2012

Alazzam M, Tidy J, Hancock BW, Osborne R, Lawrie T. First line chemotherapy for low risk gestational trophoblastic neoplasia. *Cochrane Database of Systematic Reviews* 2012, Issue 7. [DOI: 10.1002/14651858.CD007102.pub3]

Andrijono 2010

Andrijono M, Muhilal M. Prevention of post-mole malignant trophoblastic disease with vitamin A. *Asian Pacific Journal of Cancer Prevention* 2010;**11**:567–70.

Aziz 1984

Aziz MF, Kampono N, Moegni EM. Epidemiology of gestational trophoblastic neoplasm at the Dr.Cipto Mandunkusumo Hospital, Jakarta, Indonesia. *Advances in Experimental Medicine and Biology* 1984;**176**:165–75.

Bagshawe 1976

Bagshawe KD. Risk and prognostic factors in trophoblastic neoplasia. *Cancer* 1976;**38**:1373–85.

Bagshawe 1986

Bagshawe KD, Dent J, Webb J. Hydatidiform mole in England and Wales 1973–1983. *Lancet* 1986;**2**:673–7.

Bagshawe 1990

Bagshawe KD, Lawler SD, Paradinas FJ, Dent J. Gestational trophoblastic tumours following initial diagnosis of partial hydatidiform mole. *Lancet* 1990;**335**:1074–6.

Bahar 1989

Bahar AM, El Ashneh MS, Senthilselvan A. Hydatidiform mole in the elderly: hysterectomy or evacuation. *Obstetrics and Gynaecology* 1989;**29**:233–8.

Berkowitz 1985

Berkowitz RS, Goldstein DP, Bernstein MR. Natural history of partial molar pregnancy. *Obstetrics and Gynecology* 1985;**66**:677–81.

Berkowitz 1987

Berkowitz RS, Goldstein DP, Dubeshter B, Bernstein M. Management of complete molar pregnancy. *Journal of Reproductive Medicine* 1987;**32**:634–9.

Berkowitz 1995

Berkowitz RS, Goldstein DP. Gestational trophoblastic disease. *Cancer* 1995;**76**:2079–85.

Berkowitz 2009a

Berkowitz RS, Goldstein DP. Chapter 9: presentation and management of molar pregnancy. In: Hancock BW, Seckl MJ, Berkowitz RS, Cole LA editor(s). *Gestational Trophoblastic Disease*. 3rd Edition. London: International Society for the Study of Trophoblastic Disease, 2009:249–76. www.isstd.org/isstd/chapter09/files/GTD3RDCH09.pdf (accessed 21 August 2012).

Berkowitz 2009b

Berkowitz RS, Goldstein DP. Current management of gestational trophoblastic diseases. *Gynecologic Oncology* 2009;**112**:654–62.

Charry 2009

Charry RC. Chapter 16: presentation and management of molar pregnancy and gestational trophoblastic neoplasia in Latin America. In: Hancock BW, Seckl MJ, Berkowitz RS, Cole LA editor(s). *Gestational Trophoblastic Disease*. 3rd Edition. London: International Society for the Study of Trophoblastic Diseases, 2009:407-19. www.isstd.org/isstd/chapter16.html (accessed 21 August 2012).

Curry 1975

Curry SL, Hammond CB, Tyrey L, Creasman WT, Parker RT. Hydatidiform mole: diagnosis, management, and long-term follow up of 347 patients. *Obstetrics and Gynecology* 1975;**45**:1-8.

Deeks 2001

Deeks JJ, Altman DG, Bradburn MJ. Statistical methods for examining heterogeneity and combining results from several studies in meta-analysis. In: Egger M, Davey Smith G, Altman DG editor(s). *Systematic Reviews in Health Care: Meta-Analysis in Context*. 2nd Edition. London: BMJ Publication Group, 2001.

DerSimonian 1986

DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled Clinical Trials* 1986;**7**:177-88.

Fasoli 1982

Fasoli M, Ratti F, Francheschi S, La Vecchia C, Pecorelli S, Mangioni C. Management of gestational trophoblastic disease: results of a cooperative study. *Obstetrics and Gynecology* 1982;**60**(2):205-9.

Felemban 1998

Felemban AA, Bakri YN, Alkharif HA, Altuwaijri SM, Shalhoub J, Berkowitz RS. Complete molar pregnancy clinical trends at King Fahad Hospital, Riyadh, Kingdom of Saudi Arabia. *The Journal of Reproductive Medicine* 1998;**43**(1):11-3.

FIGO 2009

FIGO Committee on Gynecologic Oncology. Current FIGO staging for cancer of the vagina, fallopian tube, ovary, and gestational trophoblastic neoplasia. *International Journal of Gynecology and Obstetrics* 2009;**105**:3-4.

Fisher 2009

Fisher R. Chapter 2: genetics. In: Hancock BW, Seckl M, Berkowitz RS, Cole LA editor(s). *Gestational Trophoblastic Disease*. 3rd Edition. London: International Society for the Study of Trophoblastic Diseases, 2009:6-48. www.isstd.org/isstd/chapter02.html (accessed 21 August 2012).

Fowler 2006

Fowler DJ, Lindsay I, Seckl MJ, Sebire NJ. Routine pre-evacuation ultrasound diagnosis of hydatidiform mole: experience of more than 1000 cases from a regional referral center. *Ultrasound in Obstetrics and Gynecology* 2006;**27**:56-60.

Garner 2002

Garner EI, Lipson E, Bernstein MR. Subsequent pregnancy experience in patients with molar pregnancy and gestational trophoblastic tumor. *Journal of Reproductive Medicine* 2002;**47**(5):380-6.

Garrett 2008

Garrett LA, Garner EIO, Feltmate CM. Subsequent pregnancy outcomes in patients with molar pregnancy and persistent gestational trophoblastic neoplasia. *Journal of Reproductive Medicine* 2008;**53**:481-6.

Goldstein 1971

Goldstein DP. Prophylactic chemotherapy of patients with molar pregnancy. *Obstetrics and Gynecology* 1971;**38**(6):817-22.

Goldstein 1974

Goldstein DP. Prevention of gestational trophoblastic disease by use of actinomycin D in molar pregnancies. *Obstetrics and Gynecology* 1974;**43**:475-9.

Goldstein 1981

Goldstein DP, Berkowitz RS, Bernstein MR. Management of molar pregnancy. *Journal of Reproductive Medicine* 1981;**26**:208-12.

Goldstein 1995

Goldstein DP, Berkowitz RS. Prophylactic chemotherapy of complete molar pregnancy. *Seminars in Oncology* 1995;**22**(2):157-60.

Goldstein 2012

Goldstein DP, Berkowitz RS. Current management of gestational trophoblastic neoplasia. *Hematology and Oncology Clinics of North America* 2012;**26**:111-131.

Gul 1997

Gul T, Yilmazturk A. A review of trophoblastic diseases at the Medical School of Dicle University. *European Journal of Obstetrics, Gynaecology and Reproductive Biology* 1997;**74**:37-40.

Hancock 2009

Hancock BW. Chapter 19: difference in management and treatment: a critical appraisal. In: Hancock BW, Seckl MJ, Berkowitz, Cole LA editor(s). *Gestational Trophoblastic Disease*. 3rd Edition. London: International Society for the Study of Trophoblastic Diseases, 2009:447-59. www.isstd.org/isstd/chapter19.html (accessed 21 August 2012).

Hertz 1956

Hertz R, Li MC, Spencer DB. Effect of methotrexate therapy upon choriocarcinoma and chorioadenoma. *Proceedings of the Society of Experimental Biology and Medicine* 1956;**93**(2):361-6.

Higgins 2003

Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analysis. *BMJ* 2003;**327**:557-60.

Higgins 2011

Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Hurteau 2003

Hurteau J. Gestational trophoblastic disease: management of hydatidiform mole. *Clinical Obstetrics and Gynecology* 2003;**46**(3):557-69.

Kaye 2002

Kaye DD. Gestational trophoblastic disease following complete hydatidiform mole in Mulago Hospital, Kampala, Uganda. *African Health Sciences* 2002;**2**(2):47–51.

Koga 1968

Koga K, Maeda K. Prophylactic chemotherapy with Amethopterin for prevention of choriocarcinoma following removal of hydatidiform mole. *American Journal of Obstetrics and Gynecology* 1968;**100**(2):270–5.

Kohorn 2009

Kohorn EI. Chapter 7: the FIGO 2000 staging and risk factor scoring system for gestational trophoblastic neoplasia: a critical analysis. In: Hancock BW, Seckl MJ, Berkowitz RS, Cole LA editor(s). *Gestational Trophoblastic Neoplasia*. 3rd Edition. London: International Society for the Study of Trophoblastic Disease, 2009:205-15. www.isstd.org/isstd/chapter07.html (accessed 21 August 2012).

Lee 2009

Lee C, Smith HO, Kim SJ. Chapter 3: epidemiology. In: Hancock BW, Newlands ES, Berkowitz RS, Cole LA editor (s). *Gestational Trophoblastic Diseases*. 3rd Edition. London: International Society for the Study of Trophoblastic Diseases, 2009:49-96. www.isstd.org/isstd/chapter03.html (accessed 21 August 2012).

Lewis 1966

Lewis JL, Gore H, Hertig AT. Treatment of trophoblastic neoplasms. With rationale for the use of adjunctive chemotherapy at the time of indicated operation. *American Journal of Obstetrics and Gynaecology* 1966;**98**:716.

Palmer 1994

Palmer JR. Advances in the epidemiology of gestational trophoblastic disease. *Journal of Reproductive Medicine* 1994;**39**:155.

Parazzini 1986

Parazzini F, Vecchia LL, Pampollona S. Parental age and risk of complete and partial hydatidiform mole. *British Journal of Obstetrics and Gynaecology* 1986;**93**:582–5.

Parazzini 1988

Parazzini F, La Vecchia C, Mangili G, Caminiti C, Negri E, Cecchetti G, et al. Dietary factors and risk of trophoblastic disease. *American Journal of Obstetrics and Gynecology* 1988;**158**(1):93–9.

Park 1996

Park TK, Kim SN, Lee SK. Analysis of risk factors for postmolar trophoblastic disease: categorization of risk factors and effect of prophylactic chemotherapy. *Yonsei Medical Journal* 1996;**37**(6):412–9.

Ratnam 1971

Ratnam SS, Teoh ES, Dawood MY. Methotrexate for prophylaxis of choriocarcinoma. *American Journal of Obstetrics and Gynecology* 1971;**111**:1021.

RevMan 2011 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.1. Copenhagen:

The Nordic Cochrane Centre, The Cochrane Collaboration, 2011.

Sasaki 2009

Sasaki S, Sasaki Y. Chapter 15: clinical management of trophoblastic diseases in Japan. In: Hancock BW, Seckl MJ, Berkowitz RS, Cole LA editor(s). *Gestational Trophoblastic Disease*. 3rd Edition. London: International Society for the Study of Trophoblastic Diseases, 2009:398-406. www.isstd.org/isstd/chapter15.html (accessed 21 August 2012).

Sebire 2009

Sebire NJ, Lindsay I, Paradinas F. Chapter 4: pathology. In: Hancock BW, Seckl MJ, Berkowitz RS, Cole LA editor(s). *Gestational Trophoblastic Disease*. 3rd Edition. London: International Society for the Study of Trophoblastic Diseases, 2009:97-147. www.isstd.org/isstd/chapter04.html (21 August 2012).

Seckl 2009

Seckl MJ. Chapter 10: presentation and management of persistent gestational trophoblastic disease (GTD) and gestational trophoblastic tumours/neoplasia (GTT/GTN) in the United Kingdom. In: Hancock BW, Seckl MJ, Berkowitz RS, Cole LA editor(s). *Gestational Trophoblastic Disease*. 3rd Edition. London: International Society for the Study of Trophoblastic Diseases, 2009:277-98. www.isstd.org/isstd/chapter10.html (accessed 21 August 2012).

Seckl 2010

Seckl MJ, Sebire NJ, Berkowitz RS. Gestational trophoblastic disease. *Lancet* 2010;**376**:717–29.

Smith 2003

Smith HO. Gestational trophoblastic disease epidemiology and trends. *Clinical Obstetrics and Gynecology* 2003;**46**(3): 541–56.

Song 1987

Song H, Wu P. Hydatidiform mole in China: a preliminary survey of incidence of more than three million women. *Bulletin of the World Health Organization* 1987;**65**:507–11.

Steigrad 2003

Steigrad SJ. Epidemiology of gestational trophoblastic diseases. *Best Practice and Research Clinical Obstetrics and Gynaecology* 2003;**17**:837–47.

Takeuchi 1987

Takeuchi S. Incidence of gestational trophoblastic disease by regional registration in Japan. *Human Reproduction* 1987;**2**: 729–34.

Tham 2003

Tham B, Everard J, Tidy J, Drew D, Hancock B. Gestational trophoblastic disease in the Asian population of Northern England and North Wales. *British Journal of Obstetrics and Gynaecology* 2003;**110**:555–9.

Tidy 2009

Tidy J. Chapter 18: the role of surgery in the management of gestational trophoblastic disease. In: Hancock BW, Seckl MJ, Berkowitz RS, Cole LA editor(s). *Gestational*

Trophoblastic Disease. 3rd Edition. London: International Society for the Study of Trophoblastic Diseases, 2009: 430-46. www.isstd.org/isstd/chapter18.html (accessed 21 August 2012).

* *Indicates the major publication for the study*

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Kashimura 1986

Methods	RCT conducted in Japan. Participants recruited between 1963 and 1977. Stated as a prospective study with participants 'selected at random' to receive prophylaxis This may not be a true RCT
Participants	420 women with molar pregnancy (low and high risk) Excluded women who were referred longer than 3 weeks after evacuation, those who had received other drugs for prophylaxis (see 'Risk of bias' table below), women who had undergone hysterectomy and women diagnosed as having partial mole or hydropic degeneration
Interventions	Arm 1: methotrexate 10 mg daily (IM or oral) for 7 days, within 3 weeks of evacuation (293 women) Arm 2: no P-Chem (127 women) Women were followed up weekly with urine hCG measurements
Outcomes	GTN diagnosed by histology or Ishizuka score (a risk rating system used in Japan); side effects and subsequent pregnancy
Notes	5- to 15-year follow-up reported. Time to invasive mole diagnosis was 56.8 days in P-Chem group and 42.7 days in control group (SD not given; P = 0.6). No attrition occurred for primary outcomes. Only reported adverse effects in the P-Chem group: 27.3% experienced drug-related side effects including stomatitis (10.3%), nausea/vomiting (6.8%) and leukopenia (4.4%). Grades of toxicity were not reported but the report states that there were no severe complications or drug-related deaths. Baseline characteristics were not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	'Selected at random' suggests that this was not a truly random process. No randomisation ratio was described and yet the 2 groups were very different sizes (293 vs. 127). The authors stated that 39 patients who received other drugs besides methotrexate were excluded from the study; this also suggests a non-random study design
Allocation concealment (selection bias)	Unclear risk	Not described

Kashimura 1986 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Attrition was low for the main outcomes but high for long-term outcomes such as subsequent pregnancies
Selective reporting (reporting bias)	High risk	Baseline characteristics of the 2 groups were not compared/reported
Other bias	High risk	More women in the control group were \geq 40 years old (22% vs. 11%) and progressed to GTN (39% vs. 12%)

Kim 1986

Methods	RCT conducted in Korea. Participants recruited between 1978 and 1984
Participants	133 women with complete hydatidiform mole (both high and low risk) were randomised into 2 groups, but 62 were excluded (36 lost to follow-up, 7 had 'insufficient length of follow-up' and 19 had hysterectomy) and only 71 completed this trial (39 in the treatment group and 32 in the untreated group)
Interventions	Arm 1: methotrexate 1.0 mg/kg/day IM on days 1, 3, 5 and 7 and citrovorum factor rescue 0.1 mg/kg/day IM on days 2, 4, 6 and 8 (39/71 women including 18/31 low-risk and 21/40 high-risk women). ERPC was done on the third or fourth day of P-Chem Arm 2: no treatment other than ERPC (32 women including 13/31 low-risk and 19/40 high-risk women)
Outcomes	Efficacy: incidence of GTN Adverse effects: incidence of gastrointestinal toxicity, myelotoxicity, epithelial toxicity including rash, hair loss and mouth ulcers The number of courses required to achieve remission in cases of GTN Time to GTN diagnosis Subsequent pregnancy
Notes	Baseline characteristics were similar between the groups, including the proportion of low- and high-risk lesions. ERPC was done on the third or fourth day of P-Chem. Women were followed up weekly for until hCG was normal for 3 consecutive weeks, then monthly for 6 months, then bimonthly for 6 months, then every 6 months. The mean duration of follow-up was 19 months (SD 9.7; range 6 to 50). All women were in complete remission at study closure Pregnancy rates after molar pregnancy were similar between the 2 groups (93% vs. 94%)

	P-Chem had little effect on the rate of subsequent GTN in the low-risk group; only 2/31 low-risk women developed GTN (1 women in each study group)	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	Of 133 women treated, 62 were excluded from the study (36 were lost to follow-up, 7 had insufficient length of follow-up and 19 had a hysterectomy). Therefore the outcome data were extracted from the 71 women (39 in the treatment group and 32 in the untreated group)
Selective reporting (reporting bias)	High risk	All the pre-specified outcomes were reported. However, certain women were excluded from the analyses (those who underwent hysterectomy and those with insufficient follow-up) therefore the analyses were not by intention-to-treat
Other bias	High risk	It is unclear on what basis the participants were initially diagnosed as having CM. If prophylaxis was given based on a clinical diagnosis before ERPC, this may have resulted in women with hydropic degeneration or PM being included in the study

Limpongsanurak 2001

Methods	RCT conducted in Thailand. Participants were recruited between 1989 and 1994
Participants	Women diagnosed with high-risk CM (with histological diagnosis) within 1 week of evacuation of molar tissue. Women were considered 'high risk' if they had at least 1 of the following characteristics: initial serum hCG > 100,000 mIU/mL; uterine size larger than dates; theca lutein cysts > 6 cm; age > 40 years; or associated medical and epidemiological factors including previous GTD, toxemia, hyperthyroidism, trophoblast embolisation or disseminated intravascular coagulation 60 participants were randomised into 2 groups (30:30)
Interventions	Arm 1: IV actinomycin D (10 µg/kg) for 5 days, within 1 week of evacuation of molar tissue Arm 2: IV fluids and analgesic drugs for 5 days within 1 week of evacuation of molar tissue
Outcomes	Efficacy: incidence of GTN Adverse effects: incidence of gastrointestinal toxicity, myelotoxicity, hair loss, mouth ulcers Time to diagnosis of GTN
Notes	The gestational age at diagnosis of HM was 13.8±3.0 weeks in the intervention group and 13.6±4.2 weeks in the control group. Women were followed up for 1 year with hCG assays every 2 weeks for 3 months, then monthly for 3 months, then every 2 months up to 1 year The diagnosis of GTN was made in all women in the P-Chem group (4/4) and 12/15 women in the control group according to the following criteria: rising hCG levels for 2 weeks or a plateau for 3 weeks; persistent or recurrent vaginal bleeding with detectable hCG levels; clinical or histological evidence of invasive mole, choriocarcinoma or metastases with persistently high or rising hCG values. Histology was obtained for 3 participants 2 out of 4 women in the P-Chem group and 3 out of 15 in the control group were lost to follow-up after diagnosis of GTN, therefore 5 women with GTN received no subsequent treatment and data were insufficient to compare the number of chemotherapy courses received in each group Side effects were reported as percentages and only recorded for the P-Chem group, as follows: stomatitis (10%), nausea/vomiting (10%), oral ulcers (3.3%) and hair loss (13.3%). All adverse effects were grade 1 except for 2 patients with patchy alopecia (grade 2)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random sequence was generated by lot-drawing (information obtained by e-mail correspondence with Dr Limpongsanurak)
Allocation concealment (selection bias)	Low risk	Sealed opaque envelopes (information obtained by e-mail correspondence with Dr Limpongsanurak)

Limpongsanurak 2001 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	Described as double-blind
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Described as double-blind, but the details of outcome assessment are unclear
Incomplete outcome data (attrition bias) All outcomes	Low risk	1 woman in the P-Chem group was lost to follow-up 1 month after treatment and not included in the primary outcome analysis
Selective reporting (reporting bias)	Low risk	All pre-specified outcomes were reported
Other bias	Low risk	Baseline characteristics and risk factors for disease progression were similar between the groups

CM: complete mole; ERPC: evacuation of retained products of conception; GTD: gestational trophoblastic disease; GTN: gestational trophoblastic neoplasia; hCG: β -human chorionic gonadotrophin; IM: intramuscular; IV: intravenous; P-Chem: prophylactic chemotherapy; RCT: randomised controlled trial; SD: standard deviation.

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Geng 2011	A retrospective study evaluating characteristics and outcomes for 23 women with high-risk HM who received prophylactic chemotherapy (5-FU or dactinomycin)
Uberti 2006	A retrospective study evaluating a bolus dose of dactinomycin for prevention of persistent GTD in 29 adolescents with high-risk molar pregnancy compared with a similar control group of 31 adolescents
Uberti 2009	A retrospective study evaluating the effect of a bolus dose of dactinomycin, given 1 hour before ERPC to women with high-risk HM, on the rate of malignant transformation to GTN

5-FU: 5-fluorouracil; ERPC: evacuation of retained products of conception; GTD: gestational trophoblastic disease; GTN: gestational trophoblastic neoplasia; HM: hydatidiform mole.

Characteristics of ongoing studies [ordered by study ID]

GOG 0242

Trial name or title	Second Curettage in Treating Patients With Persistent Non-Metastatic Gestational Trophoblastic Tumor
Methods	Multicentre Phase II study (NCT00521118)
Participants	Women with histologically confirmed gestational trophoblastic neoplasia (GTN) (complete or partial hydatidiform mole) with no histologically confirmed choriocarcinoma, placental site trophoblastic tumor (PSTT), or epithelioid trophoblastic tumour (ETT) on the first curettage. Persistent, low-risk disease (based on FIGO/WHO 2002 staging and risk scoring criteria), as defined by 1 of the following criteria: Less than 10% decline in beta-human chorionic gonadotropin (hCG) levels, based on four consecutive measurements over a 3-week period (plateau); Greater than 20% rise in beta-hCG levels, based on three consecutive measurements over a 2-week period; Beta-hCG level remains elevated above normal for ≥ 6 months. WHO risk score ≤ 6 . Must have a clinically significant elevated beta-hCG level of > 20 mIU/mL. No evidence of metastatic disease beyond the uterus by pelvic examination, pelvic ultrasound, and chest x-ray. No previously treated, persistent or recurrent GTN (same gestation) that have been treated with chemotherapy
Interventions	Patients undergo a second curettage rather than standard treatment (immediate chemotherapy). Patients whose disease has transformed into choriocarcinoma, placental site trophoblastic tumour, or epithelioid trophoblastic tumour (histologically diagnosed at the second curettage) are removed from the study. All other patients undergo weekly beta-human chorionic gonadotropin (hCG) testing beginning 14 days after the second curettage and continuing until the beta-hCG level is normal. Patients then undergo further beta-hCG testing weekly for 4 weeks and then monthly for 5 months. If the level does not regress to normal, or rises, or if metastatic disease is identified, the patient is removed from the study
Outcomes	Frequency of surgical cure, defined a normal beta-human chorionic gonadotropin (hCG) level documented for 6 consecutive months AND no chemotherapy. Development of choriocarcinoma, placental site trophoblastic tumour (PSTT), or epithelioid trophoblastic tumour (ETT) histologically diagnosed at second curettage. Development of "second persistent" disease, defined as failure to achieve or maintain a normal assay, or a plateau, or a rise in the assay level after second curettage. Frequency and severity of adverse effects of second curettage, specifically uterine operative injury, haemorrhage, and infection (pelvis, fallopian tubes, and ovaries), as assessed by CTCAE version 3.0
Starting date	October 2007
Contact information	Philip J. DiSaia, Gynecologic Oncology Group
Notes	

DATA AND ANALYSES

Comparison 1. Prophylactic chemotherapy versus no prophylactic chemotherapy


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Incidence of GTN (overall)	3	550	Risk Ratio (M-H, Random, 95% CI)	0.37 [0.24, 0.57]
1.1 Methotrexate prophylaxis	2	491	Risk Ratio (M-H, Random, 95% CI)	0.39 [0.24, 0.64]
1.2 Dactinomycin prophylaxis	1	59	Risk Ratio (M-H, Random, 95% CI)	0.28 [0.10, 0.73]
2 Incidence of GTN (high-risk HM only)	2	99	Risk Ratio (M-H, Random, 95% CI)	0.29 [0.14, 0.60]
2.1 Methotrexate prophylaxis	1	40	Risk Ratio (M-H, Random, 95% CI)	0.30 [0.10, 0.95]
2.2 Dactinomycin prophylaxis	1	59	Risk Ratio (M-H, Random, 95% CI)	0.28 [0.10, 0.73]
3 Time to GTN diagnosis	2	33	Mean Difference (IV, Random, 95% CI)	28.72 [13.19, 44.24]
3.1 Methotrexate prophylaxis	1	14	Mean Difference (IV, Random, 95% CI)	30.80 [12.93, 48.67]
3.2 Dactinomycin prophylaxis	1	19	Mean Difference (IV, Random, 95% CI)	22.30 [-9.05, 53.65]
4 Number of courses of chemotherapy to cure	1		Mean Difference (IV, Random, 95% CI)	Subtotals only
4.1 Methotrexate prophylaxis	1	14	Mean Difference (IV, Random, 95% CI)	1.1 [0.52, 1.68]
5 Mortality rate	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only

Analysis I.1. Comparison I Prophylactic chemotherapy versus no prophylactic chemotherapy, Outcome I Incidence of GTN (overall).

Review: Prophylactic chemotherapy for hydatidiform mole to prevent gestational trophoblastic neoplasia

Comparison: I Prophylactic chemotherapy versus no prophylactic chemotherapy

Outcome: I Incidence of GTN (overall)




Analysis 1.2. Comparison 1 Prophylactic chemotherapy versus no prophylactic chemotherapy, Outcome 2 Incidence of GTN (high-risk HM only).

Review: Prophylactic chemotherapy for hydatidiform mole to prevent gestational trophoblastic neoplasia

Comparison: 1 Prophylactic chemotherapy versus no prophylactic chemotherapy

Outcome: 2 Incidence of GTN (high-risk HM only)




Analysis 1.3. Comparison 1 Prophylactic chemotherapy versus no prophylactic chemotherapy, Outcome 3 Time to GTN diagnosis.

Review: Prophylactic chemotherapy for hydatidiform mole to prevent gestational trophoblastic neoplasia

Comparison: 1 Prophylactic chemotherapy versus no prophylactic chemotherapy

Outcome: 3 Time to GTN diagnosis




Analysis 1.4. Comparison 1 Prophylactic chemotherapy versus no prophylactic chemotherapy, Outcome 4 Number of courses of chemotherapy to cure.

Review: Prophylactic chemotherapy for hydatidiform mole to prevent gestational trophoblastic neoplasia

Comparison: 1 Prophylactic chemotherapy versus no prophylactic chemotherapy

Outcome: 4 Number of courses of chemotherapy to cure




Analysis 1.5. Comparison 1 Prophylactic chemotherapy versus no prophylactic chemotherapy, Outcome 5 Mortality rate.

Review: Prophylactic chemotherapy for hydatidiform mole to prevent gestational trophoblastic neoplasia

Comparison: 1 Prophylactic chemotherapy versus no prophylactic chemotherapy

Outcome: 5 Mortality rate



ADDITIONAL TABLES

Table 1. Risk scoring system for the prediction of GTN in women with molar pregnancy*

Prognostic factor	Score			
	0	1	2	3
U/S diagnosis	Partial	Complete	Recurrent	
Uterine size for GA (months)	not more than 1	> 1	> 2	> 3
hCG level (mIU/mL)	< 50,000	> 50,000 to < 100,000	> 100,000 to < 1,000,000	> 1,000,000
Diameter of theca lutein cysts (cm)	-	< 6	< 6 to < 10	> 10
Patient age (years)	-	< 20	≥ 40	> 50
Medical complications**	-	≥ 1	-	-

*From [Berkowitz 1987](#)

Low risk is defined as a score of < 4; high risk is defined as a score ≥ 4

U/S: ultrasound; GA: gestational age, hCG: β -human chorionic gonadotrophin.

** hyperemesis, hyperthyroidism, pre-eclampsia, trophoblastic embolisation, disseminated intravascular coagulation.

Table 2. Comparative studies of P-Chem for hydatidiform mole

Study	Design	Participants (P-Chem)	Participants (control/no P-Chem)	Intervention	Rate of GTN (P-Chem)	Rate of GTN (control)	Comments
Koga 1968*	Case-control	107 women (HM)	42 women (HM)	Methotrexate 10 mg/day PO x 7 days given within 3 weeks of ERPC	2/107 (2%)	4/42 (10%)	No choriocarcinoma observed in the P-Chem group vs. 3/42 in the control group. Toxic side effects occurred in 84/107 women, including stomatitis (34/107) and myelosuppression (22/107)

Table 2. Comparative studies of P-Chem for hydatidiform mole (Continued)

Goldstein 1971	Prospective case-control	73 women (CM)	116 women (CM)	3 intervention arms: methotrexate 0.3 mg/kg/day x 5 days (20 women); or dactinomycin 9-12 µg/kg/day x 5 days (53 women); ERPC on day 3	6/73 (8%)	23/116 (20%)	No metastatic disease observed in the P-Chem groups. P-Chem well tolerated with minor side effects
Goldstein 1974	Prospective case-control	100 women (HM)	100 women (HM)	Dactinomycin 12 µg/kg/day x 5 days. ERPC on day 3	2/100 (2%)	16/100 (16%)	No metastatic disease observed in the P-Chem group vs. 4/100 in the control group (4%). Reversible alopecia occurred in 32% of the P-Chem group. No serious toxic reactions
Goldstein 1981	Prospective case-control	174 women (CM)	858 women (CM)	Dactinomycin 12 µg/kg/day x 5 days. ERPC on day 3	10/247 (4%)	160/858 (19%)	No metastatic disease observed in the P-Chem group vs. 34/858 (4%) in the control group. This report includes data from Goldstein 1974
Fasoli 1982	Retrospective case-control	104 women (92% CM)	250 women (CM)	Methotrexate 10 mg/day PO x 5 days every 3 weeks for 3 cycles	3/104 (3%)	23/250 (9%)	Significantly fewer high-risk women in the P-Chem group

Table 2. Comparative studies of P-Chem for hydatidiform mole (Continued)

							(1/47) vs. the control group (18/126) developed GTN (2% vs. 14%; $P < 0.05$). 2 women had severe myelosuppression and 1 had severe alopecia
Kashimura 1986*	RCT (?)	293 women (CM)	127 women (CM)	Methotrexate 10 mg/day (IM or PO) for 7 days, within 3 weeks of evacuation	22/293 (7%)	23/127 (18%)	There were 5 cases of metastatic disease in each group (1.7% vs. 3.9%, respectively) 27.3% of the P-Chem group experienced drug-related side effects including stomatitis (10.3%), nausea/vomiting (6.8%) and leukopenia (4.4%). However none were reported to be severe
Kim 1986	RCT	39/71 women (CM; 18/31 low-risk and 21/40 high-risk women)	32 women (CM)	Methotrexate 1.0 mg/kg/day IM (days 1, 3, 5, 7) and citrovorum factor rescue 0.1 mg/kg/day IM (days 2, 4, 6, 8). ERPC on day 3	4/39 (10%)	10/32 (31%)	Significantly fewer high-risk women in the P-Chem group (14%) vs. the control group (47%) developed GTN. There was no significant difference in the GTN rates of

Table 2. Comparative studies of P-Chem for hydatidiform mole (Continued)

							low-risk women between groups
Park 1996	Retrospective case-control	52 women (14 low-risk, 21 medium-risk and 17 high-risk HM)	88 women (38 low-risk, 25 medium-risk and 25 high-risk HM)	Methotrexate 1 mg/kg (days 1, 3, 5, 7) and citrovorum factor (0.1 mg/kg (days 2, 4, 6, 8) ; or dactinomycin 12 µg/kg/day x 5 days started at the time of ERPC	8/52 (15.4%)	28/88 (31.8%)	Significantly fewer high-risk women in the P-Chem group (7/17) vs. the control group (22/25) developed GTN (41% vs. 88%; P < 0.01). There was no significant difference in the GTN rates in low- and medium-risk women between groups. The time to achieve normal hCG levels was shorter in high-risk women in the P-Chem group
Limpong-sa-nurak 2001*	Double-blind RCT	30 women (high-risk CM)	30 women (high-risk CM)	Dactinomycin 10 µg/kg for 5 days, within 1 week after ERPC and histology	4/29 (15.4%)	15/30 (50%)	Mild, reversible side effects reported including stomatitis (10%) , nausea/vomiting (10%) , oral ulcers (3.3%) and hair loss (13.3%) - all grade 1 except for 2 women with grade 2 patchy alopecia

Table 2. Comparative studies of P-Chem for hydatidiform mole (Continued)

Uberti 2006	Retrospective case-control	29 adolescents (high-risk CM)	31 adolescents (high-risk CM)	Dactinomycin 1. 25 mg/m ² IV given 1 hour before ERPC	2/29 (6.9%)	9/31 (29%)	Mean risk scores and hCG levels were significantly higher and gestational age was significantly lower in the P-Chem group than the control group. Mild and transient side effects included hepatotoxicity (10%) and mild alopecia (6.8%)
Uberti 2009	Retrospective case-control	163 women (high risk, > 90% CM)	102 women (high risk, > 90% CM)	Dactinomycin 1. 25 mg/m ² IV given 1 hour before ERPC	30/163 (18.4%)	35/102 (34.3%)	Mild and transient side effects including nausea (8%), raised liver enzymes (3.7%), stomatitis (3.1%), rash (2.4%), diarrhoea (2.4%), alopecia (1.2%) and neutropenia (0.6%) were seen in 21% of the P-Chem group. Time to GTN diagnosis, subsequent drug resistance and the number of chemotherapy course to cure was similar in the 2 groups

* Three studies administered P-Chem after ERPC including [Koga 1968](#), [Kashimura 1986](#) and [Limpongsanurak 2001](#).

CM; complete mole; ERPC: evacuation of retained products of conception; GTN: gestational trophoblastic neoplasia; HM: hydatidiform mole; IM: intramuscular; IV: intravenous; P-Chem; prophylactic chemotherapy; PO: per os; RCT: randomised controlled trial.

APPENDICES

Appendix 1. MEDLINE (Ovid Web) search strategy

1 Gestational Trophoblastic Disease/
2 exp Hydatidiform Mole/
3 (hydatid* adj2 mole*).mp.
4 (molar adj2 pregnanc*).mp.
5 1 or 2 or 3 or 4
6 exp Antineoplastic Agents/
7 Antineoplastic Combined Chemotherapy Protocols/
8 Chemoprevention/
9 (chemotherap* or chemoprophyla* or chemoprevention).mp.
10 (methotrexate or arnethopterin or dactinomycin or actinomycin D or fluorouracil or etoposide).mp.
11 6 or 7 or 8 or 9 or 10
12 5 and 11
13 randomized controlled trial.pt.
14 controlled clinical trial.pt.
15 randomized.ab.
16 placebo.ab.
17 clinical trials as topic.sh.
18 randomly.ab.
19 trial.ti.
20 13 or 14 or 15 or 16 or 17 or 18 or 19
21 12 and 20
22 exp animals/ not humans.sh.
23 21 not 22

Appendix 2. EMBASE search strategy

1 trophoblastic tumor/
2 hydatidiform mole/
3 (hydatid* adj2 mole*).mp.
4 (molar adj2 pregnanc*).mp.
5 1 or 2 or 3 or 4
6 exp chemotherapy/
7 exp antineoplastic agent/
8 chemoprophylaxis/
9 (chemotherap* or chemoprophyla* or chemoprevention).mp.
10 (methotrexate or arnethopterin or dactinomycin or actinomycin D or fluorouracil or etoposide).mp.
11 6 or 7 or 8 or 9 or 10
12 5 and 11
13 crossover procedure/
14 double-blind procedure/

15 randomized controlled trial/
 16 single-blind procedure/
 17 random*.mp.
 18 factorial*.mp.
 19 (crossover* or cross over* or cross-over*).mp.
 20 placebo*.mp.
 21 (double* adj blind*).mp.
 22 (singl* adj blind*).mp.
 23 assign*.mp.
 24 allocat*.mp.
 25 volunteer*.mp.
 26 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23 or 24 or 25
 27 12 and 26
 28 (exp Animal/ or Nonhuman/ or exp Animal Experiment/) not Human/
 29 27 not 28

Appendix 3. CENTRAL search strategy

#1 MeSH descriptor Gestational Trophoblastic Disease, this term only
 #2 MeSH descriptor Hydatidiform Mole explode all trees
 #3 hydatid* near/2 mole*
 #4 molar near/2 pregnanc*
 #5 (#1 OR #2 OR #3 OR #4)
 #6 MeSH descriptor Antineoplastic Agents explode all trees
 #7 MeSH descriptor Antineoplastic Combined Chemotherapy Protocols, this term only
 #8 MeSH descriptor Chemoprevention, this term only
 #9 (chemotherap* or chemoprophyla* or chemoprevention)
 #10 (methotrexate or arnethopterin or dactinomycin or actinomycin D or fluorouracil or etoposide)
 #11 (#6 OR #7 OR #8 OR #9 OR #10)
 #12 (#5 AND #11)

WHAT'S NEW

Last assessed as up-to-date: 10 August 2012.

Date	Event	Description
21 September 2016	Amended	Contact details updated.

HISTORY

Protocol first published: Issue 3, 2008

Review first published: Issue 10, 2012

Date	Event	Description
1 April 2015	Amended	Contact details updated.
24 February 2015	Amended	Contact details updated.
11 February 2015	Amended	Contact details updated.
27 March 2014	Amended	Contact details updated.

CONTRIBUTIONS OF AUTHORS

Jing Fu, Fan He: protocol development, methodological quality assessment, retrieval of papers, data extraction, data analysis and writing the review.

Lingxia Xie, Hengxi Chen: protocol development, methodological quality assessment, data extraction and revision of the review.

Fang Fang, Taixiang Wu, Hu Lina: protocol development, content expert and revision of the review.

Tess Lawrie: methodology, data extraction, data analysis and writing.

DECLARATIONS OF INTEREST

None.

SOURCES OF SUPPORT

Internal sources

- West China Secondary Hospital, Sichuan University, China.

External sources

- This review received methodological and statistical support as part of the 10/4001/12 NIHR Cochrane Programme Grant Scheme - Optimising care, diagnosis and treatment pathways to ensure cost effectiveness and best practice in gynaecological cancer: improving evidence for the NHS, UK.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

None.

INDEX TERMS

Medical Subject Headings (MeSH)

Antineoplastic Agents [*therapeutic use]; Dactinomycin [*therapeutic use]; Gestational Trophoblastic Disease [*prevention & control]; Hydatidiform Mole [*drug therapy]; Methotrexate [*therapeutic use]; Randomized Controlled Trials as Topic

MeSH check words

Female; Humans; Pregnancy