



Cochrane
Library

Cochrane Database of Systematic Reviews

Anaesthetic interventions for prevention of awareness during surgery (Review)

Messina AG, Wang M, Ward MJ, Wilker CC, Smith BB, Vezina DP, Pace NL

Messina AG, Wang M, Ward MJ, Wilker CC, Smith BB, Vezina DP, Pace NL.
Anaesthetic interventions for prevention of awareness during surgery.
Cochrane Database of Systematic Reviews 2016, Issue 10. Art. No.: CD007272.
DOI: [10.1002/14651858.CD007272.pub2](https://doi.org/10.1002/14651858.CD007272.pub2).

www.cochranelibrary.com

TABLE OF CONTENTS

ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	10
OBJECTIVES	10
METHODS	10
RESULTS	12
Figure 1.	13
Figure 2.	16
Figure 3.	17
Figure 4.	23
Figure 5.	24
Figure 6.	25
DISCUSSION	26
AUTHORS' CONCLUSIONS	28
ACKNOWLEDGEMENTS	28
REFERENCES	29
CHARACTERISTICS OF STUDIES	56
DATA AND ANALYSES	304
Analysis 1.1. Comparison 1 Anaesthesia depth monitors (BIS and M-entropy) versus standard clinical and electronic monitoring, Outcome 1 Definite and possible awareness (Grade 4).	304
Analysis 2.1. Comparison 2 Anaesthesia depth monitors (BIS) versus standard clinical and electronic monitoring, Outcome 1 Definite and possible awareness (Grade 4).	305
Analysis 2.2. Comparison 2 Anaesthesia depth monitors (BIS) versus standard clinical and electronic monitoring, Outcome 2 Definite awareness (Grade 4).	305
Analysis 3.1. Comparison 3 Thiopentone with and without added hypnotic drugs (ketamine, etomidate), Outcome 1 Wakefulness (IFT studies) (Grades 4, 5).	306
Analysis 4.1. Comparison 4 Thiopentone and ketamine vs benzodiazepine anaesthetic techniques, Outcome 1 Awareness (Grade 4).	307
Analysis 5.1. Comparison 5 Caesarean section low- and high-dose inhaled agents, Outcome 1 Awareness (Grade 4).	307
ADDITIONAL TABLES	307
APPENDICES	308
CONTRIBUTIONS OF AUTHORS	333
DECLARATIONS OF INTEREST	333
SOURCES OF SUPPORT	334
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	334
INDEX TERMS	335

[Intervention Review]

Anaesthetic interventions for prevention of awareness during surgery

Anthony G Messina¹, Michael Wang², Marshall J Ward³, Chase C Wilker⁴, Brett B Smith⁵, Daniel P Vezina^{6,7}, Nathan Leon Pace⁸

¹The Alliance for Medical Management Education, School of Management, University of Texas at Dallas, Richardson, TX, USA. ²Clinical Psychology Unit, University of Leicester, Leicester, UK. ³Dartmouth-Hitchcock Medical Center, Lebanon, NH, USA. ⁴Clinical Toxicology III, ARUP Laboratories, Salt Lake City, UT, USA. ⁵University of Utah School of Medicine, University of Utah, Salt Lake City, UT, USA.

⁶Department of Anesthesiology, Department of Internal Medicine, Division of Cardiology, University of Utah, Salt Lake City, UT, USA.

⁷Echocardiography Laboratory, Veteran's Administration, Salt Lake City, UT, USA. ⁸Department of Anesthesiology, University of Utah, Salt Lake City, UT, USA

Contact: Anthony G Messina, The Alliance for Medical Management Education, School of Management, University of Texas at Dallas, Box 233, 1920 N. Coit Road, Richardson, TX, 75080, USA. awareness.research@gmail.com.

Editorial group: Cochrane Anaesthesia Group.

Publication status and date: New, published in Issue 10, 2016.

Citation: Messina AG, Wang M, Ward MJ, Wilker CC, Smith BB, Vezina DP, Pace NL. Anaesthetic interventions for prevention of awareness during surgery. *Cochrane Database of Systematic Reviews* 2016, Issue 10. Art. No.: CD007272. DOI: [10.1002/14651858.CD007272.pub2](https://doi.org/10.1002/14651858.CD007272.pub2).

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

General anaesthesia is usually associated with unconsciousness. 'Awareness' is when patients have postoperative recall of events or experiences during surgery. 'Wakefulness' is when patients become conscious during surgery, but have no postoperative recollection of the period of consciousness.

Objectives

To evaluate the efficacy of two types of anaesthetic interventions in reducing clinically significant awareness:

- anaesthetic drug regimens; and
- intraoperative anaesthetic depth monitors.

Search methods

We searched the Cochrane Central Register of Controlled Trials (CENTRAL, ISSUE 4 2016); PubMed from 1950 to April 2016; MEDLINE from 1950 to April 2016; and Embase from 1980 to April 2016. We contacted experts to identify additional studies. We performed a handsearch of the citations in the review. We did not search trial registries.

Selection criteria

We included randomized controlled trials (RCTs) of either anaesthetic regimens or anaesthetic depth monitors. We excluded volunteer studies, studies of patients prior to skin incision, intensive care unit studies, and studies that only randomized different word presentations for memory tests (not anaesthetic interventions).

Anaesthetic drug regimens included studies of induction or maintenance, or both. Anaesthetic depth monitors included the Bispectral Index monitor, M-Entropy, Narcotrend monitor, cerebral function monitor, cerebral state monitor, patient state index, and lower oesophageal contractility monitor. The use of anaesthetic depth monitors allows the titration of anaesthetic drugs to maintain unconsciousness.

Data collection and analysis

At least two authors independently scanned abstracts, extracted data from the studies, and evaluated studies for risk of bias. We made attempts to contact all authors for additional clarification. We performed meta-analysis statistics in packages of the R language.

Main results

We included 160 studies with 54,109 enrolled participants; 53,713 participants started the studies and 50,034 completed the studies or data analysis (or both). We could not use 115 RCTs in meta-analytic comparisons because they had zero awareness events. We did not merge 27 of the remaining 45 studies because they had excessive clinical and methodological heterogeneity. We pooled the remaining 18 eligible RCTs in meta-analysis. There are 10 studies awaiting classification which we will process when we update the review.

The meta-analyses included 18 trials with 36,034 participants. In the analysis of anaesthetic depth monitoring (either Bispectral Index or M-entropy) versus standard clinical and electronic monitoring, there were nine trials with 34,744 participants. The overall event rate was 0.5%. The effect favoured neither anaesthetic depth monitoring nor standard clinical and electronic monitoring, with little precision in the odds ratio (OR) estimate (OR 0.98, 95% confidence interval (CI) 0.59 to 1.62).

In a five-study subset of Bispectral Index monitoring versus standard clinical and electronic monitoring, with 34,181 participants, 503 participants gave awareness reports to a blinded, expert panel who adjudicated or judged the outcome for each patient after reviewing the questionnaires: no awareness, possible awareness, or definite awareness. Experts judged 351 patient awareness reports to have no awareness, 87 to have possible awareness, and 65 to have definite awareness. The effect size favoured neither Bispectral Index monitoring nor standard clinical and electronic monitoring, with little precision in the OR estimate for the combination of definite and possible awareness (OR 0.96, 95% CI 0.35 to 2.65). The effect size favoured Bispectral Index monitoring for definite awareness, but with little precision in the OR estimate (OR 0.60, 95% CI 0.13 to 2.75).

We performed three smaller meta-analyses of anaesthetic drugs. There were nine studies with 1290 participants. Wakefulness was reduced by ketamine and etomidate compared to thiopental. Wakefulness was more frequent than awareness. Benzodiazepines reduces awareness compared to thiopental, ketamine, and placebo. Also, higher doses of inhaled anaesthetics versus lower doses reduced the risk of awareness.

We graded the quality of the evidence as low or very low in the 'Summary of findings' tables for the five comparisons.

Most of the secondary outcomes in this review were not reported in the included RCTs.

Authors' conclusions

Anaesthetic depth monitors may have similar effects to standard clinical and electrical monitoring on the risk of awareness during surgery. In older studies comparing anaesthetics in a smaller portion of the patient sample, wakefulness occurred more frequently than awareness. Use of etomidate and ketamine lowered the risk of wakefulness compared to thiopental. Benzodiazepines compared to thiopental and ketamine, or higher doses of inhaled anaesthetics versus lower doses, reduced the risk of awareness.

PLAIN LANGUAGE SUMMARY

Methods to prevent people waking during surgery and remembering surgical events

Key question

We reviewed the evidence about the use of devices to adjust the amount of drugs given during anaesthesia to prevent premature waking up. We also reviewed the evidence about the choice of drugs used during anaesthesia to prevent premature waking up.

Background

Anaesthesia is the use of drugs to render a patient unconscious for painful procedures and surgery. Being anaesthetized is not the same as being asleep. Someone sleeping may be easily awakened. Someone anaesthetized should only be allowed to awake when the surgery or procedure is completed. A very small percentage of patients may wake up during anaesthesia and surgery; this is called wakefulness. Patients usually do not remember being awake after emerging from anaesthesia. However, an even smaller percentage of patients do remember or recall events from surgery afterwards. This memory is called an awareness event. If that memory is distressing, it can impair the individual's quality of life.

New devices known as anaesthetic depth monitors are being used to monitor the patient's brainwave response to anaesthetic drugs. Anaesthetic depth monitors have been compared to the usual clinical observations (e.g. fast heart rate, tearing, movement, etc.) during surgery to adjust the amount of drugs given and reduce the risk of wakefulness and awareness.

Anaesthetic drugs have many different effects on brain function. Some drugs are used alone as the sole anaesthetic. Other drugs have insufficient effect to be used as a sole anaesthetic, but are used in combination with more powerful drugs. Drugs may have different risks of the patient waking up prematurely.

Search date

The evidence is current to April 2016.

Study characteristics

We found 160 randomized controlled trials with 54,109 participants. Eighteen studies with 36,034 participants contributed evidence about devices and drugs to prevent premature waking up during surgery. Nine studies compared anaesthetic depth monitoring versus other methods to adjust drugs. Nine studies compared different drugs. There are 10 studies awaiting classification, which we will process when we update the review.

Key results

In the largest studies of anaesthetic depth monitors (five studies with 31,181 participants) there were 152 participants with possible or definite awareness (recall of surgery events after surgery). The use of anaesthetic depth monitors to adjust drugs during anaesthesia may have similar effects on the risk of awareness when compared with standard clinical and electrical monitoring. Wakefulness is reduced by ketamine and etomidate compared to thiopental. Benzodiazepines reduces awareness compared to thiopental, ketamine, and placebo. Also higher doses of inhaled anaesthetics versus lower doses reduced the risk of awareness.

Quality of evidence

The quality of the evidence was low or very low because the studies the results were not similar across studies, and there were not enough data.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. Anaesthesia depth monitors (BIS and M-entropy) versus standard clinical and electronic monitoring

Anaesthesia depth monitors (BIS and M-entropy) compared with standard clinical and electronic monitoring

Patient or population: patients with prevention of recall of events during surgery

Settings: All patients undergoing various surgical procedures in hospitals in Europe/Australia/Asia/Middle East/North America

Intervention: anaesthesia depth monitors (BIS, M-Entropy)

Comparison: standard clinical and electronic monitoring

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Standard clinical and electronic monitoring	Anaesthesia depth monitors				
Awareness Postoperative interview Follow-up: 1 to 72 days	5 per 1000	5 per 1000 (3 to 7)	OR 0.98 (0.59 to 1.62)	34,744 (9 studies)	⊕⊕⊕⊖ low ^{1,2}	—
Adverse effects of intraoperative wakefulness and/or postoperative awareness (i.e. post-traumatic stress syndrome, myocardial infarction, cardiac arrest, etc.)	—	—	—	—	—	Not defined or not identified

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

BIS: Bispectral Index; **CI:** confidence interval; **OR:** odds ratio

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹Inconsistency: downgraded one level for inconsistency of effect. Heterogeneity (I^2) was moderate (49%). There were non-overlapping 95% CIs.

2Imprecision: downgraded one level for imprecision. Although the number of participants was large (34,744), the number of events was small (173) and the upper and lower bounds of the OR 95% CI did not exclude important effects.

Summary of findings 2. Anaesthesia depth monitors (BIS) versus standard clinical and electronic monitoring

Anaesthesia depth monitors (BIS) versus standard clinical and electronic monitoring with expert panel adjudication (Grade 4 and 5) during surgery

Patient or population: patients with prevention of recall of events during surgery

Settings: All patients undergoing various surgical procedures in hospitals in Europe/Australia/Asia/Middle East/North America

Intervention: anaesthesia depth monitors (BIS)

Comparison: standard clinical and electronic monitoring

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Standard clinical and electronic monitoring	Anaesthesia depth monitors (BIS)				
Awareness: definite and possible Postoperative interview Follow-up: 1 to 72 days ⁵	5 per 1000	5 per 1000 (3 to 8)	OR 0.96 (0.35 to 2.65)	34,181 (5 studies)	⊕⊕○○ low 1,2	—
Awareness: definite Postoperative interview Follow-up: 1 to 72 days ⁶	2 per 1000	1 per 1000 (0 to 4)	OR 0.60 (0.13 to 2.75)	34,181 (5 studies)	⊕⊕○○ low 3,4	—
Adverse effects of intraoperative wakefulness and/or postoperative awareness (i.e. post-traumatic stress syndrome, myocardial infarction, cardiac arrest, etc.)	—	—	—	—	—	Not defined or not identified

*The basis for the **assumed risk** (e.g. the average control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

BIS: Bispectral Index; **CI:** confidence interval; **OR:** odds ratio

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

- 1Inconsistency: BIS definite and possible downgraded one level for inconsistency of effect. Heterogeneity (I^2) was substantial (68%). There were non-overlapping 95% CIs.
- 2Imprecision: BIS definite and possible downgraded one level for imprecision. Although the number of participants was large (34,181), the number of events was small (152) and the upper and lower bounds of the OR 95% CI did not exclude important effects.
- 3Inconsistency: BIS definite downgraded one level for inconsistency of effect. Heterogeneity (I^2) was substantial (60%). There were non-overlapping 95% CI.
- 4Imprecision: BIS definite downgraded one level for imprecision. Although the number of participants was large (34,181), the number of events was small (64) and the upper and lower bounds of the OR 95% CI did not exclude important effects.
- 5The assumed risk is the average control group event rate in the five studies: 0.0047 (78/16,749).
- 6The assumed risk is the average control group event rate in the five studies: 0.0024 (41/16,749).

Summary of findings 3. Thiopentone with and without added hypnotic drugs (ketamine, etomidate)

Thiopentone with and without added hypnotic drugs (ketamine, etomidate)

Patient or population: patients with prevention of recall of events during surgery

Settings: All patients undergoing various surgical procedures in hospitals in Europe/Australia/Asia/Middle East/North America

Intervention: anaesthetic drugs introduced after thiopentone for prevention of recall of events during surgery

Comparison: Thiopentone with and without added hypnotic drugs (ketamine, etomidate)

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Thiopentone	Anaesthetic drugs introduced after thiopentone for prevention of recall of events during surgery: Ketamine, etomidate				
Intraoperative Wakefulness Postoperative Interview Follow-up: 0 to 7 days	552 per 1000	99 per 1000 (50 to 226)	RR 0.18 (0.09 to 0.41)	141 (3 studies)	⊕⊕⊕⊕ low 1,2	—
Adverse effects of intraoperative wakefulness and/or postoperative awareness (i.e. post-traumatic stress syndrome, myocardial infarction, cardiac arrest, etc.)	—	—	—	—	—	Not defined or not identified

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **RR:** risk ratio

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹Within-study risk of bias: downgraded one level.

²Imprecision of results: downgraded one level for imprecision of effect. The high proportion of wakefulness events to sample size in these small studies was the reason for the one level downgrade compared to two levels for Comparison 4 and 5. The optimal information size threshold cannot be reached.

Summary of findings 4. Thiopentone and ketamine versus benzodiazepines (diazepam, midazolam, lorazepam)

Thiopentone and ketamine versus benzodiazepines (diazepam, midazolam, lorazepam)

Patient or population: patients with prevention of recall of events during surgery

Settings: All patients undergoing various surgical procedures in hospitals in Europe/Australia/Asia/Middle East/North America

Intervention: benzodiazepine use versus other intravenous anaesthetic techniques

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control: Ketamine, thiopentone	Benzodiazepine use versus other (control) intravenous anaesthetic techniques				
Awareness Postoperative interview Follow-up: 1 to 2 days	162 per 1000	27 per 1000 (3 to 40)	RR 0.17 (0.02 to 0.25)	291 (4 studies)	⊕⊕⊕⊕ very low ^{1,2}	—
Adverse effects of intraoperative wakefulness and/or postoperative awareness (i.e. post-traumatic stress syndrome, myocardial infarction, cardiac arrest, etc.)	—	—	—	—	—	Not defined or not identified

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **RR:** risk ratio

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹Within-study risk of bias: downgraded one level.

²Imprecision of results: downgraded two levels for imprecision of effect. The optimal information size threshold cannot be reached.

³The assumed risk is the average control group event rate in the four studies: 0.162 (16/99).

Summary of findings 5. Caesarean section with low- and high-dose inhaled agent

Caesarean section with low- and high-dose inhaled agent

Patient or population: patients with decreasing recall of events during surgery

Settings: All patients undergoing various surgical procedures in hospitals in Europe/Australia/Asia/Middle East/North America

Intervention: deeper anaesthetic techniques

Comparison: lighter anaesthetic techniques

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Lighter anaesthetic techniques	Deeper anaesthetic techniques				
Awareness Postoperative interview ¹ Follow-up: 1 day	51 per 1000	7 per 1000 (2 to 22)	RR 0.13 (0.04 to 0.43)	858 (2 studies)	⊕○○○ very low ^{1,2}	—
Adverse effects of intraoperative wakefulness and/or postoperative awareness (i.e. post-traumatic stress syndrome, myocardial infarction, cardiac arrest, etc.)	—	—	—	—	—	Not defined or not identified

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **RR:** risk ratio

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.
Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.
Very low quality: We are very uncertain about the estimate.

¹Within-study risk of bias: downgraded one level.

²Imprecision of results: downgraded two levels for imprecision of effect. The optimal information size threshold cannot be reached.

³The assumed risk is the average control group event rate in the two studies: 0.051 (21/409).

BACKGROUND

Description of the condition

The American Society of Anesthesiology (ASA) task force has defined awareness as “when a patient becomes conscious during a procedure performed under general anaesthesia and subsequently has recall of these events” (ASA 2006). Recent estimates of the number of patients having awareness under general anaesthesia in the United States have been as high as approximately 1 to 2 per 1000 anaesthetics (Sebel 2004).

People may wake up far more often during surgery than they remember after surgery (wakefulness) (Artusio 1955; Russell 1985; Russell 1993; Tunstall 1977; Appendix 1). Wakefulness and awareness cases that are not associated with pain or distress are considered, by some, to be clinically insignificant. There are no published data that define the frequency of wakefulness or awareness cases that are associated with pain and or panic. It has been reported that 10% of awareness cases are associated with pain (Jones 1994).

Awareness with pain or panic, or both, almost exclusively occurs with the use of neuromuscular blocking drugs (Cundy 1995; Lennmarken 2002; Mainzer 1979; Sandin 2000; White 1987). Neuromuscular blocking drugs in current use are also associated with significant complications other than awareness (Brull 2008; Murphy 2008; Wahl 2011).

The psychological sequelae of cases of anaesthetic awareness have been described (Appendix 1).

Description of the intervention

Two types of interventions have been proposed for preventing wakefulness and awareness: medication and specialized monitoring.

Medication

Different medications may be administered before and during general anaesthesia; these may have oral, intramuscular, intravenous, or volatile routes of administration. They include anaesthetic gases and vapours, sedatives or hypnotics, and analgesics. The choice may consist solely of the different medications that can be administered prior to the induction of anaesthesia. However, management options may also include two different protocols for general anaesthesia:

- techniques consisting of intravenous drugs only;
- traditional general anaesthesia combining both volatile anaesthetics and intravenous drugs.

Specialized monitoring

Modern anaesthesia depth monitors provide a real time electroencephalogram (EEG) for the patient in the operating theatre. Various algorithms are applied, i.e. the EEG is processed. The processed EEG is usually displayed as a unit less number scaled from 0 to 100. The value displayed is updated frequently (within seconds) with newly acquired EEG epochs. Calibration has been established by anaesthesia depth monitor manufacturers, with a value of 100 reflecting an awake state and a value of 0 reflecting cerebral electrical silence and unconsciousness (Rampil 1998). The use of anaesthesia depth monitors involves two stages.

Firstly, the monitor is placed on the patient in order to generate monitoring data. Secondly, the anaesthetist uses these data to make adjustments to the anaesthetic management of the patient.

How the intervention might work

Both types of intervention, choice of medication and use of anaesthesia depth monitors, have a common expected pathway for reducing the risk of awareness. Both are expected to produce a more intense (deeper) anaesthetic state through the selection of the type or amount of medication.

Why it is important to do this review

Patients who wake up during surgery may experience pain and distress. Besides the immediate suffering during the surgery itself, there maybe longer-term consequences. Awareness and possibly wakefulness can lead to post-traumatic stress disorder, which is a serious condition that can greatly impair quality of life. It may also cause patients to delay follow-up medical care. In some patients it may be sufficiently severe to lead to suicide (Breen 2007).

OBJECTIVES

To evaluate the efficacy of two types of anaesthetic interventions in reducing clinically significant awareness.

- anaesthetic drug regimens; and
- intraoperative anaesthetic depth monitors.

METHODS

Criteria for considering studies for this review

Types of studies

We included randomized controlled trials (RCTs).

Types of participants

We included paediatric and adult patients having all types of surgery. We excluded volunteer studies, studies of patients prior to skin incision, intensive care unit studies, and studies that only randomized different word presentations for memory tests (not anaesthetic interventions).

Types of interventions

Anaesthetic drug regimens included studies of induction or maintenance, or both. We included two types of interventions:

- anaesthetic drug regimens;
- intraoperative anaesthetic depth monitors.

Anaesthetic drug interventions included those during premedication, during induction and/or during maintenance phases of anaesthesia. Intraoperative anaesthetic depth monitors are those instruments that putatively allow anaesthesiologists to monitor the level of unconsciousness.

Types of outcome measures

Sebel et al introduced a classification system to improve the categorization of awareness complications (Sebel 2004). It was argued that more objective criteria were needed to identify a valid patient awareness report (Eger 2005). Therefore Wang and colleagues proposed a further refinement of this classification

of awareness (Wang 2012). This classification includes a six-level hierarchy from unconsciousness to consciousness that includes wakefulness, awareness, explicit and implicit memory, post-traumatic stress disorder (PTSD), and perioperative dreams and nightmares (Table 1; Appendix 1).

Primary outcomes

- Awareness or wakefulness as defined using the awareness classification system in Table 1.

The classification used in this review had not been conceived or published prior to 2012 (Wang 2012), therefore studies may not have adhered to these criteria for the determination of intraoperative state. We therefore had to infer this from study descriptions and our author survey.

We tallied details of the adjudication process from the published study or sought the details by communication with the authors. In some trials a formal two-stage process using structured interviews of patients was conducted. Potential awareness episodes were recorded in a narrative report. An independent committee of three anaesthetists, blinded to treatment group, coded the report as no awareness, possible awareness, and awareness.

Secondary outcomes

- Signs or adverse effects of intraoperative wakefulness or postoperative awareness, or both, are intraoperative patient movement, haemodynamic changes, portions of intraoperative dreams and postoperative dreams or nightmares, or both, delayed memory, full (PTSD) or partial (PTS) forms of post-traumatic stress syndrome, myocardial infarction, cardiac arrest, death, and suicide.

Search methods for identification of studies

Electronic searches

We searched the Cochrane Central Register of Controlled Trials (CENTRAL, Issue 4, 2016), PUBMED (1950 to April 2016) and MEDLINE (1950 to April 2016), and Embase (1980 to April 2016). We searched CENTRAL using the search terms found in Appendix 2. We searched MEDLINE using the search terms described in Appendix 3 and Embase using the terms found in Appendix 4, via SilverPlatter.

Searching other resources

We contacted experts in the field to identify any additional studies. We performed a handsearch from the citations found in included and excluded studies and other footnoted papers. We did not impose any language restriction. We did not search trial registries (ClinicalTrials.gov; World Health Organization (WHO) International Clinical Trials Registry Platform (ICTRP)).

Data collection and analysis

A group of the authors (AGM, NLP, MaW, CCW, and BS) independently scanned the titles and abstracts of reports identified by electronic and manual searching and by contact with experts. We evaluated full-text versions of potentially relevant studies. We used the web app Rayaan to assist in citation review (Elmagarmid 2014).

Selection of studies

A group of the authors (AGM, NLP, MaW, CCW, and BS) independently selected trials that met the inclusion criteria

by using a checklist designed for that purpose. Where there was disagreement we discussed the differences and reached a consensus. The criteria used to merge included studies in meta-analysis are described in Appendix 5.

Data extraction and management

A group of the authors (AGM, NLP, MaW, CCW, and BS) independently extracted parts of the data using a standardized study record form (see Appendix 5).

Assessment of risk of bias in included studies

A group of the authors (AGM, NLP, MaW, CCW, and BS) used Cochrane's domain-based evaluation method for assessing the risk of bias (Guyatt 2011; Higgins 2011). The assessment of risk of bias was based on the seven domains in the 'Risk of bias' table: random sequence generation (selection bias), allocation concealment (selection bias), blinding of participants, blinding of personnel (performance bias), blinding of outcome assessment (detection bias), incomplete outcome data (attrition bias), selective reporting (reporting bias), and other bias. We sent a survey to study authors to inquire about study methodology (Appendix 6). We sent this survey to authors who had email addresses in the included paper or that were readily available in a literature search of other published papers.

Measures of treatment effect

The definitions of rare and uncommon adverse events and diseases overlap (EUC 2009; HR4013; WHO 2002). Therefore, we use the terms 'rare' and 'uncommon' interchangeably in this review. Awareness is a dichotomous outcome and an uncommon or rare event. We used relative event rates, odds ratio (OR) or risk ratio (RR), as the effect size measure.

GRADE and 'Summary of findings' table methods

We used the GRADE methods to assess the quality of the body of evidence associated with specific outcomes: awareness - definite and definite and possible awareness, intraoperative wakefulness, and adverse effects of intraoperative wakefulness and/or postoperative awareness (i.e. post-traumatic stress syndrome, myocardial infarction, cardiac arrest, etc.), and to assess the quality of the evidence for the five comparisons of merged studies (Guyatt 2008). We used five domains to downgrade the quality of evidence: risk of bias, inconsistency, indirectness, imprecision, and publication bias. We used three domains to upgrade the quality of evidence: large effect, plausible confounding that could change the effect, and dose response gradient.

Unit of analysis issues

For studies of both anaesthetic drug interventions and anaesthetic depth monitors, we analysed participants in the groups to which they were randomized regardless of the actual intervention delivered, i.e. by intention-to-treat.

The unit of analysis for all outcomes was the individual participant. We defined the experimental group in a comparison as the newer treatment in each study and the control group as the older treatment.

All studies were parallel-group trials. There were no unit of analysis issues such as those arising from cross-over or cluster-randomized trials.

Dealing with missing data

To investigate the consequence of missing data, we considered using best-case or worst-case imputation. We considered including and excluding any study that appeared to have a large effect size (often the largest or earliest study) in order to assess its impact on the meta-analysis.

We described the missing data that resulted from such factors as attrition or exclusions, or both (see [Characteristics of included studies](#)). We calculated the number and percentage of missing data from each outcome group of the included studies and based our grade on the authors' account of those missing data.

Assessment of heterogeneity

We expected to find a great deal of clinical heterogeneity in the included studies, for example anaesthetic types, patient ages, etc. We considered this clinical heterogeneity when deciding whether to pool results in a meta-analysis.

To assess statistical heterogeneity, we used the I^2 statistic, which describes the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance). The importance of the observed value of I^2 depends on the magnitude and direction of effects and the strength of evidence for heterogeneity. The thresholds for interpreting I^2 heterogeneity were: moderate (30% to 60%), substantial (50% to 90%), and considerable (75% to 100%) ([Higgins 2002](#)).

Assessment of reporting biases

There were insufficient studies of the same interventions (fewer than 10) to allow the creation of funnel plots.

Data synthesis

As [Review Manager](#) has no statistical methods for very rare and uncommon events with between-study heterogeneity, we used the statistical packages in R ([R 2016](#)). For ORs, we estimated random- or mixed-effects generalized linear models in the R package [metafor](#) ([Viechtbauer 2010](#)). In some comparisons, the sparseness of data prevented logistic model estimation. In those cases, we estimated an exact fixed-effect RR ([Tian 2008](#)), in the R package [exactmeta](#) ([Yu 2013](#)). We also estimated a random-effects multivariable meta-analysis comparing the logit transformed proportions of awareness versus wakefulness in the R package [metafor](#). We declared statistical significance for $P < 0.05$ and if the 95% CI for effect sizes failed to include the line of unity. Results from the random- or mixed-effects model included an approximate 95% prediction interval (PI); this interval estimated where 95% of true outcomes fall in the hypothetical population of all possible studies ([Dean 2007](#); [Tian 2008](#)).

Results from individual studies are displayed in forest plots. Summary statistics from non-[Review Manager](#) packages are displayed in the forest plots. Statistical analyses including data sets, codes, and outputs are shown in an appendix ([Appendix 7](#)).

Subgroup analysis and investigation of heterogeneity

We considered the following subgroup analyses:

- anaesthesia depth monitoring method;
- patients at high risk of awareness;
- specific anaesthetic techniques with and without neuromuscular blocking drugs;
- implicit or explicit memory.

In a subgroup analysis, the magnitude and direction of treatment effect may be inconsistent among subgroups.

There were only sufficient data for a comparison of subgroups based on studies with high risk of awareness and alarms/alerts and total intravenous anaesthesia (TIVA).

Sensitivity analysis

We considered performing a sensitivity analysis to assess the impact of studies with a higher risk of bias. The patients lost to follow-up in these studies may be associated with higher rates of awareness, since one important sequela of awareness is a phobic avoidance of hospitals and physicians. In the context of trials, this is likely to lead to dropout during follow-up.

We did not perform a sensitivity analysis for risk of bias for the following reasons:

- [Analysis 1.1](#), [Analysis 2.1](#), and [Analysis 2.2](#) had predominantly low risk of bias for all seven domains and therefore were not suitable for risk of bias sensitivity analysis;
- [Analysis 3.1](#), [Analysis 4.1](#), and [Analysis 5.1](#) were candidates for a sensitivity analysis because of the predominantly unclear risk of bias for [Analysis 3.1](#) and the low to unclear risk of bias for [Analysis 4.1](#) and [Analysis 5.1](#). Nonetheless, we did not conduct sensitivity analyses because there were a small number of included studies in these comparisons.

RESULTS

Description of studies

See [Characteristics of included studies](#); [Characteristics of excluded studies](#).

Results of the search

Five search periods characterized this review: 2008, 2011, 2012, 2013, 2014, and 2016. We employed three databases: CENTRAL, MEDLINE, and Embase ([Appendix 2](#); [Appendix 3](#); [Appendix 4](#)). The total yield for 2008 to 2016 was 11,328 records: 1997 from CENTRAL, 3662 from MEDLINE, and 5669 from Embase plus two records from other sources. We did not search [ClinicalTrials.gov](#) or the World Health Organization (WHO) International Clinical Trials Registry Platform (ICTRP). The details of the search results can be found in [Figure 1](#).

Figure 1. Study flow diagram.





Figure 1. (Continued)



Included studies

We found 160 randomized controlled trials (RCTs) with 54,109 enrolled participants, of whom 53,713 started the studies and 50,034 completed the studies or data analysis (or both) (Appendix 1).

Sixteen of the included studies had to be translated into English: one Chinese (Zhou 2008), four Italian (Aceto 2002; Aceto 2003;

Bonato 2001; Girardi 1994), one French (Haimeur 1997), six German (Adams 1994; Blendinger 1976; Kasmacher 1996; Lehmann 1985; Lehmann 1992; Navarro 2000), two Japanese (Masuda 2002; Morimoto 2002), four Spanish (Anez 2001; Echevarria 1998; Hachero 2001; Monedero 1994), and one Turkish (Yildiz 2002). The remaining 141 were in English (Appendix 1).

Surgical risk was classified as minor in 16 (10%), moderate in 74 (46%), major in 25 (16%), and mixed in 45 (28%). There were

142 studies that could not be used in meta-analytic comparisons because 115 had zero awareness events and 27 had excessive clinical and methodological heterogeneity. Most of the 115 studies assessed awareness as a secondary outcome and, therefore, were not powered to identify awareness events. We classified 27 studies that did have awareness events as too dissimilar to merge mainly because of the lack of consensus regarding the definition of intravenous techniques ([Appendix 1](#)).

We merged 18 out of the 45 studies with awareness events in meta-analysis. These 18 studies involved 36,044 participants. In the analysis of anaesthesia depth monitoring (either Bispectral Index or M-entropy) versus standard clinical and electronic monitoring, there were nine studies of the most commonly used anaesthetics with a merged sample of 34,754 participants, which was 96% of the entire patient sample. There were nine studies, with 1290 (4%) participants, of older anaesthetics that are more commonly used in low-income countries ([WHO 2015](#)).

Trial location

One hundred and forty-two (89%) of the studies were from three continents: Europe 72 (45%), Asia 42 (26%), and North America 28 (18%). Five countries were the locations for 75 (47%) of the included studies: Germany 23 (14%), the USA 21 (13%), the UK 13 (8%), Italy 10 (6%), and India 8 (5%). The remaining countries varied between 1% and 4% ([Appendix 1](#)).

Anaesthetic interventions

Frequency of anaesthetic techniques

Different intravenous techniques were compared in 66 studies (41%). Intravenous techniques were compared to volatile agent techniques in 23 studies (14%). Volatile agent techniques were compared in 68 studies (43%). Other techniques were randomized in three studies (2%). Nitrous oxide was randomized in 13 studies (8%): it was used, but not randomized, in 69 studies (43%), and not used in 78 studies (49%). No neuromuscular blocking drugs were used in 13 (8%) of the included studies. In no study was the use of neuromuscular blocking drugs the experimental intervention ([Appendix 1](#)).

Anaesthetic depth monitoring interventions

Ninety-four (59%) of the included studies had one or more processed electroencephalogram (EEG) or auditory evoked potentials (AEP) anaesthetic depth monitors that were part of the randomized or non-randomized protocols. The Bispectral Index monitor was used in 66 (70%) of the 94 studies that included anaesthetic depth monitoring. Forty-three (65%) of the 66 studies that used a Bispectral Index monitor defined a target range against which to titrate anaesthetics. Forty-three (16.3%) used a target range of less than 50; five (11.6%) used a target range of less than 55; 28 (65.1%) used a target range of less than 60; two (4.7%) used a target range of less than 65; and one (2.3%) used a target range of less than 75 ([Appendix 1](#)).

In the five largest studies of anaesthetic depth monitors, three recruited participants expected to be at high risk of awareness ([Avidan 2008](#); [Avidan 2011](#); [Myles 2004](#)); in the other two studies a high expected risk of awareness was not described as an inclusion criterion ([Mashour 2012](#); [Zhang 2011](#)). Additionally, in the smaller trials of anaesthetic depth monitors three were not high-risk patient studies ([Gruenewald 2007](#); [Keressens 2009](#); [Mozafari 2014](#)),

and one study was in high-risk patients ([Puri 2003](#)). In the high-risk studies, there were 86 definite and possible awareness events (a rate of 0.85% in 10,147 participants); in the other five studies there were 86 definite and possible awareness events (a rate of 0.35% in 24,597 participants) ([Appendix 1](#)).

Classification of outcomes

Grade 0 is a descriptor for adequate anaesthesia ([Russell 1997](#)). Adequate anaesthesia is defined as no signs of light anaesthesia such as tachycardia, hypertension, or non-purposeful to purposeful movement. For the purpose of this review, we included participants from studies that did not display such signs but we had limited data to make this judgement; we classified six studies (4%) as such.

Grade 1 is a descriptor for intraoperative wakefulness with obliterated explicit and implicit memory ([Andrade 2008](#)). One hundred included studies (63%) reported Grade 1 outcomes. This classification was assigned by the presence of signs of light anaesthesia such as tachycardia and/or hypertension identified in haemodynamic data, BIS data and/or patient movement etc, within a study's reported results ([Appendix 1](#)).

Grade 2 is a descriptor for intraoperative wakefulness with subsequent implicit memory ([Merikle 1996](#)). Ten of the included studies (6%) reported Grade 2 outcomes.

Grade 3 is a descriptor for intraoperative wakefulness with implicit emotional memory ([Wang 2000](#)). No included study reported Grade 3 outcomes.

Grade 4 is a descriptor for patients with awareness yet resilience ([Sandin 2000](#)). Forty-three of the included studies (27%) reported Grade 4 outcomes.

Grade 5 is a descriptor for patients with awareness with emotional sequelae ([Osterman 2001](#)). Two of the included studies (1%) reported Grade 5 outcomes.

In the five comparisons of included studies reporting Grade 4 cognitive states we merged 17 (94%) of the 18 studies ([Abboud 1985](#); [Avidan 2008](#); [Avidan 2011](#); [Baraka 1989](#); [Crawford 1985](#); [Ellingson 1977](#); [Gruenewald 2007](#); [Haram 1981](#); [Keressens 2009](#); [Mashour 2012](#); [McNulty 1995](#); [Miller 1996](#); [Mozafari 2014](#); [Myles 2004](#); [Puri 2003](#); [Schultetus 1986](#); [Zhang 2009](#)), and one (6%) included study reporting Grade 5 cognitive states ([Russell 1986](#)), which were suitable for meta-analysis ([Appendix 1](#); [Table 1](#)).

Excluded studies

We excluded 315 randomized controlled trials for the reasons described in [Characteristics of excluded studies](#) and [Figure 1](#).

Studies awaiting classification

There are 10 studies awaiting classification for the reasons detailed in the [Characteristics of studies awaiting classification](#) table ([Aceto 2015](#); [Asouhidou 2015](#); [Elbadawy 2015](#); [Hoymork 2007](#); [Jiang 2016](#); [Khanjani 2014](#); [Lequeux 2014](#); [Mehmandoust 2013](#); [Rajan 2015](#); [Xie 2015](#)).

Ongoing studies

There are no ongoing studies.

Risk of bias in included studies

We were able to find 113 (70.6%) of the included study authors' email addresses and we sent them 'Risk of bias' surveys. Fifty-four (47.8%) of the authors who were sent surveys responded (survey responders), which is 33.8% (54) of the included studies (Appendix 1).

We assessed the impact of the 'Risk of bias' survey results on the change in risk of bias classification groups of high, low, or unclear risk of bias from the total included group of 160 studies in two subgroups: survey responders and non-responders.

In the 160 included studies group there were 1120 (7*160) domains and in the 18 studies included in meta-analyses there were 126 domains (Figure 2; Figure 3; Appendix 6). There was a large shift from unclear to either high or low risk of bias as a result of the response from authors (survey responders) (Appendix 6). In comparisons three to five, the lack of survey response resulted in a high percentage of unclear risk of bias domains and, therefore, downgrading of the quality of the evidence for risk of bias. More importantly, this downgrading of the quality of the evidence for risk of bias compared to the lack of downgrading for comparisons one and two was the difference between a very low quality grade compared to a low quality grade in those comparisons (Appendix 6).

Figure 2. 'Risk of bias' graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.




Figure 3. 'Risk of bias' summary: review authors' judgments about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Abboud 1985	?	?	?	+	+	+	?
Abboud 1989	?	?	?	?	+	+	?
Abboud 1995a	?	?	?	?	+	+	?
Abboud 1995b	?	?	?	+	+	+	?
Aceto 2002	?	?	?	?	+	+	?
Aceto 2003	+	?	?	?	+	+	?
Adams 1994	+	+	+	+	+	+	+
Agarwal 1977	?	?	?	?	+	+	?
Aime 2006	+	?	-	+	+	+	?
Akcali 2008	?	?	-	-	+	+	+
Anez 2001	-	?	-	?	+	+	?
Arellano 2000	+	+	-	+	+	+	?
Ashworth 1998	?	-	-	?	+	+	?
Avidan 2008	+	+	-	+	+	+	+
Avidan 2011	+	+	-	+	+	+	+
Baraka 1989	?	?	?	?	+	+	?
Baraka 1998	+	-	-	+	+	+	+
Barr 1977	+	?	+	-	+	+	?
Bauer 2004	+	?	-	?	+	+	+
Bergmann 2013	+	?	-	?	+	+	?

Figure 3. (Continued)

Bergmann 2013	+	?	-	?	+	+	?
Bestas 2004	+	-	-	+	+	+	+
Bethune 1992	?	?	?	?	+	+	?
Bhawna 2012	?	+	+	+	+	+	+
Blendinger 1976	?	?	?	?	+	+	?
Block 1991	?	?	?	?	+	+	?
Bonato 2001	?	?	-	?	+	+	?
Browne 1973	?	?	?	?	+	+	?
Bruhn 2005	+	-	-	?	+	+	?
Casati 1999	?	?	?	?	+	+	?
Celebioglu 2002	-	-	?	?	+	+	?
Celleno 1993	+	+	+	-	+	+	+
Chen 2009	?	?	?	+	+	+	?
Cheun 1987	?	?	?	?	+	+	?
Chin 2004	+	+	?	?	+	+	?
Choi 2012	+	?	-	?	+	+	?
Clyburn 1986	+	+	-	-	+	+	+
Coates 1987	?	?	?	?	+	+	?
Collins 1996	?	?	?	+	+	+	?
Crawford 1985	+	?	-	?	+	+	?
Crawford ME 1984	?	?	?	?	+	+	?
Czarko 2013	?	?	-	+	+	+	?
Deeprise 2005	+	-	-	+	+	+	+
De Kock 1995	+	+	+	+	+	+	+
Deshpande 2009	?	?	?	?	+	+	?
Dhadphale 1979	?	?	?	?	+	+	?
Drover 2002	+	?	-	-	+	+	+
Dunnett 1977	?	?	?	+	+	+	?
Echevarria 1998	+	+	?	+	+	+	+
Elhakim 2010	+	+	+	+	+	+	+
Ellingson 1977	?	?	?	?	+	+	?

Figure 3. (Continued)

Ellingson 1977	?	?	?	?	+	+	?
Fehr 2001	?	?	+	?	+	+	?
Forestier 2003	+	?	?	?	+	+	?
Fragen 1981	+	?	+	?	+	+	?
Gaitini 1995	+	?	?	?	+	+	?
Gale 2001	+	+	-	+	+	+	+
Ghaly 1988	?	?	?	?	+	+	?
Ghoneim 2000	+	?	?	+	+	+	?
Ghosh 2008	+	+	+	+	+	+	+
Girardi 1994	?	?	?	?	+	+	?
Gokce 2009	+	?	?	+	+	+	?
Goto 2000	+	+	-	+	+	+	+
Gruenewald 2007	?	?	-	+	+	+	?
Grundmann 2001	?	?	?	+	+	+	?
Gupta 1992	?	?	?	?	+	+	?
Gurman 1994	-	+	+	+	?	+	+
Hachero 2001	+	?	?	+	+	+	?
Hackner 2003	?	?	+	+	+	+	?
Haimeur 1997	?	?	?	?	+	+	?
Haram 1981	?	?	+	+	?	+	?
Hug 1988	?	?	?	?	+	+	?
Hung 1992	?	?	?	?	+	+	?
Ibraheim 2008	?	?	-	+	+	+	?
Inoue 2005	?	?	?	+	+	+	?
Jensen 1995	+	+	+	+	+	+	+
Jiahai 2012	?	?	?	?	+	+	?
Kamal 1990	?	?	+	?	+	+	?
Kasmacher 1996	?	?	?	?	+	+	?
Kerssens 2005b	+	?	-	+	+	+	?
Kerssens 2009	+	?	?	+	+	+	+
Kim 2007	?	?	?	?	+	+	?

Figure 3. (Continued)

Kim 2007	?	?	?	?	+	+	?
Kiyama 1997	?	?	+	-	+	+	?
Kreuer 2003	?	?	?	?	+	+	?
Kreuer 2005	+	?	-	+	+	+	+
Krissel 1994	?	?	?	?	+	+	?
Kudoh 1999	?	?	?	?	+	+	?
Kwon 2013	?	?	-	+	+	+	?
Lallemand 2003	+	+	+	+	+	+	+
Lam 2013	?	?	-	?	+	+	?
Lehmann 1985	+	+	+	+	+	+	+
Lehmann 1992	+	+	+	+	+	+	+
Lehmann 2007	+	-	-	-	+	+	+
Lim 1992	?	?	?	?	+	+	?
Lin 2011	?	?	?	?	+	+	?
Lindholm 2008	+	-	-	?	+	+	+
Liu 2013	+	?	?	?	+	+	?
Lu 2005	+	+	+	+	+	+	+
Maattanen 2002	?	?	-	+	+	+	?
Mashour 2012	+	+	-	+	+	+	+
Masuda 2002	?	+	-	+	?	+	+
McNulty 1995	?	?	+	?	+	+	?
Menigaux 2002	+	+	+	+	+	+	?
Mertens 2003	?	?	?	?	+	+	?
Miller 1996	+	?	+	?	+	+	?
Miranda 1992	?	?	?	+	+	+	?
Monedero 1994	?	?	?	?	+	+	?
Morimoto 2002	?	?	-	-	+	+	+
Mozafari 2014	+	?	-	?	+	+	?
Muralidhar 2008	?	?	?	?	+	+	?
Myles 1997	?	-	-	?	+	+	?
Myles 2004	+	+	?	+	+	+	?

Figure 3. (Continued)

Myles 2004	+	+	?	+	+	+	?
Myles 2007	+	?	-	+	+	+	?
Navarro 2000	?	?	?	?	+	+	?
Nayar 2009	?	?	?	+	+	+	?
Ngan 1997	+	?	+	?	+	+	?
Ngan Kee 2002	+	+	+	+	+	+	+
Oddby-Muhrbeck 1993	?	+	+	+	+	+	+
Panousis 2009	+	+	+	?	+	+	?
Pauls 2009	+	?	-	+	+	+	+
Paventi 2001	?	?	?	?	+	+	?
Pedersen 1992	?	?	+	?	+	+	?
Persec 2012	?	?	-	?	+	+	?
Piggott 1990	-	-	-	+	+	+	+
Plourde 1996	+	?	-	+	+	+	+
Puri 2003	+	?	?	?	+	+	?
Puri 2007	+	-	-	+	+	+	+
Rehberg 2007	+	?	-	-	+	+	+
Renna 2000	+	?	-	?	+	+	?
Rinaldi 2005	?	?	-	+	+	+	?
Russell 1986	-	-	-	-	+	+	+
Samarkandi 2004	-	?	?	+	+	+	?
Sareen 1997	?	?	+	?	+	+	?
Schultetus 1986	+	?	?	?	+	+	?
Schwender 1994	?	?	-	+	+	+	?
Schwender 1996	?	?	?	?	+	+	?
Shin 2012	+	?	?	?	+	+	?
Sidi 1990	?	?	?	?	+	+	?
Smith 1999	?	?	?	?	+	+	?
Song 1997	+	?	-	?	+	+	?
Sorbara 1995	?	?	?	?	+	+	?
Soyannwo 1988	?	?	?	?	+	+	?

Figure 3. (Continued)

Soyannwo 1988	?	?	?	?	+	+	?
Staikou 2013	?	?	+	?	+	+	+
Stoppe 2012	+	?	-	+	+	+	+
Stuttman 2010	+	?	-	+	+	+	+
Toft 1987	?	?	+	+	+	+	?
Toscano 2007	+	?	+	?	+	+	+
Tsai 2001	?	?	?	+	+	+	?
Tunstall 1989	?	?	?	?	+	+	?
Vakkuri 2005	+	?	-	+	+	+	+
van der Maaten 1996	+	+	?	+	+	+	+
van Leeuwen 1990	?	?	?	?	+	+	?
Wang 2013	+	?	+	+	+	+	?
White 2003	?	?	?	?	+	+	?
White 2004	?	?	-	?	+	+	?
Wong 2002	+	+	-	+	+	+	+
Wu 2001	?	?	?	?	+	+	?
Yildiz 2002	?	?	?	?	+	+	?
Yoshitani 2003	+	?	-	+	+	+	+
Zhang 2011	-	?	-	+	+	+	+
Zhou 2008	+	?	?	?	+	+	?
Zohar 2006	?	?	-	+	+	+	?

Allocation

Random sequence generation

For the 18 studies included in meta-analysis, the risk of bias was high in 11% (2/18), low in 56% (10/18), and unclear in 33% (6/18).

Random allocation concealment

For the 18 studies included in meta-analysis, the risk of bias was high in 0.0% (0/18), low in 27.8% (5/18), and unclear in 72.2% (13/18).

Blinding

Blinding of participants

For the 18 studies included in meta-analysis, the risk of bias was high in 44.4% (8/18), low in 22.2% (4/18), and unclear in 33.3% (6/18).

Blinding of outcome assessment

For the 18 studies included in meta-analysis, the risk of bias was high in 5.6% (1/18), low in 50.0% (9/18), and unclear in 44.0% (8/18).

Incomplete outcome data

For the 18 studies included in meta-analysis, the risk of bias was high in 0% (0/18), low in 94% (17/18), and unclear in 6% (1/18).

Selective reporting

For the 18 studies included in meta-analysis, the risk of bias was high in 0% (0/18), low in 100% (18/18), and unclear in 0% (0/18).

Other potential sources of bias

For the 18 studies included in meta-analysis, the risk of bias was high in 0% (0/18), low in 33% (6/18), and unclear in 67% (12/18).

Effects of interventions

See: [Summary of findings for the main comparison Anaesthesia depth monitors \(BIS and M-entropy\) versus standard clinical and electronic monitoring](#); [Summary of findings 2 Anaesthesia depth monitors \(BIS\) versus standard clinical and electronic monitoring](#); [Summary of findings 3 Thiopentone with and without added hypnotic drugs \(ketamine, etomidate\)](#); [Summary of findings 4 Thiopentone and ketamine versus benzodiazepines \(diazepam, midazolam, lorazepam\)](#); [Summary of findings 5 Caesarean section with low- and high-dose inhaled agent](#)

Primary outcomes


Awareness

1.1 Anaesthesia depth monitors (either Bispectral Index (BIS) or M-entropy) versus standard clinical parameter (Grade 4)

Awareness was an uncommon event in nine studies (Avidan 2008; Avidan 2011; Gruenewald 2007; Kerssens 2009; Mashour 2012;

Mozafari 2014; Myles 2004; Puri 2003; Zhang 2011). There were 173 occurrences among 34,744 patients (anaesthesia depth monitors 85/17,713 versus standard clinical and electronic monitoring 88/17,031), an overall event rate of about 0.5% (Analysis 1.1). These nine studies had considerable clinical and methodological heterogeneity. There was moderate statistical heterogeneity ($I^2 = 49\%$; $P = 0.04$) (Figure 4). The effect size favoured neither anaesthesia depth monitoring nor standard clinical and electronic monitoring, with little precision in the odds ratio (OR) estimate (OR 0.98, 95% confidence interval (CI) 0.59 to 1.62; 95% prediction interval (PI) 0.33 to 2.90) (Analysis 1.1). There was no difference in the odds of awareness between high-risk and non-high-risk patients ($F_{1,7} = 1.70$; $P = 0.23$). With an assumed risk for awareness using standard clinical and electronic monitoring being 5 per 1000, the corresponding risk using anaesthesia depth monitoring was 5 per 1000 (95% CI 3 to 7). The quality of the evidence was low (Summary of findings for the main comparison).

Figure 4. Meta-analysis patient awareness reports not adjudicated or adjudicated as possible and definite awareness: ADM versus standard clinical and electronic monitoring




2.1 Anaesthesia depth monitors (Bispectral Index) versus standard clinical and electronic monitoring (Grade 4)

In five of the anaesthesia depth monitoring studies a narrative report of potential awareness events identified in one or more interviews was submitted to a blinded, expert panel (Avidan 2008; Avidan 2011; Mashour 2012; Myles 2004; Zhang 2011). This panel adjudicated the outcome for each patient: no awareness, possible awareness, or definite awareness. Episodes of dreaming were also declared in two studies (Myles 2004; Zhang 2011). Expert panel adjudication was not used in the other studies in Analysis 1.1 (Gruenewald 2007; Kerssens 2009; Mozafari 2014; Puri 2003). Unpublished data were provided by the authors (Avidan 2008; Avidan 2011; Mashour 2012; Myles 2004; Zhang 2011), with details of the adjudication results (Appendix 8). There was wide variation in the results of the adjudication process.

2.1 Anaesthesia depth monitors (Bispectral Index) versus standard clinical and electronic monitoring: risk of definite and possible awareness (Grade 4)

Definite and possible awareness was an uncommon event with 152 occurrences among 34,181 participants (Bispectral Index 74/17,432 versus standard clinical and electronic monitoring 78/16,749), an overall event rate of about 0.4% (Analysis 2.1). The five studies had considerable clinical and methodological heterogeneity. There was a substantial degree of statistical heterogeneity ($I^2 = 68\%$; $P = 0.01$) (Figure 5). The effect size favoured neither Bispectral Index nor standard clinical and electronic monitoring, with little precision in the OR estimate (OR 0.96, 95% CI 0.35 to 2.65; 95% PI 0.15 to 6.41) (Analysis 2.1). The assumed risk for awareness (definite and possible awareness) using standard clinical and electronic monitoring was 5 per 1000; the corresponding risk using Bispectral Index monitoring was 5 per 1000 (95% CI 2 to 8). The quality of the evidence was low (Summary of findings 2).

Figure 5. Meta-analysis possible and definite awareness: ADM versus standard clinical and electronic monitoring (adjudication)



We subgrouped the studies by the use of alarms or alerts (or both) in the protocol and by the type of anaesthesia used (inhalation versus total intravenous anaesthesia (TIVA)). The protocols in three studies included alarms/alerts and the use of inhalation anaesthesia (Avidan 2008; Avidan 2011; Mashour 2012); the protocols in two studies did not include alarms or alerts and used predominantly

or exclusively TIVA (Myles 2004; Zhang 2011)). We repeated the meta-analysis with the studies separated into these two subgroups. There was no difference in the OR between the two subgroups ($F_{1,3} = 37$; $P = 0.16$).


We subgrouped the studies by the risk of awareness in patient recruitment. We repeated the meta-analysis with the studies separated into these two subgroups. There was no difference in the OR between the two subgroups ($F_{1,3} = 2.11$; $P = 0.24$).

2.2 Anaesthesia depth monitors (Bispectral Index) versus standard clinical and electronic monitoring: risk of definite awareness (Grade 4)

Definite awareness was a rare event with 65 occurrences among 34,181 patients (Bispectral Index 24/17,432 versus standard clinical and electronic monitoring 41/16,749), an overall event rate of about

0.2% (Analysis 2.2). The five studies had considerable clinical and methodological heterogeneity. There was a substantial degree of statistical heterogeneity ($I^2 = 60\%$; $P = 0.02$) (Figure 6). The effect size favoured Bispectral Index monitoring but with little precision in the OR estimate (OR 0.60, 95% CI 0.13 to 2.75; 95% PI 0.04 to 9.20) (Analysis 2.2). The assumed risk for awareness (definite awareness) using standard clinical and electronic monitoring was 2 per 1000; the corresponding risk using Bispectral Index monitoring was 1 per 1000 (95% CI 1 to 4). The quality of the evidence was low (Summary of findings 2).

Figure 6. Meta-analysis definite awareness: ADM versus standard clinical and electronic monitoring (adjudication)



We subgrouped the studies by the use of alarms or alerts (or both) in the protocol and by the type of anaesthesia (inhalation versus TIVA). The protocols in three studies included alarms or alerts and the use of inhalation anaesthesia (Avidan 2008; Avidan 2011; Mashour 2012); the protocols in two studies did not include alarms/alerts and used predominantly or exclusively TIVA (Myles 2004; Zhang 2011). We repeated the meta-analysis with the studies separated into these two subgroups. There was no difference in the OR between the two subgroups ($F_{1,3} = 7.21$; $P = 0.08$).

We subgrouped the studies by the risk of awareness in patient recruitment. We repeated the meta-analysis with the studies separated into these two subgroups. There was no difference in the OR between the two subgroups ($F_{1,3} = 0.43$; $P = 0.56$).

With regard to the analysis of expert adjudication of awareness in Analysis 2.1 and Analysis 2.2, as a percentage of study size the events flagged for adjudication ranged from 0.40% to 5.73%. As a percentage of study size, 'no awareness' ranged from 0.21% to 5.18%, 'possible awareness' ranged from 0.10% to 1.46%, and 'definite awareness' ranged from 0.10% to 0.53%. There was a large degree of between-study heterogeneity ($I^2 = 99\%$) in the analysis of the expert panels' adjudication of patient awareness reports (Appendix 1; Appendix 8).

Wakefulness

3.1 Thiopentone with and without added hypnotic drugs (ketamine, etomidate) (Grades 4, 5)

In four studies the isolated forearm technique was used to assess intraoperative wakefulness (Grade 0, 1, 2, or 3) by response

to complex commands (Baraka 1989; Russell 1986; Schultetus 1986; Tunstall 1989). The surgery was caesarean delivery or a gynaecological procedure. In three studies wakefulness was compared between thiopentone with and without added hypnotic drugs (ketamine, etomidate) after induction (Baraka 1989; Russell 1986; Schultetus 1986) (Analysis 3.1). There was a lower risk of wakefulness with the addition of hypnotic drugs (risk ratio (RR) 0.18, 95% CI 0.09 to 0.41). For an assumed risk of wakefulness for thiopentone of 562 per 1000, the corresponding risk will be 101 per 1000 (51 to 230) with the administration of hypnotic drugs. The quality of the evidence was low (Summary of findings 3).

Patients were questioned postoperatively concerning recall (Grade 4 or 5) (Baraka 1989; Russell 1986; Schultetus 1986; Tunstall 1989). We compared the proportion of patients with wakefulness (Grade 0, 1, 2, or 3) versus awareness (Grade 4 or 5) in a random-effects meta-analysis. Of the 254 participants studied, there were six with awareness and 90 demonstrated wakefulness. The proportion with awareness was 0.04 (95% CI 0.01 to 0.11) while the proportion with wakefulness was 0.34 (95% CI 0.25 to 0.45); these proportions were different ($F_{1,6} = 26.4$, $P = 0.0021$), with non-significant statistical heterogeneity ($P = 0.12$) (Appendix 1; Appendix 7).

Awareness

4.1 Thiopentone and ketamine versus benzodiazepines (diazepam, midazolam, lorazepam) (Grade 4)

In four studies there were 18 occurrences among 291 participants (benzodiazepines 2/192 versus thiopentone and ketamine 16/99), an overall event rate of about 6% (Ellingson 1977; Haram 1981; McNulty 1995; Miller 1996), (Analysis 4.1). There was a lower risk of awareness with the addition of benzodiazepines (RR 0.17, 95% CI 0.02 to 0.25). The assumed risk for awareness for thiopentone and ketamine was 131 per 1000; the corresponding risk using benzodiazepines was 28 per 1000 (95% CI 3 to 42). The quality of the evidence was very low (Summary of findings 4).

5.1 Caesarean section with low-dose and high-dose inhaled anaesthetic agents

There were two studies with 848 participants that compared low-dose to high-dose volatile agents (halothane, enflurane, trichloroethylene) with and without nitrous oxide during caesarean section (Abboud 1985; Crawford 1985). There were 24 occurrences among 848 participants (high-dose 3/449 versus low-dose 21/435), an overall event rate of about 3% (Analysis 5.1). There was a lower risk of awareness in the group receiving high-dose inhaled anaesthetics (RR 0.13, 95% CI 0.04 to 0.42). The assumed risk for awareness for low-dose inhaled anaesthetics was 57 per 1000; the corresponding risk using a high dose was 7 per 1000 (95% CI 2 to 23). The quality of the evidence was very low (Summary of findings 5).

We graded the quality of the evidence as low in Summary of findings for the main comparison, Summary of findings 2, and Summary of findings 3, and as very low in Summary of findings 4 and Summary of findings 5.

Secondary outcomes

We were unable to assess the secondary outcomes because they were not defined or not identified as outcomes in the included studies. Most studies would miss the diagnosis of post-traumatic

stress disorder because the postoperative interview period was usually within one month.

DISCUSSION

Summary of main results

The main finding of this review is based on 34,744 (96%) of the entire patient sample. The use of processed electroencephalogram anaesthetic depth monitoring does not reduce the risk of awareness during surgery compared to standard clinical and electronic monitoring. Bispectral Index monitors, the most commonly used anaesthetic depth monitors, did not reduce the frequency of awareness compared to standard clinical and electronic monitoring (Summary of findings for the main comparison; Summary of findings 2). Five studies comparing anaesthetic depth monitoring versus standard clinical and electronic monitoring used an expert panel to adjudicate possible events. There was significant heterogeneity in the range of patient awareness reports adjudicated as 'no awareness' by expert panels (Appendix 8).

The mapping of data within the individual randomized controlled trials (RCTs) included in this review into the broader classification system of Wang et al (Table 1) was difficult. The classification system was designed for use following surgery using the patient interview and anaesthesia record as the data source. In contrast, there are limitations to its retrospective application to patients grouped in a trial report. We assigned the highest grade(s) that were consistent with the clinical signs of anaesthetic depth contained in each study or comparison (Appendix 1). Nonetheless, the algorithm that we created allowed us to classify all of the included studies, except for six (4%) (Appendix 1). Those six studies could not be classified because the authors did not include data from ADMs (BIS) or standard clinical parameters such as hemodynamic and/or somatic data in their results sections.

In three smaller meta-analyses based on nine older studies of 1290 participants from among the entire review's patient sample, the evidence indicates that ketamine, a dissociative agent, and etomidate, an intravenous imidazole general anaesthetic, are associated with less wakefulness than the ultra-short-acting barbiturate thiopentone. The World Health Organization's list of essential medicines is based on the "...most efficacious, safe and cost-effective medicines for priority conditions..." (WHO 2015). The anaesthetics studied in this review are on this list; many of the older drugs are still in common use in developing countries. Therefore, the merged findings from the smaller, older studies are relevant to current practice: benzodiazepines reduced episodes of awareness compared to thiopentone and ketamine and a higher dose of inhaled anaesthetic agents reduced episodes of awareness compared to a lower dose (Summary of findings 3; Summary of findings 4; Summary of findings 5). Five of eight (63%) of the medications studied (ketamine, thiopentone, halothane, nitrous oxide, benzodiazepines) in these older studies are essential medications commonly used by low-income countries (WHO 2015).

As mentioned, benzodiazepines are associated with less awareness than ketamine and thiopental. However, there are no means of determining whether this is simply an amnesic effect rather than an actual increase in anaesthetic depth. This finding highlights the problematic confusion between awareness (with explicit recall) and intraoperative wakefulness without explicit recall and the

inadequacy of the criterion of postoperative recall as the definition of adequate anaesthesia. Hence, this suggests a need for a more sophisticated classification of intraoperative cognitive states (Wang 2012). In three small studies using the isolated forearm technique, wakefulness was far more frequent than awareness (Analysis 3.1; Effects of interventions; Appendix 1).

Overall completeness and applicability of evidence

Our literature search identified RCTs published over a 56-year period (1960 through 2016). We classified the entire set of 160 included studies as either included in meta-analyses (18) or not used (142) (Appendix 1; Included studies). Although the majority of studies were not merged in meta-analysis, these 142 studies contributed to the evidence contained in this review by allowing the creation of descriptive statistics (Appendix 1; Table 1). This may be helpful to future researchers studying interventions to decrease the frequency of wakefulness and awareness. There are 10 studies awaiting classification, which we will process when we update the review. In any update of this review, trial registries should also be searched.

Quality of the evidence

The patient awareness report classification system adjudicated by an expert panel used in the studies in Analysis 2.1 and Analysis 2.2 was introduced in 2004. One of the goals of this classification system was to capture more patient awareness events compared to a system that requires confirmation from intraoperative staff that patient memories of intraoperative events actually happened (Sebel 2004). We found evidence of heterogeneity between centres in the expert panels' classification of awareness from our author survey (Appendix 8).

We successfully applied the classification system of Wang et al to 96% of included studies (Wang 2012; Table 1). The classification criteria are clinical signs of light anaesthesia such as haemodynamic variables, lacrimation, and response to command, and symptoms such as the report of pain, postoperative recall and/or distress during surgery, as well as nightmares and post-traumatic stress disorder-like criteria in the postoperative period. In the 18 studies submitted for meta-analysis, all awareness events were graded 4 (awareness but resilient patient) or 5 (awareness with emotional sequelae).

We improved the quality of the evidence by identifying unpublished data for the 'Risk of bias' tables derived from our author survey (Appendix 6). The author responses to the survey provided more information about the seven risk of bias domains. This significantly changed the number of 'unclear' risk ratings to more appropriate ratings of high or low risk of bias in this review (Characteristics of included studies).

The survey allowed us to update domains for studies included in Analysis 1.1, Analysis 2.1, and Analysis 2.2. However, we received no responses from authors of studies contributing to Analysis 3.1, Analysis 4.1, or Analysis 5.1.

The risk of bias due to study funding source (anaesthesia depth monitor manufacturers and pharmaceutical companies) was unclear due to a lack of detailed disclosure information in many of the included studies. There were two types of interventions in this review: drugs and devices.

Investigator blinding was used in 31 (19%) of the 160 included studies (Appendix 6). In the drug studies that did not use blinding, performance bias may have occurred. The experimental design of anaesthesia depth monitor device trials precludes the possibility of blinding the provider.

We downgraded all comparisons in the 'Summary of findings' tables by two or three levels to a low or very low quality of evidence. The downgrades were for within-study risk of bias, inconsistency, and imprecision.

We analysed the strengths and weaknesses of the criteria that we used to merge the studies in the five comparisons in this review (Appendix 1).

Potential biases in the review process

A strength of this review was the avoidance of potential bias in the electronic literature search: we used no language restrictions. The included studies came from Europe, North America, Asia, the Middle East, Australia and New Zealand, and Africa (Appendix 1). Therefore, there was no evidence of reporting bias due to location. Included studies were written in seven languages other than English (Appendix 1), therefore there was no evidence of language bias. However, we did not search ClinicalTrials.gov or the World Health Organization (WHO) International Clinical Trials Registry Platform (ICTRP).

In addition, there were insufficient studies in each comparison to allow statistical testing for reporting bias. We did survey study authors to obtain additional information to rate the risk of bias domains, which enabled a shift from unclear to low and from unclear to high risk of bias for various domains, as reported above (Appendix 6).

Agreements and disagreements with other studies or reviews

A Cochrane review has compared the incorporation of the Bispectral Index anaesthetic depth monitor into standard practice for the management of anaesthesia (Punjasawadwong 2014). This review used search language that focused on identifying anaesthesia depth monitor (Bispectral Index) trials. The authors assessed the effects of Bispectral Index monitoring versus clinical signs and Bispectral Index monitoring versus end-tidal anaesthetic gas (ETAG) monitoring on the risk of definite awareness events; other outcomes of awakening times and anaesthetic usage were also compared. For definite awareness, they reported that the summary Peto odds ratio (OR) was significantly lower for Bispectral Index monitoring compared to clinical signs (OR 0.24, 95% confidence interval (CI) 0.12 to 0.48) (Analysis 1.1 in Punjasawadwong 2014), but not lower for Bispectral Index monitoring compared to ETAG (OR 1.13, 95% CI 0.56 to 2.26) (Analysis 1.2 in Punjasawadwong 2014). Their two analyses included five studies that used the adjudication process (Avidan 2008; Avidan 2011; Mashour 2012; Myles 2004; Zhang 2011), two studies without awareness events (Muralidhar 2008; Samarkandi 2004), and one additional study included in our Analysis 1.1 (Puri 2003). Our current review used search language focused on identifying awareness studies. Analysis 1.1 includes two additional studies not found in Punjasawadwong 2014 (Gruenewald 2007; Kerssens 2009). As a result, the reviews have unique included studies.

We did not find a difference in definite awareness rates (OR 0.60, 95% CI 0.13 to 2.75) ([Analysis 2.2](#)). Our review employed different statistical models from [Punjasawadwong 2014](#). Generally the point estimates of the individual Peto ORs in [Punjasawadwong 2014](#) ([Avidan 2008](#); [Avidan 2011](#); [Mashour 2012](#); [Myles 2004](#); [Zhang 2011](#)), were similar to the OR point estimates found here. However, the 95% CIs were wider.

Following the publication of [Punjasawadwong 2014](#), a Cochrane editorial recommended continued use of processed electroencephalogram monitors and standard clinical and electronic monitoring in research protocols to assess the ability of anaesthetic interventions to reduce the frequency of awareness ([Kettner 2014](#)).

A Cochrane review has also compared the use of the anaesthetic depth monitor spectral entropy monitoring to standard practice; one of the secondary outcomes was awareness during anaesthesia ([Chhabra 2016](#)). Eight trials with 797 participants reporting awareness were found. Of these, only one study reported awareness, with one participant in the control group having an episode ([Gruenewald 2007](#)). This study is included our review. The authors of [Chhabra 2016](#) concluded that there was insufficient evidence to support spectral entropy monitoring to detect awareness.

AUTHORS' CONCLUSIONS

Implications for practice

Based on the largest portion of evidence in this review, the use of processed electroencephalogram (EEG) anaesthesia depth monitoring may have similar effects on awareness during surgery to standard clinical and electrical monitoring; the risk of awareness is not reduced ([Summary of findings for the main comparison](#); [Summary of findings 2](#)).

Based on a smaller amount of evidence, drawn from older studies, ketamine and etomidate reduces wakefulness compared with thiopental, if this is the goal of anaesthesia. If the goal of anaesthesia is to reduce awareness, then benzodiazepines reduces awareness compared to thiopental, ketamine, and placebo. The safety, clinical efficacy, and low cost of these older anaesthetics are the reasons why they are still used in both high-income and low-income countries ([WHO 2015](#)). Clinicians who use the older anaesthetics assessed in our review can refer to the evidence displayed in [Summary of findings 3](#), [Summary of findings 4](#), and [Summary of findings 5](#).

Our confidence in the conclusions of this review is not strong. There are a large number of unclear risk of bias domains in the included studies. There is also a large degree of heterogeneity in the expert panels' adjudication of patient awareness reports ([Appendix 8](#)). In addition, the event rate is so small in the anaesthetic depth

monitoring studies that, despite the large total sample size, we downgraded the effect estimates for imprecision.

Implications for research

We agree with other investigators ([Mashour 2012](#)), that continuing to study the effects of current processed EEGs on awareness seems futile considering the rarity of the event and the heterogeneity of the effect estimates in five large trials ([Avidan 2008](#); [Avidan 2011](#); [Mashour 2012](#); [Myles 2004](#); [Zhang 2011](#)). Future studies should randomize patients between current total intravenous anaesthesia (TIVA) and potent volatile techniques, with and without muscle relaxants. Tracking wakefulness as well as awareness in future trials will allow the use of smaller sample sizes, given the increased frequency of wakefulness compared to awareness. In addition, the relationship between wakefulness and the risk of developing post-traumatic stress disorder needs further study.

Despite the controversy regarding the definition of a 'conscious' patient under anaesthesia, when surveyed anaesthetists state that if they were a patient, it would be 'unacceptable' to be paralysed, awake, and in pain even with postoperative amnesia of the event ([Girgirah 2006](#)). Hence, there is an increasing consensus amongst many anaesthetists that persistent unconsciousness (lack of wakefulness) rather than simply lack of postoperative awareness (amnesia) should be the goal of anaesthesia ([Sanders 2011](#)). The proposal of Wang et al can be used to classify both wakefulness and awareness ([Wang 2012](#); [Table 1](#)).

The proper methods to adjudicate events as being or not being awareness should also be explored.

ACKNOWLEDGEMENTS

We would like to thank Mathew Zacharias and Andrew Smith (content editors), Marialena Trivella and Cathal Walsh (statistical editors), Paul Myles, Yodying Punjasawadwong, Ian Russell, and Jackie Andrade (peer reviewers), and Tracey Lloyd (consumer representative) for their help and editorial advice during the preparation of this review.

We would also like to thank Mary McFarland from the Eccles Library at the University of Utah and Karen Hovhannisyan, MD, former Trials Search Co-ordinator, Cochrane Anaesthesia, Critical and Emergency Care Group (ACE), for their help with the literature search.

We would like to also thank the following translators: Finnish: Elena Yorgason, Spanish: Isabel Dulfano and Brett Smith, Chinese: Fen Feng, Turkish: Murat Cakia, Japanese: Jiro Kurata, German: Lore Schultheiss, Korean: Hae An Jung, Italian and French: Anthony Messina.

Finally, we would like to thank Maria Eggers (ME) and Devin B Miller (DM) for their extensive assistance in the review.

REFERENCES

References to studies included in this review

Abboud 1985 {published data only}

* Abboud TK, Kim SH, Henriksen EH, Chen T, Eisenman R, Levinson G, et al. Comparative maternal and neonatal effects of halothane and enflurane for cesarean section. *Acta Anaesthesiologica Scandinavica* 1985;**29**(7):663-8. [PUBMED: 4072589]

Abboud 1989 {published data only}

* Abboud TK, D'Onofrio L, Reyes A, Mosaad P, Zhu J, Mantilla M, et al. Isoflurane or halothane for cesarean section: comparative maternal and neonatal effects. *Acta Anaesthesiologica Scandinavica* 1989;**33**(7):578-81. [PUBMED: 2816239]

Abboud 1995a {published data only}

Abboud TK, Zhu J, Richardson M, Peres da Silva E, Donovan M. Desflurane: a new volatile anesthetic for cesarean section. Maternal and neonatal effects. *Acta Anaesthesiologica Scandinavica* 1995;**39**(6):723-6. [PUBMED: 7484023]

Abboud 1995b {published data only}

* Abboud TK, Zhu J, Richardson M, Peres Da Silva E, Donovan M. Intravenous propofol vs thiamylal-isoflurane for caesarean section, comparative maternal and neonatal effects. *Acta Anaesthesiologica Scandinavica* 1995;**39**(2):205-9. [PUBMED: 7793188]

Aceto 2002 {published data only}

Aceto P, Valente A, Adducci E, Gorgoglione M, De Cosmo G. Implicit memory, dream and auditory evoked responses during anaesthesia. *Acta Medica Romana* 2002;**40**(1):21-9.

Aceto 2003 {published data only}

Aceto P, Valente A, Gorgoglione M, Adducci E, De Cosmo G. Relationship between awareness and middle latency auditory evoked responses during surgical anaesthesia. *British Journal of Anaesthesia* 2003;**90**(5):630-5. [PUBMED: 12697591]

Adams 1994 {published data only}

Adams HA, Bauer R, Gebhardt B, Menke W, Baltes-Götz B. Total i.v. anesthesia with S-(+)-ketamine in orthopedic geriatric surgery. Endocrine stress reaction, haemodynamics and recovery. *Anaesthetist* 1994;**43**(2):92-100. [PUBMED: 8141443]

Agarwal 1977 {published data only}

Agarwal G, Sikh SS. Awareness during anaesthesia. A prospective study. *British Journal of Anaesthesia* 1977;**49**(8):835-8. [PUBMED: 889672]

Aime 2006 {published data only}

Aime I, Verroust N, Masson-Lefoll C, Taylor G, Laloe PA, Liu N, et al. Does monitoring bispectral index or spectral entropy reduce sevoflurane use?. *Anesthesia and Analgesia* 2006;**103**(6):1469-77. [PUBMED: 17122226]

Akcali 2008 {published data only}

Akcali DT, Ozkose Z, Yardim S. Do we need bispectral index monitoring during total intravenous anaesthesia for lumbar

discectomies?. *Turkish Neurosurgery* 2008;**18**(2):125-33. [PUBMED: 18597226]

Anez 2001 {published data only}

Anez C, Papaceit J, Sala JM, Fuentes A, Rull M. The effect of encephalogram bispectral index monitoring during total intravenous anaesthesia with propofol in outpatient surgery. *Revista Espanola de Anestesiologia y Reanimacion* 2001;**48**(6):264-9. [PUBMED: 11446941]

Arellano 2000 {published data only}

* Arellano RJ, Pole ML, Rafuse SE, Fletcher M, Saad YG, Friedlander M, et al. Omission of nitrous oxide from a propofol-based anaesthetic does not effect the recovery of women undergoing outpatient gynaecologic surgery. *Anesthesiology* 2000;**93**(2):332-9. [PUBMED: 10910478]

Ashworth 1998 {published data only}

Ashworth J, Smith I. Comparison of desflurane with isoflurane or propofol in spontaneously breathing ambulatory patients. *Anesthesia and Analgesia* 1998;**87**(2):312-8. [PUBMED: 9706922]

Avidan 2008 {published and unpublished data}

* Avidan MS, Zhang L, Burnside BA, Finkel KJ, Searleman AC, Selvidge JA, et al. Anesthesia awareness and the bispectral index. *New England Journal of Medicine* 2008;**358**(11):1097-108. [PUBMED: 18337600]

Avidan 2011 {published and unpublished data}

* Avidan MS, Jacobsohn E, Glick D, Burnside BA, Zhang L, Villafranca A, et al. Prevention of intraoperative awareness in a high-risk surgical population. *New England Journal of Medicine* 2011;**365**(7):591-600. [PUBMED: 21848460]

Villafranca AJ, Arenson BG, Avidan MS, Glick D, Mashour GA, Jacobsohn E. Volitional delay of self-reported outcomes: insights from a case of intraoperative awareness with explicit recall. *Anesthesia and Analgesia* 2013;**116**(2):365-7. [PUBMED: 23302976]

Baraka 1989 {published data only}

Baraka A, Louis F, Noueihid R, Diab M, Dabbous A, Sibai A. Awareness following different techniques of general anaesthesia for caesarean section. *British Journal of Anaesthesia* 1989;**62**(6):645-8. [PUBMED: 2502168]

Baraka 1998 {published data only}

Baraka A, Siddik S, Assaf B. Supplementation of general anaesthesia with tramadol or fentanyl in parturients undergoing elective caesarean section. *Canadian Journal of Anaesthesia* 1998; Vol. 45, issue 7:631-4. [PUBMED: 9717593]

Barr 1977 {published data only}

Barr AM, Moxon A, Woollam CH, Fryer ME. The effect of diazepam and lorazepam on awareness during anaesthesia for caesarian section. *Anaesthesia* 1977;**32**(9):873-8. [PUBMED: 23704]

Bauer 2004 {published data only}

Bauer M, Wilhelm W, Kraemer T, Kreuer S, Brandt A, Adams HA, et al. Impact of bispectral index monitoring on stress response and propofol consumption in patients undergoing coronary artery bypass surgery. *Anesthesiology* 2004;**101**(5):1096-104. [PUBMED: 15505444]

Bergmann 2013 {published data only}

Bergmann I, Gohner A, Crozier TA, Hesjedal B, Wiese CH, Popov AF, et al. Surgical pleth index-guided remifentanyl administration reduces remifentanyl and propofol consumption and shortens recovery times in outpatient anaesthesia. *British Journal of Anaesthesia* 2013;**110**(4):622-8. [PUBMED: 23220856]

Bestas 2004 {published data only}

Bestas A, Yasar MA, Bayar MK, Inalkac S. The effects of two different anaesthesia techniques on bispectral index values and awareness during off-pump coronary artery bypass grafting. *Journal of Clinical Monitoring and Computing* 2004;**18**:347-51. [PUBMED: 15957626]

Bethune 1992 {published data only}

Bethune DW, Ghosh S, Gray B, Kerr L, Walker IA, Doolan LA, et al. Learning during general anaesthesia: implicit recall after methohexitone or propofol infusion. *British Journal of Anaesthesia* 1992;**69**(2):197-9. [PUBMED: 1389826]

Bhawna 2012 {published data only}

Bhawna, Bajwa SJS, Lalitha K, Dhar P, Kumar V. Influence of esmolol on requirement of inhalational agent using entropy and assessment of its effect on immediate postoperative pain score. *Indian Journal of Anaesthesia* 2012;**56**(6):535-41. [PUBMED: 23325937]

Blendinger 1976 {published data only}

Blendinger I, Patschke D. Experiences with continuous infusions of etomidate in cardiac surgery. *Der Anaesthesist* 1976; Vol. 25, issue 8:391-2. [PUBMED: 949108]

Block 1991 {published data only}

Block RI, Ghoneim MM, Sum Ping ST, Ali MA. Human learning during general anaesthesia and surgery. *British Journal of Anaesthesia* 1991;**66**(2):170-8. [PUBMED: 1817616]

Bonato 2001 {published data only}

Bonato R, Sorbara C, Mapelli D, Iannizz P, Volpe B, Pittarello D, et al. Bispectral EEG analysis: a target of hypnosis during general anesthesia in cardiac surgery. *Acta Anaesthesiologica Italica/Anaesthesia and Intensive Care in Italy* 2001;**52**:83-102.

Browne 1973 {published data only}

Browne RA, Catton DV. Awareness during anaesthesia: a comparison of anaesthesia with nitrous oxide-oxygen and nitrous oxide-oxygen with innovar. *Canadian Anaesthetists' Society Journal* 1973;**20**(6):763-8. [PUBMED: 4769611]

Bruhn 2005 {published data only}

Bruhn J, Kreuer S, Bischoff P, Kessler P, Schmidt GN, Grzesiak A, et al. Bispectral index and A-line AAI index as guidance for desflurane-remifentanyl anaesthesia compared with a

standard practice group: a multi centre study. *British Journal of Anaesthesia* 2005;**94**(1):63-9. [PUBMED: 15516347]

Casati 1999 {published data only}

Casati A, Valentini G, Zangrillo A, Senatore R, Mello A, Airaghi B, et al. Anaesthesia for ultrasound guided oocyte retrieval: midazolam/remifentanyl versus propofol/fentanyl regimens. *European Journal of Anaesthesiology* 1999;**16**(11):773-8. [PUBMED: 10713871]

Celebioglu 2002 {published data only}

Celebioglu B, Pamuk AG, Aypar U, Pasaoglu I. Use of sevoflurane during cardiopulmonary bypass decreases incidence of awareness. *European Journal of Anaesthesiology* 2002;**19**(4):283-7. [PUBMED: 12074418]

Celleno 1993 {published data only}

Celleno D, Capogna G, Emanuelli M, Varrassi G, Muratori F, Costantino P, et al. Which induction drug for cesarean section? A comparison of thiopental sodium, propofol, and midazolam. *Journal of Clinical Anesthesia* 1993;**5**:284-8. [PUBMED: 8373604]

Chen 2009 {published data only}

Chen G, Buell O, Gruenewald M, Fudickar A, Scholz J, Bein B. A comparison between target-controlled and manually controlled propofol infusions in patients undergoing routine surgical procedures. *European Journal of Anaesthesiology* 2009;**26**(11):928-35. [PUBMED: 19842242]

Cheun 1987 {published data only}

Cheun JK. Intraspinal narcotic anesthesia in open heart surgery. *Journal of Korean Medical Science* 1987;**2**(4):225-9. [PUBMED: 3268179]

Chin 2004 {published data only}

Chin KJ, Yeo SW. Bispectral index values at sevoflurane concentrations of 1% and 1.5% in lower segment cesarean delivery. *Anesthesia and Analgesia* 2004;**98**(4):1140-4, table of contents. [PUBMED: 15041614]

Choi 2012 {published data only}

Choi WJ, Kim SH, Koh WU, Hwang DI, Cho SK, Park PH, et al. Effect of pre-exposure to sevoflurane on the bispectral index in women undergoing Caesarean delivery under general anaesthesia. *British Journal of Anaesthesia* 2012;**108**(6):990-7. [PUBMED: 22434266]

Clyburn 1986 {published data only}

Clyburn P, Kay NH, McKenzie PJ. Effects of diazepam and midazolam on recovery from anaesthesia in outpatients. *British Journal of Anaesthesia* 1986;**58**(8):872-5. [PUBMED: 2942163]

Coates 1987 {published data only}

Coates DP, Monk CR, Prys-Roberts C, Turtle M. Hemodynamic effects of infusions of the emulsion formulation of propofol during nitrous oxide anesthesia in humans. *Anesthesia and Analgesia* 1987;**66**(1):64-70. [PUBMED: 3492156]

Collins 1996 {published data only}

Collins SJ, Robinson AL, Holland HF. A comparison between total intravenous anaesthesia using a propofol/

alfentanil mixture and an inhalational technique for laparoscopic gynaecological sterilization. *European Journal of Anaesthesiology* 1996;**13**(1):33-7. [PUBMED: 8829934]

Crawford 1985 {published data only}

Crawford JS, Lewis M, Davies P. Maternal and neonatal responses related to the volatile agent used to maintain anaesthesia at caesarean section. *British Journal of Anaesthesia* 1985;**57**(5):482-7. [PUBMED: 3994881]

Crawford ME 1984 {published data only}

Crawford ME, Carl P, Andersen RS, Mikkelsen BO. Comparison between midazolam and thiopentone-based balanced anaesthesia for day-case surgery. *British Journal of Anaesthesia* 1984;**56**:165-9. [PUBMED: 6691877]

Czarko 2013 {published data only}

Czarko K, Kwiatosz-Muc M, Fijalkowska A, Kowalczyk M, Rutyna R. Intraoperative awareness--comparison of its incidence in women undergoing general anaesthesia for Caesarean section and for gynaecological procedures. *Anaesthesiology Intensive Therapy* 2013;**45**(4):200-4. [PUBMED: 24407896]

Deeprise 2005 {published data only}

Deeprise C, Andrade J, Harrison D, Edwards N. Unconscious auditory priming during surgery with propofol and nitrous oxide anaesthesia: a replication. *British Journal of Anaesthesia* 2005;**94**(1):57-62. [PUBMED: 15486010]

De Kock 1995 {published data only}

De Kock M, Famenne F, Deckers G, Scholtes JL. Epidural clonidine or sufentanil for intraoperative and postoperative analgesia. *Anesthesia and Analgesia* 1995;**81**(6):1154-62. [PUBMED: 7486097]

Deshpande 2009 {published data only}

Deshpande CM, Mohite SN, Kamdi P. Sufentanil vs fentanyl for fast-track cardiac anaesthesia. *Indian Journal of Anaesthesia* 2009;**53**(4):455-62. [PUBMED: 20640208]

Dhadphale 1979 {published data only}

Dhadphale PR, Jackson AP, Alseri S. Comparison of anaesthesia with diazepam and ketamine vs. morphine in patients undergoing heart-valve replacement. *Anesthesiology* 1979;**51**:200-3.

Drover 2002 {published data only}

Drover DR, Lemmens HJ, Pierce ET, Plourde G, Loyd G, Ornstein E, et al. Patient State Index: titration of delivery and recovery from propofol, alfentanil, and nitrous oxide anaesthesia. *Anesthesiology* 2002;**87**:82-9. [PUBMED: 12131107]

Dunnett 1977 {published data only}

Dunnett IA. Awareness during endotracheal intubation: a comparison of ketamine and thiopentone. *British Journal of Anaesthesia* 1977;**49**(5):491. [PUBMED: 861118]

Echevarria 1998 {published data only}

Echevarría M, Caba F, Rodríguez J, Olmedo L, Avila C, Vázquez T, et al. Explicit and implicit memory during inhalation and

intravenous anaesthesia. *Revista Espanola de Anestesiologia y Reanimacion* 1998;**45**(6):220-5. [PUBMED: 9719718]

Elhakim 2010 {published data only}

Elhakim M, Abdelhamid D, Abdelfattach H, Magdy H, Elsayed A, Elshafei M. Effect of epidural dexmedetomidine on intraoperative awareness and post-operative pain after one-lung ventilation. *Acta Anaesthesiologica Scandinavica* 2010;**54**(6):703-9. [PUBMED: 20085547]

Ellingson 1977 {published data only}

Ellingson A, Haram K, Sagen N. Ketamine and diazepam as anaesthesia for forceps delivery. A comparative study. *Acta Anaesthesiologica Scandinavica* 1977;**21**:37-40. [PUBMED: 320809]

Fehr 2001 {published data only}

Fehr SB, Zalunardo MP, Seifert B, Rentsch KM, Rohling RG, Pasch T, et al. Clonidine decreases propofol requirements during anaesthesia: effect on bispectral index. *British Journal of Anaesthesia* 2001; Vol. 86, issue 5:627-32. [PUBMED: 11575336]

Forestier 2003 {published data only}

Forestier F, Hirschi M, Rouget P, Rigal JC, Videcoq M, Girardet P, et al. Propofol and sufentanil titration with the bispectral index to provide anaesthesia for coronary artery surgery. *Anesthesiology* 2003;**99**(2):334-46. [PUBMED: 12883406]

Fragen 1981 {published data only}

Fragen RJ, Caldwell NJ. Awakening characteristics following anaesthesia induction with midazolam for short surgical procedures. *Arzneimittel-Forschung* 1981; Vol. 31, issue 12a:2261-3. [PUBMED: 7199331]

Gaitini 1995 {published data only}

Gaitini L, Vaida S, Collins G, Somri M, Sabo E. Awareness detection during caesarean section under general anaesthesia using EEG spectrum analysis. *Canadian Journal of Anaesthesia* 1995;**42**(5):377-81. [PUBMED: 7614642]

Gale 2001 {published data only}

Gale T, Leslie K, Kluger M. Propofol anaesthesia via target controlled infusion or manually controlled infusion: effects on the bispectral index as a measure of anaesthetic depth. *Journal Anaesthesia Intensive Care* 2001; Vol. 29, issue 6:579-84. [PUBMED: 11771598]

Ghaly 1988 {published data only}

Ghaly RG, Flynn RJ, Moore J. Isoflurane as an alternative to halothane for caesarean section. *Anaesthesia* 1988;**43**(1):5-7. [PUBMED: 3344947]

Ghoneim 2000 {published data only}

Ghoneim MM, Block RI, Dhanaraj VJ, Todd MM, Choi WW, Brown CK. Auditory evoked responses and learning and awareness during general anaesthesia. *Acta Anaesthesiologica Scandinavica* 2000; Vol. 44, issue 2:133-43. [PUBMED: 10695905]

Ghosh 2008 {published data only}

Ghosh I, Bithal PK, Dash HH, Chaturvedi A, Prabhakar H. Both clonidine and metoprolol modify anaesthetic depth indicators

and reduce intraoperative propofol requirement. *Journal of Anesthesia* 2008;**22**(2):131-4. [PUBMED: 18500609]

Girardi 1994 {published data only}

Girardi G, Rossi R, Cellai MP, Pieraccioli E, Novelli GP. Anesthesia with isoflurane in air and with isoflurane and nitrous oxide. *Minerva Anestesiologica* 1994;**60**(6):321-8.

Gokce 2009 {published data only}

Gokce BM, Karabiyik L, Karadenizli Y. Hypotensive anesthesia with esmolol. Assessment of hemodynamics, consumption of anesthetic drugs, and recovery. *Saudi Medical Journal* 2009;**30**(6):771-7. [PUBMED: 19526158]

Goto 2000 {published data only}

Goto T, Nakata Y, Saito H, Ishiguro Y, Niimi Y, Suwa K, et al. Bispectral analysis of the electroencephalogram does not predict responsiveness to verbal command in patients emerging from xenon anaesthesia. *British Journal of Anaesthesia* 2000;**85**(3):359-63. [PUBMED: 11103174]

Gruenewald 2007 {published data only}

Gruenewald M, Zhou J, Schloemerkerper N, Meybohm P, Weiler N, Tonner PH, et al. M-Entropy guidance vs standard practice during propofol-remifentanyl anaesthesia: a randomised controlled trial. *Anaesthesia* 2007;**62**:1124-9. [PUBMED: 17991257]

Grundmann 2001 {published data only}

Grundmann U, Silomon M, Bach F, Becker S, Bauer M, Larsen B, et al. Recovery profile and side effects of remifentanyl-based anaesthesia with desflurane or propofol for laparoscopic cholecystectomy. *Acta Anaesthesiologica Scandinavica* 2001;**45**:320-6. [PUBMED: 11207468]

Gupta 1992 {published data only}

Gupta A, Larsen LE, Sjöberg F, Lindh ML, Lennmarken C. Thiopentone or propofol for induction of isoflurane-based anaesthesia for ambulatory surgery?. *Acta Anaesthesiologica Scandinavica* 1992;**36**:670-4. [PUBMED: 1441867]

Gurman 1994 {published data only}

Gurman GM, Fajer S, Porat A, Schily M, Pearlman A. Use of EEG spectral edge as index of equipotency in a comparison of propofol and isoflurane for maintenance of general anaesthesia. *European Journal of Anaesthesiology* 1994; Vol. 11, issue 6:443-8. [PUBMED: 7851350]

Hachero 2001 {published data only}

Hachero A, Alamo F, Caba F, Echevarria M, Merino S, Gomez P, et al. Influence of bispectral index monitoring on fentanyl requirements during total intravenous anesthesia for major gynecological surgery. *Revista Espanola de Anestesiologia y Reanimacion* 2001;**48**(8):364-9. [PUBMED: 11674982]

Hackner 2003 {published data only}

Hackner C, Detsch O, Schneider G, Jelen-Esselborn S, Kochs E. Early recovery after remifentanyl-pronounced compared with propofol-pronounced total intravenous anaesthesia for short painful procedures. *British Journal of Anaesthesia* 2003; Vol. 91, issue 4:580-2. [PUBMED: 14504162]

Haimeur 1997 {published data only}

Haimeur CT, Alaoui I. General anaesthesia for caesarean section. A comparison of thiopental and ketamine. *Cahiers d'Anesthesiologie* 1997;**45**:9-13.

Haram 1981 {published data only}

Haram K, Lund T, Sagen N, Bøe OE. Comparison of thiopentone and diazepam as induction agents of anaesthesia for Caesarean section. *Acta Anaesthesiologica Scandinavica* 1981;**25**(6):470-6. [PUBMED: 7347074]

Hug 1988 {published data only}

Hug CC Jr, Hall RI, Angert KC, Reeder DA, Moldenhauer CC. Alfentanil plasma concentration v. effect relationships in cardiac surgical patients. *British Journal of Anaesthesia* 1988; Vol. 61, issue 4:435-40. [PUBMED: 3142509]

Hung 1992 {published data only}

Hung OR, Varvel JR, Shafer SL, Stanski DR. Thiopental pharmacodynamics. II. Quantitation of clinical and electroencephalographic depth of anesthesia. *Anesthesiology* 1992; Vol. 77, issue 2:237-44. [PUBMED: 1642341]

Ibraheim 2008 {published data only}

Ibraheim O, Alshaer A, Mazen K, El-Dawlaty A, Turkistani A, Alkathery K, et al. Effect of bispectral index (BIS) monitoring on postoperative recovery and sevoflurane consumption among morbidly obese patients undergoing laparoscopic gastric banding. *Middle East Journal Anesthesiology* 2008;**19**(4):819-30. [PUBMED: 18630768]

Inoue 2005 {published data only}

Inoue Y, Koga K, Sata T, Shigematsu A. Effects of fentanyl on emergence characteristics from anesthesia in adult cervical spine surgery: a comparison of fentanyl-based and sevoflurane-based anesthesia. *Journal of Anesthesia* 2005; Vol. 19, issue 1:12-6. [PUBMED: 15674509]

Jensen 1995 {published data only}

Jensen AG, Møller JT, Lybecker H, Hansen PA. A random trial comparing recovery after midazolam-alfentanil anesthesia with and without reversal with flumazenil, and standardized neurolept anesthesia for major gynecologic surgery. *Journal of Clinical Anesthesia* 1995;**7**:63-70. [PUBMED: 7772362]

Jiahai 2012 {published data only}

Jiahai M, Xueyan W, Yonggang X, Jianhong Y, Qunhui H, Zhi L, et al. Spectral entropy monitoring reduces anesthetic dosage for patients undergoing off-pump coronary artery bypass graft surgery. *Journal of Cardiothoracic and Vascular Anesthesia* 2012;**26**(5):818-21. [PUBMED: 22502772]

Kamal 1990 {published data only}

Kamal RS, Khan FA, Khan FH. Total intravenous anaesthesia with propofol and buprenorphine. *Anaesthesia* 1990;**45**(10):865-70. [PUBMED: 2240504]

Kasmacher 1996 {published data only}

Käsmacher H, Petermeyer M, Decker C. Incidence and quality of dreaming during anesthesia with propofol in comparison with enflurane. *Der Anaesthesist* 1996;**45**:146-53. [PUBMED: 8720887]

Kerssens 2005b {published data only}

Kerssens C, Ouchi T, Sebel PS. No evidence of memory function during anesthesia with propofol or isoflurane with close control of hypnotic state. *Anesthesiology* 2005;**103**:11-9. [PUBMED: 15618787]

Kerssens 2009 {published data only}

Kerssens C, Gaither JR, Sebel PS. Preserved memory function during bispectral index-guided anesthesia with sevoflurane for major orthopedic surgery. *Anesthesiology* 2009;**111**(3):518-24. [PUBMED: 19672180]

Kim 2007 {published data only}

Kim JH, Lee YS, Kim WY, Kim HJ, Chang MS, Park JY, et al. Effect of nicardipine on haemodynamic and bispectral index changes following endotracheal intubation. *Journal of International Medical Research* 2007;**35**:52-8. [PUBMED: 17408055]

Kiyama 1997 {published data only}

Kiyama S, Takeda J. Effect of extradural analgesia on the paradoxical arousal response of the electroencephalogram. *British Journal of Anaesthesia* 1997;**79**(6):750-3.

Kreuer 2003 {published data only}

Kreuer S, Biedler A, Larsen R, Altmann S, Wilhelm W. Narcotrend monitoring allows faster emergence and a reduction of drug consumption in propofol-remifentanyl anesthesia. *Anesthesiology* 2003;**99**(1):34-41. [PUBMED: 12826839]

Kreuer 2005 {published data only}

Kreuer S, Bruhn J, Stracke C, Aniset L, Silomon M, Larsen R, et al. Narcotrend or bispectral index monitoring during desflurane-remifentanyl anesthesia: a comparison with a standard practice protocol. *Anesthesia and Analgesia* 2005;**101**(2):427-34, table of contents. [PUBMED: 16037157]

Krissel 1994 {published data only}

Krissel J, Dick WF, Leyser KH, Gervais H, Brockerhoff P, Schranz D. Thiopentone, thiopentone/ketamine, and ketamine for induction of anaesthesia in caesarean section. *European Journal of Anaesthesiology* 1994;**11**:115-22. [PUBMED: 8174531]

Kudoh 1999 {published data only}

Kudoh A, Matsuki A. Middle latency auditory evoked potentials during total intravenous anesthesia with droperidol, ketamine and fentanyl. *Clinical Electroencephalography* 1999;**30**(2):79-83. [PUBMED: 10358788]

Kwon 2013 {published data only}

Kwon MA, Song J, Kim J R. Tracheal intubation with rocuronium using a "modified timing principle". *Korean Journal of Anesthesiology* 2013; Vol. 64, issue 3:218-22. [PUBMED: 23560186]

Lallemand 2003 {published data only}

Lallemand MA, Lentschener C, Mazoit JX, Bonnichon P, Manceau I, Ozier Y. Bispectral index changes following etomidate induction of general anaesthesia and orotracheal intubation. *British Journal of Anaesthesia* 2003;**91**(3):341-6. [PUBMED: 12925471]

Lam 2013 {published data only}

Lam F, Liao CC, Lee YJ, Wang W, Kuo CJ, Lin CS. Different dosing regimens for propofol induction in obese patients. *Anaesthesia* 2013; Vol. 51, issue 2:53-7.

Lehmann 1985 {published data only}

Lehmann KA, Horrichs G, Hoeckle W. The significance of tramadol as an intraoperative analgesic. A randomized double-blind study in comparison with placebo. *Der Anaesthesist* 1985;**34**(1):11-9. [PUBMED: 3883843]

Lehmann 1992 {published data only}

Lehmann KA, Krauskopf KH. Intraoperative awareness in balanced anesthesia. A literature review based on a randomized double blind study using fentanyl, pentazocine and ketamine. *Der Anaesthesist* 1992;**41**:373-85. [PUBMED: 1497125]

Lehmann 2007 {published data only}

Lehmann A, Schmidt M, Zeitler C, Kiessling AH, Isgro F, Boldt J. Bispectral index and electroencephalographic entropy in patients undergoing aortocoronary bypass grafting. *European Journal of Anaesthesiology* 2007;**27**:751-60. [PUBMED: 17241504]

Lim 1992 {published data only}

Lim BL, Low TC. Total intravenous anaesthesia versus inhalational anaesthesia for dental day surgery. *Anaesthesia and Intensive Care* 1992;**20**:475-78. [PUBMED: 1463176]

Lin 2011 {published data only}

Lin CF, Yang CY, Chao E, Lee MS, See LC. Short-term inhalation of sevoflurane during induction of general anesthesia can inhibit the A-line ARX index response to intubation: a randomized trial. *Chang Gung Medical Journal* 2011;**34**(6):599-606. [PUBMED: 2219606]

Lindholm 2008 {published data only}

Lindholm ML, Brudin L, Sandin RH. Bispectral index monitoring: appreciated but does not effect drug dosing and hypnotic levels. *Acta Anaesthesiologica Scandinavica* 2008;**52**:88-94. [PUBMED: 17976226]

Liu 2013 {published data only}

Liu N, Pruszkowski O, Leroy J E, Chazot T, Trillat B, Colchen A, et al. Automatic administration of propofol and remifentanyl guided by the bispectral index during rigid bronchoscopic procedures: a randomized trial. *Canadian Journal of Anesthesia* 2013;**60**(9):881-7. [PUBMED: 23857041]

Lu 2005 {published data only}

Lu CH, Borel CO, Wu CT, Yeh CC, Jao SW, Chao PC, et al. Combined general-epidural anesthesia decreases the desflurane requirement for equivalent A-line ARX index in colorectal surgery. *Acta Anaesthesiologica Scandinavica* 2005;**49**:1063-7. [PUBMED: 16095444]

Maattanen 2002 {published data only}

Maattanen H, Anderson R, Uusijarvi J, Jakobsson J. Auditory evoked potential monitoring with the AAITM-index during spinal surgery: decreased desflurane consumption. *Acta*

- Anaesthesiologica Scandinavica* 2002;**46**(7):882-6. [PUBMED: 12139546]
- Mashour 2012** {published and unpublished data}
 * Mashour GA, Shanks A, Tremper KK, Kheterpal S, Turner CR, Ramachandran SK, et al. Prevention of intraoperative awareness with explicit recall in an unselected surgical population: a randomized comparative effectiveness trial. *Anesthesiology* 2012;**117**(4):717-25. [Pubmed: 22990178]
- Masuda 2002** {published data only}
 Masuda T, Yamada H, Takada K, Sagata Y, Yamaguchi M, Tomiyama Y, et al. Bispectral index monitoring is useful to reduce total amount of propofol and to obtain immediate recovery after propofol anesthesia. *Masui* 2002;**51**(4):394-9. [PUBMED: 11995347]
- McNulty 1995** {published data only}
 McNulty SE, Gratch D, Costello D, Heropoulos M, Kim JY. The effect of midazolam and lorazepam on postoperative recovery after cardiac surgery. *Anesthesia and Analgesia* 1995;**81**(2):404-7. [PUBMED: 7618736]
- Menigaux 2002** {published data only}
 Menigaux C, Guignard B, Adam F, Sessler DI, Joly V, Chauvin M. Esmolol prevents movement and attenuates the BIS response to orotracheal intubation. *British Journal of Anaesthesia* 2002;**89**(6):857-62. [PUBMED: 12453930]
- Mertens 2003** {published data only}
 Mertens MJ, Olofsen E, Engbers FH, Burm AG, Bovill JG, Vuyk J. Propofol reduces perioperative remifentanyl requirements in a synergistic manner: response surface modeling of perioperative remifentanyl-propofol interactions. *Anesthesiology* 2003;**99**(2):347-59. [PUBMED: 12883407]
- Miller 1996** {published data only}
 Miller DR, Blew PG, Martineau RJ, Hull KA. Midazolam and awareness with recall during total intravenous anaesthesia. *Canadian Journal of Anaesthesia* 1996;**43**(9):946-53. [PUBMED: 8874913]
- Miranda 1992** {published data only}
 Miranda AF, Kyi W, Sivalingam N. Propofol and methohexitone for elective caesarean--a comparative study. *Medical Journal of Malaysia* 1992;**47**(4):280-6. [PUBMED: 1303480]
- Monedero 1994** {published data only}
 Monedero P, Carrascosa F, Garcia-Pedrajas F, Panadero A, Hidalgo F, Arroyo JL. Does propofol have advantages over midazolam and isoflurane? Comparative study of 2 total intravenous anesthesia techniques using midazolam and propofol, versus balanced anesthesia with isoflurane. *Revista Espanola de Anestesiologia y Reanimacion* 1994;**41**(3):156-64. [PUBMED: 8059043]
- Morimoto 2002** {published data only}
 Morimoto Y, Oka S, Mii M, Shinjo Y, Yamashita A, Gohara T, et al. Efficacy of bispectral index monitoring in improving anesthetic management, economics, and use of the operating theater. *Masui* 2002;**51**(8):862-8. [PUBMED: 12229134]
- Mozafari 2014** {published data only}
 Mozafari H, Fakhr AA, Salehi I, Moghimbegi A. The ability of bispectral-guided management compared to routine monitoring for reflecting awareness rate in patients undergoing abdominal surgery. *Iranian Red Crescent Medical Journal* 2014;**16**(9):e13584.
- Muralidhar 2008** {published data only}
 Muralidhar K, Banakal S, Murthy K, Garg R, Rani GR, Dinesh R. Bispectral index-guided anaesthesia for off-pump coronary artery bypass grafting. *Annals of Cardiac Anesthesia* 2008;**11**(2):105-10. [PUBMED: 18603750]
- Myles 1997** {published data only}
 Myles PS, Buckland MR, Weeks AM, Bujur MA, McRae R, Langley M, et al. Hemodynamic effects, myocardial ischemia, and timing of tracheal extubation with propofol-based anesthesia for cardiac surgery. *Anesthesia and Analgesia* 1997;**84**(1):12-9. [PUBMED: 8988992]
- Myles 2004** {published and unpublished data}
 * Myles PS, Leslie K, McNeil J, Forbes A, Chan MT. Bispectral index monitoring to prevent awareness during anaesthesia: the B-Aware randomised controlled trial. *Lancet* 2004;**363**:1757-63. [PUBMED: 15172773]
- Myles 2007** {published data only}
 Myles PS, Leslie K, Chan MT, Forbes A, Paech MJ, Peyton P, et al. Avoidance of nitrous oxide for patients undergoing major surgery: a randomized controlled trial. *Anesthesiology* 2007;**107**(2):221-31. [PUBMED: 17667565]
- Navarro 2000** {published data only}
 Navarro EM. Desflurane--general anesthesia for cesarean section compared with isoflurane and epidural anesthesia. *Anesthesiologie, Intensivmedizin, Notfallmedizin, Schmerztherapie* 2000;**35**(4):232-6. [PUBMED: 10830075]
- Nayar 2009** {published data only}
 Nayar R, Sahajanand H. Does anesthetic induction for Cesarean section with a combination of ketamine and thiopentone confer any benefits over thiopentone or ketamine alone? A prospective randomized study. *Minerva Anestesiologica* 2009;**75**(4):185-90. [PUBMED: 18946429]
- Ngan 1997** {published data only}
 Ngan Kee WD, Khaw KS, Ma ML, Mainland PA, Gin T. Postoperative analgesic requirement after cesarean section: a comparison of anesthetic induction with ketamine or thiopental. *Anesthesia and Analgesia* 1997;**85**(6):1294-8.
- Ngan Kee 2002** {published data only}
 Ngan Kee WD, Khaw KS, Ma KC, Wong AS, Lee BB. Randomized, double-blind comparison of different inspired oxygen fractions during general anaesthesia for Caesarean section. *British Journal of Anaesthesia* 2002;**89**(4):556-61. [PUBMED: 12393355]
- Oddby-Muhrbeck 1993** {published data only}
 Oddby-Muhrbeck E, Jakobsson J. Recall of music: a comparison between anaesthesia with propofol and isoflurane. *Acta*

Anaesthesiologica Scandinavica 1993;**37**:33-7. [PUBMED: 8424291]

Panousis 2009 {published data only}

Panousis P, Heller AR, Koch T, Litz RJ. Epidural ropivacaine concentrations for intraoperative analgesia during major upper abdominal surgery: a prospective, randomized, double-blinded, placebo-controlled study. *Anesthesia and Analgesia* 2009;**108**(6):1971-6. [PUBMED: 19448234]

Pauls 2009 {published data only}

Pauls RJ, Dickson TJ, Kaufmann AM, Cappellani RB, Ringaert KR, West M, et al. A comparison of the ability of the EEGo and BIS monitors to assess emergence following neurosurgery. *Canadian Journal of Anaesthesia* 2009;**56**(5):366-73. [PUBMED: 19340492]

Paventi 2001 {published data only}

Paventi S, Santevecchi A, Metta E, Annetta MG, Perilli V, Sollazzi L, et al. Bispectral index monitoring in sevoflurane and remifentanil anesthesia. Analysis of drugs management and immediate recovery. *Minerva Anestesiologica* 2001;**67**(6):435-9. [PUBMED: 11533541]

Pedersen 1992 {published data only}

Pedersen JE, Fernandes A, Christensen M. Halothane 2% for caesarean section. *European Journal of Anaesthesiology* 1992;**9**:319-24. [PUBMED: 1628635]

Persec 2012 {published data only}

Persec J, Persec Z, Kopljär M, Sojic N, Husedzinovic I. Effect of bispectral index monitoring on extubation time and analgesic consumption in abdominal surgery: a randomised clinical trial. *Swiss Medical Weekly* 2012; Vol. 142:w13689. [PUBMED: 23135976]

Piggott 1990 {published data only}

Piggott SE, Bogod DG, Rosen M, Rees GA, Harmer M. Isoflurane with either 100% oxygen or 50% nitrous oxide in oxygen for caesarean section. *British Journal of Anaesthesia* 1990;**65**:325-9. [PUBMED: 2121203]

Plourde 1996 {published data only}

Plourde G, Villemure C. Comparison of the effects of enflurane/N₂O on the 40-Hz auditory steady-state response versus the auditory middle-latency response. *Anesthesia and Analgesia* 1996;**82**:75-83. [PUBMED: 8712429]

Puri 2003 {published data only}

Puri GD, Murthy SS. Bispectral index monitoring in patients undergoing cardiac surgery under cardiopulmonary bypass. *European Journal of Anaesthesiology* 2003;**20**:451-6. [PUBMED: 12803261]

Puri 2007 {published data only}

Puri GD, Kumar B, Aveek J. Closed-loop anaesthesia delivery system (CLADS) using bispectral index: a performance assessment study. *Anaesthesia and Intensive Care* 2007;**35**(3):357-62. [PUBMED: 17591128]

Rehberg 2007 {published data only}

Rehberg B, Ryll C, Hadzidiakos D, Baars J. Use of a target-controlled infusion system for propofol does not improve subjective assessment of anaesthetic depth by inexperienced anaesthesiologists. *European Journal of Anaesthesiology* 2007;**24**:920-6. [PUBMED: 17588273]

Renna 2000 {published data only}

Renna M, Lang EM, Lockwood GG. The effect of sevoflurane on implicit memory: a double-blind, randomised study. *Anaesthesia* 2000;**55**(7):634-40. [PUBMED: 10919417]

Rinaldi 2005 {published data only}

Rinaldi S, Consales G, Gallerani E, Ortolani O, De Gaudio AR. A-line autoregression index monitoring to titrate inhalational anaesthesia: effects on sevoflurane consumption, emergence time and memory. *Acta Anaesthesiologica Scandinavica* 2007;**49**:692-7. [PUBMED: 15836686]

Russell 1986 {published data only}

Russell IF. Comparison of wakefulness with two anaesthetic regimens. Total i.v. v. balanced anaesthesia. *British Journal of Anaesthesia* 1986;**58**(9):965-8. [PUBMED: 3756056]

Samarkandi 2004 {published data only}

Samarkandi AH, Abdel Meguid ME, Abdullah KM, Riad W. Bispectral index monitoring and titration of anaesthetics during off-pump coronary artery bypass surgery. *Egyptian Journal of Anaesthesia* 2004;**20**:357-61.

Sareen 1997 {published data only}

Sareen J, Hudson RJ, Rosenbloom M, Thomson IR. Dose-response to anaesthetic induction with sufentanil: haemodynamic and electroencephalographic effects. *Canadian Journal of Anaesthesia* 1997;**44**(1):19-25. [PUBMED: 8988819]

Schultetus 1986 {published data only}

Schultetus RR, Hill CR, Dharamraj CM, Banner TE, Berman LS. Wakefulness during cesarean section after anesthetic induction with ketamine, thiopental, or ketamine and thiopental combined. *Anesthesia and Analgesia* 1986;**65**:723-8. [PUBMED: 3717612]

Schwender 1994 {published data only}

Schwender D, Kaiser A, Klasing S, Peter K, Pöppel E. Midlatency auditory evoked potentials and explicit and implicit memory in patients undergoing cardiac surgery. *Anesthesiology* 1994;**80**:493-501. [PUBMED: 8141445]

Schwender 1996 {published data only}

Schwender D, Daunderer M, Mulzer S, Klasing S, Finsterer U, Peter K. Spectral edge frequency of the electroencephalogram to monitor "depth" of anaesthesia with isoflurane or propofol. *British Journal of Anaesthesia* 1996;**77**:179-84. [PUBMED: 8881621]

Shin 2012 {published data only}

Shin MK, Shim HS, Yang GY, Sung WS. Effect of a target-controlled infusion of remifentanil in combination with desflurane during the "maintenance" phase of general

- anesthesia. *Korean Journal of Anesthesiology* 2012;**63**(1):25-9. [PUBMED: 22870361]
- Sidi 1990** {published data only}
 Sidi A, Halimi P, Cotev S. Estimating anesthetic depth by electroencephalography during anesthetic induction and intubation in patients undergoing cardiac surgery. *Journal of Clinical Anesthesia* 1990;**2**(2):101-7.
- Smith 1999** {published data only}
 Smith TL, Zapala D, Thompson CL, Hoye W, Kelly T. Relationship of auditory middle latency response and stem-word completion test as indicators of implicit memory formation during general anesthesia. *Journal of the American Association of Nurse Anesthetists* 1999;**67**(3):247-53. [PUBMED: 10488296]
- Song 1997** {published data only}
 Song D, Joshi GP, White PF. Titration of volatile anesthetics using bispectral index facilitates recovery after ambulatory anesthesia. *Anesthesiology* 1997;**87**(4):842-8. [PUBMED: 9357886]
- Sorbara 1995** {published data only}
 Sorbara C, Pittarello D, Rizzoli G, Pasini L, Armellin G, Bonato R, et al. Propofol-fentanyl versus isoflurane-fentanyl anesthesia for coronary artery bypass grafting: effect on myocardial contractility and peripheral hemodynamics. *Journal of Cardiothoracic and Vascular Anesthesia* 1995;**9**(1):18-23. [PUBMED: 7718750]
- Soyannwo 1988** {published data only}
 Soyannwo OA, Elegbe EO, Odugbesan CO. Effect of flunitrazepam (Rohypnol) on awareness during anaesthesia for caesarean section. *African Journal of Medicine and Medical Sciences* 1988;**17**(1):23-6. [PUBMED: 2834929]
- Staikou 2013** {published data only}
 Staikou C, Paraskeva A, Karmanioliou I, Vezakis A, Tsaroucha A. Intravenous lidocaine does not affect the anesthetic depth during rapid sequence induction and intubation as assessed by Bispectral Index monitoring: a randomized double blind study. *Archives of Medical Science* 2013; Vol. 9, issue 4:713-8. [PUBMED: 24049534]
- Stoppe 2012** {published data only}
 Stoppe C, Peters D, Fahlenkamp AV, Cremer J, Rex S, Schalte G, et al. aepEX monitor for the measurement of hypnotic depth in patients undergoing balanced xenon anaesthesia. *British Journal of Anaesthesia* 2012;**108**(1):80-8. [PUBMED: 22139005]
- Stuttman 2010** {published data only}
 Stuttman R, Jakubetz J, Schultz K, Schäfer C, Langer S, Ullmann U, et al. Recovery index, attentiveness and state of memory after xenon or isoflurane anaesthesia: a randomized controlled trial. *Anesthesiology* 2010;**100**:5. [PUBMED: 20459661]
- Toft 1987** {published data only}
 Toft P, Romer U. Comparison of midazolam and diazepam to supplement total intravenous anaesthesia with ketamine for endoscopy. *Canadian Journal of Anaesthesia* 1987;**34**(5):466-9. [PUBMED: 3311432]
- Toscano 2007** {published data only}
 Toscano A, Pancaro C, Peduto VA. Scopolamine prevents dreams during general anesthesia. *Anesthesiology* 2007;**106**(5):952-5. [PUBMED: 17457126]
- Tsai 2001** {published data only}
 Tsai PS, Huang CJ, Hung YC, Cheng CR. Effects on the bispectral index during elective caesarean section: a comparison of propofol and isoflurane. *Acta Anaesthesiologica Sinica* 2001;**39**:17-22. [PUBMED: 11407290]
- Tunstall 1989** {published data only}
 Tunstall ME, Sheikh A. Comparison of 1.5% enflurane with 1.25% isoflurane in oxygen for caesarean section: avoidance of awareness without nitrous oxide. *British Journal of Anaesthesia* 1989;**62**(2):138-43. [PUBMED: 2923765]
- Vakkuri 2005** {published data only}
 Vakkuri A, Yli-Hankala A, Sandin R, Mustola S, Hoymork S, Nyblom S, et al. Spectral entropy monitoring is associated with reduced propofol use and faster emergence in propofol-nitrous oxide-alfentanil anesthesia. *Anesthesiology* 2005;**103**(2):274-9. [PUBMED: 16052109]
- van der Maaten 1996** {published data only}
 van der Maaten JM, Epema AH, Huet RC, Hennis PJ. The effect of midazolam at two plasma concentrations of hemodynamics and sufentanil requirement in coronary artery surgery. *Journal of Cardiothoracic and Vascular Anesthesia* 1996;**10**(3):356-63. [PUBMED: 8725417]
- van Leeuwen 1990** {published data only}
 van Leeuwen L, Zuurmond WW, Deen L, Helmers HJ. Total intravenous anaesthesia with propofol, alfentanil, and oxygen-air: three different dosage schemes. *Canadian Journal of Anaesthesiology* 1990;**37**(3):282-6. [PUBMED: 2182204]
- Wang 2013** {published data only}
 Wang J, Ren Y, Zhu Y, Chen JW, Zhu MM, Xu YJ, et al. Effect of penehyclidine hydrochloride on the incidence of intra-operative awareness in Chinese patients undergoing breast cancer surgery during general anaesthesia. *Anaesthesia* 2013;**68**(2):136-41. [PUBMED: 23237605]
- White 2003** {published data only}
 White PF, Wang B, Tang J, Wender RH, Naruse R, Sloninsky A. The effect of intraoperative use of esmolol and nicardipine on recovery after ambulatory surgery. *Anesthesia and Analgesia* 2003;**97**(6):1633-8. [PUBMED: 14633533]
- White 2004** {published data only}
 White PF, Ma H, Tang J, Wender RH, Sloninsky A, Kariger R. Does the use of electroencephalographic bispectral index or auditory evoked potential index monitoring facilitate recovery after desflurane anesthesia in the ambulatory setting?. *Anesthesiology* 2004;**100**(4):811-7. [PUBMED: 15087615]
- Wong 2002** {published data only}
 Wong J, Song D, Blanshard H, Grady D, Chung F. Titration of isoflurane using BIS index improves early recovery of elderly

patients undergoing orthopedic surgeries. *Canadian Journal of Anaesthesia* 2002;**49**(1):13-8. [PUBMED: 11782323]

Wu 2001 {published data only}

Wu CC, Mok MS, Lin CS, Han SR. EEG-bispectral index changes with ketamine versus thiamylal induction of anesthesia. *Acta Anaesthesiologica Sinica* 2001;**39**(1):11-5. [PUBMED: 11407289]

Yildiz 2002 {published data only}

Yildiz K, Esmoğlu A, Doğru K, Tercan E, Boyacı A. A comparison of alfentanil and remifentanil accompanying with BIS monitoring for total intravenous anaesthesia. *Türk Anesteziyoloji ve Reanimasyon Derneği Dergisi [Journal of the Turkish Anaesthesiology and Intensive Care Society]* 2002;**30**:309-14.

Yoshitani 2003 {published data only}

Yoshitani K, Kawaguchi M, Takahashi M, Kitaguchi K, Furuya H. Plasma propofol concentration and EEG burst suppression ratio during normothermic cardiopulmonary bypass. *British Journal of Anaesthesia* 2003;**90**(2):122-6. [PUBMED: 12538365]

Zhang 2011 {published and unpublished data}

* Zhang C, Xu L, Ma YQ, Sun YX, Li YH, Zhang L, et al. Bispectral index monitoring prevent awareness during total intravenous anesthesia: a prospective, randomized, double-blinded, multi-center controlled trial. *Chinese Medical Journal* 2011;**124**(22):3664-9. [PUBMED: 22340221]

Zhou 2008 {published data only}

Zhou S-L, Zhang R, Shen N, Wei J-Q, Hei Z-Q. Comparison of induction anesthesia with sevoflurane inhalation and intravenous propofol in gynaecologic laparoscopy operation. *Zhongguo Xin Yao Za Zhi [Chinese Journal of New Drugs]* 2008;**17**(21):1880-3.

Zohar 2006 {published data only}

Zohar E, Luban I, White PF, Ramati E, Shabat S, Fredman B. Bispectral index monitoring does not improve early recovery of geriatric outpatients undergoing brief surgical procedures. *Canadian Journal of Anaesthesiology* 2006;**53**(1):20-5. [PUBMED: 16371605]

References to studies excluded from this review

Abboud 1990 {published data only}

Abboud TK, Afrasiabi A, Davidson J, Zhu J, Reyes A, Khoo N, et al. Prophylactic oral naltrexone with epidural morphine: effect on adverse reactions and ventilatory responses to carbon dioxide. *Anesthesiology* 1990;**72**(2):233-7. [PUBMED: 2105673]

Abdel-Meguid 2005 {published data only}

Abdel-Meguid ME. Bispectral index guided evaluation of the optimal midazolam induction dosing for coronary revascularization. *Egyptian Journal of Anaesthesia* 2003;**21**(2):101-4.

Abott 1980 {published data only}

Abott TR, Bentley G. Intra-operative awakening during scoliosis surgery. *Anaesthesia* 1980;**35**(3):298-302. [MEDLINE: 7396143]

Abouleish 1976 {published data only}

Abouleish E, Taylor FH. Effect of morphine-diazepam on signs of anesthesia, awareness, and dreams of patients under N2O for cesarean section. *Anesthesia and Analgesia* 1976;**55**(5):702-5. [PUBMED: 987727]

Absalom 2002 {published data only}

Absalom AR, Sutcliffe N, Kenny GN. Closed-loop control of anesthesia using bispectral index: performance assessment in patients undergoing major orthopedic surgery under combined general and regional anesthesia. *Anesthesiology* 2002;**96**(1):67-73. [PUBMED: 11753004]

Acil 2004 {published data only}

Acil M, Basgul E, Celiker V, Karagoz AH, Demir B. Perioperative effects of melatonin and midazolam premedication on sedation, orientation, anxiety scores and psychomotor performance. *European Journal of Anesthesiology* 2004;**21**(7):553-7. [PUBMED: 15318468]

Adams 1998 {published data only}

Adams DC, Hilton HJ, Madigan JD, Szerlip NJ, Cooper LA, Emerson RG. Evidence for unconscious memory processing during elective cardiac surgery. *Circulation* 1998;**98**(19 Suppl):II289-92; discussion II292-3. [PUBMED: 9852916]

Adams 2003 {published data only}

Adams HA, Meyer P, Stoppa A, Muller-Goch A, Bayer P, Hecker H. Anaesthesia for caesarean section. Comparison of two general anaesthetic regimens and spinal anaesthesia. *Der Anaesthesist* 2003;**52**(1):23-32. [PUBMED: 12577162]

Ahmad 2003 {published data only}

Ahmad S, Yilmaz M, Marcus RJ, Glisson S, Kinsella A. Impact of bispectral index monitoring on fast tracking of gynecologic patients undergoing laparoscopic surgery. *Anesthesiology* 2003;**98**(4):849-52. [PUBMED: 12657845]

Ahonen 2007 {published data only}

Ahonen J, Jokela R, Uutela K, Huiku M. Surgical stress index reflects surgical stress in gynaecological laparoscopic day-case surgery. *British Journal of Anaesthesia* 2007;**98**(4):456-61. [PUBMED: 17350969]

Alexander 1999a {published data only}

Alexander R, Booth J, Olufolabi AJ, El-Moalem HE, Glass PS. Comparison of remifentanil with alfentanil or suxamethonium following propofol anaesthesia for tracheal intubation. *Journal of Anaesthesia* 1999;**54**(11):1032-6. [PUBMED: 10540090]

Alexander 1999b {published data only}

Alexander R, Olufolabi AJ, Booth J, El-Moalem HE, Glass PS. Dosing study of remifentanil and propofol for tracheal intubation without the use of muscle relaxants. *Anaesthesia* 1999;**54**(11):1037-40. [PUBMED: 10540091]

Al-Ruzzeh 2006 {published data only}

Al-Ruzzeh S, George S, Bustami M, Wray J, Ilsley C, Athanasiou T, et al. Effect of off-pump coronary artery bypass surgery on clinical, angiographic, neurocognitive, and quality of life

outcomes: randomised controlled trial. *BMJ (Clinical research ed.)* 2006;**332**(7554):1365. [PUBMED: 16740529]

Alvarez 2000 {published data only}

Alvarez AO, Cascardo A, Albarracin Menendez S, Capria JJ, Cordero RA. Total intravenous anesthesia with midazolam, remifentanyl, propofol and cisatracurium in morbid obesity. *Obesity Surgery* 2000;**10**(4):353-60. [PUBMED: 11007629]

Andelman 2004 {published data only}

Andelman F, Zuckerman-Feldhay E, Hoffien D, Fried I, Neufeld MY. Lateralization of deficit in self-awareness of memory in patients with intractable epilepsy. *Epilepsia* 2004;**45**(7):826-33. [PUBMED: 15230708]

Anderson 2003 {published data only}

Anderson RE, Barr G, Assareh H, Jakobsson J. The AAI index, the BIS index and end-tidal concentration during wash in and wash out of sevoflurane. *Journal of Anesthesia* 2003;**58**(6):531-5. [PUBMED: 12846616]

Anderson 2004 {published data only}

Anderson RE, Barr G, Owall A, Jakobsson J. Entropy during propofol hypnosis, including an episode of wakefulness. *Journal of Anesthesia* 2004; Vol. 59, issue 1:52-6. [PUBMED: 14687099]

Andrade 2001 {published data only}

Andrade J, Englert L, Harper C, Edwards ND. Comparing the effects of stimulation and propofol infusion rate on implicit and explicit memory formation. *British Journal of Anaesthesia* 2001; Vol. 86, issue 2:189-95. [PUBMED: 11573658]

Aono 1999 {published data only}

Aono J, Mamiya K, Manabe M. Preoperative anxiety is associated with a high incidence of problematic behavior on emergence after halothane anesthesia in boys. *Acta Anaesthesiologica Scandinavica* 1999;**43**(5):542-4. [PUBMED: 10342002]

Apfelbaum 1996 {published data only}

Apfelbaum JL, Lichtor JL, Lane BS, Coalson DW, Korttila KT. Awakening, clinical recovery, and psychomotor effects after desflurane and propofol anesthesia. *Anesthesia and Analgesia* 1996;**83**(4):721-5. [PUBMED: 8831309]

Aqil 2009 {published data only}

Aqil M, Aziz-ul-Haq, Rasheed A, Altaf R. Effect of two different doses of fentanyl on intubating conditions with sevoflurane inhalation without neuromuscular blocking agents in adults. *Anaesthesia, Pain & Intensive Care* 2009;**13**(2):55-6.

Arndt 1995a {published data only}

Arndt VM, Hofmockel R, Benad G. EEG changes during propofol-alfentanil-nitrous oxide anesthesia. *Anesthesiologie und Reanimation* 1995; Vol. 20, issue 5:126-33. [PUBMED: 8526965]

Aubrun 2008 {published data only}

Aubrun F, Gaillat C, Rosenthal D, Dupuis M, Mottet P, Marchetti F, et al. Effect of a low-dose ketamine regimen on pain, mood, cognitive function and memory after major gynaecological surgery: a randomized, double-blind, placebo-controlled

trial. *European Journal of Anaesthesiology* 2008;**25**(2):97-105. [PUBMED: 17894912]

Ausems 1983 {published data only}

Ausems ME, Hug CCJ, de Lange S. Variable rate infusion of alfentanil as a supplement to nitrous oxide anesthesia for general surgery. *Anesthesia and Analgesia* 1983;**62**(11):982-6. [PUBMED: 6414340]

Ausems 1986 {published data only}

Ausems ME, Hug CC Jr, Stanski DR, Burm AG. Plasma concentrations of alfentanil required to supplement nitrous oxide anesthesia for general surgery. *Anesthesiology* 1986; Vol. 65, issue 4:362-73. [PUBMED: 3094407]

Bailey 1985 {published data only}

Bailey PL, Wilbrink J, Zwanikken P, Pace NL, Stanley TH. Anesthetic induction with fentanyl. *Anesthesia and Analgesia* 1985;**64**(1):48-53. [PUBMED: 2981489]

Balci 2006 {published data only}

Balci C, Akbulut G, Ibis AH, Arikan Y. Comparison of desflurane and sevoflurane recovery duration with BIS values. *Anestezi Dergisi [Journal of Anesthesia]* 2006;**14**(1):32-6.

Bannister 2001 {published data only}

Bannister CF, Brosius KK, Sigl JC, Meyer BJ, Sebel PS. The effect of bispectral index monitoring on anesthetic use and recovery in children anesthetized with sevoflurane in nitrous oxide. *Anesthesia and Analgesia* 2001;**92**(4):877-81. [PUBMED: 11273918]

Baraka 1990 {published data only}

Baraka A, Louis F, Dalleh R. Maternal awareness and neonatal outcome after ketamine induction of anaesthesia for Caesarean section. *Canadian Journal of Anaesthesia* 1990; Vol. 37, issue 6:641-4. [PUBMED: 2208534]

Barclay 1980 {published data only}

Barclay A, Houlton PC, Downing JW. Total intravenous anaesthesia: a technique using flunitrazepam, ketamine, muscle relaxants and controlled ventilation of the lung. *Anaesthesia* 1980;**35**(3):287-90. [PUBMED: 6104936]

Barr 1973 {published data only}

Barr AM, Wong RM. Awareness during general anaesthesia for bronchoscopy and laryngoscopy using the apnoeic oxygenation technique. *British Journal of Anaesthesia* 1973;**45**(8):894-900. [PUBMED: 4753687]

Barr 1999 {published data only}

Barr G, Jakobsson JG, Owall A, Anderson RE. Nitrous oxide does not alter bispectral index: study with nitrous oxide as sole agent and as an adjunct to i.v. anaesthesia. *British Journal of Anaesthesia* 1999;**82**(6):827-30. [PUBMED: 10562773]

Barr 2001 {published data only}

Barr G, Anderson RE, Owall A, Jakobsson JG. Being awake intermittently during propofol-induced hypnosis: a study of BIS, explicit and implicit memory. *Acta Anaesthesiologica Scandinavica* 2001;**45**(7):834-8. [PUBMED: 11472283]

Barvais 2003 {published data only}

Barvais L, Engelman E, Eba JM, Coussaert E, Cantraine F, Kenny GN. Effect site concentrations of remifentanyl and pupil response to noxious stimulation. *British Journal of Anaesthesia* 2003; Vol. 91, issue 3:347-52. [PUBMED: 12925472]

Basar 2003 {published data only}

Basar H, Ozcan S, Buyukkocak U, Akpınar S, Apan A. Effect of bispectral index monitoring on sevoflurane consumption. *European Journal of Anaesthesiology* 2003; Vol. 20, issue 5:396-400. [PUBMED: 12790212]

Becker-Blease 2006 {published data only}

Becker-Blease KA, Freyd JJ. Research participants telling the truth about their lives: the ethics of asking and not asking about abuse. *American Journal of Psychology* 2006; Vol. 61, issue 3:218-26. [PUBMED: 16594838]

Bejjani 2009 {published data only}

Bejjani G, Lequeux PY, Schmartz D, Engelman E, Barvais L. No evidence of memory processing during propofol-remifentanyl target-controlled infusion anesthesia with bispectral index monitoring in cardiac surgery. *Journal of Cardiothoracic and Vascular Anesthesia* 2009; **23**(2):175-81. [PUBMED: 19026570]

Bennett 1985 {published data only}

Bennett HL, Davis HS, Giannini JA. Non-verbal response to intraoperative conversation. *British Journal of Anaesthesia* 1985; **57**(2):174-9. [PUBMED: 3970796]

Bilotta 2007 {published data only}

Bilotta F, Caramia R, Paoloni FP, Favaro R, Araimo F, Pinto G, et al. Early postoperative cognitive recovery after remifentanyl-propofol or sufentanil-propofol anaesthesia for supratentorial craniotomy: a randomized trial. *European Journal of Anaesthesiology* 2007; Vol. 24, issue 2:122-7. [PUBMED: 16938153]

Bischoff 1997 {published data only}

Bischoff P. Intracranial pressure and anesthesia--EEG and p-EEG monitoring. *Anesthesiologie, Intensivmedizin, Notfallmedizin, Schmerztherapie* 1997; **32**(2 Suppl):122-7. [PUBMED: 9417237]

Block1991b {published data only}

Block RI, Ghoneim MM, Sum Ping ST, Ali MA. Efficacy of therapeutic suggestions for improved postoperative recovery presented during general anesthesia. *Anesthesiology* 1991; Vol. 75, issue 5:746-55. [PUBMED: 1952199]

Bogetz 1984 {published data only}

Bogetz MS, Katz JA. Recall of surgery for major trauma. *Anesthesiology* 1984; Vol. 61, issue 1:6-9. [PUBMED: 6742485]

Bogod 1990 {published data only}

Bogod DG, Orton JK, Yau HM, Oh TE. Detecting awareness during general anaesthetic caesarean section. An evaluation of two methods. *Journal of Anesthesia* 1990; Vol. 45, issue 4:279-84. [PUBMED: 2337210]

Bonhomme 2006 {published data only}

Bonhomme V, Llabres V, Dewandre PY, Brichant JF, Hans P. Combined use of Bispectral Index and A-Line Autoregressive Index to assess anti-nociceptive component of balanced anaesthesia during lumbar arthrodesis. *British Journal of Anaesthesia* 2006; Vol. 96, issue 3:353-60. [PUBMED: 16415310]

Bonke 1986 {published data only}

Bonke B, Schmitz PI, Verhage F, Zwaveling A. Clinical study of so-called unconscious perception during general anaesthesia. *British Journal of Anaesthesia* 1986; **58**(9):957-64. [PUBMED: 3756055]

Bonke 1992 {published data only}

Bonke B, Van Dam ME, Van Kleff JW, Slijper FM. Implicit memory tested in children during inhalation anaesthesia. *Journal of Anesthesia* 1992; Vol. 47, issue 9:747-9. [PUBMED: 1415970]

Bould 2007 {published data only}

Bould MD, Mahtani DG, Davies R, Roughton M, Hunter DN, Kelleher A. Bispectral index values during elective rigid bronchoscopy: a prospective observational pilot study. *Anaesthesia* 2007; **62**:438-45. [PUBMED: 17448053]

Brice 1970 {published data only}

Brice D, Hetherington R, Utting J. A simple study of awareness and dreaming during anaesthesia. *British Journal of Anaesthesia* 1970; **42**:535-42. [PUBMED: 5423844]

Brosius 2002 {published data only}

Brosius KK, Bannister CF. Oral midazolam premedication in preadolescents and adolescents. *Anesthesia and Analgesia* 2002; **94**(1):31-6, table of contents. [PUBMED: 11772796]

Bruhn 2000 {published data only}

Bruhn J, Bouillon TW, Shafer SL. Electromyographic activity falsely elevates the bispectral index. *Anesthesiology* 2000; Vol. 92, issue 5:1485-7. [10781298]

Buffett-Jerrott 2003 {published data only}

Buffett-Jerrott SE, Stewart SH, Finley GA, Loughlan HL. Effects of benzodiazepines on explicit memory in a paediatric surgery setting. *Psychopharmacology* 2003; **168**(4):377-86. [PUBMED: 12845417]

Bulach 2005 {published data only}

Bulach R, Myles PS, Russnak M. Double-blind randomized controlled trial to determine extent of amnesia with midazolam given immediately before general anaesthesia. *British Journal of Anaesthesia* 2005; **94**(3):300-5. [PUBMED: 15567810]

Burn 1963 {published data only}

Burn JM. Relaxant anesthesia for abdominal operations using hyperventilation with air. Preliminary report. *Anaesthesia* 1963; **18**:84-7. [PUBMED: 14017121]

Burrow 2001 {published data only}

Burrow B, McKenzie B, Case C. Do anaesthetized patients recover better after bispectral index monitoring?. *Journal Anaesthesia Intensive Care* 2001; Vol. 29, issue 3:239-45. [PUBMED: 11439793]

Byers 1997 {published data only}

Byers GF, Muir JG. Detecting wakefulness in anaesthetised children. *Canadian Journal of Anaesthesia* 1997;**44**(5 Pt 1):486-8. [PUBMED: 9161741]

Capitanio 1997 {published data only}

Capitanio L, Jensen EW, Filligoi GC, Makovec B, Gagliardi M, Henneberg SW, et al. On-line analysis of AEP and EEG for monitoring depth of anaesthesia. *Methods of Information in Medicine* 1997;**36**:311-4.

Chiu 2007 {published data only}

Chiu CL, Ong G, Majid AA. Impact of bispectral index monitoring on propofol administration in patients undergoing cardiopulmonary bypass. *Anaesthesia and Intensive Care* 2007;**35**(3):342-7. [PUBMED: 17591126]

Chortkoff 1995 {published data only}

Chortkoff BS, Gonsowski CT, Bennett HL, Levinson B, Crankshaw DP, Dutton RC, et al. Subanaesthetic concentrations of desflurane and propofol suppress recall of emotionally charged information. *Anesthesia and Analgesia* 1995; Vol. 81, issue 4:728-36. [PUBMED: 7574002]

Cirillo 2012 {published data only}

Cirillo V, Volpe ML, Iacono C, Crisconio P, De Robertis E, Tufano R. Is Navigator a useful tool in guiding anesthesia practice in nephropatic patients? Preliminary study. *European Journal of Anaesthesiology* 2012; Vol. 29:52.

Clark 2009 {published data only}

Clark G, Licker M, Younossian AB, Soccia PM. Titrated sedation with propofol or midazolam for flexible bronchoscopy: a randomised trial. *European Respiratory Journal* 2009;**34**(6):1277-83. [PUBMED: 19443532]

Coetzee 1998 {published data only}

Coetzee JF, van LH. Tramadol or morphine administered during operation: a study of immediate postoperative effects after abdominal hysterectomy. *British Journal of Anaesthesia* 1998;**81**(5):737-41. [PUBMED: 10193286]

Coppens 2010 {published data only}

Coppens M, Van Limmen JG, Schnider T, Wyler B, Bonte S. Study of the time course of the clinical effect of propofol compared with the time course of the predicted effect-site concentration: performance of three pharmacokinetic-dynamic models. *British Journal of Anaesthesia* 2010;**104**(4):452-8. [PUBMED: 20190259]

Cormack 1979 {published data only}

Cormack RS. Awareness during surgery--a new approach. *British Journal of Anaesthesia* 1979;**51**(11):1051-4. [PUBMED: 42425]

Crawford 1976 {published data only}

Crawford JS, James FM, 3rd, Davies P, Crawley M. A further study of general anaesthesia for Caesarean section. *British Journal of Anaesthesia* 1976;**48**(7):661-7. [PUBMED: 1016643]

Dahaba 2009 {published data only}

Dahaba AA, Bornemann H, Rehak PH, Wang G, Wu XM, Metzler H. Effect of flumazenil on bispectral index monitoring in

unpremedicated patients. *Anesthesiology* 2009;**110**(5):1036-40. [PUBMED: 19352163]

Dahaba 2010 {published data only}

Dahaba AA, Xue JX, Zhao GG, Liu QH, Xu GX, Bornemann H, et al. BIS-vista occipital montage in patients undergoing neurosurgical procedures during propofol-remifentanyl anesthesia. *Anesthesiology* 2010;**112**(3):645-51. [PUBMED: 20179501]

Davidson 2005 {published data only}

Davidson AJ, Huang GH, Czarnecki C, Gibson MA, Stewart SA, Jansen K, et al. Awareness during anesthesia in children: a prospective cohort study. *Anesthesia and Analgesia* 2005;**100**(3):653-61, table of contents. [PUBMED: 15728046]

Davies 1996 {published data only}

Davies FW, Mantzaridis H, Kenny GNC, Fisher AC. Middle latency auditory evoked potentials during repeated transitions from consciousness to unconsciousness. *Anaesthesia* 1996;**51**(25):107-13. [PUBMED: 8779362]

De Cosmo 2008 {published data only}

De Cosmo G, Congedo E, Lai C, Sgreccia M, Amato A, Beccia G, et al. Ropivacaine vs. levobupivacaine combined with sufentanil for epidural analgesia after lung surgery. *European Journal of Anaesthesiology* 2008;**25**(12):1020-5. [PUBMED: 18538053]

De Kock 2005 {published data only}

De Kock M, Lavand'homme P, Waterloos H. The short-lasting analgesia and long-term antihyperalgesic effect of intrathecal clonidine in patients undergoing colonic surgery. *Anesthesia and Analgesia* 2005;**101**(2):566-72. [PUBMED: 16037177]

Ding 1993 {published data only}

Ding Y, Fredman B, White PF. Recovery following outpatient anesthesia: use of enflurane versus propofol. *Journal of Clinical Anesthesia* 1993;**5**(6):447-50. [PUBMED: 8123267]

Ding 2007 {published data only}

Ding LA, Sun LQ, Chen SX, Qu LL, Xie DF. Modified physiological and operative score for the enumeration of mortality and morbidity risk assessment model in general surgery. *World Journal of Gastroenterology* 2007;**13**(38):5090-5. [PUBMED: 17876874]

Diz 2010 {published data only}

Diz JC, Del Rio R, Lamas A, Mendoza M, Duran M, Ferreira LM. Analysis of pharmacodynamic interaction of sevoflurane and propofol on Bispectral Index during general anaesthesia using a response surface model. *British Journal of Anaesthesia* 2010;**104**(6):733-9. [PUBMED: 20385571]

Doufas 2009 {published data only}

Doufas AG, Komatsu R, Orhan-Sungur M, Sengupta P, Wadhwa A, Mascha E, et al. Neuromuscular block differentially affects immobility and cortical activation at near-minimum alveolar concentration anesthesia. *Anesthesia and Analgesia* 2009;**109**(4):1097-104. [PUBMED: 19762737]

Downing 1976 {published data only}

Downing JW, Mahomed MC, Jeal DE, Allen PJ. Anaesthesia for caesarean section with ketamine. *Anaesthesia* 1976;**31**(7):883-92. [PUBMED: 9839]

Dressler 2007 {published data only}

Dressler I, Fritzsche T, Cortina K, Pragst F, Spies C, Rundshagen I. Psychomotor dysfunction after remifentanyl/propofol anaesthesia. *European Journal of Anaesthesiology* 2007;**24**(4):347-54. [PUBMED: 17087850]

Driscoll 2007 {published data only}

Driscoll WD, Columbia MA, Peterfreund RA. Awareness during general anesthesia: analysis of contributing causes aided by automatic data capture. *Journal of Neurosurgical Anesthesiology* 2007;**19**(4):268-72. [PUBMED: 17893580]

Eisele 1976 {published data only}

Eisele V, Weinreich A, Bartle S. Perioperative awareness and recall. *Anesthesia and Analgesia* 1976; Vol. 55, issue 4:513-8. [PUBMED: 945955]

Ekman 2004a {published data only}

Ekman A, Lindholm ML, Lennmarken C, Sandin R. Reduction in the incidence of awareness using BIS monitoring. *Acta Anaesthesiologica Scandinavica* 2004; Vol. 48, issue 1:20-6. [PUBMED: 14674969]

Ekman 2004b {published data only}

Ekman A, Brudin L, Sandin R. A comparison of bispectral index and rapidly extracted auditory evoked potentials index responses to noxious stimulation during sevoflurane anesthesia. *Anesthesia and Analgesia* 2004; Vol. 99, issue 4:1141-6, table of contents. [PUBMED: 15385365]

Eldar 1992 {published data only}

Eldar I, Lieberman N, Shiber R, Nerubay J, Perel A. Use of flumazenil for intraoperative arousal during spine fusion. *Anesthesia and Analgesia* 1992;**75**(4):580-3. [PUBMED: 1530171]

El-Kerdawy 2000 {published data only}

El-Kerdawy HM, Zalingen EE, Bovill JG. The influence of the alpha2-adrenoceptor agonist, clonidine, on the EEG and on the MAC of isoflurane. *European Journal of Anaesthesiology* 2000; Vol. 17, issue 2:105-10. [PUBMED: 10758454]

Erhan 2003 {published data only}

Erhan E, Ugur G, Gunusen I, Alper I, Ozyar B. Propofol - not thiopental or etomidate - with remifentanyl provides adequate intubating conditions in the absence of neuromuscular blockade. *Canadian Journal of Anaesthesia* 2003; Vol. 50, issue 2:108-15. [PUBMED: 12560298]

Erk 2007 {published data only}

Erk G, Ornek D, Donmez NF, Taspinar V. The use of ketamine or ketamine-midazolam for adenotonsillectomy. *International Journal of Pediatric Otorhinolaryngology* 2007; Vol. 71, issue 6:937-41. [PUBMED: 17418428]

Eroglu 2003 {published data only}

Eroglu A, Solak M, Ozen I, Aynaci O. Stress hormones during the wake-up test in scoliosis surgery. *Journal of Clinical Anesthesia* 2003; Vol. 15, issue 1:15-8. [PUBMED: 12657405]

Evans 1988 {published data only}

Evans C, Richardson PH. Improved recovery and reduced postoperative stay after therapeutic suggestions during general anaesthesia. *Lancet* 1988; Vol. 2, issue 8609:491-3. [PUBMED: 2900410]

Fahlenkamp 2010 {published data only}

Fahlenkamp AV, Krebber F, Rex S. Bispectral index monitoring during balanced xenon or sevoflurane anaesthesia in elderly patients. *European Journal of Anesthesiology* 2010;**27**(10):906-11. [PUBMED: [20671557]]

Fairley 1956 {published data only}

Fairley HB. An evaluation of local and general anaesthesia for diagnostic bronchoscopy. *Canadian Anaesthetists' Society Journal* 1956;**3**(4):366-77.

Famewo 1976 {published data only}

Famewo CE. Awareness and dreams during general anaesthesia for caesarian section a study of incidence. *Canadian Anaesthetists' Society Journal* 1976; Vol. 23, issue 6:636-9. [PUBMED: 990980]

Farang 2006 {published data only}

Farang E, Chelune GJ, Schubert A, Mascha EJ. Is depth of anesthesia, as assessed by the Bispectral Index, related to postoperative cognitive dysfunction and recovery?. *Anesthesia and Analgesia* 2006; Vol. 103, issue 3:633-40. [PUBMED: 16931673]

Filipov 2007 {published data only}

Filipov TZ, Smilov IV. Comparison of the quality of anesthesia between two techniques ascertained with bispectral index. *Anesthesiology and Intensive Care* 2007;**34**:3-13.

Fisher 2006 {published data only}

Fisher J, Hirshman E, Henthorn T, Arndt J, Passannante A. Midazolam amnesia and short-term/working memory processes. *Consciousness and Cognition* 2006; Vol. 15, issue 1:54-63. [PUBMED: 16546038]

Fitzgerald 2001 {published data only}

Fitzgerald RD, Lamm C, Oczenski W, Stimpfl T, Vycudilik W, Bauer H. Direct current auditory evoked potentials during wakefulness, anesthesia, and emergence from anesthesia. *Anesthesia and Analgesia* 2001; Vol. 92, issue 1:154-60. [PUBMED: [11133619]]

Flaishon 1997 {published data only}

Flaishon R, Windsor A, Sigl J, Sebel PS. Recovery of consciousness after thiopental or propofol. Bispectral index and isolated forearm technique. *Anesthesiology* 1997; Vol. 86, issue 3:613-9. [PUBMED: [9066327]]

Flaitz 1986 {published data only}

Flaitz CM, Nowak AJ, Hicks MJ. Evaluation of the anterograde amnesic effect of rectally administered diazepam in the sedated pedodontic patient. *ASDC Journal of Dentistry for Children* 1986; Vol. 53, issue 1:17-20. [PUBMED: [3455953]]

Flier 1986 {published data only}

Flier FJ, Smorenburg JM, van der Ent CK, Bonke B, Ruprecht J. Postanesthetic recall ability, anxiety and dreams in surgical patients. A clinical study. *Der Anaesthesist* 1986; Vol. 35, issue 10:609-12. [PUBMED: [3789390]]

Forrest 1990 {published data only}

Forrest JB, Lam L, Woo J, Rifkind A. Recovery after neuroleptanalgesia for cataract surgery. *Canadian Journal of Anaesthesia* 1990; Vol. 37, issue 4 Pt 2:S82. [PUBMED: 2193762]

Frank 2000 {published data only}

Frank T, Thieme V, Olthoff D. Clonidine within the scope of balanced inhalation anesthesia with sevoflurane--effects on pEEG parameters [Clonidin im Rahmen einer balancierten Inhalationsanasthesie mit Sevofluran--Effekte auf Parameter des pEEG.]. *Anaesthesiologie und Reanimation* 2000; **25**(2):32-6. [PUBMED: 10816895]

Fung 2008 {published data only}

Fung NY, Hu Y, Irwin MG, Chow BE, Yuen MY. Comparison between sevoflurane/remifentanyl and propofol/remifentanyl anaesthesia in providing conditions for somatosensory evoked potential monitoring during scoliosis corrective surgery. *Anaesthesia and Intensive Care* 2008; **36**(6):779-85. [PUBMED: 19115644]

Gajraj 1998 {published data only}

Gajraj RJ, Doi M, Mantzaridis H, Kenny GN. Analysis of the EEG bispectrum, auditory evoked potentials and the EEG power spectrum during repeated transitions from consciousness to unconsciousness. *British Journal of Anaesthesia* 1998; Vol. 80, issue 1:46-52. [PUBMED: [9505777]]

Gajraj 1999 {published data only}

Gajraj RJ, Doi M, Mantzaridis H, Kenny GN. Comparison of bispectral EEG analysis and auditory evoked potentials for monitoring depth of anaesthesia during propofol anaesthesia. *British Journal of Anaesthesia* 1999; Vol. 82, issue 5:672-8. [PUBMED: [10536541]]

Gan 1997 {published data only}

Gan TJ, Glass PS, Windsor A, Payne F, Rosow C, Sebel P, et al. Bispectral index monitoring allows faster emergence and improved recovery from propofol, alfentanil, and nitrous oxide anesthesia. BIS Utility Study Group. *Anesthesiology* 1997; Vol. 87, issue 4:808-15. [PUBMED: [9357882]]

Gan 1999 {published data only}

Gan TJ, Glass PS, Sigl J, Sebel P, Payne F, Rosow C, et al. Women emerge from general anesthesia with propofol/alfentanil/nitrous oxide faster than men. *Anesthesiology* May; **90**(5):1283-7. [PUBMED: 10319774]

Gazzanelli 2005 {published data only}

Gazzanelli S, Vari A, Tarquini S, Fermariello A, Caputo M, Almansour M, et al. Monitoring of consciousness with BIS during induction of anesthesia. Which muscle relaxant?. *Il Giornale di Chirurgia* 2005; **26**(4):163-9. [PUBMED: 16035253]

Ge 2003 {published data only}

Ge SJ, Zhuang XL, Wang YT, Wang ZD, Chen SL, Li HT. Performance of the rapidly extracted auditory evoked potentials index to detect the recovery and loss of wakefulness in anesthetized and paralyzed patients. *Acta Anaesthesiologica Scandinavica* 2003; Vol. 47, issue 4:466-71. [PUBMED: 12694147]

Gelman 1984 {published data only}

Gelman S, Rivas JE, Erdemir H, Oparil S, Proctor J, MacKrell T, et al. Hormonal and haemodynamic responses to upper abdominal surgery during isoflurane and balanced anaesthesia. *Canadian Anaesthetists' Society Journal* 1984; **31**(5):509-16. [MEDLINE: 6498567]

Ghabash 1996 {published data only}

Ghabash M, Matta M, Kehhaleh J. Depression of excitatory effects of propofol induction by fentanyl. *Middle East Journal of Anesthesiology* 1996; **13**(4):419-25. [PUBMED: 8713637]

Ghoneim 1988 {published data only}

Ghoneim MM, Hinrichs JV, O'Hara MW, Mehta MP, Pathak D, Kumar V, et al. Comparison of psychologic and cognitive functions after general or regional anesthesia. *Anesthesiology* 1988; Vol. 69, issue 4:507-15. [PUBMED: 3177911]

Ghoneim 1998 {published data only}

Ghoneim MM, Block RI, Dhanaraj VJ. Interaction of a subanaesthetic concentration of isoflurane with midazolam: effects on responsiveness, learning and memory. *British Journal of Anaesthesia* 1998; Vol. 80, issue 5:581-7. [PUBMED: 9691858]

Ghoneim 2007 {published data only}

Ghoneim MM, Block RI. Immediate peri-operative memory. *Acta Anaesthesiologica Scandinavica* 2007; **51**(8):1054-61. [PUBMED: 17697300]

Glass 1997 {published data only}

Glass PS, Bloom M, Kears L, Rosow C, Sebel P, Manberg P. Bispectral analysis measures sedation and memory effects of propofol, midazolam, isoflurane, and alfentanil in healthy volunteers. *Anesthesiology* 1997; Vol. 86, issue 4:836-47. [PUBMED: 9105228]

Gordon 1994 {published data only}

Gordon PC, Morrell DF, Pamm JD. Total intravenous anesthesia using propofol and alfentanil for coronary artery bypass surgery. *Journal of Cardiothoracic and Vascular Anesthesia* 1994; **8**(3):284-8. [PUBMED: 8061261]

Gregory 1969 {published data only}

Gregory GA, Eger EI, 2nd, Munson ES. The relationship between age and halothane requirement in man. *Anesthesiology* 1969; Vol. 30, issue 5:488-91. [PUBMED: 5778106]

Greif 2002 {published data only}

Greif R, Greenwald S, Schweitzer E, Laciny S, Rajek A, Caldwell JE, et al. Muscle relaxation does not alter hypnotic level during propofol anesthesia. *Anesthesia and Analgesia* 2002; Vol. 94, issue 3:604-8; table of contents. [PUBMED: 11867383]

Gross 1988 {published data only}

Gross JB, Alexander CM. Awakening concentrations of isoflurane are not affected by analgesic doses of morphine. *Anesthesia and Analgesia* 1988; Vol. 67, issue 1:27-30. [PUBMED: 3276238]

Gu 2010 {published data only}

Gu X J, Zhong T D. The effect of flurbiprofen axetil on laparoscopic cholecystectomy. *Zhonghua Yi Xue Za Zhi* 2010; **90**(11):760-2. [PUBMED: 20627022]

Guignard 2001 {published data only}

Guignard B, Coste C, Menigaux C, Chauvin M. Reduced isoflurane consumption with bispectral index monitoring. *Acta Anaesthesiologica Scandinavica* 2001; Vol. 45, issue 3:308-14. [PUBMED: 11207466]

Gunawardane 2002 {published data only}

Gunawardane PO, Murphy PA, Sleight JW. Bispectral index monitoring during electroconvulsive therapy under propofol anaesthesia. *British Journal of Anaesthesia* 2002; Vol. 88, issue 2:184-7. [PUBMED: 11878655]

Guo 2012 {published data only}

Guo Zheng-gang, Wang Xiao-yan, Lu Xu-lei, Su Xiao-jun, Hao Jian-hua. Application of Narcotrend-assisted anesthesia in-depth monitor during escharectomy and skin transplantation in burn patients with target-controlled infusion of remifentanyl hydrochloride and propofol. *Chinese Journal of Burns*. Department of Anesthesiology, the First Hospital Affiliated to the General Hospital of PLA, Beijing 100048, China., 2012:178-82.

Gupta 2006 {published data only}

Gupta A, Kaur R, Malhotra R, Kale S. Comparative evaluation of different doses of propofol preceded by fentanyl on intubating conditions and pressor response during tracheal intubation without muscle relaxants. *Paediatric Anaesthesia* 2006; Vol. 16, issue 4:399-405. [PUBMED: 16618293]

Hadzidiakos 2006 {published data only}

Hadzidiakos D, Petersen S, Baars J, Herold K, Rehberg B. Comparison of a new composite index based on midlatency auditory evoked potentials and electroencephalographic parameters with bispectral index (BIS) during moderate propofol sedation. *European Journal of Anaesthesiology* 2006; Vol. 23, issue 11:931-6. [PUBMED: 16824245]

Haessler 1993 {published data only}

Haessler R, Schwender D, Leppmeier U, Klasing S, Rindfleisch F, Peter K. Anaesthesia for coronary artery bypass grafting: opioid-analgesia combined with either flunitrazepam, propofol or isoflurane. *Acta Anaesthesiologica Scandinavica* 1993; **37**(6):532-40. [PUBMED: 8213015]

Hall 1986 {published data only}

Hall KD, Talton IH, Fox E, Klein FF, Christian C, Jobsis FF. Brain function and level of consciousness in fentanyl anesthesia in heart surgery [Gehirnfunktion und Bewusstseinslage während der Fentanylnarkose in der Herzchirurgie]. *Der Anaesthesist* 1986; **35**(4):226-30. [PUBMED: 3487258]

Hans 1998 {published data only}

Hans P, Lamy MM, Brichant JF, Cantraine F, Coussaert E, Byttebier G, et al. Emergence from target-controlled anesthesia with propofol and sufentanil in patients undergoing intracranial surgery. *Acta Anaesthesiologica Belgica* 1998; Vol. 49, issue 1:13-9. [PUBMED: 9627733]

Harris 1971 {published data only}

Harris TJ, Brice DD, Hetherington RR, Utting JE. Dreaming associated with anaesthesia: the influence of morphine premedication and two volatile adjuvants. *British Journal of Anaesthesia* 1971; Vol. 43, issue 2:172-8. [PUBMED: 5550848]

Hartridge 1963 {published data only}

Hartridge VB, Wilson RB. Balanced anesthesia for cesarean section. *American Journal of Obstetrics and Gynecology* 1963; Vol. 85:619-25. [PUBMED: 13953166]

Hartung 1986 {published data only}

Hartung, HJ, Osswald PM, Puhl R, Jacoby R. Modification of postoperative somnolence following halothane anesthesia by physostigmine. *Anesthesie, Intensivtherapie, Notfallmedizin* 1986; **21**(2):82-4. [PUBMED: 3728891]

Hashimoto 2012 {published data only}

Hashimoto A, Fujiwara Y. Relationship between bispectral index and auditory evoked potential index for propofol and midazolam during induction of general anesthesia. *British Journal of Anaesthesia*. Oxford University Press, 2012:ii116.

Hayashi 2007 {published data only}

Hayashi K, Tsuda N, Sawa T, Hagihira S. Ketamine increases the frequency of electroencephalographic bicoherence peak on the alpha spindle area induced with propofol. *British Journal of Anaesthesia* 2007; **99**(3):389-95. [PUBMED: 17621599]

Head-Rapson 1995 {published data only}

Head-Rapson AG, Devlin JC, Parker CJ, Hunter JM. Pharmacokinetics and pharmacodynamics of the three isomers of mivacurium in health, in end-stage renal failure and in patients with impaired renal function. *British Journal of Anaesthesia* 1995; Vol. 75, issue 1:31-6. [PUBMED: 7669465]

Heipertz 1986 {published data only}

Heipertz W, Vontin H, Hempel V, Eichinger V. Plasma concentration and amnesia following lormetazepam and flunitrazepam in i.v. premedication. *Anesthesie, Intensivtherapie, Notfallmedizin* 1986; Vol. 21, issue 3:113-6. [PUBMED: 3752422]

Heisterkamp 1975 {published data only}

Heisterkamp DV, Cohen PJ. The effect of intravenous premedication with lorazepam (ativan), pentobarbitone

or diazepam on recall. *British Journal of Anaesthesia* 1975;**47**(1):79-81. [PUBMED: 238548]

Heller 2005 {published data only}

Heller H, Hatami R, Mullin P, Sciacca RR, Khandji AG, Hamberger M, et al. Bilateral bispectral index monitoring during suppression of unilateral hemispheric function. *Anesthesia and Analgesia* 2005; Vol. 101, issue 1:235-41, table of contents. [PUBMED: 1597623]

Hellwagner 2003 {published data only}

Hellwagner K, Holzer A, Gustorff B, Schroegendorfer K, Greher M, Weindlmayr-Goettel M, et al. Recollection of dreams after short general anaesthesia: influence on patient anxiety and satisfaction. *European Journal of Anaesthesiology* 2003;**20**(4):282-8. [PUBMED: 12703832]

Hetem 2000 {published data only}

Hetem LA, Danion JM, Diemunsch P, Brandt C. Effect of a subanesthetic dose of ketamine on memory and conscious awareness in healthy volunteers. *Psychopharmacology* 2000; Vol. 152, issue 3:283-8. [PUBMED: 11105938]

Hirschi 2000 {published data only}

Hirschi M, Meistelman C, Longrois D. Effects of normothermic cardiopulmonary bypass on bispectral index. *European Journal of Anaesthesiology* 2000; Vol. 17, issue 8:499-505. [PUBMED: 10998033]

Honarmand 2008 {published data only}

Honarmand A, Safavi M. Prevention of propofol-induced injection pain by sufentanil: a placebo-controlled comparison with remifentanil. *Clinical Drug Investigation* 2008;**28**(1):27-35. [PUBMED: 18081358]

Hong 2008 {published data only}

Hong JY, Kang YS, Kil HK. Anaesthesia for day case excisional breast biopsy: propofol-remifentanil compared with sevoflurane-nitrous oxide. *European Journal of Anaesthesiology* 2008;**25**(6):460-7. [PUBMED: 18298873]

Horn 2009 {published data only}

Horn B, Pilge S, Kochs EF, Stockmanns G, Hock A, Schneider G. A combination of electroencephalogram and auditory evoked potentials separates different levels of anesthesia in volunteers. *Anesthesia and Analgesia* 2009;**108**(5):1512-21. [PUBMED: 19372330]

Hoymork 2003 {published data only}

Hoymork SC, Raeder J, Grimsmo B, Steen PA. Bispectral index, serum drug concentrations and emergence associated with individually adjusted target-controlled infusions of remifentanil and propofol for laparoscopic surgery. *British Journal of Anaesthesia* 2003; Vol. 91, issue 6:773-80. [PUBMED: 14633743]

Hoymork 2005 {published data only}

Hoymork SC, Raeder J. Why do women wake up faster than men from propofol anaesthesia?. *British Journal of Anaesthesia* 2005; Vol. 95, issue 5:627-33. [PUBMED: 16169889]

Huang 1988 {published data only}

Huang ST, Sum CW, Hui YL, Tan PP. Awareness of intubation with thiopental as induction agent. *Acta Anaesthesiologica Sinica* 1988; Vol. 26, issue 2:169-72. [PUBMED: 3185166]

Huang 2005 {published data only}

Huang GH, Davidson AJ, Stargatt R. Dreaming during anaesthesia in children: incidence, nature and associations. *Anaesthesia* 2005;**60**(9):854-61. [PUBMED: 16115245]

Hudetz 2007 {published data only}

Hudetz JA, Iqbal Z, Gandhi SD, Patterson KM, Hyde TF, Reddy DM, et al. Postoperative cognitive dysfunction in older patients with a history of alcohol abuse. *Anesthesiology* 2007;**106**(3):423-30. [PUBMED: 17325499]

Hughes 1994 {published data only}

Hughes JA, Sanders LD, Dunne JA, Tarpey J, Vickers MD. Reducing smoking. The effect of suggestion during general anaesthesia on postoperative smoking habits. *Anaesthesia* 1994;**49**(2):126-8. [PUBMED: 7864913]

Hughes 2009 {published data only}

Hughes TF. Modifiable midlife risk factors for late-life cognitive impairment and dementia. *Current Psychiatry Review* 2009;**5**(2):73-92. [PUBMED: 19946443]

Hutchinson 1960 {published data only}

Hutchinson R. Awareness during surgery. A study of its incidence. *British Journal of Anaesthesia* 1960;**33**:463-9. [PUBMED: 14450247]

Iannuzzi 2005 {published data only}

Iannuzzi M, Iannuzzi E, Rossi F, Berrino L, Chiefari M. Relationship between bispectral index, electroencephalographic state entropy and effect-site EC50 for propofol at different clinical endpoints. *British Journal of Anaesthesia* 2005; Vol. 94, issue 5:613-6. [PUBMED: 15734785]

Ibrahim 2001a {published data only}

Ibrahim AE, Ghoneim MM, Kharasch ED, Epstein RH, Groudine SB, Ebert TJ, et al. Speed of recovery and side-effect profile of sevoflurane sedation compared with midazolam. *Anesthesiology* 2001; Vol. 94, issue 1:87-94. [PUBMED: 11135727]

Ibrahim 2001b {published data only}

Ibrahim AE, Taraday JK, Kharasch ED. Bispectral index monitoring during sedation with sevoflurane, midazolam, and propofol. *Anesthesiology* 2001; Vol. 95, issue 5:1151-9. [PUBMED: 11684984]

Inagaki 1997 {published data only}

Inagaki Y, Shindo H, Mashimo T, Yoshiya I. The effects of epidural fentanyl on hemodynamic responses during emergence from isoflurane anesthesia and tracheal extubation: a comparison with intravenous fentanyl. *Anesthesia and Analgesia* 1997;**85**(2):328-35. [PUBMED: 9249109]

Ingelmo 2007 {published data only}

Ingelmo PM, Bendall EJ, Frawley G, Locatelli BG, Milan B, Lodetti D, et al. Bupivacaine caudal epidural anesthesia: Assessing the effect of general anesthetic technique on block onset. *Paediatric Anaesthesia* 2007;**17**(3):255-62. [PUBMED: 17263741]

Inglis 1993 {published data only}

Inglis S, Farnill D. The effects of providing preoperative statistical anaesthetic-risk information. *Journal Anaesthesia Intensive Care* 1993; Vol. 21, issue 6:799-805. [PUBMED: 8122737]

Iqbal 1985 {published data only}

Iqbal KM, Hussain P, Afzalunnessa B. Study of awareness during noninhalational anaesthesia. *Bangladesh Medical Research Council Bulletin* 1985; Vol. 11, issue 1:8-14. [PUBMED: 4026761]

Iselin-Chaves 1998 {published data only}

Iselin-Chaves IA, Flaishon R, Sebel PS, Howell S, Gan TJ, Sigl J, et al. The effect of the interaction of propofol and alfentanil on recall, loss of consciousness, and the Bispectral Index. *Anesthesia and Analgesia* 1998; Vol. 87, issue 4:949-55. [PUBMED: 9768800]

Iselin-Chaves 2000 {published data only}

Iselin-Chaves IA, El Moalem HE, Gan TJ, Ginsberg B, Glass PS. Changes in the auditory evoked potentials and the bispectral index following propofol or propofol and alfentanil. *Anesthesiology* 2000; Vol. 92, issue 5:1300-10. [PUBMED: 10781275]

Iselin-Chaves 2005 {published data only}

Iselin-Chaves IA, Willems SJ, Jermann FC, Forster A, Adam SR, Van der Linden M. Investigation of implicit memory during isoflurane anesthesia for elective surgery using the process dissociation procedure. *Anesthesiology* 2005; Vol. 103, issue 5:925-33. [PUBMED: 16249665]

Ishiyama 2005 {published data only}

Ishiyama T, Kashimoto S, Oguchi T, Yamaguchi T, Okuyama K, Kumazawa T. Epidural ropivacaine anesthesia decreases the bispectral index during the awake phase and sevoflurane general anesthesia. *Anesthesiology* 2005;**100**(3):728-32. [PUBMED: 15728060]

Isomura 2008 {published data only}

Isomura H, Hashiguchi K. Combination of dexmedetomidine and midazolam is useful for postoperative sedation in healthy young patients. *Nippon Shika Masui Gakkai Zasshi [The Journal of the Japanese Dental Society of Anesthesiology]* 2008;**36**(1):28-32.

Ittichaikulthol 2007 {published data only}

Ittichaikulthol W, Sriswasdi S, Prachanpanich N, Watcharotayangul J, Seangrungs R, Sithamwilai W. Bispectral index in assessment of 3% and 4.5% desflurane in 50% N2O for caesarean section. *Journal of the Medical Association of Thailand* 2007;**90**(8):1546-50.

Ivanov 1969 {published data only}

Ivanov SD, Nunn JF. Methods of evaluation of PCO₂ for restoration of spontaneous breathing after artificial ventilation of anaesthetized patients. *British Journal of Anaesthesia* 1969; Vol. 41, issue 1:28-37. [PUBMED: 5774302]

Jacoby 1981 {published data only}

Jacoby LL, Dallas M. On the relationship between autobiographical memory and perceptual learning. *Journal of Experimental Psychology. General* 1981; Vol. 110, issue 3:306-40. [PUBMED: 6457080]

Jaffrelot 2007 {published data only}

Jaffrelot M, Jendrin J, Floch Y, Lockey D, Jabre P, Vergne M, et al. Prevention of awakening signs after rapid-sequence intubation: a randomized study. *American Journal of Emergency Medicine* 2007;**25**(5):529-34. [PUBMED: 17543656]

Jelezov 2007 {published data only}

Jelezov C, Schmidt J, Schmitz B, Becke K, Albrecht S. EEG variables as measures of arousal during propofol anaesthesia for general surgery in children: rational selection and age dependence. *British Journal of Anaesthesia* 2007;**99**(6):845-54. [PUBMED: 17965418]

Jelicic 1993a {published data only}

Jelicic M, Bonke B, Millar K. Effect of different therapeutic suggestions presented during anaesthesia on post-operative course. *European Journal of Anaesthesia* 1993;**10**(10):343-7. [PUBMED: 11767423]

Jelicic 1993b {published data only}

Jelicic M, Asbury AJ, Millar K, Bonke B. Implicit learning during enflurane anaesthesia in spontaneously breathing patients?. *Anaesthesia* 1993;**48**(9):766-8. [PUBMED: 8214493]

Jellish 2000 {published data only}

Jellish WS, Leonetti JP, Avramov A, Fluder E, Murdoch J. Remifentanyl-based anesthesia versus a propofol technique for otologic surgical procedures. *Otolaryngology and Head and Neck Surgery* 2000; Vol. 122, issue 2:222-7. [PUBMED: 10652394]

Jellish 2009 {published data only}

Jellish WS, Leonetti JP, Buoy CM, Sincacore JM, Sawicki KJ, Macken MP. Facial nerve electromyographic monitoring to predict movement in patients titrated to a standard anesthetic depth. *Anesthesia and Analgesia* 2009;**109**(2):551-8. [PUBMED: 19608831]

Jensen 1996 {published data only}

Jensen EW, Lindholm P, Henneberg SW. Autoregressive modeling with exogenous input of middle-latency auditory-evoked potentials to measure rapid changes in depth of anesthesia. *Methods of Information in Medicine* 1996;**35**(3):256-60. [PUBMED: 8952311]

Jeon 2000 {published data only}

Jeon SY, Lim HJ. Awareness detection during a cesarean section under general anesthesia using bispectral index monitoring. *Daehan Machwigwa Haghoeji [Society of Anesthesiologists]* 2000;**39**:632-7.

Jessop 1991 {published data only}

Jessop J, Griffiths DE, Furness P, Jones JG, Sapsford DJ, Breckon DA. Changes in amplitude and latency of the P300 component of the auditory evoked potential with sedative and anaesthetic concentrations of nitrous oxide. *British Journal of Anaesthesia* 1991; Vol. 67, issue 5:524-31. [PUBMED: 1751264]

Jones 1990 {published data only}

Jones MJ, Piggott SE, Vaughan RS, Bayer AJ, Newcombe RG, Twining TC, et al. Cognitive and functional competence after anaesthesia in patients aged over 60: controlled trial of general and regional anaesthesia for elective hip or knee replacement. *BMJ* 1990; Vol. 300, issue 6741:1683-7. [PUBMED: 2390547]

Jordan 2012 {published data only}

Jordan D, Schneider G, Omerovic A, Kreuzer M, Berger S, Kochs E. Reliability to detect wakefulness and burst suppression: a new anaesthesia multimodal indicator and bis. *Journal of Neurosurgical Anesthesiology* 2012:480-1.

Jospin 2007 {published data only}

Jospin M, Caminal P, Jensen EW, Litvan H, Vallverdu M, Struys MMRF, et al. Detrended fluctuation analysis of EEG as a measure of depth of anaesthesia. *IEEE Transactions on Bio-medical Engineering* 2007;**54**(5):840-6. [PUBMED: 17518280]

Juckenhofel 1999 {published data only}

Juckenhofel S, Feisel C, Schmitt HJ, Biedler A. TIVA with propofol-remifentanyl or balanced anaesthesia with sevoflurane-fentanyl in laparoscopic operations. Hemodynamics, awakening and adverse effects. *Anaesthesist* 1999;**48**(11):807-12. [PUBMED: 10631440]

Katoh 1993 {published data only}

Katoh T, Suguro Y, Kimura T, Ikeda K. Morphine does not affect the awakening concentration of sevoflurane. *Canadian Journal Anaesthesia* 1993;**40**(9):825-8. [PUBMED: 8403176]

Katoh 1994 {published data only}

Katoh T, Uchiyama T, Ikeda K. Effect of fentanyl on awakening concentration of sevoflurane. *British Journal of Anaesthesia* 1994;**73**(3):322-5. [PUBMED: 7946857]

Kavey 1979 {published data only}

Kavey NB, Ahshuler KZ. Sleep in herniorrhaphy patients. *American Journal of Surgery* 1979; Vol. 138, issue 5:683-7. [PUBMED: 227283]

Kennedy 1985 {published data only}

Kennedy DJ, Ogg TW. Alfentanil and memory function. A comparison with fentanyl for day case termination of pregnancy. *Anaesthesia* 1985;**40**(6):537-40. [PUBMED: 3927763]

Kenny 1999 {published data only}

Kenny GN, Mantzaridis H. Closed-loop control of propofol anaesthesia. *British Journal of Anaesthesia* 1999; Vol. 83, issue 2:223-8. [PUBMED: 10618933]

Kerssens 2002 {published data only}

Kerssens C, Lubke GH, Klein J, van der Woerd A, Bonke B. Memory function during propofol and alfentanil anaesthesia:

predictive value of individual differences. *Anesthesiology* 2002; Vol. 97, issue 2:382-9. [PUBMED: 12151928]

Kerssens 2003 {published data only}

Kerssens C, Klein J, Bonke B. Awareness: monitoring versus remembering what happened. *Anesthesiology* 2003; Vol. 99, issue 3:570-5. [PUBMED: 12960540]

Kerssens 2005a {published data only}

Kerssens C, Hamann S, Peltier S, Hu XP, Byas-Smith MG, Sebel PS. Attenuated brain response to auditory word stimulation with sevoflurane: a functional magnetic resonance imaging study in humans. *Anesthesiology* 2005; Vol. 103, issue 1:11-9. [PUBMED: 15983451]

Kertai 2011 {published data only}

Kertai MD, Palanca BJ, Pal N, Burnside BA, Zhang L, Sadiq F, et al. Bispectral index monitoring, duration of bispectral index below 45, patient risk factors, and intermediate-term mortality after noncardiac surgery in the B-Unaware Trial. *Anesthesiology* 2011;**114**(3):545-56. [PUBMED: 21293252]

Kestin 1990 {published data only}

Kestin IG, Harvey PB, Nixon C. Psychomotor recovery after three methods of sedation during spinal anaesthesia. *British Journal of Anaesthesia* 1990; Vol. 64, issue 6:675-81. [PUBMED: 2116156]

Kevin 2002 {published data only}

Kevin LG, Cunningham AJ, Bolger C. Comparison of ocular microtremor and bispectral index during sevoflurane anaesthesia. *British Journal of Anaesthesia* 2002; Vol. 89, issue 4:551-5. [PUBMED: (Print)12393354]

Khandwala 2008 {published data only}

Khandwala M, Ahmed S, Goel S, Simmons IG, McLure HA. The effect of hyaluronidase on ultrasound-measured dispersal of local anaesthetic following sub-Tenon injection. *Eye (London, England)* 2008;**22**(8):1065-8. [PUBMED: 17525774]

Kiernan 1995 {published data only}

Kiernan BD, Dane JR, Phillips LH, Price DD. Hypnotic analgesia reduces R-III nociceptive reflex: further evidence concerning the multifactorial nature of hypnotic analgesia. *Pain* 1995; Vol. 60, issue 1:39-47. [PUBMED: 7715940]

Kim 1978 {published data only}

Kim CL. Awareness during cardiopulmonary bypass. *American Association of Nurse Anesthetists Journal* 1978; Vol. 46, issue 4:373-83. [PUBMED: 362790]

Kim 2010 {published data only}

Kim HT, Heo HE, Kwon YE, Lee MJ. Effect of remifentanyl on consumption of sevoflurane in entropy monitored general anaesthesia. *Korean Journal of Anesthesiology* 2010;**59**(4):238-43. [PUBMED: 21057612]

Kliempt 1999 {published data only}

Kliempt P, Ruta D, Ogston S, Landeck A, Martay K. Hemispheric-synchronisation during anaesthesia: a double-blind randomised trial using audiotapes for intra-operative

nociception control. *Anaesthesia* 1999; Vol. 54, issue 8:769-73. [PUBMED: 10460529]

Kocaman 2007 {published data only}

Kocaman Akbay B, Demiraran Y, Sezen GY, Akcali G, Somunkiran A. Use of the bispectral index to predict a positive awareness reaction to laryngeal mask airway-Fastrach insertion and intubation. *Acta Anaesthesiologica Scandinavica* 2007;**51**(10):1368-72. [PUBMED: 17944640]

Kokki 2007 {published data only}

Kokki, H, Wennervirta J, Laisalmi M, Vakkuri A. Regular tramadol use does not effect the propofol dose requirement for induction of anaesthesia. *European Journal of Anaesthesiology* 2007;**24**(9):776-81. [PUBMED: 17462119]

Kotiniemi 1996 {published data only}

Kotiniemi LH, Ryhanen PT, Moilanen IK. Behavioural changes and children's memories after intravenous, inhalation and rectal induction of anaesthesia. *Paediatric Anaesthesia* 1996;**6**(3):201-7. [PUBMED: 8732611]

Kreuer 2001 {published data only}

Kreuer, S, Biedler A, Larsen R, Scotho S, Altmann S, Wilhelm W. The Narcotrend--a new EEG monitor designed to measure the depth of anaesthesia. A comparison with bispectral index monitoring during propofol-remifentanil-anaesthesia. *Der Anaesthesist* 2001;**50**(12):921-5. [PUBMED: 11824075]

Lapidus 2007 {published data only}

Lapidus LJ, Rosfors S, Ponzer S, Levander C, Elvin A, Larfars G, et al. Prolonged thromboprophylaxis with dalteparin after surgical treatment of achilles tendon rupture: a randomized, placebo-controlled study. *Journal of Orthopaedic Trauma* 2007;**21**(1):52-7. [PUBMED: 17211270]

Latto 1977 {published data only}

Latto IP, Waldron BA. Anaesthesia for Caesarean section. Analysis of blood concentrations of halothane using 0.2% or 0.65% halothane with 50% nitrous oxide in oxygen. *British Journal of Anaesthesia* 1977;**49**(4):371-8. [PUBMED: 889651]

Lefoll-Masson 2007 {published data only}

Lefoll-Masson C, Fermanian C, Aime I, Verroust N, Taylor G, Laloe PA, et al. The comparability of bispectral index and state entropy index during maintenance of sufentanil-sevoflurane-nitrous oxide anesthesia. *Anesthesia and Analgesia* 2007;**105**(5):1319-25, table of contents. [PUBMED: 17959961]

Lehmann 2000 {published data only}

Lehmann A, Zeitler C, Thaler E, Isgro F, Boldt J. Comparison of two different anesthesia regimens in patients undergoing aortocoronary bypass grafting surgery: sufentanil-midazolam versus remifentanil-propofol. Retractions. [J Cardiothorac Vasc Anesth. 2011]. *Journal of Cardiothoracic and Vascular Anesthesia* 2000;**14**(4):416-20. [PUBMED: 10972607]

Lehmann 2002 {published data only}

Lehmann A, Boldt J, Thaler E, Piper S, Weisse U. Bispectral index in patients with target-controlled or manually-controlled infusion of propofol. Retraction in Bispectral index in patients

with target-controlled or manually-controlled infusion of propofol: Retraction. [Anesth Analg. 2011]. *Anesthesia and Analgesia* 2002;**95**(3):638-44. [PUBMED: 12198052]

Lehmann 2003 {published data only}

Lehmann A, Karzau J, Boldt J, Thaler E, Lang J, Isgro F. Bispectral index-guided anesthesia in patients undergoing aortocoronary bypass grafting. Retraction in Bispectral index-guided anesthesia in patients undergoing aortocoronary bypass grafting: Retraction. [Anesth Analg. 2011]. *Anaesthesia and Analgesia* 2003;**96**:336-43. [PUBMED: 12538174]

Lequeux 2003 {published data only}

Lequeux PY, Cantraine F, Levarlet M, Barvais L. Absence of explicit and implicit memory in unconscious patients using a TCI of propofol. *Acta Anaesthesiologica Scandinavica* 2003;**47**(7):833-7. [PUBMED: 12859304]

Leslie 2007 {published data only}

Leslie K, Skrzypek H, Paech MJ, Kurowski I, Whybrow T. Dreaming during anesthesia and anesthetic depth in elective surgery patients: a prospective cohort study. *Anesthesiology* 2007;**106**(1):33-42. [PUBMED: 17197843]

Levine 1993 {published data only}

Levine MF, Hartley EJ, Macpherson BA, Burrows FA, Lerman J. Oral midazolam premedication for children with congenital cyanotic heart disease undergoing cardiac surgery: a comparative study. *Canadian Journal of Anaesthesia* 1993;**40**(10):934-8. [PUBMED: 8222032]

Liao 2010 {published data only}

Liao R, Li JY, Liu GY. Comparison of sevoflurane volatile induction/maintenance anaesthesia and propofol-remifentanil total intravenous anaesthesia for rigid bronchoscopy under spontaneous breathing for tracheal/bronchial foreign body removal in children. *European Journal of Anaesthesiology* 2010;**27**(11):930-4. [PUBMED: 20683333]

Liou 1994 {published data only}

Liou CM, Kang HM, Lai HC, Liu YC, Hung CJ, Wang SJ, et al. Will epidural with light general anesthesia increase the incidence of awareness with recall or dream postoperatively. *Acta Anaesthesiologica Sinica* 1994;**32**(4):229-36. [PUBMED: 7894918]

Liu 2005 {published data only}

Liu N, Chazot T, Huybrechts I, Law-Koune JD, Barvais L, Fischler M. The influence of a muscle relaxant bolus on bispectral and datex-ohmeda entropy values during propofol-remifentanil induced loss of consciousness. *Anesthesia and Analgesia* 2005;**101**(6):1713-8. [PUBMED: 16301247]

Liu 2006 {published data only}

Liu N, Chazot T, Genty A, Landais A, Restoux A, McGee K, et al. Titration of propofol for anesthetic induction and maintenance guided by the bispectral index: closed-loop versus manual control: a prospective, randomized, multicenter study. *Anesthesiology* 2006;**104**:686-95. [PUBMED: 16571963]

Liu 2009 {published data only}

Liu SH, Wei W, Ding GN, Ke JD, Hong FX, Tian M. Relationship between depth of anesthesia and effect-site concentration of propofol during induction with the target-controlled infusion technique in elderly patients. *Chinese Medical Journal* 2009;**122**(8):935-40. [PUBMED: 19493418]

Lopez 2007 {published data only}

Lopez U, Habre W, Laurencon M, Haller G, Van der Linden M, Iselin-Chaves IA. Intra-operative awareness in children: the value of an interview adapted to their cognitive abilities. *Anaesthesia* 2007;**62**(8):778-89. [PUBMED: 17635425]

Low 2007 {published data only}

Low D, Ng I, Ng WH. Awake craniotomy under local anaesthesia and monitored conscious sedation for resection of brain tumours in eloquent cortex--outcomes in 20 patients. *Annals of the Academy of Medicine, Singapore* 2007;**36**(5):326-31. [PUBMED: 17549278]

Luginbuhl 2007 {published data only}

Luginbuhl M, Ypparila Wolters H, Rufenacht M, Petersen Felix S, Korhonen I. Heart rate variability does not discriminate between different levels of haemodynamic responsiveness during surgical anaesthesia. *British Journal of Anaesthesia* 2007;**98**(6):728-36. [PUBMED: 17468493]

Luginbuhl 2010 {published data only}

Luginbühl M, Schumacher PM, Vuilleumier P, Vereecke H, Heyse B, Bouillon TW, et al. Noxious stimulation response index: a novel anesthetic state index based on hypnotic-opioid interaction. *Anesthesiology* 2010;**112**(3):872-80. [PUBMED: 20216387]

Lyons 1991 {published data only}

Lyons G, Macdonald R. Awareness during caesarean section. *Anaesthesia* 1991;**46**(1):62-4. [PUBMED: 1996761]

Magni 2009 {published data only}

Magni G, Rosa IL, Melillo G, Savio A, Rosa G. A comparison between sevoflurane and desflurane anesthesia in patients undergoing craniotomy for supratentorial intracranial surgery. *Anesthesia and Analgesia* 2009;**109**(2):567-71. [PUBMED: 19608833]

Mahomedy 1976 {published data only}

Mahomedy MC, Downing JW, Jeal DE, Allen PJ. Ketamine for anaesthetic induction at Caesarean section. *South African Medical Journal* 1976;**50**(22):846-8. [PUBMED: 935971]

Malek 2009 {published data only}

Malek J, Hess L, Votava M, Marecek F, Kurzova A. The combination of dexmedetomidine, fentanyl, midazolam and atropine may result in dangerous respiratory depression and hypoxemia. *European Journal of Anaesthesiology* 2009;**26**(Suppl 45):136.

Malek 2010 {published data only}

Malek B, Seghlis R, Dabbagh A. Awareness during anesthesia in elective abdominal surgery in opium abusers. *Tehran University Medical Journal* 2010;**67**(10):731-5.

Malviya 2009 {published data only}

Malviya S, Galinkin JL, Bannister CF, Burke C, Zuk J, Popenhagen M, et al. The incidence of intraoperative awareness in children: childhood awareness and recall evaluation. *Anesthesia and Analgesia* 2009;**109**(5):1421-7. [PUBMED: 19713260]

Martorano 2008 {published data only}

Martorano PP, Aloj F, Baietta S, Fiorelli A, Munari M, Paccagnella F, et al. Sufentanil-propofol vs remifentanil-propofol during total intravenous anesthesia for neurosurgery. A multicentre study. *Minerva Anestesiologica* 2008;**74**(6):233-43.

Mathews 2008 {published data only}

Mathews DM, Gaba V, Zaku B, Neuman GG. Can remifentanil replace nitrous oxide during anesthesia for ambulatory orthopedic surgery with desflurane and fentanyl?. *Anesthesia and Analgesia* 2008;**106**(1):101-8, table of contents. [PUBMED: 18165562]

Maybauer 2007 {published data only}

Maybauer DM, Geldner G, Blobner M, Puhringer F, Hofmockel R, Rex C, et al. Incidence and duration of residual paralysis at the end of surgery after multiple administrations of cisatracurium and rocuronium. *Anaesthesia* 2007;**62**(1):12-7. [PUBMED: 17156221]

Mayer 2007 {published data only}

Mayer J, Boldt J, Schellhaass A, Hiller B, Suttner SW. Bispectral index-guided general anesthesia in combination with thoracic epidural analgesia reduces recovery time in fast-track colon surgery. Retraction in Bispectral index-guided general anesthesia in combination with thoracic epidural analgesia reduces recovery time in fast-track colon surgery: Retraction. [Anesth Analg. 2011]. *Anesthesia and Analgesia* 2007;**104**(5):1145-9, tables of contents. [PUBMED: 17456665]

Mellema 2010 {published data only}

Mellema A, Borner M, Joppich R, Burmester S, Schutte JK, Wappler F. Comparison of inhalational versus intravenous anaesthetic techniques for elective spinal surgery under comparable anaesthetic depth using BIS monitoring. A randomized, prospective, clinical trial. *Perioperative Medizin* 2010;**2**(2):86-93.

Messahel 2003 {published data only}

Messahel FM, Al-Qahtani AS. Awareness during surgery. *Saudi Medical Journal* 2003;**24**(9):967-70. [PUBMED: 12973479]

Messahel 2007 {published data only}

Messahel FM, Gregorio MJ. Intraoperative awareness--a three year prospective study using bispectral index monitor (BIS). *Middle East Journal of Anesthesiology* 2007;**19**(3):587-94. [PUBMED: 18044286]

Mi 1998 {published data only}

Mi WD, Sakai T, Takahashi S, Matsuki A. Haemodynamic and electroencephalograph responses to intubation during induction with propofol or propofol/fentanyl. *Canadian Journal of Anaesthesia* 1998; Vol. 45:19-22. [PUBMED: 9466021]

Mi 1999 {published data only}

Mi WD, Sakai T, Singh H, Kudo T, Kudo M, Matsuki A. Hypnotic endpoints vs. the bispectral index, 95% spectral edge frequency and median frequency during propofol infusion with or without fentanyl. *European Journal of Anaesthesiology* 1999;**16**(1):47-52. [PUBMED: 10084101]

Mirakhur 1986 {published data only}

Mirakhur RK, Lavery GG, Gibson FM, Clarke RS. Intubating conditions after vecuronium and atracurium given in divided doses (the priming technique). *Acta Anaesthesiologica Scandinavica* 1986;**30**(5):347-50. [PUBMED: 2876574]

Moerman 1995 {published data only}

Moerman A, Herregods L, Foubert L, Poelaert J, Jordaens L, D'Hont L, et al. Awareness during anaesthesia for implantable cardioverter defibrillator implantation. Recall of defibrillation shocks. *Anaesthesia* 1995; Vol. 50, issue 8:733-5. [PUBMED: 7645710]

Motsch 1996 {published data only}

Motsch J, Wandel C, Neff S, Martin E. A comparative study of the use of sevoflurane and propofol in ambulatory surgery. *Der Anaesthesist* 1996;**45**(Suppl 1):57-62. [PUBMED: 8775105]

Mourisse 2007 {published data only}

Mourisse J, Lerou J, Struys M, Zwarts M, Boon L. Multi-level approach to anaesthetic effects produced by sevoflurane or propofol in humans: 1. BIS and blink reflex. *British Journal of Anaesthesia* 2007;**98**(6):737-45. [PUBMED: 17519261]

Moustafa 2008 {published data only}

Moustafa AM, Negmi HH, Rabie ME. The combined effect of ketamine and remifentanyl infusions as total intravenous anaesthesia for scoliosis surgery in children. *Middle East Journal of Anaesthesiology* 2008;**19**(5):1151-68. [PUBMED: 18637615]

Nakagawa 2001 {published data only}

Nakagawa M, Mammoto T, Sakai T, Kishi Y, Mashimo T. Premedication modifies the quality of sedation with propofol during regional anaesthesia. *Canadian Journal of Anaesthesia* 2001;**48**(3):284-7. [PUBMED: 11305831]

Nelskyla 2001 {published data only}

Nelskyla KA, Yli-Hankala AM, Puro PH, Korttila KT. Sevoflurane titration using bispectral index decreases postoperative vomiting in phase II recovery after ambulatory surgery. *Anesthesia and Analgesia* 2001;**93**(5):1165-9. [PUBMED: 11682388]

Ng 1974 {published data only}

Ng KH, Gurubatham AI. Awareness during caesarean section under general anaesthesia. *Medical Journal of Australia* 1974;**2**(21):774-6. [PUBMED: 4453236]

Nishijima 1999 {published data only}

Nishijima S, Konno M, Sakurada H. The effects of propofol anaesthesia with or without the use of nitrous oxide on the intraoperative involuntary movement, the postoperative awareness and vomiting. *Masui* 1999;**48**(11):1216-9. [PUBMED: 10586554]

O'Sullivan 1988 {published data only}

O'Sullivan EP, Childs D, Bush GH. Peri-operative dreaming in paediatric patients who receive suxamethonium. *Anaesthesia* 1988;**43**(2):104-6. [PUBMED: 3190825]

Oikkonen 1994 {published data only}

Oikkonen M. Propofol vs isoflurane for gynaecological laparoscopy. *Acta Anaesthesiologica Scandinavica* 1994;**38**(2):110-4. [PUBMED: 8171944]

Onaka 1998 {published data only}

Onaka M, Yamamoto H, Akatsuka M, Mori H. Anaesthetic management by continuous total intravenous anaesthesia. *Masui* 1998;**47**(10):1200-6. [PUBMED: 9834591]

Oudenaarden 1979 {published data only}

Oudenaarden JJ. Etomidate as an induction agent in minor operative procedures. *Current Medical Research and Opinion* 1979;**6**(1):30-4. [PUBMED: 378555]

Padmanabhan 2009 {published data only}

Padmanabhan U, Leslie K, Eer AS, Maruff P, Silbert BS. Early cognitive impairment after sedation for colonoscopy: the effect of adding midazolam and/or fentanyl to propofol. *Anesthesia and Analgesia* 2009;**109**(5):1448-55. [PUBMED: 19617584]

Panousis 2007 {published data only}

Panousis P, Heller AR, Burghardt M, Bleyl JU, Koch T. The effects of electromyographic activity on the accuracy of the Narcotrend monitor compared with the Bispectral Index during combined anaesthesia. *Anaesthesia* 2007;**62**(9):868-74. [PUBMED: 17697211]

Philbin 1990 {published data only}

Philbin DM, Rosow CE, Schneider RC, Koski G, D'Ambra MN. Fentanyl and sufentanil anaesthesia revisited: how much is enough?. *Anesthesiology* 1990;**73**(1):5-11. [PUBMED: 2141773]

Philipp 2002 {published data only}

Philipp A, Wiesenack C, Behr R, Schmid FX, Birnbaum DE. High risk of intraoperative awareness during cardiopulmonary bypass with isoflurane administration via diffusion membrane oxygenators. *Perfusion* 2002;**17**(3):175-8.

Piano 2007 {published data only}

Piano G, Ketteler ER, Prachand V, DeValk E, Van Ha TG, Gewertz BL, et al. Safety, feasibility, and outcome of retrievable vena cava filters in high-risk surgical patients. *Journal of Vascular Surgery* 2007;**45**(4):784-8. [PUBMED: 17398387]

Pomfrett 2009 {published data only}

Pomfrett CJ, Dolling S, Anders NR. Delta sleep-inducing peptide alters bispectral index, the electroencephalogram and heart rate variability when used as an adjunct to isoflurane anaesthesia. *European Journal of Anaesthesiology* 2009;**26**(2):128-34. [PUBMED: 19142086]

Pompeo 2007 {published data only}

Pompeo E, Tacconi F, Mineo D, Mineo TC. The role of awake video-assisted thoracoscopic surgery in spontaneous

pneumothorax. *Journal of Thoracic and Cardiovascular Surgery* 2007;**133**(3):786-90. [PUBMED: 17320585]

Porter 2008 {published data only}

Porter R, Booth D, Gray H, Frampton C. Effects of the addition of remifentanyl to propofol anesthesia on seizure length and postictal suppression index in electroconvulsive therapy. *Journal of ECT* 2008;**24**(3):203-7. [PUBMED: 18772705]

Pryor 2010 {published data only}

Pryor KO, Reinsel RA, Mehta M, Li Y, Wixted JT, Veselis RA. Visual P2-N2 complex and arousal at the time of encoding predict the time domain characteristics of amnesia for multiple intravenous anesthetic drugs in humans. *Anesthesiology* 2010;**113**(2):313-26. [PUBMED: 20613477]

Qi 2014 {published data only}

Qi S, Wu X Q, Lu X B. Application of transcutaneous electrical acupoint stimulation combined with infusion of propofol in anodynia bronchoscopy. *Zhongguo Zhen Jiu [Chinese Acupuncture & Moxibustion]* 2014;**NA**:483-6.

Rabiee 2012 {published data only}

Rabiee SM, Alijanpour E, Naziri F, Alreza H, Esmaeili V. A comparison of depth of anesthesia and hemodynamic variables with sodium thiopental and propofol as induction agents for cesarean section. *Journal of Babol University of Medical Sciences* 2012:7-12.

Reinhart 1985 {published data only}

Reinhart K, Dallinger-Stiller G, Dennhardt R, Heinemeyer G, Eyrich K. Comparison of midazolam, diazepam and placebo i.m. as premedication for regional anaesthesia. A randomized double-blind study. *British Journal of Anaesthesia* 1985;**57**(3):294-9. [PUBMED: 3156614]

Russell 1997 {published data only}

Russell IF, Wang M. Absence of memory for intraoperative information during surgery under adequate general anaesthesia. *British Journal of Anaesthesia* 1997;**78**(1):3-9. [PUBMED: 9059196]

Russell 2001 {published data only}

Russell IF, Wang M. Absence of memory for intra-operative information during surgery with total intravenous anaesthesia. *British Journal of Anaesthesia* 2001;**86**(2):196-202. [PUBMED: 11573659]

Samuelsson 2007 {published data only}

Samuelsson P, BruDin L, Sandin RH. Late psychological symptoms after awareness among consecutively included surgical patients. *Anesthesiology* 2007;**106**(1):26-32. [17197842]

Schneider 2005 {published data only}

Schneider G, Hollweck R, Ningler M, Stockmanns G, Kochs EF. Detection of consciousness by electroencephalogram and auditory evoked potentials. *Anesthesiology* 2005;**103**(5):934-43. [PUBMED: 16249666]

Schraag 1998 {published data only}

Schraag S, Mohl U, Bothner U, Georgieff M. Clinical utility of EEG parameters to predict loss of consciousness and response to skin incision during total intravenous anaesthesia. *Anaesthesia* 1998;**53**(4):320-5. [PUBMED: 9613295]

Schroeck 2010 {published data only}

Schroeck H, Fecho K, Abode K, Bailey A. Vocal cord function and bispectral index in pediatric bronchoscopy patients emerging from propofol anesthesia. *Pediatric Pulmonology* 2010;**45**(5):494-9. [PUBMED: 20425858]

Schulz 2007 {published data only}

Schulz U, Keh D, Barner C, Kaisers U, Boemke W. Bispectral index monitoring does not improve anesthesia performance in patients with movement disorders undergoing deep brain stimulating electrode implantation. *Anesthesia and Analgesia* 2007;**104**(6):1481-7, table of contents. [PUBMED: 17513646]

Schwender 1991a {published data only}

Schwender D, Keller I, Schlund M, Klasing S, Madler C. Acoustic evoked potentials of medium latency and intraoperative wakefulness during anesthesia maintenance using propofol, isoflurane and flunitrazepam/fentanyl. *Anaesthesist* 1991;**40**(4):214-21. [PUBMED: 2058823]

Schwender 1991b {published data only}

Schwender D, Keller I, Daschner B, Madler C. Sensory information processing during general anesthesia-- acoustic-evoked 30-40 Hz oscillations and intraoperative wakefulness during cesarean section [Sensorische Informationsverarbeitung während Allgemeinanästhesie--Akustisch evozierte 30-40 Hz-Oszillation und intraoperative Aufwachreaktion während Sectio caesarea]. *Anesthesiologie, Intensivmedizin, Notfallmedizin, Schmerztherapie: AINS* 1991;**26**(1):17-24. [PUBMED: 1863679]

Schwender 1993 {published data only}

Schwender D, Muller A, Madler M, Faber-Zullig E, Ilmberger J. Recovery of psychomotor and cognitive functions following anesthesia. Propofol/alfentanil and thiopental/isoflurane/alfentanil. *Anaesthesist* 1993;**42**(9):583-91. [PUBMED: 8214530]

Schwender 1994 {published data only}

Schwender D, Faber-Zullig E, Klasing S, Poppel E, Peter K. Motor signs of wakefulness during general anaesthesia with propofol, isoflurane and flunitrazepam/fentanyl and midlatency auditory evoked potentials. *Anaesthesia* 1994;**49**(6):476-84. [PUBMED: 8017589]

Schwender 1997 {published data only}

Schwender D, Daunderer M, Mulzer S, Klasing S, Finsterer U, Peter K. Midlatency auditory evoked potentials predict movements during anesthesia with isoflurane or propofol. *Anesthesia and Analgesia* 1997;**85**(1):164-73. [PUBMED: 9212142]

Schwieger 1989 {published data only}

Schwender D, Faber-Zullig E, Klasing S, Poppel E, Peter K. Motor signs of wakefulness during general anaesthesia with propofol, isoflurane and flunitrazepam/fentanyl and midlatency auditory

evoked potentials. *Anaesthesia* 1994;**49**(6):476-84. [PUBMED: 8017589]

Schwieger IM, Hug CCJ, Hall RI, Szlam F. Is lower esophageal contractility a reliable indicator of the adequacy of opioid anesthesia?. *Journal of Clinical Monitoring* 1989;**5**(3):164-9. [PUBMED: 2769314]

Schwieger 1991 {published data only}

Schwieger IM, Hall RI, Hug CCJ. Assessing the adequacy of fentanyl anesthesia: plasma concentrations and lower esophageal contractility. *Acta Anaesthesiologica Scandinavica* 1991;**35**(3):227-34. [PUBMED: 2038929]

See 2007 {published data only}

See JJ, Lew TWK, Kwek TK, Chin KJ, Wong MFM, Liew QY, et al. Anaesthetic management of awake craniotomy for tumour resection. *Annals of the Academy Medicine, Singapore* 2007;**36**(5):319-25. [PUBMED: 17549277]

Shariffuddin 2007 {published data only}

Shariffuddin II, Wang CY. Randomised crossover comparison of the Ambu AuraOnce Laryngeal Mask with the LMA Classic laryngeal mask airway in paralysed anaesthetised patients. *Anaesthesia* 2008;**63**(1):82-5. [PUBMED: 18086075]

Shiau 2007 {published data only}

Shiau JM, Chen WH, Yang YL, Su HP, Wu YH, Tseng CC. Earlier cessation of desflurane supply in closed-circuit anesthesia reduces emergence time in patients undergoing breast surgery. *Acta Anaesthesiologica Taiwanica* 2007;**45**:21-6. [PUBMED: 17424755]

Shimohata 2007 {published data only}

Shimohata T, Shinoda H, Nakayama H, Ozawa T, Terajima K, Yoshizawa H, et al. Daytime hypoxemia, sleep-disordered breathing, and laryngopharyngeal findings in multiple system atrophy. *Archives of Neurology* 2007;**64**(6):856-61. [PUBMED: 17562934]

Short 1991 {published data only}

Short TG, Chui PT. Propofol and midazolam act synergistically in combination. *British Journal of Anaesthesia* 1991;**67**(5):539-45. [PUBMED: 1751266]

Sidiropoulou 2008 {published data only}

Sidiropoulou T, Buonomo O, Fabbi E, Silvi MB, Kostopanagiotou G, Sabato AF, et al. A prospective comparison of continuous wound infiltration with ropivacaine versus single-injection paravertebral block after modified radical mastectomy. *Anesthesia and Analgesia* 2008;**106**(3):997-1001, table of contents. [PUBMED: 18292452]

Sintavanuruk 2010 {published data only}

Sintavanuruk K, Pongruekdee S, Taharavanich R, Laosuwan S, Charuluxananan S. Comparative study of effective-site target controlled infusion with standard bolus induction of propofol for laryngeal mask airway insertion. *Asian Biomedicine* 2010;**4**(1):177-82.

Skaja 2006 {published data only}

Skaja D, Fgraff M. Bispectral index monitoring during volatile anaesthesia for optimal selection of inhalation anaesthetics in lumbar laminotomy. *Anaesthesiology Intensive Therapy* 2006;**38**:72-5.

Sosner 2010 {published data only}

Sosner P, Rigouzzo A, Sabourdin N, Laude D, Constant I. Non-invasive investigation of cardiovascular autonomic nervous system activity under anesthesia controlled by EEG: a comparative study between sevoflurane and propofol, in children. European Society of Cardiology, ESC Congress 2010. 2010; Vol. 31:1014.

Spaulding 1984 {published data only}

Spaulding BC, Choi SD, Gross JB, Apfelbaum JL, Broderson H. The effect of physostigmine on diazepam-induced ventilatory depression: a double-blind study. *Anesthesiology* 1984;**61**(5):551-4. [PUBMED: 6437287]

Stonell 2006 {published data only}

Stonell CA, Leslie K, He C, Lee L. No sex differences in memory formation during general anesthesia. *Anesthesiology* 2006;**105**(5):920-6. [PUBMED: 17065885]

Struys 2001 {published data only}

Struys MM, De Smet T, Versichelen LF, Van De Velde S, Van den Broecke R, Mortier EP. Comparison of closed-loop controlled administration of propofol using Bispectral Index as the controlled variable versus "standard practice" controlled administration. *Anesthesiology* 2001;**95**(1):6-17. [PUBMED: 11465585]

Suarez 1994 {published data only}

Suarez M, Mendoza CU, Castaneda R, Lux E, Rufino C, Sanchez R. Incidence of memory of transoperative events during total intravenous general anesthesia with propofol. *Archives of Medical Research* 1994;**25**(4):413-7.

Suliman 2007 {published data only}

Suliman S, Ericksen T, Labuschgne P, de Wit R, Stein DJ, Seedat S. Comparison of pain, cortisol levels, and psychological distress in women undergoing surgical termination of pregnancy under local anaesthesia versus intravenous sedation. *BMC Psychiatry* 2007;**7**:24. [PUBMED: 17565666]

Terblanche 2008 {published data only}

Terblanche N, Coetzee JF. A comparison of induction of anaesthesia using two different propofol preparations. *Southern African Journal of Anaesthesia and Analgesia* 2008;**14**(6):25-9.

Ting 2004 {published data only}

Ting CK, Hu JS, Teng YH, Chang YY, Tsou MY, Tsai SK. Desflurane accelerates patient response during the wake-up test for scoliosis surgery. *Canadian Journal of Anaesthesia* 2004;**51**(4):393-7. [PUBMED: 15064271]

Toraman 2013 {published data only}

Toraman F, Ustalar Ozgen S, Sayin Kart J, Ariturk C, Erkek E, Guclu P, et al. Target Controlled Infusion (TCI) of fentanyl and midazolam in coronary artery bypass surgery: effects

on anaesthesia depth and drug consumption. [Turkish]. *Gogus-Kalp-Damar Anestezi ve Yogun Bakim Dernegi Dergisi* 2013;**19**:113-7.

Treggiari 2009 {published data only}

Treggiari MM, Romand JA, Yanez ND, Deem SA, Goldberg J, Hudson L, et al. Randomized trial of light versus deep sedation on mental health after critical illness. *Critical Care Medicine* 2009;**37**(9):2527-34. [PUBMED: 19602975]

Trillo 2009 {published data only}

Trillo L, Garca C, Ramn J, Sadurni M, Molt L. Conference: European Anaesthesiology Congress, EUROANAESTHESIA 2009 Milan Italy. *Euroanaesthesia* 2009;**26**:138.

Tufano 2000 {published data only}

Tufano R, Palomba R, Lambiase G, Giurleo LG. The utility of bispectral index monitoring in general anesthesia. *Minerva Anestesiologica* 2000;**66**(5):389-93. [PUBMED: 10965722]

Tunstall 1981 {published data only}

Tunstall ME, Hawksworth GM. Halothane uptake and nitrous oxide concentration. Arterial halothane levels during Caesarean section. *Anaesthesia* 1981;**36**(2):177-82. [PUBMED: 7212227]

Turan 2010 {published data only}

Turan A, Kasuya Y, Govinda R, Obal D, Rauch S, Dalton JE, et al. The effect of aminophylline on loss of consciousness, bispectral index, propofol requirement, and minimum alveolar concentration of desflurane in volunteers. *Anesthesia and Analgesia* 2010;**110**(2):449-54. [PUBMED: 19955506]

Turner 1969 {published data only}

Turner DJ, Wilson J. Effect of diazepam on awareness during caesarean section under general anaesthesia. *British Medical Journal* 1969;**2**(5659):736-7. [PUBMED: 4891882]

Ueyama 1986 {published data only}

Ueyama H, Tashiro C, Mori T, Yoshiya I. Risk factors for awareness and dreams during general anesthesia. *Masui* 1986;**35**(4):593-8. [PUBMED: 3747105]

Valtonen 1988 {published data only}

Valtonen M, Kanto J, Klossner J. Anaesthesia for cardioversion: a comparison of propofol and thiopentone. *Canadian Journal of Anaesthesia* 1988;**35**(5):497-83. [PUBMED: 3048754]

Vanacker 2002 {published data only}

Vanacker B, Van Geldre L. A randomized study of the efficacy and recovery of remifentanyl-based and alfentanil anaesthesia with desflurane or sevoflurane for gynecological surgery. *Acta Anaesthesiologica Scandinavica* 2002;**53**(1):21-6. [PUBMED: 11975425]

Vanacker 2007 {published data only}

Vanacker BF, Vermeyen KM, Struys MMRF, Rietbergen H, Vandermeersch E, Saldien V, et al. Reversal of rocuronium-induced neuromuscular block with the novel drug sugammadex is equally effective under maintenance anesthesia with propofol or sevoflurane. *Anesthesia and Analgesia* 2007;**104**(3):563-8. [PUBMED: 17312209]

Vann 2007 {published data only}

Vann MA, Ogunnaike BO, Joshi GP. Sedation and anesthesia care for ophthalmologic surgery during local/regional anesthesia. *Anesthesiology* 2007;**107**(3):502-8. [PUBMED: 17721254]

Velly 2007 {published data only}

Velly LJ, Rey MF, Bruder NJ, Gouvitsos FA, Witjas T, Regis JM, et al. Differential dynamic of action on cortical and subcortical structures of anesthetic agents during induction of anesthesia. *Anesthesiology* 2007;**107**(2):202-12. [PUBMED: 17667563]

Wanatabe 1984 {published data only}

Watanabe A, Namiki A, Omote T, Takahashi T, Ichinose H. Evaluation of isolated forearm technique for detection of wakefulness during endotracheal intubation procedures. *Masui* 1984;**33**(8):870-4. [PUBMED: 6512958]

Wang 2005 {published data only}

Wang Y, Yue Y, Sun YH, Wu AS, Wu QW, Zhang YQ, et al. Investigation and analysis of incidence of awareness in patients undergoing cardiac surgery in Beijing, China. *Chinese Medical Journal* 2005;**118**(14):1190-4. [PUBMED: 16117864]

Wang 2007 {published data only}

Wang SM, Constable RT, Tokoglu FS, Weiss DA, Freyle D, Kain ZN. Acupuncture-induced blood oxygenation level-dependent signals in awake and anesthetized volunteers: a pilot study. *Anesthesia and Analgesia* 2007;**105**(2):499-506. [PUBMED: 17646512]

Watanabe 1998 {published data only}

Watanabe A, Namiki A, Ujike Y, Watanabe H, Aoki M. Wakefulness during the induction with high-dose fentanyl and oxygen anesthesia. *Journal of Anesthesia* 1998;**2**(2):165-9. [PUBMED: 15236075]

Weber 2005 {published data only}

Weber F, Pohl F, Hollnberger H, Taeger K. Impact of the Narcotrend Index on propofol consumption and emergence times during total intravenous anaesthesia with propofol and remifentanyl in children: a clinical utility study. *European Journal of Anaesthesiology* 2005;**22**(10):741-7. [PUBMED: 16211731]

Weber 2009 {published data only}

Weber F, Kriek N, Blusse H. Conference: European Anaesthesiology Congress, EUROANAESTHESIA 2009 Milan Italy; Influence of a muscle relaxant bolus on BIS, Cerebral State Index (CSI) and Spectral Edge Frequency (SEF) in children under propofol anaesthesia.. *Euroanaesthesia* 2009;**26**:144.

Wellisch 2012 {published data only}

Wellisch OM, Kronenfeld M, Saberito D, Pagala M, Choueka J, Gupta P. A prospective randomized controlled trial comparing the efficacy and safety of dexmedetomidine versus propofol TIVA in conjunction with regional block for shoulder surgery in beach chair sitting position. *Regional Anesthesia and Pain Medicine* 2012;**0**(6):1.

Wihelm 2000 {published data only}

Wihelm W, Grundmann U, Aken HV, Haus EM, Larsen R. A multicenter comparison of isoflurane and propofol as adjuncts to remifentanil-based anesthesia. *Journal of Clinical Anesthesia* 2000;**12**(2):129-35. [PUBMED: 10818327]

Wilson 1970 {published data only}

Wilson J, Lewis SA, Jenkinson JL. Electroencephalographic investigation of awareness during anaesthesia. *British Journal of Anaesthesia* 1970;**42**(9):804-5. [PUBMED: 5470077]

Wu 2005 {published data only}

Wang Y, Yue Y, Sun YH, Wu AS, Wu QW, Zhang YQ, et al. Investigation and analysis of incidence of awareness in patients undergoing cardiac surgery in Beijing, China. *Chinese Medical Journal* 2005;**118**(14):1190-4. [PUBMED: 16117864]

Xu 2009 {published data only}

Xu L, Wu AS, Yue Y. The incidence of intra-operative awareness during general anesthesia in China: a multi-center observational study. *Acta Anaesthesiologica Scandinavica* 2009;**53**(7):873-82. [PUBMED: 19496761]

Yan 2005 {published data only}

Yan M, Wang Y, Hu XS, Cheng W, Liu ZM. Target-controlled infusion of remifentanil and propofol during operation with suspension laryngoscopy. *Zhejiang Da Xue Xue Bao Yi Xue Ban [Journal of Zhejiang University Medical Sciences]* 2005;**34**(6):557-61, 565. [PUBMED: 16331821]

Yan 2014 {published data only}

Yan J, Lin CQ, Zhang CZ, Ou JY, Luo FR, Zhang SL, et al. Effects of assisted-electroacupuncture on recovery of fast tracking anesthesia in mPCNL. *Zhongguo Zhen Jiu [Chinese Acupuncture & Moxibustion]* 2014;**34**(4):385-8.

Yang 1994 {published data only}

Yang LH, Lin S M, Lee WY, Liu CC. Intraoperative transcranial electrical motor evoked potential monitoring during spinal surgery under intravenous ketamine or etomidate anaesthesia. *Acta Neurochirurgica* 1994;**127**(3-4):191-8. [PUBMED: 7942202]

Yeh 2009 {published data only}

Yeh YC, Lin TF, Chang HC, Chan WS, Wang YP, Lin CJ, et al. Combination of low-dose nalbuphine and morphine in patient-controlled analgesia decreases incidence of opioid-related side effects. *Journal of the Formosan Medical Association* 2009;**108**(7):548-53. [PUBMED: 19586828]

Yi 2008 {published data only}

Yi C, Jee D. Influence of the anaesthetic depth on the inhibition of the oculocardiac reflex during sevoflurane anaesthesia for paediatric strabismus surgery. *British Journal of Anaesthesia* 2008;**101**(2):234-8. [PUBMED: 18524784]

Yildiz 2007 {published data only}

Yildiz TS, Solak M, Toker K. Comparison of laryngeal tube with laryngeal mask airway in anaesthetized and paralysed patients. *European Journal of Anaesthesiology* 2007;**24**(7):620-5. [PUBMED: 17376250]

Zhang 2009 {published data only}

Zhang GH, Sun L. Peri-intubation hemodynamic changes during low dose fentanyl, remifentanil and sufentanil combined with etomidate for anesthetic induction. *Chinese Medical Journal* 2009;**122**(19):2330-4. [PUBMED: 20079135]

Zohar 2007 {published data only}

Zohar E, Noga Y, Davidson E, Kantor M, Fredman B. Perioperative patient safety: correct patient, correct surgery, correct side--a multifaceted, cross-organizational, interventional study. *Anesthesia and Analgesia* 2007;**105**(2):443-7. [PUBMED: 17646503]

References to studies awaiting assessment
Aceto 2015 {published data only}

Aceto P, Perilli V, Lai C, Sacco T, Modesti C, Luca E, et al. Minimum alveolar concentration threshold of sevoflurane for postoperative dream recall. *Minerva Anestesiologica* 2015; Vol. 81, issue 11:1201-9.

Asouhidou 2015 {published data only}

Asouhidou I, Trikoupi A. Esmolol reduces anesthetic requirements thereby facilitating early extubation; a prospective controlled study in patients undergoing intracranial surgery. *BMC Anesthesiology* 2015;**15**(11):172.

Elbadawy 2015 {published data only}

Elbadawy AM, Khidr AM, Saleh AA. Comparative study of implicit memory during bispectral index guided total intravenous anesthesia versus sevoflurane inhalation anesthesia. *Egyptian Journal of Anaesthesia* 2015;**31**(1):9-13.

Hoymork 2007 {published data only}

Hoymork SC, Hval K, Jensen EW, Raeder J. Can the cerebral state monitor replace the bispectral index in monitoring hypnotic effect during propofol/remifentanil anaesthesia?. *Acta Anaesthesiologica Scandinavica* 2007;**51**(2):210-6. [PUBMED: 17330331]

Jiang 2016 {published data only}

Jiang A, Chen LJ, Wang YX, Li MC, Ding YB. The effects of different methods of anaesthesia for laparoscopic radical gastrectomy with monitoring of entropy. *Anticancer Research* 2016;**36**(3):1305-8.

Khanjani 2014 {published data only}

Khanjani S, Naghibi K, Azarnoush H. Awareness and Apgar score in elective Cesarean section under general anesthesia with propofol or Isoflurane: a prospective, randomized, double-blinded clinical trial study. *Advances in Biomedical Engineering Research* 2014;**3**:234.

Lequeux 2014 {published data only}

Lequeux PY, Hecquet F, Bredas P. Does anesthetic regimen influence implicit memory during general anesthesia?. *Anesthesia and Analgesia* 2014;**119**(5):1174-9.

Mehmandoost 2013 {published data only}

Mehmandoost MK, Naghibi K. Comparative study of the effects of two anesthetic methods with propofol and isoflurane on mother's awareness during the operation and Apgar score in the newborns delivered by elective cesarean section. *Journal of Isfahan Medical School*; <http://english.mui.ac.ir/content/iums-journals> 2013;**31**(228):265-73.

Rajan 2015 {published data only}

Rajan S, Hassain A, Puthenveetil N, Kumar L. Efficacy and safety of low-dose ketamine as an adjunct analgesic and amnesic during caesarean section under general anaesthesia. *Indian Journal of Anaesthesia* 2015;**59**(10):653-7.

Xie 2015 {published data only}

Xie M, Lan ZX, Liu YF. Effects of sevoflurane and propofol on preoperative implicit and explicit memories in general anaesthesia patients of elective surgery: a randomized controlled trial. *Chinese Journal of Evidence-Based Medicine* 2015;**15**(9):1000-4.

Additional references
Andrade 2008

Andrade J, Deeprose C, Barker I. Awareness and memory function during paediatric anaesthesia. *British Journal of Anaesthesia* 2008; Vol. 100, issue 3:389-96. [PUBMED: 18222981]

Artusio 1955

Artusio JF Jr. Ether analgesia during major surgery. *JAMA* 1955;**157**(1):33-6.

ASA 2006

ASA Task Force. Practice advisory for intraoperative awareness and brain function monitoring: a report by the American Society of Anesthesiologists Task Force on intraoperative awareness. *Anesthesiology* 2006;**104**(4):847-64.

Benhamou 2003

Benhamou D, Ghosh C, Mercier FJ. A randomized sequential allocation study to determine the minimum effective analgesic concentration of levobupivacaine and ropivacaine in patients receiving epidural analgesia for labor. *Anesthesiology* 2003;**99**(6):1383-6. [PUBMED: 14639153]

Breen 2007

Breen T. Lawsuit: Wide-Awake Surgery Led to Death. Retrieved September 19, 2016. Fox News Channel April 10, 2007.

Brull 2008

Brull SJ, Naguib M, Miller RD. Residual neuromuscular block: rediscovering the obvious. *Anesthesia and Analgesia* 2008; Vol. 107, issue 1:11-4.

Chhabra 2016

Chhabra A, Subramaniam R, Srivastava A, Probhakar H, Kalaivani M, Paranjape S. Spectral entropy monitoring for adults and children undergoing general anaesthesia. *Cochrane Database of Systematic Reviews* 2016, Issue 3. [DOI: [10.1002/14651858.CD010135.pub2](https://doi.org/10.1002/14651858.CD010135.pub2)]

Cundy 1995

Cundy JM. Post traumatic stress disorders (Letter). *British Journal of Anaesthesia* 1995; Vol. 75:501-2. [PUBMED: 7488502]

Dean 2007

Dean CB, Nielsen JD. Generalized linear mixed models: a review and some extensions. *Lifetime Data Analysis* 2007;**13**(4):497-512.

Eger 2005

Eger EI, Sonner JM. How likely is awareness during anesthesia?. *Anesthesia and Analgesia* 2005;**100**(5):1544-5. [PUBMED: 15845735]

Elmagarmid 2014

Elmagarmid A, Fedorowicz A, Hammady H, Ilyas I, Khabsa M, Ouzzani M. Rayyan: a systematic reviews web app for exploring and filtering searches for eligible studies for Cochrane Reviews. Evidence-Informed Public Health: Opportunities and Challenges. Abstracts of the 22nd Cochrane Colloquium; 2014 21-26 Sep; Hyderabad, India. John Wiley & Sons, 2014.

EUC 2009

EUC. Useful Information on Rare Diseases from an EU Perspective. Retrieved 19 May 2009. European Commission: Health & Consumer Protection Directorate-General. Luxembourg: Commission Européenne, 2009.

Girgirah 2006

Girgirah K, Kinsella SM. Propofol and memory. *British Journal of Anaesthesia* 2006; Vol. 97, issue 5:746-7; author reply 747-8. [PUBMED: 17032664]

Guyatt 2011

Guyatt GH, Oxman AD, Vist G, Kunz R, Brozek J, Alonso-Coello P, et al. GRADE guidelines: 4. Rating the quality of evidence--study limitations (risk of bias). *Journal of Clinical Epidemiology* 2011;**64**(4):407-15.

Higgins 2002

Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine* 2002;**21**(11):1539-58. [PUBMED: 12111919]

Higgins 2011

Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

HR4013

HR4013. Public Law 107-280. An act to amend the Public Health Service Act to establish an Office of Rare Diseases at the National Institutes of Health, and for other purposes. United States Government Printing Office. Washington, D.C., 2002; Vol. H.R. 4013, issue H. Rept. 107-543:AE 2.110:107-280.

Jones 1994

Jones JG. Perception and memory during general anaesthesia. *British Journal of Anaesthesia* 1994;**73**(1):31-7. [PUBMED: 8038054]

Kettner 2014

Kettner S. Not too little, not too much: delivering the right amount of anaesthesia during surgery. *Cochrane Database of Systematic Reviews* 2014;**6**:ED000084.

Lacassie 2004

Lacassie HJ, Lacassie HP, Muir HA. Is the minimum local analgesic concentration method robust enough?. *Anesthesiology* 2004;**101**(2):550; author reply 550-1. [PUBMED: 15277943]

Lenmarken 2002

Lenmarken C, Bildfors K, Enlung G, Samuelsson P, Sandin R. Victims of awareness. *Acta Anaesthesiologica Scandinavica* 2002; Vol. 46, issue 3:229-31. [PUBMED: 11939910]

Mainzer 1979

Mainzer J Jr. Awareness, muscle relaxants and balanced anaesthesia. *Canadian Anaesthetists' Society Journal* 1979;**26**(5):386-93. [PUBMED: 385118]

Malek 2010a

Malek J, Marecek F, Hess L. A combination of dexmedetomidine with ketamine and opioids results in significant inhibition of hemodynamic changes associated with laparoscopic cholecystectomy and in prolongation of postoperative analgesia. *Rozhledy V Chirurgii* 2010;**89**:275-81.

Mashour 2009

Mashour G, Tremper KK, Avidan MS. Protocol for the "Michigan Awareness Control Study": a prospective, randomized, controlled trial comparing electronic alerts based on bispectral index monitoring or minimum alveolar concentration for the prevention of intraoperative awareness. *BMC Anesthesiology* 2009;**9**(7):1.

Merikle 1996

Merikle PM, Daneman M. Memory for unconsciously perceived events: evidence from anesthetized patients. *Consciousness & Cognition* 1996; Vol. 5, issue 4:525-41. [PUBMED: 9063614]

Murphy 2008

Murphy GS, Szokol JW, Marymont JH, Greenberg SB, Avram MJ, Vender JS. Residual neuromuscular blockade and critical respiratory events in the postanesthesia care unit. *Anesthesia and Analgesia* 2008; Vol. 107, issue 1:130-7. [1526-7598: (Electronic)]

Myles 2015 (personal communication)

Myles PS. Professor/Director of Anaesthesia and Perioperative Medicine, Alfred Hospital and Monash University, Melbourne, Australia. Letter to: Anthony Messina 5 April 2015.

Osterman 2001

Osterman JE, Hopper J, Heran WJ, Keane TM, van der Kolk BA. Awareness under anesthesia and the development of posttraumatic stress disorder. *General Hospital Psychiatry* 2001; Vol. 23, issue 4:198-204. [PUBMED: 11543846]

Punjasawadwong 2014

Punjasawadwong Y, Phongchiewboon A, Bunchungmongkol N. Bispectral index for improving anaesthetic delivery and postoperative recovery. *Cochrane Database of Systematic Reviews* 2014, Issue 6. [DOI: [10.1002/14651858.CD003843.pub3](https://doi.org/10.1002/14651858.CD003843.pub3)]

R 2016 [Computer program]

R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria, 2016.

Rampil 1998

Rampil IJ. A primer for EEG signal processing in anesthesia. *Anesthesiology* 1998;**89**(4):980-1002. [PUBMED: 9778016]

Review Manager [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Russell 1985

Russell IF. Balanced anaesthesia: does it anaesthetize?. *Anesthesia and Analgesia* 1985; Vol. 64, issue 9:941-2. [PUBMED: 4025858]

Russell 1993

Russell IF. Midazolam-alfentanil: an anaesthetic? An investigation using the isolated forearm technique. *British Journal of Anaesthesiology* 1993;**70**(1):42-6. [PUBMED: 8431332]

Sanders 2011

Sanders RD, Sleigh JW. Prevention of intraoperative awareness. *New England Journal of Medicine* 2011;**365**(21):2033; author reply 2034-5.

Sandin 2000

Sandin RH, Enlund G, Samuelsson P, Lenmarken C. Awareness during anaesthesia: a prospective case study. *Lancet* 2000; Vol. 355, issue 9205:707-11. [PUBMED: 10703802]

Schulz 2010

Schulz KF, Altman DG, Moher D, CONSORT Group. CONSORT 2010 statement: Updated guidelines for reporting parallel group randomised trials. *BMJ* 2010;**340**(c):332.

Sebel 2004

Sebel PS, Bowdle TA, Ghoneim MM, Rampil IJ, Padilla RE, Gan TJ, et al. The incidence of awareness during anesthesia: a multicenter United States study. *Anesthesia and Analgesia* 2004; Vol. 99, issue 3:833-9, table of contents. [PUBMED: 15333419]

Tian 2008

Tian L, Cai T, Pfeffer M, Piankov N, Cremieux P, Wei LJ. Exact and efficient inference procedure for meta-analysis and its application to the analysis of independent two by two tables with all available data but without artificial continuity correction. *Biostatistics* 2008;**10**(2):275-81. [PUBMED: 18922759]

Tunstall 1977

Tunstall ME. Detecting wakefulness during general anaesthesia for caesarean section. *British Medical Journal* 1977;**1**(6072):1321. [PUBMED: 861598]

Vakkuri 2000

Vakkuri A, Yli-Hankala A, Llundgren L. BIS-monitored depth of anaesthesia in laparoscopic tubal ligation. Memory and Awareness in Anaesthesia IV. Proceedings of the 4th International Symposium on Memory and Awareness in Anaesthesia. London: Imperial College Press, 2000:304-11.

Viechtbauer 2010

Viechtbauer W. Conducting meta-analyses in {R} with the {metafor} package. *Journal of Statistical Software* 2010;**36**(3):1-48. [1548-7660]

Villafranca 2013

Villafranca AJ, Arenson BG, Avidan MS, Glick D, Mashour GA, Jacobsohn E. Volitional delay of self-reported outcomes: insights from a case of intraoperative awareness with explicit recall. *Anesthesia and Analgesia* 2013;**116**(2):365-7. [PUBMED: 23302976]

Wahl 2011

Wahl RU, Wurpts G, Merk HF. Perioperative anaphylaxis caused by rocuronium allergy in a child. *Hautarzt* 2011;**62**(11):812-4.

Wang 2000

Wang M. The psychological consequences of awareness during surgery. In: Jordan C, Vaughan DJA, Newton DEF editor(s). Memory and Awareness in Anaesthesia IV. London: Imperial College Press, 2000:315-24.

Wang 2012

Wang M, Messina AG, Russell IF. The topography of awareness: a classification of intra-operative states. *Anaesthesia* 2012;**67**(11):1197-201. [PUBMED: 23033820]

White 1987

White DC. Consciousness Awareness and Pain in General Anaesthesia. 7. London: Oxford: Butterworths, 1987.

WHO 2002

WHO. Pharmacovigilance: Frequency of adverse drug reactions. World Health Organization (http://www.who.int/medicines/areas/quality_safety/safety_efficacy/trainingcourses/definitions.pdf). Council of International Organizations of Medical Science (CIOMS), 2002.

WHO 2015

WHO. Annex 1, 19th WHO Model List of Essential Medicines (April 2015). Explanatory notes. World Health Organization (WHO) (http://www.who.int/medicines/publications/essentialmedicines/EML2015_8-May-15.pdf) 2015.

Wilson 1980

Wilson ME. Awareness in general anaesthesia. *British Medical Journal* 1980;**280**:1270.

Yu 2013 [Computer program]

Yu Y, Tian L. Exactmeta: Exact fixed effect meta analysis. R package version 1.0-1 [<http://CRAN.R-project.org/package=exactmeta>]. Stanford University, 2013.

References to other published versions of this review
Messina 2008

Messina AG, Ward MJ, Pace NL. Anaesthetic interventions for prevention of awareness during surgery. *Cochrane Database of Systematic Reviews* 2008, Issue 3. [DOI: [10.1002/14651858.CD007272](https://doi.org/10.1002/14651858.CD007272)]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES
Characteristics of included studies [ordered by study ID]
Abboud 1985

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: USA Sex: female Age: mean 27, 24, 24, 26, 25 ASA: not stated ("healthy") Procedure: caesarean delivery ("elective") Study size: 81
Interventions	Randomized portion of anaesthetic: volatile agent doses

Anaesthetic interventions for prevention of awareness during surgery (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Abboud 1985 (Continued)

- Intervention 1: maintenance: general anaesthesia 50% N₂O (N = 16): control
- Intervention 2: maintenance: general anaesthesia 0.25% halothane + 50% N₂O (N = 16)
- Intervention 3: maintenance: general anaesthesia 0.5% halothane + 50% N₂O (N = 18)
- Intervention 4: maintenance: general anaesthesia 0.5% enflurane + 50% N₂O (N = 18)
- Intervention 5: maintenance: general anaesthesia 1.0% enflurane + 50% N₂O (N = 13)

Outcomes	<p>Primary outcomes: Quote: "adverse neonatal effects of sub anesthetic doses of halothane or enflurane."</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4</p> <p>Quote: "Of the patients who had N₂O alone [Intervention 1], 12.5% (2/16) had awareness versus none in the other groups"</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of potent inhalational technique (volatile agent)/ N₂O yes/hypnotic/supplemental narcotics/muscle relaxants induction yes/maintenance unclear</p> <p>Anaesthesia induction: 4 mg/kg thiopental + succinylcholine 1.5 mg/kg + curare 3 mg; anaesthesia maintenance after delivery: N₂O + narcotics</p> <p>No titration anaesthetic strategy for light anaesthesia defined in the paper</p> <p>Comment: no sample size calculation (power analysis) provided in paper</p> <p>Time of outcome determination: 24 h post partum</p> <p>Method of outcome determination: interview</p> <p>Email sent jsl3nov42@webtv.net 3 November 2010; as with most RCTs that were published years ago, it was difficult to find current contact information for most authors</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "patients were randomly assigned"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "All patients were interviewed 24 h post-partum to determine the incidence of awareness by a person who was unaware of the anesthetic management of the mother"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data

Abboud 1985 (Continued)

Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information provided

Abboud 1989

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: USA Sex: female Age: mean 28 ASA: not stated ("healthy") Procedure: caesarean delivery ("elective primary or repeat, or arrest of dilatation") Study size: 60
Interventions	Randomized portion of anaesthetic: volatile agent doses Intervention 1: maintenance: 0.5% isoflurane (N = 20) Intervention 2: maintenance: 1 % isoflurane (N = 20) Intervention 3: maintenance: 0.5% halothane (N = 20) 50% N ₂ O and O ₂ was added in each case
Outcomes	Primary outcomes: cbc chemistry urine analysis etc.; the maternal and neonatal effects of isoflurane or halothane for caesarean section Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "...none of the patients developed intraoperative awareness"
Notes	Non-randomized portion of anaesthetic: parts of potent inhalational technique (volatile agent)/ N₂O yes/hypnotic/supplemental narcotics/muscle relaxants induction yes/unclear maintenance Dryad topic from this RCT: light and deep anaesthesia and operating room turnover 1940s to current time Time of outcome determination: not stated Method of outcome determination: not stated ROB survey emailed jsl3nov42@webtv.net

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "patients were randomly assigned"

Abboud 1989 (Continued)

Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist and postoperative interviewer: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: no information provided. Although Cochrane guidelines allow review authors to make judgements about domain grades based on previous studies by the author (Abboud 1985), our editorial team prefers that we only assess what is written in the current study
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information provided

Abboud 1995a

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: USA Sex: female Age: mean 28 to 29 ASA: not stated Procedure: caesarean delivery Study size: 75
Interventions	Randomized portion of anaesthetic: volatile agent doses Pre-delivery: induction/maintenance Intervention 1: desflurane end-tidal 3% (N = 25) Intervention 2: desflurane end-tidal 6% (N = 25) Intervention 3: enflurane end-tidal 0.6% (N = 25)
Outcomes	Primary outcomes: cbc chemistry urine analysis etc. Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "none of the patients developed intraoperative awareness."
Notes	Non-randomized portion of anaesthetic: parts of potent inhalational technique (volatile agent)/ N₂O yes/hypnotic/supplemental narcotics/muscle relaxants induction yes/unclear maintenance

Abboud 1995a (Continued)

Rapid sequence induction regimen thiopental + succinylcholine

 Anaesthesia maintenance: before delivery N₂O 50% and O₂; after delivery N₂O 67% + butorphanol 1 mg to 2 mg and **reduced doses of desflurane and enflurane**

 ROB survey emailed jsl3nov42@webtv.net
Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "patients were randomly assigned"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist and postoperative interviewer: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: anaesthesiologist and postoperative interviewer: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information provided

Abboud 1995b

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: USA Sex: female Age: mean 30 ASA: not stated Procedure: caesarean delivery ("repeat, failure to progress, malpresentation, placenta praevia (non-bleeding)") Study size: 74
Interventions	Randomized portion of anaesthetic: parts of IV MCI Intervention 1: induction: propofol 1.5 mg to 2.5 mg/kg; maintenance: propofol 0.2 mg/kg/min and reduced to 0.05 mg/kg/min (N = 37)

Abboud 1995b (Continued)

 intervention 2: induction: thiamylal 3 mg to 4 mg/kg; maintenance: 0.25% to 0.75% **isoflurane** (N = 37)

Outcomes	Primary outcomes: cbc chemistry urine analysis etc. Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Quote: "One patient in the isoflurane group reported awareness of the surgical procedure when interviewed 24 h later (Table 1)"
Notes	<p>Non-randomized portion of anaesthetic: parts of IV N₂O yes/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance unclear</p> <p>Anaesthesia induction: group specific induction regimen + succinylcholine 1.5 mg/kg; anaesthesia maintenance: before delivery N₂O 50%; after delivery N₂O 67% + butorphanol 1 mg to 2 mg</p> <p>No titration anaesthetic strategy for light anaesthesia defined in the paper</p> <p>Time of outcome determination: 24 h post partum</p> <p>Method of outcome determination: interview</p> <p>ROB survey emailed jsl3nov42@webtv.net</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "patients were randomly assigned"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist and postoperative interviewer: no information provided Comment: difficult to blind anaesthesiologist to inhalational agents. Older study when very light anaesthesia used (all groups) compared to today's standards. That low-dose inhalational agents had a significant effect on awareness rates is consistent with other studies. Therefore, the lack of blinding unlikely to impact the outcomes (Cochrane Handbook table 8.5.c).
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "All patients were interviewed 24-h post-partum to determine the incidence of awareness by a person who was unaware of the anesthetic management of the mother"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information provided

Aceto 2002

Methods	Study design: randomized parallel groups
---------	--

Aceto 2002 (Continued)

	Study dates: not stated
Participants	<p>Country: Italy</p> <p>Sex: both</p> <p>Age: NA "no statistically significant differences among the group regarding age"</p> <p>ASA: I-II</p> <p>Procedure: elective laparoscopic cholecystectomy</p> <p>Study size: 32</p>
Interventions	<p>Randomized portion of anaesthetic: volatile agent types ±N₂O</p> <p>Intervention 1: maintenance: (Group A) sevoflurane + air (FIO₂ 40%) (N = 8)</p> <p>Intervention 2: maintenance: (Group B) sevoflurane + N₂O (60%) in oxygen 40% (N = 8)</p> <p>Intervention 3: maintenance: (Group C) isoflurane + air (FIO₂ 40%) (N = 8)</p> <p>Intervention 4: maintenance: (Group D) isoflurane + N₂O (60%) in oxygen 40% (N = 8)</p> <p>Inhalation dose not specified except for 1 MAC prior to MLAER recordings</p>
Outcomes	<p>Primary outcomes: Quote: "latency of the Pa waves in patients with or without subconscious awareness before and during anaesthesia"</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 2</p> <p>Quote: "In the postoperative interview none of the patients was able to recollect explicit memories of intra-operative events"</p> <p>Quote: "One of the patients from isoflurane group (Group C) showed implicit memory of the intra-operative tape story"</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent: N₂O no/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance PRN/ADM: MLAERs</p> <p>Anaesthesia induction: thiopental sodium 5 mg/kg, fentanyl 3 mg/kg and vecuronium 0.08 mg/kg. Anaesthesia maintenance: see above and boluses of fentanyl 2 µg/kg and vecuronium PRN clinical necessity</p> <p>Comment: the authors: 4 interventions MAC equivalents</p> <p>"MLAERs were recorded before anaesthesia, at 1 MAC mechanisms of dreams"</p> <p>"We think that, during anaesthesia, dreams imply a mental state in the transition zone between explicit and implicit memory (subconscious awareness)"</p> <p>Time of outcome determination: "An interview was conducted in the hospital about 24 h after awakening for assessing explicit and implicit memory"</p> <p>Method of outcome determination: explicit recall test and implicit free association test; author's survey sent 26 November 2013 gdecosmo@rm.unicatt.it</p>
Risk of bias	
Bias	Authors' judgement Support for judgement
Random sequence generation (selection bias)	Unclear risk Quote: "For the maintenance of anaesthesia patients were randomly..."

Aceto 2002 (Continued)

Comment: although Cochrane policy allows review authors to use study authors' methods from other RCT papers that they have published, this approach was discouraged by the editors of this Cochrane review. The same authors in [Aceto 2003](#) were more specific in their description of randomization and it had a low risk rating

Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: information provided is inadequate
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "In the postoperative interview none of the patients was able to recollect explicit memories of intra-operative events"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information provided

Aceto 2003

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Italy Sex: both Age: 18 to 70 ASA: I and II Procedure: elective laparoscopic cholecystectomy Study size: 40
Interventions	Randomized portion of anaesthetic: volatile agent types ±N₂O Intervention 1: maintenance: (Group A) sevoflurane + air (FIO ₂ 40%) (N = 10) Intervention 2: maintenance: (Group B) sevoflurane + N ₂ O (60%) in oxygen 40% (N = 10) Intervention 3: maintenance: (Group C) isoflurane + air (FIO ₂ 40%) (N = 10) Intervention 4: maintenance: (Group D) isoflurane + N ₂ O (60%) in oxygen 40% (N = 10)
Outcomes	Primary outcomes: Quote: "latency of the Pa waves in patients with or without subconscious awareness before and during anaesthesia."

Aceto 2003 (Continued)

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 2

Quote: "In the postoperative interview, none of the patients were able to recollect explicit memories of intraoperative events"

Quote: "One of the male patients from the isoflurane+air group (Group C) showed implicit memory of the intraoperative tape story"

Quote: "A dream-like process, related to the story played during anaesthesia, occurred in one of the female patients in the sevoflurane+nitrous oxide group"

Notes

Non-randomized portion of anaesthetic: parts of volatile agent: N₂O no/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance PRN/ADM MLAERs

Anaesthesia induction: thiopental sodium 5 mg/kg, fentanyl 3 mg/kg, and vecuronium bromide 0.08 mg/kg. Anaesthesia maintenance: see above + boluses of fentanyl 2 µg/kg were given according to clinical necessity.

MLAERs were recorded

Subconscious awareness:

Quote: "The two patients with subconscious awareness were in the group with a Pa latency increase less than the 10th percentile. This cut-off showed a sensitivity of 100% and a specificity of 95%, with a positive predictive value of 75% and a negative predictive value of 100% in distinguishing patients with subconscious awareness...increase in Pa latency was related to subconscious awareness"

Time of outcome determination: an interview was conducted in the hospital about 24 h after awakening for assessing explicit and implicit memory

Method of outcome determination: explicit recall test and implicit free association test

Author's survey sent 26 November 2013 gdecosmo@rm.unicatt.it

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "For the maintenance of anaesthesia, patients were assigned, using randomisation tables, to one of four anaesthetic regimen groups..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: information provided is inadequate
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: Aceto 2002
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria

Aceto 2003 (Continued)

Other bias	Unclear risk	Comment: no information provided
------------	--------------	----------------------------------

Adams 1994

Methods	Study design: double-blind, randomized parallel groups Study dates: Quote: "ca. 1991 - 1993" (ROB survey)
Participants	Country: Germany Sex: both Age: greater than 60; mean 68 ASA: I-II Procedure: elective hip or knee replacement Study size: 37 (40 enrolled)
Interventions	<p>Randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia/induction/dissociative agent MCI</p> Intervention 1: induction: 1 mg/kg S-(+)-ketamine; maintenance: 2 mg/kg/h S-(+)-ketamine (N = 20) Intervention 2: induction: 2 mg/kg racemic ketamine; maintenance: 4 mg/kg/h racemic ketamine (N = 20)
Outcomes	Primary outcomes: endocrine stress response - recovery - haemodynamic reaction Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "no intraoperative awareness was reported." Quote: "After racemic ketamine, 1 patient remembered a negative dream and 1 patient a positive dream. In the S (+)-group, 1 positive dream was reported"
Notes	<p>Non-randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia/N₂O no: narcotics and/or hypnotics bolus/muscle relaxants induction yes/maintenance unclear</p> Anaesthetic induction: 0.1 mg/kg midazolam + 0.5 mg atropine + 2 mg vecuronium + 1.5 mg/kg suxamethonium + 0.1 mg/kg vecuronium; anaesthetic maintenance: see above interventions Time of outcome determination: before discharge Method of determination: interview Translator: Lore Schultheiss adams.ha@mh-hannover.de ROB survey response 24 January 2011

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "were investigated in a double-blind, randomised design...."

Adams 1994 (Continued)

		ROB survey response by Dr Adams: "Random by computer program with random numbers. Random numbers were opened not earlier then the study was finished"
Allocation concealment (selection bias)	Low risk	ROB survey response: "Double blinded study to patient and anaesthetist as well. Random numbers were opened not earlier then the study was finished"
Blinding of participants and personnel (performance bias) All outcomes	Low risk	ROB survey response: "Patient, Anesthesiologist, Awareness outcome assessor (interviewer)"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	ROB survey response: "Patient, Anesthesiologist, Awareness outcome assessor (interviewer)"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "Three patients in the ketamine-racemate group showed severe arterial hypertension and were withdrawn from the study" Comment: imbalanced exclusion of 3/40 patients at high risk of awareness, but no significant difference between groups (3/20 vs 0/20): Peto OR 8.23 (0.81 to 84.07)
Selective reporting (reporting bias)	Low risk	ROB survey response: "Study protocol is not more available (study before 2000). The published record included all expected data and outcomes, no selective reporting. Non significant results were reported to show missing differences between the groups"
Other bias	Low risk	ROB survey response: "Limited number of patients"

Agarwal 1977

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: India Sex: both Age: 15 to 50 ASA: not stated Procedure: Quote: "elective surgical operations": appendectomy, exploratory laparotomy, gastro-jejunoscopy and vagotomy, partial gastrectomy, pyelo-nephrolithotomy, nephrectomy, herniorrhaphy, suprapubic cystolithotomy, fallopian tube ligation Study size: 138
Interventions	Randomized portion of anaesthetic: IV: premedication Intervention 1: premedication: atropine 0.6 mg Intervention 2: premedication: atropine 0.6 mg + diazepam 0.15 mg/kg Intervention 3: premedication: atropine 0.6 mg + pethidine 1.5 mg/kg
Outcomes	Primary outcomes: postoperative recall of intraoperative awareness: Quote "ability premedication to lessen frequency unpleasant recall when patients premedicated with narcotics before nitrous oxide-oxygen-relaxant anaesthesia": class 1

Agarwal 1977 (Continued)

Quote: "No patient in any of the three groups reported pain during surgery or awareness of the surroundings"

Notes

Non-randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia: N₂O yes/narcotic/hypnotic/muscle relaxants induction yes/maintenance yes

Anaesthesia induction: thiopental 7 mg/kg + suxamethonium 1 mg/kg + lignocaine 4%; anaesthesia maintenance: O₂:N₂O 2:1 + curare 0.6 mg/kg then bolus to maintain apnoea; reversed neostigmine + atropine 1.2 mg

Time of outcome determination: in operating room, 30 to 45 min after surgery, on ward 4 to 8 h after surgery

Method of outcome determination: interview

Comment: no awareness reports is interesting in setting of only O₂/N₂O/curare major surgery and randomization was premed; these results are contrary to the merged studies of comparison 4 (benzodiazepines vs other drugs)

No email could be found for author survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "patients were allocated randomly to one of three groups..."
Allocation concealment (selection bias)	Unclear risk	Comment: as above
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist and postoperative interviewer: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: anaesthesiologist and postoperative interviewer: no information provided Quote: "No patient in any of the three groups reported pain during surgery or awareness of the surroundings"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information provided

Aime 2006

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: France

Anaesthetic interventions for prevention of awareness during surgery (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Aime 2006 (Continued)

Sex: both
 Age: 18 to 80
 ASA: I-III
 Procedure: Quote: "elective abdominal, gynaecologic, urologic, or orthopaedic surgery expected to last at least 1 h"
 Study size: 140 enrolled 125 results = 15 dropouts; 15/140 (~ 10%)

Interventions
Randomized portion of anaesthetic: volatile agent types: ADM: BIS vs SCPs

Intervention 1: sufentanil and sevoflurane - **standard clinical practices (SCPs)** N = 60 - 54 = 6 exclusions

Intervention 2: sufentanil and sevoflurane - **BIS 40 to 60** N = 40 - 34

Intervention 3: sufentanil and sevoflurane - **spectral entropy 40 to 60** N = 40 - 37 = 3 exclusions

Outcomes

Primary outcomes: sevoflurane consumption

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Quote: "No patient reported intraoperative recall"

Notes
Non-randomized portion of anaesthetic: parts of volatile agent: N₂O yes/narcotics/hypnotics bolus MCI/muscle relaxants induction yes/maintenance yes

Premedication: 100 mg hydroxyzine

Induction: propofol 2 mg to 3 mg/kg + sufentanil 0.2 µg to 0.3 µg/kg + atracurium 0.5 mg/kg

Maintenance: sevoflurane in 60% N₂O with oxygen 1 L/min + sufentanil 0.15 to 0.20 µg/kg/h and 5 µg bolus + atracurium 0.3 mg/kg/h + propofol 50 mg to 100 mg bolus if needed

Time of outcome determination: 1 and 3 days postoperatively

Method of determination: standardized interview

Quote: "The primary end point of this study was defined as the reduction in sevoflurane consumption"

Comment: this RCT was not powered for awareness - inadequate sample size

Author's survey sent on 26 November 2013 to: m.fischler@hopital-foch.org; second request sent 18 September 2014; author response on 19 September 2014: "Our methodology was that postop interviewer was blinded to intraop data and to randomization"

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "...were randomly allocated to one of three groups, the standard practice group, the BIS-guided group, or the spectral entropy-guided group, using a randomization list performed with computer generated random numbers."
Allocation concealment (selection bias)	Unclear risk	Comment: insufficient information provided
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "In the standard practice group, the screen monitor was customized to make BIS and Entropy values invisible to the attending anesthesiologist. In the BIS and in the spectral entropy guided groups, only the guiding parameter was displayed to the users"

Aime 2006 (Continued)

		Comment: anaesthesiologist not blinded to the randomized anaesthetics and both groups not blinded to SCPs even though both control groups blinded to ADM
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "...all patients were visited in the postanaesthesia care unit and on the first and third postoperative days and interviewed about intraoperative recall using a standardized interview..." Comment: author response on 19 September 2014: "Our methodology was that postop interviewer was blinded to intraoperative data and to randomization"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: patients who had clinical signs of intraoperative arousal were excluded. Awareness was a secondary outcome and the imbalance of excluded patients as they relate to the risk of awareness was inadvertently high risk in order to remove the confounding effect of propofol administration on sevoflurane consumption Comment: SCP: 3/60 high-risk awareness exclusions; ADM (BIS and Entropy): 2/80 high-risk awareness exclusions; imbalanced between groups but not significantly different Peto OR 0.49 (0.08 to 2.93); similar results with all types exclusions: 9/80 vs 6/60, 1.14 (0.39 to 3.35) Quote: "For dichotomous outcome data, the proportion of missing outcomes compared with observed event risk enough to induce clinically relevant bias in intervention effect estimate" (Higgins 2011)
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information provided

Akcali 2008

Methods	Study design: randomized parallel groups Study dates: Quote: "2001-2002" (ROB bias survey)
Participants	Country: Turkey Sex: both Age: 42 ASA: I-II Procedure: lumbar discectomy Study size: 56
Interventions	Randomized portion of anaesthetic: TIVA bolus ADM (BIS) vs SCP Intervention 1: remifentanyl bolus (1 µg/kg) + propofol (2 mg/kg) SCP), N = 28 Intervention 2: remifentanyl bolus + propofol titrated BIS 45 to 65, N = 28
Outcomes	Primary outcomes: effect of Bispectral Index (BIS) monitoring on haemodynamic parameters, drug consumption Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Akali 2008 (Continued)

Quote: "None of the patients remembered intubation, positioning and incision moments but haemodynamic reactions were recorded in some cases"

Notes

Non-randomized portion of anaesthetic: parts of TIVA/N₂O no/narcotics/hypnotics bolus MCI/ muscle relaxants induction yes/maintenance yes

Anaesthesia induction: remifentanil bolus (1 µg/kg) + propofol (2 mg/kg) or (1 mg/kg + 10 mg boluses to achieve BIS of 45 to 65) + pancuronium (0.1 mg/kg) + intubation, additional muscle relaxants given 1 to 2 twitch response to TOF

Anaesthesia maintenance: 33% O₂ and 67% **air** propofol infusion (4 mg/kg/h) + remifentanil (0.1 µg/kg/min)

Quote: "In order to avoid observer bias, the control group was studied first while a BIS monitor was attached to the patient but the screen was blinded to the anesthesiologist by covering by a card"

Time of outcome determination: first postoperative day

Method of outcome determination: interview

Survey response: 27 March 2011 from Didem Akcali didemakcali@yahoo.com

Email sent on 26 November 2013 to clarify randomization process

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The first 28 patients scheduled for surgery were randomised to control group. The last 28 patients were used as BIS group as first control group was studied in order to avoid observer bias." (Email bias survey, see notes) Quote: "These patients were randomly assigned to one of two study groups." (author's survey)
Allocation concealment (selection bias)	Unclear risk	Comment: no information
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "Patient" (author's survey) Quote: "first the control group was studied in order to avoid observer bias. The BIS monitor screen was blinded by a card and important instances were marked without seeing the monitor screen. Afterwards BIS group was studied." (author's survey)
Blinding of outcome assessment (detection bias) All outcomes	High risk	Quote: "Patient" (author's survey)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Quote: "The study protocol is available and all of the study's pre-specified (primary and secondary) outcomes that are of interest in the review have been reported in the pre-specified way." (Author's survey)
Other bias	Low risk	Quote: "there was no bias. The study was performed as planned without any source of bias." (Author's survey)

Anez 2001

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Spain Sex: unclear Age: group A 43.05 ± 15.2, group B 38.1 ± 14.7 ASA: I and II Procedure: vascular, venous, ortho outpatient Study size: 40, completed 39
Interventions	Randomized portion of anaesthetic: TIVA ADM (BIS) vs SCP Intervention 1: BIS blinded, N = 20 Intervention 2: BIS open (40 to 60), N = 20
Outcomes	Primary outcomes: effect of Bispectral Index (BIS) monitoring on propofol consumption, awakening, time to discharge, haemodynamic parameters, drug consumption Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 No awareness reports
Notes	Non-randomized portion of anaesthetic: parts of intravenous (IV) TIVA anaesthesia/N₂O no/narcotics and/or hypnotics bolus TCI/muscle relaxants induction yes/maintenance unclear Premedication: midazolam 0.03 µg/kg + atropine 0.01 µg/kg; anaesthesia induction: alfentanil 10 µg/kg, boluses 5-10 µg, rocuronium 0.2 mg/kg (smaller than usual intubating dose) TCI propofol initial dose 3.5 µg to 4 µg/mL/min, LMA; anaesthesia maintenance: propofol 2 µg/mL ambulatory surgery/propofol administered intravenously through a computerized system (Diprifuso®) and laryngeal mask Translated by Brett Smith; survey sent: canez@galenics.com

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	Quote: translation: "sequential randomisation: picked first 20 into one group and 2nd 20 into the second group"
Allocation concealment (selection bias)	Unclear risk	Comment: rule: no information
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: randomization was performed for BIS open or closed BIS blinded to SCP group but SCP open to both groups
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: anaesthesiologist and postoperative interviewer: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data

Anez 2001 (Continued)

Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information provided

Arellano 2000

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Canada Sex: female Age: 18 to 55 ASA: I and II Procedure: elective ambulatory termination of pregnancy or gynaecologic laparoscopy Study size: 1490, completed 1207, dropouts 283 (dropout 19%)
Interventions	Randomized portion of anaesthetic: parts of TIVA no N₂O Intervention 1: maintenance: propofol N ₂ O O ₂ 65% to 35% Intervention 2: maintenance: propofol alone TIVA O ₂ 100%
Outcomes	Primary outcomes: time to home readiness, postoperative recovery Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Quote: "We did not observe any cases of awareness in the TIVA...the incidence of intraoperative awareness is not greater than 0.4% inpatients undergoing outpatient gynaecologic surgery with our TIVA protocol...The aggregated incidence of awareness in the N ₂ O group was 0.13%...we did not encounter this unacceptably high incidence of recall as our protocol specified a starting propofol infusion rate of > 160 µg kg ⁻¹ min ⁻¹ with subsequent reductions if indicated clinically" Quote: "...one patient in this study reported intraoperative awareness (laparoscopy, N ₂ O group). The attending anaesthesiologist noted that this was likely caused by a kinked intravenous line that interrupted the flow of propofol for a short period" Comment: not powered for awareness
Notes	Non-randomized portion of anaesthetic: parts of IV and TIVA: N₂O yes and no/narcotics/hypnotics bolus MCI/muscle relaxants induction yes/maintenance yes TP anaesthesia induction: fentanyl 0.7 mg/kg + 20 mg lidocaine and 2.0 mg/kg propofol TP maintenance: propofol 20 mg boluses at signs of light anaesthesia Laparoscopy induction: fentanyl 1.5 mg/kg + d-tubocurarine 3 mg + 20 mg lidocaine + 2.0 mg/kg propofol with propofol titrated to loss of lid reflex + succinylcholine 1.5 mg/kg vecuronium 0.075 mg to 0.1 mg/kg Laparoscopy maintenance: propofol 100 µg to 200 µg/kg/min + propofol 20 mg boluses if needed Time of outcome determination: 24 h Method of outcome determination: interview "...these two surgical procedures represented the vast majority of ambulatory gynaecologic surgery performed at the institutions studied, patients undergo-

Arellano 2000 (Continued)

ing other ambulatory gynaecologic procedures were not studied to reduce heterogeneity in study population"

Email sent to arellano@is.dal.ca and arel111@yahoo.com on 26 November 2013

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Patients were randomly allocated by computer-generated random numbers in blocks of four..."
Allocation concealment (selection bias)	Low risk	Quote: "Patients were allocated to either the TIVA or N ₂ O group when the anaesthesiologist opened the sealed opaque envelopes at induction of anaesthesia"
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "... The anaesthesiologists were not blinded to treatment allocation to ensure safe anaesthetic care. Biased administration of the anaesthetics and un-blinding of the research assistants were prevented by the following...." Comment: based on the criteria for this domain we graded it high risk but would use discretion if RCT merged into meta-analysis
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Subjects were assessed by a blinded research assistant every 20 min until they reached it score 9. In addition, the same research assistant interviewed patients by telephone 24 hours after discharge ..."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: imbalanced but not significantly different between groups: N ₂ O (40/740) vs TIVA O ₂ (32/750), Peto OR 1.28 (0.80 to 2.06)
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Ashworth 1998

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: UK Sex: female and male Age: 18 to 70 ASA: I-II Procedure: ambulatory surgery Study size: 90
Interventions	Randomized portion of anaesthetic: parts of IV vs parts of volatile agent Intervention 1: maintenance: isoflurane ET 0.6%, then 0.25% to 1%; N = 30 Infusion of propofol, 160 µg/kg/min then 50 µg to 200 µg/kg/min; N = 30

Anaesthetic interventions for prevention of awareness during surgery (Review)

Ashworth 1998 (Continued)

Intervention 3: desflurane ET 3.6% then 1.4% to 6% PRN; N = 30

Outcomes	<p>Primary outcomes: postoperative recovery; respiratory complications; heart rate (h) (A) and mean arterial blood pressure; episodes of purposeful movement</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4</p> <p>Quote: "The IV anaesthetic propofol was associated with more patient movement (without awareness) during surgery"</p> <p>Comment: no awareness events</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of IV vs parts of volatile agent N₂O yes/narcotics/hypnotics bolus/muscle relaxants induction no/maintenance no</p> <p>Induction: a single anaesthesiologist, fentanyl 1 µg/kg/lidocaine 20 mg and propofol 20 mg to 40 mg every 10 S, LMA was inserted after adequate jaw relaxation. No muscle relaxant used, N₂O 67%/O₂ 6 L/min for 10 min then 2 L/min</p> <p>Quote: "In spontaneously breathing patients, inadequate anaesthesia is manifested by purposeful movements, which allows corrective action to be taken promptly. As a result, awareness is unlikely, and none of our patients recalled intraoperative events. When muscle relaxants and controlled ventilation are used, however, awareness is a greater risk because inadequate anesthesia can occur without obvious changes in clinical signs"</p> <p>No email found</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "An open-label, prospective study design was used involving random allocation to one of three parallel groups for the maintenance of anesthesia (isoflurane, propofol, or desflurane) with blind assessment of postoperative recovery"
Allocation concealment (selection bias)	High risk	Quote: "Participants or investigators enrolling participants could possibly foresee assignments and thus introduce selection bias, such as allocation based on: Using an open random allocation schedule (e.g. a list of random numbers)" (Higgins 2011)
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: see above
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "No patient subsequently had recall of any intraoperative event on direct postoperative questioning"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information

Avidan 2008

Methods	Study design: randomized parallel groups Study dates: September 2005 to October 2006
Participants	Country: USA Sex: both Age: mean age 59.5 to 59.2 at high risk of awareness ASA: I-IV Procedure: not identified Study size: 2000 entered study and 1941 completed possible interviews
Interventions	Randomized portion of anaesthetic: volatile agent types ADM BIS vs ETAG Intervention 1: BIS guided anaesthesia (target BIS range, 40 to 60), N = 1000 Intervention 2: ETAG guided anaesthesia (target ETAG 0.7 to 1.3 MAC), enrolled 2000, completed study 1941
Outcomes	Primary outcomes: awareness/wakefulness as defined using our "Classification of intra-operative cognitive states" Table 1: class 4 Comment: 16 awareness reports by patients = 4 definite + 5 possible + 7 no awareness: expert panel classification. 12/16 patient awareness reports NOT used in calculation of awareness rate (definite awareness) Definite and possible BIS 10/967 vs SCP 6/974 Definite: BIS 2/967 vs SCP 2/974
Notes	Non-randomized portion of anaesthetic: parts of volatile agent: N₂O unclear/narcotics/hypnotics/muscle relaxants induction unclear/maintenance unclear Anaesthesia induction: left to discretion of anaesthetist Anaesthesia maintenance: left to discretion of anaesthetist Comment: high risk of anaesthesia awareness/isoflurane, sevoflurane or desflurane/supplemental N ₂ O permitted Time of outcome determination: within 24 h, 1 to 3 days, and 72 days postoperative period Survey response: 27 September 2011 Michael Avidan, avidanm@anest.wustl.edu Email sent to author to clarify missing data issues: 7 December 2013: Michael Avidan, avidanm@anest.wustl.edu . Dr Avidan responded. Senior authors developed consensus on ROB domain emailed to authors and response acknowledging our assessment sent/received 28 December 2013. Email sent 21 October 2014 adjudicated awareness reports and response 21 October 2014; 14 November 2014 adjudicated awareness reports for review copies of reports not published in RCT. Dr Avidan responded 14 November 2014
Risk of bias	
Bias	Authors' judgement Support for judgement
Random sequence generation (selection bias)	Low risk Quote: "The design was a single-center, prospective study, in which 2000 patients underwent pre-randomization electronically in blocks of 100, with 50

Avidan 2008 (Continued)

		patients assigned to a BIS-guided protocol and 50 to an ETAG-guided protocol"
		Quote from survey: "Computer generated randomization. Block randomized in blocks of 100 (50 in each group per block). Unique case identifier and information on group allocation (BIS or ETAG) was sealed in opaque envelopes"
Allocation concealment (selection bias)	Low risk	Quote from survey: "The opaque envelope was opened by a research coordinator only after the patient (who had already signed consent) was transferred to the operating room"
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "The anesthesiologists were not blinded to treatment allocation to ensure safe anaesthetic care" Quote from paper: "formed consent. The anesthesia practitioners were aware of the assignments of the patients , but the patients, the postoperative interviewers, the expert reviewers, and the statistician were not"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patient, Awareness outcome assessor (interviewer), Statistician, expert reviewers who determined whether awareness was definite, possible or not awareness"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: 2000 patients enrolled, in table 1, trial enrolment, there were 33 BIS and 26 ETAG patients excluded after randomization. 1905/1941 (98%) completed 1 to 3 awareness interviews and 36/1941 (2%) completed no awareness interviews Senior authors' consensus: ... because of the balanced event rates and sample sizes, even though the reasons for a portion of the dropouts are not known, there would be little impact on the reported outcome from the study if these patients had been interviewed for awareness 33/2000 vs 26/2000, Peto OR 1.27 (0.76 to 2.13) Email sent to author to clarify these issues: 7 December 2013: Michael Avidan, avidanm@anest.wustl.edu
Selective reporting (reporting bias)	Low risk	Quote: "The study protocol was not published in the public domain. The trial was registered on clinical trials.com (NCT00281489)"
Other bias	Low risk	Quote: "We know of no bias related to the study design"

Avidan 2011

Methods	Study design: randomized parallel groups Study dates: May 2008 through May 2010
Participants	Country: USA Sex: both Age: mean 60 ± 14.2, 61 ± 14.4; high risk of awareness ASA: I-IV (> 80% patients ASA III-IV) Procedure: elective surgery

Avidan 2011 (Continued)

Study size: 6041 enrolled, 49,000 screened, 5801 included in trial, 5713 completed >= 1 interview (primary outcome analysis) (98.3%)

Interventions	<p>Randomized portion of anaesthetic: volatile agent types ADM BIS vs ETAG</p> <p>Intervention 1: BIS guided anaesthesia (target BIS range, 40 to 60), N = 2861</p> <p>Intervention 2: ETAG guided anaesthesia (target ETAG 0.7 to 1.3 MAC), N = 2852</p>
Outcomes	<p>Primary outcomes: awareness/wakefulness as defined using our "Classification of intra-operative cognitive states" Table 1: class 4</p> <p>Comment: see discussion section on classification</p> <p>Quote: "...With some awareness events apparently occurring with BIS values below 60, decreasing anesthetic concentration solely on the basis of a BIS value of less than 60 is not recommended...rare event such as awareness, unidentified risk factors such as genetic resistance to anesthetic agents could have been unequally distributed between the BIS and ETAC groups despite randomization and could have confounded the results"</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent: N₂O unclear/narcotics/hypnotics/muscle relaxants induction unclear/maintenance unclear</p> <p>BIS or Anaesthetic Gas to Reduce Explicit Recall (BAG-RECALL)</p> <p>General anaesthesia with isoflurane, sevoflurane or desflurane</p> <p>Patients at high risk were defined as those with at least one risk factor (Table 1)</p> <p>Anaesthesia induction: left to discretion of anaesthetist</p> <p>Anaesthesia maintenance: left to discretion of anaesthetist</p> <p>Time of outcome determination: modified Brice questionnaire, within 72 h after surgery and at 30 days after extubation</p> <p>Method of outcome determination: interview</p> <p>Survey response: 27 September 2011 Michael Avidan, avidanm@anest.wustl.edu; see Avidan 2008 for subsequent email communications; email sent 21 October 2014 adjudicated awareness reports and response 21 October 2014; 14 November 2014 adjudicated awareness reports for review copies of reports not published in RCT. Responded 14 November 2014; emails January to February 2015: adjudicated awareness reports</p> <p>A delayed report of an awareness event was published in 2013 (Villafranca 2013)</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	<p>Quote: "6100 pre-randomization designations were generated electronically in blocks of 100, divided equally between the groups"</p> <p>Quote from survey: "Computer generated randomization. Block randomized in blocks of 100 (50 in each group per block), randomization was generated at the coordinating site. Unique case identifier and information on group allocation (BIS or ETAG) was sealed in opaque envelopes"</p>
Allocation concealment (selection bias)	Low risk	Quote: "Labels indicating BIS group or ETAC group were sealed in opaque, numbered envelopes"

Avidan 2011 (Continued)

		Quote from survey: "The opaque envelope was opened by a research coordinator only after the patient (who had already signed consent) was transferred to the operating room"
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "Patients in the ETAC group had monitors configured to conceal the BIS number, so the anesthesia practitioners were unaware of the BIS values. The practitioners in both groups could view the ETAC. Summaries of the BIS and ETAC protocols" Comment: see this domain's comments in Avidan 2008 Comment: anaesthesia practitioners ETAG group using monitor configuration omits BIS number, so unaware of BIS values; both groups view ETAG concentrations
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"Patient, Awareness outcome assessor (interviewer), Statistician, expert reviewers who determined whether awareness was definite, possible"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: see Avidan 2008 (46/2907) vs (50/2902), Peto OR 0.92 (0.61 to 1.37) Email sent to author to clarify this issue: 25 November 2013: Michael Avidan, avidanm@anest.wustl.edu
Selective reporting (reporting bias)	Low risk	Quote from survey: "The study protocol is available and was published in the journal BMC Anesthesiology (http://www.ncbi.nlm.nih.gov/pubmed/19948045), and the study's pre-specified (primary and secondary) outcomes that are of interest in the review have been reported in the pre-specified way. The distressing experience of awareness was reported as a post hoc analysis"
Other bias	Low risk	Quote from survey: "We know of no important source of bias related to study design"

Baraka 1989

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Lebanon Sex: female Age: mean age 31 ASA: not available, "healthy" Procedure: caesarean section Study size: 50
Interventions	Randomized portion of anaesthetic: parts of intravenous (IV) vs parts of volatile agent/N₂O ± Intervention 1: induction: thiopentone 4 mg/kg; maintenance: N ₂ O 50% + halothane 0.5% in O ₂ , N = 10 Intervention 2: induction: thiopentone 4 mg/kg; maintenance: halothane 1% in O ₂ , N = 10

Baraka 1989 (Continued)

Intervention 3: induction: ketamine 1.5 mg/kg; maintenance: 50% N₂O + 0.5% **halothane** in O₂, N = 10

Intervention 4: induction: ketamine 1.5 mg/kg; maintenance: 1% **halothane** in O₂, N = 10

Intervention 5: induction: **ketamine** 1.5 mg/kg; maintenance: 100% O₂, N = 10

Outcomes

Primary outcomes: APGAR, blood loss maternal artery PO₂ (kPa), PCO₂ (kPa), umbilical vein PO₂ (kPa), PCO₂ (kPa)

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4

Quote: "Awareness following induction of anaesthesia was detected using isolated forearm technique"

Comment: in this review this is referred to as "wakefulness"; "recall" is used by the authors as patient memory of postoperative events (awareness)

Table 1 p646: "awareness" (wakefulness) 18 (14 in thiopentone and 4 in ketamine groups) patients, "recall" (awareness) 2 patients, intraoperative dreams 1, intraoperative hallucination 1 (classified as dream; both dreams in ketamine group)

"Two of the 20 thiopentone patients had recall, confined to the time of delivery"

Notes

Non-randomized portion of anaesthetic: parts of intravenous (IV) vs parts of volatile agent/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance yes

Anaesthetic induction: suxamethonium 1.5 mg/kg

After delivery - anaesthetic maintenance: 66 N₂O in O₂ + fentanyl 3 µg/kg/alcuronium 0.25 mg/kg

Time of outcome determination: following recovery and the next day

Method of outcome determination: interview

"Awareness following induction of anaesthesia was detected using the isolated forearm technique...Following tracheal intubation, the patient's right hand was clasped by the anaesthetist and the patient was asked to squeeze and relax her hand three times successively... Because hand movement may occur spontaneously or as a response to surgical stimulation, the test was considered positive only if the patient squeezed and relaxed her hand according to instructions"

ROB survey see [Baraka 1998](#): the Cochrane Anaesthesia, Critical and Emergency Care Group's editorial team prefers not to extrapolate to previously published RCTs in assigning ROB risk

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The 50 patients were allocated randomly to five equal groups"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: insufficient information provided

Baraka 1989 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no dropout
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Baraka 1998

Methods	Study design: randomized parallel groups Study dates: "January 1997 till October 1997" (email bias survey, see notes)
Participants	Country: Lebanon Sex: female Age: mean 30 ASA: I-II Procedure: elective caesarean section Study size: 40
Interventions	Randomized portion of anaesthetic: parts of TIVA: narcotic vs narcotic Intervention 1: induction with 100 mg tramadol + tramadol 50 mg as needed, N = 20 Intervention 2: induction with 100 mg fentanyl + fentanyl 50 µg, N = 20
Outcomes	Primary outcomes: similar to Baraka 1989 Secondary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Quote: "No patient in the fentanyl group had any intraoperative recall, while two patients in the tramadol group recalled crying of their newborns at the delivery time; no pain was experienced at that time"
Notes	Non-randomized portion of anaesthetic: parts of TIVA: N₂O yes/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance yes Anaesthesia induction: 100 mg tramadol or 100 mg fentanyl + thiopental 3 mg/kg + succinylcholine 1.5 mg/kg Anaesthesia maintenance: N ₂ O 50% + vecuronium 0.1 mg/kg; post partum: N ₂ O 50% Time of outcome determination: following recovery and day after operation Method of outcome determination: interview Survey response: 10 March 2011 Boutros Assaf, nbassaf@cyberia.net.lb

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Baraka 1998 (Continued)

Random sequence generation (selection bias)	Low risk	Quote: "paper draw" (email bias survey) Quote: "Patients were randomly allocated into two groups"
Allocation concealment (selection bias)	High risk	"The anesthesiologist was not blinded" (email bias survey)
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: see below
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "the anaesthesiologist was not blinded...Patient, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	"study protocol not available" (email bias survey) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote: "none"

Barr 1977

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: UK Sex: female Age: 28 ASA: not available Procedure: caesarean section Study size: 220 enrolled, 37 dropouts, 183 completed study
Interventions	Randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia: maintenance: post-delivery: hypnotics Intervention 1: post-delivery: lorazepam 2 mg, N = 110 enrolled; 93 results Intervention 2: post-delivery: diazepam 10 mg, N = 110; 90 results
Outcomes	Primary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Quote: "A total of seven patients had pleasant dreams, seven had unpleasant dreams and three had actual recall"
Notes	Non-randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia (Liverpool technique): N₂O yes/narcotics/hypnotics bolus/NMBs induction yes/maintenance yes

Anaesthetic interventions for prevention of awareness during surgery (Review)

Barr 1977 (Continued)

Anaesthesia induction: sodium **thiopentone** to max of 300 mg or 4 mg/kg + suxamethonium or **pancuronium**; anaesthesia maintenance: N₂O in oxygen 2:1. If signs of lightness **PRN**: incremental doses **pethidine** 25 mg max dose (**Liverpool technique**). Post-delivery medication: ergometrine 0.5 mg + pethidine 0.5 mg + study drug

Comment: see discussion of **retrograde amnesia**

Quote: "unpleasant dreams indicate inadequate depth of anaesthesia"

Time of outcome determination: second or third postoperative day

Method of outcome determination: interview

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The trial ampoules...were distributed from a table of random numbers"
Allocation concealment (selection bias)	Unclear risk	Comment: as above
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Immediately following delivery the anaesthetist injected ergometrine 0.5 mg, pethidine 25 mg, and the contents of a trial ampoule into the intravenous infusion. The trial ampoules contained lorazepam 2 mg in 2 ml, or diazepam 10 mg in 2 ml; these were distributed from a table of random numbers"
Blinding of outcome assessment (detection bias) All outcomes	High risk	"Postoperative interview. The patient was interviewed on the second or third postoperative day by one of the authors. The questions asked are listed in Table 1"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: 17% dropouts; lorazepam 93 (110 - 93 = 17 exclusions, 110 - 90 = 20 exclusions) and diazepam 90 completed study; Peto OR 0.82 (0.41 to 1.67) Quote: "Thirty-seven of the 220 patients originally admitted to the trial were later excluded because of deviation from the protocol, incomplete records, broken ampoules, or lost forms; as far as the authors are able to ascertain none of these patients was aware during the procedure"
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Bauer 2004

Methods	Study design: randomized parallel groups Study dates: "December 2000 - February 2002" (email bias survey, see notes)
Participants	Country: Germany Sex: both Age: 60 ASA: not available Procedure: elective coronary artery bypass grafting

Bauer 2004 (Continued)

Study size: 40

Interventions	<p>Randomized portion of anaesthetic: parts of TIVA TCI vs MCI and ADM BIS 40 to 50</p> <p>Intervention 1 (TCI group): background infusion of remifentanyl (0.3 µg/kg/min) and propofol at a target concentration of 3 µg/mL using target-controlled infusion (TCI, Diprifusor), N = 20</p> <p>Intervention 2 (BIS group): general anaesthesia background infusion of remifentanyl (0.3 µg/kg/min) and propofol titrated to maintain a BIS value of 40 to 50, N = 20</p>
Outcomes	<p>Primary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Quote: "None of the patients reported awareness during a standardized interview"</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of TIVA: N₂O no/narcotics/hypnotics bolus MCI TCI/ muscle relaxants induction yes/maintenance yes/ADM no</p> <p>Anaesthesia induction: remifentanyl infusion (0.1 µg/kg/min) + propofol at target concentration of (3 µg/mL) + atracurium plus additional boluses: ensure relaxation throughout surgery</p> <p>Anaesthesia maintenance: remifentanyl infusion (0.3 µg/kg/min) + propofol titrated in group specific manner (see above) + atracurium boluses as needed; controlled ventilation; BIS group was titrated to BIS < 60</p> <p>Time of outcome determination: first and third postoperative day</p> <p>Quote: "Total intravenous anesthesia using propofol- remifentanyl effectively attenuates the neurohumoral stress"</p> <p>Method of outcome determination: standardized interview</p> <p>Survey response: 2 February 2011, Michael Bauer michael.bauer@med.uni-jena.de</p> <p>Re-emailed 26 December 2013 re allocation</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "computer-generated randomisation list" (email survey)
Allocation concealment (selection bias)	Unclear risk	Quote: "sealed envelopes; opened immediately prior to induction of anaesthesia" (email bias survey) Comment: unclear if envelope opaque as per Cochrane guidelines (Higgins 2011)
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "patient" (email survey)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "None of the patients reported intraoperative recall during the standardised interview on postoperative days 1 and 3"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "data complete" (email survey)

Bauer 2004 (Continued)

Selective reporting (reporting bias)	Low risk	Quote: "Study protocols are not available but report includes all expected outcomes" (email survey) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote: "none" (email survey)

Bergmann 2013

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Germany Sex: male/female Age: 18 to 75: mean 44 to 48 ASA: ASA I and II (90%) to III (10%) Procedure: orthopaedic: shoulder/knee/ankle Study size: 170 entered; 151 completed study
Interventions	Randomized portion of anaesthetic: parts of TIVA ADM (SPI (plethysmograph)) vs SCP "The propofol dose was adjusted according to entropy in both groups" Intervention 1 (SPI group): remifentanyl adjusted surgical pleth index (SPI), SPI: 20 - 50, (N = 76) Intervention 2 (control group): remifentanyl adjusted clinical parameters (SCPs), (N = 75) Propofol dose was adjusted SE entropy in an identical manner in both groups
Outcomes	The primary endpoints of the study were differences between the groups in the recovery times and the consumption of anaesthetic drugs Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Secondary endpoints were the occurrence of complications such as intraoperative awareness, nausea and vomiting, postoperative pain, patient satisfaction with the anaesthesia, shivering, and haemodynamic stability "No patient reported intraoperative awareness"
Notes	Non-randomized portion of anaesthetic: parts of TIVA: N₂O no/narcotics/hypnotics bolus MCI/ NMBs induction PRN/maintenance PRN SR and CV ADM (Entropy) Quote: "surgical pleth index (SPI) is an index based on changes in plethysmographic characteristics that correlate with the balance between the sympathetic and parasympathetic nervous system" All patients: IV midazolam (1 mg to 3 mg), total IV technique remifentanyl/propofol: bolus remifentanyl (1 mg/kg), infusion 0.2 mg/kg/min, propofol 2 mg/kg/min until entropy SE < 60 ; propofol 4 mg/kg/h, than adjusted entropy target range between 40 to 60 ; infusion reduced 10 min before end of the operation SE value allowed increase 60 and 65. The airway was secured whenever possible with a laryngeal mask airway for lower limb surgery, and with a tracheal tube for shoulder operations; mivacurium (0.2 mg/kg) for intubation; AP was lowered with titrated 5 mg IV doses urapidil Quote: "SPI is a dimensionless number between 0 (low stress) and 100 (high stress) that is calculated from the heart rate (HR) and the pulse wave amplitude obtained with a finger clip...SPI = 100 - (0.33 × HBI _{norm} + 0.67 × PPG _{norm})...The precise description and calculation of the algorithm is provided elsewhere.22"

Bergmann 2013 (Continued)

 ROB survey: we emailed on 21 March 2015, ingo.bergmann@med.uni-goettingen.de
Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "After randomization using a computer-generated list, the patients were assigned to one of the two study groups"
Allocation concealment (selection bias)	Unclear risk	Comment: no information
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "In the SPI group, the remifentanyl dose was adjusted according to the SPI, while in the control group, it was administered according to standard clinical criteria. The propofol dose was adjusted according to SE entropy in an identical manner in both groups"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "Telephone interviews were conducted on the evening after the operation, and on the first and second postoperative days"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "One hundred and fifty-one of the 170 recruited patients were included in the final analysis. Nineteen patients were excluded due to incomplete data sets or retraction of consent. The anthropometric data are shown in Table 1. Seventy-six patients were in the group with SPI-guided remifentanyl administration and 75 patients in the control group" Comment: no high-risk dropouts; 170 recruited, all exclusions no significant difference between groups: SCP (10/85) vs SPI (9/85), 1.13 (0.43 to 2.91)
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information

Bestas 2004

Methods	Study design: randomized parallel groups Study dates: "2003" (email survey)
Participants	Country: Turkey Sex: both Age: 59.6 ASA: not available Procedure: off-pump CABG surgery Study size: 50
Interventions	Randomized portion of anaesthetic: IV bolus MCI ADM (BIS) vs SCP Intervention 1: induction: fentanyl (7 µg to 15 µg/kg) + propofol (2 mg to 2.5 mg/kg); maintenance: fentanyl (15 µg/kg/h) + propofol (6 mg/kg/h), N = 25 Intervention 2: induction: fentanyl (7 µg to 15 µg/kg) + midazolam (0.1 mg to 0.3 mg/kg); maintenance: fentanyl (15 µg/kg/h) + midazolam 0.1 mg/kg/h, N = 25

Bestas 2004 (Continued)

Comment: depth of anaesthesia was based on clinical signs. BIS values were not shown

Outcomes	<p>Primary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>"No patients were noted to recall the sounds presented during the operation and the preoperative events...no patients reported to have heard anything or had any dreams intraoperatively"</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of IV:/N₂O no/narcotics/hypnotics bolus MCI/muscle relaxants induction yes/maintenance yes</p> <p>Anaesthesia induction: fentanyl (7 µg to 15 µg/kg) + group-specific study drug (see above) + vecuronium (0.1 mg/kg). Anaesthesia maintenance: fentanyl (15 µg/kg/h) + group-specific study drug (see above) + vecuronium 0.02 mg/kg as needed</p> <p>Time of outcome determination: 24 h after surgery</p> <p>Method of outcome determination: standardized questionnaire</p> <p>Author responded to email survey</p> <p>Survey response: 24 January 2011 Azize Bestas, abestas@firat.edu.tr</p> <p>Email sent to clarify allocation on 26 November 2013 to Azize Bestas, abestas@firat.edu.tr</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "randomly divided into two groups" Quote (email bias survey): "A random numbers table was used"
Allocation concealment (selection bias)	High risk	Quote (email bias survey): "Anesthesia was administered to patients by an anaesthesiologist not blinded to the study allocation (blinded for BIS values)"
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "Patient, Anesthesiologist, Awareness outcome assessor (interviewer)" Comment: although many authors define ADM vs SCP as a blinded method, SCP is exposed to both groups
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patient, Anesthesiologist, Awareness outcome assessor (interviewer)"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no dropouts
Selective reporting (reporting bias)	Low risk	Quote: "There is no any important risk of bias." (Email bias survey, see notes) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote: "There is no any important risk of bias." (Email bias survey, see notes)

Bethune 1992

Methods	Study design: randomized parallel groups
---------	--

Bethune 1992 (Continued)

Study dates: not stated

Participants	Country: England Sex: both Age: 60.6 ASA: not available Procedure: coronary artery surgery Study size: 44 enrolled 43 completed study
Interventions	Randomized portion of anaesthetic: parts of IV Intervention 1: maintenance: propofol 2 mg/kg/h, N = 22 Intervention 2: maintenance: methohexitone 1.5 mg/kg/h, N = 22
Outcomes	Primary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 2 "No patient had explicit recall of any events during the period when the tape was played. The patients in the propofol group who heard the tape during surgery had significant implicit recall of the word associations..."
Notes	Non-randomized portion of anaesthetic: parts of IV (opioid-based anaesthesia)/N₂O no/narcotics/hypnotics bolus MCI/muscle relaxants induction yes/maintenance yes Anaesthesia induction: midazolam (2 mg) + fentanyl (15 µg/kg) + pancuronium (0.14 mg/kg) Anaesthesia maintenance: drug infusion; atracurium 50 mg and droperidol 5 mg were administered on initiation of cardiopulmonary bypass (CPB) Time of outcome determination: between 36 and 48 h after surgery Method of outcome determination: interview Comment: this implicit memory was lost when re-tested in ICU with sedation No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "...were allocated randomly..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: pre- and postoperative interviewer was blinded to assignment - implicit study tape
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "Data from one patient in the propofol group were mislaid and had been omitted from the analysis"

Bethune 1992 (Continued)

Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Bhawna 2012

Methods	Study design: randomized parallel groups Study dates: January 2007 to December 2008
Participants	Country: India Sex: male and female Age: 25 and 65 years of age ASA: I-II Procedure: lower abdominal surgeries Study size: 50
Interventions	Randomized portion of anaesthetic: cardiac drug vs placebo Intervention 1: (N = 25) esmolol infusion Intervention 2: (N = 25) saline infusion Entropy: 40 to 60 Quote: "...loading dose of randomly selected study drug infusion (0.5 mg/kg) over 5 min, 20 min before induction followed by a continuous infusion of the study drug at 0.5 mg/kg/min till the closure of skin incision"
Outcomes	Primary outcomes: "...absolute isoflurane requirement between the two groups to keep the entropy values between 40 and 60 during the peri-operative period" Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "No incidence of intra-operative awareness was reported by any patient from either of the groups"
Notes	Non-randomized portion of anaesthetic: parts of volatile agent: N₂O no/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance PRN/ADM: entropy Induction: fentanyl (3.0 µg/kg)/propofol (1.25 mg to 2.0 mg/kg) titrated entropy 40 to 60/atracurium (0.5 mg/kg) then PRN/oxygen (FIO ₂) at 0.4/isoflurane/target RE/SE entropy sukhminder_bajwa2001@yahoo.com 5 February 2015 response

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote from ROB survey response: "Simple randomization"
Allocation concealment (selection bias)	Low risk	Quote from ROB survey response: "By use of sequentially numbered opaque sealed envelopes (SNOSE)"
Blinding of participants and personnel (performance bias)	Low risk	Quote from ROB survey response: "Patient, Anesthesiologist, Awareness outcome assessor (interviewer)"

Anaesthetic interventions for prevention of awareness during surgery (Review)

Bhawna 2012 (Continued)

All outcomes		Comment: infusions used same rate
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote from ROB survey response: as above
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote from ROB survey response: "No such attrition/exclusions were reported in the study"
Selective reporting (reporting bias)	Low risk	Quote from ROB survey response: "All specified outcomes have been well addressed in the study"; "No significant difference in MAC was observed at intervals other than 30 min, 40 min and 105 min intervals" Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote from ROB survey response: "Study did not have any bias potential as the design of the study was conceived after thorough discussion among all researchers"

Blendinger 1976

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Germany Sex: both Age: not stated ASA: not stated Procedure: open heart surgery Study size: 15 in test group; control group not stated, assume 15 as per translator
Interventions	Randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia MCI Intervention 1: control group no etomidate : average fentanyl dose 30 µg/kg/min Intervention 2: experimental group average etomidate dose 2.55 mg/kg ; average fentanyl dose 12 µg/kg/min
Outcomes	Primary outcomes: anaesthesia consumption, postoperative recovery parameters Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: none of the subjects reported any memories
Notes	Non-randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia: N₂O yes/narcotics and/or hypnotics bolus MCI/muscle relaxants induction yes/maintenance yes Muscle relaxation, intubation; maintenance: controlled respiration/N ₂ O (2:1) fentanyl was intermittently injected Comment: high-dose fentanyl anaesthesia with and without etomidate supplementation Comment: details of anaesthetic unclear to translator; translator Lore Schultheiss No email address available for ROB survey

Blending 1976 (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: unclear to translator
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no apparent dropout
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Block 1991

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: USA Sex: female Age: mean 34-38-35 (by group) ASA: I-II-III Procedure: operations on the fallopian tubes for infertility, vertical banding gastroplasty for morbid obesity, total abdominal hysterectomy, cholecystectomy, other small groups Study size: 72
Interventions	Randomized portion of anaesthetic: parts of IV + N₂O vs parts of volatile agent Intervention 1: inhalation group: Isoflurane ET 1.0 MAC/70% N ₂ O/O ₂ , N = 12 Isoflurane ET 1.3 MAC/70% N ₂ O/O ₂ , N = 24 Isoflurane ET 1.5 MAC/70% N ₂ O/O ₂ , N = 12 Bolus PRN: 0.75 µg/kg fentanyl/equivalent sufentanil: systolic BP 15% baseline Intervention 2; 70% N ₂ O/O ₂ 2 doses of opioids, N = 24 Induction: fentanyl 7.5 µg induction/equivalent opioids, thiopentone

Block 1991 (Continued)

 Maintenance: 70% N₂O/O₂

Outcomes	Primary/secondary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: no implicit/explicit memory Quote: "determine if learning occurs during general anaesthesia"
Notes	<p>Non-randomized portion of anaesthetic: parts of IV + N₂O vs parts of volatile agent N₂O yes/narcotics/hypnotics bolus MCI/muscle relaxants induction yes/maintenance unclear</p> Thiopentone 3.0 mg/kg Bolus fentanyl 2.5 µg/kg for systolic BP > 15% baseline or patient moved Induction/maintenance: alfentanil 50 µg/kg (equivalent dose other opioids) then 1 µg to 1.5 µg/kg/min; 70% N ₂ O/O ₂ , infusion: assumed sufentanil 1 µg, fentanyl 5 µg, and alfentanil 25 µg to be equivalent doses No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Patients were allocated blindly and randomly to one of two anesthesia methods..."
Allocation concealment (selection bias)	Unclear risk	Comment: see above
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: insufficient information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Bonato 2001

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Italy

Bonato 2001 (Continued)

Sex: female and male
Age: mean age 64 and 63 years (range 28 to 75)
ASA: cardiac surgery: II-IV
Procedure: cardiac surgery
Study size: 93

Interventions

Randomized portion of anaesthetic: ADM BIS 60 vs 40: implicit memory test

Control group: N = 20
73 subjects (experimental group)
Group A: N = 35
Group B: N = 38
Intervention 1: BIS 60 (word list played)
Intervention 2: BIS 40

Outcomes

Primary/secondary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 2
Quote: "significant intraoperative implicit memory was found ($P < 0.005$), but no patient had spontaneous or directed recall of intraoperative events"

Notes

Non-randomized portion of anaesthetic: parts of TIVA: N₂O no/narcotics/hypnotics bolus MCI/muscle relaxants induction yes/maintenance yes
Remifentanyl-propofol-vecuronium-vecuronium, CPB
The mean dose of propofol group A was 1.9 mg/kg/h, group B 2.3 mg/kg/h; remifentanyl mean dose required in the 2 groups is the same: 0.45 mg/kg/min
Translated by AM
No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "...prospective, randomised study"
Allocation concealment (selection bias)	Unclear risk	Comment: inadequate information
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: anaesthesiologists targeted anaesthesia to 2 different endpoints, BIS 60 and 40
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "a standardised interview was conducted to determine the extent of intraoperative explicit memory and a Word Stem"

Bonato 2001 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing data
Selective reporting (reporting bias)	Low risk	Comment: primary outcome: implicit memory; secondary outcome: awareness Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information

Browne 1973

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Canada Sex: female Age: 20 to 76 ASA: I-II Procedure: gynaecological operations Study size: 112
Interventions	Randomized portion of anaesthetic: parts of intravenous (IV) (neurolapt) anaesthesia/induction Intervention 1: induction: thiopentone (250 mg to 500 mg), N = 56 Intervention 2: induction: innovar 1 mL to 2 mL (fentanyl-droperidol) + thiopentone 250 to 500 mg, Innovar (0.5 mL to 1 mL), PRN signs light anaesthesia, N = 56
Outcomes	Primary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Quote: "Recall of music appeared to be definite in two patients in Group I" Quote: "This study is concerned with the assessment of the patient's awareness for events occurring during and after surgery, when using two types of light anaesthesia"
Notes	Non-randomized portion of anaesthetic: parts of intravenous (IV) (neurolapt) anaesthesia: N₂O yes/narcotics and/or hypnotics bolus muscle relaxants induction yes/maintenance PRN Maintenance: 60% N ₂ O/curare or pancuronium PRN; patients were exposed to a fixed auditory stimulus (music) Comment: Innovar had less movement; see Dryad Time of outcome determination: within 48 h Method of outcome determination: interview No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The patients were then divided at random into two groups"

Browne 1973 (Continued)

Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: insufficient information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Bruhn 2005

Methods	Study design: randomized parallel groups Study dates: Quote: "2001-2003" (email bias survey)
Participants	Country: Germany Sex: both Age: 18 to 80 ASA: I-III Procedure: minor surgery > 1 h Study size: 200
Interventions	Randomized portion of anaesthetic: volatile agent types ADM (BIS) vs SCP Intervention 1: BIS controlled desflurane-remifentanyl, N = 71 Intervention 2: AAI controlled desflurane-remifentanyl, N = 58 Intervention 3: clinical parameters controlled desflurane-remifentanyl; maintenance: desflurane increase 0.5% if needed, N = 71
Outcomes	Primary outcomes: recovery times and drug consumption: BIS vs SCP Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Quote: "No patient complained of intraoperative recall" 20 patients remembered dreaming during surgery
Notes	Non-randomized portion of anaesthetic: parts of volatile agent: N₂O no/narcotics/hypnotics bolus MCI/muscle relaxants induction yes/maintenance no/ADM BIS or AAI

Bruhn 2005 (Continued)

Premedication: midazolam 7.5 mg

Induction: remifentanil 0.4 µg/kg/min + propofol 2 mg/kg + cis-atracurium 0.1 mg/kg + remifentanil increased 0.05 µg/kg/min if needed

Maintenance: remifentanil reduced to 0.2 µg/kg/min + desflurane 3% ET in O₂ and air. No more neuromuscular blocking agents were given intraoperatively

Time of outcome determination: 1 and 3 days postoperatively

Method of determination: interview

Primary author responded to email bias survey 31 March 2011. See bias table below for responses

Survey response: 31 March 2011 J. Bruhn [j.bruhn@anes.umcn.nl](mailto:bruhn@anes.umcn.nl)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The patients were randomised by drawing lots from a closed box." (Email bias survey)
Allocation concealment (selection bias)	High risk	Quote: "Anaesthesiologist: Due to the study design, the anaesthesiologist had to know to what group a patient was randomised." (Email bias survey)
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "Patient" (email bias survey, see notes)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "In post-operative period, investigators interviewed patients about awareness with an informal interview." (Email bias survey)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "The outcome data were complete, there were no exclusions from the analysis." (Email bias survey)
Selective reporting (reporting bias)	Low risk	Quote: "...non-significant results from the study, i.e. no differences between the groups were reported." (Email bias survey) Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information (Email bias survey)

Casati 1999

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Italy Sex: "healthy women" Age: 18 to 70 ASA: I-II

Casati 1999 (Continued)

Procedure: ultrasound guided oocyte retrieval for in vitro fertilization procedures
 Study size: 60

Interventions	<p>Randomized portion of anaesthetic: parts of IV</p> <p>Intervention 1: induction/maintenance: propofol/fentanyl</p> <p>Intervention 2: induction/maintenance: midazolam/remifentanyl</p>
Outcomes	<p>Primary outcomes: recovery characteristics 2 anaesthesia protocols</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4</p> <p>Quote: "Four patients in the midazolam/remifentanyl group (13%) would not accept the same anaesthetic procedure for further in vitro fertilization treatment due to intra-operative awareness, while all propofol/fentanyl patients were prepared to accept the same procedure again (P < 0.05)"</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of IV: N₂O no/narcotics/hypnotics bolus MCI/muscle relaxants induction no/maintenance no</p> <p>Midazolam/remifentanyl group, (N = 30) IV midazolam (0.05 mg/kg) infusion remifentanyl; propofol/fentanyl group, N = 30, IV fentanyl (1.5 mg/kg)/propofol (1 mg/kg) than propofol (2 mg to 4 mg/kg/h). SR 50% oxygen if pulse oxygen sat. <= 90% manually assisted using an oral airway-facemask</p> <p>No email address available for ROB survey</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "were then randomly divided in two groups according"
Allocation concealment (selection bias)	Unclear risk	Comment: inadequate information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: inadequate information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: inadequate information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information provided

Celebioglu 2002

Methods	Study design: randomized parallel groups
---------	--

Anaesthetic interventions for prevention of awareness during surgery (Review)

Celebioglu 2002 (Continued)

Study dates: April to September 1999

Participants	Country: Turkey Sex: both Age: 54, 51 ASA: NA Procedure: open heart surgery Study size: 59
Interventions	<p>Randomized portion of anaesthetic: maintenance: CPB: (IV) vs volatile agent (inhalational anaesthesia): music explicit/implicit memory test</p> Intervention 1: maintenance during CPB: fentanyl 10 µg/kg + dehydrobenzperidol 5 mg + 5 mg boluses PRN, N = 30 Intervention 2: maintenance during CPB: fentanyl 10 µg/kg + sevoflurane 2%, N = 29
Outcomes	Primary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Group 1: 5 cases of awareness: 5/30 (17%) Group 2: 0 cases of awareness: 0/29 Awareness 5/59 (9%)
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent/N₂O yes + narcotics + muscle relaxant(s) induction yes/maintenance explicit yes/implicit memory test</p> <p>** IV vs sevoflurane only during CPB rest of case sevoflurane ***Classify as IV techniques compared to volatile techniques because of the unique phase of anaesthesia, CPB</p> Comment: difficult to classify type of anaesthetic technique: IV agents vs volatile during CPB with volatile technique pre and post CPB Induction: etomidate (0.3 mg/kg) + dehydrobenzperidol (5 mg) + fentanyl (5 µg/kg) + vecuronium (0.1 mg/kg) Maintenance: sevoflurane 2% in 50% N ₂ O + fentanyl (5 µg/kg) + additional vecuronium was given during surgery Quote: "...three different audiotapes were played during the operation: one containing radio static and assorted noises such as ringing telephones during the pre-bypass period, a tape of a famous Turkish folk music singer during the actual period of bypass, followed by classical music during the post bypass period" Time of outcome determination: 8 and 24 h after surgery Method of outcome determination: interview Bilge Celebioglu, bilgesel@superonline.com.tr survey 4 February 2013 - failed to deliver

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	Quote: "Random allocation was done by assigning the first 10 patients to Group 1, the next 10 patients to Group 2, and so on"

Celebioglu 2002 (Continued)

Allocation concealment (selection bias)	High risk	Quote: "Random allocation was done by assigning the first 10 patients to Group 1, the next 10 patients to Group 2, and so on"
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Celleno 1993

Methods	Study design: randomized parallel groups Study dates: "1992" (email survey)
Participants	Country: Italy Sex: female Age: mean 31 ASA: Quote: "Ninety healthy women at term ..." implied: I and II Procedure: elective caesarean section Study size: 90
Interventions	Randomized portion of anaesthetic: volatile agent types: IV induction hypnotics Intervention 1: induction: thiopental 15 mg/kg, N = 30 Intervention 2: induction: propofol 2.4 mg/kg, N = 30 Intervention 3: induction: midazolam 0.3 mg/kg, N = 30
Outcomes	Primary outcomes: maternal/fetal anaesthesia drug consumption, APGAR, surgical/anaesthesia times, quality of anaesthesia assessed: standard somatic, sympathetic, and haemodynamic signs. Tearing, sweating, or opening eyes considered to indicate light plane of anaesthesia Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "None of the patients reported any recall of the intraoperative period" Comment: dreams during surgery were reported in 17% of the propofol group and 3% of the thiopental group
Notes	Non-randomized portion of anaesthetic: parts of volatile agent: N₂O yes/narcotics/hypnotics bolus/muscle relaxants induction ys/maintenance PRN Anaesthesia induction: group specific induction regimen (see above) + succinylcholine 1.5 mg/kg + intubation

Celleno 1993 (Continued)

Anaesthesia maintenance 50% N₂O in oxygen + 0.75% isoflurane; after the umbilical cord clamped: N₂O 60% + infusion of oxytocin 0.02 U/mL + atracurium as necessary

Time of outcome determination: first postoperative day

Method of outcome determination: interview

Survey response: 24 January 2011, Capogna dipartimento.anestesia@gruppegarofalo.com

Email about postoperative interviewer 21 January 2013

Emailed again to clarify postoperative blinding 4 March 2013

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Patients were randomized, according to a computerized randomization code, to one of three groups of 30 patients each to receive thiopental 15 mg/kg, propofol 12.4 mg/kg, or midazolam 0.3 mg/kg" Quote: "Computerized randomization code" (email survey)
Allocation concealment (selection bias)	Low risk	Quote: "The investigator was not involved in the anesthesia administration and not aware of the induction agent used" (email survey)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Two anesthesiologists were assigned to each patient, one who was blinded to the induction drug administered to the mother and was conducting the investigation and another who was administering the anesthesia. A neurologist and technician-neurologist performing the maternal electroencephalogram (EEG), as well as the neonatologist who performed the neonatal examinations, also were blinded to the maternal treatment"
Blinding of outcome assessment (detection bias) All outcomes	High risk	Quote: "The day after surgery, all patients were asked about the occurrence of intraoperative dreams, awareness, or discomfort at the time of injection of the induction drug" Quote: "Patient, Anesthesiologist" (email survey) Email about postoperative interviewer 21 January 2013 Emailed again to clarify postoperative blinding 4 March 2013
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "No patient was excluded from the study or the analysis, main outcome data are published" (email survey)
Selective reporting (reporting bias)	Low risk	Quote: "The original study protocol and the original data are no more available: we performed this study more than 15 years ago and the meantime we also moved to another hospital!" (email survey) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Comment: none identified by author response to survey

Chen 2009

Methods Study design: randomized parallel groups

Chen 2009 (Continued)

	Study dates: not provided
Participants	Country: Germany Sex: both Age: 25 to 56 ASA: I and II Procedure: ENT surgery Study size: 50
Interventions	Randomized portion of anaesthetic: parts of TIVA TCI vs MCI Intervention 1: TCI induction/maintenance propofol 4 µg/mL than remifentanil 0.3 µg/kg/min, N = 25 Intervention 2: MCI induction/maintenance bolus propofol 2 mg/kg than continuous propofol 5 mg/kg/h + remifentanil 0.3 µg/kg/min, N = 25 Auditory-evoked potential/BIS
Outcomes	Primary outcomes: anaesthesia and recovery parameters, propofol and remifentanil consumption Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: no recall
Notes	Non-randomized portion of anaesthetic: parts of TIVA: N₂O no/narcotics/hypnotics bolus MCI/ muscle relaxants induction yes/maintenance no: ADM: BIS/MLAEPs Induction: propofol 20 mg (if BIS not < 60 during induction) + rocuronium 0.6 mg/kg, no other relaxants were injected during maintenance of anaesthesia. Maintenance: maintain BIS 40 to 60 and MAP within 20% of baseline values Total dose table: page 931. Total propofol dosages and anaesthesia times available Time of outcome determination: on the day after surgery Method of outcome determination: interview Correspondence to Berthold Bein, MD Email: bein@anesthesie.uni-kiel.de

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Patients were randomly allocated to one of two groups by opening of a sealed envelope"
Allocation concealment (selection bias)	Unclear risk	Quote: "Patients were randomly allocated to one of two groups by opening of a sealed envelope" Comment: not downgraded for not using the word "opaque"
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided

Chen 2009 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor "On the day after surgery, all patients were interviewed about awareness and memory during the perioperative period by an anaesthesiologist unaware of the type of anaesthesia performed"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Cheun 1987

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Korea Sex: both Age: NA ASA: NA Procedure: open heart surgery Study size: 180
Interventions	Randomized portion of anaesthetic: regional anaesthesia: spinal Intervention 1: spinal anaesthesia 0.1 mg/kg of morphine, N = 60 Intervention 2: spinal anaesthesia 1.5 mg/kg of meperidine to 55, N = 60 Intervention 3: spinal anaesthesia mixture of morphine and meperidine to 30 patients, N = 60 To eliminate intraoperative awareness, lorazepam 0.1 mg/kg or diazepam 10 mg to 20 mg/kg
Outcomes	Primary outcomes: haemodynamics (blood pressure) and duration of anaesthesia Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: Table 4. Complications list 16 awareness events but probably data entry error and that was myalgias and awareness is 0 not myalgias; text zero events Quote: "No one complained of intraoperative awareness, and almost all of the patients were comfortable during the surgery and recovery periods"
Notes	Non-randomized portion of anaesthetic: parts of TIVA anaesthesia/N₂O no: muscle relaxants yes - both induction/maintenance Adjuvants: diazepam 10 mg to 20 mg or lorazepam 4 mg to 8 mg Premedication: morphine 0.1 mg IM lorazepam 3 mg to 5 mg glycopyrrolate 0.005 mg IM Anaesthesia induction: thiopental 5 mg/kg + succinylcholine 1 mg/kg/, 100% O ₂ maintenance: 100% O ₂ , lorazepam, pancuronium 0.1 mg/kg, 0.1 mg/kg or diazepam 0.4 mg/kg for intraoperative amnesia No titration anaesthetic strategy for light anaesthesia given in paper

Cheun 1987 (Continued)

Time of outcome determination: first postoperative day
Method of outcome determination: interview

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Open heart surgery patients, who were suitable for spinal anaesthesia were randomly selected"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Chin 2004

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Singapore Sex: female and male Age: parturients ASA: I-II Procedure: caesarean delivery surgery Study size: 23 enrolled and completed study
Interventions	Randomized portion of anaesthetic: volatile agent types Intervention 1: maintenance: ET 1% sevoflurane Intervention 2: ET 1.5% sevoflurane Successful BIS outcome N = 12; unsuccessful BIS outcome N = 11, N = 23

Chin 2004 (Continued)

The up-down allocation method randomized each subsequent patient's ET dose based on the previous patients ET dose being judged as successful or unsuccessful based on BIS criteria explained in ROB table

Outcomes

Primary outcomes: anaesthesia and recovery parameters, APGAR, successful BIS response to randomized ET sevoflurane and relationships between sevoflurane concentration and other outcome variables
Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Quote: "There were no instances of intraoperative recall"

Notes

Non-randomized portion of anaesthetic: parts of volatile agent: N₂O yes/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance yes: ADM BIS

IV sodium thiopental 4 mg/kg and succinylcholine 1 mg to 1.5 mg/kg intubation

Maintained N₂O 3 L/min/O₂ 3 L/min. Atracurium bolus of 30 mg + 10 mg PRN

Patients were mechanically ventilated end-tidal concentration 0.5 MAC in 50%

Quote: "Our aim was to determine the BIS values achieved with the equivalent end-tidal concentration of sevoflurane and to determine if a larger concentration would consistently maintain **BIS values 60**"

N₂O throughout surgery. Morphine 0.1 mg to 0.15 mg/kg was after delivery

ROB survey 7 January 2014. Email: gasgetJie@yahoo.co.uk

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The end-tidal concentration (ETC) of sevoflurane administered to each patient was decided using an up down sequential allocation design, wherein each patient's dose was determined by the previous patient's BIS response" Comment: this "up down sequential method" is debated in (Benhamou 2003 ; Lacassie 2004)
Allocation concealment (selection bias)	Low risk	Comment see above
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment see above Comment: sent author ROB survey to clarify 7 January 2014
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "None of the patients had any recall of intra-operative events"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information

Choi 2012

Methods	Study design: randomized parallel groups Study dates: December 2010 and February 2011
Participants	Country: USA and Korea Sex: female Age: mean 31 to 32 ASA: I-II Procedure: caesarean delivery Study size: 70 enrolled, 64 recruited, 61 completed study
Interventions	Randomized portion of anaesthetic: volatile agent types Intervention 1: (N = 15) control 1 Intervention 2: (N = 15) control 2 Intervention 3: (N = 16) pre-sevoflurane 1 group - 1.2 to 1.3 vol (control 2) end-tidal sevoflurane; but were also pre-exposed to 1 vol% sevoflurane for the final 1 min of the pre-oxygenation period Intervention 4: (N = 15) pre-sevoflurane 2 groups - 1.0 to 1.1 vol (control 1) end-tidal sevoflurane; but were also pre-exposed to 1 vol% sevoflurane for the final 1 min of the pre-oxygenation period
Outcomes	Primary outcomes: BIS values, non-invasive arterial pressure, and heart rate Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "no patient reported intraoperative recall"
Notes	Non-randomized portion of anaesthetic: parts of volatile agent: N₂O yes/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance yes: ADM: BIS recorded Induction: thiopental sodium 4 mg/kg/trachea intubated after muscle paralysis with succinylcholine 1.5 mg/kg, with rocuronium 0.5 mg/kg muscle relaxation. Anaesthesia was maintained with either 1.0 to 1.1 or 1.2 to 1.3 vol% ET sevoflurane/50% N ₂ O/oxygen. After delivery, N ₂ O/O ₂ :2:1/sevoflurane ET 0.8 vol% minimize uterine tocolysis. Midazolam 0.05 mg to 0.1 mg/kg and fentanyl 1.0 mg to 3.0 mg/kg/ sedation or analgesia to the end of surgery. Lung ventilation ET CO ₂ 30 to 35 mmHg ROB survey. We emailed sjinwoo@hotmail.com on 22 March 2015; multiple emails sent in 2013

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "This prospective, randomized, controlled trial... registered with the Clinical Research Information Service (code number KCT0000069)" Quote: "Patients were randomly divided into one of the four groups using a computer-generated randomization schedule; the randomized block size was eight (http://www.randomization.com); randomization was achieved using sealed envelopes prepared by our consultant statistician"
Allocation concealment (selection bias)	Unclear risk	Comment: using sealed envelopes
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "The anesthetist controlling sevoflurane concentration was blinded to the BIS value" Comment: BIS not randomized; sevoflurane concentrations were randomized

Choi 2012 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "Each patient was asked on discharge from the postoperative care unit and 24 h after the operation, whether she could recall any events during the operation"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "Of the 70 subjects initially enrolled in the study, six were excluded because they requested regional anaesthesia. Thus, 64 subjects were randomized into the four groups, and 61 completed the study..." Comment: no high-risk exclusions Comment: 2 exclusions from control group 2/30 vs 1 from experimental group 1/30: not significantly different, Peto OR 0.48 (0.05 to 4.85)
Selective reporting (reporting bias)	Low risk	Comment: as above Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information

Clyburn 1986

Methods	Study design: randomized parallel groups Study dates: "From memory, during 1984. Completed by end of 1984" (email bias survey, see notes)
Participants	Country: UK Sex: female Age: mean: 29 and 26 ASA: I-II Procedure: minor gynaecological surgery Study size: 60
Interventions	Randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia Intervention 1: induction: midazolam 70 µg/kg, N = 30 Intervention 2: induction: diazepam 150 µg/kg, N = 30
Outcomes	Primary outcomes: pain on injection, nausea/vomiting, recovery time Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: no awareness "Patients were asked to volunteer comments on unpleasant aspects of the procedure"
Notes	Non-randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia: N₂O yes/narcotics and/or hypnotics bolus/muscle relaxants induction no/maintenance no Induction: etomidate 300 µg/kg + fentanyl 1.5 µg/kg + group-specific study drug (see above) Maintenance: 66% N ₂ O in 34% O ₂ with intermittent boluses of etomidate as needed Quote: "Relaxation was deemed unsatisfactory in eight of the 60 patients, but in each this was corrected by a further increment of etomidate." Survey response: 17 January 2011 Paul Clyburn clyburn@cf.ac.uk

Clyburn 1986 (Continued)

Emailed author 26 December 2013 to clarify allocation

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The patients were randomly allocated to one of two anaesthetic techniques" Quote: "I believe we used randomisation tables with sealed envelopes." (email survey)
Allocation concealment (selection bias)	Low risk	Quote: "Sequential envelope opening" (email survey) Dr Clyburn responded to email survey requesting clarification of allocation process on 28 December 2013 Quote: "Yes we used opaque envelopes"
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "No blinding occurred" (email survey)
Blinding of outcome assessment (detection bias) All outcomes	High risk	Quote: "No blinding occurred" (email survey)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "There were no exclusions or attrition of patients studied" (email survey)
Selective reporting (reporting bias)	Low risk	Quote: "Protocol no longer available but text does report all expected outcomes" (email bias survey, see notes) Quote: "Basically, there were no differences between the two studied groups i.e. negative findings only" (email survey) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote: "Not aware of any other sources of bias" (email survey)

Coates 1987

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: UK Sex: female and male Age: 39 to 57 ASA: I-II Procedure: Quote: "body surface surgery" Study size: 17

Coates 1987 (Continued)

Interventions	<p>Randomized portion of anaesthetic: parts of IV: MCI hypnotic</p> <p>Intervention 1: maintenance: infusion propofol 54 µg/kg/min, N = 9</p> <p>Intervention 2: propofol 108 µg/kg/min, N = 8</p>
Outcomes	<p>Primary outcomes: haemodynamic effects</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Quote: "None of the patients, including those who had moved in response to the first incision or the introduction of IPPV, had any recall of these events or suggestion of awareness during the procedure"</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of IV: N₂O yes/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance PRN</p> <p>Premedicated with morphine, 0.15 mg/kg. Anaesthesia was induced: propofol, 2 mg/kg/succinylcholine 1 mg/kg for intubation and then 0.3 mg/kg PRN during surgery/intermittent supplement 67% N₂O. Positive pressure ventilation (IPPV) to maintain the end-tidal CO₂ at 4.8% to 5.4%</p> <p>No email address available for ROB survey</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Normotensive patients (ASA I or II) scheduled for body surface surgery were randomly allocated to group 1 (propofol infusion rate: (54 mcg/kg/min) or group 2 (108 mcg/kg/min)"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: no information
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "Patients were questioned about their experience during induction, maintenance and recovery with specific attention to the possibility of awareness"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information provided; no ROB survey data

Collins 1996

Methods	<p>Study design: randomized parallel groups</p> <p>Study dates unknown</p>
---------	--

Collins 1996 (Continued)

Participants	Country: Australia Sex: female Age: 25 to 39 ASA: I-II Procedure: laparoscopic gynaecological surgery Study size: 30
Interventions	<p>Randomized portion of anaesthetic: parts of TIVA vs parts of volatile agent + N₂O</p> <p>Intervention 1: maintenance: ventilated oxygen/air mix FIO₂ = 0.3 and infusion of propofol (110 mg/mL) mixed with 4 mL of alfentanil (0.5 mg/mL), starting rate 1.5 mL/kg/h, N = 15</p> <p>Intervention 2: maintenance: oxygen/N₂O FIO 0.3 isoflurane starting with inspired 1% adjusted pert response to surgical stimulation, N = 15</p>
Outcomes	Primary outcomes: recovery characteristics Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 "No patient experienced any awareness"
Notes	<p>Non-randomized portion of anaesthetic: parts of TIVA vs parts of volatile agent N₂O no/narcotics/hypnotics bolus MCI/muscle relaxants induction yes/maintenance unclear</p> <p>Induction: propofol 2 mg to 3 mg/kg-1 + alfentanil 7.5 µg kg + vecuronium 75 µg kg/intubation + ketorolac 30 mg LV + droperidol 0.5 mg + glycopyrronium bromide 3 µg kg; residual muscle paralysis was reversed</p> <p>No email address available for ROB survey</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Thirty patients undergoing laparoscopic gynaecological sterilization, as day-cases, were randomly allocated to receive either total intravenous anaesthesia (TIVA) with a propofol and alfentanil mixture or a standard inhalational technique"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor "On arrival in recovery, a second investigator, blinded to the type of anaesthesia received, assessed the times to eye opening, obeying command/hand squeeze) and orientation (giving the correct date of birth)...Patients were specifically asked about awareness"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "One patient in each group was given post-operative pethidine prior to the assessments and they were therefore excluded from the final analysis" Comment: 2/30 = 7% possibly high-risk awareness balanced exclusion
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria

Collins 1996 (Continued)

Other bias	Unclear risk	Comment: insufficient information provided
------------	--------------	--

Crawford 1985

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: England Sex: female Age: parturients ASA: not stated Procedure: emergency caesarean section Enrolled and completed study size: 237 + 540 = 777
Interventions	Randomized portion of anaesthetic: volatile agent types Intervention 1: maintenance: trichloroethylene 0.2 vol. %, N = 135 Intervention 2: maintenance: trichloroethylene 0.3 vol. %, N = 128 Intervention 3: maintenance: halothane (0.2 vol. %), N = 129 Intervention 4: maintenance: halothane (0.3 vol. %), N = 129 Intervention 5: maintenance: halothane (0.4 vol. %), N = 127 Intervention 6: maintenance: halothane (0.5 vol. %), N = 129
Outcomes	Primary outcomes: intra and postoperative parameters specific to C-sections: anaesthesia consumption and time to surgical/anaesthesia endpoints such as entry of uterus and delivery of fetus, blood loss, surgical/anaesthesia duration, fetal wellbeing parameters and maternal recovery parameters (table IV, table V) Secondary outcome: awareness and unpleasant dreams (table VI), awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Comment: Table VI: 22 awareness events; dreams: 13 Low dose: trichloroethylene 0.2 + H 0.2 + H 0.3 = 19 aware/393 vs high dose: trichloroethylene 0.3 + H 0.4 + H 0.5 = 3 aware/384 Quote: "Our results clearly show that trichloroethylene 0.2 vol.%, halothane 0.2 vol.% or halothane 0.3 vol.% [low dose]...produces an unacceptably high incidence of "awareness plus unpleasant dreams"
Notes	Non-randomized portion of anaesthetic: parts of volatile agent (parts of potent volatile technique)/nitrous/hypnotic/muscle relaxants (succinylcholine infusion) induction yes/maintenance yes Induction: hyoscine 0.4 mg + thiopentone 250 mg to 300 mg + suxamethonium 100 mg Maintenance: O ₂ 8 L/min + nitrous 4 L/min + infusion suxamethonium 1 mg/mL No titration anaesthetic strategy for light anaesthesia Trichloroethylene 8/263 = 3% The incidence of maternal awareness unpleasant dreams was unacceptably high in the lower concentrations were used, and it is recommended that either trichloroethylene 0.3 vol. % or halothane 0.4 or 0.5 vol. % be used

Crawford 1985 (Continued)

Comment: since the mechanism of intraoperative dreams is controversial, we simulated a reduction of 50% in patient awareness/dream reports balanced across both groups as a method to exclude a portion of dream reports and the significant difference between low and high-dose inhalation agents persisted (2/384 vs 9/393), Peto OR 0.28 (95% CI 0.09 to 0.93)

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The choice of volatile agent was made by reference to a series of randomised numbers..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: inadequate information
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided Quote: "On the day following operation, the mother was asked if she had had any dreams during the operation. If the answer was affirmative, the possibility of "awareness" was pursued...The entire series showed a significantly higher incidence of awareness or unpleasant dreams, or both, when.."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Crawford ME 1984

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Denmark Sex: female Age: 27 and 25 ASA: I-II Procedure: termination of pregnancy (day case surgery) Study size: 100
Interventions	Randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia Intervention 1: induction: midazolam 0.2 mg/kg + midazolam 0.1 mg/kg if needed; maintenance: midazolam 0.15 mg/kg, fentanyl 0.1 mg, and/or both as needed, N = 50 Intervention 2: induction: thiopentone 4 mg/kg + supplementary dose (2 mg/kg) as needed; maintenance: thiopentone 3 mg/kg, fentanyl 0.1 mg, and/or both as needed, N = 50

Crawford ME 1984 (Continued)

Outcomes Primary outcomes: haemodynamic effects, anaesthetic consumption, nausea, vomiting and dizziness, degree of amnesia

Awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Quote: "no patient was able to remember any part of the operation"

Notes **Non-randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia: N₂O yes/narcotics and/or hypnotics bolus/muscle relaxants induction no/maintenance no**

Non-randomized anaesthetics: premedication droperidol 0.5 mg IM, both groups 30 min before surgery. Group A (N = 50), induction midazolam 0.2 mg/kg IV, midazolam 0.1 mg/kg, midazolam 0.15 mg/kg and/or fentanyl 0.10 mg: PRN eyelash reflex present 3 min after the initial dose, after loss consciousness, fentanyl 0.15 mg, maintenance 67% N₂O/oxygen. PRN: movement, peripheral vasoconstriction, lacrimation, sweating, change heart rate or arterial pressure), Group B (N = 50), induction thiopentone 4 mg/kg IV and PRN 2 mg/kg after 3 min, fentanyl 0.15 mg IV; maintenance: 67% N₂O/oxygen, supplemented by thiopentone 3 mg/kg and/or fentanyl 0.10 mg

Comment: balanced anaesthesia describes IV technique in this RCT

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "allocated randomly to one of two groups (A and B)"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Czarko 2013

Methods Study design: randomized parallel groups
Study dates: not stated

Participants Country: Poland
Sex: female

Czarko 2013 (Continued)

Age: aged: 31 to 50
 ASA: I-II
 Procedure: elective gynaecological procedures, short gynaecological procedures, and caesarean section patients
 Study size: 337

Interventions

Randomized portion of anaesthetic: TIVA (TCI) vs parts volatile agent + infusion narcotics (TCI)

Intervention 1: (N = 51) group A, **TIVA/TCI** remifentanyl incremental doses to plasma 8.5 ng mL⁻¹, then infusion propofol to of 8 µg mL⁻¹. Maintained infusion remifentanyl/propofol pre-set doses of 3 ng to 6 ng/mL⁻¹ and 2 µg to 4 µg/mL⁻¹

Intervention 2: (N = 95) group B — thiopentone (5 mg/kg-1), fentanyl (3 µg to 5 µg/kg-1), cisatracurium (0.1 mg/kg-1), **sevoflurane** (1 to 2 vol%)

Intervention 3: (N = 16) group C — propofol (2 mg/kg-1), fentanyl (3 µg to 5 µg/kg-1) **sevoflurane** (1 to 2 vol%), cisatracurium

Intervention 4: (N = 175) group D **intravenous (IV)** thiopentone (5 mg/kg-1), suxamethonium (1 mg/kg-1), fentanyl (3 µg to 5 µg/kg-1) and cisatracurium (0.05 mg/kg-1) after foetus extraction

All patients: **N₂O** and O₂; FIO₂ 0.33; auditory evoked potentials: **AEP** target range: 15 to 25

Group A, B, C: **elective gynaecological** procedures; Group D: **caesarean section**

Quote: "All of the patients received a mixture of N₂O and O₂; FIO₂ was maintained at the level of 0.33"

Outcomes

Primary: outcomes: intraoperative sensations

Secondary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4

Quote: "3 cases, the descriptions of intraoperative events suggested intraoperative awareness"

Comment: **sensations usually associated with intraoperative dreams** were reported by 14% (46/337); this is evidence that dreams maybe related to light planes of anaesthesia as suggested by other authors over decades

Notes

Non-randomized portion of anaesthetic: TIVA vs parts of volatile agent/N₂O/infusion of narcotics (TCI) + muscle relaxants induction yes/maintenance yes/ADM AEP (15 to 25)

In group A total intravenous anaesthesia: target controlled infusion (**TCI**), cisatracurium, (0.1 mg/kg-1), infusion of remifentanyl incremental doses until reaching plasma concentration 8.5 ng/mL⁻¹, infusion propofol to plasma concentration 8 µg/mL⁻¹. TCI maintained infusion remifentanyl/propofol pre-set doses 3 ng to 6 ng/mL⁻¹ and 2 µg to 4 µg/mL⁻¹; remaining groups, combined general anaesthesia: Group B - thiopentone (5 mg/kg-1), fentanyl (3 µg to 5 µg/kg-1), cisatracurium (0.1 mg/kg-1), sevoflurane (1 to 2 vol%)

Group C - propofol (2 mg/kg-1), fentanyl (3 µg to 5 µg/kg-1), sevoflurane (1 to 2 vol%); group D — thiopentone (5 mg/kg-1), suxamethonium (1 mg/kg-1), fentanyl (3 µg to 5 µg/kg-1) and cisatracurium (0.05 mg/kg-1) **after foetus extraction**. All received N₂O and O₂; FIO₂ 0.33; sleep depth: measuring auditory evoked potentials: AEP monitor, A-line auditory evoked potential index was kept within range 15 to 25

Comment: Annexe: questionnaire for assessment of intraoperative awareness: 22 questions assess recall/pain/dreams/tactile/auditory sensations

Quote: "...Group A received **total intravenous anaesthesia with TCI**, and groups B, C and D received **balanced anaesthesia**. The depth of anaesthesia was monitored with an AEP monitor. Blinded structured"

Comment: see Dryad topic sensations associated with dreams

Czarko 2013 (Continued)

 ROB survey. We emailed on 22 March 2015 kate.czarko@wp.pl
Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "patients were enrolled in the study and randomly allocated to 4 groups according to the type of general anaesthesia performed...."
Allocation concealment (selection bias)	Unclear risk	Comment: no information
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: no blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patients were surveyed three times: 2 h after anaesthesia and on post-anaesthesia days 7 and 30 (by phone). Postoperative surveys were conducted by a person not involved in anaesthesia"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no dropouts
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information

De Kock 1995

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: NA Sex: female and male Age: 35.3 ± 9.2 ASA: NA Procedure: abdominal surgery colic resection Study size: 40
Interventions	Randomized portion of anaesthetic: regional anaesthesia: epidural: 2 infusion rates Intervention 1: induction epidural clonidine (4 µg/kg in 10 mL) infused in 20 min followed by a 2 µg/kg infusion (5 mL/h) during 12 h (Group 1) Intervention 2: induction epidural sufentanil (0.5 µg/kg in 10 mL) in 20 min followed by a 0.25 µg/kg infusion (5 mL/h) during 12 h (Group 2)
Outcomes	Primary outcomes: efficacy and side effects of epidural clonidine and sufentanil

De Kock 1995 (Continued)

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4

Comment: 2 awareness reports Group 1 (clonidine)

Notes

Non-randomized portion of anaesthetic: IV N₂O propofol MCI/muscle relaxants: both induction/maintenance

Before anaesthesia, epidural catheter inserted L1-L2; induction and maintenance: propofol/N₂O, propofol bolus (0.5 mg/kg) PRN if inadequate, then bolus sufentanil 0.035 µg/kg. Neuromuscular block: infusion atracurium (5 pg/kg/min). Atracurium infusion was discontinued at the beginning of the closure of the peritoneum. Anaesthesia was maintained with a **propofol infusion** of 3 mg/kg/h and 50% N₂O in oxygen. Neuromuscular block maintained infusion atracurium (5 pg/kg/min)

ROB survey email sent 3 January 2014 decock@anes.ucl.ac.be; response 10 January 2014

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	ROB survey response by Dr Marc de Kock: "computer generated randomization list" Quote: "At this time, patients were randomly assigned to receive either epidural clonidine (Group 1) or epidural sufentanil..."
Allocation concealment (selection bias)	Low risk	ROB survey response: "study drugs were prepared by an anesthesiologist not involved in patient's care and were blinded to the anesthesiologist in charge"
Blinding of participants and personnel (performance bias) All outcomes	Low risk	ROB survey response: "Patient, Anesthesiologist, Awareness outcome assessor (interviewer)" Quote: "At this time, patients were randomly assigned to receive either epidural clonidine (Group 1) or epidural sufentanil (Group 2)"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Comment: see above
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "One patient in Group 2 presented with an immediate postoperative respiratory depression that required prolonged ventilatory support and was therefore excluded from the postoperative study protocol..." Comment: respiratory depression: epidural narcotic effect: sign of deep anaesthesia hence a low risk of awareness
Selective reporting (reporting bias)	Low risk	ROB survey response: "the study protocol is available and all the outcome data are reported" Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	ROB survey response: "no particular bias"

Deeprise 2005

Methods

Study design: randomized parallel groups
 Study dates: not stated

Deeprise 2005 (Continued)

Participants	Country: UK Sex: both Age: 16 to 72 ASA: I Procedure: orthopaedic Study size: 64 enrolled, 62 completed study
Interventions	<p>Randomized portion of anaesthetic: parts of IV/narcotics (fentanyl vs no fentanyl) induction/implicit memory word test</p> Intervention 1: induction: fentanyl 1.5 µg/kg + "sleep" dose of propofol, N = 32 Intervention 2: induction: no fentanyl + propofol, N = 30
Outcomes	Primary outcomes: relationship between BIS (mean-BIS and max-BIS) and memory scores Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 2 Comment: no patients reported awareness; implicit memory was present in both groups Quote: "Patients were interviewed for awareness using the structured interview of intraoperative events (based on Russell and Wang, 1997). There was no evidence for spontaneous or prompted recall"
Notes	<p>Randomized portion of anaesthetic: parts of IV: induction: hypnotic/maintenance: propofol TCI nitrous/SR: LMA/no muscle relaxants/ADM BIS open not target</p> Induction: see interventions (' sleep ' dose of propofol)/narcotics(randomized group yes/no fentanyl); maintenance: 3 mg to 6 mg/kg/h infusion rate + nitrous 66% and O ₂ 33%; no titration anaesthetic strategy for light anaesthesia given in paper. Maintained TCI target-controlled infusion 3 mg and 9 mg/kg-1 h-1: clinical judgement. N ₂ O 66% and oxygen 33% spontaneously laryngeal mask. BIS not blinded but also not used to guide titration of anaesthetic Quote: "Priming remained above zero when data from the six patients with above-chance performance on the yes-no recognition test and the 25 patients with max-BIS >60 were excluded from the analysis" Comment see Dryad: evidence for and against implicit memory c.deeprise@shfac.uk emailed 2011 Jackie Andrade jackie.andrade@plymouth.ac.uk emailed 1 February 2013 Responded to survey: 4 June 2013 Catherine Deeprise catherine.deeprise@plymouth.ac.uk

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Patients were assigned randomly to a fentanyl or no-fentanyl study group to which the experimenter (CD) was blinded" Quote survey response by Dr Deeprise: "Simple randomization in which either "fentanyl" or "no-fentanyl" was drawn from a bag for each patient. A total of 64 (32 per group) slips of paper indicating "fentanyl" or "no-fentanyl" were created before the study commenced. The sample size was based on power calculations (see point 7 below)"
Allocation concealment (selection bias)	High risk	Experimenter (CD) was blinded to anaesthetic assignment Anaesthetist was not blinded to anaesthetic protocol, but was not involved in memory testing" Comment: experimenter did intra and postoperative management of protocol not the anaesthesiologist

Deeprise 2005 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote survey response by Dr Deeprise: "The experimenter was blinded to anaesthetic technique. The study completed the proposed sample size based on our pilot work using the auditory word stem completion tasks and previous published research (Deeprise and Andrade, 2004)" Comment: the protocol for this review (Messina 2008), includes implicit memory tests in the classification of outcomes (class 2). But, interventions to decrease wakefulness/awareness are defined as anaesthetic regimens and ADMs. The anaesthesiologist, although blinded to the word tests, was not blinded to the randomized anaesthetics.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor: "Patients were assigned randomly to a fentanyl or no-fentanyl study group to which the experimenter (CD) was blinded" Comment: experimenter did postoperative interviews
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment we analysed 2 total and 1 high-risk awareness dropout(s) from the no fentanyl group and found no significant difference between groups: (2/30) vs (0/32), Peto OR 8.18 (0.50 to 133.94); (1/30) vs (0/32) Peto OR 7.90 (0.16 to 398.87)
Selective reporting (reporting bias)	Low risk	Quote: "The study is described in full in my Phd thesis which presents the rationale for the expected outcomes as described in the published paper. There are no unreported outcome variables" Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Comment: underpowered study which was identified by author ROB survey response as a source of other bias is a source of imprecision, which is not covered in this domain. No other sources of internal biases identified by author Author sent ROB table grades for comment 7 December 2013 catherine.deeprise@plymouth.ac.uk

Deshpande 2009

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: India Sex: female and male Age: 15 to 50 ASA: >= III Procedure: open heart surgery: valvular and simple congenital heart disease surgery Study size: 100, 98
Interventions	Randomized portion of anaesthetic: volatile agent types: IV narcotics Intervention 1: sufentanil group: induction sufentanil 0.5 µg/kg; maintenance: sufentanil 0.1 µg/kg PRN Intervention 2: fentanyl group: induction fentanyl 3 µg/kg; maintenance: fentanyl 1 µg/kg

Deshpande 2009 (Continued)

Outcomes	<p>Primary outcomes: anaesthetic consumption and recovery parameters, mean ventilation time between groups</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4</p> <p>Comment: sufentanil group 1 patient awareness event at sternotomy</p>
----------	--

Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent: N₂O yes/narcotics/hypnotics bolus MCI/muscle relaxants induction yes/maintenance yes</p> <p>Induction: midazolam 0.05 mg/kg sleep dose thiopental vecuronium 0.1 mg/kg for intubation; maintenance: N₂O isoflurane ET 0.8% to 1% midazolam and vecuronium PRN before and after CPB with goal of stable haemodynamics; CPB: infusion propofol 4 mg to 5 mg/kg/h; sufentanil 0.1 µg/kg PRN or fentanyl 1 µg/kg, max doses for case sufentanil fentanyl midazolam 1 and 6 µg/kg and 5 mg respectively</p> <p>Emailed ROB survey 2 January 2014 and asked Dr Deshpande to clarify the reason for referral of awareness patient to psychiatrist for an accurate classification category Table 1 desh56@hotmail.com</p>
-------	---

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"...prospective randomized double blind study..."
Allocation concealment (selection bias)	Unclear risk	Comment: inadequate information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: inadequate information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: inadequate information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	<p>Comment: 100 patients, 50 in each group, "another 2 patients were excluded from study because of prolonged ventilation >6 hours secondary to surgical complication"</p> <p>Comment: no significant difference between groups if exclusions were in one or both groups (2/50 vs 0/50), Peto OR 7.54 (0.47 to 122.28)</p>
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information provided

Dhadphale 1979

Methods	<p>Study design: randomized parallel groups</p> <p>Study dates: not stated</p>
Participants	<p>Country: USA</p> <p>Sex: both</p>

Dhadphale 1979 (Continued)

Age: 49, 52
 ASA: not given
 Procedure: mitral or aortic valve replacement
 Study size: 32 enrolled, 30 completed study

Interventions	Randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia Intervention 1: induction: diazepam 0.4 mg/kg + ketamine 2 mg/kg, N = 16 Intervention 2: induction: morphine 3 mg/kg, N = 16
Outcomes	Primary outcomes: circulatory responses (heart rate, mean arterial blood pressure, PaCO ₂) Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 No intraoperative awareness occurred in either group
Notes	Non-randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia: N₂O yes/narcotics and/or hypnotics bolus MCI/muscle relaxants induction yes/maintenance yes Premedication: morphine 5 mg to 10 mg + scopolamine 0.2 mg to 0.4 mg; induction: succinylcholine 1.5 mg/kg; maintenance: nitrous 50% + pancuronium 0.1 mg/kg + ketamine 1 mg/kg/h No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "randomly selected patients were given..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "All patients were seen postoperatively and questioned specifically about possible awareness..."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "Two deaths occurred, both in Group I." Comment: group 1 (ketamine) patients at increased risk of awareness (2/16) were not significantly different from morphine group (0/16), Peto OR 7.90 (0.47 to 132.20)
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Drover 2002

Methods	Study design: randomized parallel groups
---------	--

Anaesthetic interventions for prevention of awareness during surgery (Review)

Drover 2002 (Continued)

Study dates: 18 May 1999 to 4 January 2000 (ROB survey)

Participants	Country: USA Sex: males and females Age: 18 to 80 years ASA: I-III Procedure: elective surgical procedures scheduled for at least 30 min Study size: 347 (102 controls + 245 randomized), 306 completed study (82 controls + 224 randomized patients)
Interventions	Randomized portion of anaesthetic: parts of IV: ADM: PSI (25 to 50) Intervention 1: PSI guidance, N = 123 Intervention 2: standard practice guidelines, N = 122
Outcomes	Primary outcomes: drug dosage, recovery times Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "There was no incidence of reported awareness or memories in any patient in any group"
Notes	Non-randomized portion of anaesthetic: parts of IV: N₂O yes/narcotics/hypnotics bolus MCI/NMBs induction PRN/maintenance PRN: SR/LMA or CV/ETT ADM EEG Premedication: midazolam 1 mg to 2 mg; induction: propofol 1 mg to 3 mg/kg + alfentanil $\leq 30 \mu\text{g}/\text{kg}$; maintenance: propofol 140 $\mu\text{g}/\text{kg}/\text{min}$ + alfentanil 0.5 $\mu\text{g}/\text{kg}/\text{min}$ + nitrous 50%; muscle relaxation PRN Quote: "...consciousness, patients breathed via a laryngeal mask airway or a muscle relaxant was administered and an endotracheal tube placed." Comment: the isolated-arm technique was used to evaluate patient movement after administration of the neuromuscular blocking agent. The authors used a historic control group in the comparison with the experimental group. Comment: wakefulness defined by response to complex command was not used; somatic responses only were recorded; also, patients were not given additional doses of anaesthesia until autonomic responses stopped but until target range of ADM reached Author responded to the email bias survey response: 8 August 2011, David Drover ddrover@stanford.edu Email sent to author to clarify these issues ddrover@stanford.edu 7 December 2013 Email sent to clarify these issues: ddrover@stanford.edu 7 December 2013; responded 8 December 2013

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Randomization was determined prior to start of the study was by the coordinating company. Randomization was performed using MS Excel and randomization assignments were placed in sealed envelopes that were opened once the subject entered the study." (Email bias survey, see notes)
Allocation concealment (selection bias)	Unclear risk	Quote from ROB survey response: "The assignment was in a sealed envelope and was not opened until anesthesia was to begin..." Comment: await author clarification on the use opaque, sealed envelopes

Drover 2002 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	High risk	<p>Comment: readers can view a detailed description by the author, Dr Drover</p> <p>Quote from ROB survey response: "Patient, Anesthesiologist... For the SPC group, anesthesiologists... were blinded to the PSI information... PSI group were guided by the PSI measure...One of the goals of the study was to see change in behavior of the anesthesiologist with respect to drug use. If the anesthesiologist was blinded, he could not see the output from the monitor, otherwise, if unblinded, he had full access to the monitor. The patient was always blinded"</p> <p>Comment: both groups not blinded to SCPs</p>
Blinding of outcome assessment (detection bias) All outcomes	High risk	<p>Quote: "Postoperative interview was not conducted by a blinded investigator. Awareness was a secondary endpoint." (email bias survey)</p>
Incomplete outcome data (attrition bias) All outcomes	Low risk	<p>Comment: Table 1 enrolled 347 is correct and methods section 306 is an error per ROB survey response; 347 enrolled - 306 completed study = 41/347 = 12% dropout; but, 347 - 102 controls and PSI training group = 245 randomized to PSI vs SCP - 41 total dropouts - 20 controls/training dropouts = 21/245 (9%) dropouts study sample. Reasons for not completing protocol: SCP vs PSI: no high-risk dropouts</p> <p>Comment: Table 1: SCP exclusions 10/122 vs PSI 11/123, Peto OR 1.10 (0.45 to 2.68): no significant difference between groups</p> <p>Email sent to clarify these issues: ddrover@stanford.edu 7 December 2013; Dr. Drover responded 8 December 2013</p>
Selective reporting (reporting bias)	Low risk	<p>"No" (email bias survey)</p> <p>Comment: awareness outcome part of inclusion criteria</p>
Other bias	Low risk	<p>Quote: "One of the goals of the study was to see change in behavior of the anesthesiologist with respect to drug use"</p>

Dunnett 1977

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: USA Sex: both Age: adult patients ASA: not given, "good health" Procedure: no surgery identification Study size: 77
Interventions	<p>Randomized portion of anaesthetic: parts of volatile agent (balanced) with intravenous (IV) induction agents</p> <p>Intervention 1: induction: thiopentone 3 mg/kg, N = 38 Intervention 2: induction: ketamine 2 mg/kg, N = 39</p>
Outcomes	Primary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4

Dunnett 1977 (Continued)

Quote: "One patient in the thiopentone group remembered an unpleasant sensation of a tube being put into his throat and associated this with gagging"

Notes

Non-randomized portion of anaesthetic: parts of volatile agent (balanced) N₂O yes/narcotics and/or hypnotics bolus/muscle relaxants induction yes/maintenance yes

Premedication: tubocurarine 3 mg; induction: intervention + suxamethonium 1 mg/kg; maintenance: nitrous in O₂ + **halothane** + pancuronium; no anaesthetic strategy for light anaesthesia described

Quote from discussion section, see Dryad: "The possibility of **retrograde amnesia**..."

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Patients were allocated at random to induction of anaesthesia with either a barbiturate or ketamine..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "The same interviewer was used throughout the series and was not aware which induction agent had been used in a given patient"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Echevarria 1998

Methods

Study design: randomized parallel groups
 Study dates: "17 months" (email bias survey, see notes)

Participants

Country: Spain
 Sex: male and female
 Age: 18 to 60
 ASA: I-II
 Procedure: elective surgery
 Study size: 100 enrolled, 98 completed study

Interventions

Randomized portion of anaesthetic: TIVA MCI vs volatile agent types

Intervention 1: desflurane/N₂O induction: atropine 0.5 mg + fentanyl 2 µg/kg + atracurium 0.5 mg/kg; maintenance: nitrous 60% + desflurane, N = 25

Echevarria 1998 (Continued)

Intervention 2: isoflurane/N₂O induction: atropine 0.5 mg + fentanyl 2 µg/kg + atracurium 0.5 mg/kg; maintenance: nitrous 60% + isoflurane, N = 25

Intervention 3: **IV anaesthesia** fentanyl/N₂O induction: atropine 0.5 mg + fentanyl 2 µg/kg + atracurium 0.5 mg/kg; maintenance: nitrous 60% + fentanyl 3 µg/kg, N = 25

Intervention 4: **total intravenous anaesthesia** induction: alfentanil 15 µg/kg + propofol; maintenance: O₂ + propofol 6 mg/kg/h + alfentanil 50 µg/kg then 1.5 µg/kg/min, N = 25

All groups: taped music followed by an order requiring a non-verbal response

Outcomes	<p>Primary outcomes: evidence of implicit memory (taped music followed by an order requiring a non verbal response)</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 2</p> <p>Comment: no explicit memories of intraoperative events; evidence for implicit memory in isoflurane group</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent: N₂O yes/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance unclear/music memory and non-verbal response</p> <p>Music not word test (Isabel Dufano - translator)</p> <p>Author responded to the email bias survey on 22 January 2011. Responses are recorded in the 'Risk of bias' table. Author's responses were translated by translator Brett Smith/translator Isabelle Dufano</p> <p>Survey response: 22 January 2011, Mercedes Echevarria mercedes.etxeba@terra.es</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Comment: author ROB survey response that sequence allocation was performed by using Statgraphics software
Allocation concealment (selection bias)	Low risk	Comment: author ROB survey response that an anaesthesiologist who was blinded to the anaesthetic administration performed the allocation concealment and postoperative interview
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: difficult to blind infusion pump sound vs inhaled agent
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Comment: see above
Incomplete outcome data (attrition bias) All outcomes	Low risk	<p>Quote: "In recruiting only 1 patient refused to participate in the study. During the study one patient was excluded from group 3 (fentanyl and nitrous oxide) due to hospital discharged at 12 hours." (email bias survey, see notes)</p> <p>Comment: dropout unrelated to risks of awareness</p>
Selective reporting (reporting bias)	Low risk	<p>Quote ROB survey response: "The article described the entire protocol and results of the analysis of both research, as well as secondary outcomes of the study." (email bias survey, see notes)</p> <p>Comment: awareness outcome part of inclusion criteria</p>

Echevarria 1998 (Continued)

Other bias	Low risk	Quote: "No" (email bias survey, see notes)
------------	----------	--

Elhakim 2010

Methods	Study design: randomized parallel groups Study dates: "2008-2009" (email ROB survey)
Participants	Country: Egypt Sex: male Age: 40 to 60 ASA: II-III Procedure: thoracic surgery with one-lung ventilation Study size: 50
Interventions	<p>Randomized portion of anaesthetic: regional anaesthesia: epidural: induction</p> Intervention 1: post-induction/maintenance: epidural: dexmedetomidine 1 µg/kg + bupivacaine 0.5% 30 mg to 40 mg (Group DB), N = 25 Intervention 2: post- induction/maintenance: epidural: bupivacaine 0.5% 30 mg to 40 mg (Group B), N = 25
Outcomes	Primary outcomes: compare analgesic effect, BIS, haemodynamics, blood gases Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Quote: "Two patients (8%) in group B reported possible intraoperative awareness"
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent isoflurane: N₂O no/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance unclear/ADM BIS recorded</p> Anaesthesia induction: fentanyl 3 µg/kg + thiopental 3 mg to 5 mg/kg + pancuronium 0.1 mg/kg; anaesthesia maintenance: isoflurane 0.3% to 0.5% ET; anaesthetic/epidural drugs adjusted BIS: 40 and 60 Time of outcome determination: 24 h, 72 h, 30 days Method of outcome determination: interview Authors report definite and possible cases of awareness. Note Table A1 2 patient awareness reports adjudicated as possible Author responded to the email bias survey on 12 February 2011: "Two (8%) patients in group B reported dreams or nightmares (see Appendix 1). In two patients (cases 1 and 2), the initial BIS after induction of anaesthesia was 37 and 39, the average BIS during the surgical procedure was 44 and 48 and the highest recorded value was 51 and 53, respectively." Survey response: 12 February 2011, Prof. Dr. Mokhtar Elhakim mokhtar.elhakim@gmail.com

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Elhakim 2010 (Continued)

Random sequence generation (selection bias)	Low risk	Quote ROB survey response Dr Elhakim: "Patients were randomly divided into two equal groups using computer-generated random numbers with the closed-sealed envelope, to receive either control or study drugs"
Allocation concealment (selection bias)	Low risk	Quote ROB survey response Dr Elhakim: "To insure the study was blinded, the anaesthetist who prepared or administered the study drugs was not involved in patient care"
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote ROB survey: "Patient, Anesthesiologist, Awareness outcome assessor (interviewer)" Comment: both epidurals contained local anaesthetic. Hence, blinding possible.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Intraoperative awareness in the period from induction of anaesthesia till recovery was assessed by experts who were unaware of the patient group assignments"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "All our patients completed the study." (Email ROB survey)
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote ROB survey: "The main limitation of the present study is the short nature of the study period, and the small number of cases that were reviewed, decreasing the confidence in our results and conclusion"

Ellingson 1977

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Norway Sex: female Age: 18 to 33 years ASA: not given Procedure: forceps delivery Study size: 26
Interventions	Randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia: dissociative vs benzodiazepine Intervention 1: induction: ketamine 2 mg/kg maintenance: after delivery supplemental doses of ketamine (1 mg/kg) as needed, N = 13 Intervention 2: induction: rapid IV injection of diazepam 30 mg maintenance: N ₂ O/O ₂ (6 + 2 litres) to increase analgesia + local anaesthetic for episiotomy and suture, N = 13
Outcomes	Primary outcomes: assessment newborn; awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Comment: awareness reports: ketamine group 4/13 vs diazepam 0/13 7 bad dreams; 3 pleasant dreams in ketamine group: dream content contained peculiar dreams and visual disturbances: sensation of rotating room and presence of many people; one patient felt that her child was a rabbit

Ellingson 1977 (Continued)

Notes

Non-randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia: N₂O yes/narcotics and/or hypnotics bolus/muscle relaxants induction no/maintenance no

Premedication: atropine 0.6 mg

Induction: see interventions; mask cases

Maintenance: none

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Twenty-six patients, in whom forceps delivery was indicated, were allocated at random into two groups of 13 each"
Allocation concealment (selection bias)	Unclear risk	Comment: only above information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: personnel knew treatment group assignment
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Fehr 2001

Methods	Study design: randomized, double-blind, placebo-controlled Study dates: not stated
Participants	Country: Switzerland Sex: male and female Age: 18 to 70 ASA: I-II Procedure: superficial surgical procedures expected to last at least 45 min Study size: 50
Interventions	Randomized portion of anaesthetic: cardiac drug vs placebo

Fehr 2001 (Continued)

Intervention 1: maintenance: clonidine 4 µg/kg, N = 25
 Intervention 2: maintenance: placebo in 0.9% NaCl 100 mL, N = 25

Outcomes	<p>Primary outcomes: clonidine's impact on depth of anaesthesia as measured by BIS, BIS guided propofol concentration; implicit memory word test</p> <p>Secondary: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Quote: "No patient had any free recall either of the presented items or of any other pre- or intra-operative events."</p> <p>Comment: no evidence of explicit or implicit memory</p>	
Notes	<p>Non-randomized portion of anaesthetic: parts of TIVA N₂O no/hypnotic (propofol TCI) infusions MCI/narcotics infusion (remifentanil MCI): muscle relaxants maintenance yes induction/maintenance unclear/ADM BIS recorded</p> <p>Induction: propofol (TCI pump target plasma concentration incremental steps until patient unconscious) + rocuronium 0.6 mg/kg + 40% oxygen</p> <p>Maintenance: propofol (target concentration) + remifentanil MCI between 0.01 µg to 1 µg/kg/min maintain BIS and haemodynamic stability</p> <p>Notes: atropine 0.5 mg was administered if the heart rate fell below 40 beats per min</p> <p>Survey was sent to Donat Spahn, corresponding author, on 14 January 2011. No response</p>	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "...we allocated the patients randomly..."
Allocation concealment (selection bias)	Unclear risk	Comment: see above quote
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Comment: anaesthesiologist: no information provided. However, clonidine or placebo was infused after a steady state was reached post intubation; the use of placebo is a form of blinding
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Forestier 2003

Methods	Study design: randomized parallel groups Study dates: September 1999 to December 2000
Participants	Country: France Sex: male and female Age: mean 61 (SD 9) years ASA: II-III Procedure: coronary artery bypass grafting Study size: 110 intention-to-treat, 111 enrolled
Interventions	<p>Randomized portion of anaesthetic: parts of TIVA: sufentanil TCI</p> <p>Intervention 1: maintenance: sufentanil effect site concentrations (Ce) of 0.5 µg/mL decreased by a third after sternotomy, N = 21</p> <p>Intervention 2: maintenance: sufentanil effect site concentrations (Ce) of 0.75 µg/mL decreased by a third after sternotomy, N = 23</p> <p>Intervention 3: maintenance: sufentanil effect site concentrations (Ce) of 1.0 µg/mL decreased by a third after sternotomy, N = 23</p> <p>Intervention 4: maintenance: sufentanil effect site concentrations (Ce) of 1.25 µg/mL decreased by a third after sternotomy, N = 21</p> <p>Intervention 5: maintenance: sufentanil effect site concentrations (Ce) of 1.5 µg/mL decreased by a third after sternotomy, N = 23</p> <p>propofol and sufentanil, both administered by computer-controlled infusion, were titrated on the bispectral index (BIS) values</p>
Outcomes	<p>Primary outcomes: cardiovascular stability, time to tracheal extubation, patient satisfaction</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Comment: awareness not defined as a dichotomous variable but a continuous one. The authors used a scoring system found in the appendix. Intraoperative dreams scored as awareness; dreams reported but not clear whether intraoperative or postoperative. We had difficulty interpreting this scoring system in regard to the number of patient awareness/dream reports that the author are defining as consistent with recall/dreams. Author contacted for clarification.</p> <p>Comment: score > 3 consistent with recall during surgery; page 342 indicates that the range of scores identified recall/dream in group 1 and 3 (maximal score = 3), group 2 (max = 7), group 4 (max = 4), group 5 (max = 6). We have assumed that Table 6 represents 4 patients who scored > 3 that the authors defined as on the spectrum of recall/dreams; we will revise in updated version if the authors clarify</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of TIVA: N₂O no/narcotics/hypnotics bolus MCI/ muscle relaxants induction yes/maintenance PRN/ADM BIS</p> <p>Induction: sufentanil 5 µg/mL in saline + propofol 1.5 µg/mL (TCI) + intubation BIS < 60 pancuronium 0.1 mg/kg; maintenance: propofol 1 µg/mL (TCI); pancuronium PRN</p> <p>Comment: classify as TIVA as per rules Appendix 8; might be informative to correlate the target blood levels of propofol and sufentanil with wakefulness using IFT</p> <p>Email survey sent 4 February 2013 to d.longrois@chu-nancy.fr - email failed</p> <p>Second email sent on 7 December 2013 for ROB table feedback</p>

Risk of bias

Forestier 2003 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "computer generated random numbers in closed envelopes opened when the patient arrived in the OpR"
Allocation concealment (selection bias)	Unclear risk	Quote: "computer generated random numbers in closed envelopes opened when the patient arrived in the OpR"
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: all protocol violations but one (occurred prior to induction and was excluded from study) were included in outcomes analysis; both postoperative patient deaths occurred after the first awareness interview; therefore, no high-risk awareness patients excluded Author sent characteristics and ROB tables for comment 7 December 2013: d.longrois@chu-nancy.fr . There was no response from this author to our survey.
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Fragen 1981

Methods	Study design: randomized parallel groups Study dates: 1981
Participants	Country: USA Sex: female Age: 19 to 60 ASA: I-II Procedure: short gynaecological operations Study size: 99
Interventions	<p>Randomized portion of anaesthetic: IV: premedication: narcotic vs benzodiazepine vs barbiturate vs placebo</p> Intervention 1: premedication: fentanyl 1.5 µg/kg + induction: midazolam 0.175 mg/kg, N = 25 Intervention 2: premedication: saline 1.5 µg/kg (placebo) + induction: midazolam 0.175 mg/kg, N = 26 Intervention 3: premedication: fentanyl 1.5 µg/kg + thiopental 3.75 mg/kg, N = 25 Intervention 4: premedication: saline 1.5 µg/kg (placebo) + thiopental 3.75 mg/kg, N = 23 Maintenance: doses equal to one-fourth the initial induction dose were given IV when signs of awakening or movement appeared

Fragen 1981 (Continued)

Outcomes Primary outcomes: speed/quality of recovery (awakening characteristics)

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Comment: anterograde amnesia means no recall of intraoperative events as per author survey response (see notes)

(Above from table 3 in study)

Notes **Non-randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia: N₂O yes/narcotics and/or hypnotics bolus/muscle relaxants induction no/maintenance no**

Induction: midazolam 0.175 mg/kg or thiopental 3.75 mg/kg. If 2 of 3 consecutive patients receiving midazolam remained awake after 0.175 mg/kg, the dose was increased by 0.025 mg/kg such that most patients in the midazolam + saline group were induced with a dose of 0.25 mg/kg. Maintenance: N₂O 67%

Time of outcome determination: 24 h postoperative

Method of outcome determination: recall of pictures shown the previous day

Comment: ROB survey Fragen states that **anterograde amnesia** indicates that no patient had recall of intraoperative events

Survey response: 17 January 2011, RJ Fragen r-fragen@northwestern.edu

Third email sent to Robert Fragen rfragen@yahoo.com 4 February 2013

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "According to a random number table, fentanyl...or an equivalent volume of saline was given..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Comment: Dr Fragen's response to ROB survey stated that the anaesthesiologist was blinded to the premedication randomized drug
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Gaitini 1995

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Israel Sex: female Age: 44 ASA: I Procedure: elective caesarean section Study size: 50
Interventions	Randomized portion of anaesthetic: IV parts of volatile agent induction Intervention 1: induction: thiopentone (4 mg/kg), N = 25 Intervention 2: induction: ketamine (1 mg/kg), N = 25
Outcomes	Primary outcomes: correlation between EEG separately for those patients with responsiveness to verbal commands and for those without Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 2 Quote: "None of the patients recalled anything of the surgery" Comment: implicit memory: 13/25 thiopentone, 5/25 ketamine
Notes	Non-randomized portion of anaesthetic: parts of volatile agent: N₂O yes/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance unclear/EEG SEF90 Anaesthesia induction: succinylcholine (1.5 mg/kg) Anaesthesia maintenance: 50% N ₂ O 50% oxygen with end-tidal halothane 0.5 MAC; relationship between EEG (spectral edge frequency 90 - SEF90) and occurrence of awareness Time of outcome determination: 24 h postoperatively Method of outcome determination: interview Comment: IFT used, awareness = wakefulness, recall = explicit memory Quote: "In the thiopentone group, 13 of 25 patients (52%), moved their hands in response to the anaesthetist's instruction, before delivery. The pre-delivery movements in response to command occurred at an average SEF90 value of 18.09 + 3.1 Hz. In the ketamine group, five of 25 patients (24%) moved their hands in response to command before delivery at an average SEF90 of 12.0" No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The patients were assigned by a randomised code to receive either thiopentone (4 mg/kg) or ketamine (1 mg/kg) for induction of anaesthesia"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided

Gaitini 1995 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Gale 2001

Methods	Study design: randomized parallel groups Study dates: June 2000 to January 2001
Participants	Country: Australia Sex: both Age: 44 ASA: I-III Procedure: elective surgery Study size: 40
Interventions	Randomized portion of anaesthetic: parts of IV: induction: MCI vs TCI hypnotic (propofol) Intervention 1: induction: manually controlled infusion of propofol at ~2 mg/kg at 1200 mL/h maintenance: propofol 3 to 10 mg/kg, N = 20 Intervention 2: induction: target controlled infusion of propofol of 3 µg to 8 µg/mL; maintenance: propofol 2 µg to 5 µg/mL, N = 20
Outcomes	Primary outcomes: BIS median absolute performance error; the total dose of propofol infused and the number of changes in target concentration or infusion rate were also noted Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "No patient reported awareness of intraoperative events"
Notes	Non-randomized portion of anaesthetic: parts of IV: N₂O yes/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance yes ADM BIS Anaesthesia induction: midazolam 0.03 µg/kg + fentanyl 2 µg/kg + rocuronium. Anaesthesia maintenance: nitrous 66% propofol, rocuronium, fentanyl as needed; BIS recorded

Gale 2001 (Continued)

 Survey response: 4 February 2011, Kate Leslie kate.leslie@mh.org.au
Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Randomization to TCI or MCI was achieved using random number tables and sealed opaque envelope allocation after consent had been obtained." (email bias survey, see notes)
Allocation concealment (selection bias)	Low risk	Quote: "Sealed opaque envelope allocation after consent had been obtained" (email bias survey, see notes)
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "Patient, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patient, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "40 patients consented and were randomised. There were no attrition after consent or after randomisation. Data for all main outcomes was complete. These details were not reported in the paper." (email bias survey, see notes)
Selective reporting (reporting bias)	Low risk	Quote: "The study protocol is not available, but the protocol is clear from the published report." (email bias survey, see notes) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote: "none" (email bias survey, see notes)

Ghaly 1988

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: England Sex: female Age: 28 Procedure: elective caesarean section for cephalopelvic disproportion Study size: 50
Interventions	Randomized portion of anaesthetic: parts of volatile agent Intervention 1: maintenance: halothane 0.5%, N = 25 Intervention 2: maintenance: Isoflurane 0.75%, N = 25
Outcomes	Primary outcomes: anaesthesia recovery, APGAR scores, blood gases

Ghaly 1988 (Continued)

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Comment: none of the mothers complained of intraoperative dreams or awareness

Notes

Non-randomized portion of anaesthetic: parts of volatile agent N₂O yes/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance yes

Anaesthesia induction: thiopentone (3 mg to 5 mg/kg) + suxamethonium 100 mg. Anaesthesia maintenance: 50:50 N₂O:O₂ + group-specific volatile agent; post-delivery: alfentanil 1.0 mg IV/N₂O 70%/suxamethonium 0.1% rate just sufficient to abolish diaphragmatic movement

Time of outcome determination: first postoperative day

Method of outcome determination: interview

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "They were randomly allocated to either the isoflurane or halothane group"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Ghoneim 2000

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: USA Sex: both Age: 23 ASA: I-III

Ghoneim 2000 (Continued)

Procedure: elective surgery

Study size: 180, 179

Interventions

Randomized portion of anaesthetic: parts of IV vs parts of volatile agent: opioid bolus vs opioid infusion MCI vs volatile agent

Opioid bolus: intervention 1: induction: fentanyl 7.5 µg/kg, after 5 min, thiopental sleep dose (incremental dose until fall asleep), maintenance: 70% N₂O/O₂ and muscle relaxant; 2.5 µg/kg fentanyl supplements as needed, N = 100

Opioid infusion: intervention 2: induction: alfentanil 50 µg/kg + alfentanil 1.5 µg/kg/min; after 90 seconds, thiopental sleep dose (incremental dose until fall asleep), maintenance: 70% N₂O/O₂ and muscle relaxant; alfentanil 1.0 µg to 1.5 µg/kg/min titrated to patient response to noxious stimuli, N = 40

Volatile agent: isoflurane 0.3%: intervention 3: induction: fentanyl 1 µg/kg after 5 min, thiopental sleep dose (incremental dose until fall asleep), maintenance: 70% N₂O/O₂ and muscle relaxant; isoflurane 0.3% ET + fentanyl up to 1 µg/kg/h as needed, N = 16

Isoflurane 0.7%: intervention 4: similar to the preceding group except that isoflurane maintained 0.7% ET, N = 24

Outcomes

Primary outcomes: implicit memories

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4

Quote: "Six patients showed explicit recall of intraoperative events: All received the opioid bolus regimen"

Quote: "**None** of the patients gave evidence of **awareness** when they were asked initially the **simple four questions** mentioned above. However, with further **prompting cue questions** they gave evidence of awareness"

Notes

Non-randomized portion of anaesthetic: parts of IV vs parts of volatile agent N₂O yes/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance yes/ADM AER recorded

Anaesthesia induction: thiopental sleep dose + nitrous 70% with muscle relaxant

Anaesthesia maintenance: N₂O 70% in oxygen

Comment: 10 tapes (5 of each story) were randomized to maintain blinded conditions and assessed the AER. Each story contained target words

Comment: opioid anaesthesia is a light anaesthetic with more dreams consistent with Utting's view

Comment: extended interview identifies more patient awareness reports than the standardized interview (p133 see for interview protocol)

Time of outcome determination: at most 4 days postoperatively

ROB survey: we emailed on 16 June 2015

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "decided to allocate most patients to the opioid bolus regimen (N=100) and a lesser number to the opioid infusion regimen (N=40), isoflurane 0.7% (N=24) and isoflurane 0.3% regimen (N=16), using randomization tables"

Ghoneim 2000 (Continued)

Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor: "Ten tapes (five of each story) were randomized to maintain blinded conditions for the investigators, who interviewed the patients postoperatively and assessed the AER...The research assistant was also "blinded" to the method of anesthesia and to the specific story which was played"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "one male receiving isoflurane 0.7% had to be eliminated from the AER analyses because of equipment failure"
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Ghosh 2008

Methods	Study design: randomized parallel groups Study dates: Quote: "2005-06" (email bias survey, see notes)
Participants	Country: India Sex: 86 male, 4 female Age: mean 29.1 ASA: I-II Procedure: peripheral nerve repair surgery Study size: 90
Interventions	Randomized portion of anaesthetic: cardiac drug vs placebo Intervention 1: premedication (1 h before surgery): group I: metoprolol 100 mg, N = 30 Intervention 2: premedication: Group II: placebo control, N = 30 Intervention 3: premedication: Group III: clonidine 200 µg, N = 30
Outcomes	Primary outcomes: anaesthetic consumption, haemodynamic effects Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "... none of the patients had free recall of any intraoperative event"
Notes	Non-randomized portion of anaesthetic: parts of IV: N₂O no/narcotics/hypnotics bolus MCI/muscle relaxants induction no/maintenance no/ADM: BIS

Ghosh 2008 (Continued)

Anaesthesia induction: fentanyl (2 µg/kg) + propofol until loss of response; anaesthesia maintenance: fentanyl infusion (1 µg/kg/h) + propofol infusion titrated to BIS 40 to 60 + N₂O in oxygen (2:1) **LMA, BIS 40 to 50**

Time of outcome determination: 2 h postoperatively

Method of outcome determination: interview

Survey response: 17 February 2011, Parmod Kumar Bithal bithal.parmod@gmail.com

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "After we had obtained approval from the institutional ethics committee and informed consent from the patients, they were divided into three groups, of 30 each, by the use of computer-generated block randomisation numbers" Quote: "Computer generated randomisation sequence of 90 patients in 3 blocks of 30 patients each" (email bias survey, see notes)
Allocation concealment (selection bias)	Low risk	Quote: "Identical opaque sealed envelopes were used for the study drugs. All envelopes were labelled and contained drug according to sequence generated by computer. Envelopes were prepared by person not a part of investigating team" (email bias survey, see notes)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Patient, Anaesthesiologist, Awareness outcome assessor"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patient, Anaesthesiologist, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "All patients completed the study and none was excluded from final analysis" (email bias survey, see notes)
Selective reporting (reporting bias)	Low risk	Quote: "All primary and secondary end points have been reported" (email bias survey, see notes) "No results however insignificant have been selectively withheld from publication" (email bias survey, see notes) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote: "None that we are aware of in the post operative period all" (email bias survey, see notes)

Girardi 1994

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Italy

Girardi 1994 (Continued)

Sex: female and male
 Age: 40 to 54
 ASA: I
 Procedure: saphenectomy surgery
 Study size: 51

Interventions
Randomized portion of anaesthetic: volatile agent types

Intervention 1: group 1 (26 patients), 5% isoflurane in air, by **mask**; maintenance: group 1, 2% isoflurane in air

Intervention 2: group 2 (25 patients), 3% isoflurane and 60% N₂O, by **mask**; maintenance: group 2, 1.2% isoflurane and 60% N₂O

Outcomes

Primary outcomes: control the depth, the quality of recovery, recovery time
 Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Comment: no patient recall of surgical events

Notes
Non-randomized portion of anaesthetic: parts of volatile agent: N₂O no/narcotics/hypnotics bolus/muscle relaxants induction no/maintenance no/ADM EEG and Evans test

Induction: thiopental (3.5 mg/kg), atracurium (0.6 mg/kg) IV

Comment: manual mask induction then ventilator with equipotent isoflurane dose; goal was to use inhalational agents only; authors conclude that **pure inhalational** anaesthesia achieves the goals of anaesthesia as long as you use the **appropriate concentrations**

Quote: "Isoflurane anesthesia in air, in adequate concentrations, provides a sufficient level of analgesia, hypnosis, amnesia, without clinical side effects"

Standardized interview 1 h after anaesthesia and 24 h later

Both groups: EEG compressed spectral array, and clinical signs of pain by Evans test

Translated by AM

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Patients were randomly assigned to: group..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided; no ROB survey data
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: no information provided
Blinding of outcome assessment (detection bias)	Unclear risk	Comment: no information provided

Girardi 1994 (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information provided

Gokce 2009

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Turkey Sex: both Age: 29 ASA: I Procedure: elective septorhinoplasty Study size: 40
Interventions	Randomized portion of anaesthetic: cardiac drug + IV vs IV anaesthesia: hypotensive anaesthesia: induction/maintenance Comment: remifentanil-propofol induced hypotensive anaesthesia (group RP) or remifentanil-propofol-esmolol induced hypotensive anaesthesia (group RP-E): hypotensive technique with safe lower limits of 80 systolic and mean 50 mmHg , both groups titrated to SNAP index score 40% to 60% Intervention 1: No esmolol : induction: remifentanil bolus mg/kg + infusion 0.1 µg to 0.5 µg/kg/min; maintenance: remifentanil 0.1 µg to 0.5 µg/kg/min (control), N = 20 Intervention 2: Yes esmolol : as intervention 1 plus: induction: esmolol bolus 100 µg to 300 µg/kg/min infusion; maintenance: esmolol 100 µg to 300 µg/kg/min, N = 20
Outcomes	Primary outcomes: recovery times, haemodynamic effects, anaesthetic consumption Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Comment: 1 awareness patient in group RP
Notes	Non-randomized portion of anaesthetic: parts of IV N₂O no/narcotics/hypnotics bolus MCI/muscle relaxants induction yes/maintenance unclear PRN/ADM: SNAP (40% to 60%) Premed: midazolam (0.03 mg/kg IV); induction: propofol bolus (2 mg to 2.5 mg/kg) + rocuronium (0.6 mg/kg) + O ₂ 50% + rocuronium (table 2); maintenance: propofol 4 mg to 10 mg/kg/h+ remifentanil infusion (0.1 µg to 0.5 µg/kg/h) Quote: "Depth of anaesthesia was measured by depth of anaesthesia (SNAP index score) (Nicolet Biomedical, VIASYS Healthcare, Madison, Wisconsin, USA)" Time of outcome determination: in recovery room

Gokce 2009 (Continued)

Method of outcome determination: interview time NA

Dr Lale Karabrysk, email: karabiyik@gazi.edu.tr

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The patients were equally assigned via computer-generated random numbers"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor: "After the extubation, the patients were transferred to the postanaesthesia care unit (PACU), where further recordings were carried out by an independent observer blinded to the anaesthetic regimen"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Goto 2000

Methods	Study design: randomized parallel groups Study dates: Quote: "From July 7 1998 to January 25, 1999" (email bias survey, see notes)
Participants	Country: Japan Sex: female Age: 38 to 56 ASA: I and II Procedure: elective total abdominal or vaginal hysterectomy Study size: 20
Interventions	Randomized portion of anaesthetic: volatile agent types Intervention 1: maintenance: xenon: 56% xenon (0.8 MAC), N = 10 Intervention 2: maintenance: isoflurane: 1.0% to 1.5% in 6 L/min flow of oxygen, N = 10 BIs in both groups
Outcomes	Primary: concentration of anaesthetic: BIS SEF95 values

Goto 2000 (Continued)

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Comment: zero patient awareness reports; 4 patients had **wakefulness** at BIS 50 with xenon post end surgery

Notes

Non-randomized portion of anaesthetic: regional anaesthesia: epidural: induction/maintenance: parts of volatile agent: N₂O no/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance unclear/ADM: BIS recorded

Anaesthesia induction: **epidural** 10 mL mepivacaine 1.5% with 1:200,000 epinephrine, propofol 2.5 mg/kg IV + 5% sevoflurane + vecuronium 10 mg IV; anaesthesia maintenance: see interventions; **muscle relaxant** reversed at end surgery

Time of outcome determination: 2 h postoperative

Method of outcome determination: interview

Comment: after surgery...asked to open their eyes and squeeze the anaesthetists hand. For xenon, in some patients this occurred at BIS < 50. None remembered wakefulness

Survey response: 2 March 2011, Takahisa Goto takigoto@yokohama-cu.ac.jp

Survey questionnaire 25 February 2011

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The envelop method. Before starting the whole study, we made 11 and 9 cards with Xe or Iso, respectively, and put each card to an identical envelop and sealed. Before induction of anaesthesia, the anaesthetist selected one envelop, opened it, and gave either xenon or isoflurane according to what the card said. We made a little more cards of xenon than those of isoflurane simply because xenon was a newer drug" (email survey)
Allocation concealment (selection bias)	Low risk	Quote: "As described in the randomized method, we decided allocation just before induction of anesthesia. After randomization, we did not tell the participants whether they would be receiving xenon or isoflurane. Similarly, the investigator who enrolled the participants had no means to know in advance if they would be allocated to xenon or isoflurane. The anesthetists who actually conducted anesthesia were NOT blinded to the anesthetics given. Blinding was virtually impossible" (email survey)
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "Patient, Awareness outcome assessor (interviewer)" (email survey)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patient, Awareness outcome assessor (interviewer)" (email survey)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "No attrition or exclusions were made during the study" (email survey)
Selective reporting (reporting bias)	Low risk	Quote: "Yes, our study is free of selective reporting. We reported all the results we obtained" (email survey)

Goto 2000 (Continued)

Comment: awareness outcome part of inclusion criteria

Other bias	Low risk	Quote: "I cannot think of any other potential bias" (email survey)
------------	----------	--

Gruenewald 2007

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Germany Sex: female Age: 33, 38 ASA: I and II Procedure: elective gynaecological laparoscopy Study size: 72
Interventions	Randomized portion of anaesthetic: TIVA ADM (M-entropy) vs SCP Intervention 1: standard clinical practice Intervention 2: M-entropy monitoring 40 and 60
Outcomes	Primary outcomes: consumption of anaesthetics, recovery times Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Comment: 1 patient in SCP reported awareness
Notes	Non-randomized portion of anaesthetic: parts of TIVA: N₂O no/narcotics/hypnotics bolus MCI/ muscle relaxants induction yes/maintenance unclear/ADM BIS recorded Anaesthesia induction: propofol (2 mg/kg) + remifentanyl (0.3 µg to 0.5 µg/kg/min) + rocuronium (0.6 mg/kg); maintenance: propofol + remifentanyl Comment: blinding anaesthetist, learning bias according to 2 opposing views of Roizen and Lindholm Time of outcome determination: first postoperative day Method of outcome determination: interview Email: gruenewald@anaesthesie.uni-kiel.de ; ROB survey - we emailed on 11 April 2015

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Randomisation to the two treatment groups (standard practice or entropy) was done by opening a sealed envelope"
Allocation concealment (selection bias)	Unclear risk	Quote: "Randomisation to the two treatment groups (standard practice or entropy) was done by opening a sealed envelope"

Gruenewald 2007 (Continued)

		Comment: unclear if envelope opaque as per Cochrane guidelines (Higgins 2011)
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: both groups used BIS and M-entropy; in SCP group both were blinded and in experimental group entropy was not blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor: "On the first postoperative day, all patients were asked by a blinded anaesthetist if they had any memory or awareness during different stages..."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "All patients were included into the final analysis"
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Grundmann 2001

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Germany Sex: male and female Age: 23 to 65 ASA: I and II Procedure: elective laparoscopic cholecystectomy Study size: 50
Interventions	Randomized portion of anaesthetic: parts of TIVA vs parts of volatile agent Intervention 1: maintenance: Group R/P: propofol 4 mg/kg/h, N = 25 Intervention 2: maintenance: Group R/D: desflurane 3% ET, N = 25
Outcomes	Primary outcomes: haemodynamic responses, recovery profile Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "None of the patients showed any signs of wakefulness in response to surgical procedures or had postoperative recall of intraoperative events"
Notes	Non-randomized portion of anaesthetic: parts of TIVA vs parts of volatile agent N₂O no + infusion of narcotics MCI + muscle relaxant(s) induction yes/maintenance unclear Premed: 10 mg diazepam; induction: remifentanil infusion MCI (0.5 µg/kg/min) until patient felt dazed + propofol IV (2 mg/kg) + atracurium (0.5 mg/kg) + ventilation oxygen 2 L/min in air Anaesthesia maintenance: remifentanil infusion MCI reduced 0.25 µg/kg/min

Grundmann 2001 (Continued)

Comment: see Dryad author's definition of anaesthesia

Time of outcome determination: after recovery

Method of outcome determination: interview

ROB survey. We emailed on 11 April 2015

 Dr. Grundmann email: aiugru@krzsun.med-rz.uni-sb.de
Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Fifty patients (ASA I–II, 23–65 yr) were randomly assigned"
Allocation concealment (selection bias)	Unclear risk	Comment: inadequate information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor: "...the patients were directly transferred to the postanesthesia care unit (PACU), where further recordings were done by an independent, blinded observer, who was unaware of the administered anaesthesia regimen"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Gupta 1992

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Sweden Sex: both Age: 17 to 49, mean: 33.2 ± 10, 27.5 ± 6 ASA: I and II Procedure: outpatient arthroscopic procedures of the knee Study size: 30
Interventions	Randomized portion of anaesthetic: volatile agent types: IV induction hypnotics Intervention 1: induction: thiopentone 5 mg to 6 mg/kg, N = 14

Gupta 1992 (Continued)

Intervention 2: induction: propofol 2 mg to 3 mg/kg, N = 16

Outcomes	Primary outcomes: psychomotor recovery; differences between isoflurane and propofol groups at each measurement, dream frequency Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: assessment for dreams and awareness was in methods. Results section reports only dreams does not mention awareness; awareness reported
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent: N₂O no/narcotics/hypnotics bolus/muscle relaxants induction no/maintenance no</p> <p>Anaesthesia induction: alfentanil 0.25 mg; maintenance: isoflurane (0.5% to 2%) in oxygen and air + alfentanil 0.25 mg plus additional doses every 15 min; spontaneous breathing with mask; ventilation was controlled if there was apnoea exceeding 30 s</p> <p>Time of outcome determination: prior to discharge</p> <p>Method of outcome determination: interview</p> <p>No email address available for ROB survey</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The patients were randomly divided into two groups"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided Quote: "The patients were interviewed prior to discharge as to whether they had any dreams or were aware during the procedure..."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Gurman 1994

Methods	Study design: randomized parallel groups Study dates: "year of 1992" (email bias survey, see notes)
---------	--

Gurman 1994 (Continued)

Participants	Country: Israel Sex: male and female Age: mean 18 to 70 ASA: I-II Procedure: surgical procedures lasting at least 30 min Study size: 48 enrolled; 43 completed study
Interventions	<p>Randomized portion of anaesthetic: parts of TIVA vs parts of volatile agent</p> Intervention 1: maintenance: propofol varying between 3 mg to 6 mg/kg/h, N = 24 Intervention 2: maintenance: isoflurane 1% to 1.75%, N = 24 Both isoflurane and propofol dosage as well as the amount of fluids administered were adjusted in order to maintain a SEF between 8 Hz and 12 Hz and mean blood pressure and heart rate within normal limits
Outcomes	Primary outcomes: haemodynamic effects, recovery effects, differences anaesthetics consumed, SEF variation Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "No patients in either group showed any evidence of awareness during maintenance of anaesthesia (by raising the isolated arm) or by postoperative recall of intra-operative events"
Notes	<p>Non-randomized portion of anaesthetic: parts of TIVA vs parts of volatile agent N₂O yes/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance PRN/ADM: SEF recorded</p> Premedication: oral diazepam 5 mg to 10 mg Induction: vecuronium 1 mg + fentanyl 0.2 mg + thiopentone 5 mg/kg + suxamethonium 5 mg/kg Maintenance: N ₂ O/O ₂ 2:1 litre/min + vecuronium and fentanyl when needed Author responded to the email bias survey on 25 January 2011 Survey response: 25 January 2011, Gabriel M. Gurman MD gurman@bgu.ac.il Emailed author on 8 December 2013 for feedback about characteristics table and ROB domains; responded 12 December 2013: "I have nothing to comment"

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	Quote: "the patients were randomised into two groups according to the last digit of their identity number" Quote: "last digit ID number" (email survey)
Allocation concealment (selection bias)	Low risk	Quote: "There was only one of us (MS) who dealt with randomisation and he did not take part to the results analysis" (email survey)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Patient, Anaesthesiologist, Awareness outcome assessor (interviewer)" (email survey)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Above

Gurman 1994 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Unclear risk	<p>Comment: (5/48) 11% of study group dropped out but unclear what that distribution was across the 2 groups</p> <p>Quote: "Three were eventually excluded due to lack of complete data. Two other cases were eliminated because of the need for naloxone to treat prolonged postoperative respiratory depression"</p> <p>Comment: naloxone group not at high risk of awareness</p> <p>Comment: unclear risk because we cannot determine if imbalanced exclusions</p> <p>If 5 exclusions in 1 group that is a significant difference, Peto OR 0.11 (0.02 to 0.70)</p>
Selective reporting (reporting bias)	Low risk	<p>Quote: "The study protocol is not available but it is clear that the published report includes all expected outcomes, including those that were pre-specified." (email survey)</p> <p>Comment: awareness outcome part of inclusion criteria</p>
Other bias	Low risk	<p>Quote: "No problems like those described above" (email bias survey, see notes)</p>

Hachero 2001

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Spain Sex: female Age: 18 to 65 ASA: I-II Procedure: major gynaecological surgery Study size: 40
Interventions	<p>Randomized portion of anaesthetic: ADM (BIS) vs SCP</p> <p>Intervention 1: BIS: maintenance: propofol doses to keep BIS maintained between 40 to 60, N = 20</p> <p>Intervention 2: standard clinical practices (SCP): maintenance: propofol 10 mg/kg/h for 5 min, 8 mg/kg/h 5 min, 6 mg/kg/h until end + fentanyl 150 µg or 75 µg when needed, N = 20</p>
Outcomes	<p>Primary outcomes: anaesthetic consumption, recovery times, haemodynamic effects, BIS changes</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Quote: "En ningun caso las pacientes refirieron recuerdo explicito intraoperatorio." Translation: "In no case patients reported explicit recall intraoperatively"</p>
Notes	<p>Non-randomized portion of anaesthetic: TIVA/N₂O no/induction: muscle relaxants: infusions propofol/induction yes/maintenance yes</p>

Hachero 2001 (Continued)

Premedication: midazolam 0.05 mg/kg + atropine 0.01 mg/kg + dehydrobenzperidol 40 µg/kg + fentanyl 2 µg/kg

Induction: propofol 2 mg to 2.5 mg/kg + mivacurium 0.2 mg/kg

Maintenance: mivacurium 0.5 mg/kg/h + fentanyl 75 µg to 150 µg PRN BP/h > 20% baseline

Translated by Brett Smith

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Mediante una tabla generada por ordenador las pacientes fueron asignadas de formas aleatoria a dos grupos..." Translation: "Using a computer-generated table patients were randomly assigned to two groups..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Translation: assessor: "upon arriving in the PACU a nurse that was unaware of the anaesthetic group that they belonged to about their pain...The possibility of intraoperative awareness was investigated in all patients through an interview given 24 hours post-op"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Hackner 2003

Methods	Study design: randomized parallel groups Study dates: NA
Participants	Country: Germany Sex: both Age: 18 to 73 ASA: I-III Procedure: elective panendoscopy, microlaryngoscopy, or tonsillectomy Study size: 44 enrolled, 43 completed study
Interventions	Randomized portion of anaesthetic: parts of TIVA

Hackner 2003 (Continued)

Intervention 1 maintenance: **propofol-pronounced**: propofol **100 µg/kg/min**, remifentanil **0.15 µg/kg/min**, N = 22

Intervention 2 maintenance: **remifentanil-pronounced**: propofol **50 µg/kg/min**, remifentanil **0.45 µg/kg/min**, N = 22

Outcomes	<p>Primary outcomes: halogenated agents and the acid base status, cbc chemistry, urine analysis etc; values of MAP, HR, SpO₂, and BIS</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>"...No intra-operative awareness with recall was reported..."</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of TIVA: N₂O no/narcotics/hypnotics bolus MCI/muscle relaxants induction yes/maintenance PRN/ADM: BIS</p> <p>Anaesthesia induction: remifentanil (0.4 mg/kg ± 1 over 30 sec), followed by propofol (2.0 mg/kg ± 1 over 30 sec) and mivacurium (0.2 mg/kg ± 1); maintenance: propofol: remifentanil combo (100 µg/kg/min:0.15 µg/kg/min) or (50 µg/kg/min:0.45 µg/kg/min), mivacurium PRN surgical requirements. BIS > 55</p> <p>Time of outcome determination: 24 h postoperative</p> <p>Method of outcome determination: interview recovery room and on the day after anaesthesia</p> <p>ROB survey. We emailed on 12 April 2015</p> <p>Email: o.detsch@lrz.tum.de</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Before induction, patients were randomised to receive either..."
Allocation concealment (selection bias)	Unclear risk	Comment: insufficient information provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: anaesthesiologist/assessor: "The anaesthetist providing anaesthesia and the interviewer recording all variables were blinded to the drug concentrations"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: anaesthesiologist/assessor: "The anaesthetist providing anaesthesia and the interviewer recording all variables were blinded to the drug concentrations"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "one patient was excluded because of changes in the surgical procedure"
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Haimeur 1997

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Maroc Sex: female Age: mean 27 to 30 in 3 groups ASA: NA Procedure: caesarean section surgery Study size: 30
Interventions	Randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia vs volatile agent types Intervention 1: induction: thiopental 4 mg/kg maintenance: N₂O /O ₂ 50% + halothane 0.5%; N = 10 Intervention 2: induction: ketamine 1 mg/kg maintenance: N₂O /O ₂ 50%; N = 10 Intervention 3: induction: ketamine 1 mg/kg maintenance: N₂O /O ₂ 50% + halothane 0.5%; N = 10
Outcomes	Primary outcomes: wakefulness , haemodynamic parameters, time to loss of consciousness between thiopental and ketamine, incisions to delivery time, APGAR scores Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 5 Comment: positive wakefulness: group 1: 4 events, group 2: 2 events, group 3: 1 event; 2 patient awareness reports and 1 postoperative nightmare report; see Dryad for translation
Notes	Non-randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia vs volatile agent types: N₂O yes/narcotics and/or hypnotics bolus/muscle relaxants induction yes/maintenance unclear/ADM: response to command (IFT) Vecuronium for intubation Isolated forearm technique Translated by Anthony Messina No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "For induction...are randomly distributed in one of the following three groups "
Allocation concealment (selection bias)	Unclear risk	Comment: inadequate information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: inadequate information provided
Blinding of outcome assessment (detection bias)	Unclear risk	Comment: inadequate information provided

Haimeur 1997 (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information provided; no ROB survey data

Haram 1981

Methods	Study design: randomized parallel groups Study dates: not stated	
Participants	Country: Norway Sex: women Age: 19 to 44 ASA: not given Procedure: elective caesarean section Study size: 97 enrolled, 82 completed surgery, 79 had complete data for analysis In 2 groups of 43 and 39 women, respectively	
Interventions	Randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia Intervention 1: thiopentone induction 3 mg/kg + additional bolus doses (25 mg to 50 mg) if needed; N = 43 Intervention 2: diazepam induction 0.3 mg/kg + additional bolus doses (5 mg to 15 mg); N = 39	
Outcomes	Primary outcomes: APGAR scores, acid-base balance; times from injection of the induction agent to sleep; loss of ciliary reflexes Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Comment: Table 3: group 1: awareness 5: awareness with noise 1, pain 1, unpleasant dreams 3, no awareness 35 = 40 = N Group 2: awareness 0: awareness with noise 0, pain 0, unpleasant dreams 0, no awareness 39 = N; the term wakefulness is used to describe postoperative memory of intraoperative events, awareness	
Notes	Non-randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia: N₂O yes/narcotics and/or hypnotics bolus/muscle relaxants induction yes/maintenance yes Induction: atropine 0.6 mg + N ₂ O/O ₂ 4:2 + boluses hypnotic sedative drugs Maintenance: suxamethonium 50 mg to 100 mg + N ₂ O/O ₂ 4:2 + analgesic (pethidine 50 mg to 100 mg IV) after delivery No email address available for ROB survey	
Risk of bias		
Bias	Authors' judgement	Support for judgement

Haram 1981 (Continued)

Random sequence generation (selection bias)	Unclear risk	Quote: "They were randomly allocated to two groups"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "The anaesthetist providing anaesthesia and the interviewer recording all variables were blinded to the drug concentrations"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor: "On first postoperative day, the patients were interviewed by one of the authors (KH), who was not aware which induction agent had been used"
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	<p>Comment: enrolled 97 minus 15 excluded due to obstetrical criteria; 97 started surgery, 82 completed surgery; 79 had complete study data for analysis; hence (18/97) 19% dropouts but no dropout for reasons that defined them as high-risk awareness cases: dropouts for SCP data (unable to collect); no awareness interview dropouts</p> <p>Comment: inadequate information to determine if exclusions were balanced between groups</p>
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Hug 1988

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: USA Sex: female/male Age: 58 to 61 ASA: NA Procedure: CABG Study size: 19
Interventions	Randomized portion of anaesthetic: TIVA (MCI) Alfentanil infusion rate (prime/maintenance (µg/kg/min)) Intervention 1: (N = 6) 60/4.5 µg/kg/min Intervention 2: (N = 4) 60/5.4 µg/kg/min Intervention 3: (N = 5) 72/6.6 µg/kg/min Intervention 4: (N = 4) 86/7.8 µg/kg/min
Outcomes	Primary: outcomes: plasma concentration of alfentanil, somatic and haemodynamic responses Secondary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Hug 1988 (Continued)

Quote: "No patient had recall of intraoperative events when interviewed 1 to 3 days after surgery"

Notes

Non-randomized portion of anaesthetic: TIVA/N₂O no/infusion of narcotics (MCI) + muscle relaxants induction yes//maintenance yes PRN

Lorazepam 0.08 mg/kg (intervention 1), 0.04 mg/kg (intervention 2 to 4), PO premed

Metocurine 0.05 mg/kg while breathing 100% oxygen, suxamethonium and PRN after intubation

Comment: no MR use after induction to view somatic response

Quote: "The highest plasma concentration of alfentanil to prevent response to a stimulus other than tracheal intubation was different between the two studies ($P < 0.05$). We conclude that alfentanil alone is insufficient to suppress haemodynamic and somatic motor responses to noxious stimulation during CABG and that the role of premedication is significant..." "Rigidity on induction of anaesthesia occurred in 14 of the 19 patients and was promptly relieved by suxamethonium. Suxamethonium was again required in 12 patients to facilitate opening of the sternum"

No email address

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "chosen randomly from one of three different options (table II)"
Allocation concealment (selection bias)	Unclear risk	Comment: inadequate information
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: inadequate information
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: inadequate information
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no dropouts
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information

Hung 1992

Methods

Study design: randomized parallel groups
 Study dates: not stated

Participants

Country: Canada, USA
 Sex: male
 Age: aged: 30 to 79
 ASA: I-II
 Procedure: not specified

Hung 1992 (Continued)

Study size: 26

Interventions	Randomized portion of anaesthetic: IV hypnotic (thiopental) (TCI) infusion serum levels Intervention 1: thiopental serum concentrations 10 µg to 30 µg/mL 5/26 had noxious stimulation because they were arousable and responded to verbal command Intervention 2: higher, randomly assigned target serum concentration of 40 µg to 90 µg/mL tracheas of 6 could not be intubated due to inability to intubate without muscle relaxants	
Outcomes	Primary: outcomes: association thiopental serum concentrations and clinical signs anaesthetic depth as defined by EEG and several perioperative stimuli (verbal command, tetanic nerve stimulation, trapezius muscle squeeze, and laryngoscopy) Secondary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "When interviewed 24 h postoperatively, none of the subjects could recall the events that occurred during the study"	
Notes	Non-randomized portion of anaesthetic: ADM EEG recorded No post-intubation anaesthetic technique described: classified as "other" technique Quote: "A positive response was recorded if purposeful extremity movement or coughing was observed the probability of no movement to each stimulus was characterized using logistic regression. The biphasic thiopental concentration-EEG relationship and the isoelectric EEG at the high serum thiopental concentrations needed to prevent purposeful movement responses limit the utility of the EEG as a measure of anaesthetic depth when thiopental is used alone ...The movement responses may be associated with spinal (brain stem in the case of laryngoscopy and intubation) reflexes to peripheral noxious stimuli. However, they also may be associated with light anaesthesia and inadequate cortical CNS suppression, since most of these movement responses were associated with an increase of mean arterial pressure and heart rate. It is not possible for us to separate the cortical from spinal components of a movement or cough response....Laryngoscopy followed by intubation can be considered the most noxious stimulus that has been quantitated with available methodology in humans. Because of the concurrent peri operative use of anaesthetic drugs with specific actions... traditional clinical signs of anaesthetic depth such as movement and hemodynamic responses to noxious stimuli become less interpretable " No email survey	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The first randomly assigned target serum concentration of 10-30 mcg/ml was maintained ..."
Allocation concealment (selection bias)	Unclear risk	Comment: inadequate information
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: inadequate information
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: inadequate information

Hung 1992 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no dropouts
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information

Ibraheim 2008

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Saudi Arabia Sex: both Age: 41, 39 ASA: I-II Procedure: laparoscopic gastric banding Study size: 30
Interventions	Randomized portion of anaesthetic: volatile agent types/ADM Intervention 1: maintenance: sevoflurane BIS between 40 to 60 , N = 15 Intervention 2: maintenance: sevoflurane SCP, N = 15
Outcomes	Primary outcomes: recovery times, anaesthetic consumption Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "There was no recall or dreaming reported by any patient in Recovery room or 24 hours after surgery in both groups"
Notes	Non-randomized portion of anaesthetic: parts of volatile agent: N₂O no/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance yes Induction: fentanyl 2 µg/kg + propofol 1.5 mg to 2.0 mg/kg + succinylcholine 1.0 mg to 1.5 mg/kg; maintenance: sevoflurane 2% in 2 L/min + fentanyl 100 µg + atracurium Time of outcome determination: discharge from recovery room and 24 h postoperative Method of determination: interview Email: osamaibraheim@yahoo.com ; awaiting email bias survey response

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Patients were randomly assigned to two groups"

Ibraheim 2008 (Continued)

Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "BIS Group: the anaesthesiologist had access to the monitor and adjusted the concentration of sevoflurane to achieve a target BIS in the range 40-60. Non BIS (control group) the anaesthesiologist adjusted the sevoflurane concentration purely according to the clinical signs" Comment: anaesthesiologists were not blinded to inhalational agent nor to SCPs
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor: "Blinded study personnel recorded..."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Inoue 2005

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: USA Sex: men and women Age: 37 to 76 ASA: I-II Procedure: cervical spine surgery Study size: 75
Interventions	Randomized portion of anaesthetic: parts of IV vs parts of volatile agent: TCI vs supplemental volatile vs solely volatile Intervention FP 1: induction and maintenance: fentanyl 50 µg intermittent boluses + propofol 1.5 µg to 3.5 µg/mL TCI , both titrated against clinical signs, N = 25 Intervention Fs 2: induction: fentanyl 100 µg + propofol 1 mg to 3 mg/kg; maintenance: fentanyl 50 µg intermittent boluses + supplementary sevoflurane (0.5% to 1.0% ET), N = 25 Intervention S 3: induction: fentanyl 100 µg + propofol 1 mg to 2 mg/kg; maintenance: solely with sevoflurane 1.5% to 2.5% ET, N = 25
Outcomes	Primary outcomes: the time to extubation, bucking scores, and pain scores Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 0 Quote: "There were no light anesthesia responses such as hazardous involuntary movements, bucking, or awareness during surgery"

Inoue 2005 (Continued)

Comment: no awareness events; all patients were paralysed for induction and a portion of the sample were paralysed during maintenance

Notes

Non-randomized portion of anaesthetic: parts of IV/N₂O yes vs parts of volatile agent: N₂O yes/ muscle relaxants induction yes/maintenance yes

Premedication: 25 mg hydroxyzine + 0.5 mg atropine; anaesthesia induction: vecuronium; maintenance: vecuronium as needed + nitrous 60%, O₂ 40%

Time of outcome determination: 24 h postoperative
 Method of outcome determination: interview

Emailed author 27 December 2010 for risk of bias information; no response received

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Patients were randomly assigned into one of three groups"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor: "the extent of awareness and readiness for the neurological examination were assessed using a predetermined scoring scale by a nurse observer blinded to the method of anesthesia"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Jensen 1995

Methods	Study design: randomized parallel groups Study dates: 1987 (email bias survey, see notes)
Participants	Country: Denmark Sex: female Age: mean 42 to 47 ASA: I-II Procedure: major elective gynaecologic surgery Study size: 80, 74 complete data postoperative
Interventions	Randomized portion of anaesthetic: parts of: neurolept (IV) vs TIVA reversal flumazenil vs no reversal flumazenil ±N₂O

Jensen 1995 (Continued)

TIVA using midazolam-alfentanil intervention 1 and 2, with or without reversal with flumazenil vs standardized neurolept anaesthesia with **N₂O**, intervention 3

Intervention 1: **TIVA** induction: alfentanil 50 µg/kg + midazolam 150 µg/kg + alfentanil 10 µg/kg/min 10 min, 2 µg/kg/min 20 min + midazolam 8 µg/kg/min 10 min, 3.3 µg/kg/min 20 min; maintenance: alfentanil 1 µg/kg/min + midazolam 2 µg/kg/min + atracurium 7 µg/kg/min; **reversal flumazenil 2 mg**, N = **20**

Intervention 2: **TIVA** induction: alfentanil 50 µg/kg + midazolam 150 µg/kg + alfentanil 10 µg/kg/min 10 min, 2 µg/kg/min 20 min + midazolam 8 µg/kg/min 10 min, 3.3 µg/kg/min 20 min; maintenance: alfentanil 1 µg/kg/min + midazolam 2 µg/kg/min + atracurium 7 µg/kg/min; 20 mg, N = **20 - 1 = 19**

Reversal placebo

Intervention 3: **neurolept** induction: fentanyl 5 µg/kg + droperidol 250 µg/kg + thiopental 4 mg/kg and supplemental doses 50 mg; maintenance: **N₂O** in 33% O₂ + fentanyl and droperidol doses as needed, N = **40 - 1 = 39**

Outcomes

Primary outcomes: recovery characteristics

Quote: "Recovery after Midazolam-Alfentanil Anaesthesia with and without Reversal with Flumazenil, and Standardized Neurolept Anaesthesia"

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4

Comment: the dynamic nature of postoperative memory was demonstrated by a patient who had immediate postoperative recall of skin incision as a "burning sensation", but the next day the patient "failed to recall the event." This is evidence that RCTs that interview patients after the immediate post-operative period underestimate the frequency of awareness

Comment: naloxone needed in 11/40 patients in TIVA groups with and without reversal and 0/40 patients in neurolept group. Significantly more naloxone used in TIVA groups 1 and 2 (11/40) vs neurolept group (0/40), Peto OR 9.87 (2.79 to 34.97)

Notes

Non-randomized portion of anaesthetic: parts of: neurolept (IV) vs TIVA/N₂O no/narcotics/hypnotics bolus MCI/muscle relaxants induction yes/maintenance yes

Premedication: midazolam 0.1 mg/kg; induction/maintenance: group-specific maintenance regimen. See above. Intubation and relaxation: atracurium 0.5 mg/kg IV, infusion 7 µg/kg/min: TOF ratio, aim one tactile twitch until skin suture, end of surgery/residual neuromuscular blockade reversed atropine 1 mg and neostigmine 2.5 mg IV

Time of outcome determination: before discharge

Method of outcome determination: interview

Emailed author on 27 December 2010 for risk of bias information

Author responded to the email bias survey on 18 January 2011

Survey response: 18 January 2011, Anders G. Jensen anders.gadegaard.jensen@ouh.regionsyddanmark.dk

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The patients were randomized by sealed envelopes into three groups" Quote: "Sealed, numbered envelopes were used. Patients were given the next number when they were enrolled in the study. Envelopes prepared and randomisation performed in advance by a person not participating in the study." (Email bias survey, see notes)

Jensen 1995 (Continued)

Allocation concealment (selection bias)	Low risk	<p>Quote: "The Anaesthesiologist knew, that the patient was given either TIVA (Group 1 and 2) or Neurolept anaesthesia (Group 3). Placebo or flumazenil was drawn from blinded, numbered ampoules. Hence, the Anaesthesiologist was unaware of allocation to group 1 or 2"</p> <p>Quote: "Outcome was assessed by an investigator blinded to group allocation. This investigator did not have access to the patients files." (Email survey)</p> <p>Comment: anaesthesiologists not blinded to anaesthetic technique but blinded to reversal drug vs placebo drug</p>
Blinding of participants and personnel (performance bias) All outcomes	Low risk	<p>Quote: "Patient, Anaesthesiologist, Awareness outcome assessor (interviewer)" (email survey)</p> <p>Quote: "There were 3 groups in the study: 1) TIVA with reversal with flumazenil 2) TIVA with reversal with placebo 3) Standardized neurolept anaesthesia Quote: "The Anaesthesiologist knew, that the patient was given either TIVA (Group 1 and 2) or Neurolept anaesthesia (Group 3). Placebo or flumazenil was drawn from blinded, numbered ampoules. Hence, the Anaesthesiologist was unaware of allocation to group 1 or 2." survey</p> <p>Comment: complicated protocol in regard to blinding and allocation. We will not downgrade because both TIVA groups were blinded to placebo vs study drug and group 3 was blinded to groups 1 and 2.</p>
Blinding of outcome assessment (detection bias) All outcomes	Low risk	<p>Quote: "Outcome was assessed by an investigator blinded to group allocation. This investigator did not have access to the patients files"</p>
Incomplete outcome data (attrition bias) All outcomes	Low risk	<p>Comment: from table 6 and Dr Jensen's response to ROB survey: 2/40 patients from TIVA groups and 4/40 patients from neurolept groups excluded due to lack of data; this is imbalanced between groups but not significantly different, Peto OR 0.49 (0.09 to 2.56)</p> <p>Author sent characteristics and ROB tables for comment 12/7 and 8/13: Anders G. Jensen anders.gadegaard.jensen@ouh.regionssyddanmark.dk</p>
Selective reporting (reporting bias)	Low risk	<p>Quote: "The study was performed more than 20 years ago. The study protocol is no longer available" (email bias survey, see notes)</p> <p>Quote: "All results from our study were reported. That includes the non-significant ones" (email survey)</p> <p>Comment: awareness outcome part of inclusion criteria</p>
Other bias	Low risk	<p>Quote: "I do not find any risk of bias" (email survey)</p>

Jiahai 2012

Methods	<p>Study design: randomized parallel groups</p> <p>Study dates: not stated</p>
Participants	<p>Country: China</p> <p>Sex: female/male</p> <p>Age: mean 59 to 60</p> <p>ASA: unknown but must be \geq III</p> <p>Procedure: OPCAB surgery</p>

Jiahai 2012 (Continued)

Study size: 70

Interventions	<p>Randomized portion of anaesthetic: TIVA ADM (Entropy) vs SCP</p> <p>Intervention 1: (N = 35) entropy values (45 to 55) visible (the entropy group)</p> <p>Intervention 2: (N = 35) without the entropy values visible (the control group)</p>
Outcomes	<p>Primary outcomes: adrenocorticotrophic hormone (ACTH) and cortisol levels</p> <p>Secondary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Quote: "None of the patients in the 2 groups reported intraoperative recall in the postoperative interview"</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of TIVA: N₂O no/narcotics/hypnotics bolus MCI/ muscle relaxants induction yes/maintenance yes</p> <p>Induction midazolam, 0.1 mg/kg; etomidate, 0.1 mg/kg; sufentanil, 1 µg/kg, intubation pancuronium, 0.1 mg/kg. After intubation, infusion propofol, 4 mg to 8 mg/kg/h, and sufentanil, 0.5 µg to 2.0 µg/kg/h, ventilated ET CO₂ 32 mmHg to 42 mmHg. Additional pancuronium, 0.03 mg/kg given; filling pressures and fluid balance maintained lactated Ringer's solution and 6% hydroxyeth. starch; entropy group, propofol infusion rate titrated SE value 45 to 55; bolus propofol, 20 to 50 mg PRN abrupt SE increase; sufentanil infusion adjusted so RE-SE difference remain within 10 U</p> <p>Ma Jiahai, MD, Email: mjh-214@163.com</p> <p>ROB survey: we emailed mjh-214@163.com on 22 March 2015</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "prospective, randomized, controlled study was conducted on 70 patients undergoing first-time OPCAB surgery"
Allocation concealment (selection bias)	Unclear risk	Comment: no information
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: no information
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "A standardized questionnaire (Appendix 1) to measure explicit intraoperative recall was completed immediately after tracheal extubation and 3 days later"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no dropouts
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information

Kamal 1990

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Pakistan Sex: both Age: 41 ASA: I-II Procedure: cholecystectomy Study size: 36
Interventions	Randomized portion of anaesthetic: parts of TIVA: narcotic dose Intervention 1: induction buprenorphine 2.5 µg/kg bolus, N = 18 Intervention 2: induction buprenorphine 5 µg/kg bolus, N = 18
Outcomes	Primary outcomes: duration of anaesthesia, arterial blood pressure and heart rate data, incidence of side effects within the 2 groups Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "None of the patients reported any awareness"
Notes	Non-randomized portion of anaesthetic: parts of TIVA: N₂O no/narcotics/hypnotics bolus MCI/ muscle relaxants induction yes/maintenance yes Premedication: diazepam 0.15 mg/kg; anaesthesia induction: propofol (1 mg/kg) followed by 10 min infusion at (10 mg/kg/h) followed by 10 min infusion at (8 mg/kg/h) + pancuronium (0.1 mg/kg) + group-specific bolus of buprenorphine (agonist-partial antagonist (narcotic)) neuromuscular blockade at the end of surgery was reversed with neostigmine Anaesthesia maintenance: propofol infusion (6 mg/kg/h) Comment: see Dryad: advantages of total intravenous anaesthesia Time of outcome determination: 2 h postoperative Method of outcome determination: interview No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote "patients were randomly allocated into two groups of 18 each"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias)	Low risk	Quote: anaesthesiologist: "An anaesthetist unconnected with the study gave a bolus of intravenous buprenorphine so that the observer was blinded to the dose received"

Kamal 1990 (Continued)

All outcomes

Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Kasmacher 1996

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Germany Sex: both ASA: I and II Study size: 230 Procedure: minor elective surgery
Interventions	Randomized portion of anaesthetic: TIVA vs volatile agent types ("balanced anaesthesia") Intervention 1: induction: etomidate 2 mg/kg; maintenance: 0.8% to 1.5% enflurane, N = 109 Intervention 2: induction: propofol 2 mg/kg + after saturation phase of ~ 10 mg/kg/h, then followed with 5 mg to 6 mg/kg/h N = 121
Outcomes	Primary: dreaming, haemodynamics Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: no evidence intraoperative awareness Sensory experiences during propofol anaesthesia are not stimulus-related perceptions or awareness but dreams similar to normal ones Dreams: 60% (73/121) vs propofol 11% (12/109) enflurane
Notes	Non-randomized portion of anaesthetic: parts of TIVA vs parts of volatile agent types ("balanced anaesthesia"): N₂O no/narcotics/hypnotics bolus MCI/muscle relaxants induction yes/maintenance yes ADM BIS recorded Maintenance both groups: supplemented fentanyl, vecuronium and N ₂ O ("balanced anaesthesia"); BIS Maintenance both groups: enflurane 0.8% to 1.5%, supplemented fentanyl, vecuronium and N ₂ O ("balanced anaesthesia")

Kasmacher 1996 (Continued)

Comment: Kasmacher's view: propofol intraoperative dreams: unrelated to surgery; intraoperative dream incidence: $.60 \times 121 = 73$ propofol; $11 \times 109 = 12$ enflurane; no email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "...randomly assigned to one of two groups..." p147
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Kerssens 2005b

Methods	Study design: randomized parallel groups Study dates: 2001-2002
Participants	Country: USA Sex: both ASA: I and II Study size: 106 - 90 = 16 dropouts Procedure: elective (ambulatory) surgery
Interventions	Randomized portion of anaesthetic: TIVA vs volatile agent types Group 1: maintenance of anaesthesia with propofol (propofol group, N = 48) or Group 2: isoflurane (isoflurane group, N = 42) Word stem completion test
Outcomes	Primary: word test implicit memory associated with preoperative anxiety score

Kerssens 2005b (Continued)

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 0

Quote: "None of the patients consciously recalled the intraanesthetic period on interview...absence of both implicit and explicit memory function in this study"

Notes

Non-randomized portion of anaesthetic: parts of TIVA vs parts of volatile agent types: N₂O yes/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance yes ADM BIS 50 to 55

Induction: lidocaine (1 mg/kg), fentanyl (2 g/kg) and propofol (2 mg/kg). Succinylcholine (1.5 mg/kg) intubation. Maintenance: neuromuscular blockade was maintained at the discretion of the anaesthesiologist, using vecuronium PRN train-of-four at 1:4. N₂O in oxygen (FIO₂ 0.4). Anaesthetics titrated as close to BIS 50 to 55 (mean BIS during word presentation in the trauma study 54). Additional fentanyl (50 g 100 g) if heart rate or blood pressure changed as described previously

Survey response from ckerssens@simpleC.com 10 March 2011

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	ROB survey Dr Kerssens: "a computer-generated assignment to both study list (1 in 4) and memory test (different versions)"
Allocation concealment (selection bias)	Unclear risk	ROB survey Dr Kerssens: "Everything was computer programmed. All we did as investigators was type in patient study number and the computer determined which list to play plus test to use. It was completely blinded. Everybody knew words would be played but WHICH words were unknown to all." Comment: no information regarding anaesthetic allocation
Blinding of participants and personnel (performance bias) All outcomes	High risk	ROB survey Dr Kerssens: "Patient, Anesthesiologist, Awareness outcome assessor (interviewer)" Comment: blinding for word list not anaesthetics
Blinding of outcome assessment (detection bias) All outcomes	Low risk	ROB survey Dr Kerssens: " Patient, Anesthesiologist, Awareness outcome assessor (interviewer)"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "...excluded 16 patients because of anesthetic (drug) protocol violations (n 8), because critical stimuli could not be presented during anesthesia (n 4), or because patients had left the hospital before they were tested (n 4), resulting in 90 evaluable patients. Study group size was chosen to approximate the same number of patients (n 96) as in our previous study" ROB survey Dr Kerssens: "Outcome data was complete. Exclusions are listed in the paper but did not affect anesthesia treatment groups differentially"
Selective reporting (reporting bias)	Low risk	ROB survey Dr Kerssens: "yes"
Other bias	Unclear risk	ROB survey Dr Kerssens: "No, all results were reported"

Kerssens 2009

Methods

Study design: randomized parallel groups

Kerssens 2009 (Continued)

Study dates: Quote: "January 2004 to February 2007" (email bias survey, see notes)

Participants

Country: USA
Sex: both
Age: 61.2, 63.9
Procedure: joint replacement surgery
Study size: 167 enrolled, 128 completed study

Interventions

Randomized portion of anaesthetic: volatile agent types: ADM: BIS 50 to 60 vs SCPs
Intervention 1: BIS monitor used to guide anaesthetic, maintained between **50 to 60**, N = 67
Intervention 2: BIS not seen, standard clinical signs to guide anaesthetic, N = 61

Outcomes

Primary outcomes: effect of BIS-guided anaesthesia vs SCP on memory recognition function
Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1
Quote: "When interviewed postoperatively, three patients (2.3%) reported recall of the time period between falling asleep and waking up from anaesthesia"
Comment: 2 patients were in BIS group and 1 in SCP group. There was evidence of implicit memory in the BIS guided group compared to the SCP group.

Notes

Non-randomized portion of anaesthetic: parts of volatile agent: N₂O no/narcotics/hypnotics bolus MCI/muscle relaxants induction yes/maintenance PRN
Anaesthesia induction: propofol 2 mg/kg + fentanyl 3 µg/kg + vecuronium bromide (0.1 mg/kg) tracheal intubation, additional doses as necessary; maintenance: **sevoflurane** in oxygen using standard ventilation parameters + fentanyl 50 µg to 100 µg + esmolol 0.5 mg/kg + phenylephrine 100 µg + fentanyl 50 µg to 100 µg/kg as needed. Physiologic parameters recorded: BIS, end-tidal gas concentrations (but actual ET data reported) (every 5 s) and vital signs (every 3 min)
Time of outcome determination: 6 h postoperative
Method of outcome determination: interview questions
Comment: see Dryad topics BIS and implicit memory vs SCPs and dreams
Survey response: 10 March 2011, Chantal Kerssens ckerssens@simpleC.com
Author sent characteristics and ROB tables for comment 7 December 2013: ckerssens@simpleC.com

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Patients were randomly assigned to one of two anaesthetic management groups using a computer-generated list linking subject study numbers to group assignment"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias)	Unclear risk	Comment: no information provided

Kerssens 2009 (Continued)

All outcomes

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor: "Outcome assessors were blinded to study group allocation and tested patients postoperatively for recall and recognition memory. Recall was assessed approximately 6 h after surgery with five questions"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "Numbers are reported in the paper. The ratios and reasons for exclusion/attrition were comparable between groups." (email bias survey, see notes) Author sent characteristics and ROB tables for comment 7 December 2013: ckerssens@simpleC.com
Selective reporting (reporting bias)	Low risk	Quote: "True" (email bias survey, see notes) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote: "Yes. We stopped enrolment/testing when the main study -unrelated to memory function - reached it's target enrolment number." (email bias survey, see notes)

Kim 2007

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: South Korea Sex: both Age: 38 ASA: I Procedure: elective orthopaedic or gynaecological surgery Study size: 40
Interventions	Randomized portion of anaesthetic: cardiac drug vs placebo Intervention 1: induction: saline control (N = 20) Intervention 2: induction: nicardipine 15 µg/kg (N = 20)
Outcomes	Primary outcomes: effect of IV nicardipine on haemodynamic and BIS responses Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "No patient in either group had any recall of the procedure"
Notes	Non-randomized portion of anaesthetic: parts of volatile agent: N₂O yes/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance unclear/ADM: BIS recorded Anaesthesia induction: thiopental (5 mg/kg) + fentanyl (1.5 µg/kg) + rocuronium (0.6 mg/kg). 30 sec post-induction saline or nicardipine (15 µg/kg) administered Anaesthesia maintenance: 1.0% inspired concentration of sevoflurane and 50% N ₂ O in O ₂ BIS recorded Time of outcome determination: in recovery room

Kim 2007 (Continued)

Method of outcome determination: interview
 Bias survey sent via email to author. No response received
 Dr Yoon-Sook Lee yslee4719@gmail.com
 Re-sent 13 April 2015

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The patients were randomly allocated, by sealed envelope assignment, into two groups..."
Allocation concealment (selection bias)	Unclear risk	Comment: sealed-envelope technique COMMENT: as described in previous tables
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Kiyama 1997

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Japan Sex: female Age: 44 to 48 years mean each group ASA: I Procedure: gynaecological surgery: total abdominal hysterectomy Study size: 20
Interventions	Randomized portion of anaesthetic: regional anaesthesia: epidural: induction/maintenance Intervention 1: with pre-incisional (before induction) epidural: 1.5% lignocaine with 1:200,000 epi 20 mL after test dose: T4 dermatome to pinprick; N = 10

Kiyama 1997 (Continued)

Intervention 2: maintenance: 15 minutes post-incisional epidural: 1.5% lignocaine with 1:200,00 epi 20 mL after test dose; N = 10

Outcomes	Primary outcomes: effects surgical stimuli on EEG; haemodynamic effects and EEG variables Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: no patient-reported intraoperative recall in the immediate postoperative period and 24 h later
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent: N₂O yes/narcotics/hypnotics bolus I/muscle relaxants induction yes/maintenance unclear ADM EEG recorded</p> Pre-induction: epidural placed; induction: thiopentone 5 mg/kg + isoflurane ET 1%/N ₂ O 40% vecuronium 0.1 mg/kg; maintenance: then isoflurane 1.0% ET and N ₂ O/O ₂ 50%; controlled ventilation Comment: no burst suppression on EEG during study; no recall; interview POD 1 No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "...allocated randomly..."
Allocation concealment (selection bias)	Unclear risk	Comment: above
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Comment: blinded to epidural drug but anaesthesiologists could deduce which intervention group they were managing based on haemodynamic changes (table 2)
Blinding of outcome assessment (detection bias) All outcomes	High risk	Comment: not blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information provided

Kreuer 2003

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Germany Sex: male/female

Kreuer 2003 (Continued)

Age: 18 to 80

ASA: I, II or III

Procedure: minor orthopaedic

Study size: 120

Interventions
Randomized portion of anaesthetic: ADM (BIS 50 Narcotrend D0) vs SCP

Intervention 1: monitor type: Narcotrend target D0

Intervention 1: monitor type: BIS target 50

Intervention 1: monitor type: standard clinical parameter (SCP)

Outcomes

Primary outcomes: anaesthetic consumption, recovery times (eye opening/extubation)
 Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Quote: "Finally, all patients were visited in the postanaesthesia care unit and on the first and third post-operative day and were interviewed about intraoperative recall"

Quote: "No patient reported intraoperative recall"

Notes
Non-randomized portion of anaesthetic: parts of TIVA N₂O no: propofol–remifentanil TCI MCI/muscle relaxants induction yes/maintenance unclear (atracurium)

Premedicated diazepam 0.15 mg/kg PO

Induction/maintenance non-randomized portion of anaesthetic: TIVA

Anaesthesia induction: remifentanil MCI 0.4 µg/kg/min + 5 min later, propofol **TCI** 3.5 µg/mL + atracurium 0.1 mg/kg intubated 3 min later, ventilated ET CO₂ 35 mmHg, remifentanil reduced 0.2 µg/kg/min, whereas propofol TCI adjusted

BIS vs Narcotrend target values vs SCP

Comment: see Dryad topic inadequate anaesthesia

Time of outcome determination: first and third day postoperative. Method of outcome determination: interview

Email sent to secondary author, Wilhelm, on 1 April 2011: no response Dr. Wilhelm: wolfram.wilhelm@uniklinik-saarland.de

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "were randomized to receive a propofol–remifentanil anaesthetic controlled by Narcotrend, by BIS®, or solely by clinical parameters"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: insufficient information provided

Kreuer 2003 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Kreuer 2005

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Germany Sex: both Age: 18 to 80 ASA: I-III Procedure: minor orthopaedic surgery expected to last at least 1 h Study size: 120
Interventions	Randomized portion of anaesthetic: volatile agent types: ADM: BIS 50/Narcotrend D0 vs SCPs Intervention 1: induction and maintenance: SP (N = 40) - control Intervention 2: induction and maintenance: BIS 50 (N = 40) Intervention 3: induction and maintenance: Narcotrend D0 (N = 40)
Outcomes	Primary outcomes: Quote: "The primary end-point of this study was defined as the time taken to spontaneous opening of eye" Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "Complaints of intraoperative recall were not reported"
Notes	Non-randomized portion of anaesthetic: parts of volatile agent: N₂O no/narcotics/hypnotics bolus/muscle relaxants induction yes/maintenance unclear All patients were premedicated with midazolam 7.5 mg orally Induction/maintenance non-randomized portion of anaesthetic: balanced anaesthesia defined by IV (propofol/remifentanyl) and inhalation (desflurane) Remifentanyl infusion at 0.4 µg/kg + 2 mg/kg propofol, oxygen was given by face mask ventilation + 0.5 mg/kg atracurium, trachea was intubated, ventilated ET CO ₂ 35 mmHg, maintenance: remifentanyl reduced 0.2 µg/kg/min, desflurane adjusted: BIS vs Narcotrend target values vs SCP Comment: see Dryad inadequate anaesthesia

Kreuer 2005 (Continued)

Bias survey not sent to author due to incorrect email address

Care Medicine, Sascha Kreuer, MD email to sascha.kreuer@uniklinik-saarland.de

ROB survey. We re-emailed on 13 April 2015; responded 13 April 2015 sascha.kreuer@uks.eu

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "adult patients were randomised to receive...After enrolment, patients were randomised by drawing lots from a closed box" ROB survey Dr Kreuer: "Patients were randomised by drawing lots from a closed box"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided. RCTs that are included in this review and indicate that CONSORT criteria are part of their methods, have unblinded anaesthesiologists. Hence, we cannot assume that other ROB domains are low risk without ROB survey input ROB survey response: "In all patients, irrespective of the individual Group assignment, both BIS values and Narcotrend Levels were continuously recorded in intervals of 5 min by a second independent investigator (LA). In the Standard practice group, both monitors were covered behind a curtain and invisible for the attending anaesthesiologist (CS), whereas, in the EEG groups, either only the Narcotrend or only the BIS monitor was uncovered"
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: anaesthesia titrated to ADMs x 2 vs SCP ROB survey response: "Patient, Anaesthesiologist, Awareness outcome assessor (interviewer)"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Recovery times were recorded by a blinded investigator (MS)"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data ROB survey response: "No Patient was excluded from data Analysis"
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria ROB survey response: "The study protocol is available. All results are reported in the manuscript"
Other bias	Low risk	Comment: as above ROB survey response: "There is no Bias like described above"

Krissel 1994

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Germany Sex: female

Krissel 1994 (Continued)

ASA: not given, "healthy women"

Age: 30

Procedure: elective caesarean section

Study size: 75

Interventions	<p>Randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia vs volatile agent types: induction/maintenance</p> <p>Intervention 1: (Group A) Induction: thiopentone (4 mg/kg) maintenance: ventilation 2 + 2 litre/min N₂O/O₂ mix with 0.8% enflurane (N = 25)</p> <p>Intervention 2: (Group B) Induction: thiopentone (2 mg/kg) + ketamine maintenance: (0.5 mg/kg) ventilation 2 + 2 litre/min N₂O/O₂ mix with 0.8% enflurane (N = 25)</p> <p>Intervention 3: (Group C) Induction: ketamine (1 mg/kg) maintenance: no enflurane (N = 25)</p>	
Outcomes	<p>Primary outcomes: maternal blood pressure, neonate muscle tone, UID intervals</p> <p>Awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4</p> <p>Quote: "Intra-operative consciousness was reported only by patients in the thiopentone group [1 patient awareness report], unpleasant dreams only in the ketamine group and pleasant dreams in both the ketamine and the thiopentone/ketamine groups"</p>	
Notes	<p>Non-randomized portion of anaesthetic: parts of IV vs parts of volatile agent N₂O yes/narcotics/hypnotics bolus/muscle relaxant induction yes/maintenance yes</p> <p>Induction/maintenance non-randomized portion of anaesthetic:</p> <p>Anaesthesia induction: alcuronium (0.1 mg pre-curization). Group-specific induction regimen + succinylcholine (1.5 mg/kg) + alcuronium (0.1 mg/kg)</p> <p>After delivery: all groups: nitrous 67% O₂ 33% + enflurane 1% to 1.5% with alcuronium and ketamine 0.25 mg/kg if needed</p> <p>Note the various definitions of inhalation vs IV</p> <p>Time of outcome determination: 24 h postoperative</p> <p>Method of outcome determination: interview</p> <p>Comment: no email found for author; bias survey not sent</p>	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "...randomly allocated..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided

Krissel 1994 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Kudoh 1999

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Japan Sex: both Age: 25 ASA: NA Procedure: orthopaedic Study size: 40
Interventions	Randomized portion of anaesthetic: parts of TIVA MCI vs bolus maintenance Intervention 1: (group A) maintenance: fentanyl infusion (5 µg/kg/h) for the first 60 min and 3 µg/kg/h for the next 90 min; (N = 20) Intervention 2: (group B) maintenance: fentanyl bolus 50 µg to 100 µg at signs of light anaesthesia; (N = 20)
Outcomes	Primary outcomes: 1 - TIVA (ketamine/droperidol/fentanyl) impact on MLAEP and explicit memory; 2 - impact of differences in infusion methods on dream frequency Awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: no explicit memories; several patients had intraoperative dreaming; group A = 5 dreams group B = 10 (see table 1)
Notes	Non-randomized portion of anaesthetic: parts of TIVA N₂O no/narcotics/hypnotics bolus/muscle relaxant induction yes/maintenance unclear PRN/ADM: MLAEP recorded Total intravenous anaesthesia: droperidol, ketamine, and fentanyl: middle latency auditory evoked potentials Anaesthesia induction: ketamine (2 mg/kg) + fentanyl (5 µg/kg) + droperidol (0.1 mg/kg) + suxamethonium (1 mg/kg) intubation Anaesthesia maintenance: group-specific regimen (see above) + ketamine (2 mg/kg/h) Comment: study of explicit memory: affirmative message and auditory stimulation by pop music during surgery using a tape recorder

Kudoh 1999 (Continued)

Comment: see Dryad: dreams associated with lower levels anaesthesia

Comment: see Dryad topic advantages TIVA

Time of outcome determination: 1 to 3 days postoperative

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: groups were randomized. No discussion of method of randomization
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Kwon 2013

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Korea Sex: female/male Age: 18 to 70; mean 40 to 45 ASA: I-II Procedure: elective surgery Study size: 40
Interventions	<p>Randomized portion of anaesthetic: muscle relaxants: rocuronium vs succinylcholine: maintenance: no information</p> <p>Intervention 1: (N = 20) 0.6 mg/kg rocuronium followed by 1.5 mg/kg propofol: intubated just after confirming loss of consciousness</p> <p>Intervention 2: (N = 20) 1.5 mg/kg propofol and 1.5 mg/kg succinylcholine: intubated 1 minute after injecting succinylcholine</p>

Kwon 2013 (Continued)

Intubation condition, timing of events, and complications were recorded

Outcomes	<p>Primary outcome: acceptable intubation conditions</p> <p>Secondary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Quote: "None of the patients complained awareness of the intubation procedure or had respiratory difficulty during a postoperative interview"</p>
Notes	<p>Non-randomized portion of anaesthetic: hypnotic sedative/narcotic/lidocaine induction agents, muscle relaxants yes/maintenance: no information; other techniques</p> <p>Induction: all patients received 1.5 µg/kg fentanyl intravenously with pre-oxygenation for 2 minutes and were randomized to receive 0.6 mg/kg rocuronium followed by 1.5 mg/kg propofol or 1.5 mg/kg propofol and 1.5 mg/kg succinylcholine</p> <p>Maintenance: no information</p> <p>Comment: no information about maintenance phase of the study; Jaegyok Song, MD, email: drjack@nate.com</p> <p>ROB survey: we emailed drjack@nate.com on 22 March 2015</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Patients were randomly assigned to receive rocuronium (group 1) or succinylcholine (group 2)"
Allocation concealment (selection bias)	Unclear risk	Comment: no information
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: no apparent blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "All patients were interviewed by a blinded investigator 6-24 h after surgery"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no dropouts
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information

Lallemand 2003

Methods	<p>Study design: randomized parallel groups</p> <p>Study dates: March, April 2002 survey</p>
Participants	Country: France

Lallemand 2003 (Continued)

Sex: both
Age: 30
ASA: I
Procedure: elective surgery
Study size: 30

Interventions

Randomized portion of anaesthetic: parts of volatile agent: IV hypnotic agents: induction

Intervention 1: etomidate 0.2 mg/kg (N = 10)
Intervention 2: etomidate 0.3 mg/kg (N = 10)
Intervention 3: etomidate 0.4 mg/kg (N = 10)

Outcomes

Primary outcomes: times to disappearance of the eyelash reflex, to a decrease in the BIS to 50, and to tracheal intubation were compared. The BIS values 30 s following tracheal intubation, and mean arterial pressure (MAP) and heart rate (h) at all time points
Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1
Quote: "No awareness was recorded"

Notes

Non-randomized portion of anaesthetic: parts of volatile agent: N₂O yes/narcotics/hypnotics bolus/muscle relaxant induction yes/maintenance unclear/ADM: BIS

Anaesthesia induction: etomidate (by group) + rocuronium 0.6 mg/kg
Anaesthesia maintenance: sufentanil + isoflurane + BIS <= 50 for **intubation**
Comment nitrous not identified in methods but described in results: "One patient who received etomidate 0.4 mg kg ± 1 did not require any additional drug administration after sufentanil 10 mg had been given in accordance with the study design, and underwent surgery with a BIS value of 30. He recovered consciousness with no recall when nitrous oxide was discontinued"
Quote: "The number of patients for the present study was based on the fact that similar investigations used at the most 30 patients...."
Time of outcome determination: day after surgery
Method of outcome determination: interview
Survey response: 5 February 2011, C Lentschener claude.lentschener@cch.aphp.fr
Author sent characteristics and ROB table grades for comment 8 December 2013: claude.lentschener@cch.aphp.fr

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote ROB survey: "... nurse not involved in the assessment (a) performed the randomisation ... sealed opaque envelopes that contained the group assignments according to a previously computer-generated random list"
Allocation concealment (selection bias)	Low risk	Comment: sealed-envelope technique

Lallemand 2003 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Patient, Anaesthesiologist, Awareness outcome assessor (interviewer)...(b) prepared ...syringes containing... etomidate; and (c) activated the pump infusing...etomidate dose...anaesthetist...took care of BIS monitoring and tracheal intubation, but remained unaware of the etomidate dose administered"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patient, Anaesthesiologist, Awareness outcome assessor (interviewer)" (email survey)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: ROB survey: "...One patient in the 0.2 mg kg ⁻¹ etomidate group was still aware 10 minutes following etomidate administration. This patient was withdrawn from further assessment....that this patient had a high BIS value 10 min following etomidate administration. ..." Comments: this excluded patient is at increased risk of awareness. However, there was a significant difference between groups regarding dropouts: (1/10) etomidate 0.2 mg vs etomidate 0.3 and 0.4 mg (0/20), Peto OR 0.05 (0.00 to 3.18) Author sent characteristics and ROB table grades for comment 8 December 2013: claude.lentschener@cch.aphp.fr
Selective reporting (reporting bias)	Low risk	Quote: "I don't understand this query. The study protocol is clear. Every necessary information is provided. If you wish any additional information in this respect tell me in an additional mail. Sorry. CL" (email bias survey, see notes) Comment: see rules for grading selective reporting for a list of examples of this type of ROB Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote: "In my opinion, there is not any risk of bias Investigators were blinded to group allocation. Data were honestly collected. All recorded data are produced. Excluded patient is reported. I don't see any problem; neither did the Editor" (email survey)

Lam 2013

Methods	Study design: randomized parallel groups Study dates: 1 March 2012 to 31 May 2012
Participants	Country: Taiwan (Republic of China) Sex: female/male Age: 18 to 59 ASA: I-II Obese patients with a body mass index (BMI) of 30 kg/m ² or greater Procedure: bariatric surgery (laparoscopic gastric banding, sleeve gastrectomy, and Roux-en-Y bypass surgery) Study size: 40 enrolled; 38
Interventions	Randomized portion of anaesthetic: parts of volatile agent: IV hypnotic agents: induction I nduction single bolus propofol 2 mg/kg calculated from: Intervention 1 - total body weight (TBW ; 20 patients)

Lam 2013 (Continued)

 Intervention 2 - corrected body weight 60% (**CBW60**; 18 patients)

Outcomes	<p>Primary outcomes: hypotension, complete blood count, coagulation profile, liver function, renal function, electrolytes, total cholesterol, triglyceride, high-density lipoprotein, uric acid, or blood sugar data, BIS values</p> <p>Secondary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Quote: "none reported intraoperative awareness or recall"</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent: N₂O no/narcotics/hypnotics bolus/muscle relaxant induction yes/maintenance yes/ADM: BIS recorded</p> <p>All received: fentanyl (2 µg to 3 µg/LBW) and lidocaine (100 mg) The onset of loss of consciousness: patient dropped the syringe... asked to open...eyes and the eye-lash test by the blinded medical personnel... bolus propofol loss consciousness/succinylcholine (1 mg/TBW): endotracheal intubation. Anaesthesia maintained: 3% sevoflurane 50% oxygen-air mixture (6 L/min) first 5 minutes... BIS monitor</p> <p>Cisatracurium to maintain neuromuscular blockade</p> <p>Quote: "We also defined a bispectral index (BIS) value of greater than 60 as indicative of possible intra-operative awareness"</p> <p>ROB survey: we emailed on 2 April 2015; email address: lin.soon@gmail.com (C-S Lin)</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "We performed a prospective, randomized controlled study to evaluate the clinical efficacy of two different dosages of propofol"
Allocation concealment (selection bias)	Unclear risk	Comment: no information
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "Except for the anesthesiologist who administered the induction dose of propofol, all other medical personnel and the patients were blinded to the propofol induction dose protocol." Email survey: requested details 2 April 2015
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "On questioning the patients during the postoperative interview, none reported intraoperative awareness or recall"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "Two patients were excluded because of incomplete data collection or because they had undergone multiple endotracheal intubation attempts." (CBW 2/20 vs 0/20 TBW). Although imbalanced, there were no significant difference between groups, Peto OR 7.79 (0.47 to 129.11)
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information

Lehmann 1985

Methods	Study design: randomized parallel groups Study dates: Quote: "1984" (email bias survey, see notes)
Participants	Country: Germany Sex: both Age: 37, 41, 40 ASA: I and II Procedure: elective orthopaedic or abdominal surgery Study size: 40
Interventions	Randomized portion of anaesthetic: parts of volatile agent: IV hypnotic agents: maintenance: supplemental narcotic vs placebo Intervention 1: maintenance: placebo infusion (0.9% NaCl) (N = 20) intervention 2: maintenance: tramadol infusion 13 mg/kg/h for 20 min, then 1 mg/kg/h (N = 20)
Outcomes	Primary outcomes: relative cumulative enflurane times (tramadol, placebo), frequency enflurane given as PRN dose, MAP, HR, postoperative recovery time and pain scales, anaesthesiologists judgement of the quality of anaesthesia Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Comment: see Dryad spontaneous memory; dreams associated with memory; partial amnesia, pain yet satisfied with surgery
Notes	Premedication: fentanyl 0.1 mg + atropine 0.5 mg + droperidol 5 mg Non-randomized portion of anaesthetic: parts of volatile agent: N₂O no/narcotics/hypnotics bolus MCI/muscle relaxant induction yes/maintenance yes Induction: methohexitone 100 mg + succinylcholine 1 mg/kg + pancuronium 4 mg Maintenance: N ₂ O/O ₂ 79:21, 4 breaths/min and enflurane (0.5% to 1.5 vol.%) PRN Comment: see Dryad topic for awareness and lower level of anaesthesia; SCPs not predictive of awareness; translated by Lora Schulteiss Survey response: 4 March 2011, Klaus A klaus.uni-koeln.de

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "computer generated randomisation list...the randomisation plan was designed by the manufacturer "Firma Gruenthal", and was not accessible to any persons involved in the clinical study. only after the study was finished, the codes were revealed" Computer-generated randomization list (email bias survey, see notes)
Allocation concealment (selection bias)	Low risk	Quote: "coded ampoules according to randomisation plan, provided by Gruenthal; randomised music presentation according to an own computer generated randomisation list." (email bias survey, see notes)

Lehmann 1985 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Patient, Anaesthesiologist, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patient, Anaesthesiologist, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "no dropouts for any parameter" (email bias survey, see notes)
Selective reporting (reporting bias)	Low risk	Quote: "no" (email bias survey, see notes) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote: "I can not remember any bias." (email bias survey, see notes)

Lehmann 1992

Methods	Study design: randomized parallel groups Study dates: Quote: "1991" (email bias survey, see notes)
Participants	Country: Germany Sex: female Age: 27 to 66 ASA: I and II Procedure: elective gynaecologic surgeries Study size: 60
Interventions	Randomized portion of anaesthetic: parts of IV vs parts of volatile light: balanced anaesthesia I IV induction narcotics vs dissociate agent Intervention 1: induction: fentanyl (5 µg/kg); maintenance: 2 µg/kg/h (N = 20) - control Intervention 2: induction: pentazocine (2 mg/kg); maintenance: 0.8 mg/kg/h (N = 19) Intervention 3: induction: ketamine (2 mg/kg); maintenance: 0.8 mg/kg/h (N = 20)
Outcomes	Primary outcomes: vegetative parameters relative cumulative enflurane times (F, P, K), frequency enflurane given as PRN dose, MAP, HR, postoperative recovery time and pain scales, retrospective anaesthesiologist's judgement of the quality of anaesthesia Secondary outcome: incidence of dreams and recollection of music Awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Comment: see Dryad topic dreams precede awareness, definitions of awareness and memory
Notes	Randomized portion of anaesthetic: parts of IV vs parts of volatile light: balanced anaesthesia I (N₂O yes narcotics + muscle relaxant + PRN volatile inhalation agent)/muscle relaxant induction yes/maintenance unclear

Lehmann 1992 (Continued)

Premedication: pethidine 1 mg/kg IM + promethazine 1.5 mg/kg IM + atropine 0.5 mg IM, 60 min

Anaesthesia induction: alcuronium (2 mg + 8 mg) + methohexital (1.5 mg/kg) + succinylcholine (1 mg/kg) + bolus of randomized narcotic (fentanyl or pentazocine)/anaesthetic (ketamine) bolus and infusion; maintenance: N₂O:O₂ 75:25 + **inadequate** anaesthesia: **enflurane** (0.5% to 2 % vol) for short periods if insufficient anaesthesia indicated; muscle relaxant reversed: neostigmine/atropine

Comment: see Dryad definition: balanced anaesthesia see Appendix 8, definitions of negative dreams, wakefulness, recall of music or words

Interviewed after extubation and POD 1

Lore Schultheiss was translator

Survey response: 4 March 2011, Klaus A klaus.uni-koeln.de

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "...randomised double blind study to evaluate...." Quote: "same design and evaluation as in previous message, concerning PubMed PMID: 3883843 but no help from industry (which means, we prepared all the solutions our self according to a self generated computer randomisation list; the syringes with either fentanyl, pentazocine or ketamine were given to the anaesthesiologist by a colleague not involved in performing or evaluating the study)" (email bias survey, see notes)
Allocation concealment (selection bias)	Low risk	Quote: "see PubMed PMID: 3883843 'coded ampoules according to randomisation plan, provided by Gruenthal; randomised music presentation according to an own computer generated randomisation list'" (email bias survey, see notes)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Comment: yes blind data collection: research nurse administered the randomized drugs
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patient, Anaesthesiologist, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no dropouts
Selective reporting (reporting bias)	Low risk	Quote: "no" (email bias survey, see notes)
Other bias	Low risk	Quote: "I cannot remember any bias" (email bias survey, see notes)

Lehmann 2007

Methods	Study design: randomized parallel groups Study dates: "2004" (email bias survey, see notes)
Participants	Country: Germany

Lehmann 2007 (Continued)

Sex: both
 Age: 65
 Procedure: coronary artery bypass grafting
 Study size: 66

Interventions
Randomized portion of anaesthetic: ADM vs SCP

Intervention 1: BIS 50 (45 to 55) N = 33
 Intervention 2: BIS 40 (35 to 45) N = 33
 Simultaneously, state entropy and response entropy were recorded

Outcomes

Primary outcomes: Quote: "designed and powered to compare differences in the values of BIS and spectral entropy"
 Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1
 Comment: no recall

Notes
Non-randomized portion of anaesthetic: parts of IV N₂O no/narcotics/hypnotics bolus MCI/muscle relaxant induction yes/maintenance yes ADM entropy measured

BIS 50 group induction: midazolam (0.07 mg/kg) + sufentanil (1 µg/kg) + pancuronium (0.1 mg/kg); maintenance: sufentanil 1.5 µg to 2 µg/kg/h + midazolam 0.03 mg to 0.07 mg/kg and sufentanil 0.5 µg to 1 µg/kg as needed
BIS 40 group induction: midazolam (0.1 mg/kg) + sufentanil (1.5 µg/kg) + pancuronium (0.1 mg/kg); maintenance: sufentanil 0.5 µg to 1.5 µg/kg/h + midazolam 0.05 µg to 0.1 µg/kg and sufentanil 1 µg to 2 µg/kg as needed
 Anaesthesia maintenance: O₂ in air 50% + pancuronium 0.03 mc/kg as needed
 The spectral **entropy** parameters RE and SE were **measured**
 Comment: see Dryad topic reduction in inotropic support
 Time of outcome determination: third day after surgery
 Method of outcome determination: interview
 Survey response: 18 January 2011, andreaa@klilu.de

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "flipping coin" (email bias survey, see notes)
Allocation concealment (selection bias)	High risk	Quote: "No concealment" (email bias survey, see notes) Comment: the randomization is to an open BIS endpoint of 40 and 50 and anaesthesia is targeted to a specific BIS endpoint value of 40 or 50. As in other RCTs merged into a meta-analysis in this review, there would be no downgrade for this method of allocation in determining the quality of evidence.

Lehmann 2007 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "Patient" (email bias survey, see notes) Comment: impact on quality of evidence the same as above
Blinding of outcome assessment (detection bias) All outcomes	High risk	Quote: "Patient" (email bias survey, see notes)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "Data were complete in all patients." (email bias survey, see notes)
Selective reporting (reporting bias)	Low risk	Quote: "Study protocol is available; primary and secondary outcomes have been reported. Study was not designed to detect awareness" (email bias survey, see notes) Quote: "Hemodynamics, mixed venous oxygen saturation were recorded but not reported" (email bias survey, see notes)
Other bias	Low risk	Quote: "BIS and entropy values were manually recorded" (email bias survey, see notes)

Lim 1992

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Singapore Sex: both Age: 18 ASA: I and II Procedure: removal of impacted or buried teeth Study size: 50
Interventions	Randomized portion of anaesthetic: parts of TIVA vs parts of volatile agent Intervention 1: induction: (TIVA) propofol (2 mg/kg) at 40 mg/10 sec maintenance: propofol infusion reduced to 6 mg/kg/h by reducing flow manually 2 mg/kg/h every 10 minutes + propofol bolus (0.5 mg/kg) no narcotics as needed oxygen/air , N = 25 Intervention 2: induction: (thiopentone/isoflurane/N₂O), thiopentone (4 mg/kg); maintenance: 66% N₂O in oxygen + 0.5% isoflurane , N = 25
Outcomes	Primary outcomes: recovery characteristics Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: no awareness reported
Notes	Non-randomized portion of anaesthetic: parts of TIVA vs parts of volatile agent N₂O yes/narcotics/hypnotics bolus MCI/muscle relaxant induction yes/maintenance unclear Induction: suxamethonium (1.0 mg to 1.5 mg/kg) + atracurium (0.5 mg/kg) + supplementary dose (0.1 mg/kg)

Lim 1992 (Continued)

 Air/oxygen with TIVA and N₂O with isoflurane

Method of outcome determination: interview

Bias survey sent to author

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "scheduled for surgical removal of impacted or buried teeth were randomly allocated to receive either TIVA using propofol"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "The patients were interviewed by an anaesthetic trainee for complications of anaesthesia ..."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Lin 2011

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Taiwan Sex: female and male Age: mean 56 ± 15 ASA: I-II Procedure: orthopaedic surgery: total hip replacement (THR) (N = 20), the Girdlestone procedure (N = 2), revision of a THR (N = 6), revision of a total knee replacement (N = 1), or knee arthroscopy (N = 1) Study size: 30 enrolled and completed study
Interventions	Randomized portion of anaesthetic: volatile agent types Intervention 1: induction: sevoflurane group inhaled 6% sevoflurane and 4 L/min O ₂ for 3 minutes before intubation, N = 15

Lin 2011 (Continued)

Intervention 2: induction: **non-sevoflurane** group was given **4 L/min O₂ alone** with **IV induction**, N = 15

AAI used in both groups

Outcomes	<p>Primary outcomes: differences in BP, HR, and AAI during intubation between groups</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Comment: no awareness events reported</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent: N₂O no/narcotics/hypnotics bolus MCI/muscle relaxant induction yes/maintenance unclear/ADM AAI recorded</p> <p>Both groups: induction: 3 µg/kg fentanyl, 4 mg/kg thiamylal, and 0.2 mg/kg cis-atracurium and maintenance: 2.5% sevoflurane and 4 L O₂</p> <p>ROB survey. Email: yangcy@adm.cgmh.org.tw, mazuifeng@adm.cgmh.org.tw 7 January 2014</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Subjects were assigned to either the sevoflurane or non-sevoflurane group, according to designations randomly selected by the researcher from a pool that contained 15 assignments to the sevoflurane group and 15 to the non-sevoflurane group..."
Allocation concealment (selection bias)	Unclear risk	Comment: inadequate information
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Quote: "All study procedures were prepared and performed by an anaesthesiologist and an anaesthetic nurse. The same doctor administered anaesthesia for all surgical procedures. The other researcher was responsible for recording the study results for each patient and monitoring vital signs"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "they were interviewed to determine their state of awareness during the intubation and operative procedures. No patients in either group were aware of these events"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information

Lindholm 2008

Methods	<p>Study design: randomized parallel groups</p> <p>Study dates: Quote: "030423-041126" (email bias survey, see notes)</p>
Participants	Country: Sweden

Lindholm 2008 (Continued)

Sex: both
 Age: 50
 ASA: I-III
 Procedure: non-cardiac surgery
 Study size: 320

Interventions	<p>Randomized portion of anaesthetic: ADM vs SCP BIS 40 to 60</p> <p>Intervention 1: SCP blinded to BIS, N = 110</p> <p>Intervention 2: open to BIS 40 to 60, N = 110</p> <p>Induction: automatic closed-loop titration of propofol (closed loop group, N = 20)</p> <p>Manually target controlled infusion of propofol (target control infusion (TCI) group, N = 20)</p>
Outcomes	<p>Primary outcomes: BIS levels, anaesthetic gas consumption, fentanyl use, and subjective opinions on utility and reliability</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Comment no explicit recall</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent: N₂O yes/narcotics/hypnotics bolus MCI TCI/muscle relaxant induction yes/maintenance yes</p> <p>Anaesthesia induction: thiopental or propofol; anaesthesia maintenance: fentanyl + sevoflurane in N₂O; relaxant anaesthesia; for both groups, remifentanyl was administered by a TCI target 2 ng/mL as propofol infusion 1.5 µg/L anxiolysis</p> <p>Comment: see Dryad topic learning bias</p> <p>Survey response: 14 February 2011, Maj-Lis Lindholm MajLisL@LTKalmar.se</p> <p>Clarification missing data email sent 27 December 2013</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	<p>Quote: "A closed envelope was drawn for each study patient, containing a note saying "open" which meant access to the BIS-monitoring, or "closed" meaning that the monitor should be covered." (email bias survey, see notes)</p> <p>Quote: "It is impossible to blind anyone to the fact that the BIS data is available or not." (email bias survey, see notes)</p> <p>Quote: "The sequence of treatments was determined in blocks of 10 (five manual TCI and five closed-loop group) using a random number generator..."</p> <p>Comment: induction regimen is low-risk randomization protocol; technically, the RCT is designed primarily to track the impact of learning with BIS open and closed</p>
Allocation concealment (selection bias)	High risk	<p>Quote: "In the first part of the study, the ten first patients for each nurse were anaesthetized without available BIS..." (email bias survey, see notes)</p>

Lindholm 2008 (Continued)

		Comment: Cochrane policy requires downgrade to high risk despite the evidence provided in this RCT that the learning bias is not always related to non-blinded anaesthesiologists and despite author's statement above
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "It is impossible to blind anyone to the fact that the BIS data is available or not." (email bias survey, see notes)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: no information
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "One patient was not possible to reach because he had no telephone....The only exclusions made were done after the case was completed, and if so, only because of technical failure... " (email bias survey, see notes) Comment: 1 patient could not be reached for interview to assess awareness 1/320 = 0.3% dropout frequency Comment: for awareness outcome, although this lack of follow-up could be for non-relevant reasons, this could possibly be a high-risk awareness case avoiding interaction with medical staff. The patient had no phone. There was no difference between both groups with 1 dropout.
Selective reporting (reporting bias)	Low risk	Quote: "The study protocol is available and was followed." "No we did not" (email bias survey, see notes) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote: "We can not see any obvious bias in the study" (email bias survey, see notes)

Liu 2013

Methods	Study design: randomized parallel groups Study dates: NA
Participants	Country: USA/France Sex: M/F Age: 49 to 73 range ASA: I-IV Procedure: rigid bronchoscopic procedures Study size: 70 enrolled; 67 completed study
Interventions	Randomized portion of anaesthetic: parts of TIVA: MCI vs BIS dual loop: induction/maintenance Intervention 1: (N = 34) manual target-controlled infusion of propofol and remifentanyl (manual TCI group) BIS 40 to 60 Intervention 2: (N = 33) dual-loop group with automatic titration guided by the BIS (dual-loop group) 40 to 60
Outcomes	Primary outcomes: Quote: "The primary outcome was the time spent with adequate anaesthesia, defined by a BIS value of 40-60, expressed as a percentage of time during the maintenance period" Secondary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Liu 2013 (Continued)

"No case of intraoperative awareness was detected"

Notes

Non-randomized portion of anaesthetic: part TIVA: N₂O no/narcotics/hypnotics bolus/muscle relaxant induction yes/maintenance yes

Comment: indications for muscle relaxants in bronchoscopy

M. Fischler MD email: m.fischler@hopital-foch.org

ROB survey: we emailed m.fischler@hopital-foch.org on 22 March 2015

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Eligible patients were assigned by random number generation in a 1:1 ratio to either the manual TCI or the dual-loop group"
Allocation concealment (selection bias)	Unclear risk	Comment: no information
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: both groups titrated to BIS 40 to 60
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "In the PACU and on the day following the procedure, we visited and interviewed all patients regarding intraoperative recall to seek possible intraoperative explicit awareness with a standardized questionnaire"
Incomplete outcome data (attrition bias) All outcomes	Low risk	<p>Comment: 70 enrolled; 67 completed study</p> <p>Quote: "Seventy patients were included, 35 in each group. Three patients were excluded for prolonged absence of BIS signal (one patient) and incomplete data collection (two patients). The analysis thus relates to 34 patients in the manual TCI group and 33 patients in the dual-loop group."</p> <p>Comment: no high-risk awareness exclusions, imbalanced exclusions not significantly different between groups: (2/35) vs (1/35), Peto OR 1.99 (0.20 to 19.75)</p>
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information

Lu 2005

Methods

Study design: randomized parallel groups

Study dates: not given

Participants

Country: France

Sex: both

Age: 58, 59

ASA: I-III

Lu 2005 (Continued)

Procedure: elective minor or major surgery

Study size: 164

Interventions	<p>Randomized portion of anaesthetic: regional anaesthesia: epidural: induction</p> <p>Intervention 1: Group GE: induction: 15 mL of 2% lidocaine epidurally, N = 82</p> <p>Intervention 2: Group GS: induction: saline epidural, N = 82</p>
Outcomes	<p>Primary outcomes: differences in the desflurane concentration between groups</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Comment: no patient-reported intraoperative awareness</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent N₂O no + supplemental narcotics (balanced anaesthesia)/muscle relaxants induction yes/maintenance unclear/PRN ADM AAI 20 ± 5</p> <p>Anaesthesia induction: thiopental (5 mg/kg) + rocuronium (0.6 mg/kg)</p> <p>Anaesthesia maintenance: desflurane titrated in 100% oxygen with target AAI 20 ± 5</p> <p>Time of outcome determination: in recovery room and before discharge</p> <p>Method of outcome determination: interview</p> <p>Survey response: 26 January 2011, Chih-Shung Wong w82556@gmail.com</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	<p>Comment: randomly allocated, by selection of sealed envelopes</p> <p>Quote: "Patient assignments were generated using a computer random number generator and stored in sealed envelopes before initiation of the study protocol." (email bias survey, see notes)</p>
Allocation concealment (selection bias)	Low risk	<p>Comment: sealed envelopes</p> <p>Quote: "The pharmacy department independently prepared sterile syringes of either epidural lidocaine or epidural saline with assigned random numbers. The envelopes were opened immediately after induction of anaesthesia." (email bias survey, see notes)</p> <p>Comment: Cochrane policy emphasizes the need for opaque, sealed envelopes; we graded it low risk because even if the investigators could see through the envelope they could not know in advance the association of a random number and the drug, which could not be differentiated between placebo and lidocaine</p>
Blinding of participants and personnel (performance bias) All outcomes	Low risk	<p>Quote: "Patient, Anaesthesiologist, Awareness outcome assessor (interviewer)." (email bias survey, see notes)</p> <p>Quote: "The anaesthesiologist in charge of the anaesthesia was not aware of the epidural study drug or who controlled the desflurane titration during the whole procedure. The epidural solutions were prepared and injected by a different anaesthesiologist who did not participate in the anaesthesia care or evaluation of the patients"</p>

Lu 2005 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patient, Anaesthesiologist, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "All patients of both groups completed the study protocol" (email bias survey, see notes)
Selective reporting (reporting bias)	Low risk	Quote: "The study protocol was clear. No other non-significant results was available" (email bias survey, see notes)
Other bias	Low risk	Quote: "Although the anaesthesiologist in charge of the anaesthesia was not aware of the epidural study drugs, he also could predict it with the changes of vital signs and the desflurane titration" (email bias survey, see notes)

Maattanen 2002

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Sweden Sex: female and male Age: 44 and 49 ± 13 and 10 ASA: I-II Procedure: elective open spine surgery Study size: 30
Interventions	Randomized portion of anaesthetic: ADM vs SCP Intervention 1: Group I (N = 15), maintenance: desflurane and 50% N ₂ O, was titrated with a target AAI-index of 20 ± 5 Intervention 2: Group II (N = 15), maintenance: desflurane and 50% N ₂ O, titrated according to routine clinical signs (SCP) : heart rate, blood pressure, sweating and tears. No pre-defined fixed MAC-multiple was administered
Outcomes	Primary outcomes: anaesthetic consumption determination (desflurane vaporiser was filled and weighed and re-weighed at end) Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "no patients showed signs of awareness or had any recall postoperatively"
Notes	Non-randomized portion of anaesthetic: parts of inhalation or volatile agent/N₂O yes + supplemental narcotics (balanced anaesthesia)/muscle relaxant induction yes/maintenance unclear/random ADM (AAI: 20) vs SCP AAI-index of 20 ± 5 ADM (AAI: 20) vs SCP: inhalational: desflurane/N ₂ O Premed betamethasone 8 mg IV, induction: propofol as needed, fentanyl 0.5 mg and vecuronium/intubation, controlled ventilation, fentanyl 50 µg PRN and then 2% desflurane if needed

Maattanen 2002 (Continued)

Fentanyl 50mg PRN both groups if need haemodynamic control after sequential 2% increase in desflurane...No other drugs were given during the desflurane anaesthesia

Comment see Dryad topic definition Evan's score and AAI targets with and without muscle relaxants

ROB survey emailed 03 January 2014 jan.jacobsson@mm-medical.se; delivery failed

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Patients were randomised by the envelope technique"
Allocation concealment (selection bias)	Unclear risk	Comment: see above
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "The same anaesthetist (JJ), with 1 year's experience with the A-line monitor, performed all anaesthesia's and, by the nature of the study, was <i>non-blinded</i> . The postoperative nursing staff was blinded"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "grading by nurses otherwise not involved in the study...interviewed in the recovery room...explicit memories from surgery or anaesthesia"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information provided

Mashour 2012

Methods	Study design: randomized parallel groups Study dates: May 2008 until May 2010
Participants	Country: USA Sex: both Age: > 18 years ASA: I-IV Procedure: all surgical cases Study size: enrolled 21,601 completed study 18,836
Interventions	<p>Randomized portion of anaesthetic: ADM (BIS < 60) vs SCP (MAC > 0.5) parts inhalation (majority 98%), parts TIVA (minority 2% cases)</p> <p>Alert if the bispectral index value (BIS) > 60 and/or if the age-adjusted minimum alveolar concentration (MAC) < 0.5, BIS group N = 9460, anaesthetic concentration (MAC) group N = 9376</p>

Mashour 2012 (Continued)

Inhalational or intravenous technique for any surgical case

Outcomes	Primary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Comment: modified intention-to-treat analysis: definite awareness 11/9376 or 0.12% (95% CI 0.07% to 0.21%) Combined incidence of definite and possible awareness cases: 0.08% BIS monitoring 0.20% MAC group 0.38% no intervention group	
Notes	<p>Non-randomized portion of anaesthetic: ADM (BIS < 60) vs SCP (MAC > 0.5) parts inhalation (majority 98%), parts TIVA (minority 2% cases): N₂O unclear/narcotics/hypnotics bolus MCI unclear/muscle relaxant induction yes/maintenance unclear/random (BIS) > 60</p> <p>The study was terminated because of futility</p> <p>Discretion of anaesthesiologist</p> <p>Inhalation technique 98% and TIVA 2% of cases</p> <p>Brice interview: screened one time 28 to 30 days after surgery by telephone or written form of the interview</p> <p>Comment: see Dryad topic definition Michigan Awareness Classification and other conclusions from RCT; gmashour@med.umich.edu: multiple email communications from 31 January 2013 to 21 February 2015. Topics: ROB survey, clarification of missing data, adjudicated awareness reports</p>	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	<p>Quote: "Randomization was performed using a random-number, computer-generated block scheme based on even or odd operating room number"</p> <p>Quote: "A detailed description of the experimental protocol for the Michigan Awareness Control Study (ClinicalTrials.gov No. NCT00689091) has been previously reported (Mashour 2009). The conduct of the study and the reporting of results followed the Consolidated Standards of Reporting Trials guidelines" (Schulz 2010)</p>
Allocation concealment (selection bias)	Low risk	Comment: Schulz 2010
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: SCPs exposed to both groups
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Blinded, trained interviewers used the modified Brice interview..."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: 13% (1395/10770) % ETAG interview NOT completed-dropout ETAG vs 13% (1371/10831) % BIS interview NOT completed - dropout BIS Peto OR 0.97 (0.90 to 1.05)

Mashour 2012 (Continued)

Selective reporting (reporting bias)	Low risk	Quote author ROB response: "...we reported both significant and non-significant findings" Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote: as was detailed in the article, the study was terminated due to futility

Masuda 2002

Methods	Study design: randomized parallel groups Study dates: 1999 to 2000
Participants	Country: Japan Sex: both Age: aged 18 to 65 ASA: I-II (Quote: "were free from hypertension and obesity (BMI < 28)") Procedure: elective surgery for 2 to 3 h Study size: 46
Interventions	Randomized portion of anaesthetic: ADM BIS (40 to 60) vs SCP Intervention 1: blinded to BIS (40 to 60) , N = 20 Intervention 2: SCP, N = 19
Outcomes	Primary outcomes: propofol infusion rates, total amount of propofol, recovery times for both groups Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: in regard to awareness test: "We used a standard questionnaire. We did not find any suspected cases of intraoperative awareness"
Notes	Non-randomized portion of anaesthetic: parts of TIVA N₂O no induction/muscle relaxant induction/maintenance PRN Premed: atropine (0.01 mg/kg) and midazolam (0.05 mg/kg, max 3 mg). Induction: fantanyl (1 µg to 2 µg/kg), vecuronium (0.1 mg/kg), propofol (1.5 mg to 2 mg/kg) fantanyl (at each anaesthesiologist's discretion), vecuronium (added by 1 mg to 2 mg targeting the TOF count of 1). Maintenance: propofol (decreased from 10 mg to 6 mg/kg/h and titrated between 4 mg to 6 mg/kg/h targeting the BIS value 40 to 60), balance of anaesthetics that were not randomized Neither of the induction doses of propofol nor the total doses of fantanyl differed between the BIS and the control groups as indicated in their Table 2. Only the total propofol consumption and the average rate of propofol infusion showed a significant difference as indicated in their Table 3. Translated by Jiro Kurata MD Survey response: 1 September 2011, Messina, Kurata, Masuda jkurata@plum.plala.or.jp

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Masuda 2002 (Continued)

Random sequence generation (selection bias)	Unclear risk	Quote: "We randomly allocated all the patients that underwent elective surgery for 2-3 hours, aged 18-65 years, had the ASA PS of 1-2, and were free from hypertension and obesity (BMI < 28). She did not mention the method of randomizations. They did not find any statistical differences between the BIS and control groups (Table 2)."
Allocation concealment (selection bias)	Low risk	Quote: "The other anaesthesiologists than myself allocated an attending anaesthesiologist to each case. (I did not do such allocation.) Each attending anaesthesiologist was handed a data sheet to fill in during the case without being informed of the purpose of the study. (My residents and I made a preoperative visit to each patient, obtained informed consent, and prepared for BIS monitoring)" Anaesthesiologists were not informed of either the purpose of the study or the group allocation of their patients
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: See ROB rule
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: 1) Patients: blinded (they were not able to see the presence or absence of a BIS monitor) 2) Attending anaesthesiologists: they were not blinded to the usage of a BIS monitor 3) Interviewers: blinded. They did not manage the cases in the OpR
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Quote: "We excluded 7 patients in total: 1) One who had a BIS value less than 80 on arriving at the OpR 2) Two who bled more than 2000 gram 3) Three whose operation lasted less than 1 h 4) One whose operation lasted longer than 5 h (All those reasons are detailed in the text)" 7/46 = 15% Comment: excluded patients due to bleeding are at increased risk of awareness; we do not know if the exclusions were balanced in both groups
Selective reporting (reporting bias)	Low risk	Quote: "...we reported all the results including those that did not reach statistical significance"
Other bias	Low risk	Quote: "We closed the study after enrolling all the 46 patients as had been planned. We had a significantly larger number of female patients in both groups, but observed no statistical difference between the groups in a M/F ratio..."

McNulty 1995

Methods Study design: randomized parallel groups

McNulty 1995 (Continued)

Study dates: not provided

Participants	Country: USA Sex: both Age: 48 to 78 ASA: III and IV Procedure: elective coronary artery bypass or valve replacement Study size: 96
Interventions	<p>Randomized portion of anaesthetic: parts of volatile light (volatile PRN dosing): balanced anaesthesia I IV induction benzodiazepines vs placebo</p> Intervention 1: maintenance (CPB): midazolam low dose 0.05 mg/kg, N = 25 Intervention 2: maintenance (CPB): midazolam high dose 0.1 mg/kg, N = 10 Intervention 3: maintenance (CPB): lorazepam low dose 0.05 mg/kg, N = 26 Intervention 4: maintenance (CPB): lorazepam high dose 0.1 mg/kg, N = 10 Intervention 5: maintenance (CPB): placebo, N = 25
Outcomes	Primary outcomes: recovery times Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Quote: "there were three patients in the placebo group who had sensory experiences consistent with intraoperative awareness. Two of these patients described sensations that were consistent with awareness during CPB (rewarming and cardioversion). The third patient reported awareness during laryngoscopy only"
Notes	<p>Randomized portion of anaesthetic: parts of volatile light: balanced anaesthesia I (N₂O yes narcotics + muscle relaxant + PRN volatile inhalation agent)/muscle relaxants induction yes/maintenance yes</p> Premedication: morphine 0.05 mg to 0.10 mg/kg IM + scopolamine 0.2 mg to 0.3 mg IM + diazepam 0.08 mg to 0.10 mg/kg OS 90 minutes before induction Anaesthesia induction: fentanyl 35 µg to 50 µg/kg + pancuronium 0.1 mg/kg; anaesthesia maintenance: fentanyl + enflurane PRN to treat HTN up to 5 min prior to CPB and after CPB Comment: balanced anaesthesia definitions, see Lehmann 1992 Time of outcome determination: no evidence Method of outcome determination: interview Bias survey sent but not response received
Risk of bias	
Bias	Authors' judgement Support for judgement
Random sequence generation (selection bias)	Unclear risk Quote: "The study drugs were randomized and prepared by the pharmacy in unlabeled syringes"

McNulty 1995 (Continued)

Allocation concealment (selection bias)	Unclear risk	No information provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Comment: syringes were unlabelled Quote: "All individuals involved in the patient's care were blinded to the identity of the study drug"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: insufficient information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Menigaux 2002

Methods	Study design: randomized parallel groups Study dates: not given
Participants	Country: France Sex: both Age: 18 to 70 ASA: I-II Procedure: elective non-cranial surgery Study size: 50
Interventions	Randomized portion of anaesthetic: cardiac drug vs placebo Intervention 1: induction esmolol 1 mg/kg + 250 µg/kg/min, N = 25 Intervention 2: saline solution of equal volume, N = 25
Outcomes	Primary outcomes: changes in BIS, heart rate and MAP with orotracheal intubation (DBIS, DHR, and DMAP, respectively) Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "None of the patients reported recall of intraoperative events"
Notes	Non-randomized portion of anaesthetic: parts of TIVA/N₂O no/muscle relaxant induction yes/maintenance unclear/TCI/IFT/ADM: BIS recorded Anaesthesia induction: propofol was administered via a computer-assisted, TCI , infusion; propofol 4 mg/mL (i.e. 2 mg/kg bolus followed by 225 mg/kg/min) during the entire study period + vecuronium 0.1 mg/kg

Menigaux 2002 (Continued)

IFT: patient movement to intubation; positive response gross purposeful movement arm with the tourniquet attached within 1 min intubation; study ended 16 min after starting the propofol

Anaesthesia maintenance: no information

Time of outcome determination: day after operation

Method of outcome determination: interview POD 1

ROB survey. Email: marcel.chauvin@apr.ap-hop-paris.fr. We emailed on 16 April 2015

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Group allocations were based on computer-generated codes"
Allocation concealment (selection bias)	Low risk	Quote: "Patients and personnel involved in patient management and data collection were unaware of the group assignment"
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Patients and personnel involved in patient management and data collection were unaware of the group assignment"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patients and personnel involved in patient management and data collection were unaware of the group assignment"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided (email bias survey sent)

Mertens 2003

Methods	Study design: randomized parallel groups Study dates: not given
Participants	Country: Netherlands Sex: female Age: 18 to 65 ASA: I or II Procedure: lower abdominal surgery Study size: 30
Interventions	Randomized portion of anaesthetic: parts of TIVA TCI: induction/maintenance

Mertens 2003 (Continued)

Intervention 1: propofol target concentration 2 µg/mL + remifentanyl, N = 10

Intervention 1: propofol target concentration 4 µg/mL + remifentanyl, N = 10

Intervention 1: propofol target concentration 6 µg/mL + remifentanyl, N = 10

3 laryngoscopies in protocol: attempt identify 1 responder and 1 non responder

Outcomes	Primary outcomes: mean measured blood, propofol, and blood remifentanyl concentrations, BIS, spectral edge frequency, systolic and diastolic blood pressures, and heart rate Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "Twenty-four hours postoperatively, the patients were interviewed to evaluate possible side effects and any recall of intraoperative events...None of the patients reported awareness of intraoperative events"	
Notes	<p>Non-randomized portion of anaesthetic: parts of TIVA/N₂O no: muscle relaxant induction yes/maintenance yes ADM BIS recorded/TCI</p> <p>Anaesthesia induction: propofol infusion group-specific target concentrations (TCI) + atracurium (0.4 mg/kg) + remifentanyl infusion target concentration 2 ng/mL</p> <p>Anaesthesia maintenance: group-specific propofol infusion + remifentanyl infusion adjusted to maintain LOC; BIS monitor; inadequate anaesthesia accepted if verified by all 3 observers. To facilitate identification of somatic responses, atracurium given at the minimal dose necessary for surgery (train-of-four levels 1 to 3)</p> <p>Time of outcome determination: 24 h postoperative</p> <p>Method of outcome determination: interview</p> <p>ROB survey. Email: m.j.mertens@lumc.nl. We emailed on 16 April 2015: email address rejected.</p>	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The patients were randomly assigned to one of three study groups to receive, in a double-blind manner"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "One patient in group C had to be excluded from the study due to improper handling of the blood samples" Comment: 33% (1/30) dropout; imbalanced among groups but this was the highest dose group so likelihood of awareness should be lowest. The difference between groups was not significant (1/20 vs 0/10) Peto OR 4.48 (0.07 to 286.49)

Mertens 2003 (Continued)

Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Miller 1996

Methods	Study design: randomized parallel groups Study dates: not given
Participants	Country: Canada Sex: both Age: 18 to 65 ASA: I-II Procedure: arthroscopic knee or laparoscopic procedures Study size: 90
Interventions	<p>Randomized portion of anaesthetic: parts of TIVA: pre-induction hypnotic vs placebo</p> Intervention 1: pre-induction saline 0.1 mL/kg, N = 21 Intervention 2: pre-induction midazolam 15 µg/kg, N = 24 Intervention 3: pre-induction midazolam 30 µg/kg, N = 23 Intervention 4: pre-induction midazolam 45 µg/kg, N = 22
Outcomes	Primary outcomes: effects of midazolam-propofol: evaluated amount propofol required to induce anaesthesia and infusion rates propofol during maintenance and SCPs Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Quote: "The study was discontinued prematurely, as six patients unexpectedly experienced intraoperative awareness with recall (4/21 = 19.1% PLAC vs 2/69 2.9% midazolam groups, P = 0.038)"
Notes	<p>Non-randomized portion of anaesthetic: parts of TIVA bolus MCI/N₂O no/muscle relaxant induction yes/maintenance yes/placebo</p> Surgical day care procedures: unpremedicated, anaesthesia induction: atracurium 0.03 µg/kg then 0.1 mL/kg study drug + alfentanil 20 µg/kg + propofol up to 1 mg/kg + propofol 10 mg IV bolus PRN + atracurium 0.47 mg/kg + intubation Anaesthesia maintenance: air/oxygen 2:1 + propofol 100 µg/kg/min (range: 80 to 200 µg/kg/min PRN light anaesthesia) + propofol 300 µg/kg when infusion rate changed + alfentanil infusion (0.5 µg/kg/min) prior to skin incision + atracurium to maintain 1 to 2 twitches TOF Time of outcome determination: 24 to 48 h postoperative Method of outcome determination: interview Quote: " The study was terminated prematurely due to an unexpectedly high incidence of intraoperative awareness with recall "

Miller 1996 (Continued)

Comment: authors state patients with awareness with pain were satisfied with anaesthetic. This patient response contrasts with what anaesthesiologists as patients prefer (Girgira 2006); no email found

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Patients were then block-randomised (in five blocks of 20) into one of four study groups, according to a computer-generated randomisation schedule"
Allocation concealment (selection bias)	Unclear risk	Quote: "Allocation concealment was achieved with the use of sealed envelopes" Comment: opaque not mentioned
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Blinding was established by preparing midazolam in coded syringes, on the morning of surgery, in concentrations of either 0.15, 0.30, or 0.45 mg.ml ⁻¹ (for groups M-15, M-30 and M-45, respectively). This allowed for study drug/saline preparations to be delivered in a volume of 0.1 ml. kg ⁻¹ iv, in order to prevent group identification"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: insufficient information provided
Other bias	Unclear risk	Comment: insufficient information provided

Miranda 1992

Methods	Study design: randomized parallel groups Study dates: not given
Participants	Country: Malaysia Sex: female Age: 33.2, 31.45 ASA: I Procedure: elective caesarean section Study size: 30
Interventions	Randomized portion of anaesthetic: parts of volatile: balanced anaesthesia II IV induction Intervention 1: induction methohexitone 1%, N = 15

Miranda 1992 (Continued)

Intervention 2: induction propofol 1%, N = 15

Outcomes	<p>Primary outcomes: recovery parameters</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Comment: there was no recall in all patients</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile: balanced anaesthesia II (volatile agent continuous dosing + supplemental narcotics ±muscle relaxant)/N₂O yes/muscle relaxant induction yes/maintenance yes</p> <p>Premedication: ranitidine 150 mg 12 h before + ranitidine 50 mg + metaclopropamide 10 mg</p> <p>Anaesthesia induction: suxamethonium 1 mg/kg</p> <p>Anaesthesia maintenance: N₂O 60% in oxygen 40% + enflurane 1% + atracurium 0.4 mg/kg</p> <p>Post-delivery syntocinon 10 units + fentanyl 3 µg/kg</p> <p>Time of outcome determination: day after surgery</p> <p>Method of outcome determination: interview</p> <p>No email address available for ROB survey</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Patients were randomised into two groups..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	<p>Quote: assessor: "Both the recovery staff and the doctor at the recovery were 'blinded' as regards to the induction agent"</p> <p>Quote: assessor: "The medical officer was not aware of the induction agents used"</p>
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Monedero 1994

Methods	Study design: randomized parallel groups Study dates: not given
Participants	Country: Spain Sex: male and female Age: 18 to 60 ASA: I-II Procedure: elective breast, lumbar, gynaecological surgery Study size: 63
Interventions	<p>Randomized portion of anaesthetic: parts of TIVA vs parts of balanced anaesthesia II (volatile agent + supplemental narcotics ±muscle relaxant)</p> Intervention 1: induction: midazolam perfusion 0.3 mg/kg/h (Group M); maintenance: midazolam perfusion 0.12 mg/kg/h, N = 21 Intervention 2: induction: propofol 2.5 mg/kg (group P); maintenance: propofol perfusion 7 mg/kg/h + pre-incision dose 1.5 mg/kg/h (Group P), N = 21 Intervention 3: induction: thiopental 3 mg/kg (group I); maintenance: isoflurane 1.15% (Group I), N = 21
Outcomes	Primary outcomes: intraoperative signs of inadequate anaesthesia and recovery parameters Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "No patients manifested having memories(awareness) of the intra-operative period"
Notes	<p>Non-randomized portion of anaesthetic: parts of TIVA vs parts of volatile: balanced anaesthesia II (volatile agent continuous dosing + supplemental narcotics ±muscle relaxant)/N₂O no/muscle relaxant induction yes/maintenance yes/MCI</p> TIVA (group M and P) vs inhalational (group 1): midazolam or propofol infusion vs isoflurane 1.15%: alfentanil/vecuronium Induction: all patients 50 µg/kg alfentanil and vecuronium bromide 0.12 mg/kg/h Maintenance: the 3 groups also received one pre-incision dose of alfentanil 25 µg/kg and post-incision perfusion at 60 µg/kg/h. alfentanil changed 20 µg/kg/h PRN After surgery group M flumazenil 0.5 mg IV and a perfusion of flumazenil 0.5 mg over 60 min Quote: "Conclusions: None of the TIVA techniques proved superior in all the parameters studied during anaesthetic maintenance when compared with <i>balanced</i> isoflurane/alfentanil, although the propofol-alfentanil combination was found to be superior to that of midazolam-alfentanil. After anaesthesia, however, recovery was better with the association of propofol-alfentanil and adverse side effects were fewer. Flumazenil at the doses used was ineffective for preventing re sedation due to midazolam" Brett Smith: translator from Spanish No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote from translator Brett Smith: "...Whereas there is no indication of randomization in the abstract (English or Spanish), within the body of the paper

Monedero 1994 (Continued)

I found that the authors did indeed randomize the patients to the three different intervention groups. Here is where I located the reference: under the section heading "Pacientes y metodos".

Quote: "...Los pacientes se distribuyeron de forma aleatoria ('The patients were allocated in a random way') en 3 grupos de 21 pacientes: midazolam (grupo M). propofol (grupo P) o induccion con tiopental (grupo I)"

Allocation concealment (selection bias)	Unclear risk	Quote from translator Brett Smith: "the authors did report allocation concealment of patients to the 3 intervention groups (M, P, I), although they don't say how they randomized the allocation, i.e. random number generator, etc. As far as the other biases, I didn't see any evidence of blinding...authors didn't report any participant attrition..."
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: see above
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: see above
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: see above
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information provided

Morimoto 2002

Methods	Study design: randomized parallel groups Study dates: data discarded
Participants	Country: Japan Sex: male and female Age: 56 ± 9, 53 ± 12 (range 18 to 70) ASA: I-II Procedure: mixed type elective surgery Study size: 60, 14 excluded, 46 completed study, 23% dropout
Interventions	Randomized portion of anaesthetic: ADM BIS (40 to 60) vs SCP Intervention 1: SCP blinded to BIS, N = 21 Intervention 2: open to BIS (40 to 60) , N = 25 Total N completed study = 46, 14 dropouts
Outcomes	Primary outcomes: anaesthesia consumption and recovery times

Morimoto 2002 (Continued)

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Comment: no awareness

Notes

Non-randomized portion of anaesthetic: parts of volatile agent/N₂O yes + supplemental narcotics + muscle relaxant induction yes/maintenance unclear

Quote: "Sixty patients (ASA physical status 1 or 2) undergoing various surgical procedures under sevoflurane/nitrous oxide anaesthesia were studied..... In the BIS group, sevoflurane concentration was adjusted to achieve target BIS values between **40-60** during surgery and 60-75 during the final.."

Premed: atropine (0.5 mg) plus midazolam (2 mg, if < 60 years old) or hydroxyzine (50 mg, if > = 60 years old). Induction: thiopental (5 mg/kg), vecuronium (0.1 mg/kg). Maintenance: sevoflurane (as shown in the abstract)/**N₂O**, balance of anaesthetics that were not randomized, fentanyl (control: 129 ± 64; BIS: 132 ± 80 µg), vecuronium (control: 16 ± 5; 14 ± 4 mg), pentazocine (control: 2.4 ± 5.6; 4.2 ± 6.9 mg)

Quote: "Table 1 shows the discharge criteria from PACU. Table 2 shows demographics. The line means Age, Sex(male/female), Weight, Anaesthetic time, Operation time, Blood loss, Total doses of fentanyl, Total doses of vecuronium, Doses of sevoflurane, Incidence of Hypertension, Incidence of bradycardia or hypotension, Incidence of nausea and vomiting, Total doses of pentazocine in PACU. The left column shows control group and right column shows BIS group"

Quote: "The important limitation of this study is not double blinded design. The anaesthesiologists [were] aware [of] the patients' group during the surgery"

Comment: nonetheless, the control group were blinded to BIS values

Jiro Kurata MD translator

Survey response: 1 September 2011, Messina, Kurata, Morimoto jkurata@plum.plala.or.jp

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "We used envelope method"
Allocation concealment (selection bias)	Unclear risk	Comment: no information
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "The study group was blinded to the patient not to anaesthesiologists and interviewer" "The patients were randomly assigned to two groups of which anaesthesia was carried out with (BIS group) or without (control group) monitoring BIS, and in the latter, anaesthesiologist was blinded to the BIS values" Comment: blinded to ADM in SCP group but ADM group is not blinded to both ADM and SCP
Blinding of outcome assessment (detection bias) All outcomes	High risk	Quote: "The study group was blinded to the patient not to anaesthesiologists and interviewer"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: 14 patients were excluded because the operation time was shorter than 2 h or longer than 6 h 14/60 = 23%; BIS 21 = N and SCP = 25 = N; this is imbalanced across groups but there was no significant difference between groups with respect to these exclusions: (21/30 vs 25/30) Peto OR 0.48 (0.15 to 1.57)

Morimoto 2002 (Continued)

Selective reporting (reporting bias)	Low risk	Comment based on the translator's verbal and written response: no Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Comment based on the translator's verbal and written response: no

Mozafari 2014

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Iran Sex: male/female Age: mean 47 and 48; range 18 to 65 ASA: I-III Procedure: elective abdominal surgery (laparoscopy, cholecystectomy) Study size: 392 enrolled, 333 completed study
Interventions	Randomized portion of anaesthetic: ADM BIS values (target range: 45 to 65) vs SCP: volatile agent Intervention 1: BIS monitoring (N = 163) BIS values (target range: 45 to 65) Intervention 2: routine monitoring (N = 170)
Outcomes	Primary outcomes: BIS values and vital parameters including systolic blood pressure (SBP), diastolic blood pressure (DBP), heart rate (HR), and SPO ₂ Secondary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Quote: "The overall incidence of awareness in the BIS and routine monitoring groups were 5.5% and 4.1%, which was not significantly different" Comment: total awareness events 16
Notes	Randomized portion of anaesthetic: parts of volatile agent/N₂O yes + supplemental narcotics + muscle relaxant induction yes/maintenance unclear Anaesthesia induced: sufentanil 0.1 µg to 0.2 µg/kg, thiopental 3 mg to 5 mg/kg and atracurium 0.5 mg/kg maintained isoflurane or halothane/ N₂O/BIS Comment: the reported awareness events from this RCT suggests that there is something unusual about the method of administration of anaesthesia compared to other RCTs in this review from other countries with similar interventions but lower awareness rates or the criteria or protocol that other studies use to identify or include patient awareness reports is different. The difference maybe related to 1) the percentage of illiterate patients in study and/or 2) the validated awareness questionnaire specific to Persian culture (Malek 2010a) Quote: "However, it seems that the incidence of this phenomenon and its complications are exactly dependent on the quality of postoperative interview by specialists. It has been shown that detection of awareness depends on the technique, timing and structure of interview ..." Comment: Brice interview is NOT validated Author: Amir Asadi Fakhr, Department of Anesthesiology, School of Paramedicine, Hamadan University of Medical Sciences, Hamadan, IR Iran. Tel: +98-9183159883, Email: asadi@umsha.ac.ir ROB survey. We emailed asadi@umsha.ac.ir on 22 March 2015; no response 17 April 2015

Mozafari 2014 (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "We considered about 196 samples for each group. Patients were allocated to BIS monitoring (n = 163) and routine monitoring (n = 170) groups using the permuted block randomization method"
Allocation concealment (selection bias)	Unclear risk	Comment: no information
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: personnel knew treatment group assignment
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "In addition, information related to the awareness during anesthesia was collected by a special questionnaire including formalized set of open-ended questions"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "We considered about 196 samples for each group. Patients were allocated to BIS monitoring (n = 163) and routine monitoring (n = 170) groups using the permuted block randomization method. Nevertheless, 30 and 26 persons disagreed to participate in the study (BIS monitoring n = 163 and routine monitoring n = 170)" Comment: exclusions before surgery started; no exclusions between groups that started and finished surgery
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information

Muralidhar 2008

Methods	Study design: randomized parallel groups Study dates: not given
Participants	Country: India Sex: both Age: mean 50 ASA: III-IV Procedure: off-pump coronary artery bypass grafting (CABG) Study size: 40
Interventions	Randomized portion of anaesthetic: parts of TIVA vs parts of volatile agent ADM vs SCP intervention 1: isoflurane BIS open vs closed (SP), N = 20 intervention 2: propofol BIS open vs closed (SP), N = 20

Muralidhar 2008 (Continued)

Group A (I - no BIS): isoflurane in O₂; ETAC 1% to 1.2%

Group B (I - BIS): **isoflurane** inspired concentration adjusted BIS **50 ± 5**

Group C (P - no BIS): maintained propofol infusion (after a 50mg bolus) at 6 mg to 8 mg/kg/h to sternotomy and 4 mg to 6 mg/kg/h thereafter

Group D (P - BIS): maintained (after a 50 mg bolus) titrated to a BIS value of **50 ± 5**

Outcomes	<p>Primary outcomes: anaesthesia consumption</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Quote: "The patients who experienced recall were reevaluated and determined by a senior author and in his judgement none of the awareness reports were valid"</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of TIVA N₂O no vs parts of volatile agent + supplemental narcotics + muscle relaxant induction yes/maintenance unclear</p> <p>General anaesthesia induced, 100% O₂ facemask, fentanyl 2 µg/kg, midazolam 100 µg/kg/sleep dose of thiopentone</p> <p>Endotracheal intubation/pancuronium 0.15 mg/kg and mechanical ventilation</p> <p>Quote: "If patient suggested that he suffered from awareness under anaesthesia he/she was visited by the senior author and the attending anaesthesiologist to discuss, explain the peri-operative events, answer patient's questions sympathetically and refer the patient for psychological counselling, if necessary"</p> <p>Comment: senior author was sole adjudicator of patient awareness reports. He classified all reports as not valid awareness</p> <p>No email address available for ROB survey</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "randomly divided into four groups by a sealed envelope technique"
Allocation concealment (selection bias)	Unclear risk	Quote: "randomly divided into four groups by a sealed envelope technique"
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria

Muralidhar 2008 (Continued)

Other bias	Unclear risk	Comment: insufficient information provided
------------	--------------	--

Myles 1997

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Australia Sex: female and male Age: 64 ± 10 ASA: NA Procedure: CABG surgery Study size: 124 patients (34 with poor ventricular function), 119
Interventions	<p>Randomized portion of anaesthetic: parts of TIVA vs parts of volatile agent + supplemental narcotics</p> Intervention 1: TIVA : maintenance: propofol-based (5 mg/kg/h prior to sternotomy, than 3 mg/kg/h thereafter; N = 58) Intervention 2: inhalation balanced anaesthesia : enflurane-based 0.2% to 1.0%, fentanyl boluses (5 µg/kg) sternotomy, CPB; midazolam 0.1 mg/kg CPB, enflurane BP.85 MAP; no enflurane on CPB (N = 66)
Outcomes	Primary outcomes: recovery parameters (e.g. average time to extubation) Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: no awareness events reported
Notes	<p>Non-randomized portion of anaesthetic: parts of TIVA/N₂O no vs parts of volatile agent + supplemental narcotics + muscle relaxant induction yes/PRN maintenance</p> Induction: fentanyl 15 µg/kg and midazolam 0.05 mg/kg enflurane group: additional bolus of fentanyl 5 µg/kg prior to sternotomy and fentanyl 10 µg/kg with midazolam 0.1 mg/kg at (CPB) Comment: identical induction technique, IV midazolam 0.05 mg/kg/fentanyl 15 µg/kg. Endotracheal intubation/muscle relaxation/IV pancuronium 0.12 mg/kg. PRN: IV vecuronium 2 mg to -4 mg Maintenance of anaesthesia differed 5 patients excluded: due to intraoperative complications ROB survey sent p.myles@alfred.org.au

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Myles 1997 (Continued)

Random sequence generation (selection bias)	Unclear risk	Quote: "we randomized patients after stratification according to the surgeon's angiographic assessment of contractility ...Randomization was determined by a table of random numbers"
Allocation concealment (selection bias)	High risk	Quote: "An acknowledged deficiency of our study was that the investigators were not blinded to treatment allocation ...We retained details of intraoperative drug treatment until after extubation, so that ICU staff were unaware of group allocation , and so decisions were minimally biased ..."
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: inhalation vs TIVA
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: see above Quote: "within 3 days of surgery and queried about any recall of intraoperative events"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: based on our reading of the results section, we assume that the 129 patients that started surgery and the 5 exclusions before the end of surgery were distributed as follows: propofol (4/62) vs enflurane (1/67), Peto OR 3.75 (0.63 to 22.27); hence, although the exclusions, which were due to death or change in surgical plans due to severe disease, were imbalanced, there was no significant difference between groups
Selective reporting (reporting bias)	Low risk	Comment: primary outcome: recovery parameters (e.g. average time to extubation); secondary outcome: awareness; no ROB survey data Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information provided

Myles 2004

Methods	Study design: randomized parallel groups Study dates: not given
Participants	Country: Australia, China Sex: both Age: mean 58 BIS and 57 SCP High risk of awareness ASA: I-IV (> 80% patients ASA III-IV) ASA: I-V Procedure: elective and emergency Study size: enrolled 2503, completed study 2463
Interventions	Randomized portion of anaesthetic: ADM BIS 40 to 60 vs SCP: parts inhalation (53% of cases) or TIVA techniques (43% of cases) Intervention 1: BIS monitor, maintain BIS 40 to 60 (N = 1225)

Myles 2004 (Continued)

Intervention 2: SCP No BIS monitor (N = 1238)

Outcomes	<p>Primary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4</p> <p>Quote: "There were 22 reports of confirmed or possible awareness in the BIS group (1.8%) and 27 reports in the routine care group (2.2%; P = 0.49) "</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of TIVA MCI (43% of cases) vs parts of volatile agent (57% of cases)/N₂O yes + supplemental narcotics + muscle relaxant induction yes/PRN maintenance</p> <p>Anaesthesia induction/maintenance: anaesthetic drugs: clinical judgement</p> <p>Comment: Table 4: TIVA was used on 43% in BIS group and 42% in SCP group. So balanced anaesthesia (inhaled agents with IV agents) used in 57% and 58%, respectively</p> <p>Time of outcome determination: 2 to 4 h, 24 to 36 h, and 30 PODs</p> <p>Method of outcome determination: structured interview</p> <p>Comment: 22% (11/49) of definite and possible adjudicated patient awareness reports had pain</p> <p>Comment: Table 3: 50% (18/36) adjudicated possible awareness reports are intraoperative dream reports. Some of these may have been in postoperative period; up to 50% (9/18) of these dreams could be interpreted as being associated with pain/distress; see Dryad appendix</p> <p>Quote: "Dreamt about aliens and thought aliens had taken over the operation (theatre staff had had a conversation about aliens during surgery)"</p> <p>Comment: this one intraoperative dream report was corroborated by OpRstaff. It is proof that patient dream reports that do not contain content consistent with events during surgery should not be interpreted as proof that they do not represent fragments of intraoperative memory as part of a portion of patient reports of intraoperative dreams</p> <p>Paul Myles p.myles@alfred.org.au on 14 January 2011: ROB survey not received. Multiple email communications about the following topics were exchanged from 14 January 2011 to 5 April 2015: adjudication of patient awareness reports by expert panel</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "perioperative care were unaffected. After consent was obtained and immediately before induction of anaesthesia the anaesthetist rang the co-ordination centre to obtain a computer-generated random group allocation"
Allocation concealment (selection bias)	Low risk	Comment: see random sequence allocation
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: insufficient information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor: "Follow-up was undertaken by a blinded observer, with a structured interview, and an independent endpoint adjudication committee was established"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: 2503 enrolled - 40 excluded before surgery started = 2463 started surgery; 65 patients (BIS group 30, routine care group 35) did not provide any

Myles 2004 (Continued)

interview data, mainly because of critical illness or death in the postoperative period

Peto OR: not interviewed:

(30/1225 vs 35/1238) 100.0%; 0.86 (0.53 to 1.41)

number of patients able to be interviewed at each of the three time periods were:

(23/1252 vs 17/1251) 100.0%; 1.36 (0.73 to 2.53)

Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Myles 2007

Methods	Study design: randomized parallel groups Study dates: April 2003 and November 2004
Participants	Country: worldwide Sex: both Age: > 18, average age 55.8, 54.6 ASA: I-IV Procedure: major surgery expected to last at least 2 h Study size: 2050, 2012
Interventions	Randomized portion of anaesthetic: N₂O vs N₂O-free Intervention 1: maintenance oxygen 80% + nitrogen 20% (N = 997) Intervention 2: maintenance: N ₂ O 70% + oxygen 30% (N = 1015) control
Outcomes	Primary outcomes: duration of hospital stay Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Comment: 2 awareness reports in N ₂ O group
Notes	Non-randomized portion of anaesthetic: N₂O vs N₂O-free: parts of TIVA and parts of volatile agent + supplemental narcotics + muscle relaxant induction yes/PRN maintenance ADM BIS subgroup recorded Anaesthesia induction: standard anaesthetic care, anaesthetic depth adjusted clinical judgement and, if available, bispectral index monitoring Comment: restricted the secondary analysis to the N ₂ O-free group to minimize the possibility of selection (Berksonian) bias . Additional regression analyses: explore the effect of possible covariate imbalance Comment: despite the decrease in postoperative complications, did not observe a meaningful difference in duration of hospital stay between groups

Myles 2007 (Continued)

Comment: N₂O-free anaesthetic: less myocardial infarction and death, but could be **atype II error because study not of sufficient size/rare in unselected patients**: follow-up trial in 7000 patients at risk of coronary artery disease (the ENIGMA II trial)

Time of outcome determination: 24 h and 30 days postoperative

Method of outcome determination: structured questionnaire

Bias email survey sent. No response received

ROB survey sent p.myles@alfred.org.au

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Patients were randomly assigned to ...using a computer- generated code, accessed via an automated telephone voice recognition service..."
Allocation concealment (selection bias)	Unclear risk	Quote: "Treatment assignment was stratified by site and elective/emergency status of the surgery, using permuted blocks"
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "All patients otherwise received standard anaesthetic care and monitoring. Choice of other anaesthetic drugs and intravenous fluids was at the discretion of the attending anaesthesiologist"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "At the end of the procedure, the intraoperative case report form and documentation of group identity were faxed to the data management centre and then placed in an opaque envelope by the anaesthesiologist. The envelope was then sealed to ensure blinding of research staff conducting the postoperative follow-ups.... All patients were seen by a research assistant on the day after surgery to assess their quality of recovery and to detect awareness" Quote: assessor: "Patients and observers were blind to group identity"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: figure 1: 4035 undergoing surgery > 2 h GA, 1696 excluded before surgery for non-medical reasons, 3187 patient eligible for study, 2274 excluded before surgery after eligible: no obvious factor for exclusion increased risk of awareness, 2050 randomized, 1020 assigned N ₂ O-free group: 23 developed exclusion criterion: 997 assessed for primary endpoint (of whom 5 received N ₂ O): 23/1020 = 2.3%, 1030 assigned N ₂ O group: 15 developed exclusion criterion: 1015 assessed for primary endpoint (of whom 9 did not receive N ₂ O): 15/1030 = 1.5%. If we assess these exclusions as an outcome in data analysis table using Peto OR: (23/1020) vs 15/1030), Peto OR 1.55 (0.82 to 2.95). That is, there is no significant difference between the 2 groups
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Navarro 2000

Methods Study design: randomized parallel groups
 Study dates: not stated

Navarro 2000 (Continued)

Participants	Country: Germany Sex: female Age: 25 to 26 ± 5 to 5.5 ASA: I-II Procedure: caesarean section Study size: 75 enrolled and completed study and data collection
Interventions	Randomized portion of anaesthetic: volatile vs volatile Intervention 1: (N = 25) desflurane 2.5 % Intervention 2: (N = 25) isoflurane 0.5 % Intervention 3: (N = 25) epidural 15 mL ropivacaine 7.5 mg/mL with fentanyl 100 µg
Outcomes	Primary: outcomes: recovery time (recovery period), intraoperative haemodynamic changes, blood loss, APGAR scores, neurologic adaptive capacity scores, umbilical vein blood gas analysis Secondary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "The patients were interviewed about intraoperative awareness 24 and 48 h after the operation. None of them reported awareness during the operation"
Notes	Non-randomized portion of anaesthetic: parts of volatile agent/N₂O/muscle relaxant induction yes/maintenance yes Thiopental (4 mg/kg IV) and succinylcholine (1.0 mg to 1.5 mg/kg) IV N ₂ O to 50% in oxygen (O ₂), pancuronium Alfentanil 15 µg to 25 µg/kg after birth PRN, oxytocin, neostigmine, and atropine antagonized muscle relaxants ROB survey no email address available
Risk of bias	
Bias	Authors' judgement Support for judgement
Random sequence generation (selection bias)	Unclear risk Quote: "...were randomly divided into two groups of 25 patients who received either desflurane or isoflurane"
Allocation concealment (selection bias)	Unclear risk Comment: inadequate information
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk Comment: inadequate information
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk Comment: inadequate information
Incomplete outcome data (attrition bias) All outcomes	Low risk Comment: no dropouts

Navarro 2000 (Continued)

Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information

Nayar 2009

Methods	Study design: randomized parallel groups Study dates: November 2005 to December 2007
Participants	Country: India Sex: female Age: 24 ASA: I-II Procedure: caesarean section Study size: 60
Interventions	Randomized portion of anaesthetic: parts of volatile agent (IV induction agents of inhalational anaesthesia) Intervention 1: induction: thiopentone (5 mg/kg) (N = 20) Intervention 2: induction: ketamine (1 mg/kg) (N = 20) Intervention 3: induction: ketamine (0.5 mg/kg) + thiopentone (2.5 mg/kg) (N = 20)
Outcomes	Primary outcomes: APGAR, haemodynamics Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "None of our patients reported intraoperative awareness or dreams"
Notes	Randomized portion of anaesthetic: parts of volatile agent/N₂O yes + supplemental narcotics + muscle relaxant induction yes/maintenance yes Premedication: ranitidine 150 mg + 30 mL of 0.3 M sodium citrate Anaesthesia induction: study specific drug (see above) + suxamethonium (1.5 mg/kg) Anaesthesia maintenance: ventilation with N ₂ O, oxygen, and halothane (0.5%), vecuronium Time of outcome determination: 3 h and 24 h postoperative Method of outcome determination: interview Email survey sent, no response received ROB survey: Dr Nayar anesthesia62@hotmail.com We emailed on 18 April 2015.

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Nayar 2009 (Continued)

Random sequence generation (selection bias)	Unclear risk	Quote: "following which they were randomly assigned to 3 groups..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor: "Each patient was visited by an anaesthetist blinded to patient group..."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Ngan 1997

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Ngan: Hong Kong, China Sex: female Age: 29 to 34 ASA: ASA I-II Procedure: caesarean section surgery Study size: 40 (38 statistical analysis)
Interventions	Randomized portion of anaesthetic: parts of volatile agent (IV induction agents of inhalational anaesthesia) + supplemental narcotics + muscle relaxant induction yes/maintenance yes Intervention 1: induction: thiopental 4 mg/kg (N = 20) Intervention 2: ketamine 1 mg/kg (N = 20)
Outcomes	Primary outcomes: difference in 24 h morphine consumption, umbilical arterial and venous blood gases, APGAR scores Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "No patients had recall of intraoperative events or unpleasant dreams"

Ngan 1997 (Continued)

Notes

Non-randomized portion of anaesthetic: parts of volatile agent N₂O yes + supplemental narcotics + muscle relaxant induction yes/maintenance yes

Rapid-sequence induction using study drug, succinylcholine 1.5 mg/kg, ventilated 50% N₂O in oxygen ET CO₂ 4.2%. Isoflurane ET 0.5%, atracurium. **After delivery** 10 IU oxytocin and morphine 0.15 mg/kg, N₂O increased to 70%, and isoflurane reduced ET 0.3%

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Patients were randomized by drawing of shuffled coded envelopes"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Comment: no evidence of blinding Quote: "... Anesthesiologist not involved with patient assessments calculated each induction dose, diluted it to 20 mL with saline, and covered the syringe with opaque adhesive tape"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "No patient reported recall of intraoperative events"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: 2 dropouts ketamine group (2/20) vs thiopental group (0/20); uneven distribution of high-risk dropouts but no significant difference between groups, Peto OR 7.79 (0.47 to 129.11) Quote: "patients in the ketamine group were excluded, one because of massive intraoperative haemorrhage that required hysterectomy and postoperative ventilation and one because of technical problems with the PCA device"
Selective reporting (reporting bias)	Low risk	Comment: primary outcome: morphine consumption; secondary outcome: awareness; no ROB survey data Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information provided; no ROB survey data

Ngan Kee 2002

Methods	Study design: randomized parallel groups Study dates: "Oct-1998 to Apr-2001" (email bias survey, see notes)
Participants	Country: China Sex: female Age: 26 to 43 ASA: I and II

Ngan Kee 2002 (Continued)

Procedure: elective caesarian section

Study size: 60

Interventions	Randomized portion of anaesthetic: parts of volatile agent vary dose/N₂O yes vs no Intervention 1: maintenance sevoflurane 0.6% ET + O ₂ 30% + N ₂ O 70% (N = 20) Intervention 2: maintenance sevoflurane 1% ET + O ₂ 50% + N ₂ O 50% (N = 20) Intervention 3: maintenance sevoflurane 2% ET + O ₂ 100% (N = 20)	
Outcomes	Primary outcomes: difference in umbilical arterial and venous blood gases, APGAR scores Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "No patient reported recall of intraoperative events. One patient in Group 50 reported experiencing intraoperative dreams but was not distressed by this"	
Notes	Non-randomized portion of anaesthetic: parts of volatile agent/N₂O no (IV induction agents of inhalational anaesthesia)+ supplemental narcotics + muscle relaxant induction yes/PRN maintenance Premedication: ranitidine 150 mg before surgery + sodium citrate 30 mL 0.3 M Anaesthesia induction: thiopental 4 mg/kg + succinylcholine 1.5 mg/kg + atracurium as needed (PRN); anaesthesia maintenance: see above interventions Morphine 0.15 mg/kg + oxytocin 10 IU post-delivery Author responded to the email bias survey on 21 January 2011. Responses are recorded in the 'Risk of bias' table. Survey response: 21 January 2011, Ngan Kee, Warwick Dean warwick@cuhk.edu.hk	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Computer-generated randomisation code held in sealed opaque envelopes" (email bias survey, see notes)
Allocation concealment (selection bias)	Low risk	Comment above (email bias survey, see notes)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Separate investigators applied intervention (oxygen) and administered anaesthesia while different investigators performed data collection. Anaesthesia machine turned away from investigators collecting data so they were unable to see settings"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"Patient, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data Quote: "No attrition or exclusions" (email bias survey, see notes)
Selective reporting (reporting bias)	Low risk	Quote: "yes" (email bias survey, see notes). Unclear - probably meant "no" language barrier.

Ngan Kee 2002 (Continued)

Comment: awareness outcome part of inclusion criteria

Other bias

Low risk

Quote: "yes" (email bias survey, see notes)

Comment same as selection bias

Oddby-Muhrbeck 1993

Methods

Study design: randomized parallel groups

Study dates: 1991-1992

Participants

Country: Sweden

Sex: female

Age: 30, 32

ASA: I

Procedure: laparoscopy

Study size: 60

Interventions

Randomized portion of anaesthetic: parts of TIVA propofol vs parts of volatile agent isoflurane/IFT/music memory

Intervention 1: maintenance: propofol 10 mg/kg, 8 mg/kg, and 5 mg to 8 mg/kg as needed (N = 30)

Intervention 2: maintenance: isoflurane 1.5 to 2 MAC 3 to 6 min then set as needed (N = 30)

Patients 2 groups randomly assigned 3 **subgroups** listening: **soft music**, hard rock music or **no** music at all. **Isolated forearm** technique detect insufficient anaesthesia.

12 patients in each group were randomly assigned hear one piece of **soft music** (Cavatine by Stanley Myer). Another 12 patients one piece of **hard rock music** (Prowler by Iron Maiden). 6 patients in each group played **blank tape - control groups**

Outcomes

Primary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Comment: no patient in any of the groups had memories of operation or moved the isolated arm in response to command or surgery

Comment: none of the 60 patients moved the isolated arm in response to commands or surgery

Quote: "...two patients picked the correct piece of music. It is impossible to say if they were guessing, dreaming or if they really did recall the music that was played"

Notes

Non-randomized portion of anaesthetic: parts of TIVA/N₂O yes vs parts of volatile agent + supplemental narcotics + muscle relaxants induction yes/maintenance yes

Premedication: midazolam IM (4 mg below 70 kg and 5 mg above)

Anaesthesia induction: fentanyl 1.5 µg to 2 µg/kg + propofol 1 mg to 3 mg/kg + suxamethonium chloride 1 mg to 1.5 mg/kg

Anaesthesia maintenance: 66% **N₂O** in O₂ + atracurium 0.15 mg to 2.0 mg/kg/fentanyl PRN

Comment: important issues in this paper are 1-4 stages of unconsciousness and TIVA defined to include N₂O: see Dryad topic

Oddby-Muhrbeck 1993 (Continued)

Time of outcome determination: 24 h postoperative

Method of outcome determination: questionnaire

This paper design with volunteers and breakdown of music guessing should be model to compare other music papers

 Survey response: 21 January 2011, jan jakobsson jan.jakobsson@ki.se

 Survey response: 8 August 2011, jan jakobsson jan.jakobsson@ki.se

Survey response: 21 January 2013 re. method of randomization

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Thirty patients, randomly selected, received propofol infusion..."
Allocation concealment (selection bias)	Low risk	Quote survey: "Envelopes with group assignment: Envelopes provided randomly and a nurse otherwise not involved in the study broke an envelope after that the patient had been screened and provided written and oral consent"
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Patient, Anaesthesiologist, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patient, Anaesthesiologist, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "Eight patients were excluded because the laparoscopy was followed by laparotomy or other surgical procedures such as tubotomy through the laparoscope." Comment: 13% dropout (8/60); dropout related to surgical findings
Selective reporting (reporting bias)	Low risk	Quote: "The study protocol is not available but it is clear that the published report include all expected outcomes" (email bias survey, see notes) Quote: "No all results associated to the study protocol are reported" (email bias survey, see notes) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote: "No obvious bias" (email bias survey, see notes)

Panousis 2009

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Germany Sex: both

Panousis 2009 (Continued)

Age: 51 to 66

ASA: II-III

Procedure: major abdominal surgery

Study size: 45, 43

Interventions	<p>Randomized portion of anaesthetic: regional anaesthesia: epidural: local anaesthetic, ropivacaine vs placebo</p> <p>Intervention 1: Group 1: pre-induction: ropivacaine 10 mL of 0.5% every 60 min + sufentanil (0.5 µg/mL), N = 15</p> <p>Intervention 2: Group 2: pre-induction: ropivacaine 10 mL of 0.2% every 60 min+ sufentanil (0.5 µg/mL), N = 13</p> <p>Intervention 3: Group 3: pre-induction: placebo (saline) control group, N = 15</p>
Outcomes	<p>Primary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1)</p> <p>Quote: "During the postoperative interviews, no patient reported intraoperative awareness or recall"</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent/N₂O yes + supplemental narcotics + muscle relaxant induction yes/PRN maintenance BIS 55, even without changes in the PRST</p> <p>Anaesthesia induction: propofol (1.5 mg/kg) + sufentanil (0.5 µg/kg) + rocuronium (0.5 mg/kg)</p> <p>Comment: see Dryad topic definition PSRT score</p> <p>Anaesthesia maintenance: desflurane in 60% N₂O at 1 MAC + additional remifentanyl; desflurane administration was adjusted to the actual demand under continuous BIS and PRST score monitoring</p> <p>An increase in BIS 55, even without changes in the PRST score, was treated by increasing the end-tidal concentration of desflurane until a maximum end-tidal desflurane concentration of an age-adapted 1 MAC</p> <p>Time of outcome determination: 2 h and 2 days postoperative</p> <p>Method of outcome determination: Brice questionnaire</p> <p>Bias survey sent via email to author. No response received yet.</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The solutions were prepared by a nurse according to a computer derived block randomization list"
Allocation concealment (selection bias)	Low risk	Quote: "Syringes were delivered to the operating room (OpR) merely labelled with "TEAMAC-Study" and the consecutive patient number"
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "The attending anaesthesiologist was blinded to patient randomization"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "To detect intraoperative awareness or recall, patients were interviewed using the Brice questionnaire immediately after surgery, and 2 days later when they were discharge..."

Panousis 2009 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "Because of protocol violations in terms of remifentanil administration at desflurane levels <1 MAC, two patients from Group 2 were withdrawn from the database" Comment: 4.5% (2/45) imbalanced dropout rate across groups, but protocol violation was due to giving more remifentanil than required; hence, no increase awareness risk associated with dropouts
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Pauls 2009

Methods	Study design: randomized parallel groups Study dates: "October 10, 2007 - April 1, 2008"(email bias survey, see notes)
Participants	Country: Canada Sex: both Age: 49.9, 59.6 ASA: I-IV Procedure: neurosurgery Study size: 23, 21
Interventions	Randomized portion of anaesthetic: ADMs: EEG vs BIS Intervention 1: only EEG monitor visible to anaesthesiologist for maintenance, N = 8 Intervention 2: only BIS monitor visible to anaesthesiologist for maintenance, N = 8 Continuously measured/time to burst suppression
Outcomes	Primary outcomes: an assessment of quality of emergence Quote (ClinicalTrials.gov): "This study is designed to test the hypothesis that the EEGo monitor will be superior to the BIS monitor to assess emergence following neuroanesthesia" Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "...none of the patients reported evidence of intraoperative awareness at 24 h post-surgery"
Notes	Non-randomized portion of anaesthetic: parts of volatile agent/N₂O no + supplemental narcotics MCI + muscle relaxant induction yes/PRN maintenance Premedication: midazolam 1 mg Anaesthesia induction: remifentanil 1 µg/kg + propofol 1.5 mg to 2.5 mg/kg + lidocaine 1.5 mg/kg + rocuronium 0.6 mg/kg Anaesthesia maintenance: desflurane 0.5 to 1.5 MAC + remifentanil 0.05 µg to 0.1 µg/kg/min + propofol up to 50 µg/kg/min + phenylephrine 0.1 µg to 0.5 µg/kg/min + morphine 0.1 mg/kg

Pauls 2009 (Continued)

Time of outcome determination: 24 h postoperative

Comment: see Dryad discussion differences BIS vs EEG

Method of outcome determination: interview

Quote: "Author responded to the email bias survey on 12 February 2011. Responses are recorded in the 'Risk of bias' table

Survey response: 12 February 2011, Wac Mutch wacmutch@shaw.ca

<https://clinicaltrials.gov/ct2/show/NCT00443807>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Both the BIS monitor and the EEG signals were...measured...the clinician was given access to only one of the monitors, as randomly allocated by a third party with a coin toss to either EEGo (heads) or BIS (tails)"
Allocation concealment (selection bias)	Unclear risk	Quote: "Not assigned until all patients were induced with both monitors applied and functional. Then randomized as above" (email bias survey, see notes)
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "Patient, Anesthesiologist, Awareness outcome assessor (interviewer)" (email bias survey, see notes) Comment: not blinded to SCPs
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Twenty-four hours following surgery, a survey of patient satisfaction and a query regarding awareness were completed by an anesthesiologist blinded to the mode of EEG monitoring"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "All 21 recruited patients completed the protocol... cerebral aneurysm clipping...MVD for trigeminal neuralgia. The two surgical groups behaved very differently on emergence...were not compared with the MVD patients to assess emergence criteria..." (email bias survey, see notes) Comment: cerebral aneurysm clipping surgery can not be assessed for awareness due to surgery on brain; there was no dropout in regard to awareness assessment
Selective reporting (reporting bias)	Low risk	Quote: "The study protocol was listed at ClinicalTrials.gov with identifier NCT00443807." (email bias survey, see notes) Quote: "There was commentary contained within the study of an episode of patient arousal." (email bias survey, see notes) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote: "Each patient had both monitors applied. They were compared post-hoc with no inherent bias" (email bias survey, see notes)

Paventi 2001

Methods Study design: randomized parallel groups

Paventi 2001 (Continued)

	Study dates
Participants	Country: Italy Sex: both Age: mean: 42, 48 (18 to 75) ASA: I-IV Procedure: abdominal surgery (greater than 30 minutes) Study size: 90
Interventions	Randomized portion of anaesthetic: ADM (BIS) vs SCP intervention 1: BIS40 to 60, N = 45 intervention 2: SCP (BIS blinded), N = 45
Outcomes	Primary outcomes: anaesthetic consumption, recovery times Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: outcomes: no explicit memory (awareness)
Notes	Non-randomized portion of anaesthetic: parts of volatile agent/N₂O no, air yes + infusion of narcotics MCI + muscle relaxant(s) induction yes/maintenance yes Premed: diazepam 5 mg to 10 mg PO; anaesthetics given per anaesthesia All patients: induction and maintenance: remifentanyl 1 (µg/kg) and TPS (4 mg to 8 mg/kg) vecuronium 0.1 mg/kg for intubation and for maintaining neuromuscular blockade during maintenance Maintenance: remifentanyl mean 0.4 µg/kg/min and sevoflurane 0.5 MAC air 50%; after skin closure at the end of surgery, residual neuromuscular blockade was reversed Email survey sent to author, paventi@iol.it , no response received

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "randomly allocated"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias)	Low risk	Comment: no missing outcome data

Paventi 2001 (Continued)

All outcomes

Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Pedersen 1992

Methods	Study design: randomized parallel groups Study dates: not given
Participants	Country: Denmark Sex: female Age: 31, 29 ASA: NA Procedure: emergency or elective caesarean section Study size: 51
Interventions	Randomized portion of anaesthetic: volatile agent doses \pm N₂O Intervention 1: maintenance oxygen 2 litres/min + N₂O 2 litres/min + halothane 0.5% , N = 26 Intervention 2: maintenance oxygen 4 litres/min + halothane 2% maintenance changed to oxygen 2 litres/min + N₂O 2 litres/min + halothane 0.5% , N = 25
Outcomes	Primary outcomes: difference in the duration of anaesthesia, recovery time Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Comment: 15% of the mothers receiving 0.5% halothane had some recall. No maternal reminiscence was seen using 2% halothane.
Notes	Non-randomized portion of anaesthetic: IV hypnotic/PRN narcotic/muscle relaxant parts of volatile (potent inhalation) technique/N₂O no/muscle relaxants induction yes/PRN maintenance Pre-oxygenation atropine 0.5 mg + oxygen 10 litres/min x 3 mins, manual hand controlled ventilation , anaesthesia induction: thiopentone 4 mg to 5 mg/kg + suxamethonium 1.5 mg/kg/intubation, maintenance (randomized inhalational agents) pre-delivery; post cord clamp , Syntocinon infusion (40 IU/1 L solution)+ pethidine + gallamine PRN Time of outcome determination: day after operation Method of outcome determination: interview No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Pedersen 1992 (Continued)

Random sequence generation (selection bias)	Unclear risk	Quote: "the choice of anesthesia was made by drawing an envelope randomizing the patient to one of...the volatile anaesthetic mixture was administered according to the randomized pre-induction allocation"
Allocation concealment (selection bias)	Unclear risk	Comment: insufficient information provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "The patients were manually ventilated by an anaesthesiologist who was completely unaware of the anaesthetic gas mixture delivered"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: no information
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Persec 2012

Methods	Study design: randomized parallel groups Study dates: February to July 2011
Participants	Country: Croatia Sex: 21 male and 19 female Age: median 65 years (25 to 84 years) ASA: II-III Procedure: major abdominal surgery (explorative laparotomy and colostomy/colectomy) Study size: 45, 40
Interventions	Randomized portion of anaesthetic: ADM (BIS) vs SCP Intervention 1: routine anaesthesia care: non BIS-guided group Intervention 2: BIS-guided anaesthesia 50 and 60
Outcomes	Primary outcomes: BIS levels, measurement of heart rate (HR), systolic blood pressure (sBP), end-tidal CO ₂ (ET CO ₂), operation time, and extubation time Secondary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 "None of the patients reported explicit awareness"
Notes	Non-randomized portion of anaesthetic: volatile (potent inhalation) + N₂O technique: IV hypnotic/PRN narcotic/muscle relaxant both induction/maintenance The operation was performed under general anaesthesia using midazolam (0.15 mg/kg), fentanyl (2 µg/kg), and vecuronium (0.1 mg/kg) to facilitate endotracheal intubation, and 1.5 to 2.5 MAC of sevoflu-

Persec 2012 (Continued)

rane, **N₂O** 50% in oxygen, boluses of fentanyl and vecuronium for maintenance. Intraoperatively, after induction doses of fentanyl, anaesthesia was mainly balanced with sevoflurane

Correspondence: Jasminka Persec, MD, PhD, Clinical Department of Anesthesiology, Reanimatology and Intensive Care Medicine, University Hospital Dubrava, Av. G. Suska 6, HR-Zagreb, Croatia

ROB survey. We emailed jpersec@net.amis.hr on 22 March 2015.

ClinicalTrials.gov (NCT01470898) - no information on web page <https://clinicaltrials.gov/ct2/results?term=NCT01470898>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The investigation was also registered on ClinicalTrials.gov (NCT01470898)...According to a computer generated randomisation list... " Comment: ClinicalTrials.gov (NCT01470898) - no information on web page https://clinicaltrials.gov/ct2/results?term=NCT01470898
Allocation concealment (selection bias)	Unclear risk	Comment: see above
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "The non BIS-guided group ... was blinded to the anaesthesiologist in charge. All values were recorded by the younger anaesthesiologist, who was not involved in the anaesthesia maintenance. All other aspects of peri-operative treatment were similar, including choice of anaesthetic agents and monitoring" Comment: as explained in previous RCT domains, the SCP group is NOT blinded to both groups. We use this as an example that indicates that specific aspects of RCT methods can result in high risk domains in the ROB table. Hence, unless explicitly described we do not assume that trial registration leads to low ROB domain grades
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: as above
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "Out of 45 patients recruited for the study, 5 patients were excluded because of an inoperative malignant process found intra-operatively"
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: see above

Piggott 1990

Methods	Study design: randomized parallel groups Study dates: "Late 1988 - late 1989" (email bias survey, see notes)
Participants	Country: Wales Sex: female

Piggott 1990 (Continued)

Age: not given

ASA: NA

Procedure: elective or emergency caesarean section

Study size: 200, 197

Interventions
Randomized portion of anaesthetic: volatile agent doses \pm N₂O

Intervention 1: maintenance isoflurane 1.8% in 100% oxygen was reduced to 1.2% after 5 min, N = 100

 Intervention 2: maintenance isoflurane 1.2% in 50% N₂O in oxygen was reduced to 0.6% after 5 min, N = 100

Outcomes

Primary outcomes: umbilical venous and arterial blood, samples

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Comment: there were no instances of awareness

 Comment: 4 patients (2 in each group) recalled **dreaming** on direct questioning (**2 had moved the isolated arm and 2 had not**)

 Quote: "One patient (group 50) reported a **sensation of painful pressure, but believed she had been dreaming. She had not moved her arm.**"

Comment: the patient in group 50 could also have been classified as awareness, 2 moved their arms presumably not in response to complex command; many would call this non-purposeful movement; see discussion of non-purposeful movement

 Comment: this is evidence that a **portion of dreams are a variant of intraoperative memory**
Notes
Non-randomized portion of anaesthetic: volatile (randomized) (potent inhalation) technique continuous then PRN: IV hypnotic/PRN narcotic/muscle relaxants both induction/maintenance

Anaesthesia induction: thiopentone 3 mg to 4 mg/kg + suxamethonium 1.5 mg/kg + pancuronium 4 mg to 6 mg

Anaesthesia maintenance: isoflurane increased 50% for 5 min added as needed

Quote: "...no taped message was played to the anaesthetized patients, and they were not asked beforehand to move the hand if they felt pain, as it was considered that this would add to existing stress"

Comment: see Dryad topic non-purposeful vs purposeful movement with IFT and topic of air oxygen paralysis only in neonates

Time of outcome determination: postoperative before discharge

Method of outcome determination: interview

Author responded to the email bias survey on 7 February 2011. Responses are recorded in the 'Risk of bias' table.

 Survey response: 7 February 2011, David George Bogod david.bogod@nottingham.ac.uk
Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	Quote: "Alternate groups for alternate weeks...Randomisation was far from ideal by current standards!" (email bias survey, see notes)

Piggott 1990 (Continued)

Allocation concealment (selection bias)	High risk	Quote: "Participants unaware of allocation sequence. Investigators aware" (email bias survey, see notes)
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "Patient, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patient, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "Uncertain so long after the study. Not fully addressed in the paper. 197 out of 200 analysed for indication for Caesarean section, and 100% of neonates followed-up. From what I recall, data were very largely complete, as subjects were in-patients during follow-up" (email bias survey, see notes) Comment: unclear about distribution of 3 patient dropouts Comment: assume the most imbalanced variant and there is no significant difference between groups: (3/50 vs (0/50), Peto OR 7.70 (0.78 to 75.76)
Selective reporting (reporting bias)	Low risk	Quote: "Protocol no longer available (this was 31 years ago!). I can only say that, from what I recall, it was followed fully" (email bias survey, see notes) Quote: "All results reported" (email bias survey, see notes) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Comment: no exclusions listed Quote: "Randomisation was far from ideal by current standards! Other forms of bias were, to the best of my knowledge so long after the event, excluded"

Plourde 1996

Methods	Study design: randomized parallel groups Study dates: Quote: "Jan to Oct 1991" (email bias survey, see notes)
Participants	Country: Canada Sex: female Age: 19 to 50 ASA: I-II Procedure: reduction mammoplasty Study size: 12
Interventions	Randomized portion of anaesthetic: volatile agent doses \pmN₂O Intervention 1: maintenance N ₂ O (66% end-tidal (ET)) enflurane 0.5% ET, N = 4 Intervention 2: maintenance N ₂ O (66% end-tidal (ET)) enflurane 0.8% ET, N = 4 Intervention 3: maintenance N ₂ O (66% end-tidal (ET)) enflurane 1.1% ET, N = 4

Plourde 1996 (Continued)

Outcomes	<p>Primary outcomes: 40 Hz ASSR/AML, AMLR amplitude, latency, or phase data</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Comment: no patient had any recollection of intraoperative events</p>
----------	---

Notes	<p>Non-randomized portion of anaesthetic: volatile (randomized) (potent inhalation) technique: IV hypnotic/PRN narcotic/muscle relaxants both induction/maintenance ADM AMLR recorded</p> <p>Anaesthesia induction: fentanyl 3 µg/kg + thiopental 3 mg to 5 mg/kg + vecuronium 0.08 mg/kg</p> <p>Anaesthesia maintenance: fentanyl 1.0 µg/kg as needed + vecuronium 0.01 mg/kg</p> <p>Time of outcome determination: 4 to 8 weeks postoperative</p> <p>Method of outcome determination: interview</p> <p>Neostigmine 40 µg/kg + glycopyrrolate 10 µg/kg used to reverse muscle paralysis</p> <p>Primary author responded to email bias survey. Responses are recorded in 'Risk of bias' table below.</p> <p>Comment: study was not powered to assess for awareness but electrophysiology (author's response to survey)</p> <p>Survey response: 13 February 2011, Gilles Plourde gilles.plourde@mcgill.ca</p>
-------	---

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	<p>Quote paper: "four patients per concentration; random assignment..."</p> <p>Quote survey: "Assignment (concentration of enflurane, recording order i.e. ASSR or AMLR first) written on cards which were shuffled and individually places in a sealed envelope by the department secretary who was familiar with these procedures" (email bias survey, see notes)</p>
Allocation concealment (selection bias)	Unclear risk	Quote: "Anaesthesiologist did not keep track of cases done. So each envelope brought a surprise" (email bias survey, see notes)
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "Patient, Awareness outcome assessor (interviewer), Data analyst" (email bias survey, see notes)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patient, Awareness outcome assessor (interviewer), Data analyst" (email bias survey, see notes)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "One patient was excluded because of poor baseline AMLR" (email bias survey, see notes)
Selective reporting (reporting bias)	Low risk	<p>Quote: "Main goal of study was electrophysiology. Awareness was only outcome examined...Yes. Some findings did not reach significance" (email bias survey, see notes)</p> <p>Comment: awareness outcome part of inclusion criteria</p>
Other bias	Low risk	Quote: "No obvious bias" (email bias survey, see notes (email))

Puri 2003

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: India Sex: both Age: 18 to 70 ASA: NA but by type of surgery ASA III-IV Procedure: valve replacement or coronary artery grafting under cardiopulmonary bypass Study size: 30
Interventions	Randomized portion of anaesthetic: volatile (potent inhalation) ADM (BIS) vs SCP Intervention 1: BIS (45 to 55), N = 14 Intervention 2: SCP, N = 16
Outcomes	Primary outcomes: haemodynamic disturbances, time to recovery Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Quote: "None of the patients in the study group had recall of intraoperative events when interviewed on the first postoperative day, while one patient in the control group had awareness during the sternotomy"
Notes	Non-randomized portion of anaesthetic: volatile (potent inhalation) + N₂O technique: IV hypnotic/PRN narcotic/muscle relaxants both induction/maintenance (CPB) Anaesthesia induction: morphine 0.2 mg/kg 1 + midazolam 0.05 mg/kg + thiopental + vecuronium 0.08 mg/kg Maintenance induction titrated to keep BIS between 45 and 55 emergence 65 to 75 + morphine 0.05 mg to 0.1 mg/kg as needed, N = 15 Anaesthesia maintenance: isoflurane + N ₂ O (66% O ₂ before CPB) + morphine 0.025 mg/kg/h + morphine 3 mg + midazolam 1 mg + vecuronium 0.5 mg Quote: "...while in the study group, the anaesthesiologist adjusted the vasoactive drugs to maintain pressure while keeping the BIS constant " Comment: this is the method cardiac anaesthesiologists use to maintain what they judge to be an appropriate depth of anaesthesia without lightening the anaesthetic as the main or supplemental intervention to treat hypotension. This cause and method of avoiding awareness is rarely described in papers. Comment: senior author adjudicated patient awareness reports as not valid without expert panel; studying this phenomena Time of outcome determination: 24 h after operation Method of outcome determination: interview gdpuri007@hotmail.com

Risk of bias

Puri 2003 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "...were randomized into either a study group (using BIS) or a control group (no BIS) using computer generated numbers...."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: insufficient information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: insufficient information provided Quote: "Every patient was interviewed to determine any recall on the first postoperative day after the trachea had been extubated"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Puri 2007

Methods	Study design: randomized parallel groups Study dates: "Feb 2003 to Dec 2004" (email bias survey, see notes)
Participants	Country: India Sex: both Age: 39 Procedure: elective non-cardiac surgery Study size: 40
Interventions	Randomized portion of anaesthetic: TIVA + CLADS vs TIVA + ADM: BIS 50 Intervention 1: closed loop anaesthesia delivery system (CLADS): propofol 5.03 ± 1.68 mg/kg-1, h-1 Intervention 2: manual infusion control: 7.33 ± 2.07 mg/kg-1, h-1 propofol titrated to BIS = 50
Outcomes	Primary outcomes: induction time (the time required to achieve target BIS after start of infusion), induction dose, minimum BIS within 30 seconds of induction, total dose propofol, median duration of closed loop control or manual control, and median time interval between end of closed-loop control (or end propofol infusion in manual control) and extubation Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: no awareness cases

Puri 2007 (Continued)

Notes

Non-randomized portion of anaesthetic: parts of TIVA/N₂O no: narcotics MCI/muscle relaxants induction yes/maintenance yes (CPB)

Anaesthesia induction: fentanyl (2 µg/kg) prior to induction than fentanyl 1 µg/kg-1, h-1 continuous infusion for the duration of surgery; endotracheal intubation/0.1 mg/kg vecuronium

Anaesthesia maintenance: fentanyl infusion (1 µg/kg/h)

Time of outcome determination: at discharge, 1 day, and 1 week postoperative

Method of outcome determination: modified Brice questionnaire

Survey response: 16 February 2011, Dr Goverdhan Dutt Puri gdpuri007@hotmail.com

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote ROB survey response, Dr Puri: "Computer generated" (email bias survey, see notes)
Allocation concealment (selection bias)	High risk	Quote ROB survey response, Dr Puri: "Anaesthesiologist conducting anaesthesia knew of the group the patient belong. But anaesthesiologist assessing awareness post-op did not know the group to which patient belong" (email bias survey, see notes)
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: given the closed vs open loop systems used, like Pedersen 1992 , it is possible to blind the anaesthesiologist but requires extra manpower
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote ROB survey response, Dr Puri: "Patient, Awareness outcome assessor (interviewer), Data analyst...But anaesthesiologist assessing awareness post-op did not know the group to which patient belong" (email bias survey, see notes)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote ROB survey response, Dr Puri: "all data reported"
Selective reporting (reporting bias)	Low risk	Quote ROB survey response, Dr Puri: "...don't think any non significant results of any importance to the study relevant were excluded" (email bias survey, see notes) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote ROB survey response, Dr Puri: "free of bias as all patients had anaesthesia to be controlled on target BIS numbers" (email bias survey, see notes)

Rehberg 2007

Methods	Study design: randomized parallel groups Study dates: "5/2005-10/2005" (email bias survey, see notes)
Participants	Country: Germany Sex: both

Rehberg 2007 (Continued)

Age: 49, 55

ASA: I-III

Procedure: elective minor urological or gynaecological surgery

Study size: 92

Interventions	<p>Randomized portion of anaesthetic: TIVA: MCI vs TCI propofol</p> <p>Intervention 1: induction propofol MCI no restriction of dose; maintenance propofol no dose restriction; both manual TIVA N = 46 - 2 (2/46 = 5%) (unplanned ICU stay = 44)</p> <p>Intervention 2: induction propofol TCI 7 mg to 10 mg/mL; maintenance propofol TCI 2.5 mg to 4.5 mg/mL; N = 46 - 4 (4/46 = 9%) (unplanned ICU stay = 42)</p>
Outcomes	<p>Primary outcomes: proportion of priming words identified and the BS values</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Quote: "none of our patients expressed explicit memory"</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of TIVA/N₂O no/muscle relaxants induction yes/maintenance no/BIS blinded</p> <p>Midazolam (0.1 mg/kg) premed; induction: fentanyl bolus 1 mg to 3 mg/kg, neuromuscular blockers only induction if intubation; LMA used also; maintenance: manual dosing of fentanyl</p> <p>Bispectral index recorded in a blinded manner</p> <p>Quote: "Neuromuscular blockers were given only during anaesthesia induction to facilitate intubation, and no further intraoperative doses were given"</p> <p>Quote: "Participating anaesthesiologists were allowed to perform total IV propofol anaesthesia supplemented by fentanyl but without nitrous oxide"</p> <p>Time of outcome determination: 12 h to 24 h postoperative</p> <p>Method of outcome determination: interview</p> <p>Benno Rehberg benno.rehberg-klug@hcuge.ch</p> <p>8 December 2013; responded 9 December 2013, 10 March 2011, 24 January 2011 re. muscle relaxant use; dropout rate, characteristics table</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote from Dr. Rehberg: "computer-generated list of consecutive allocation to the groups, no blocks" (email bias survey, see notes)
Allocation concealment (selection bias)	Unclear risk	Quote from Dr. Rehberg: "concealed envelopes, opened immediately before induction" (email bias survey, see notes) Comment: unclear if sealed and opaque
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "Anaesthesiologists, who were blinded towards the BIS..." Comment: SCP not blinded

Rehberg 2007 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	High risk	Quote from Dr. Rehberg: "Potential bias: Assessors were not blinded to treatment" (email bias survey)
Incomplete outcome data (attrition bias) All outcomes	Low risk	<p>Comment: outcome data are missing in both intervention groups, imbalanced as percentage units because of unplanned ICU transfer. TIVA: (2/46 = 5%) vs manual controlled infusion (4/46 = 9%); but no significant difference between both groups in dropouts, Peto OR 9.85 (0.51 to 188.36); severely ill patients not interviewed are at higher risk of awareness due to their unstable intraoperative clinical condition (see secondary outcomes), which frequently results in the anaesthesiologist lightening the anaesthetic as part of the treatment of that unstable condition that is often manifest by a common final pathway of hypotension</p> <p>Author sent characteristics and ROB table grades for comment 8 December 2013: Benno Rehberg benno.rehberg-klug@hcuge.ch</p> <p>Quote from Dr. Rehberg 9 December 2013: "I agree with your assessment in general. Although none of the patients admitted to the ICU were unstable during surgery, it is difficult to assess their risk of awareness in comparison to the other patients in retrospect (which was not our primary outcome)"</p> <p>Quote: "Based on performing a statistical analysis and information from Dr. Rehberg we have changed the grade to low risk"</p>
Selective reporting (reporting bias)	Low risk	<p>Quote from Dr. Rehberg: "The study protocol is available, but written in German language. However, the published report includes all pre-specified outcomes" (email bias survey, see notes)</p> <p>Quote: "The main (and other) outcome reported was actually non-significant" (email bias survey, see notes)</p> <p>Comment: awareness outcome part of inclusion criteria</p>
Other bias	Low risk	<p>Quote from Dr. Rehberg: "Potential bias: Assessors were not blinded to treatment." (email bias survey)</p> <p>Comment: that bias is accounted for in the "Blinding of outcome assessment (detection bias)" domain. However, no other risk identified.</p>

Renna 2000

Methods	Study design: randomized parallel groups Study dates: not given
Participants	Country: UK Sex: female Age: mean 41.7 ASA: NA Procedure: minor gynaecological surgery Study size: 48
Interventions	Randomized portion of anaesthetic: volatile agent different doses/word test: implicit memory/positive and neutral suggestion

Renna 2000 (Continued)

Intervention 1: sevoflurane 1.2% ET + either word list A or B + either positive or neutral suggestion, N = 16

Intervention 2: sevoflurane 1.5% ET + either word list A or B + either positive or neutral suggestion, N = 16

Intervention 3: sevoflurane 2% ET + either word list A or B + either positive or neutral suggestion, N = 16

Outcomes

Primary outcomes: primary outcome implicit memory; secondary outcome: awareness

BIS, word lists, neutral suggestion

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 2

Quote: "...there was **evidence of implicit memory**...There was no evidence of a therapeutic effect of positive suggestion...The bispectral index ...not... statistical significance ...indicator... susceptibility to priming....no patient had recall of priming or intra-operative events"

Notes

Non-randomized portion of anaesthetic: volatile agent/N₂O no/100% O₂/facemask/no muscle relaxants/ADM BIS recorded

Breathing induction and maintenance: sevoflurane and **O₂ only** with facemask; at each of the 3 target ET points in induction **word tests** and suggestion transmitted by headphones

Quote: "Anaesthesia was **induced** by inhalation of **sevoflurane in oxygen via a facemask**. At loss of eyelid reflex...After the target end-tidal concentration...stable...bi spectral index ...noted, headphones ...common two-syllable words ...prime implicit memory... then ...neutral ('You will be having your operation today') or the positive ('You will feel great after the operation!') suggestion...Surgical anaesthesia ...established ...operation performed. **At no point during surgery did patients receive drugs other than sevoflurane and oxygen**"

Comment: one of the **few volatile agent only anaesthetics**

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Allocation was randomised by drawing a code from a hat, which ensured 16 patients in each concentration group and equal numbers of A and B, and of positive and neutral suggestion, within each concentration group"
Allocation concealment (selection bias)	Unclear risk	Quote: "Each patient was allocated to one of three groups characterised by their target end-tidal sevoflurane concentration (1.2, 1.5 or 2%), then to a subgroup, A or B, to balance the test of perceptual facilitation, and finally, to a second subgroup for positive or neutral suggestion (a stimulus requiring semantic processing)..."
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: not blinded to sevoflurane doses
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "...Two to three hours after the operation, each patient was interviewed and asked if they remembered hearing anything while they were asleep"

Renna 2000 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "One patient was not included in the analysis because she required propofol to control her movements during the excitement phase of inhalational induction" Comment: dropout rate 1/48 (2%); high-risk awareness dropout; imbalanced across groups but no significant difference between groups: sevoflurane 1.2% (1/16) vs sevoflurane 2.5% and 2% (0/32), Peto OR 0.05 (0.00 to 3.18)
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Rinaldi 2005

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Italy Sex: both Age: 49, 52 ASA: I-II Procedure: major abdominal surgery Study size: 100
Interventions	Randomized portion of anaesthetic: volatile (potent inhalation) technique ADM AAI (AAI 20 ± 5) vs SCP Intervention 1: sevoflurane titrated AAI (AAI 20 ± 5) + fentanyl 1 µg/kg PRN, N = 50 Intervention 2: sevoflurane titrated clinical signs (SCPs) + fentanyl 1 µg/kg PRN, N = 50
Outcomes	Primary outcomes: sevoflurane consumption, time to recovery, memory Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: no patients showed signs of awareness or had any recall postoperatively Quote: "No patients experienced explicit memory and there was not significant difference in implicit memory between Groups A and B (P = 0.107)...11 patients had intraoperative dreams and patients had implicit memory of word tests"
Notes	Non-randomized portion of anaesthetic: volatile (potent inhalation) technique/N₂O no: IV hypnotic/PRN narcotic/muscle relaxants both induction/maintenance Premed: diazepam 0.1 mg/kg; induction: fentanyl 2 µg/kg + propofol 1.5 mg to 2 mg/kg + atracurium 0.6 mg/kg + 3 min FIO ₂ 80% + intubation Anaesthetic maintenance: sevoflurane oxygen/air (titrated to randomized groups); atracurium 0.4 mg/kg/h up to 30 min before the end of surgery/morphine 0.07 mg/kg plus ketorolac 0.4 mg/kg as preventive analgesia, and ondansetron 8 mg for postoperative nausea, maintain target AAI 205 Time of outcome determination: 2 h postoperative

Rinaldi 2005 (Continued)

Method of outcome determination: interview

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Patients were randomised using the closed envelope technique in two groups"
Allocation concealment (selection bias)	Unclear risk	Quote: "Patients were randomised using the closed envelope technique in two groups" Comment: we assume this is referring to allocation
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "anaesthetist... performed all anaesthetic procedures and, by the nature of the study, was not blinded"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Implicit and explicit memory has been evaluated by the physician whose voice had been recorded for the implicit memory test. This physician did not perform or neither know the anaesthetic technique, so memory testing was blinded"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Russell 1986

Methods	Study design: randomized parallel groups Study dates: "1982-1984" (email bias survey, see notes)
Participants	Country: UK Sex: female Age: 19 to 74 ASA: NA Procedure: gynaecological surgery Study size: 55
Interventions	Randomized portion of anaesthetic: TIVA vs "inhaled nitrous oxide with IV fentanyl increments ("balanced" regimen)" otherwise described as intravenous anaesthesia Intervention 1: TIVA: etomidate 100 µg/kg/min for 10 min then...O ₂ /air, 100 mc/kg/min for rest of procedure + fentanyl bolus, N = 25

Russell 1986 (Continued)

 Intervention 2: N₂O + fentanyl bolus ("balanced"), N = 30

Outcomes

Primary outcomes: wakefulness; time to recovery

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

 Comments: no recall, 3 dreams, 7% wakeful in the etomidate group, and one 9 dreams; 44% occurrence of recall in the N₂O group

Notes

Non-randomized portion of anaesthetic: TIVA vs "inhaled nitrous oxide with IV. fentanyl increments ("balanced" regimen)" otherwise described as intravenous anaesthesia: muscle relaxants induction yes/maintenance PRN

Premedication: temazepam 10 mg to 20 mg + fentanyl 250 µg

Anaesthesia induction: thiopentone suxamethonium 1 mg to 1.5 mg/kg

 Anaesthesia maintenance: N₂O:O₂ 2:1 vecuronium 4 mg + vecuronium 2 mg as needed + fentanyl 100 µg as needed

 When patient responded to command: thiopentone 100 mg + fentanyl 100 µg or etomidate 5 mg + 100 mc fentanyl; nitrous required significantly more fentanyl boluses and hence is defined as the **"light" anaesthesia group**

 Comment: **balanced anaesthesia** is defined as inhalational agents with small-dose narcotics; others define it as a regimen based on inhaled N₂O with IV fentanyl increments; TIVA anaesthetic is defined as based on a 2-stage infusion of etomidate plus increments of fentanyl; other definitions of balanced anaesthesia: [Lallemand 2003](#); [Lehmann 1985](#); [Lehmann 1992](#); [Myles 1997](#)

Comment: see Dryad topics: definition of light and inadequate anaesthesia; isolated forearm technique (IFT)

 Comment: wakefulness during anaesthesia has been recognized for many years since [Artusio 1955](#) described this state in spontaneously breathing patients under ether anaesthesia. Using the IFT, [Wilson 1980](#) found some 30% of patients were wakeful

Time of outcome determination: before discharge

Method of outcome determination: interview

Author responded to the email bias survey on 4 February 2011. Responses are recorded in the 'Risk of bias' table

Survey response: 4 February 2011, Ian F i.f.@hull.ac.uk

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	<p>Quote: "Patients were randomised according to whether their unit number was odd or even...no one has control over the unit number assigned to a patient. It is true that the unit number was known before inclusion in the study, but since all patients meeting the entry criteria were included and received their appropriate anaesthetic, I do not think this method of randomisation introduced bias" (email bias survey, see notes)</p> <p>Comment: other experts disagree with Dr. Russell Myles 2015 (personal communication). Despite the 4/7 domains in this RCT having a 'high' risk, we have given this RCT an overall ROB rating that is acceptable for use in our meta-analysis in this review</p>

Russell 1986 (Continued)

Allocation concealment (selection bias)	High risk	Quote: "I was the sole investigator and anaesthetist. Allocation could not be concealed" (email bias survey, see notes) Comment: if the anaesthesiologist is not blinded to the drugs then the lack of allocation concealment will not change the impact of the unblinded state of the anaesthesiologist on the ROB
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "Patient" (email bias survey)
Blinding of outcome assessment (detection bias) All outcomes	High risk	Comment: not blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: 3 patients had partial data due to uncontrolled hypertension; 2 in one group, 1 in the other. All patients (N = 55) were included in the awareness evaluation (email bias survey, see notes)
Selective reporting (reporting bias)	Low risk	Quote: "The study protocol is no longer available as this was over 25 years ago. But all of the study's pre-specified (primary and secondary) outcomes that are of interest in the review have been reported" (email bias survey, see notes) Quote: "Yes. While all the primary outcomes were significant, some secondary outcomes were not significant" (email bias survey, see notes) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Comment: from author survey responses: the author believes that for reasons explain above, there were no other biases

Samarkandi 2004

Methods	Study design: randomized parallel groups Study dates: not given
Participants	Country: Saudi Arabia Sex: both Age: 58 Procedure: off-pump coronary bypass surgery Study size: 40
Interventions	Randomized portion of anaesthetic: ADM BIS 40 to 60 vs SCP Intervention 1: SCP, N = 20 Intervention 2: BIS 40 to 60, N = 20
Outcomes	Primary outcomes: anaesthetic consumption, haemodynamics Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Samarkandi 2004 (Continued)

Comment: no recorded event of intraoperative awareness in both groups; no intraoperative dreams

Notes

Non-randomized portion of anaesthetic: TIVA or volatile I? Balanced anaesthesia I (narcotics + muscle relaxant + PRN volatile inhalation agent)/N₂O no/muscle relaxants both induction/maintenance

Anaesthesia induction: sufentanil 1 µg to 1.5 µg/kg + midazolam 0.05 mg to 0.1 mg/kg + rocuronium 0.9 mg/kg

Anaesthesia maintenance: sufentanil 0.2 µg/kg/h + midazolam 1.5 µg/kg/h + rocuronium 0.5 mg/kg/h + **sevoflurane PRN**

Anaesthesia was induced with sufentanil 1 µg to 1.5 µg/kg, midazolam 0.05 mg to 0.1 mg/kg and rocuronium 0.9 mg/kg then a maintenance sufentanil 0.2 µg/kg/h, midazolam 1.5 µg/kg/h and rocuronium 0.5 mg/kg/h **supplemented with sevoflurane as required**. Induction doses as well as anaesthetic maintenance supplementation doses were guided by the BIS range 40 to 60; Group 2 control group, only clinical judgement was used to titrate

Comment: postoperative standardized questionnaire uses the memory of intraoperative dreams as the starting point to assess explicit memory as do other authors like [Pedersen 1992](#)

Comment: author seems to state that the administration of a **volatile agent was used for inadequate anaesthesia (PRN)** in both groups; emailed author but no response for clarification of this issue; if this is the case, this is another example of the variable definitions of **TIVA**

Comment: see appendix for discussion of the definitions of balanced anaesthesia

Time of outcome determination: 2 days postoperative

Method of outcome determination: standardized questionnaire

Dr. Mohamed Essam Abdel-Meguid, email: memeguid@hotmail.com ROB survey. We emailed on 19 April 2015: recipient failed permanently.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	Quote: "Randomisation was performed using patient's medical record number, being odds related to Group I and evens related to Group II"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor: "Postoperatively, patients were visited on the 2nd postoperative day by one of the medical staff who was blinded about the grouping and they were asked to answer a standardized questionnaire including"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Sareen 1997

Methods	Study design: randomized parallel groups Study dates: not given
Participants	Country: Canada Sex: both Age: 63 ± 9; 65 ± 7 ASA: not given Procedure: elective CABG Study size: 34
Interventions	Randomized portion of anaesthetic: TIVA or "balanced" narcotic portion: 2 doses compared Intervention 1 induction: Group L: sufentanil 3 µg/kg, N = 17 Intervention 2 induction: Group H: sufentanil 15 µg/kg, N = 17
Outcomes	Primary outcomes: haemodynamics, EEG Quote: "effect of a five-fold variation in sufentanil dose on the haemodynamic and electroencephalographic (EEG) response to anaesthetic induction and tracheal intubation" Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "no patient had awareness"
Notes	Non-randomized portion of anaesthetic: TIVA or volatile I "balanced" anaesthesia I (narcotics + muscle relaxant + PRN volatile inhalation agent)/N₂O no/muscle relaxants both induction/maintenance/ADM EEG spectral edge LIFESCAN recorded Premedication: 60 µg/kg lorazepam PO induction and maintenance: sufentanil + vecuronium 0.15 mg/kg, 0.15 mg/kg vecuronium, Neurometrics (Lifescan) aperiodic analysis used as ADM Comment: nomenclature between RCTs is confusing Time of outcome determination: NA Method of outcome determination: interview Email survey sent to author, no response received ithomson@sbrc.umanitoba.ca

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Patients were randomly assigned to one of two groups"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias)	Low risk	Quote: "Sufentanil was prepared by our pharmacy in concentrations of either 10 mcg/ml or 50 mcg/ml and administered in a double blind fashion"

Sareen 1997 (Continued)

All outcomes

Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Schultetus 1986

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: USA Sex: female Age: 23.5, 25.5 ASA: I-II Procedure: elective caesarean section Study size: 36
Interventions	Randomized portion of anaesthetic: IV: opioid - intravenous technique Intervention 1: induction ketamine 1 mg/kg, N = 12 Intervention 2: induction thiopental 4 mg/kg, N = 13 Intervention 3: induction ketamine 0.5 mg/kg + thiopental 2 mg/kg, N = 11
Outcomes	Primary outcomes: haemodynamics, fetal blood gas tensions, APGAR Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: 3 patients (8%): awareness : 1 in thiopental group and 2 in combination group. 4 had intra-operative dreams : 2 thiopental and 2 combination groups; 12 followed commands (wakefulness): 1 from ketamine, 7 thiopental, 4 combinations; 17 reaching movements : 9 from thiopental and 8 from combinations
Notes	Non-randomized portion of anaesthetic: IV: opioid - intravenous technique/muscle relaxants yes both induction/maintenance/N₂O yes Preoperatively sodium citrate 30 mL of 0.3 M + d-tubocurarine chloride 0.07 mg/kg; induction: succinylcholine 2 mg/kg than continuous infusion succinylcholine; anaesthesia maintenance: 70% N ₂ O in O ₂ + succinylcholine chloride 2 mg/min; after delivery fentanyl (100 µg) was given

Schultetus 1986 (Continued)

Comment: "balanced" means "inhaled" agent N₂O not volatile, by our rules define this as IV anaesthesia

Time of outcome determination: postoperative

Method of outcome determination: interview

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "scheduled for elective cesarean section and general anaesthesia were assigned by a randomized code to receive..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Schwender 1994

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Germany Sex: both Age: 57.4, 61.6, 61.2, 56.8 ASA: NA Procedure: elective cardiac surgery Study size: 45
Interventions	Randomized portion of anaesthetic: parts of IV: opioid vs TIVA vs parts of balanced anaesthesia II (volatile agent + supplemental narcotics ± muscle relaxant) induction/maintenance:

Schwender 1994 (Continued)

IV vs TIVA vs volatile agent continuous administration (balanced II) + music implicit memory tests/MCI

Intervention 1: induction flunitrazepam 0.01 mg/kg + fentanyl 0.01 mg/kg; maintenance **flunitrazepam** 1.2 mg/h, N = 10

Intervention 2: induction etomidate 0.25 mg/kg and fentanyl 0.005 mg/kg; maintenance **isoflurane** (0.6 to 1.2 vol%), N = 10

Intervention 3: induction etomidate 0.25 mg/kg and fentanyl 0.005 mg/kg; maintenance **propofol** 4 mg to 8 mg/kg/h, N = 10

Intervention 4: the group 4 patients were **not exposed to the audiotape** and those patients were assigned **randomly to 1 of the 3 anaesthetic** regimes, N = 15

Audiotape with implicit memory task using the Robinson Crusoe story

Outcomes

Primary outcomes: implicit memory interview, MLAEP

Quote: "The main goals of this study were to determine whether explicit or implicit memory is present during cardiac surgery, and if so, whether memory functions can be related to specific characteristics of MLAEP."

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 2

Quote: "No patient had a clear explicit memory of intraoperative events. However, there were statistically significant differences in the incidence of implicit recall among the groups. Five patients in the flunitrazepam- fentanyl group, 1 patient in the isoflurane-fentanyl group, 1 patient in the propofol-fentanyl group, and no patient in the control group showed an implicit memory"

Comment: implicit recall was related to continued presence of MLAEP

Notes

Randomized portion of anaesthetic: nitrous N/parts of IV: opioid vs TIVA vs parts of balanced anaesthesia II (volatile agent + supplemental narcotics ±muscle relaxant)/muscle relaxants yes both induction/maintenance: ADM MLAEP

IV vs TIVA vs volatile agent continuous administration (balanced II) ± music implicit memory tests/ADM MLAEP recorded

Premedication: benzodiazepine (flunitrazepam 1 mg to 2 mg orally) 45 to 60 min before anaesthesia. Anaesthesia maintenance: **high-dose fentanyl analgesia** 1.2 mg/h + pancuronium 0.1 mg/kg. In addition, **group 1** received **flunitrazepam** (1.2 mg/h), **group 2 isoflurane** (0.6 to 1.2 vol%), and **group 3 propofol** (4 to 8 mg/kg/h) to maintain general anaesthesia, pancuronium 0.1 mg/kg. The patients of **group 4** (n = 15) were **randomly assigned** to 1 of the 3 anaesthetic regimes. **MALEP**

Comment: see Dryad topic: details regarding this RCT's control group; implicit memory test, portions of novel Robinson Crusoe used as a parable; and associated with "Friday" and postoperative recovery

Comment: the ability to retrieve the **memory of implicit tasks** are a **more sensitive measure of memory during anaesthesia than explicit tasks**

Time of outcome determination: 3 to 5 hours in postoperative period

Method of outcome determination: interview

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Schwender 1994 (Continued)

Random sequence generation (selection bias)	Unclear risk	Quote: "those patients were assigned randomly to one of the anaesthetic regimes..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: anaesthesiologist: see below
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor: "All experimental evaluations were conducted under double-blind conditions: neither the patients nor the interviewer knew which anaesthetic had been used or whether an audio tape had been presented" Comment: this is an example where the term "double-blind" cannot be assumed to indicate the anaesthesiologist was blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Schwender 1996

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Germany Sex: both Age: 21 to 79 ASA: I-II Procedure: elective gynaecological or urological laparotomy Study size: 47
Interventions	Randomized portion of anaesthetic: IV vs volatile agent Comment: as per rules: N₂O used hence IV not TIVA anaesthesia Intervention 1: induction: epidural + thiopentone 5 mg/kg; maintenance isoflurane 0.4 to 1.2 vol%, N = 23 Intervention 2: induction: epidural + propofol 2 mg/kg; maintenance propofol 3 mg to 5 mg/kg/h, N = 24
Outcomes	Primary outcomes: sensitivity of movement/specificity adequate anaesthesia associated with threshold levels of SEF 90

Schwender 1996 (Continued)

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Quote: "No patient had explicit recall of intraoperative events"

Notes

Non-randomized portion of anaesthetic: regional anaesthesia: epidural and GETGA parts of TIVA vs parts of volatile agent/N₂O yes + supplemental narcotics + muscle relaxants yes both induction/maintenance ADM SEF recorded

Comment: as per rules: N₂O used hence IV not TIVA anaesthesia

Premedication: oral clorazepate (benzodiazepine) 45 to 60 minutes before surgery; anaesthesia induction: epidural + randomized inhalation vs IV + vecuronium 0.1 mg/kg + **N₂O** 50% + O₂

Anaesthesia maintenance: see intervention maintenance; no muscle relaxants for maintenance

Comment: PSRT>2 or spontaneous movements defined light anaesthesia

Comment: sensitivity 72% is number successfully predicted movements/total movements; **specificity 82%** number correctly predicted situations of adequate anaesthesia/number situations of adequate anaesthesia threshold 14 Hz GETA + epidural to minimize pain

Comment: there was **no correlation between haemodynamic data and patient movement**

Comment **epidural blocks spinal cord and ablates reflex movement**

Comment: see Dryad topic purposeful movement in setting of NO reflex movement

Comment: see Dryad topic: **wide range of BIS levels** associated with transition into **unconsciousness and consciousness** under anaesthesia

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "patients allocated randomly to 2 groups"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Shin 2012

Methods	<p>Study design: randomized parallel groups</p> <p>Study dates: not stated</p>
Participants	<p>Country: Korea</p> <p>Sex: female and male</p> <p>Age: 37-38</p> <p>ASA: I</p> <p>Procedure: elective orthopaedic or extremity surgeries</p> <p>Study size: 90</p>
Interventions	<p>Randomized portion of anaesthetic: parts of volatile agent 3 doses ±infusion of narcotics (TCI)</p> <p>Intervention 1: maintenance: 4% to 6% inspired desflurane + target-controlled concentration (TCI) 1 ng/mL, N = 30</p> <p>Intervention 2: maintenance: 4% to 6% inspired desflurane + (TCI) 2 ng/mL remifentanil, N = 30 - 2 = 28</p> <p>Intervention 3: 7% to 9% inspired desflurane only without remifentanil infusion</p> <p>BIS both groups, N = 30</p>
Outcomes	<p>Primary outcomes: Quote: "optimal target-controlled concentration of remifentanil combined with desflurane, by using a more widely and decreasing end-tidal concentration of desflurane"</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Comment: awareness events: none</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent/N₂O no + muscle relaxants yes for both induction/maintenance/ADM BIS recorded</p> <p>Induction propofol 2 mg/kg and lidocaine 0.5 mg/kg, vecuronium 0.15 mg/kg IV intubation vecuronium 0.02 mg/per stimulator algorithm</p> <p>Comment: inhalation vs inhalation (balanced anaesthesia): desflurane vary ET with varied infusion rate remifentanil: BIS both groups</p> <p>Comment: infusion remifentanil allows a wider range of volatile gent: see Dryad topic</p> <p>Emailed Dr Shin, smkeun311@yahoo.co.kr, the ROB survey 3 January 2014</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Randomization into one of the three groups was based on an Excel random-number generation"
Allocation concealment (selection bias)	Unclear risk	<p>Comment: inadequate information provided</p> <p>Quote: "...were registered and randomly allocated to receive either a target-controlled concentration"</p>

Shin 2012 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: no evidence of blinding. TIVA vs volatile agent.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: inadequate information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: 2 patients excluded: systolic blood pressure was below 90 mmHg (1 in group R2) and BIS > 60 >= 2 min (1 in group R2) treated immediately. Post-operative interview: no awareness report; following up with the awareness interview in these 2 excluded patients allowed us to grade the missing data as low risk. There was no significant difference between groups due to exclusions (2/60 vs 0/30), Peto OR 4.56 (0.24 to 87.68) (P = 0.31)
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information provided

Sidi 1990

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Israel and USA Sex: female and male Age: 18 to 70 ASA: III or IV Procedure: elective cardiac surgery with cardiopulmonary bypass (CPB): CABG, aortic valve replacement with/without mitral valve replacement, and mitral valve replacement with tricuspid valve valvuloplasty Study size: 32, 29
Interventions	Randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia for intubation only: ADM: EEG/CSA vs SCP Intervention 1: induction: fentanyl 50 µg/kg with no EEG monitoring, N = 16 Intervention 2: induction: fentanyl 25 to 50 µg based on changes in EEG , N = 16
Outcomes	Primary outcomes: EEG associated haemodynamic changes secondary to induction/laryngoscopy Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "No patient in either group had recall"
Notes	Non-randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia for intubation only/N₂O no: muscle relaxants induction yes

Sidi 1990 (Continued)

Premed: morphine 10 mg IM/diazepam 10 mg PO; induction:midazolam (0.015 to 0.03 mg/kg), and fentanyl (25 to 50 µg/kg)/pancuronium (0.15 mg/kg) 100% O₂, volatile anaesthetics were not used before endotracheal intubation

Comment: IV anaesthesia intubation titrated to fentanyl; dose based on EEGs vs control: fixed dose; IV vs IV: ADM: EEG: CSA: open/closed

Comment: authors describe technique as IV anaesthesia for induction; no post intubation description of maintenance anaesthesia

Inadequate anaesthesia protocol (see Dryad)

No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "16 were randomly assigned to be monitored with continuous EEG starting immediately before anesthetic induction. The remaining patients were monitored as the others but without EEG"
Allocation concealment (selection bias)	Unclear risk	Comment: inadequate information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: inadequate information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "None of the patients in the two groups reported recall of induction, intubation, or intraoperative events"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: if hypertension persisted or tachycardia occurred before or during induction, the patient was excluded from the study EEG group 2 exclusions with missing data; SCP group (fixed fentanyl dose) 3 exclusions 1 hypertension peri-intubation, 2 died on CPB; NS (2/16 vs 3/16), Peto OR 0.63 (0.10 to 4.13) (P = 0.63); NS also for high-risk awareness analysed as a subgroup: (0/16 vs 3/16) Peto OR 0.12 (0.01 to 1.22) (P = 0.07)
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: inadequate information provided

Smith 1999

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: USA Sex: male and female ASA: I-II

Smith 1999 (Continued)

Age: 18 to 65

Procedure: any elective surgery

Study size: 101 enrolled, 81 completed surgery 20% (20/101) dropped out

Interventions	<p>Randomized portion of anaesthetic: parts of intravenous (IV) - TIVA vs inhalational anaesthesia: midazolam vs sevoflurane</p> <p>Intervention 1: maintenance: midazolam 0.540 µg/kg + fentanyl 4.5 µg/kg/h + N₂O 50%, N = 23</p> <p>Intervention 2: intervention 1 + word test, N = 28</p> <p>Intervention 3: maintenance: 1.4% ET sevoflurane + fentanyl + 50% N₂O, N = 25</p> <p>Intervention 4: intervention 3 + word test given and correlated with, N = 25</p> <p>Auditory middle latency response (AMLR) recorded all groups</p>
Outcomes	<p>Primary outcomes: response from stem-word completion test</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4</p> <p>Comment: awareness: 2 midazolam group: 1 heard auditory clicks, 1 felt drill pressing into his leg</p> <p>Comment: implicit memory (defined: completion of word stem test) compared to control group: patient scores: sevoflurane no relationship (P = 0.07); midazolam (P = 0.07)</p>
Notes	<p>Non-randomized portion of anaesthetic: N₂O yes/parts of intravenous (IV) vs inhalational anaesthesia: midazolam vs sevoflurane/muscle relaxants induction yes/PRN maintenance: AMLR recorded: implicit memory test</p> <p>IV group: induction: lidocaine 20%, propofol 1.5 mg to 2.5 mg/kg, fentanyl 1.0 µg to 3.5 µg/kg, succinylcholine 1.5 mg/kg or atracurium 0.4 mg/kg/intubation; maintenance: midazolam 0.54 µg/kg, fentanyl 4.5 µg/h, 50% N₂O 50% O₂, all infusions except atracurium PRN</p> <p>Inhalational group: maintenance: sevoflurane 1.4% ET, fentanyl 1.4 µg/kg/h, N₂O 50%</p> <p>Comment: see Dryad topics: AMLRs association to implicit memory, defined IV anaesthesia includes N₂O</p> <p>No email address available for ROB survey</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "randomly assigned"
Allocation concealment (selection bias)	Unclear risk	Comment: no information
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Quote: "double blinded"
Blinding of outcome assessment (detection bias)	Unclear risk	Quote: "double blinded"

Smith 1999 (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	Low risk	<p>Comment: 18% of patients enrolled were excluded from statistical analysis: of 101, 12 eliminated due technical issues related to AMLR recording and memory testing; 6 excluded due to lack of preoperative implicit memory tests (groups not specified) - these do not impact awareness rate; 2 midazolam patients excluded: postoperative explicit memory (awareness)</p> <p>There is no significant difference between groups if we analyse the awareness rates by adding the 2 patient awareness reports to the midazolam group in an intention-to-treat analysis (N = 101, Peto OR 7.39 (0.46 to 119.86)) and after the non-high awareness risk patients were excluded (N = 89, Peto OR 6.79 (0.42 to 110.77), N = 83, Peto OR 6.75 (0.41 to 110.34))</p>
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information

Song 1997

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: USA Sex: female Age: 27, 28 Procedure: laparoscopic tubal ligation Study size: 60
Interventions	<p>Randomized portion of anaesthetic: parts of volatile agent/ADM (BIS 60) vs SCP</p> <p>Groups I and II: maintenance: desflurane, 2% to 5%, BIS open (60) vs closed (SCP), N = 30</p> <p>Group III and IV: maintenance: sevoflurane, 0.7% to 2%, BIS open (60) vs closed (SCP), N = 30</p>
Outcomes	Primary outcomes: times to awakening and extubation Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "None of the patients reported recall of intraoperative events when questioned at the time of discharge from the hospital"
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent/N₂O/supplemental narcotics/muscle relaxants yes for both induction/maintenance/ADM (BIS) vs SCP</p> <p>Control groups (Groups I and III), volatile anaesthetics administered: standard clinical practice, anaesthesiologists blinded to BIS value</p> <p>Induction: midazolam (2 mg), fentanyl 1 µg/kg, propofol 2 mg/kg, succinylcholine, 1 mg/kg, and lidocaine 4% (4 mL)</p> <p>Maintenance: group assignment + N₂O 1 L/min (65%), in oxygen, 0.7 L/min, mivacurium used</p>

Song 1997 (Continued)

In the control groups (blinded to the BIS) (Groups I and III): volatile anaesthetics were adjusted SCP

In the BIS-titrated groups (Groups II and IV): the volatile anaesthetics were titrated to **BIS 60**

All patients were mechanically ventilated

Comment: BIS 60 as titration endpoint used in [Vakkuri 2000](#)

See Dryad topic light anaesthesia with BIS; underpowered studies and type II error

Emailed ROB survey pwhite@medner.swmed.edu 03 January 2014 - failed

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "computer-generated random numbers table"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "One of the criticisms of this study is the possibility of bias as a result of the lack of a double-blind design. However, this clinical investigation was conducted in the context of standard clinical practice, and routine blinding procedures would not be appropriate"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Sorbara 1995

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Italy Sex: both Age: 37 to 70 ASA: NA Procedure: CABG: normal-moderately impaired LV function (ejection fraction ~40%) as assessed by pre-operative LV cineangiography and LV end-diastolic pressure <= 18 mmHg Study size: 30

Sorbara 1995 (Continued)

Interventions	<p>Randomized portion of anaesthetic: parts of volatile agent</p> <p>Intervention 1: maintenance before sternotomy propofol 3 mg/kg/h (N = 15)</p> <p>Intervention 2: maintenance before sternotomy isoflurane 0.6% (IF group) (N = 15)</p>
Outcomes	<p>Primary outcomes: assessment of LV contractility</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Comment: no patient of either group experienced dreaming or had any recall of operative events when questioned</p>
Notes	<p>Non-randomized portion of anaesthetic: N₂O yes/parts of volatile agent + supplemental narcotics + muscle relaxants: yes both induction/maintenance</p> <p>Premedication flunitrazepam 2 mg, orally 90 minutes before surgery</p> <p>Anaesthesia induction: thiopental 1 mg/kg, fentanyl 20 µg/kg, and vecuronium 0.1 mg/kg</p> <p>Maintenance after sternotomy boluses (5 µg/kg) fentanyl (max maintenance dose 30 µg/kg): prophylactic: blunt/intense periods pain/autonomic stimulation (e.g. sternal splitting and spread, aortic mobilization, cannulation, decannulation, and sternal closure) 0.025 mg/kg pancuronium/h</p> <p>Comment: light anaesthesia techniques resulted from the need to use a ceiling dose of IV moderate to long acting anaesthetics in order to extubate the patient at the end of the case; ceiling dose</p> <p>Time of outcome determination: 2 days and 1 week postoperative</p> <p>Method of outcome determination: interview</p> <p>ROB survey sent via email to secondary author on 17 January 2011. No response received yet.</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "On the day before surgery, the patients were randomly allocated to either the PF (N = 15) or IF (N = 15) group"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "All patients were questioned 2 days and 1 week postoperatively about dreaming or recall of operative events"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria

Sorbara 1995 (Continued)

Other bias	Unclear risk	Comment: insufficient information provided
------------	--------------	--

Soyannwo 1988

Methods	Study design: randomized parallel groups Study dates: NA
Participants	Country: Nigeria Sex: female Age: 28-29 ASA: all patients were classified as ASA I or II Procedure: elective or emergency caesarian section Study size: 150
Interventions	<p>Randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia: pethidine (meperidine) (25 mg) and pethidine (25 mg) plus flunitrazepam (benzodiazepine)</p> Intervention 1: maintenance Group A - 25 mg pethidine (control), N = 50 Intervention 2: maintenance Group B - 25 mg pethidine + flunitrazepam 0.015 mg/kg body weight (experiment subcat 1), N = 50 Intervention 3: maintenance Group C - 25 mg pethidine + flunitrazepam 0.030 mg/kg body weight (experiment subcat 2), N = 50
Outcomes	Primary outcomes: Quote: "Effect of flunitrazepam (Rohypnol) on awareness during anaesthesia for Caesarian section". Awareness/wakefulness as defined using an awareness classification system (see Table 1): class 4 Comment: awareness - Group A 14/50 (28%), Group B - 10/50 (20%), Group C - 2/50 (4%); total awareness incidence 17% (26/150) Quote: "...six felt pain at the beginning of surgery while three felt pain throughout the procedure"
Notes	<p>Non-randomized portion of anaesthetic: N₂O yes/parts of intravenous (IV) anaesthesia: N₂O/narcotic/hypnotic/muscle relaxants yes both induction/maintenance</p> Anaesthesia induction: 250 mg thiopentone sodium intravenously + 0.6 mg atropine + 100 mg suxamethonium chloride. Anaesthesia maintenance: N₂O and oxygen 41:21/min, and pancuronium bromide or fazadinium bromide in appropriate doses. After delivery pethidine (analgesic) + flunitrazepam Comment: the technique using intravenous agents and N ₂ O is not described by author as intravenous technique etc. The incidence of awareness during obstetric anaesthesia: see Dryad topic frequency awareness with nitrous-based techniques Time of outcome determination: within 72 h Method of outcome determination: questionnaire No email address available for ROB survey

Risk of bias

Soyannwo 1988 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "They were randomly divided into three groups of 50 patients each"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Staikou 2013

Methods	Study design: randomized parallel groups Study dates: 3 November 2008 to 14 July 2011
Participants	Country: Greece Sex: male/female Age: 20 to 70 years old ASA: I-II Procedure: surgery under general anaesthesia (specific type not described) Study size: 84 84 enrolled - 78 started - 72 completed study
Interventions	Randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia for intubation only: lidocaine vs placebo (saline) Intervention 1: normal saline Intervention 2: lidocaine 1.5 mg/kg
Outcomes	Primary outcomes: the impact of lidocaine pretreatment on BIS values on rapid sequence induction and intubation (RSII) Secondary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: "none of the patients reported awareness/recall of the procedure"
Notes	Non-randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia for intubation/N₂O yes only: maintenance: no information BIS recorded

Staikou 2013 (Continued)

No post-intubation anaesthetic technique described: classified as "other" technique

Propofol 2 mg/kg, lidocaine or normal saline and rocuronium 1 mg/kg, trachea was intubated; difficult intubation excluded

ET CO₂ 35 to 40 mmHg and sevoflurane 1% ET/N₂O-oxygen mixture (FIO₂: 0.45). Opioids spared during the study period; **BIS scores recorded** by investigator blinded to patient's allocation group

Chryssoula Staikou MD email: c_staikou@yahoo.gr; ROB survey email sent 25 January 2015; email query re: maintenance anaesthetic, signs of light anaesthesia 22 March 2015; response on 24 March 2015

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The study is also registered in the ClinicalTrials.gov protocol registration system (NCT01238718)" Quote from ROB survey response: "Patients were randomly assigned by the use of sealed envelopes describing the group of assignment"
Allocation concealment (selection bias)	Unclear risk	Quote from ROB survey response: "Patients were randomly assigned by the use of sealed envelopes describing the group of assignment"
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote from ROB survey response: "The patients received either lidocaine 1.5 mg/kg or normal saline, both prepared... an independent investigator ...The anaesthesiologist (other investigator) was blinded to group allocation...Recordings were made by the anaesthesiologist who was blinded to patient's allocation group" Comment: assume that the syringes are blinded
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote from ROB survey response: "...investigators interviewed patients about awareness during surgery with an informal interview..."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: 84 patients enrolled, 6 excluded: 4 for technical reasons and 2 for airway problems during intubation that made them high-risk awareness dropouts. The 2 were balanced between both groups; hence no downgrade, Peto OR 0.89 (0.05 to 14.63)
Selective reporting (reporting bias)	Low risk	Quote from ROB survey response: "The study protocol is available and all of the study's pre-specified (primary and secondary) outcomes that are of interest in the review have been reported in the pre-specified way." Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote from ROB survey response: "all results were reported. There may be some limitations in the study: the lag time of the used monitor...we do not consider that it affected our results... timing of administration of the drug (lidocaine = intervention) ...clearly described in the manuscript ..."

Stoppe 2012

Methods	Study design: randomized parallel groups Study date: unknown
---------	---

Stoppe 2012 (Continued)

Participants	<p>Country: Germany Sex: both Age: 40 ASA: I and III Procedure: elective abdominal surgery: gynaecologic or urologic abdominal surgery Study size: 42 (2 excluded: 40 statistical analysis)</p>
Interventions	<p>Randomized portion of anaesthetic: volatile agent types (sevoflurane vs xenon)</p> <p>Intervention 1: sevoflurane (1 to 1.4 vol% MAC) titrated SCPs, N = 21</p> <p>Intervention 2: xenon (53 to 56 vol% MAC) titrated AEP, N = 21</p> <p>AEPs compared to BIS</p>
Outcomes	<p>Primary outcomes: impact of xenon and sevoflurane on auditory-evoked potentials as assessed by aepEX monitor</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Quote: "After clinical observation and postoperative Brice interview, no signs of awareness were recorded"</p>
Notes	<p>Non-randomized portion of anaesthetic: N₂O no/parts of volatile agent + infusion of narcotics + muscle relaxant(s) induction yes/maintenance unclear/ADM (BIS/AEP) vs SCP recorded simultaneously</p> <p>Balanced volatile inhalation (narcotic supplementation, remifentanil infusion): ADM (AEP) vs SCP: xenon vs sevoflurane (AeP compared BIS)/remifentanil infusion</p> <p>Induction: propofol/remifentanil infusion/rocuronium; maintained by remifentanil (0.15 µg/kg/m) and either sevoflurane (1 to 1.4 vol% MAC) or xenon (53 to 56 vol% MAC)</p> <p>Intraoperative awareness Brice questionnaire at 2 h and 12 h after anaesthesia</p> <p>Primary outcome: "auditory-evoked potentials as assessed by aepEX monitor...xenon...aepEX-derived values: compared: BIS and control group with sevoflurane"</p> <p>Secondary endpoints: relevant outcome and recovery parameters: "Aldrete and Myles score and assessment: intraoperative awareness"</p> <p>ROB survey response mcoburn@ukaachen.de</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Comment: the study was registered at the European Medicines Agency (EudraCT number: 2008-004132-20) and at ClinicalTrials.gov (NCT number: 00793663). Nonetheless, there are no details about the method of randomized allocation and other domains ; ROB survey: "computer based randomization"
Allocation concealment (selection bias)	Unclear risk	Comment: ADMs are blinded to anaesthesiologist; description of sealed opaque envelope or similar concealment protocols not clear; emailed on 20 April 2015 for clarification
Blinding of participants and personnel (performance bias)	High risk	Quote: "The performing attending anaesthetist was blinded to the BIS and aepEX values" Comment: not blinded to SCPs

Stoppe 2012 (Continued)

All outcomes

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "...Brice questionnaire was performed at 2 and 12 h after end of anaesthesia by an independent physician"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: 1 high-risk awareness dropout in xenon group is not significantly different from control group if we assume that dropout would have submitted an awareness report, Peto OR 7.39 (0.15 to 372.38)
Selective reporting (reporting bias)	Low risk	Comment: ROB survey: states that there was no selective reporting Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote ROB survey: "1. xenon anaesthesia ...not commonly used... 3...BIS monitoring ...shown to provide reliable measurement of hypnotic depth, the validity of this monitoring remains controversial. Therefore ...AepEX monitoring does not necessarily reflect a true assessment of anaesthetic depth but can only state that hypnotic depth monitoring using aepEx was comparable to BIS measurement and clinical evaluation" Comment: these can be described as possible causes of diversity rather than form of internal bias (other bias)

Stuttman 2010

Methods	Study design: randomized parallel groups Study date: Quote: "1/2003 - 12/2004" (email bias survey, see notes)
Participants	Country: Germany Sex: both Age: 40 ASA: I and II Procedure: lumpectomy, mammoplasty, liposuction, arthroscopy Study size: 61
Interventions	Randomized portion of anaesthetic: volatile agent types Intervention 1: maintenance: xenon (ET 63%) (N = 31) Intervention 2: maintenance: isoflurane in N ₂ O (ET 0.6%) (N = 30)
Outcomes	Primary outcomes: compare xenon with isoflurane anaesthesia impact on early cognitive recovery with the syndrome short test (SST) Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 0 Quote: "no patient reported awareness on the visit one day postoperatively"
Notes	Non-randomized portion of anaesthetic: N₂O yes/parts of volatile agent + infusion of narcotics MCI + muscle relaxant(s) induction yes/maintenance none/BIS 40

Stuttman 2010 (Continued)

Anaesthesia induction: propofol (1 mg to 2 mg/kg IV) + propofol, 0.003 mg/kg fentanyl and 0.6 mg/kg + rocuronium

Anaesthesia maintenance: see interventions; infused propofol/fentanyl (0.0015 mg/kg BW) PRN. Rocuronium bromide was not repeated during the operation

Time of outcome determination: 1 day after surgery

Method of outcome determination: interview

ROB survey email sent 21 February 2011

Email survey response: 25 February 2011, Stuttman ralph.stuttman@bergmannstrost.com

Author sent characteristics and ROB table grades for comment 8 December 2013: ralph.stuttman@bergmannstrost.com

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Patients were randomised in variable blocks of 4-8 patients in order to balance the groups using simple self-programmed software..."
Allocation concealment (selection bias)	Unclear risk	Quote: "...cards were in envelopes signed with numbers to recognize different block size. Randomization was done by Rolf Lefering, IFOM in cologne, a very experienced statistical experts in medical studies." (email bias survey, see notes) Comment: unclear if opaque, sealed envelopes Comment: no details regarding the 7 domains at http://www.controlled-trials.com/search?q=01110844
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "Patient, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patient, Awareness outcome assessor (interviewer)" (email bias survey, see notes) Comment: assessor: the investigator measuring the SST pre- and postoperatively was a medical assistant and was blinded for the inhalational anaesthetic utilized
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "...one patient was excluded because of intra-operative hypertension. The method to conduct anaesthesia was changed in this case" (email bias survey, see notes) Comment: there is no significant difference in awareness rates between both groups if we assume the hypertensive dropout at high risk of awareness is placed in the experimental or control group; when Peto OR analysis is applied there is no significant difference with exclusion in either group: Peto OR 0.13 (0.00 to 6.60) Author sent characteristics and ROB table grades for comment 8 December 2013: ralph.stuttman@bergmannstrost.com
Selective reporting (reporting bias)	Low risk	Quote: "the study protocol is available and all primary and secondary outcome data were recorded completely in the pre-specified way..." (email bias survey, see notes)

Stuttman 2010 (Continued)

Trial registration: the trial was registered with the number ISRCTN01110844 at <http://www.controlled-trials.com/isrctn/pf/01110844>

Comment: awareness outcome part of inclusion criteria

Other bias	Low risk	Quote: "there was...no baseline imbalance because a block design was used to select patients. The test to prove cognitive function ...A learning effect in the test was compensated by using two different test designs. In our opinion there was no important risk of bias." (email bias survey, see notes)
------------	----------	--

Toft 1987

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Denmark Sex: both Age: 52.8, 53.5 ASA: not given Procedure: bronchoscopy, mediastinoscopy or laryngoscopy, or combinations of these procedures Study size: 50
Interventions	Randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia (neurolept i.e. ketamine dissociative) induction Intervention 1: induction infusion ketamine 250 mg + midazolam 12.5 mg in 250 mL of 0.9% NaCl, N = 25 Intervention 2: induction infusion ketamine 250 mg + diazepam 20 mg in 250 mL of 0.9% NaCl, N = 25
Outcomes	Primary outcomes: compare midazolam with diazepam using total intravenous anaesthesia with ketamine/benzodiazepine-fentanyl: anaesthetic consumption, duration, time to recovery Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: "No awareness during anaesthesia was reported"
Notes	Non-randomized portion of anaesthetic: N₂O no/parts of intravenous (IV) anaesthesia (neurolept i.e. ketamine dissociative) muscle relaxants induction yes/maintenance unclear Premedicated with intramuscular morphine 7.5 mg and scopolamine 0.3 mg, pancuronium 0.01 mg/kg and fentanyl 100 µg to 150 µg 5 minutes before the induction of anaesthesia Anaesthesia induction: interventions 1 and 2, succinylcholine 1.5 mg/kg; maintenance:ketamine-benzodiazepine solution + fentanyl 50-100µg PRN as indicated by HR/BP;MR Time of outcome determination: postoperative before discharge Method of outcome determination: interview ROB survey/questionnaire email was sent to author on 14 February 2011 to palle.toft@ouh.regionsyddanmark.dk : no response received

Toft 1987 (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Patients were allocated in a random and double-blind fashion two groups of 25 patients each"
Allocation concealment (selection bias)	Unclear risk	Comment: above
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "The patient, the anaesthetist and the observer were all blinded"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "The patient, the anaesthetist and the observer were all blinded" Quote: "Before leaving the hospital the patients were asked by one of the investigators if they had experienced any dreaming, visual disturbances, or awareness during the anaesthetic"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Toscano 2007

Methods	Study design: randomized parallel groups Study date: "2000" (email bias survey, see notes)
Participants	Country: Italy Sex: female Age: 27 ASA: I and II Procedure: minor gynaecologic surgery Study size: 100
Interventions	Randomized portion of anaesthetic: IV: premedication Intervention 1: pre-medication scopolamine IM (2.5 µg/kg IM) (N = 50) Intervention 2: pre-medication atropine IM (10 µg/kg IM) (N = 50)
Outcomes	Primary outcomes: incidence of dreams, anaesthetic consumption Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Toscana 2007 (Continued)

Comment: although the paper has no mention of patients being asked about recall of intraoperative events after surgery or awareness, the author states in the author's response to our ROB survey: "none had awareness after asking them"

Notes

Non-randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia: N₂O/SR

Anaesthesia induction: propofol bolus (2.5 mg/kg)

Anaesthesia maintenance: propofol infusion (12 mg/kg/h) N₂O propofol as a 2.5 mg/kg bolus, followed by 12 mg/kg

Time of outcome determination: 20 min and 6 h after surgery

Ventilation was assisted manually 70% N₂O/O₂ with **facemask** until the end of surgery

Comment: see Dryad topic - propofol activates basal forebrain cholinergic

Method of outcome determination: interview

Survey response: 14 February 2011, Carlo Pancaro carlopancaro@hotmail.com

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "computer generated, then envelopes" (email bias survey, see notes)
Allocation concealment (selection bias)	Unclear risk	Quote: "computer generated, then envelopes" (email bias survey, see notes)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "syringe was prepared by someone else. Not from people taking care of the patients" (email bias survey, see notes) Quote: "Patient, Anaesthesiologist" (email bias survey, see notes) Quote: "In all subjects, anaesthesia was induced by a second anaesthesiologist, blinded to the premedication drug used"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "Patient, Anaesthesiologist" (email bias survey, see notes) Comment: paper indicates blinded observer for dream interview; survey response does not address this
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: 3% (3/100) dropout rate Quote: "2 patients received inhalational agents, 1 patient received morphine intraoperatively;...these three patients were excluded from the analysis since inhalational agents and morphine can suppress dream activity. However, none of these three patients reported any dreams or awareness" (email bias survey, see notes) Comment: since author assessed the dropouts for awareness, there are no missing data in regard to this outcome
Selective reporting (reporting bias)	Low risk	Quote: "we reported everything" (email bias survey, see notes) Quote: "none had awareness after asking them" Comment: awareness outcome part of inclusion criteria

Toscano 2007 (Continued)

Other bias	Low risk	Quote: "free of bias" (email bias survey, see notes)
------------	----------	--

Tsai 2001

Methods	Study design: randomized parallel groups Study dates: NA
Participants	Country: Taiwan Sex: female Age: 22 to 46 ASA: I and II Procedure: elective caesarean section Study size: 24
Interventions	<p>Randomized portion of anaesthetic: parts of volatile agent anaesthesia: post-delivery: volatile agent vs hypnotic: propofol</p> Intervention 1: maintenance post-delivery isoflurane ET 0.5, N = 12 Intervention 2: maintenance post-delivery propofol 8 mg/kg/h, N = 12
Outcomes	Primary outcomes: BIS values, haemodynamics Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "None of the patients in either group subsequently reported recall of events during th entire course of delivery"
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent anaesthesia: N₂O/narcotic/hypnotic/muscle relaxants yes for both induction/maintenance: BIS target < 75</p> <p>Volatile agent technique supplemented post-delivery with propofol vs isoflurane; classify as other - sevoflurane pre-delivery then post-delivery random propofol vs isoflurane</p> Anaesthesia induction: pentothal (4 mg/kg) + succinylcholine (1 mg/kg) + atracurium (0.5 mg/kg). Anaesthesia maintenance: isoflurane (ET 1.15%) in 50% N ₂ O/50% O ₂ + post-delivery: fentanyl 3 µg/kg + droperidol 5 mg + 67% N ₂ O in O ₂ + atracurium group-specific drug (see above) Comment: this is very light anaesthesia with BIS target < 75 BIS target < 75 , if BIS >= 75 > 5 mins anaesthesia increased; if MAP < 65 >= 3 min after RX bolus 500 mL fluid, decrease isoflurane or propofol; if uterine contraction poor increase oxytocin Time of outcome determination: day after surgery Method of outcome determination: interview Email for ROB table sent to author on 22 February 2011. No email response received yet.

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Tsai 2001 (Continued)

Random sequence generation (selection bias)	Unclear risk	Quote: "...patients were randomly assigned to either of two groups..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor: "The effects ... were detected by a helper ... who was blinded to the anaesthetic given"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "Therefore, there was no exclusion of patients who were initially included in the study..." Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Tunstall 1989

Methods	Study design: randomized parallel groups Study dates: not given
Participants	Country: Scotland Sex: female Age: not given, mean ASA: NA Procedure: elective and semi-elective caesarian section Study size: 113
Interventions	Randomized portion of anaesthetic: volatile agent types: enflurane vs isoflurane Pilot study completed; followed by this protocol: Intervention 1: maintenance 3 min enflurane 3% + enflurane 1.5% + post-delivery enflurane 0.5%, N = 50 Intervention 2: maintenance 3 min isoflurane 3% + isoflurane 1.25% + post-delivery isoflurane 0.5%, N = 63
Outcomes	Primary outcomes: effects withdrawing N₂O for increased inspired concentration: enflurane or isoflurane; depth anaesthesia: isolated forearm technique: wakefulness (as first defined by Tunstall) Awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Tunstall 1989 (Continued)

Quote: "No mother experienced awareness (that is, postoperative recall) and no mother experienced dreams. Wakefulness was observed in 24 patients who received enflurane and in 23 who received isoflurane at the 2-min interval..."

Comment: 48% (24/50) enflurane group and 37% (23/63) isoflurane group

Notes

Non-randomized portion of anaesthetic: N₂O /parts of potent inhalational technique (volatile agent)/hypnotic/supplemental narcotics/muscle relaxants induction yes/maintenance yes/IFT: response to command (wakefulness)/N₂O yes - no pre-delivery and yes post-delivery: study - enflurane vs isoflurane

Anaesthesia induction: thiopentone 250 mg + suxamethonium 100 mg + oxygen 100%; anaesthesia maintenance: atracurium 25 mg to 30 mg + post-delivery oxytocin 10 units + 66% N₂O + metoclopramide 10 mg + papaveretum 20 mg

Comment: conclusion of RCT: the new volatile agents studied, 1.5% enflurane/1.25% isoflurane in oxygen only, were acceptable as **sole maintenance agent**

Comment: see Dryad topic **movement scores figures 1-4**; if the anaesthetist's hand was gripped without being released on the command to open and shut the hand, it was classified as reflex grip

Time of outcome determination: before discharge

Method of outcome determination: interview

Comment: see Dryad topic movement scores figures 1-4

Email: m.e.tunstall@abdn.ac.uk

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "...allocated randomly to receive either enflurane or isoflurane"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Vakkuri 2005

Methods	<p>Study design: randomized parallel groups</p> <p>Study dates: Quote: "8/7/2002 - 27/2/2003" (email bias survey, see notes)</p>
Participants	<p>Country: Finland, Sweden, Norway</p> <p>Sex: both</p> <p>ASA: I-III</p> <p>Age: 18 to 80</p> <p>Procedure: elective surgery procedures 45 min < time < 150 min</p> <p>Study size: 335</p>
Interventions	<p>Randomized portion of anaesthetic: ADM (Entropy 45 to 65) vs SCP (target ADM < 65)</p> <p>Intervention 1: entropy values shown group, N = 160</p> <p>Intervention 2: entropy values shown not control group, N = 160</p>
Outcomes	<p>Primary outcomes: Quote: "hypothesis that intraoperative monitoring of entropy would decrease propofol consumption during propofol–nitrous oxide–alfentanil anaesthesia"</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Quote: "None of the patients reported any anaesthesia- or surgery- related memories in the two post-operative interviews"</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of IV MCI/N₂O/muscle relaxants induction yes/ maintenance PRN: alfentanil/propofol/N₂O/N₂O/endotracheal tube or laryngeal mask/muscle relaxants when "appropriate"</p> <p>Induction: alfentanil \geq 30 μg/kg + propofol 1.0 mg to 2.5 mg/kg; maintenance: mixture of oxygen (35% to 50%) and N₂O (50% to 65%), infusions: alfentanil max. dose 30 μg/kg/h + propofol max dose 9 mg/kg/h</p> <p>Inadequate anaesthesia: alfentanil and propofol boluses and/or muscle relaxant choice of anaesthesiologist when considered "appropriate" when entropy indices increased, suggesting impending awakening; note the use of muscle relaxants in the protocol to treat a clinical judgement of impending awakening rather than vasoactive drugs and more anaesthesia; this is a commonly used protocol</p> <p>Comment: see discussion facial frontal muscle activation less sensitive to the effects of neuromuscular blocking drugs than are the hand muscles. Part of the depth of anaesthesia monitoring was defining the degree of paralysis</p> <p>Time of outcome determination: in PACU and 24 h postoperative</p> <p>Method of determination: interview</p> <p>Author survey email sent (Yli-Hankal) on 27 January 2011 to arvi.yli-hankala@uta.fi, no response received. Survey response: 15 February 2011, Anne Vakkuri anne.vakkuri@hus.fi. Author sent characteristics and ROB table grades for comment 8 December 2013: anne.vakkuri@hus.fi; email topic incomplete data 14 June 2015</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Vakkuri 2005 (Continued)

Random sequence generation (selection bias)	Low risk	Quote: "...randomly assigned, according to computer-generated random numbers, into the control group or the entropy group" (email bias survey, see notes)
Allocation concealment (selection bias)	Unclear risk	<p>Quote: "Each study site was provided with a sufficient number of closed randomisation envelopes. With sequential coding, the subjects were treated in blocks of 10 (5 patients in each group). The envelopes were opened in the operating room immediately before the induction of anaesthesia"</p> <p>Quote (email bias survey, see notes): "Each patient got their own randomization number sealed in an envelope. The study nurse opened the envelope after the patient had been recruited to the study, prior to anaesthesia induction"</p> <p>Comment: authors describe much detail but omit the word "opaque"; we will upgrade if informed by authors</p>
Blinding of participants and personnel (performance bias) All outcomes	High risk	<p>Quote: "Patient, Awareness outcome assessor (interviewer)" (email bias survey)</p> <p>Comment: ROB survey response indicates anaesthesiologist not blinded to anaesthetics administered. However, because they were blinded to ADM in control group, that meets our criteria for low risk</p> <p>Quote: "The enrolled patients were randomized to receive propofol–nitrous oxide–alfentanil anaesthesia either with Entropy values shown (entropy group) or with entropy values not shown (control group)"</p> <p>Comment: SCP exposed to both groups</p>
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Patient, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Incomplete outcome data (attrition bias) All outcomes	Low risk	<p>Quote: "Seventeen patients were excluded: 14 because of lack of registered data, 1 because of violation of the inclusion criteria, 1 because of accidental use of a potent inhalational agent, and 1 due to respiratory arrest during the emergence phase. The data from 368 patients (48 historical controls, 160 controls, and 160 entropy patients) were included in final analyses" (email bias survey, see notes)</p> <p>Comment: no high-risk awareness exclusions; no details about the distribution between groups; however, extreme imbalance would not be significant</p> <p>17/160 vs 0/160, Peto OR 8.21 (3.10 to 21.78)</p>
Selective reporting (reporting bias)	Low risk	<p>Quote: "The study protocol is available and all of the study's pre-specified (primary and secondary) outcomes that are of interest in the review have been reported in the pre-specified way. Regarding the historical control patients, only significant differences between historical controls and control group patients were reported. This is stated in the paper." (email bias survey, see notes)</p> <p>Comment: awareness outcome part of inclusion criteria</p>
Other bias	Low risk	Quote: "The study is free of other bias." (email bias survey, see notes)

van der Maaten 1996

Methods Study design: randomized parallel groups

van der Maaten 1996 (Continued)

Study dates: Quote: "in 1994" (email bias survey, see notes)

Participants	Country: the Netherlands Sex: both Age: average 61.4, 59.3 ASA: I-II Procedure: coronary artery bypass grafting: exclusion criteria were left main coronary artery disease. Impaired left ventricular (LV) function (ejection fraction < 40%). LV end-diastolic pressure > 18 mmHg Study size: 20, 18
Interventions	<p>Randomized portion of anaesthetic: parts of TIVA: hypnotic (midazolam) infusions TCI and MCI: maintenance</p> Intervention 1: maintenance midazolam (TCI) target plasma concentration of 150 ng/mL , 2 µg/kg/min during 85 minutes + midazolam MCI 1.25 µg/kg/min thereafter, N = 10 Intervention 2: maintenance target plasma concentration 300 ng/mL midazolam, 10 µg/kg/min , during 15 minutes, then 4 µg/kg/min for the next 70 minutes + midazolam 2.5 µg/kg/min thereafter, N = 10
Outcomes	Primary outcomes: haemodynamic data, sufentanil and midazolam plasma concentrations Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "Intraoperative awareness was not reported"
Notes	<p>Non-randomized portion of anaesthetic: N₂O no/parts of TIVA hypnotic (midazolam) infusions/narcotics infusion (sufentanil): muscle relaxants induction yes/maintenance unclear</p> Preoperative cardiac medication, consisting of β-blockers, calcium entry blockers, nitrates, and antihypertensive agents, was continued and administered on the morning of surgery Pre-medicated: with morphine sulphate 0.15 mg/kg Anaesthesia induction : midazolam 0.1 mg/kg + sufentanil, 2.5 µg/kg + pancuronium 0.1 mg/kg; anaesthesia maintenance : sufentanil 1 µg to 2 µg/kg/h + sufentanil 50 µg as needed + nitroglycerin 0.25 µg to 3 µg/kg/min as needed + sufentanil 1 µg/kg + phenylephrine 50 µg to 100 µg as needed Time of outcome determination: 3 days postoperative Method of outcome determination: interview Survey response: 18 January 2011, 31 August 2011, J van der Maaten j.m.a.a.van.der.maaten@an-est.umcg.nl

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Patients were randomly assigned to receive midazolam at a target plasma concentration of 150 ng/mL (group 1) or 300 ng/mL (group 2)" Quote: "Computer-generated block randomization (10x group1 and 10x group2)." (email bias survey, see notes)
Allocation concealment (selection bias)	Low risk	Quote survey: "The concentration of the study drug (midazolam, 1mg/ml) in the syringe was the same for both study groups"

van der Maaten 1996 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Quote survey: "Considering the difference in infusion rate for both groups, the anaesthesiologist could have been aware of the assignment to one group or the other " (email bias survey, see notes) Comment: rule; infusions can not be blinded due to differences in infusion rates
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Three days after surgery, the patients were interviewed to detect recollection of auditory or other forms of awareness of intraoperative events..." Quote: "Patient, Awareness outcome assessor (interviewer)" (email bias survey, see notes)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "There were no missing values and no attrition, exclusions or re-inclusions. Data set was complete, including the plasma drug concentration samples" Quote (email bias survey, see notes): "There were 10 patients in each study group. All data were complete, except for the samples for midazolam and sufentanil plasma concentrations there were two missing values in each sample set (2 missing values on 110 samples for the midazolam and sufentanil respectively)" Comment: there was no significant difference between groups: (2/10 vs 0/10), Peto OR 8.26 (0.48 to 142.43)
Selective reporting (reporting bias)	Low risk	Quote: "The study protocol is available and outcomes of interest have been reported" (email bias survey, see notes) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote: "Free of other bias" (email bias survey, see notes)

van Leeuwen 1990

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: the Netherlands Sex: both Age: aged 15 to 72 years ASA: I and II Procedure: orthopaedic or ophthalmic operations Study size: 30
Interventions	Randomized portion of anaesthetic: parts of TIVA: hypnotic (propofol) infusions MCI: maintenance Intervention 1: maintenance propofol 2 mg/kg/h, N = 10 Intervention 2: maintenance propofol 3 mg/kg/h, N = 10 Intervention 3: maintenance propofol 4 mg/kg/h, N = 10

van Leeuwen 1990 (Continued)

Outcomes Primary outcomes: cardiovascular parameters between groups; duration of anaesthesia, mean total doses (induction and maintenance) of both propofol and alfentanil

Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1

Quote: "Awareness did not occur in any patient"

Notes **Non-randomized portion of anaesthetic: N₂O no parts of TIVA hypnotic (propofol) infusions MCI/narcotics infusion (alfentanil): maintenance/muscle relaxants induction yes/maintenance no**

All patients received premedication with diazepam 10 mg orally 1 to 1.5 h before surgery. Anaesthesia induction: propofol 2 mg/kg + alfentanil 10 µg/kg/min + vecuronium bromide 0.1 mg/kg. A single dose of vecuronium 0.1 mg/kg was used as the muscle relaxant; anaesthesia maintenance: alfentanil infusion 1 µg/kg/min + propofol 20 mg/kg and alfentanil 1 mg was administered as needed. Oxygen in air (FIO₂ 0.35). Neuromuscular block was reversed with atropine + neostigmine + naloxone 0.2 mg to 0.4 mg, if respiratory depression was present

Time of outcome determination: day and month after operation

Method of outcome determination: interview

Email survey sent to author, no response received

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The patients were randomly allocated into three groups of ten patients each, Groups A, B, and C. All..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided Comment: different infusion rates difficult to blind to anaesthesiologist (ROB survey van der Maaten 1996)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "The patients were questioned by the anaesthetist about intraoperative awareness. This was done in the recovery room, the day after and one month after the operation"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Wang 2013

Methods Study design: randomized parallel groups
Study dates: January 2010 to October 2011

Wang 2013 (Continued)

Participants	Country: China Sex: female Age: aged 15 to 75 years, 52.9 (9.8), 53.3 (9.6) ASA: I-III Procedure: breast cancer surgery including modified radical mastectomy, total mastectomy, lumpectomy, breast-conserving surgery and breast reconstruction Study size: 920 enrolled, 908 completed study
Interventions	Randomized portion of anaesthetic: parts of TIVA: premedication (phencyclidine) vs placebo Intervention 2: premedication (N = 456) phencyclidine (PHC) 0.01 mg/kg Intervention 2: premedication (N = 452) placebo (saline) BIS-guided total intravenous anaesthesia
Outcomes	Primary outcomes: awareness/wakefulness as defined using an awareness classification system (see Table 1) Quote: "The primary outcome was to evaluate the effect of PHC on intra-operative awareness." Quote: "A committee of three experts , blinded to the study conditions, independently scrutinised all reported recollections...PHC group, none of the patients had recall of intra-operative events (0%), saline group, five of 452 patients reported intra-operative awareness (1.1%)"
Notes	Non-randomized portion of anaesthetic: N₂O no/parts of TIVA hypnotic (propofol TCI and bolus midazolam)/narcotics (bolus sufentanil): maintenance/muscle relaxants induction yes/maintenance PRN/BIS 40 to 60 General anaesthesia was induced by propofol (TCI) plasma target-controlled infusion (a target plasma concentration of 3.5 to 4.5 lg/mL 1)/ bolus midazolam (0.03 mg/kg 1)/ bolus sufentanil (0.3 lg/kg 1)/intubation 0.2 mg/kg 1 cisatracurium BIS < 45. TCI adjusted maintain BIS 40 to 60 /neuromuscular blockade: cisatracurium PRN Interventions: cardiovascular instability: if blood pressure deviated > 30% baseline value for > 5 min Correspondence to: ZM Tan Email: zmtan166@163.com ROB survey: we emailed zmtan166@163.com on 22 March 2015

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Random assignment of patients was established by computer-generated codes. Allocation concealment was established by placing the randomisation sequence in consecutively numbered, opaque envelopes"
Allocation concealment (selection bias)	Unclear risk	Comment: unclear if sealed; await author's response to ROB survey
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "The PHC group received 0.01 mg.kg ⁻¹ PHC intravenously, whereas the saline group patients received saline intravenously as placebo. Penehyclidine hydrochloride or saline solutions were prepared in a syringe by the first anaesthesiologist, who was also responsible for subject grouping. The PHC was diluted (1 mg in 1 ml) ...This second anaesthesiologist was also responsible for the anaesthetic manage"
Blinding of outcome assessment (detection bias)	Low risk	Quote: "The third anaesthesiologist served as the postoperative interviewer. Awareness was defined as recall of intra-operative events"

Wang 2013 (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "Patients' data were not included if: two consecutive recorded BIS values were outside the target range 40–60 ; the time of surgery was longer than 10 h or shorter than 30 min; and patients required ephedrine or atropine because of circulatory instability. In all, data from 12 patients were not analysed (four in the PHC group, eight in the saline group), which left 908 patients' data (456 in the PHC group and 452 in the saline group) available for analysis (Fig. 1)" Comment: Figure 1: PHC group: excluded 4: BIS < 40 N = 1, > 60 N = 1, ephedrine N = 2, atropine N = 0; saline group: excluded 8: BIS < 40 N = 3, > 60 N = 2, ephedrine N = 2, atropine N = 1; high-risk awareness exclusions: BIS > 60 N = 1 for PHC and 2 for saline groups; 2% (1/460) vs 4% (2/460); since the saline group has increased awareness events and high-risk awareness exclusions vs PHC group, there is no downgrade In addition, there was no significant difference between groups, Peto OR 0.75 (0.17 to 3.32)
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: no information

White 2003

Methods	Study design: randomized parallel groups Study dates:
Participants	Country: USA Sex: female Age: 37, 41, 40 Procedure: gynaecologic laparoscopic procedures Study size: 45
Interventions	Randomized portion of anaesthetic: parts of volatile agent (desflurane): cardiac medication vs placebo: esmolol/nicardipine/saline Intervention 1: induction: saline 5 mL, maintenance: infusion saline, N = 15 Intervention 2: induction: esmolol 50 mg + saline 1 mL, maintenance infusion esmolol, N = 15 Intervention 3: induction: esmolol 50 mg + nicardipine 1 mg, maintenance: infusion esmolol, N = 15 infusion saline and esmolol started at a rate of 0.005 mL/kg 1 to 2 min before the skin incision
Outcomes	Primary outcomes: times to awakening and discharge home Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "Although the study was not adequately powered to assess awareness under anaesthesia, no patient in any of the groups reported recall of intraoperative events"

White 2003 (Continued)

Notes

Non-randomized portion of anaesthetic: parts of volatile agent (desflurane)/N₂O/hypnotic (propofol) bolus narcotics (fentanyl) + muscle relaxant(s) (vecuronium) induction yes/PRN maintenance ADM (BIS) blinded

Anaesthesia induction: fentanyl 1.5 µg/kg + propofol 2 mg/kg + vecuronium 0.12 mg/kg + group-specific study drug; anaesthesia maintenance: desflurane 2% (inspired) and N₂O 67% in oxygen + vecuronium 1 mg to 2 mg PRN; BIS **blinded**

Pneumoperitoneum maintained at 15 to 20 mmHg/Trendelenburg position; MAP maintained 15% of the pre-induction baseline value by increasing/decreasing the infusion rate study medication 50% to 100%

Email survey sent, no response received

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "...were randomly assigned to 1 of 3 treatment groups..."
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "Finally, none of the patients in the study reported recall of intraoperative events when questioned at the time of discharge"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

White 2004

Methods	Study design: randomized parallel groups Study dates: not given
Participants	Country: USA Sex: female Age: 48, 54, 50 ASA: I-III Procedure: gynaecologic laparoscopic surgery

White 2004 (Continued)

Study size: 60

Interventions	<p>Randomized portion of anaesthetic: ADM (AAI 15 to 25, N = 20 BIS 50 to 60) vs SCP</p> <p>Intervention 1: maintenance: SCP, blinded to BIS or AAI, N = 20</p> <p>Intervention 2: maintenance BIS 50 to 60, N = 20</p> <p>Intervention 3: maintenance AAI 15 to 25, N = 20</p> <p>Value in the range of 50 to 60 or 15 to 25, respectively</p>
Outcomes	<p>Primary outcomes: relationship between BIS and PSI values during the induction and emergence periods, BIS and PSI values and the probability of unconsciousness</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Quote: "None of the patients reported recall of intraoperative events when questioned at the time of discharge from the hospital or during the follow-up telephone interview at 24 h after surgery"</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent (desflurane)/N₂O/hypnotic (propofol) bolus narcotics (fentanyl) + muscle relaxant(s) (vecuronium) induction yes/maintenance yes</p> <p>Premedication: midazolam 2 mg; anaesthesia induction: propofol 1.5 mg to 2.5 mg/kg + fentanyl 1 µg to 1.5 µg/kg + succinylcholine 1 mg to 1.5 mg/kg; anaesthesia maintenance: desflurane 3% with 60% N₂O in oxygen 1.5 L/min/1 L/min + cisatracurium 10 mg to 20 mg + esmolol 10 mg as needed + neostigmine 0.05 mg/kg + glycopyrrolate 0.01 mg</p> <p>Clinical signs of excessive anaesthetic effect (e.g. a decrease in MAP 20% pre-incision value), desflurane decreased by 2%; residual neuromuscular blockade reversed</p> <p>Comment: correlation coefficients for eye opening with respect to AAI/BIS and desflurane ET concentration poor</p> <p>Time of outcome determination: before discharge and 24 h postoperative</p> <p>Method of outcome determination: interview</p> <p>Email survey sent, no response received. ROB survey: we emailed Dr White paul.white@utsouthwestern.edu on 30 April 2015; email rejected: "Delivery to the following recipient failed permanently"</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "...were randomly assigned to one of three groups"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: SCP exposed to both groups
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "At the time of discharge from the hospital and during the follow-up telephone interview at 24 h after surgery, patients were asked whether they recalled any events during the intraoperative period"

White 2004 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Wong 2002

Methods	Study design: randomized parallel groups Study date: "1998-2000" (email bias survey, see notes)
Participants	Country: Canada Sex: both Age: 76, 71 ASA: I-III Procedure: elective orthopaedic knee or hip replacement surgery Study size: 68
Interventions	Randomized portion of anaesthetic: ADM (BIS 50 to 60) vs SCP Intervention 1: BIS guided titration of anaesthesia (BIS 50 to 60), N = 34 Intervention 2: standard clinical practice of anaesthesia (SP), N = 34
Outcomes	Primary outcomes: BIS values, haemodynamics, anaesthetic consumption, recovery time Awareness/wakefulness as defined using an awareness classification system (see Table 1) Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 0 Quote: "None of the patients reported awareness" No patient had intraoperative dreams
Notes	Non-randomized portion of anaesthetic: parts of volatile agent (isoflurane)/N₂O/hypnotic (propofol/midazolam) bolus narcotics (fentanyl) + muscle relaxant(s) (rocuronium) induction yes/maintenance yes Anaesthesia induction: propofol 1 mg to 2 mg/kg + fentanyl 2 mg to 3 mg/kg + midazolam 1 mg + rocuronium 0.6 mg/kg Anaesthesia maintenance: isoflurane and 60% to 70% N ₂ O in oxygen at 3 L/min for 5 minutes, then decreased to 1.5 L/min Reversal of neuromuscular blockade 5 minutes prior to the discontinuation of inhalational agents Comment see Dryad topic for light anaesthesia protocol Comment: haemodynamic stability, rapid recovery and quick discharge time are the 3 most common parameters that many of the included RCTs use to guide their dosing of anaesthesia to patients

Wong 2002 (Continued)

Time of outcome determination: 72 h + 14 days postoperative

Method of outcome determination: interview

 Survey emailed to jean_wong@yahoo.com on 6 December 2013

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "...block randomization with concealed varying block sizes was performed with computer generated random numbers" Survey: "computer generated random numbers with varying block sizes"
Allocation concealment (selection bias)	Low risk	Quote: "The assignments were kept in sealed envelopes by a research assistant not involved with the study" Comment: await author response as to whether envelope was sealed
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "In the standard practice (SP) group... anaesthesiologist was blinded to the BIS value... " Comment: SCP exposed to both groups
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "At the 72-h interview, the patients were asked a series of questions relating to whether they had any recall of intraoperative events (Appendix)" Quote: "Patient, Awareness outcome assessor (interviewer)"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "all patients were included" 68 patients were enrolled in this study; however, 8 patients (3 from the SP group, and 5 from the BIS group) were excluded from the analysis for protocol violations. 2 patients had bipolar disorder; 2 patients received propofol near the end of surgery; 2 patients received excessive fentanyl near the end of surgery; and 1 patient desaturated necessitating discontinuation of N ₂ O intraoperatively There was no difference in demographic data between the 2 groups (Table I)
Selective reporting (reporting bias)	Low risk	Quote: "all pre-specified outcomes of interest were reported" Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote survey: "no other risk of bias"

Wu 2001

Methods	Study design: randomized parallel groups Study date: not given
Participants	Country: Taiwan Sex: female Age: 36 ASA: I and II

Wu 2001 (Continued)

Procedure: elective gynaecologic surgeries

Study size: 40

Interventions	<p>Randomized portion of anaesthetic: induction hypnotic agents ketamine (dissociative agent) vs thiamylal (barbiturate)</p> <p>Intervention 1: ketamine (1.5 mg/kg IV) induction, N = 20</p> <p>Intervention 2: thiamylal (5 mg/kg IV) induction, N = 20</p>
Outcomes	<p>Primary outcomes: BIS values, haemodynamics</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Quote: "No patient reported recall, delirium, hallucination or awareness under anaesthesia as questioned in the Post Anaesthesia Care Unit (PACU) when they were fully awake"</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of volatile agent (isoflurane)/N₂O/muscle relaxants: yes both induction (succinylcholine)/maintenance (atracurium PRN)/BIS recorded</p> <p>Comment: anaesthesia induction: interventions 1 and 2 titrated: clinical signs/BIS recorded/succinylcholine (1 mg/kg)</p> <p>Comment: anaesthesia maintenance: isoflurane 1.5 MAC in N₂O 3 L/min and O₂ 2 L/min + atracurium PRN</p> <p>BIS monitor used only to collect data not to manage patient</p> <p>Email survey sent to author, no response received</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The patients were then divided randomly into two groups: Group K"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: "No patient reported recall, delirium, hallucinations or awareness under anaesthesia..."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Yildiz 2002

Methods	<p>Study design: randomized parallel groups</p> <p>Study date: not given</p>
Participants	<p>Country: Turkey</p> <p>Sex: both</p> <p>Age: 42</p> <p>ASA: I-II</p> <p>Procedure: laparoscopic cholecystectomy</p> <p>Study size: 50</p>
Interventions	<p>Randomized portion of anaesthetic: parts of TIVA anaesthesia/narcotic (alfentanil and remifentanil)</p> <p>Intervention 1: induction: alfentanil 10 mg/kg, N = 25</p> <p>Intervention 2: induction: remifentanil 1.5 mg/kg, N = 25</p> <p>Cards with pictures presented to patients</p>
Outcomes	<p>Primary outcomes: BIS values, haemodynamics, "delayed memory recall test"</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 2</p> <p>Quote: "After gained consciousness none of the patients in both groups did not have any awareness according to the meetings with the patients"</p> <p>Comment: class 2: negative explicit recall, positive implicit memory for word/image/music stimuli but no emotional sequelae; both groups had "delayed memory recall test [implicit memory?]" of cards with images and there were significantly fewer "error points" with the remifentanil vs alfentanil group</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of TIVA anaesthesia: air/narcotic (randomized: alfentanil and remifentanil)/hypnotic (propofol)/muscle relaxants induction yes/maintenance unclear (atracurium)/BIS recorded</p> <p>Anaesthesia induction: (randomized: alfentanil (10 µg/kg) or remifentanil (1 µg/kg))/propofol (2 mg/kg)/atracurium (0.6 mg/kg); anaesthesia maintenance: (randomized: alfentanil (1 µg/kg/min) or remifentanil (0.25 µg/kg/min))/propofol (9 mg/kg/h)/ventilation O₂:air 1:1, residual neuromuscular block antagonized/BIS monitor used</p> <p>Time of outcome determination: 24 h postoperative</p> <p>Method of outcome determination: interview</p> <p>Learning test: delayed memory test - pictures</p> <p>Murat translator</p> <p>No email address to send survey</p>
Risk of bias	
Bias	Authors' judgement Support for judgement

Yildiz 2002 (Continued)

Random sequence generation (selection bias)	Unclear risk	Quote: "...randomly separated into two groups"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: anaesthesiologist: no information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Yoshitani 2003

Methods	Study design: randomized parallel groups Study dates: "From October, 1998 to July, 1999." (email bias survey, see notes)
Participants	Country: Japan Sex: both Age: 37 to 85 ASA: not stated Procedure: cardiac surgery Study size: 45
Interventions	Randomized portion of anaesthetic: parts of IV: hypnotic (propofol) infusions TCI Intervention 1: maintenance: propofol 4 mg/kg/h, N = 15 Intervention 2: maintenance: propofol 5 mg/kg/h, N = 15 Intervention 3: maintenance: propofol 6 mg/kg/h, N = 15
Outcomes	Primary outcomes: BIS values, anaesthetic consumption, haemodynamic parameters Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Quote: "No patient reported awareness of intraoperative events"

Yoshitani 2003 (Continued)

Notes

Non-randomized portion of anaesthetic: parts of intravenous (IV) anaesthesia: N₂O/narcotic (fentanyl)/hypnotic (propofol)/muscle relaxants induction yes/maintenance yes (vecuronium): EEG recorded

Anaesthesia induction: fentanyl 10 mg/kg + propofol 3 µg/mL + vecuronium 0.2 mg/kg

Anaesthesia maintenance: fentanyl 5 µg/kg/h (to a total of 30 µg/kg) + oxygen 50% and N₂O 50%

The EEG was monitored continuously from induction of anaesthesia to emergence

Author responded to the email bias survey on 3 February 2011. Responses are recorded in the 'Risk of bias' table.

Survey response: 3 February 2011, Kenji Yoshitani ykenji@kfz.biglobe.ne.jp

Emailed author 29 December 2013 to clarify allocation concealment

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Patients were randomly assigned to receive propofol at 4 (Group A), 5 (Group B), or 6 mg/kg/h IV (Group C) using the sealed envelope technique in block randomization with 15 in each group"
Allocation concealment (selection bias)	Unclear risk	Quote: "We used sealed envelope method." (email bias survey, see notes) Comment: emailed author 29 December 2013 to clarify; await author response regarding the use of opaque envelopes
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "This study was an open label study. To change propofol dosage according to the assignment was difficult to blind. However, patients did not know the group assignment"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote survey: "Patient, Awareness outcome assessor (interviewer)"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "All outcome was completed in the study. There was no exclusion because of single center small study" (email bias survey, see notes)
Selective reporting (reporting bias)	Low risk	Quote: "The study protocol is available and all of the study's pre-specified outcomes that are of interest in the review have been reported in the pre-specified way" (email bias survey, see notes) Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	Quote: "We did not have any risk of bias. We did not have a design related to the specific study design, stop early and so on"

Zhang 2011

Methods	Study design: randomized parallel groups Study dates: November 2008 to November 2010
---------	---

Zhang 2011 (Continued)

Participants	Country: China Sex: both Age: ≥ 18 years ASA: I to III with significantly sicker patients in BIS guided group ($P < 0.01$) Procedure: Type of operation: 1 = neurosurgery, 2 = craniofacial and cervical surgery, 3 = heart surgery, 4 = gynaecologic and obstetrics surgery, 5 = chest and abdomen surgery, 6 = urinary surgery, 7 = spine and limb surgery, 8 = others Study size: enrolled 5309, completed study 5228 patients
Interventions	Randomized portion of anaesthetic: ADM (BIS 40 to 60) vs SCP: TIVA TIVA: BIS guided (40 to 60), N = 2309 BIS blinded, N = 2919
Outcomes	Primary outcomes: awareness/wakefulness as defined using our 'Classification of intra-operative cognitive states' Table 1: class 4 BIS-guided group vs BIS blinded (control group) Confirmed awareness: 4/2919 (0.14%) vs 15/2309 (0.65%) ($P = 0.002$, OR 0.21, 95% CI 0.07 to 0.63) Possible awareness: 4/2919 (0.14%) vs 6/2309 (0.26%) ($P = 0.485$) Dreaming: 3.1% vs 3.1% ($P = 0.986$)
Notes	Non-randomized portion of anaesthetic: parts of TIVA/N₂O unclear hypnotic (midazolam + propofol infusion)/narcotics: other anaesthetics (analgesics and muscle relaxants): discretion of the anaesthetist Comment: TIVA not defined No premedication Induction: midazolam + propofol infusion Maintenance: propofol infusion and other anaesthetics (analgesics and muscle relaxants): discretion of the anaesthetist Comment: this is one of the most detailed stratifications of patient awareness reports in this review Comment: see Dryad topic: explanation of awareness events Conclusion: authors agree with Ghoneim 2009 and Schwender 1995: "light anaesthesia was the main reason for awareness" Comment: authors describe one of the most complete lists of causes of awareness Email communications: yueyun@hotmail.com , zhch1127@sina.com : 31 January 2013 through 14 November 2014; topics: ROB survey, missing data, adjudicated awareness reports including non-published data; 13 June 2015 N ₂ O

Risk of bias

Zhang 2011 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	ROB survey response by Dr Zhang, Quote: "Randomization was carried out in each individual center through computer-generated random numbers to develop randomized program..." Comment: there is an unexplained large discrepancy in the sample sizes for the 2 interventions: 2309 vs 2919
Allocation concealment (selection bias)	Unclear risk	ROB survey response by Dr Zhang: "Awareness investigators and patients were blinded to the group allocation. We have independent investigators who did not know the patient allocation and did not participate in anesthesia procedure. The anesthesiologists were not blinded, for they have to maintain anesthesia depth by BIS or not according to the allocation"
Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: SCPs exposed to both groups and infusion rates difficult to blind
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Interviewers and patients were blinded to the group allocation"
Incomplete outcome data (attrition bias) All outcomes	Low risk	ROB survey response: "Outcome data was collected from 5309 patients. Fifty-four cases were withdrawn because the information of group allocation was unavailable (without awareness cases), another 21 patients were excluded due to younger than 18 years (11 cases in Group A, and 10 cases in Group B), and a further 6 patients were excluded because of failure to be interviewed at any of the two times (2 cases in Group A, 2 cases in Group B; 1 patient died postoperatively, operation was cancelled in 1 case after anaesthesia induction). Thus, a total of 5228 cases were enrolled in the final statistical analysis with 2919 cases in Group A and 2309 cases in Group B"
Selective reporting (reporting bias)	Low risk	Quote: "No" Comment: awareness outcome part of inclusion criteria
Other bias	Low risk	ROB survey response: "There was one bias related to our study design that the anesthesiologists were not blinded to the group allocation"

Zhou 2008

Methods	Study design: randomized parallel groups Study dates: NA
Participants	Country: China Sex: female Age: mean age 39 ± 7 ASA: I-II Procedure: gynaecologic laparoscopy Study size: 45

Zhou 2008 (Continued)

Interventions	<p>Randomized portion of anaesthetic: parts of TIVA (TCI propofol/propofol bolus) vs parts of volatile agent (sevoflurane)</p> <p>Intervention 1: induction and maintenance: sevoflurane 6%: induction; 4%: maintenance MAC, N = 15</p> <p>Intervention 2: induction: TCI propofol 3 mg/mL, N = 15</p> <p>Intervention 3: induction: propofol bolus 2 mg/kg, N = 15</p>
Outcomes	<p>Primary outcomes: BIS values and haemodynamics peri-intubation</p> <p>Awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1</p> <p>Quote: "no adverse reaction concerning memory was observed in three groups"</p>
Notes	<p>Non-randomized portion of anaesthetic: nitrous N/parts of TIVA vs volatile agent (sevoflurane) + narcotics (sufentanil) + muscle relaxant(s) (rocuronium) induction yes/maintenance unclear/ADM BIS 60</p> <p>Induction: sufentanil 0.5 µg/kg + sevoflurane 6 MAC + rocuronium bromide 0.6 mg/kg/BIS</p> <p>Comment: intubation at BIS 60 is example of light anaesthesia</p> <p>Zancong Shen translator</p> <p>ROB survey sent yueyun@hotmail.com</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote survey: "Patients were randomized by their assigned random numbers, however after randomization each group was given different treatment drugs...And or anesthesia depth monitors like BIS (BIS is the key depth monitoring endpoint (60) when intubation was applied for all patients),Narcotrend, AAI, etc. All 45 patients had in GYN laparoscopy"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Comment: difficult to blind infusion-based anaesthetic and volatile agent
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Comment: assessor: no information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	<p>Comment: no missing data</p> <p>Quote: "They followed up with all patients post surgery for any adverse reaction related to memory, but did not find any)"</p>
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Zohar 2006

Methods	Study design: randomized parallel groups Study dates: NA
Participants	Country: Israel and USA Sex: female Age: geriatric > 65 ASA: Procedure: short elective transurethral surgical procedures Study size: 50
Interventions	Randomized portion of anaesthetic: ADM (BIS 50 to 60) vs SCP 1) Standard practice (control) group, N = 25 2) Experimental (BIS) group (50 to 60), N = 25
Outcomes	Primary outcomes: anaesthetic and analgesic requirements, times to eye opening, and other recovery parameters Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class 1 Comment: no awareness events reported
Notes	Randomized portion of anaesthetic: parts of volatile agent/hypnotic/(propofol)/(sevoflurane)/N₂O/narcotic (fentanyl)/spontaneous respiration, laryngeal mask airway: hence, no muscle relaxant(s) No preanaesthetic medication Induction: fentanyl 1.0 µg to 1.5 µg/kg + propofol 1.5 mg to 2.0 mg/kg Maintenance: sevoflurane 1.5% with 60% N ₂ O in oxygen, fentanyl 25 µg IV PRN tachypnoea; LMA; BIS 50 to 60 Comment: supplemental "rescue" doses of fentanyl (25 µg IV) were significantly more in SCP vs BIS group Comment: see Dryad topic calculation of sevoflurane minimum alveolar concentration (MAC) during the maintenance period No email address available for ROB survey

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "written informed consent, 50 geriatric outpatients were enrolled in this prospective, randomized, assessor-blinded study involving two treatment groups"
Allocation concealment (selection bias)	Unclear risk	Comment: no information provided

Zohar 2006 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	High risk	Comment: SCPs exposed to both groups
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: assessor: "Early recovery endpoints were recorded at one minute intervals following discontinuation of the maintenance anaesthetics by a 'blinded' observer"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Comment: no missing outcome data
Selective reporting (reporting bias)	Low risk	Comment: awareness outcome part of inclusion criteria
Other bias	Unclear risk	Comment: insufficient information provided

Acronyms and abbreviations

AAI = A-line ARX index; ADM = anaesthesia depth monitor; AEP = auditory evoked potential; AER = auditory evoked response; AMLR auditory middle-latency response; AP = arterial pressure; APGAR = Appearance, Pulse, Grimace, Activity, and Respiration developed in 1952 by an anaesthesiologist named Virginia Apgar; AML = auditory middle latency response; ASA = American Society of Anesthesiology; ASSR = auditory steady-state response; BAG-RECALL = BIS or Anaesthetic Gas to Reduce Explicit Recall trial; BIS = Bispectral Index; BP = blood pressure; CABG = coronary artery graft surgery; cbc = complete blood count; CBW = corrected body weight; CI = confidence interval; CLADS = closed-loop anaesthesia delivery system; CPB = cardiopulmonary bypass; CV = cardiovascular; DBIS, DBP, DHR, DMAP = changes in BIS, heart rate and MAP; EEG = electroencephalogram; ENT = ear, nose and throat; ET = end-tidal; ETAG = end-tidal anaesthesia gas; ETT = endotracheal tube; FIO₂ = fraction of inspired oxygen; GA = general anaesthesia; GETA = general endotracheal anaesthesia; GETGA = general endotracheal general anaesthesia; h = hours; HBI = pulse plethysmography; mmHg = millimetres of mercury; h = hour; HR = heart rate; HTN = hypertension; Hz = hertz; ICU = intensive care unit; IFT = isolated forearm technique; IM = intramuscular; IPPV = intermittent positive pressure ventilation; IU = international unit; IV = intravenous; L = litres; LMA = laryngeal mask airway; LOC = loss of consciousness; LV = left ventricular; kg = kilogram; MAC = minimal alveolar concentration; µg = micrograms; MCI = manually controlled infusion; mg = milligram; min = minute; µg/kg/min = micrograms per kilogram per minute; MLAEP = middle latency auditory evoked potentials; MLAER = middle latency auditory evoked responses; MVD = microvascular decompression; N = sample size; NA = not available; NaCl: saline; NS not significant; N₂O = nitrous oxide; NMB = neuromuscular blocking agents; OPCAB = off pump coronary artery bypass; OR = odds ratio; OpR = operating room; Pa = waveform part of auditory evoked potentials; PACU = post-anaesthesia care unit; PHC = phencyclidine; PO = per ora (by mouth); POD = postoperative day; PPG = photo plethysmography; PRN = pro re nata (as the occasion arises) 'as needed'; PRST = P (systolic blood Pressure), R (heart Rate), S (Sweating) and T (Tears); PSI = pound per square inch; RCT = randomized controlled trial; RE = response entropy; ROB = risk of bias; RSII[AM1] = rapid sequence induction and intubation; S = S- (+)-ketamine is more potent than its racemic mixture; SBP = systolic blood pressure; SCP = standard clinical practice; SD = standard deviation; SE = state entropy; SEF = spectral edge frequency; SNAP = sensory nerve action potentials; SNOSE = sequentially numbered, opaque, sealed envelopes; SPI = surgical pleth index; SR = spontaneous respiration; sys = systolic; TBW = total body weight; TCI = target-controlled infusions; THR = total hip replacement; TIVA = total intravenous anaesthesia; TOF = train-of-four; TP termination of pregnancy; TPS Paventi 2000 no definition; UID = uterine incision to delivery interval; vs = versus

** acronym used twice

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Abboud 1990	Not general anaesthesia
Abdel-Meguid 2005	No awareness test
Abott 1980	No randomization of medications
Abouleish 1976	No randomization of medications

Study	Reason for exclusion
Absalom 2002	No randomization of medications
Acil 2004	No awareness test
Adams 1998	No randomization of medications
Adams 2003	Not general anaesthesia. Survey response: 24 January 2011, Prof. Dr med. Hans Anton Adams adams.ha@mh-hannover.de
Ahmad 2003	No awareness test
Ahonen 2007	No awareness test
Al-Ruzzeh 2006	No awareness test
Alexander 1999a	No awareness test
Alexander 1999b	No awareness test
Alvarez 2000	No randomization of medications
Andelman 2004	No randomization of medications
Anderson 2003	No randomization of medications
Anderson 2004	No randomization of medications
Andrade 2001	No randomization of medications
Aono 1999	No randomization of medications
Apfelbaum 1996	Volunteers not surgery
Aqil 2009	No awareness test
Arndt 1995a	Not randomized
Aubrun 2008	No awareness test
Ausems 1983	No randomization of medications
Ausems 1986	No awareness test
Bailey 1985	Wake up test
Balci 2006	No awareness test
Bannister 2001	No awareness test
Baraka 1990	No randomization of medications
Barclay 1980	No randomization of medications
Barr 1973	No randomization of medications

Study	Reason for exclusion
Barr 1999	Volunteers not surgery
Barr 2001	Volunteers not surgery
Barvais 2003	No randomization of medications
Basar 2003	No awareness test
Becker-Blease 2006	Volunteers not surgery
Bejjani 2009	No randomization of medications
Bennett 1985	No randomization of medications
Bilotta 2007	No awareness test-POCD
Bischoff 1997	Expert commentary about awareness: EEG review
Block1991b	No awareness test-therapeutic suggestion results only
Bogetz 1984	No randomization of medications
Bogod 1990	No randomization of medications
Bonhomme 2006	No awareness test
Bonke 1986	No randomization of medications
Bonke 1992	No awareness test
Bould 2007	No randomization of medications
Brice 1970	No randomization of medications
Brosius 2002	No awareness test
Bruhn 2000	Expert commentary about awareness: case report
Buffett-Jerrott 2003	No awareness test - complicated memory test but no evidence awareness of intraoperative events assessed
Bulach 2005	No awareness test
Burn 1963	No randomization of medications
Burrow 2001	No awareness test
Byers 1997	No randomization of medications
Capitania 1997	No awareness test; no randomization of medications
Chiu 2007	No awareness test
Chortkoff 1995	Volunteers not surgery

Study	Reason for exclusion
Cirillo 2012	Unable to find either the citation or full-text paper in PubMed (despite <i>European Journal of Anaesthesiology</i> 2012 search), Embase or Cochrane databases
Clark 2009	No awareness test
Coetzee 1998	No randomization of medications for anaesthesia but narcotic randomized at wound closure
Coppens 2010	No awareness test
Cormack 1979	No randomization of medications
Crawford 1976	No randomization of medications
Dahaba 2009	No awareness test
Dahaba 2010	No randomization of medications
Davidson 2005	No randomization of medications
Davies 1996	No randomization of medications
De Cosmo 2008	No awareness test
De Kock 2005	No awareness test
Ding 1993	No awareness test
Ding 2007	Not general anaesthesia - retrospective study of postoperative complications
Diz 2010	No awareness test
Doufas 2009	No randomization of medications during surgery; not performed after skin incision
Downing 1976	Expert commentary about awareness - retrospective review records for patient awareness reports
Dressler 2007	No randomization of medications
Driscoll 2007	No randomization of medications
Eisele 1976	No randomization of medications
Ekman 2004a	No randomization of medications
Ekman 2004b	No awareness test
El-Kerdawy 2000	No awareness test
Eldar 1992	Wake up test
Erhan 2003	No awareness test
Erk 2007	No awareness test
Eroglu 2003	No awareness test

Study	Reason for exclusion
Evans 1988	No randomization of medications by anaesthetic but only word test
Fahlenkamp 2010	No awareness test
Fairley 1956	No randomization of medications
Famewo 1976	No randomization of medications
Farag 2006	No awareness test
Filipov 2007	No awareness test in abstract - unable to get full paper through University of Utah Library (they tried through all possible sources)
Fisher 2006	No randomization of medications
Fitzgerald 2001	Volunteers not surgery
Flaishon 1997	No randomization of medications
Flaitz 1986	No randomization of medications
Flier 1986	No randomization of medications
Forrest 1990	Not general anaesthesia
Frank 2000	No awareness test
Fung 2008	Wake up test performed
Gajraj 1998	Not general anaesthesia
Gajraj 1999	Not general anaesthesia
Gan 1997	No awareness test
Gan 1999	No awareness test
Gazzanelli 2005	No randomization of medications
Ge 2003	No randomization of medications
Gelman 1984	No awareness interview
Ghabash 1996	No awareness test
Ghoneim 1988	No awareness test
Ghoneim 1998	Volunteers not surgery
Ghoneim 2007	No randomization of medications
Glass 1997	Volunteers not surgery
Gordon 1994	No randomization of medications

Study	Reason for exclusion
Gregory 1969	No randomization of medications
Greif 2002	Volunteers not surgery
Gross 1988	No awareness test
Gu 2010	No awareness test
Guignard 2001	No randomization of medications
Gunawardane 2002	No randomization of medications
Guo 2012	Not general anaesthesia
Gupta 2006	No awareness test
Hadzidiakos 2006	No randomization of medications - only word test
Haessler 1993	No awareness test
Hall 1986	No awareness test
Hans 1998	No awareness test
Harris 1971	No randomization of medications
Hartridge 1963	No randomization of medications
Hartung 1986	No awareness test
Hashimoto 2012	Unable to find either the citation or full-text paper in PubMed, Embase or Cochrane databases; peer reviewed study not published
Hayashi 2007	No awareness test
Head-Rapson 1995	No randomization of medications
Heipertz 1986	No awareness test
Heisterkamp 1975	Not general anaesthesia
Heller 2005	No randomization of medications
Hellwagner 2003	No awareness test
Hetem 2000	Volunteers not surgery
Hirschi 2000	No randomization of medications
Honarmand 2008	No awareness test
Hong 2008	No awareness test
Horn 2009	No awareness test; volunteers not surgery

Study	Reason for exclusion
Hoymork 2003	No randomization of medications
Hoymork 2005	No randomization of medications
Huang 1988	No randomization of medications
Huang 2005	No randomization of medications - randomized by child positive awareness
Hudetz 2007	No awareness test
Hughes 1994	No randomization of medications
Hughes 2009	Expert commentary about awareness
Hutchinson 1960	No randomization of medications
Iannuzzi 2005	Expert commentary about awareness
Ibrahim 2001a	Not general anaesthesia
Ibrahim 2001b	Not general anaesthesia
Inagaki 1997	No awareness test
Ingelmo 2007	No awareness test
Inglis 1993	No awareness test
Iqbal 1985	No randomization of medications
Iselin-Chaves 1998	Volunteers not surgery
Iselin-Chaves 2000	Volunteers not surgery
Iselin-Chaves 2005	No randomization of medications
Ishiyama 2005	Not general anaesthesia
Isomura 2008	Not general anaesthesia - sedated for 20 hours on average in the ICU after maxillofacial surgery (translated Jiro Kurata MD)
Ittichaikulthol 2007	No awareness test in abstract - unable to get full paper through University of Utah Library (they tried through all possible sources)
Ivanov 1969	No randomization of medications
Jacoby 1981	Volunteers not surgery
Jaffrelot 2007	No awareness test
Jelezcov 2007	No awareness test
Jelicic 1993a	No randomization of medications
Jelicic 1993b	No randomization of medications

Study	Reason for exclusion
Jellish 2000	No awareness test
Jellish 2009	No awareness test
Jensen 1996	No awareness test
Jeon 2000	No randomization of medications: IFT
Jessop 1991	Expert commentary about awareness: editorial
Jones 1990	No awareness test
Jordan 2012	Unable to find citation and/or abstract or full paper after searching other databases; emailed author with no response
Jospin 2007	No awareness test
Juckenhofel 1999	No awareness test
Katoh 1993	No awareness test
Katoh 1994	No awareness test
Kavey 1979	No randomization of medications
Kennedy 1985	No awareness test
Kenny 1999	No randomization
Kerssens 2002	No randomization of medications
Kerssens 2003	No randomization and pre-incision (IFT)
Kerssens 2005a	Volunteers not surgery
Kertai 2011	No awareness test
Kestin 1990	Not general anaesthesia
Kevin 2002	No randomization of medications
Khandwala 2008	Unrelated study
Kiernan 1995	Volunteers not surgery
Kim 1978	No randomization of medications
Kim 2010	No awareness test
Kliempt 1999	No randomization of medications
Kocaman 2007	No randomization of medications
Kokki 2007	No awareness test

Study	Reason for exclusion
Kotiniemi 1996	No awareness test as performed in adults only behavioural changes
Kreuer 2001	No randomization of medications
Lapidus 2007	Unrelated study
Latto 1977	No randomization of medications
Lefoll-Masson 2007	No randomization of medications
Lehmann 2000	Paper retracted; survey response: email a@klilu.de
Lehmann 2002	Paper retracted; survey response: email a@klilu.de
Lehmann 2003	Paper retracted; survey response: 21 January 2011 a@klilu.de
Lequeux 2003	No awareness test evaluation - memory tapes randomized; survey response: 10 February 2011 Lequeux pilequeu@ulb.ac.be
Leslie 2007	No randomization of medications
Levine 1993	No awareness test
Liao 2010	No awareness test
Liou 1994	No randomization of medications
Liu 2005	No awareness test
Liu 2006	No awareness assessment; ROB survey response m.fischler@hopital-foch.org
Liu 2009	No awareness test
Lopez 2007	No randomization of medications
Low 2007	Not general anaesthesia; no awareness test
Luginbuhl 2007	No awareness test
Luginbuhl 2010	No awareness test
Lyons 1991	No randomization of medications
Magni 2009	No awareness test
Mahomedy 1976	No randomization of medications
Malek 2009	No awareness test - abstract only available
Malek 2010	No randomization of medications
Malviya 2009	No randomization - children 5 to 15 - expert adjudication
Martorano 2008	No awareness test

Study	Reason for exclusion
Mathews 2008	No awareness test
Maybauer 2007	No awareness test
Mayer 2007	No awareness test
Mellema 2010	No awareness test
Messahel 2003	No randomization of medications
Messahel 2007	No randomization of medications
Mi 1998	No awareness test
Mi 1999	No awareness test
Mirakhur 1986	No awareness test
Moerman 1995	No randomization of medications
Motsch 1996	No awareness test
Mourisse 2007	No awareness test
Moustafa 2008	No awareness test
Nakagawa 2001	No awareness test and not general anaesthesia
Nelskyla 2001	No awareness test
Ng 1974	No randomization of medications
Nishijima 1999	No awareness test: "quality of awareness" by how fast the patients became aware and oriented after discontinuation of general anaesthetics. Translator Jiro Kurata MD.
O'Sullivan 1988	No randomization
Oikkonen 1994	No awareness test
Onaka 1998	No awareness test; abstract only
Oudenaarden 1979	No awareness test no randomization
Padmanabhan 2009	Not general anaesthesia
Panousis 2007	No randomization of medications
Philbin 1990	No awareness test
Philipp 2002	No awareness test - CPB membrane associated with 50% reduction isoflurane level
Piano 2007	No randomization of medications
Pomfrett 2009	No awareness test

Study	Reason for exclusion
Pompeo 2007	Unrelated study
Porter 2008	No randomization of medications
Pryor 2010	No awareness test and volunteers not surgery
Qi 2014	Unable to find either the citation or full-text paper in PubMed, Embase or Cochrane databases; unable to identify authors communication information
Rabiee 2012	Unable to find either the citation or full-text paper in PubMed, Embase or Cochrane databases; emailed author with no response
Reinhart 1985	Not general anaesthesia
Russell 1997	No randomization of medications; ROB survey response i.f.russell@hull.ac.uk : randomized only for audio message; only one anaesthetic used
Russell 2001	No randomization of medications ; ROB survey response i.f.russell@hull.ac.uk : randomized audio message
Samuelsson 2007	No randomization of medications
Schneider 2005	Wake up test
Schraag 1998	No awareness test
Schroeck 2010	No randomization of medications
Schulz 2007	No awareness test
Schwender 1994	No awareness test
Schwender 1991a	No randomization of medications
Schwender 1991b	No randomization of medications
Schwender 1993	No awareness test
Schwender 1997	No randomization of medications
Schwieger 1989	No randomization of medications
Schwieger 1991	No randomization of medications
See 2007	No awareness test
Shariffuddin 2007	No awareness test
Shiau 2007	No randomization of medications or ADMs - randomization at end of surgery (desflurane stopped before closure)
Shimohata 2007	Unrelated study - sleep apnoea under anaesthesia and death
Short 1991	No awareness test

Study	Reason for exclusion
Sidiropoulou 2008	No awareness test
Sintavanuruk 2010	No awareness test
Skaja 2006	No awareness test
Sosner 2010	No awareness test abstract only available
Spaulding 1984	Not general anaesthesia; volunteers
Stonell 2006	No randomization only word test. Survey response: 4 February 2011, Kate Leslie kate.leslie@mh.org.au
Struys 2001	No awareness test
Suarez 1994	No randomization
Suliman 2007	Not general anaesthesia; no randomization
Terblanche 2008	No awareness test
Ting 2004	Wake up test
Toraman 2013	Unable to find either the citation or full-text paper in PubMed, EMBASE or Cochrane databases
Treggiari 2009	ICU not surgery
Trillo 2009	No awareness test
Tufano 2000	No awareness test: Quote: "There was no significant differences in the incidence of intraoperative responses between groups." email address not available to send survey to clarify the author's definition of "intraoperative responses"
Tunstall 1981	No awareness test
Turan 2010	No awareness test
Turner 1969	No randomization of medications
Ueyama 1986	No randomization of medications
Valtonen 1988	Not general anaesthesia - sedation for cardioversion
Vanacker 2002	No randomization of medications
Vanacker 2007	No awareness test
Vann 2007	Not general anaesthesia
Velly 2007	No awareness test
Wanatabe 1984	No randomization of medications
Wang 2005	No randomization of medications

Study	Reason for exclusion
Wang 2007	Volunteers not surgery
Watanabe 1998	Volunteers not surgery
Weber 2005	No aware test - indirect signs only in paediatrics
Weber 2009	No awareness test
Wellisch 2012	Unable to find either the citation or full-text paper in PubMed, Embase or Cochrane databases
Wihelm 2000	No awareness test
Wilson 1970	Expert commentary about awareness: letter to editor
Wu 2005	No randomization of medications
Xu 2009	No randomization of medications
Yan 2005	No awareness test: Zancong translator quote: "Awareness was checked based on OAAS criteria post surgery. The paper does not mention follow up with pts on any memory during the surgery"
Yan 2014	No awareness test - translation issue: no assessment patient reports of awareness rather "awareness" was translated instead of "consciousness or awake, etc.": "The recovery time of awareness and extubation time in the treatment group"
Yang 1994	No awareness test; no randomization
Yeh 2009	No awareness test; post-surgery randomization
Yi 2008	No awareness test
Yildiz 2007	No awareness test
Zhang 2009	No awareness test
Zohar 2007	No awareness test

Acronyms and abbreviations

ADM = anaesthesia depth monitor; CPB = cardiopulmonary bypass; EEG = electroencephalogram; ICU = intensive care unit; IFT = isolated forearm technique; OAAS = Observer's Assessment of Alertness/Sedation; POCD = postoperative cognitive dysfunction; ROB = risk of bias

Characteristics of studies awaiting assessment [ordered by study ID]

Aceto 2015

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: pending full-text review Sex: female and male Age: pending full-text review Procedure: elective thyroidectomy

Aceto 2015 (Continued)

Study size: 130

Interventions	<p>Randomized portion of anaesthetic: TIVA vs volatile</p> <p>Maintenance: auditory recording was presented to patients during anaesthesia maintenance</p> <p>Intervention 1: BIS-guided group in which sevoflurane MAC was adjusted on the basis of BIS values</p> <p>Intervention 2: haemodynamic parameters (HP)-guided group in which MAC was adjusted based on HP</p>
Outcomes	<p>Primary outcomes: "...whether Bispectral Index (BIS)-guided anaesthesia might decrease sevoflurane minimum alveolar concentration (MAC) when compared with haemodynamically-guided anaesthesia, and to search for a MAC threshold useful for preventing arousal, dream recall and implicit memory"</p> <p>Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class pending full text review</p>
Notes	<p>Non-randomized portion of anaesthetic: parts of IV vs parts of volatile agent N₂O yes/narcotics/hypnotics bolus/NMBs induction no/maintenance no</p> <p>Anaesthesia was induced with propofol 2 mg kg⁽⁻¹⁾, fentanyl 3 µg kg⁽⁻¹⁾ and cis-atracurium 0.15 mg/kg⁽⁻¹⁾</p> <p>Dream recall and explicit/implicit memory were investigated upon awakening and approximately after 24 h</p> <p>RESULTS: mean sevoflurane MAC during auditory presentation was similar in the 2 groups (0.85 ± 0.16 and 0.87 ± 0.17 (P = 0.53) in BIS-guided and HP-guided groups, respectively). Frequency of dream recall was similar in the 2 groups: 27% (N = 17) in BIS-guided group, 18% (N = 12) in HP-guided group, P = 0.37</p> <p>In both groups, dream recall was less probable in patients anaesthetized with MAC values >= 0.9 (area under ROC curve = 0.83, sensitivity = 90%, and specificity = 49%)</p> <p>Conclusion: BIS-guided anaesthesia was not able to generate different MAC values compared to HP-guided anaesthesia</p> <p>Independent of the guide used for anaesthesia, a sevoflurane MAC over 0.9 was required to prevent postoperative dream recall</p>

Asouhidou 2015

Methods	<p>Study design: randomized parallel groups</p> <p>Study dates: not stated</p>
Participants	<p>Country: pending full-text review</p> <p>Sex: female and male</p> <p>Age: pending full-text review</p> <p>ASA: pending full-text review</p> <p>Procedure: craniotomy for aneurysm clipping or tumour dissection were randomly</p> <p>Study. size: 42</p>
Interventions	<p>Randomized portion of anaesthetic:</p>

Asouhidou 2015 (Continued)

Intervention 1: propofol, N=21

Intervention 2: sevoflurane, N=21

esmolol group received 500 µg/kg of esmolol bolus 10 min before induction of anaesthesia, followed by additional 200 µg/kg/min of esmolol

inspired concentration of sevoflurane and the infusion rate of propofol were adjusted in order to maintain a BIS value between 40–50

Bispectral Index-BIS and cardiac output

Outcomes

Primary outcomes: effect of esmolol on the consumption of propofol and sevoflurane in patients undergoing craniotomy

Secondary outcome: Awareness/wakefulness as defined using an awareness classification system (see Table 1) class pending full-text review

Notes

Non-Randomized portion of anaesthetic: parts of IV vs parts of volatile agent N20:U/ narcotics/hypnotics Bolus/NMBsinduction Y/maintenance U/ADM: Y: Bispectral Index-BIS

Anaesthesia was induced with propofol, fentanyl and a single dose of cis-atracurium, followed by continuous infusion of remifentanyl and either propofol or sevoflurane

Esmolol group: doses of propofol and sevoflurane were 18-50 µg/kg/min and 0.2-0.5 MAC

Control group: 100-150 µg/kg/ and 0.9-2.0 MAC (P = 0.000 for both groups)

All procedures were anaesthesiologically uneventful with no episodes of intraoperative emerge

Conclusions: Esmolol is effective not only in attenuating intraoperative haemodynamic changes related to sympathetic overdrive but also in minimizing significant propofol and sevoflurane requirements without compromising the haemodynamic status

ClinicalTrials.gov Identifier: NCT02455440 . Registered 26 May 2015

Elbadawy 2015

Methods

Study design: randomized parallel groups

Study dates: not stated

Participants

Country: pending full-text review

Sex: female and male

Age: pending full-text review

ASA: I and II

Procedure: orthopaedic procedures

Study size: 40

Interventions

Randomized portion of anaesthetic:

Intervention 1: TIVA, N = 20

Intervention 2: volatile anaesthesia, N = 20

BIS-guided (40 to 60)

Elbadawy 2015 (Continued)

Outcomes Primary outcomes: no implicit memory of previously introduced auditory material

Secondary outcome: no awareness/wakefulness as defined using an awareness classification system (see Table 1) class pending full-text review

Notes **Non-randomized portion of anaesthetic: parts of IV vs parts of volatile agent N₂O unclear/narcotics/hypnotics bolus/NMBs induction yes/maintenance yes/ADM: yes: BIS-guided (40 to 60)**

Anaesthesia was induced with propofol and maintained with propofol, fentanyl, and cis-atracurium. In Group II anaesthesia was induced with propofol and maintained with sevoflurane, fentanyl, and cis-atracurium

Explicit memory was evaluated by asking 4 standard questions regarding intraoperative awareness. Free recall and recognition tests for implicit memory testing were carried out 30 min and 120 min after recovery.

BIS-controlled anaesthesia appears to abolish implicit memory

No PubMed citation

Hoymork 2007

Methods Study design: randomized parallel groups

Study dates: not stated

Participants Country: Norway and Denmark

Sex: female and male

Age: NA

ASA: I and II

Procedure: laparoscopy or breast/surface surgery

Study size: 55

Interventions Quote: "The patients were randomly allocated to whether BIS or CSM was placed on the upper part of the forehead or closer to the eyebrows, where an eventual EMG influence could be more pronounced"

Outcomes Quote: "None of our patients reported any recall from the operation, and clinical awakening was never observed during surgery, despite one episode of movement"

Notes Quote: "In conclusion, the cerebral state monitor was a satisfactory alternative to BIS for monitoring hypnotic effect in 87% of our patients. In 13% of the patients, CSI displayed values indicating an awake state despite clinical sleep, all correctly identified with the BIS. Our study was done with the very first version of the CSM, while the BIS monitor has undergone several revisions. Clinical studies in other patient populations, undergoing different anaesthetic regimens, are warranted"

Jiang 2016

Methods Study design: randomized parallel groups

Study dates: not stated

Jiang 2016 (Continued)

Participants	Country: pending full-text review Sex: female and male Age: pending full-text review ASA: pending full-text review Procedure: laparoscopic radical gastrectomy Study size: 100
Interventions	<p>Randomized portion of anaesthetic: total intravenous anaesthesia (TIVA) and combined intravenous and inhaled anaesthesia (CIIA)</p> Intervention 1: TIVA, propofol and remifentanyl: target controlled infusion (TCI), N = 50 Intervention 2: CIIA, sevoflurane and continuous infusion of remifentanyl after anaesthesia induction, N = 50 State entropy (SE) maintained in the range of 45 to 60
Outcomes	Primary outcomes: concentrations: epinephrine, norepinephrine and dopamine; durations of surgical operation, breathing recovery, extubation, awakening, and postoperative orientation recovery recorded; and 48 H postoperative adverse reactions: time patient becomes calm for 5 min after entering the operating theatre (T0); upon completion of pneumoperitoneum (PPT) (T1); 15 min after PPT (T2); intraoperative detection (T3), immediately after extubation (T4); and 15 min after extubation (T5) Secondary outcome: no awareness/wakefulness as defined using an awareness classification system (see Table 1): class pending full-text review
Notes	<p>Non-randomized portion of anaesthetic: parts of IV vs parts of volatile agent N₂O unclear/narcotics/hypnotics bolus/NMBs induction no/maintenance no/ADM: yes: entropy indices</p> Conclusion: at the same anaesthetic depth, the CIIA method outperforms the TIVA method in suppressing the stress response and obtaining smooth awakening after laparoscopic radical gastrectomy for patients with gastric cancer; therefore, the CIIA method has a better anaesthetic effect

Khanjani 2014

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: pending full-text review Sex: female and male Age: pending full-text review ASA: pending full-text review Procedure: caesarean section Study size: 90
Interventions	<p>Randomized portion of anaesthetic:</p> Intervention 1: inhalation (isoflurane) maintenance: isoflurane 1 MAC

Khanjani 2014 (Continued)

Intervention 2: intravenous protocol (propofol) propofol 100 µg/kg/minute
Bispectral Index (BIS) between 45 and 60

Outcomes

Primary outcomes: APGAR score of newborns
Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class pending full-text review
Quote: "...four cases of confirmed awareness were found in the propofol group and three cases in the Isoflurane group (8/9% vs. 6/7%), but the Apgar scores were comparable between the two groups"

Notes

Non-randomized portion of anaesthetic: parts of IV vs parts of volatile agent N₂O yes/narcotics/hypnotics bolus/NMBs induction no/maintenance no
Induction: propofol and succinylcholine

Lequeux 2014

Methods

Study design: randomized parallel groups
Study dates: not stated

Participants

Country: pending full-text review
Sex: female and male
Age: pending full-text review
ASA: pending full-text review
Procedure: pending full-text review
Study size: 120

Interventions

Randomized portion of anaesthetic:
Intervention 1: high-dose: remifentanil effect-site concentration (in ng/mL) was always double that of propofol (in µg/mL)
Intervention 2: low-dose: half of above
Patients in these 2 groups were played a list of 20 words via headphones during surgery
Intervention 3: control for memory tests: not played any words during anaesthesia

Outcomes

Primary outcomes: no implicit learning of intraoperative auditory stimuli
Secondary outcome: no awareness/wakefulness as defined using an awareness classification system (see Table 1): class pending full-text review

Notes

Non-randomized portion of anaesthetic: parts of IV vs parts of volatile agent N₂O yes/narcotics/hypnotics bolus/NMBs induction no/maintenance no
All patients were anaesthetized with a target-controlled infusion of propofol and remifentanil, targeting a bispectral index (BIS) value of 50
Conclusions: could not demonstrate the presence of implicit or explicit memorization under propofol-remifentanil anaesthesia either with a low- or a high-dose opioid anaesthetic regimen

Mehmandoost 2013

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: Iran Sex: female Age: 18 to 35 years ASA: ASA I-II Procedure: caesarean section Study size: 90
Interventions	Intervention 1: propofol (100 µg/kg/min) Intervention 2: isoflurane 1 MAC (minimum alveolar concentration) Both groups titrated to BIS 45 to 60
Outcomes	Quote: "There was not a significant difference between two groups in incidence of awareness"
Notes	Awaiting Persian translation

Rajan 2015

Methods	Study design: randomized parallel groups Study dates: not stated
Participants	Country: pending full-text review Sex: female and male Age: pending full-text review ASA: pending full-text review Procedure: caesarean section Study size: 40
Interventions	Randomized portion of anaesthetic: pre-induction: hypnotic vs placebo Just before induction of anaesthesia Intervention 1: 0.25 mg/kg ketamine, N = 20 Intervention 2: 5 ml normal saline intravenously (IV), N = 20
Outcomes	Primary outcomes: efficacy and safety of low-dose ketamine, used as an adjunct analgesic and amnesic Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class pending full-text review

Rajan 2015 (Continued)

Quote: "Higher number of the patients in Group C had intraoperative lacrimation as compared to Group K (50% vs. 0%, P < 0.001). Ten percent of the patients in Group C had hallucinations/recall of intraoperative events while none of the patients in Group K experienced the same, but the difference was statistically insignificant (P = 0.487)"

Notes

Non-randomized portion of anaesthetic: parts of volatile agent N₂O yes/narcotics/hypnotics bolus/NMBs induction yes/maintenance unclear

After intubation, patients were ventilated with O₂ and N₂O (40%:60%) with 0.7% end-tidal isoflurane. Fentanyl and midazolam were given following delivery of the baby

Results: Pre-induction haemodynamic parameters and those recorded at 1 min after induction were comparable in both groups. However, heart rate and systolic blood pressure recorded after intubation (at 3, 5, 7, 9, 12, 15, 20, 30 and 45 min after induction) showed significantly high values in Group C (P < 0.05). Mean arterial pressure also showed a similar pattern

Umbilical vein pO₂, pCO₂ and pH were comparable in both groups. In Group C, intraoperative lacrimation (50% vs 0%) and hallucinations/recall of intraoperative events (10% vs 0%) were high

Conclusion: IV ketamine 0.25 mg/kg can be safely used as an adjunct analgesic and amnesic to attenuate haemodynamic responses during caesarean section under GA without affecting the foetal outcome

Xie 2015

Methods

Study design: randomized parallel groups
Study dates: December 2013 to May 2014

Participants

Country: pending full-text review
Sex: female and male
Age: pending full-text review
ASA: pending full-text review
Procedure: elective surgery
Study size: 150

Interventions

Randomized portion of anaesthetic: TIVA vs volatile: implicit memory tests

Intervention 1: induced and maintained with sevoflurane, N = 50
Intervention 2: induced and maintained with propofol, N = 50
Intervention 1 and 2: given a list of test materials to remember and listen before the anaesthesia
Intervention 3: given the same test materials, and received test with the PDP in 12 to 36 hours before surgery, N = 50

Outcomes

Primary outcomes: effects of sevoflurane and propofol on preoperative implicit memories
Secondary outcome: awareness/wakefulness as defined using an awareness classification system (see Table 1): class pending full-text review

Notes

Non-randomized portion of anaesthetic: parts of IV vs parts of volatile agent N₂O yes/narcotics/hypnotics bolus/NMBs induction unclear/maintenance unclear

Xie 2015 (Continued)

Conclusion: propofol and sevoflurane can decrease the score of explicit memory after anaesthesia within 12 to 36 hours, and there are no significant differences in explicit memory between the 2 drugs. Both propofol and sevoflurane can decrease the score of implicit memory, but the influence of sevoflurane on the implicit memory is less than propofol within 12 to 36 hours

No PubMed citation

Acronyms and abbreviations

ADM = anaesthesia depth monitor; ASA = American Society of Anesthesiology; BIS = Bispectral Index; cis = cisatracurium is one of the 10 isomers of the parent molecule, atracurium; CIIA = combined intravenous and inhaled anaesthesia; GA = general anaesthesia; H = hours; HP = haemodynamic parameters; IV = intravenous; kg = kilogram; MAC = minimal alveolar concentration; µg = micrograms; min = minute; mL = millilitres; N = sample size; ng = nanograms; NMB = neuromuscular blocking agents; PDP = parallel distributed processing; PPT = pneumoperitoneum; ROC = receiver operating characteristic; SE = state entropy; T = time; TCI = target-controlled infusions; TIVA = total intravenous anaesthesia; vs = versus

DATA AND ANALYSES

Comparison 1. Anaesthesia depth monitors (BIS and M-entropy) versus standard clinical and electronic monitoring

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Definite and possible awareness (Grade 4)			Other data	No numeric data

Analysis 1.1. Comparison 1 Anaesthesia depth monitors (BIS and M-entropy) versus standard clinical and electronic monitoring, Outcome 1 Definite and possible awareness (Grade 4).

Study	ADM Events	N	Definite and possible awareness (Grade 4)		OR 95% CI	Totals	Summary OR 95% CI; P value I ² ; heterogeneity P value
			SCP Events	N			
Avidan 2008	6	967	3	974	2.02 (0.50 to 8.10)	—	—
Avidan 2011	20	2861	8	2852	2.50 (1.10 to 5.69)	—	—
Gruenewald 2007	0	37	1	35	0.31 (0.01 to 7.78)	—	—
Kerssens 2009	2	67	1	61	1.85 (0.16 to 20.89)	—	—
Mashour 2012	18	9460	19	9376	0.94 (0.49 to 1.79)	—	—
Mozafari 2014	9	163	7	170	1.36 (0.49 to 3.74)	—	—
Myles 2004	22	1225	27	1238	0.82 (0.46 to 1.45)	—	—
Puri 2003	0	14	1	16	0.36 (0.01 to 9.47)	—	—
Zhang 2011	8	2919	21	2309	0.30 (0.13 to 0.68)	85/17,713 vs 88/17,031	0.98 (0.59 to 1.62); P = 0.93 I ² = 49%; P = 0.04

Comparison 2. Anaesthesia depth monitors (BIS) versus standard clinical and electronic monitoring

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Definite and possible awareness (Grade 4)			Other data	No numeric data

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2 Definite awareness (Grade 4)			Other data	No numeric data

Analysis 2.1. Comparison 2 Anaesthesia depth monitors (BIS) versus standard clinical and electronic monitoring, Outcome 1 Definite and possible awareness (Grade 4).

Study	ADM Events	N	Definite and possible awareness (Grade 4)			Alarms/alerts ----- Inhalation vs TIVA	Subgroups ----- Summary OR 95% CI	Totals ----- Summary OR 95% CI; P value I ² ; heterogeneity P value Subgroup interaction P value
			SCP Events	N	OR 95% CI			
Avidan 2008	6	967	3	974	2.02 (0.50 to 8.10)	Alarms ----- Inhalation	—	—
Avidan 2011	20	2861	8	2852	2.50 (1.10 to 5.69)	Alarms ----- Inhalation	—	—
Mashour 2012	18	9460	19	9376	0.94 (0.49 to 1.79)	Alarms and alerts ----- Inhalation	Alarms and/or alerts inhalational ----- 1.51 (0.45 to 5.07)	—
Myles 2004	22	1225	27	1238	0.82 (0.46 to 1.45)	None ----- Majority TIVA	—	—
Zhang 2011	8	2919	21	2309	0.30 (0.13 to 0.68)	None ----- Exclusively TIVA	No alarms TIVA ----- 0.55 (0.15 to 1.96)	74/17,432 vs 78/16,749 ----- 0.96 (0.35 to 2.65); P = 0.93 I ² = 68%; P = 0.01; P = 0.17 (subgroup)

Analysis 2.2. Comparison 2 Anaesthesia depth monitors (BIS) versus standard clinical and electronic monitoring, Outcome 2 Definite awareness (Grade 4).

Study	ADM Events	N	Definite awareness (Grade 4)			Subgroup ----- Summary OR 95% CI	Subgroups ----- Summary OR 95% CI	Totals ----- Summary OR 95% CI; P value I ² ; heterogeneity P value Subgroup interaction P value
			SCP Events	N	OR 95% CI			
Avidan 2008	2	967	2	974	1.01 (0.14 to 7.17)	Alarms ----- Inhalation	—	—

Study	ADM Events	N	Definite awareness (Grade 4)			Subgroup	Subgroups	Totals
			SCP Events	N	OR 95% CI			
						Summary OR 95% CI	Summary OR 95% CI	Summary OR 95% CI; P value I ² ; heterogeneity P value Subgroup interaction P value
Avidan 2011	8	2861	2	2852	4.00 (0.85 to 18.83)	Alarms ----- Inhalation		
Mashour 2012	8	9460	11	9376	0.72 (0.29 to 1.79)	Alarms and alerts ----- Inhalation	Alarms and/or alerts inhalational ----- 1.10 (0.31 to 3.97)	
Myles 2004	2	1225	11	1238	0.18 (0.04 to 0.82)	None ----- Majority TIVA		
Zhang 2011	4	2919	15	2309	0.21 (0.07 to 0.63)	None ----- Exclusively TI-VA	No alarms TIVA ----- 0.20 (0.04 to 0.96)	24/17,432 vs 41/16,749 ----- 0.60 (0.13 to 2.75); P = 0.40 I ² = 60%; P = 0.02; P = 0.08 (subgroup)

Comparison 3. Thiopentone with and without added hypnotic drugs (ketamine, etomidate)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Wakefulness (IFT studies) (Grades 4, 5)			Other data	No numeric data

Analysis 3.1. Comparison 3 Thiopentone with and without added hypnotic drugs (ketamine, etomidate), Outcome 1 Wakefulness (IFT studies) (Grades 4, 5).

Study	Ketamine, etomidate Events	N	Wakefulness (IFT studies) (Grades 4, 5)		RR 95% CI	Totals	Summary RR 95% CI; P value
			Thiopentone Events	N			
Baraka 1989	4	30	14	20	0.19 (0.06 to 0.45)	—	—
Russell 1986	2	30	11	25	0.10 (0.03 to 0.55)	—	—
Schultetus 1986	5	23	7	13	0.39 (0.14 to 1.04)	11/83 vs 32/58	0.18 (0.09 to 0.41); P = 10 ⁻⁷

Comparison 4. Thiopentone and ketamine vs benzodiazepine anaesthetic techniques

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Awareness (Grade 4)			Other data	No numeric data

Analysis 4.1. Comparison 4 Thiopentone and ketamine vs benzodiazepine anaesthetic techniques, Outcome 1 Awareness (Grade 4).

Study	Benzodiazepines Events	N	Awareness (Grade 4)		RR 95% CI	Totals	Summary RR 95% CI; P value
			Ketamine, thiopentone Events	N			
Ellingson 1977	0	13	4	13	0.00 (0.00 to 1.12)	—	—
Haram 1981	0	39	5	40	0.00 (0.00 to 0.83)	—	—
McNulty 1995	0	71	3	25	0.00 (0.00 to 0.56)	—	—
Miller 1996	2	69	4	21	0.12 (0.02 to 0.83)	2/192 vs 16/99	0.17 (0.02 to 0.25); P < 10 ⁻⁷

Comparison 5. Caesarean section low- and high-dose inhaled agents

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Awareness (Grade 4)			Other data	No numeric data

Analysis 5.1. Comparison 5 Caesarean section low- and high-dose inhaled agents, Outcome 1 Awareness (Grade 4).

Study	High dose Events	N	Awareness (Grade 4)		RR 95% CI	Totals	Summary RR 95% CI; P value
			Low dose Events	N			
Abboud 1985	0	65	2	16	0.00 (0.00 to 0.80)	—	—
Crawford 1985	3	384	19	393	0.13 (0.04 to 0.51)	3/449 vs 21/409	0.13 (0.04 to 0.43); P = 0.0001

ADDITIONAL TABLES
Table 1. Classification of intraoperative cognitive states

Grade	Intraoperative state	Intraoperative state	Immediate postoperative state	Late postoperative state (> month)	Descriptor	Exemplar study or review
0	Unconscious	No signs light anaesthesia, no response to command	No recall	No recall	Adequate anaesthesia	Russell 1997

Table 1. Classification of intraoperative cognitive states (Continued)

1	Conscious	Signs light anaesthesia/response to command	No recall	No recall or emotional sequelae	Intraoperative <i>wakefulness</i> with obliterated explicit and implicit memory	Andrade 2008
2	Conscious; word stimuli presented	Signs/response to command	No explicit recall, implicit memory for word stimuli	No explicit recall, implicit memory for word stimuli but no emotional sequelae	Intraoperative <i>wakefulness</i> with subsequent <i>implicit memory</i>	Merikle 1996
3	Conscious	Signs/response to command	No recall	PTSD/nightmares but no explicit recall	Intraoperative <i>wakefulness</i> with <i>implicit emotional memory</i>	Wang 2000
4	Conscious	Signs/response to command	Explicit recall with or without pain	Explicit recall but no emotional sequelae	<i>Awareness</i> but resilient-patient	Sandin 2000
5	Conscious	Signs/response to command	Explicit recall with distress and/or pain	PTSD/nightmares with explicit recall	<i>Awareness</i> with <i>emotional sequelae</i>	Osterman 2001

First presented at the 7th International Symposium on Memory and Awareness in Anaesthesia, Munich, Germany, March 2008. Revised in publication [Wang 2012](#). Permission granted by John Wiley & Sons Inc. on 17 June 2014.
 PTSD = post-traumatic stress disorder

APPENDICES

Appendix 1. Dryad website link

Data available from the Dryad Digital Repository: <http://dx.doi.org/10.5061/dryad.2vn65>

Appendix 1: Associated Dryad word document title within each section of review

Background section

Description of the condition

Psychological sequelae&Light anaesthesia.docx

Cardiovascular Function & Pathologies

PTSD Diagnostic Criteria.docx

Wakefulness endnote 2014 annotate.docx

Why it is important to do this review section

Psychological sequelae&Light anaesthesia.docx

Methods

Types of outcomes

Refinement of Sebel classification system.docx

Results

Description of studies

Included studies

Sample size N=160.xlsx

Language.xlsx

surgical risk.xlsx

primary.secondary.outcome.xlsx

Nomenclature.docx

ROB.results.xlsx

Excluded studies

Excluded RCTs.315.xlsx

Trial location

Continents.countries.xlsx

Anaesthetic interventions

Frequencies of anaesthetics techniques

Anesthesia Techniques.xlsx

MR.use.xlsx

ADM interventions

ADM all types.xlsx

Classification of outcomes

Classification.Wang.Messina.Ward.grade

Response to complex command.xlsx

Effect of interventions

Primary outcomes

Awareness

patient awareness reports.xlsx

Wakefulness

wakeful v wakeful.aware v wakeful.xlsx

Discussion

Summary of main results

Classification.Wang.Messina.Ward.grade.xlsx

Thiopental studies.docx

Overall completeness and applicability of evidence

RCTs not used in comparisons.xlsx

Quality of the evidence

Quality of the evidence.docx

ACE130 reviewer#1.defense RCT merge criteria.summary.docx

ACE 130 reviewer#1.defense RCT merge criteria.details.docx

ACE 130 Peer reviewer.#2.docx

ACE 130 reviewer #2.MWa response.classification table.docx

ACE 130 reviewer #3 peer review.docx

ACE 130 consumer TL.am.docx

Potential biases in the review process

Potential biases in the review process.docx

Appendix 2. Search strategy for CENTRAL

- #1 MeSH descriptor: [General Surgery] explode all trees
- #2 MeSH descriptor: [Surgical Procedures, Operative] this term only
- #3 MeSH descriptor: [Anesthesia] this term only
- #4 MeSH descriptor: [Autonomic Agents] this term only
- #5 MeSH descriptor: [Anesthetics] this term only
- #6 MeSH descriptor: [Muscle Relaxants, Central] explode all trees
- #7 MeSH descriptor: [Neuromuscular Blocking Agents] explode all trees
- #8 surgical stimulation* or an?esth*.ti,ab or ((neuromuscular or nondepolarizing) near blocking) or muscle relaxant*.ti,ab
- #9 #1 or #2 or #3 or #4 or #5 or #6 or #7 or #8
- #10 MeSH descriptor: [Wakefulness] explode all trees
- #11 MeSH descriptor: [Memory] explode all trees
- #12 MeSH descriptor: [Awareness] explode all trees
- #13 MeSH descriptor: [Attention] explode all trees
- #14 (memor* or wakefulness or awareness or attention or awakesness or wake up threshold):ti,ab
- #15 #10 or #11 or #12 or #13 or #14
- #16 MeSH descriptor: [Immobilization] explode all trees
- #17 MeSH descriptor: [Stress Disorders, Post-Traumatic] explode all trees
- #18 MeSH descriptor: [Paralysis] explode all trees
- #19 MeSH descriptor: [Unconsciousness] explode all trees
- #20 MeSH descriptor: [Consciousness] explode all trees
- #21 MeSH descriptor: [Dreams] explode all trees
- #22 MeSH descriptor: [Panic] this term only
- #23 MeSH descriptor: [Amnesia] this term only
- #24 paraliz* or dream*.ti,ab or post?traumatic*.ti,ab or (an?esthetic near depth) or (((evok* near potential*) or eeg or electroencephalogra*) and processed)
- #25 #16 or #17 or #18 or #19 or #20 or #21 or #22 or #23 or #24
- #26 #9 and (#15 or #25)

Appendix 3. Search strategy for MEDLINE (Ovid SP)

1. General Surgery/ or anaesthesia/ or Surgical Procedures, Operative/ or Autonomic Agents/ or Anesthetics/ or Muscle Relaxants, Central/ or Neuromuscular Blocking Agents/ or surgical stimulation*.mp. or an?esth*.ti,ab. or ((neuromuscular or nondepolarizing) adj6 blocking).mp. or muscle relaxant*.ti,ab.
2. Wakefulness/ or Memory/ or Awareness/ or Attention/ or (memor* or wakefulness or awareness or attention or awakesness or wake up threshold).mp.
3. Immobilization/ or Stress Disorders, Post-Traumatic/ or (((evok* adj3 potential*) or eeg or electroencephalogra*) and processed).mp. or Paralysis/ or Unconsciousness/ or Consciousness/ or Dreams/ or Panic/ or Amnesia/ or paraliz*.mp. or dream*.ti,ab. or post?traumatic*.ti,ab. or (an?esthetic adj6 depth).mp.
4. 1 and (2 or 3)
5. ((randomized controlled trial or controlled clinical trial).pt. or clinical trials as topic.sh. or (random* or double-blind* or placebo).ti,ab.) not (animals not (humans and animals)).sh.
6. 4 and 5

The definitions of the following abbreviations are: [tw] is the PubMed tag for text word, [pt] is the PubMed tag for publication type, and [mh] is the PubMed tag for the MeSH term.

Appendix 4. Search strategy for EMBASE (Ovid SP)

1. surgery/ or surgical-patient/ or anaesthesia/ or "agents interacting with transmitter, hormone or drug receptors"/ or anesthetic-agent/ or muscle-relaxant-agent/ or neuromuscular-blocking-agent/ or surgical stimulation*.ti,ab. or an?esth*.ti,ab. or ((neuromuscular or nondepolarizing) adj3 blocking).ti,ab. or muscle relaxant*.ti,ab.
2. wakefulness/ or memory/ or awareness/ or attention/ or (memor* or wakefulness or awareness or attention or awakesness or wake up threshold).ti,ab.
3. immobilization/ or posttraumatic-stress-disorder/ or posttraumatic-stress-disorder/ or (((evok* adj3 potential*) or eeg or electroencephalogra*) adj3 processed).ti,ab. or paralysis/ or unconsciousness/ or consciousness/ or dream/ or panic/ or amnesia/ or paraliz*.ti,ab. or dream*.ti,ab. or post?traumatic*.ti,ab. or (an?esthetic adj3 depth).ti,ab.
4. 1 and (2 or 3)

5. ((randomized controlled trial or controlled clinical trial).pt. or randomized.ab. or placebo.ab. or randomly.ab. or trial.ti.) not (animals not (humans and animals)).sh.
 6. 4 and 5

Appendix 5. Selection criteria and checklists

Selection criteria	Yes	No	Unclear/not stated or quote from paper
Study design randomized (if no, exclude)			
Drug randomized			
Brain monitor randomized			
Learning test randomized (if yes, exclude)			
Study assessed for awareness (if no/unclear, exclude)			
Study done after skin incision but before end of surgery (if no, exclude)			
ICU study (if yes, exclude)			
Blinding (patient, physician, postoperative interviewer)			
See 'Risk of bias' table			
Abbreviations for checklist table			
After skin inc	After skin incision: randomization and duration of surgery is after not before skin incision		
PMID	PubMed identification number for each study		

1 10

Checklist: Data extraction 1A

PMID	Author	Title	Date-yr	Drug or brain monitor randomized	Randomized quote in database	Aware test	Aware quote in database	After skin inc	Category rev1
------	--------	-------	---------	----------------------------------	------------------------------	------------	-------------------------	----------------	---------------

Checklist: Data extraction 2

Author	Year	IV vs IV 1,0 yes no	IV vs Inhal (balanced) 1,0	Regional + GA 1,0	Regional + GA description	Inhal vs Inhal (balanced) 1,0	Multiple anaesthesia techniques	Surgery	Induc random 1,0
--------	------	---------------------	----------------------------	-------------------	---------------------------	-------------------------------	---------------------------------	---------	------------------

Checklist: Data extraction 3

IFT 1.0	Wang chart classification	Country	Sample size, N	Explicit memory events #	% Awareness	Implicit IFT Wakefulness #	% Wakefulness	Intraoperative dream events #	% Intraoperative dreams
---------	---------------------------	---------	----------------	--------------------------	-------------	----------------------------	---------------	-------------------------------	-------------------------

Checklist: Data extraction 1B

Reviewer 1	Category rev 2	Reviewer 2	Category rev am	Reviewer am	CATEGORY FINAL	Characteristics table	Bias table	Data chart
------------	----------------	------------	-----------------	-------------	----------------	-----------------------	------------	------------

(Continued)

Main random 1,0	Premed random 1,0	Regional random + GA 1.0	Cardiac and/or placebo random 1,0	Cardiac and/or placebo random description	CPB random 1,0	OB Pre deliv. random 1,0	ADM 1,0	ADM random 1,0	Type of ADM
Postop dreams	% Postop dreams	Implicit word/music testing	% Implicit memory	Movement events #	% Movement	Check classification/bias table 1,0	Check data table 1,0		
Checklist: Data extraction 1C									
Random standard clinical and electronic monitoring	TCI infusions 1.0	Muscle relax. Induc 1,0	Muscle relax. Main 1,0	Spontaneous Resp. 1,0 vs Controlled Resp.					

Appendix 6. Author survey

A survey was sent to the authors of all of the papers included in this review in order to obtain information not stated in the papers and, if possible, to obtain more detailed information regarding the methodology of the included studies. The survey questions were as follows:

1. Can you define the dates during which patients were enrolled and studied?
2. What method of randomization did you use?
3. How could participants and investigators enrolling participants not foresee assignment? Or How did you conceal to what groups patients were randomized to the anaesthetists?
4. Indicate whether or not blinding occurred for patient, anaesthetist, Awareness outcome assessor (interviewer), other, or if no blinding occurred?
5. Describe the completeness of outcome data for each main outcome, including attrition and exclusions from the analysis. State whether attrition and exclusions were reported, the numbers in each intervention group (compared with total randomized participants), reasons for attrition/exclusions where reported, and any re-inclusions in analyses performed by the review authors.
6. The study protocol is available and all of the study's pre-specified (primary and secondary) outcomes that are of interest in the review have been reported in the pre-specified way. The study protocol is not available but it is clear that the published reports include all expected outcomes, including those that were pre-specified (convincing text of this nature may be uncommon).
7. Did you not report any non-significant results from your study?
8. There is at least one important risk of bias? For example, the study had a potential source of bias related to the specific study design used; or stopped early due to some data-dependent process (including a formal-stopping rule); or had extreme baseline imbalance; or has been claimed to have been fraudulent; or had some other problem.
9. In the postoperative period, did investigators interview patients about awareness during surgery with either a standard questionnaire or an informal interview? Did any patients spontaneously report awareness during surgery?
10. After skin incision, was the study performed during all or part of the surgery? If only part, when?

Cochrane ROB policies:

Cochrane Handbook 2008, Table 8.5.C "Either participants or some key study personnel were not blinded, but outcome assessment was blinded and the non-blinding of others unlikely to introduce bias."

Risk of bias survey				
1120				
= 160 * 7				
	High	Low	Unclear	Total
Survey responders N = 54	53	281	44	378
Survey responders N = 54	378	378	378	
Survey responders N = 54	14.0%	74.3%	11.6%	100.0%
<hr/>				
Survey non-responders N = 106	31	286	425	742
Survey non-responders N = 106	742	742	742	
Survey non-responders N = 106	4.2%	38.5%	57.3%	100.0%
				1120

(Continued)

All patients (yes + no survey responders) N = 160	86	564	470	1120
All patients (yes + no survey responders) N = 160	1120	1120	1120	
All patients (yes + no survey responders) N = 160	7.7%	50.4%	42.0%	100.0%
<hr/>				
Meta-analysis N = 18				
126 domains				
= 18*7				
Meta-analysis N = 6 responders	High	Low	Unclear	
Meta-analysis N = 6 responders	9	31	2	42
Meta-analysis N = 6 responders	42	42	42	
Meta-analysis N = 6 responders	21.4%	73.8%	4.8%	100%
<hr/>				
Meta-analysis N = 12 non-responders	High	Low	Unclear	
Meta-analysis N = 12 non-responders	3	37	44	84
Meta-analysis N = 12 non-responders	84	84	84	
Meta-analysis N = 12 non-responders	3.6%	44.0%	52.4%	100%
				126
<hr/>				
All merged patients (yes + no survey responders) N = 18	30	62	34	126
All merged patients (yes + no survey responders) N = 18	126	126	126	
All merged patients (yes + no survey responders) N = 18	24%	49%	27%	

# RCTs merged	Comparison	# domains (# RCTs*7)	High	Low	Unclear	# domains (# RCTs*7)	Evidence downgraded for ROB
	C1		7	42	14	63	
	C1	= 9*7	63	63	63		
9	C1	63	11.1%	66.7%	22.2%	100.0%	NO
	C2	= 5*7	5	27	3	35	
5	C2	35	35	35	35		
			14.3%	77.1%	8.6%		NO
			High	Low	Unclear		
	C3		3	9	9	21	
	C3	= 3*7	21	21	21		
3	C3	21	14.3%	42.9%	42.9%	100.0%	YES
	C 4		0	12	16	28	
	C 4	= 4*7	28	28	28		
4	C 4	28	0.0%	42.9%	57.1%	100.0%	YES

(Continued)

	C 5	= 2*7	1	6	7	12	
2	C 5	14	14	14	14		
			7.1%	42.9%	50.0%	100.0%	YES
18 unique RCTs							

Appendix 7. R Data, Code, Output

The effectiveness of anaesthetic interventions for prevention of wakefulness and awareness during and after surgery

NLP

R Project started October 21, 2014

This analysis run on Fri Jun 17 14:36:23 2016.

checkpoint

setup

snapshot

Analysis 1.1

df.1.1

```
## studynames exp.y exp.n ctl.y ctl.n Year risk
## 1 Avidan 2008 6 967 3 974 2008 High
## 2 Avidan 2011 20 2861 8 2852 2011 High
## 3 Gruenewald 2007 0 37 1 35 2007 Not High
## 4 Kerssens 2009 2 67 1 61 2009 Not High
## 5 Mashour 2012 18 9460 19 9376 2012 Not High
## 6 Mozafari 2014 9 163 7 170 2014 Not High
## 7 Myles 2004 22 1225 27 1238 2004 High
## 8 Puri 2003 0 14 1 16 2003 High
## 9 Zhang 2011 8 2919 21 2309 2011 Not High
```

summary(df.1.1)

```
## studynames exp.y exp.n ctl.y
## Length:9 Min. : 0.000 Min. : 14 Min. : 1.000
## Class :character 1st Qu.: 2.000 1st Qu.: 67 1st Qu.: 1.000
## Mode :character Median : 8.000 Median : 967 Median : 7.000
## Mean : 9.444 Mean :1968 Mean : 9.778
## 3rd Qu.:18.000 3rd Qu.:2861 3rd Qu.:19.000
## Max. :22.000 Max. :9460 Max. :27.000
## ctl.n Year risk
## Min. : 16 Min. :2003 High :4
## 1st Qu.: 61 1st Qu.:2007 Not High:5
## Median : 974 Median :2009
## Mean :1892 Mean :2009
## 3rd Qu.:2309 3rd Qu.:2011
## Max. :9376 Max. :2014
```

```
df.1.1.rma <- rma.uni(ai = exp.y, ci = ctl.y, n1i = exp.n, n2i = ctl.n, measure = "OR",
data = df.1.1, slab = studynames, method = "ML")
summary(df.1.1.rma)
```

```
##
## Random-Effects Model (k = 9; tau^2 estimator: ML)
##
## logLik deviance AIC BIC AICc
## -10.9729 13.1464 25.9458 26.3403 27.9458
##
## tau^2 (estimated amount of total heterogeneity): 0.2416 (SE = 0.2379)
## tau (square root of estimated tau^2 value): 0.4916
## I^2 (total heterogeneity / total variability): 49.04%
## H^2 (total variability / sampling variability): 1.96
##
## Test for Heterogeneity:
## Q(df = 8) = 16.0004, p-val = 0.0424
```

```
##
## Model Results:
##
## estimate se zval pval ci.lb ci.ub
## -0.0220 0.2559 -0.0860 0.9314 -0.5235 0.4795
##
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

predict(df.1.1.rma, transf = exp, digits = 2)

## pred ci.lb ci.ub cr.lb cr.ub
## 0.98 0.59 1.62 0.33 2.90

anova.rma(df.1.1.rma)

##
## Test of Moderators (coefficient(s) 1):
## QM(df = 1) = 0.0074, p-val = 0.9314

# Figure 4 forest(df.1.1.rma, atranf = exp, refline = 0, addcred = T)

Analysis 1.1 subgroups Risk

df.1.1.risk.rma <- rma.uni(ai = exp.y, ci = ctl.y, n1i = exp.n, n2i = ctl.n,
measure = "OR", data = df.1.1, slab = studynames, method = "ML", mods = ~risk,
knha = T)
summary(df.1.1.risk.rma)

##
## Mixed-Effects Model (k = 9; tau^2 estimator: ML)
##
## logLik deviance AIC BIC AICc
## -10.1621 11.5248 26.3242 26.9159 31.1242
##
## tau^2 (estimated amount of residual heterogeneity): 0.1551 (SE = 0.1844)
## tau (square root of estimated tau^2 value): 0.3938
## I^2 (residual heterogeneity / unaccounted variability): 34.75%
## H^2 (unaccounted variability / sampling variability): 1.53
## R^2 (amount of heterogeneity accounted for): 35.81%
##
## Test for Residual Heterogeneity:
## QE(df = 7) = 13.3148, p-val = 0.0648
##
## Test of Moderators (coefficient(s) 2):
## F(df1 = 1, df2 = 7) = 1.7001, p-val = 0.2335
##
## Model Results:
##
## estimate se tval pval ci.lb ci.ub
## intrcpt 0.2942 0.3412 0.8621 0.4172 -0.5127 1.1010
## riskNot High -0.6121 0.4695 -1.3039 0.2335 -1.7222 0.4980
##
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

anova.rma(df.1.1.rma, df.1.1.risk.rma)

## df AIC BIC AICc logLik LRT pval QE tau^2
## Full 3 26.3242 26.9159 31.1242 -10.1621 13.3148 0.1551
## Reduced 2 25.9458 26.3403 27.9458 -10.9729 1.6216 0.2029 16.0004 0.2416
## R^2
## Full
## Reduced 35.81%
```

```
predict(df.1.1.risk.rma, transf = exp, digits = 2, newmods = 0:1)
```

```
## pred ci.lb ci.ub cr.lb cr.ub
## 1 1.34 0.60 3.01 0.39 4.60
## 2 0.73 0.34 1.56 0.22 2.42
```

Analysis 2.1

```
df.2.1
```

```
## studynames exp.y exp.n ctl.y ctl.n Year type risk
## 1 Avidan 2008 6 967 3 974 2008 Alarms High
## 2 Avidan 2011 20 2861 8 2852 2011 Alarms High
## 3 Mashour 2012 18 9460 19 9376 2012 Alarms Not High
## 4 Myles 2004 22 1225 27 1238 2004 TIVA High
## 5 Zhang 2011 8 2919 21 2309 2011 TIVA Not High
```

```
summary(df.2.1)
```

```
## studynames exp.y exp.n ctl.y
## Length:5 Min. : 6.0 Min. : 967 Min. : 3.0
## Class :character 1st Qu.: 8.0 1st Qu.:1225 1st Qu.: 8.0
## Mode :character Median :18.0 Median :2861 Median :19.0
## Mean :14.8 Mean :3486 Mean :15.6
## 3rd Qu.:20.0 3rd Qu.:2919 3rd Qu.:21.0
## Max. :22.0 Max. :9460 Max. :27.0
## ctl.n Year type risk
## Min. : 974 Min. :2004 Alarms:3 High :3
## 1st Qu.:1238 1st Qu.:2008 TIVA :2 Not High:2
## Median :2309 Median :2011
## Mean :3350 Mean :2009
## 3rd Qu.:2852 3rd Qu.:2011
## Max. :9376 Max. :2012
```

```
df.2.1.rma <- rma.uni(ai = exp.y, ci = ctl.y, n1i = exp.n, n2i = ctl.n, measure = "OR",
data = df.2.1, slab = studynames, method = "ML", knha = T)
summary(df.2.1.rma)
```

```
##
## Random-Effects Model (k = 5; tau^2 estimator: ML)
##
## logLik deviance AIC BIC AICc
## -5.5848 10.8575 15.1697 14.3885 21.1697
##
## tau^2 (estimated amount of total heterogeneity): 0.3323 (SE = 0.3151)
## tau (square root of estimated tau^2 value): 0.5764
## I^2 (total heterogeneity / total variability): 68.35%
## H^2 (total variability / sampling variability): 3.16
##
## Test for Heterogeneity:
## Q(df = 4) = 14.3557, p-val = 0.0062
##
## Model Results:
##
## estimate se tval pval ci.lb ci.ub
## -0.0361 0.3645 -0.0992 0.9258 -1.0481 0.9758
##
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
predict(df.2.1.rma, transf = exp, digits = 2)
```

```
## pred ci.lb ci.ub cr.lb cr.ub
## 0.96 0.35 2.65 0.15 6.41
```

```

anova.rma(df.2.1.rma)

##
## Test of Moderators (coefficient(s) 1):
## F(df1 = 1, df2 = 4) = 0.0098, p-val = 0.9258

# Figure 5 forest(df.2.1.rma, atranf = exp, refline = 0, addcred = T)

Analysis 2.1 subgroups TIVA

df.2.1.type.rma <- rma.uni(ai = exp.y, ci = ctl.y, n1i = exp.n, n2i = ctl.n,
measure = "OR", data = df.2.1, slab = studynames, method = "ML", mods = ~type,
knha = T)
summary(df.2.1.type.rma)

##
## Mixed-Effects Model (k = 5; tau^2 estimator: ML)
##
## logLik deviance AIC BIC AICc
## -3.6018 6.8915 13.2037 12.0320 37.2037
##
## tau^2 (estimated amount of residual heterogeneity): 0.0919 (SE = 0.1506)
## tau (square root of estimated tau^2 value): 0.3032
## I^2 (residual heterogeneity / unaccounted variability): 34.90%
## H^2 (unaccounted variability / sampling variability): 1.54
## R^2 (amount of heterogeneity accounted for): 72.33%
##
## Test for Residual Heterogeneity:
## QE(df = 3) = 7.5974, p-val = 0.0551
##
## Test of Moderators (coefficient(s) 2):
## F(df1 = 1, df2 = 3) = 3.3692, p-val = 0.1638
##
## Model Results:
##
## estimate se tval pval ci.lb ci.ub
## intrcpt 0.4127 0.3806 1.0843 0.3576 -0.7985 1.6240
## typeTIVA -1.0129 0.5518 -1.8355 0.1638 -2.7690 0.7432
##
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

anova.rma(df.2.1.rma, df.2.1.type.rma)

## df AIC BIC AICc logLik LRT pval QE tau^2
## Full 3 13.2037 12.0320 37.2037 -3.6018 7.5974 0.0919
## Reduced 2 15.1697 14.3885 21.1697 -5.5848 3.9660 0.0464 14.3557 0.3323
## R^2
## Full
## Reduced 72.33%

predict(df.2.1.type.rma, transf = exp, digits = 2, newmods = 0:1)

## pred ci.lb ci.ub cr.lb cr.ub
## 1 1.51 0.45 5.07 0.32 7.11
## 2 0.55 0.15 1.96 0.11 2.71

Analysis 2.1 subgroups Risk

df.2.1.risk.rma <- rma.uni(ai = exp.y, ci = ctl.y, n1i = exp.n, n2i = ctl.n,
measure = "OR", data = df.2.1, slab = studynames, method = "ML", mods = ~risk,
knha = T)
summary(df.2.1.risk.rma)

##

```

```

## Mixed-Effects Model (k = 5; tau^2 estimator: ML)
##
## logLik deviance AIC BIC AICc
## -4.2051 8.0980 14.4101 13.2385 38.4101
##
## tau^2 (estimated amount of residual heterogeneity): 0.1613 (SE = 0.1993)
## tau (square root of estimated tau^2 value): 0.4016
## I^2 (residual heterogeneity / unaccounted variability): 48.60%
## H^2 (unaccounted variability / sampling variability): 1.95
## R^2 (amount of heterogeneity accounted for): 51.46%
##
## Test for Residual Heterogeneity:
## QE(df = 3) = 9.9456, p-val = 0.0190
##
## Test of Moderators (coefficient(s) 2):
## F(df1 = 1, df2 = 3) = 2.1102, p-val = 0.2423
##
## Model Results:
##
## estimate se tval pval ci.lb ci.ub
## intrcpt 0.3480 0.4202 0.8282 0.4683 -0.9893 1.6853
## riskNot High -0.9211 0.6340 -1.4527 0.2423 -2.9389 1.0968
##
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

anova.rma(df.2.1.rma, df.2.1.risk.rma)

## df AIC BIC AICc logLik LRT pval QE tau^2
## Full 3 14.4101 13.2385 38.4101 -4.2051 9.9456 0.1613
## Reduced 2 15.1697 14.3885 21.1697 -5.5848 2.7595 0.0967 14.3557 0.3323
## R^2
## Full
## Reduced 51.46%

predict(df.2.1.risk.rma, transf = exp, digits = 2, newmods = 0:1)

## pred ci.lb ci.ub cr.lb cr.ub
## 1 1.42 0.37 5.39 0.22 9.01
## 2 0.56 0.12 2.55 0.08 4.08

```

Analysis 2.2

df.2.2

```

## studynames exp.y exp.n ctl.y ctl.n Year type risk
## 1 Avidan 2008 2 967 2 974 2008 Alarms High
## 2 Avidan 2011 8 2861 2 2852 2011 Alarms High
## 3 Mashour 2012 8 9460 11 9376 2012 Alarms Not High
## 4 Myles 2004 2 1225 11 1238 2004 TIVA High
## 5 Zhang 2011 4 2919 15 2309 2011 TIVA Not High

```

summary(df.2.2)

```

## studynames exp.y exp.n ctl.y
## Length:5 Min. :2.0 Min. : 967 Min. : 2.0
## Class :character 1st Qu.:2.0 1st Qu.:1225 1st Qu.: 2.0
## Mode :character Median :4.0 Median :2861 Median :11.0
## Mean :4.8 Mean :3486 Mean : 8.2
## 3rd Qu.:8.0 3rd Qu.:2919 3rd Qu.:11.0
## Max. :8.0 Max. :9460 Max. :15.0
## ctl.n Year type risk
## Min. : 974 Min. :2004 Alarms:3 High :3
## 1st Qu.:1238 1st Qu.:2008 TIVA :2 Not High:2

```



```
## Median :2309 Median :2011
## Mean :3350 Mean :2009
## 3rd Qu.:2852 3rd Qu.:2011
## Max. :9376 Max. :2012

df.2.2.rma <- rma.uni(ai = exp.y, ci = ctl.y, n1i = exp.n, n2i = ctl.n, measure = "OR",
data = df.2.2, slab = studynames, method = "ML", knha = T)
summary(df.2.2.rma)

##
## Random-Effects Model (k = 5; tau^2 estimator: ML)
##
## logLik deviance AIC BIC AICc
## -7.6377 9.7555 19.2753 18.4942 25.2753
##
## tau^2 (estimated amount of total heterogeneity): 0.6684 (SE = 0.7169)
## tau (square root of estimated tau^2 value): 0.8176
## I^2 (total heterogeneity / total variability): 59.88%
## H^2 (total variability / sampling variability): 2.49
##
## Test for Heterogeneity:
## Q(df = 4) = 11.9667, p-val = 0.0176
##
## Model Results:
##
## estimate se tval pval ci.lb ci.ub
## -0.5172 0.5503 -0.9399 0.4005 -2.0451 1.0107
##
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

predict(df.2.2.rma, transf = exp, digits = 2)

## pred ci.lb ci.ub cr.lb cr.ub
## 0.60 0.13 2.75 0.04 9.20

anova.rma(df.2.2.rma)

##
## Test of Moderators (coefficient(s) 1):
## F(df1 = 1, df2 = 4) = 0.8833, p-val = 0.4005

# Figure 6 forest(df.2.2.rma, atranf = exp, refline = 0, addcred = T)
```

Analysis 2.2 subgroups TIVA

```
df.2.2.type.rma <- rma.uni(ai = exp.y, ci = ctl.y, n1i = exp.n, n2i = ctl.n,
measure = "OR", data = df.2.2, slab = studynames, method = "ML", mods = ~type,
knha = T)
summary(df.2.2.type.rma)

##
## Mixed-Effects Model (k = 5; tau^2 estimator: ML)
##
## logLik deviance AIC BIC AICc
## -4.5182 3.5165 15.0364 13.8647 39.0364
##
## tau^2 (estimated amount of residual heterogeneity): 0 (SE = 0.2302)
## tau (square root of estimated tau^2 value): 0
## I^2 (residual heterogeneity / unaccounted variability): 0.00%
## H^2 (unaccounted variability / sampling variability): 1.00
## R^2 (amount of heterogeneity accounted for): 100.00%
##
## Test for Residual Heterogeneity:
```

```
## QE(df = 3) = 3.5165, p-val = 0.3186
##
## Test of Moderators (coefficient(s) 2):
## F(df1 = 1, df2 = 3) = 7.2090, p-val = 0.0747
##
## Model Results:
##
## estimate se tval pval ci.lb ci.ub
## intrcpt 0.0976 0.4028 0.2422 0.8242 -1.1844 1.3796
## typeTIVA -1.7078 0.6361 -2.6850 0.0747 -3.7320 0.3164 .
##
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

anova.rma(df.2.2.rma, df.2.2.type.rma)

## df AIC BIC AICc logLik LRT pval QE tau^2
## Full 3 15.0364 13.8647 39.0364 -4.5182 3.5165 0.0000
## Reduced 2 19.2753 18.4942 25.2753 -7.6377 6.2389 0.0125 11.9667 0.6684
## R^2
## Full
## Reduced 100%

predict(df.2.2.type.rma, transf = exp, digits = 2, newmods = 0:1)

## pred ci.lb ci.ub cr.lb cr.ub
## 1 1.10 0.31 3.97 0.31 3.97
## 2 0.20 0.04 0.96 0.04 0.96
```

Analysis 2.2 subgroupsRisk

```
df.2.2.risk.rma <- rma.uni(ai = exp.y, ci = ctl.y, n1i = exp.n, n2i = ctl.n,
measure = "OR", data = df.2.2, slab = studynames, method = "ML", mods = ~risk,
knha = T)
summary(df.2.2.risk.rma)
```

```
##
## Mixed-Effects Model (k = 5; tau^2 estimator: ML)
##
## logLik deviance AIC BIC AICc
## -7.2858 9.0517 20.5716 19.3999 44.5716
##
## tau^2 (estimated amount of residual heterogeneity): 0.5201 (SE = 0.6159)
## tau (square root of estimated tau^2 value): 0.7212
## I^2 (residual heterogeneity / unaccounted variability): 53.13%
## H^2 (unaccounted variability / sampling variability): 2.13
## R^2 (amount of heterogeneity accounted for): 22.19%
##
## Test for Residual Heterogeneity:
## QE(df = 3) = 10.7061, p-val = 0.0134
##
## Test of Moderators (coefficient(s) 2):
## F(df1 = 1, df2 = 3) = 0.4309, p-val = 0.5584
##
## Model Results:
##
## estimate se tval pval ci.lb ci.ub
## intrcpt -0.1299 0.8451 -0.1537 0.8876 -2.8193 2.5594
## riskNot High -0.7749 1.1805 -0.6564 0.5584 -4.5318 2.9821
##
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

anova.rma(df.2.2.rma, df.2.2.risk.rma)
```

```
## df AIC BIC AICc logLik LRT pval QE tau^2
## Full 3 20.5716 19.3999 44.5716 -7.2858 10.7061 0.5201
## Reduced 2 19.2753 18.4942 25.2753 -7.6377 0.7038 0.4015 11.9667 0.6684
## R^2
## Full
## Reduced 22.19%
```

```
predict(df.2.2.risk.rma, transf = exp, digits = 2, newmods = 0:1)
```

```
## pred ci.lb ci.ub cr.lb cr.ub
## 1 0.88 0.06 12.93 0.03 30.13
## 2 0.40 0.03 5.58 0.01 13.21
```

Analysis 3.1

```
## studlab exp.y exp.n ctl.y ctl.n
## 1 Baraka 1989 4 30 14 20
## 2 Russell 1986 2 30 11 25
## 3 Schultetus 1986 5 23 7 13
```

```
## studlab exp.y exp.n ctl.y
## Length:3 Min. :2.000 Min. :23.00 Min. : 7.00
## Class :character 1st Qu.:3.000 1st Qu.:26.50 1st Qu.: 9.00
## Mode :character Median :4.000 Median :30.00 Median :11.00
## Mean :3.667 Mean :27.67 Mean :10.67
## 3rd Qu.:4.500 3rd Qu.:30.00 3rd Qu.:12.50
## Max. :5.000 Max. :30.00 Max. :14.00
## ctl.n
## Min. :13.00
## 1st Qu.:16.50
## Median :20.00
## Mean :19.33
## 3rd Qu.:22.50
## Max. :25.00
```

```
## study= 1
## study= 2
## study= 3
```

```
## $ci.fixed
## constant inverse-variance fisher asymptotical-MH range
## est 0.19186687 0.17906059 0.15066071 2.286242e-01 NA
## lower CI 0.08790225 0.08790225 0.08790225 1.265683e-01 1e-03
## upper CI 0.40738028 0.41020410 0.43052661 4.129708e-01 1e+03
## p 0.00000000 0.00000000 0.00000000 1.001218e-06 NA
##
## $study.ci
## est lower CI upper CI p limit
## study 1 0.19375713 0.06246219 0.4473941 0.001088307 1
## study 2 0.09804554 0.02744120 0.5452620 0.004248856 1
## study 3 0.39061504 0.13970328 1.0443160 0.067075978 1
##
## $precision
## [1] "+/- 0.00690430312741843"
```

Analysis 3.1 meta regression

```
explicit.df
```

```
## studlab cognitiveState events count logitEstimator logitVariance
## 1 Baraka 1989 wakefulness 18 50 -0.57536.... 0.08680556
## 2 Baraka 1989 awareness 2 50 -3.17805.... 0.52083333
## 3 Russell 1986 wakefulness 13 55 -1.17272.... 0.10073260
## 4 Russell 1986 awareness 1 55 -3.98898.... 1.01851852
```

```
## 5 Schultetus 1986 wakefulness 12 36 -0.69314.... 0.12500000
## 6 Schultetus 1986 awareness 3 36 -2.39789.... 0.36363636
## 7 Tunstall 1989 wakefulness 47 113 -0.33950.... 0.03642811
## 8 Tunstall 1989 awareness 0 113 -5.42495.... 2.00881057
```

```
summary(explicit.df)
```

```
## studlab cognitiveState events count
## Length:8 wakefulness:4 Min. : 0.00 Min. : 36.0
## Class :character awareness :4 1st Qu.: 1.75 1st Qu.: 46.5
## Mode :character Median : 7.50 Median : 52.5
## Mean :12.00 Mean : 63.5
## 3rd Qu.:14.25 3rd Qu.: 69.5
## Max. :47.00 Max. :113.0
## logitEstimator logitVariance
## Min. :-5.4249 Min. :0.03643
## 1st Qu.:-3.3808 1st Qu.:0.09725
## Median :-1.7853 Median :0.24432
## Mean :-2.2213 Mean :0.53260
## 3rd Qu.:-0.6637 3rd Qu.:0.64525
## Max. :-0.3395 Max. :2.00881

##
## Multivariate Meta-Analysis Model (k = 8; method: REML)
##
## Variance Components:
##
## outer factor: studlab (nlvls = 4)
## inner factor: cognitiveState (nlvls = 2)
##
## estim sqrt fixed
## tau^2 0.0612 0.2473 no
## rho -0.9566 no
##
## Test for Residual Heterogeneity:
## QE(df = 6) = 10.0016, p-val = 0.1246
##
## Test of Moderators (coefficient(s) 2):
## F(df1 = 1, df2 = 6) = 26.4014, p-val = 0.0021
##
## Model Results:
##
## estimate se tval pval ci.lb
## intrcpt -0.6536 0.1870 -3.4946 0.0129 -1.1112
## cognitiveStateawareness -2.5263 0.4917 -5.1382 0.0021 -3.7294
## ci.ub
## intrcpt -0.1959 *
## cognitiveStateawareness -1.3232 **
##
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

## pred ci.lb ci.ub cr.lb cr.ub
## 1 0.3422 0.2476 0.4512 0.1959 0.5263
## 2 0.0399 0.0145 0.1053 0.0123 0.1217
```

Analysis 4.1

```
df.4.1
```

```
## studlab exp.y exp.n ctl.y ctl.n
## 1 Ellingson 1977 0 13 4 13
## 2 Haram 1981 0 39 5 40
```

```
## 3 McNulty 1985 0 71 3 25
```

```
## 4 Miller 1996 2 69 4 21
```

```
summary(df.4.1)
```

```
## studlab exp.y exp.n ctl.y
## Length:4 Min. :0.0 Min. :13.0 Min. :3.00
## Class :character 1st Qu.:0.0 1st Qu.:32.5 1st Qu.:3.75
## Mode :character Median :0.0 Median :54.0 Median :4.00
## Mean :0.5 Mean :48.0 Mean :4.00
## 3rd Qu.:0.5 3rd Qu.:69.5 3rd Qu.:4.25
## Max. :2.0 Max. :71.0 Max. :5.00
## ctl.n
## Min. :13.00
## 1st Qu.:19.00
## Median :23.00
## Mean :24.75
## 3rd Qu.:28.75
## Max. :40.00
```

```
exact.4.1 <- meta.exact(data = mat.4.1, type = 'risk ratio')
```

```
## study= 1
## study= 2
## study= 3
## study= 4
```

```
exact.4.1
```

```
## $ci.fixed
## constant inverse-variance fisher asymptotical-MH range
## est 0.20701413 0.16943378 0.12246162 0.1043379866 NA
## lower CI 0.02398833 0.02398833 0.02398833 0.0314276549 1e-03
## upper CI 0.26546056 0.24774221 0.22490546 0.3463960486 1e+03
## p 0.00000000 0.00000000 0.00000000 0.0002228273 NA
##
## $study.ci
## est lower CI upper CI p limit
## study 1 0.0000000 0.0000000 1.1162513 0.03824597 1
## study 2 0.0000000 0.0000000 0.8268940 0.03851101 1
## study 3 0.0000000 0.0000000 0.5636235 0.02942011 1
## study 4 0.1226559 0.0239384 0.8332003 0.02974091 1
##
## $precision
## [1] "+/- 0.00690430312741843"
```

Analysis 5.1

```
df.5.1
```

```
## studlab exp.y exp.n ctl.y ctl.n
## 1 Abboud 1985 0 65 2 16
## 2 Crawford JS 1985 3 384 19 393
```

```
summary(df.5.1)
```

```
## studlab exp.y exp.n ctl.y
## Length:2 Min. :0.00 Min. : 65.0 Min. : 2.00
## Class :character 1st Qu.:0.75 1st Qu.:144.8 1st Qu.: 6.25
## Mode :character Median :1.50 Median :224.5 Median :10.50
## Mean :1.50 Mean :224.5 Mean :10.50
## 3rd Qu.:2.25 3rd Qu.:304.2 3rd Qu.:14.75
## Max. :3.00 Max. :384.0 Max. :19.00
## ctl.n
```

```
## Min. : 16.0
## 1st Qu.:110.2
## Median :204.5
## Mean :204.5
## 3rd Qu.:298.8
## Max. :393.0

exact.5.1 <- meta.exact(data = mat.5.1, type = "risk ratio")

## study= 1
## study= 2

exact.5.1

## $ci.fixed
## constant inverse-variance fisher asymptotical-MH range
## est 0.13121999 0.13121999 0.13121999 0.1431659640 NA
## lower CI 0.03999447 0.03999447 0.03999447 0.0453311444 1e-03
## upper CI 0.43052661 0.42756289 0.42169650 0.4521503599 1e+03
## p 0.00010000 0.00010000 0.00010000 0.0009238782 NA
##
## $study.ci
## est lower CI upper CI p limit
## study 1 0.0000000 0.00000000 0.8020027 0.036672683 1
## study 2 0.1314158 0.03981611 0.5139695 0.001851141 1
##
## $precision
## [1] "+/- 0.00690430312741843"

save.image("Data/implicit.RData")
```

Appendix 8. Adjudication of awareness events

Study	Study size	Number of events	Judged	Judged	Judged
		To be adjudicated	No awareness	Possible awareness	Definite awareness
Avidan 2008	1941	16 (0.82%)	7 (0.36%)	5 (0.26%)	4 (0.21%)
Avidan 2011	5713	50 (0.88%)	22 (0.38%)	18 (0.32%)	10 (0.18%)
Mashour 2012	18,836	76 (0.40%)	39 (0.21%)	18 (0.10%)	19 (0.10%)
Myles 2004	2463	61 (2.48%)	12 (0.49%)	36 (1.46%)	13 (0.53%)
Zhang 2011	5228	300 (5.73%)	271 (5.18%)	10 (0.19%)	19 (0.36%)
Totals	34,181	503 (1.47%)	351 (1.03%)	87 (0.23%)	65 (0.19%)

Study	Number of events	Judged	Judged	Judged
	To be adjudicated	No awareness	Possible awareness	Definite awareness

(Continued)

Avidan 2008	16	7 (43.8%)	5 (31.3%)	4 (25.0%)
Avidan 2011	50	22 (44.0%)	18 (36.0%)	10 (20.0%)
Mashour 2012	76	39 (51.3%)	18 (23.7)	19 (25.0%)
Myles 2004	61	12 (19.7%)	36 (59.0%)	13 (21.3%)
Zhang 2011	300	271 (90.3%)	10 (3.3%)	19 (6.3%)
Totals	503	351 (69.9%)	87 (15.3%)	65 (12.9%)

Details of the adjudication process were tallied from the published study or were sought by communication with the authors.

In the two-stage process, structured interviews of patients were conducted one or more times. Potential awareness episodes were recorded in a narrative report.

An independent committee of three anaesthetists, blinded to treatment group, coded the report as no awareness, possible awareness, and awareness.

- As a percentage of study size, events flagged for adjudication ranged from 0.40% to 5.73%.
- As a percentage of study size, no awareness ranged from 0.21% to 5.18%; possible awareness ranged from 0.10% to 1.46%; definite awareness ranged from 0.10% to 0.53%.

A random-effects meta-analysis of the logit transformed proportions of 'no awareness' events of all adjudicated events was performed in metafor ([Viechtbauer 2010](#)).

Similar models were also estimated for the proportion of potential awareness episodes and definite awareness events.

Average proportions, 95% CIs and 95% prediction intervals (PI) were obtained in the meta-analysis.

- Average proportion of events flagged for adjudication 0.013(95% CI 0.005 to 0.034; 95% PI 0.001 to 0.119).
- Average proportion of adjudicated events judged no awareness 0.53(95% CI 0.25 to 0.79; 95% PI 0.06 to 0.95).
- Average proportion of adjudicated events judged awareness 0.17(95% CI 0.10 to 0.28; 95% PI 0.05 to 0.46).
- Between-study statistical heterogeneity was very high ($I^2=99\%$, 96%, 80%) for events flagged, events adjudicated 'no awareness' and events adjudicated definite awareness.

The [Zhang 2011](#) study had a greater proportion of events adjudicated and events not judged awareness. A sensitivity meta-analysis excluding [Zhang 2011](#) was also estimated.

Averages, 95% CIs and 95% prediction intervals (PI) were obtained in the meta-analysis.

- Average proportion of events flagged for adjudication 0.009(95% CI 0.004 to 0.019; 95% PI 0.002 to 0.047).
- Average proportion of adjudicated events judged no awareness: 0.39(95% CI 0.25 to 0.55; 95% PI 0.14 to 0.71).
- Average proportion of adjudicated events judged awareness 0.22(95% CI 0.17 to 0.29; 95% PI 0.17 to 0.29).
- Between-study statistical heterogeneity remained very high ($I^2=96\%$, 78%) for events flagged and events adjudicated 'no awareness'.
- Between-study statistical heterogeneity was eliminated ($I^2=0\%$) for events adjudicated definite awareness.

Explicit recall is a subjective experience. Corroboration by operating theatre staff that matches specifics of recall with intraoperative events has been reported.

It is very difficult to be certain of the veracity of explicit recall without this corroboration.

Lacking external corroboration, suggestions have been made for improving and revising the adjudication process ([Myles 2015 \(personal communication\)](#)).

- Membership of adjudication panels should be taken from diverse clinical/research disciplines.
- Prerequisites for adjudication panel membership should be established, e.g. years of clinical experience, avoidance of conflicts of interest.

- Establishment of consensus guidelines of the diagnostic criteria including external corroboration if available.
- Workflow standards for the panel should be established, e.g. blinding, clear majority voting, independent ascertainment/categorization.
- Evaluating agreement (e.g. kappa statistic) amongst the review panel members would be worthwhile.

Adjudicate Analysis

NLP

RStudio Project started December 12, 2014

This analysis was run Fri Jun 17 17:10:49 2016.

checkpoint

snapshot

prepareData

List data

```
## slab studySize adjudicated notAwareness definiteAwareness
## 1 Avidan 2008 1941 16 7 4
## 2 Avidan 2011 5713 50 22 10
## 3 Mashour 2012 18836 76 39 19
## 4 Myles 2004 2463 61 12 13
## 5 Zhang 2011 5228 300 271 19
```

prepareMeta

List effect sizes

```
## slab studySize adjudicated notAwareness definiteAwareness
## 1 Avidan 2008 1941 16 7 4
## 2 Avidan 2011 5713 50 22 10
## 3 Mashour 2012 18836 76 39 19
## 4 Myles 2004 2463 61 12 13
## 5 Zhang 2011 5228 300 271 19
## adjudicated.yi adjudicated.vi notAwareness.yi notAwareness.vi
## 1 -4.7901 0.0630 -0.2513 0.2540
## 2 -4.7297 0.0202 -0.2412 0.0812
## 3 -5.5087 0.0132 0.0526 0.0527
## 4 -3.6732 0.0168 -1.4069 0.1037
## 5 -2.7989 0.0035 2.2348 0.0382
## definiteAwareness.yi definiteAwareness.vi
## 1 -1.0986 0.3333
## 2 -1.3863 0.1250
## 3 -1.0986 0.0702
## 4 -1.3063 0.0978
## 5 -2.6939 0.0562
```

meta1

Display meta analysis models. Zhang 2011 data excluded.

```
## [1] "Adjudicated Events Meta Analysis"
##
## Random-Effects Model (k = 4; tau^2 estimator: REML)
##
## tau^2 (estimated amount of total heterogeneity): 0.5679 (SE = 0.4863)
## tau (square root of estimated tau^2 value): 0.7536
## I^2 (total heterogeneity / total variability): 96.39%
## H^2 (total variability / sampling variability): 27.68
##
## Test for Heterogeneity:
## Q(df = 3) = 112.3584, p-val < .0001
##
```



```

## Model Results:
##
## estimate se zval pval ci.lb ci.ub
## -4.6745 0.3859 -12.1142 <.0001 -5.4308 -3.9182 ***
##
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

## [1] "Not Awareness Event Meta Analysis"

##
## Random-Effects Model (k = 4; tau^2 estimator: REML)
##
## tau^2 (estimated amount of total heterogeneity): 0.3433 (SE = 0.3711)
## tau (square root of estimated tau^2 value): 0.5859
## I^2 (total heterogeneity / total variability): 78.04%
## H^2 (total variability / sampling variability): 4.55
##
## Test for Heterogeneity:
## Q(df = 3) = 14.0155, p-val = 0.0029
##
## Model Results:
##
## estimate se zval pval ci.lb ci.ub
## -0.4553 0.3372 -1.3500 0.1770 -1.1163 0.2057
##
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

## [1] "Definite Awareness Event Meta Analysis"

##
## Random-Effects Model (k = 4; tau^2 estimator: REML)
##
## tau^2 (estimated amount of total heterogeneity): 0 (SE = 0.0909)
## tau (square root of estimated tau^2 value): 0
## I^2 (total heterogeneity / total variability): 0.00%
## H^2 (total variability / sampling variability): 1.00
##
## Test for Heterogeneity:
## Q(df = 3) = 0.5511, p-val = 0.9075
##
## Model Results:
##
## estimate se zval pval ci.lb ci.ub
## -1.2233 0.1679 -7.2868 <.0001 -1.5524 -0.8943 ***
##
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

predict1

List prediction intervals. Zhang 2011 data excluded.

```

## [1] "Adjudicated %"

## pred ci.lb ci.ub cr.lb cr.ub
## 0.0092 0.0044 0.0195 0.0018 0.0467

## [1] "Not Awareness Event %"

## pred ci.lb ci.ub cr.lb cr.ub
## 0.3881 0.2467 0.5512 0.1443 0.7047

## [1] "Definite Awareness Event %"

```

```
## pred ci.lb ci.ub cr.lb cr.ub
## 0.2273 0.1747 0.2902 0.1747 0.2902
```

meta2

Display meta analysis models. Zhang 2011 data included.

```
## [1] "Adjudicated Events Meta Analysis"
```

```
##
## Random-Effects Model (k = 5; tau^2 estimator: REML)
##
## tau^2 (estimated amount of total heterogeneity): 1.1330 (SE = 0.8174)
## tau (square root of estimated tau^2 value): 1.0644
## I^2 (total heterogeneity / total variability): 98.81%
## H^2 (total variability / sampling variability): 84.05
##
## Test for Heterogeneity:
## Q(df = 4) = 543.2804, p-val < .0001
##
## Model Results:
##
## estimate se zval pval ci.lb ci.ub
## -4.2933 0.4808 -8.9290 <.0001 -5.2357 -3.3509 ***
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
## [1] "Not Awareness Event Meta Analysis"
```

```
##
## Random-Effects Model (k = 5; tau^2 estimator: REML)
##
## tau^2 (estimated amount of total heterogeneity): 1.7454 (SE = 1.3066)
## tau (square root of estimated tau^2 value): 1.3211
## I^2 (total heterogeneity / total variability): 95.83%
## H^2 (total variability / sampling variability): 23.99
##
## Test for Heterogeneity:
## Q(df = 4) = 126.0925, p-val < .0001
##
## Model Results:
##
## estimate se zval pval ci.lb ci.ub
## 0.0975 0.6080 0.1603 0.8726 -1.0942 1.2891
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
## [1] "Definite Awareness Event Meta Analysis"
```

```
##
## Random-Effects Model (k = 5; tau^2 estimator: REML)
##
## tau^2 (estimated amount of total heterogeneity): 0.4153 (SE = 0.3781)
## tau (square root of estimated tau^2 value): 0.6444
## I^2 (total heterogeneity / total variability): 80.67%
## H^2 (total variability / sampling variability): 5.17
##
## Test for Heterogeneity:
## Q(df = 4) = 26.1814, p-val < .0001
##
## Model Results:
##
```

```
## estimate se zval pval ci.lb ci.ub
## -1.5624 0.3276 -4.7688 <.0001 -2.2045 -0.9202 ***
##
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

predict2

List prediction intervals. Zhang 2011 data included.

```
## [1] "Adjudicated %"
```

```
## pred ci.lb ci.ub cr.lb cr.ub
## 0.0135 0.0053 0.0339 0.0014 0.1188
```

```
## [1] "Not Awareness Event %"
```

```
## pred ci.lb ci.ub cr.lb cr.ub
## 0.5244 0.2508 0.7840 0.0599 0.9502
```

```
## [1] "Definite Awareness Event %"
```

```
## pred ci.lb ci.ub cr.lb cr.ub
## 0.1733 0.0993 0.2849 0.0484 0.4637
```

CONTRIBUTIONS OF AUTHORS

Anthony G Messina (AGM), Michael Wang (MaW), Marshall J Ward (MJW), Chase C Wilker (CCW), Brett B Smith (BS), Daniel P Vezina (DV), Nathan Leon Pace (NLP)

Conceiving the review: AGM

Co-ordinating the review: AGM

Undertaking manual searches: AGM, NLP, MaW, CCW, BS

Screening search results: AGM, NLP, PMaW, CCW, BS

Organizing retrieval of papers: AGM, NLP, MaW, CCW, BS

Screening retrieved papers against inclusion criteria: AGM, NLP, MaW, CCW, BS

Appraising quality of papers: AGM, NLP, MaW, BS, CCW

Abstracting data from papers: AGM, NLP, MaW, BS, CCW

Writing to authors of papers for additional information: AGM, NLP, MaW, CCW, BS

Providing additional data about papers: AGM, NLP, MaW, MJW, DV

Obtaining and screening data on unpublished studies: AGM, NLP, CCW

Data management for the review: AGM, NLP, MaW, CCW

Entering data into Review Manager (RevMan): NLP, AGM, MaW, CCW

RevMan statistical data: NLP, AGM, CCW, DV

Other statistical analysis not using RevMan: NLP, AGM, CCW, DV

Double entry of data: AGM, NLP, MaW, CCW

Interpretation of data: NLP, AGM, MaW, CCW, MJW, DV

Statistical inferences: NLP, AGM, NP, CCW, MJW, DV

Writing the review: AGM, NLP, MaW, MJW, CCW, NP, DV

Securing funding for the review: not applicable

Performing previous work that was the foundation of the present study: AGM, NLP

Guarantor for the review (one author): AGM

Person responsible for reading and checking review before submission: AGM

DECLARATIONS OF INTEREST

Anthony G Messina: none known.

Michael Wang was involved in the design, conduct, or publication of potentially eligible studies for this Cochrane review ([Russell 1997](#); [Russell 2001](#)). Prof Wang has received funding from Abbvie for invited lectures and meeting attendance, including accidental anaesthetic awareness events. He has in the past been loaned two Bispectral Index monitors for research purposes by Aspect Medical/Covidian.

Marshall J Ward: none known.

Chase C Wilker: none known.

Brett B Smith: none known.

Daniel P Vezina: board member and executive position since January 2012 with a Big Data analytics private company (MedAnalytics), and a congestive heart failure clinical solution private company since June 2008 (Guardsman Scientific).

Nathan Leon Pace: none known.

SOURCES OF SUPPORT

Internal sources

- None, Other.
- University of Utah, USA.

Institutional salary

- University of Leicester, UK.

Institutional salary

- Veterans Administration, USA.

Institutional salary

External sources

- No sources of support supplied

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

Background:

This has been updated since the protocol ([Messina 2008](#)), with new sections in RevMan. We have added new paragraphs in the Dryad appendix about the psychological sequelae of cases of anaesthetic awareness, complications from muscle relaxant use, intraoperative dreams, onset of explicit memory in the infant, lack of intraoperative wakefulness, post-traumatic stress disorder, and delayed memory amongst other topics that were not in the protocol ([Appendix 1](#)).

Methods section:

More authors were added to the review. We updated many of the methods sections. Assessment of heterogeneity was interpreted in the context of recently published criteria for the assessment of inconsistency ([Guyatt 2011](#)). Statistical meta-analyses were done in the R language using the exactmeta and metafor packages ([Tian 2008](#); [Viechtbauer 2010](#)).

The web app Rayyan was used to assist in citation review ([Elmagarmid 2014](#)).

Outcomes:

We have changed the sensitivity analysis mentioned in the protocol from risk of bias domains to subgroup analyses based on clinical characteristics.

We added an analysis of the adjudication process for patient awareness reports by expert panels in [Analysis 1.1](#), [Analysis 2.1](#), and [Analysis 2.2](#).

We have changed the secondary outcomes in the review:

Secondary outcomes

- Signs or adverse effects of intraoperative wakefulness and/or postoperative awareness are intraoperative patient movement, haemodynamic changes, portions of intraoperative dreams and postoperative dreams and/or nightmares, delayed memory, full (PTSD) or partial (PTS) forms of post-traumatic stress syndrome, myocardial infarction, cardiac arrest, death, and suicide.

Search methods for identification of studies:

The search strategies have been revised and expanded.

INDEX TERMS**Medical Subject Headings (MeSH)**

*Anesthesia; *Anesthetics; Benzodiazepines [administration & dosage]; Consciousness Monitors; Electroencephalography [methods]; Etomidate; Intraoperative Awareness [*prevention & control]; Ketamine; Mental Recall; Monitoring, Intraoperative [instrumentation] [*methods]; Randomized Controlled Trials as Topic; Thiopental [administration & dosage]; Wakefulness

MeSH check words

Humans