



Published in final edited form as:

Nat Rev Chem. 2019 January ; 3(1): 15–34. doi:10.1038/s41570-018-0059-x.

Modern applications of low-valent early transition metals in synthesis and catalysis

Evan P. Beaumier[#], Adam J. Pearce[#], Xin Yi See[#], and Ian A. Tonks^{*}

Department of Chemistry, University of Minnesota, Minneapolis, MN, USA.

[#] These authors contributed equally to this work.

Abstract

Low-valent early transition metals are often intrinsically highly reactive as a result of their strong propensity toward oxidation to more stable high-valent states. Harnessing these highly reducing complexes for productive reactivity is potentially powerful for C-C bond construction, organic reductions, small-molecule activation and many other reactions that offer orthogonal chemoselectivity and/or regioselectivity patterns to processes promoted by late transition metals. Recent years have seen many exciting new applications of low-valent metals through building new catalytic and/or multicomponent reaction manifolds out of classical reactivity patterns. In this Review, we survey new methods that employ early transition metals and invoke low-valent precursors or intermediates in order to identify common themes and strategies in synthesis and catalysis.

Early transition metals are excellent candidates for the design and implementation of new catalytic reactions: they are earth abundant, often highly reactive and frequently exhibit different structures and orthogonal reactivity compared with late transition metals^{1,2}. However, owing to their electropositive, oxophilic³ nature, the organometallic chemistry of early transition metals has historically been dominated by complexes in high-valent states (FIG. 1). By contrast, there are considerably fewer examples of the application of low-valent early transition metals: such complexes are often highly reducing and require strong stabilizing ligands, such as strong π -acceptors or bulky cyclopentadienyls.

Nonetheless, there is great interest in harnessing the highly reactive nature of low-valent early transition metals for practical synthesis and catalysis. There are myriad elegant examples of stoichiometric N₂ activation^{4–17}, reductive coupling^{18–22} and radical reactions^{23–25} mediated by low-valent early transition metals. The majority of this reactivity is dependent on the fact that highly reduced early metals can activate otherwise unreactive

^{*} itonks@umn.edu.

Author contributions E.P.B., A.J.P. and X.Y.S. researched data for the article. All authors made substantial contributions to discussion of content, writing, reviewing and editing of the manuscript before submission.

Competing interests

The authors declare no competing interests.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

molecules, wherein metal oxidation and the formation of strong metal-ligand bonds provide a large thermodynamic driving force.

Although many seminal contributions to the field were carried out decades ago, there has been a renaissance in the field over the past 15 years enabled by new ligand designs, reagents and approaches. Early transition metals are well positioned to have considerable impact in a wide array of fields, from small-molecule activation and energy storage to photoredox catalysis and other new organic methods. This Review covers new applications of low-valent early transition metals²⁶ (classified here as groups 3, 4 and 5, excluding V because it does not predominantly exist in its highest oxidation state) in synthesis and catalysis. In particular, this Review highlights how the unique properties of low-valent metals can access new mechanisms and enable challenging chemical transformations inaccessible by their late transition metal counterparts.

Generation of low-valent complexes

The most common routes for generating low-valent early transition metal complexes involve the reduction of metal halides with strong alkali metals^{27–32}; reduction with other (often weaker) reducing metals, such as Mg (REFS^{33,34}), Mn (REF.³⁵), Al (REFS^{36,37}) or Zn (REF.³⁸) (providing more functional group tolerance); or p-H abstraction^{39–43} from a metal alkyl (Fig. 2a). Many of these reductions can be performed in situ to directly generate reactive low-valent intermediates or in the presence of π -accepting ligands to generate isolable or transiently stable ‘masked’ species that can liberate a reactive low-valent metal fragment upon loss of the re-accepting ligand. For example, the Negishi reagent, Cp₂Zr(η^2 -butene) (**1**), is synthesized by the treatment of Cp₂ZrCl₂ with ⁿBuLi or ⁿBuMgBr, wherein the Cp₂ZrⁿBu₂ intermediate undergoes β -H abstraction to liberate one equivalent of butane along with the product^{44–46}. Cp₂Ti(TMS₂C₂) (**2**) (TMS₂C₂ = bistrimethylsilylacetylene) and related Ti and Zr analogues can be synthesized by reduction of the corresponding high-valent chlorides with Mg in the presence of the π -accepting alkyne TMS₂C₂ (REF⁴⁷), while substituted hafnocene analogues have been synthesized using Li as a reductant^{48,49}.

Stable low-valent coordination complexes of the early transition metals that lack strongly stabilizing ligands are uncommon. However, examples continue to emerge that challenge the limits of stability (FIG. 2b). For example, a suite of divalent group 3 amides has recently been reported^{50–53}, including the first crystallographically characterized Sc^{II} compound⁵², extending the chemistry of low-valent Sc and Y beyond metallocene derivatives^{54,55}. The Sc^{II} amide complex **3** exhibits remarkably different reactivity from other reduced group 3 and lanthanide complexes, for the first time revealing end-on, reversible binding of N₂ to form the bridged bimetallic **4** (REF⁵³). Bulky aromatic heterocycles derived from phosphalkynes, which are much poorer donors than Cp-based ligands, can also stabilize Sc^{II} complexes⁵⁶. An improved synthesis and full electronic characterization of py₄TiCl₂ (**5**)⁵⁷, a rare example of a Ti^{II} coordination complex, have recently been reported, building upon earlier studies by several groups^{58,59}. The chemistry of low-valent early transition metal isocyanide^{60–62} and arene complexes^{63,64} has also continued to expand, including the recent synthesis of the formally Ta⁰ complex **6** (REF⁶⁰).

The high reactivity of strong alkali metal reductants places an inherent restriction on the solvent and reagent compatibility of reactions that employ them as reductants. Considerable progress has been made in designing new reactions and reductants that overcome this challenge. One creative workaround that was recently introduced is to reduce compounds by passing them through columns of KC_8 , limiting (or at least fine tuning) the contact time of the reagent and solvent with the strong reductant⁶⁵. Another strategy is to use inner-sphere reductants that are less prone to over-reduction. In this context, Tsurugi and Mashima have promoted the use of organosilicon reductants, such as 3,6-bis(trimethylsilyl)-1,4-cyclohexadiene^{66–68} (**7**) and 1,4-bis(trimethylsilyl)-1,4-dihydropyrazine^{69–71} derivatives (for example, **8**), to reduce group 4 and group 5 metal halides (Fig. 2c) for homogeneous catalytic applications. In the case of **8**, the steric bulk of the tetramethylpyrazine by-product prevents its coordination to the reduced metal catalyst, which is critical for productive chemistry. These organosilicon reductants derive their reduction potential (ranging from +0.83 V to –0.40 V versus Cp_2Fe) from the aromatization of the 1,4-diene core. These reductants have been used by the broader community for wide-ranging applications, from the in situ reduction of Ti precursors in chemical vapour deposition⁷² to the in situ generation of Ti^{III} for reductive umpolung reactions⁷³. Similarly, commercially available tris(trimethylsilyl)silane has been used to generate Nb^{III} cyclotrimerization catalysts in situ⁷⁴. A major advantage of these silicon-based reductants is that the by-products are simple to remove: Me_3SiCl is less likely than alkali metal halides to coordinate to electrophilic metals, and the aromatic by-products can be removed in vacuo. Metal salt by-products are often difficult and/or tedious to remove and may interfere with productive chemistry of the reduced complex; thus, these procedures offer an important advancement for applications of reduced early metal chemistry.

Alternative reduction methods of early transition metals include single-electron transfers from weak reductants such as citrate⁷⁵ or benzylamine⁷⁶. Hydrogenolysis of metal alkyls has also been used to synthesize low-valent Nb (REF⁷⁷) and Zr (REF⁷⁸) arene complexes. Redox non-innocent ligands can also engender redox reactivity with early transition metals without formally changing the metal oxidation state^{67,68,79–82}.

Reductive coupling reactions

Reductive coupling of unsaturated organics by d^2 early transition metals has been exploited for a vast array of synthetically useful C-C bond forming reactions. Typically, these reactions involve activation of an alkene or an alkyne to form either a metallacyclopropane or metallacyclopentene, respectively. These species can be considered as 1,2-dianion equivalents and can be intercepted sequentially by two different electrophiles. Often, the resulting metallacycles can be further functionalized, forming the basis for a diverse set of coupling reactions, such as the Negishi^{21,22,45}, Fagan-Nugent^{19,83–85}, Kulinkovich^{43,86–88}, Pauson-Khand^{89–94} and several other stoichiometric coupling reactions of alkynes, alkenes, alkyl halides and carbonyl compounds^{18,20}. Although most d^2 early metals exhibit this type of reactivity, low-valent Ti and Zr complexes — in particular, Cp_2M^0 ($M = Ti, Zr$) and $(O^iPr)_2Ti^{II}$ fragments — have seen the most development in this field, and as such the use of such reagents in the total synthesis of natural products has flourished^{23,95–97}.

Ti(OⁱPr)₄/RM-mediated reductive coupling reactions.

The discovery of the main group metal alkyl reduction of early metals⁴⁵ inspired many reductive coupling reactions using titanium. In 1989, Kulinkovich reported the Ti(OⁱPr)₄-catalysed cyclopropanation of esters with Grignard reagents to form cyclopropanols⁸⁶ (FIG. 3a). Shortly after, the Kulinkovich reaction was translated to amides (de Meijere modification), which results in the formation of cyclopropylamines by a similar mechanism⁹⁸. Since these initial reports, several improvements have been made to this chemistry. Intermolecular alkene couplings can be achieved by using cyclohexyl or cyclopentylmagnesium chloride to activate Ti(OⁱPr)₄, as the resulting Ti-cyclopentene (or cyclohexene) adduct **9** can be displaced by monosubstituted or tethered alkenes to make a new alkene adduct **10** (REFS^{99, 100}) (FIG. 3a, bottom). This discovery has helped dramatically expand the scope of Kulinkovich reactions, as alkene starting materials are more widely available than Grignard reagents. The Kulinkovich cyclopropanation has an intrinsic diastereoselectivity for the cis-1,2-dialkylated cyclopropanol when using Grignard reagents larger than ethyl, and the use of chiral TADDOL (a,a,a',a'-tetraaryl-2,2-disubstituted-1,3-dioxolane-4,5-dimethanol) ligands has allowed for good enantioselectivity^{88,101,102}. This diastereoselectivity can be reversed when cyclopropanating homoallylic alcohols as a result of coordinating directing group effects⁹⁹ or remote stereocentre influences¹⁰³.

A recent modification of the Kulinkovich-de Meijere reaction allows for the synthesis of carbocyclic amino ketones mediated by Ti(OⁱPr)₄/RMgBr (REF¹⁰⁴) (FIG. 3b, top). By using allyl-substituted lactams such as **11**, the cyclopropanation step of the de Meijere mechanism can be arrested, as the bridgehead nitrogen atom of the alkene-inserted intermediate **12** cannot π -bond with the neighbouring carbon (Bredt's rule)¹⁰⁵. This ultimately leads to cyclic aminoketone **13** upon protic workup. Increasing the ring size of the lactam relieves ring strain in the intermediate **12**; thus, larger ring systems are no longer affected by Bredt's rule and typical de Meijere cyclopropylamine generation proceeds. Similarly, interrupted Kulinkovich-de Meijere reactions yield hydroxyl-substituted lactams (**16**) with homoallylic imide (**14**) substrates^{106–108}, which do not undergo cyclopropanation owing to resonance stabilization of the metallacyclic intermediate **15** (FIG. 3b, bottom).

Micalizio has shown that regioselectivity and stereoselectivity in the reductive coupling of various unsaturated compounds can be induced through the use of alkoxide directing groups (FIG. 3c) — a strategy similar to that employed by Sharpless for asymmetric alkene epoxidation¹⁰⁹. This is an advancement from the limited selectivity obtained through Fagan-Nugent chemistry: substrate-directed metallacycle-mediated cross-couplings, although requiring prefunctionalization with potentially undesirable functional groups, allow for the regiospecific assembly of target molecules. By contrast, undirected cross-couplings of the same type typically give statistical mixtures in unbiased systems^{110,111}.

Regioselective and stereoselective coupling of alkenes with imines is possible when using Ti(OⁱPr)₄ⁿBuLi as opposed to Ti(OⁱPr)₄/RMgBr (REFS^{112,113}). Using BuLi as the metal alkyl reductant rather than a Grignard reagent grants dramatically enhanced stability to the titanacyclopropane intermediate as a result of the interaction of LIOⁱPr with the metal centre,

forming a tentative ‘ate’ complex¹¹³. Homoallylic alcohols cross-couple with aromatic imines to give 1,5-aminoalcohols using $\text{Ti}(\text{O}^i\text{Pr})_4/\text{RMgBr}$ (REF¹⁰⁹), but the employment of $\text{Ti}(\text{O}^i\text{Pr})_4/\text{BuLi}$ allows for both aromatic and aliphatic imines to be used successfully in the reaction¹¹⁴. Imines (**17**) couple with allylic alcohols (**18**) to form stereodefined homoallylic amines (**19**)¹¹⁵ (FIG. 3c, top). Alkene stereoselectivity is achieved in these reactions through coordination of the deprotonated alcohol to the preformed Ti-imine complex, which undergoes syn-carbometallation (**20**) and syn-elimination through boat-like transition states (**21** and **22**) (FIG. 3c, top right). Primary homoallylic amines are accessible through similar coupling reactions with aldehydes in the presence of LiHMDS (lithium hexamethyldisilazide), which forms the TMS-imine complex in situ. These primary homoallylic amines have been applied as starting materials for the synthesis of complex bicycles¹¹⁶ and pyridines¹¹⁷.

Substrate-directed selective alkene-alkyne-alkyne formal [2+2+2] reactions are also possible. For example, alkoxide-directed annulations can give hydroindane derivatives (**26** and **27**) and angularly substituted decalins^{118–122} (FIG. 3c, bottom). The reactions proceed through alkoxide-directed insertion of an alkyne into the titanacyclopropene **23** to make the bicyclic metallacyclopentadiene **24**. The chemoselectivity of this insertion is driven by the electronic bias of the $\alpha\text{-SiMe}_3$ group: the Ti-C(TMS) bond is more difficult to insert into than the Ti-C(Ph) bond of **23**. Next, [4+2] cycloaddition of the pendent alkene forms a bridged bicyclic metallacyclopentene **25**, which can undergo allylic isomerization and be quenched through protonation with allylic transposition (**26**) or oxidation (**27**). The choice of alcohol for the quench affects the selectivity for either cis-hydroindanes or trans-hydroindanes, although the origin of this effect remains unclear¹¹⁹.

Other early-transition-metal-mediated reductive coupling reactions.

Metallacycle **28**, formed by reducing Cp^*TiCl_3 ($\text{Cp}^* = \text{C}_5\text{Me}_5^-$) in the presence of dimethylbutadiene, can react with two equivalents of isocyanide to give an asymmetric metal-bound bis(imine) **29** (REF¹²³) (FIG. 4a). Furthermore, upon heating **29** at 55°C in pyridine, cyclopentenimine **30** is extruded, and $\text{Cp}^*\text{TiCl}(\text{py})(\text{N}^i\text{Bu})$ is formed. The result is a unique formal [4+1] coupling reaction alongside cleavage of an isocyanide triple bond, such that the C atom is inserted into both C-H bonds at the 2 position to give an exo-methylene. By contrast, the reaction with the hafnium analogue of **28** results in **31**, which was initially misidentified as a cyclic amidine complex¹²⁴ but instead is the result of imine C-C coupling from the Hf analogue of **29**. The Cp^*_2Zr analogue can selectively insert one equivalent of isocyanide (or CO) into the metal-butadiene bond, allowing for sequential reactions to be carried out¹²⁵.

$\text{Cp}^R_2\text{M}(\text{TMS}_2\text{C}_2)$ ($\text{M} = \text{Ti}, \text{Zr}$; $\text{R} = \text{H}, \text{alkyl}$) can act as a masked source of $\text{Cp}^R_2\text{M}^{\text{II}}$, easily liberating TMS_2C_2 upon treatment with a variety of reagents⁴⁷. Recently, Rosenthal has discovered that $\text{Cp}^*_2\text{M}^{\text{II}}$ fragments can couple nitriles to form metalla-2,5-diazacyclopentadienes, which can undergo further reaction and exchange one nitrile partner for another unsaturated substrate (for example, a different nitrile, CO_2 or H_2)^{126–128}. Nitrile-nitrile couplings are quite uncommon, although they have been observed with Ti^{III} complexes much earlier¹²⁹. Taking advantage of this reaction manifold, Tilley has recently

used $\text{Cp}_2\text{Ti}(\text{TMS}_2\text{C}_2)$ for the coupling of tethered dinitriles, such as **32**, to diazametallacyclopentadienes, such as **33** (REF¹⁰⁰). Once coupled, **33** can be reacted further with electron-deficient alkynes, H_3O^+ or ECl_2 ($\text{E} = \text{SO}, \text{POPh}, \text{S}, \text{Se}, \text{Te}$) to generate pyrazines (**34**), diones (**35**) or diazoles (**36**) (FIG. 4b). Although this work focused primarily on installing these moieties in polycyclic aromatic hydrocarbons, [2+2+2] pyrazine synthesis at an early transition metal had not been previously observed, providing new insight on routes to incorporate nitriles more effectively into reductive coupling reactions mediated by early transition metals. In a similar vein, stoichiometric $\text{Cp}_2\text{Zr}(\text{TMS}_2\text{C}_2)$ has been used as a general platform for C-C coupling and can be used to couple tethered diynes to make polycyclic aromatic hydrocarbons^{130–133}. $\text{Cp}_2\text{M}^{\text{II}}$ fragments can also be used to construct metallacyclocumulenes and other highly strained metallacycles^{47,49,134–142}. Here, the steric bulk of the Cp^* ligands imbue stability, which enables easy isolation of the metal complexes that feature exotic bonding motifs. Structural characterization of the products of reductive metallacyclization of 1,3-butadiynes reveals that the rich chemistry is strongly dependent on the terminal diyne substituents. Similarly, Beckhaus has coupled unsaturated organics with group 4 metal pentafulvene complexes^{143–148}, presenting an interesting approach to the synthesis of exotic appended Cp-type ligands. Likewise, fulvenes can be homocoupled to give ansa-metallocene complexes from group 4 metal chlorides and $^n\text{BuLi}$ (REFS^{149,150}).

There are fewer examples of group 5 metals facilitating reductive coupling reactions. Alkyne-stabilized masked Ta^{III} hydrides can stoichiometrically insert isonitriles and alkynes into the Ta-H bond, followed by C-C reductive coupling of the bound alkyne to give butadienyl and azadiene fragments¹⁵¹. Dimethyltantalum corrole complexes can react with CO to give coupled diacetyl complexes¹⁵². This process is presumed to occur via insertion of two CO molecules into the two Ta-Me bonds, followed by reductive elimination of the two acyl ligands to give diacetyl, which then binds to the resulting low-valent tantalum as the ene-diolate form to formally oxidize Ta^{III} to Ta^{V} .

Catalytic alkyne cyclotrimerization is also common with low-valent group 4–5 transition metals^{34,110,153}. There are two limiting mechanisms for cyclotrimerization: triple insertion followed by reductive elimination through a metallacycloheptatriene (**37**) or [4+2] cycloaddition of the 3rd unsaturated partner to make a metallanorbornadiene (**38**) (FIG. 4c, left). Recent mechanistic work on a ditantalum system indicates that the [4+2] pathway is more likely¹⁵⁴. Remarkably, alkyne cyclotrimerization reactions can be intercepted with other unsaturated substrates; for example, Obora has shown that thermally stable low-valent Nb catalysts can chemoselectively insert alkenes^{155–157} and nitriles¹⁵⁸ to give cross-coupled products¹⁵⁹ (FIG. 4c, right).

Functionalization and defunctionalization

Hydrogenation.

Catalytic hydrogenation of unsaturated substrates is challenging with d^2 early transition metals because of their propensity to promote reductive coupling. Additionally, there are very few examples of direct H_2 activation by d^2 group 5 metals^{160–166}. Nonetheless, there are several elegant examples of arene and alkene hydrogenation and alkyne semi-hydrogenation catalysed by d^2 group 5 metals^{162,167,178}.

β -Diketiminato Nb^{III} imido complexes promote remarkably selective semi-hydrogenation of alkynes¹⁷⁶ (FIG. 5a, top). Treatment of (BDI)Nb(N'Bu)(CO)₂ (**39**) (BDI = 2,6-diisopropylbenzene-p-diketiminato) with 1-phenyl-1-propyne yields the metallacyclopropene complex **33**, which does not undergo reductive coupling owing to steric encumbrance around the metal centre¹⁷⁹. Consequently, it was found that **40** can catalyse the semi-hydrogenation of phenylpropyne to Z-p-methylstyrene under H₂/CO. Two mechanistic proposals were made (FIG. 5a, right): through alkyne-CO ligand exchange, followed by σ -bond metathesis of the metallacyclopropene of **40** with H₂ to yield a Nb^V vinyl hydride (**41**), from which C-H reductive elimination then liberates Z- β -methylstyrene and regenerates an arene-bound Nb^{III} (**42**) (path A), or through 1,2-H₂ addition across the imido of **40** to make a Nb amido hydride (**43**), which can then undergo reductive H migration to vinyl amide **44** followed by α -elimination to **42** (path B). The rate and selectivity for semi-hydrogenation are highly CO-dependent, which may provide a tuneable handle in future systems. Higher CO concentration maintained selectivity but slowed the reaction, indicating that CO-alkyne exchange is kinetically relevant. However, lower loadings led to the formation of more fully hydrogenated propylbenzene, indicating that CO also plays a role in efficient product displacement.

Fortier recently described the synthesis of a new masked Ti^{II} synthon, **45** (FIG. 5a, bottom). This complex undergoes a relatively rare example of early transition metal d² C-H oxidative addition^{160,180,194} followed by β -H abstraction to extrude H₂ and make a new cyclometallated complex **46** (REF¹⁹⁵). Remarkably, this reaction is reversible, opening up the possibility that these complexes might be useful as hydrogenation catalysts. In fact, **45** also performs a unique example of early-transition-metal-promoted transfer hydrogenation¹⁹⁶ (FIG. 5a, bottom left), hydrogenating cyclohexene with concomitant production of **46**, with the ligand serving as the terminal hydrogen donor. This interesting system can also catalytically hydrogenate cyclic alkenes and cyclic dienes, although the mechanism by which this occurs — transfer hydrogenation or a more classical dihydride mechanism — remains unknown.

Dehydrogenation.

There are several examples of C-H oxidative addition to low-valent early transition metals^{160,180–194,197,198}, including several that lead to stoichiometric dehydrogenation^{188–190}. The majority of dehydrogenation reactions promoted by early transition metals are, however, predicated on high-valent, electrophilic metal reactivity, such as 1,2-C-H addition across metal-ligand multiple bonds^{199–201}. The first catalytic, selective dehydrogenation of alkanes was recently demonstrated with Ti catalyst **47** [REF.²⁰²] [FIG. 5b]. This system takes advantage of 1,2-C-H addition of the alkane C-H bond to a Ti alkylidyne **50**, followed by β -H abstraction of the resultant Ti-alkyl-alkylidene (**51**) to formally reduce Ti, making a new Ti^I alkyl-alkene adduct **52** (REFS^{40–41–203–204}). Remarkably, this system is rendered catalytic by addition of a methylene transfer agent, phenyldibenzophosphole methylene ylide, H₂CP(C₁₂H₈)Ph (**48**), which serves as an oxidant to reoxidize the Ti^{II} fragment (**53**) back to a Ti^{IV} alkylidene (**54**)²⁰². The structure of this reagent is critical, as deleterious phenyl migration from a Ti^{II}-ylide adduct will occur with other simple arylphosphorus ylides²⁰⁴. Although turnover numbers in this system are poor

and phosphorus ylides are impractical terminal oxidants, this system is remarkable in its selectivity, yielding only terminal alkene products. This selectivity results from the unique mechanism, which — unlike late transition metal alkane dehydrogenation catalysts — does not involve metal hydride species, which could isomerize the alkene.

There are many low-valent Ti and Zr complexes capable of the catalytic dehydrogenation of amine boranes, particularly using metallocene derivatives^{205–210}. In a recent elegant application, a Cp*₂TiCl₂^mBuLi catalytic system has been shown to promote an unusual step-growth dehydropolymerization of MeNH₂(BH₃), enabling the first synthesis of high-molar-mass poly(N-methylaminoborane) by an early transition metal²¹¹. This system was also successful in the catalytic dehydropolymerization of several other primary amine borane derivatives. Despite computational evidence for a M^{II}/M^{IV} redox mechanism for amine borane dehydrogenation²¹², there is conflicting experimental evidence that indicates alternate mechanisms may also be involved. For example, several M^{IV} precatalysts such as M(NMe₂)₄ (M = Ti, Zr)²⁰⁸, [Pr₂Si(NDIPP)₂] Zr(NMe₂)₂-LiCl(THF)₃ (DIPP = 2,6-diisopropylphenyl)²¹³ and K⁵-(Me₃SiNCH₂CH₂)₂N(CH₂CH₂SiMe₂CH₂) Zr (REF¹⁹⁶) may promote dehydrogenation via c-bond metathesis routes, while in other cases, M^{III} metallocene species have been observed in situ²¹⁴.

Hydrodefluorination.

C-F bond dissociation energies (BDEs) range from 110 to 130 kcal mol⁻¹ fluoro-organics are not easily biodegraded but instead are often incinerated^{217,218}. Thus, there is considerable impetus to design catalytic methods for C-F bond degradation^{219,220}. Although C-F bond activation has been observed across the transition metal series^{221–229}, early transition metals are exceptionally fluorophilic, making them excellent candidates for hydrodefluorination reactions. There are numerous examples in which low-valent early transition metals can mediate C-F bond activation to generate M-F species, including examples with Ti (REFS^{230–233}), Zr (REFS^{233–239}), Hf (REF²⁴⁰), Nb (REFS^{241–243}) and Sc (REF²⁴⁴). The challenge in these systems is rendering them catalytic, as the resultant M-F species is often a considerable thermodynamic sink²⁴⁵.

In the first demonstration of catalytic C-F bond activation by an early metal, Cp₂TiF₂ was shown to defluorinate perfluorinated cycloalkanes to produce perfluorinated arenes in the presence of excess Al⁰ with HgCl₂ (REF²⁴⁶) (FIG. 5c, top). In the same study, it was found that Cp₂ZrF₂ similarly catalyses this reaction in the presence of Mg⁰ and HgCl₂, although in a rather exothermic fashion. This result was followed by mechanistic work using Cp*₂ZrH₂. Under an atmosphere of H₂, it was found that Cp*₂ZrH₂ reacted with primary, secondary and tertiary C-F bonds to yield the corresponding alkanes and Cp*₂ZrHF (REF²⁴⁷). Intriguingly, mechanistic evidence supported a radical mechanism in which in situ generated Cp*₂ZrH abstracted F% despite the fact that the relative rate of C-F reactivity was 1° > 2° > 3° — the reverse of the pattern seen in typical radical reactions. In support of the radical mechanism, added reductant increased the rate of 1-fluorohexene hydrodefluorination threefold, while radical inhibitors such as 9,10-dihydroanthracene inhibit the rate of defluorination. Radical clock experiments of Cp*₂ZrH₂ with cyclopropylcarbinyl fluoride provided Cp*₂ZrHF and ring-opened Cp*₂Zr(“Bu)H as opposed to methylcyclopropane. It

was suggested that the steric bulk of Cp^*ZrH_2 may be the origin of the observed rate trends^{237–247–248}.

There are also several notable examples of Nb-catalysed hydrodefluorination. Mixtures of NbCl_5 and LiAlH_4 catalyse the hydrodefluorination of aryl fluorides and benzylic fluorides^{249–250} (FIG. 5c, middle). In these reactions, trifluoromethyl groups were reduced fully to methyl groups under mild conditions. In the case of aryl fluoride reductions, it was proposed that the reaction proceeds through a $(\text{DME})\text{NbCl}_3(\text{arene})$ ($\text{DME} = 1,2$ -dimethoxyethane) intermediate that undergoes nucleophilic substitution by hydride, but the mechanism of these reductions has not been fully established. Arnold and Bergman recently examined a more well-defined system for aryl fluoride hydrodefluorination using a d^2 $(\text{BDI})\text{Nb}^{\text{III}}(\text{N}^t\text{Bu})$ fragment²⁵¹ (FIG. 5c, bottom). Mechanistic studies in this system indicate that $(\text{BDI})\text{Nb}^{\text{III}}(\text{N}^t\text{Bu})(\text{arene})$ (**57**) can oxidatively add Ar-F to generate $(\text{BDI})\text{Nb}^{\text{V}}(\text{N}^t\text{Bu})\text{F}(\text{Ar})$ (**55**), which can be transmetallated with silanes, making a strong Si-F bond and a $(\text{BDI})\text{Nb}^{\text{V}}(\text{N}^t\text{Bu})\text{H}(\text{Ar})$ species capable of C-H reductive elimination. In both the NbCl_5 and $(\text{BDI})\text{Nb}$ systems, catalytic turnover is achieved by using a terminal reductant (Si, Al) that is more fluorophilic than Nb.

Catalytic single-electron transfer

Single-electron transfers employing low-valent early transition metal reductants are frequently used in organic synthesis. The Nugent-RajanBabu reagent^{24–25}, Cp_2TiCl , is used in radical ring-opening-cyclization cascades and is the most common early transition metal reagent for single-electron transfer²⁵². These types of cyclization cascades have been used in numerous total syntheses of natural products^{23–253–254}. Cp_2TiCl is easily generated via in situ reduction of commercially available Cp_2TiCl_2 by Zn or Mn dust and has recently been touted as a ‘green’ reagent for chemical synthesis^{255–257}. This reagent typically exists as the chloride-bridged dimer, $[\text{Cp}_2\text{TiCl}]_2$, and forms the active monomeric species through a solvent-assisted equilibrium²⁵⁸.

Gansäuer expanded the use of the Nugent-RajanBabu reagent in the context of radical ring opening of epoxides by developing a catalytic procedure that uses a weak acid (2,4,6-collidine hydrochloride) to protonate the generated titanium alkoxide (**59**) and a reductant (Zn or Mn) to regenerate Cp_2TiCl (REF^{35}) (FIG. 6a, left). 2,4,6-Collidine hydrochloride is an ideal acid for epoxide reactions; it is strong enough to protonate the Ti-alkoxide bonds but weak enough to not open the epoxide ring independently. Furthermore, the liberated 2,4,6-collidine conjugate base is sterically congested, disfavoring association with titanium that would hinder catalysis. In their initial report, Mn was the preferred reductant in this reaction, as the MnCl_2 by-product is insufficiently Lewis acidic to promote epoxide ring opening, although later catalytic applications have successfully used Zn. With appropriate solvent and counteranion choice, cationic $[\text{Cp}_2\text{Ti}]$ fragments can also be used as catalysts²⁵⁹. This catalytic single-electron reduction of epoxides has been used to functionalize arenes, alkenes and alkynes^{259–263} (FIG. 6a, right). Remarkably, these radical reactions are able to proceed enantioselectively when using large, terpene-derived cyclopentadienyl ligands (**60**) on titanium²⁶³.

Cp_2TiCl can also be used to catalyse redox umpolung reactions for the coupling of nitriles or activated alkenes with carbonyls or imines^{73,264–274} (FIG. 6b). These reactions share similar mechanistic features to classical epoxide and azirine ring-opening reactions; owing to the oxophilicity of Ti, treatment of many types of carbonyl derivatives with Ti^{III} species results in the formation of Ti^{IV} carbonyl radicals. These Ti carbonyl radicals can further react with myriad electrophiles^{273,275}. In a detailed mechanistic study of a Ti^{III} -catalysed acyloin-type cross-coupling of ketones and nitriles, it was found that a bimolecular and bimetallic radical recombination mechanism that circumvented free radical intermediates was operational²⁷⁴ (FIG. 6b, bottom). Umpolung-like conjugate additions have also been achieved with hemiaminals, yielding quaternary carbons-a to amines, and coupling or reduction of α,β -unsaturated carbonyls is also possible^{276–278}.

Combining the above approaches, both Gansauer and Lin have each demonstrated that three-membered ring rupture can also be promoted by α -carbonyl reduction in N-acyl aziridines^{279,280} and cyclopropylketones²⁸¹ (FIG. 6c). In the case of N-acyl aziridines (**61**), treatment with Ti^{III} sources results in ring opening that, in contrast to typical aziridine ring openings, produces the more substituted carbon radical **62**. These reactions take advantage of the biradical character^{282–288} of the resulting Ti^{IV} azaenolate or enolate **62**; radical reductive couplings of the ring-opened azaenolate or enolate with alkene electrophiles can be performed catalytically in the presence of reductant (Mn^0 or Zn^0) and weak organic acid (collidine HBr or NEt_3HCl) to form amines (path A), while redox-neutral [3+2] cycloadditions with electrophilic alkenes can also be achieved in a catalytic fashion (path B).

$\text{Cp}_2\text{Ti}^{\text{III}}$ pinacolates, generated through Cp_2TiCl -promoted pinacol coupling of benzaldehyde followed by further reduction, undergo facile deoxygenation to form alkenes. This reaction challenges the classical McMurry coupling mechanism, in which Ti^{II} or Ti^0 complexes were thought to promote carbonyl coupling and/or olefination²⁸⁹.

Single-electron strategies in small-molecule activation and photoredox catalysis.

Reduced early transition metal complexes have historically been excellent platforms for small-molecule activation because of their strong reduction potentials and propensity to form thermodynamically stable metal-ligand multiple bonds. However, these same features become a challenge when attempting to render these reactions catalytic. There are myriad examples of stoichiometric N_2 activation with low-valent early transition metals. Given that this field has been thoroughly reviewed^{4–17}, here, we focus on only several examples that highlight emerging new strategies toward catalytic applications of small-molecule activation — in particular, proton-coupled electron transfers (PCETs) in early transition metals and their applications toward N_2 reduction and water oxidation.

Recent work using group 4 metallocene amide derivatives has found PCET to be a promising route to ammonia formation via M-N hydrogenolysis^{290,291} (FIG. 7a). This route has the potential to overcome the incredible free energy inefficiency of nitrogen reduction involving separated sources of protons and electrons (for example, $[\text{HOEt}_2] + [\text{BArF}_{24}] + \text{KC}_8$). Experimental and computational studies have demonstrated that coordination of NH_3 to $(\eta^5\text{-C}_5\text{Me}_4\text{SiMe}_3)_2\text{M}$ ($\text{M} = \text{Ti}, \text{Zr}$) fragments decreases the N-H bond dissociation free energy (BDFE) by $>40 \text{ kcal mol}^{-1}$ compared with free NH_3 . In general, the degree of N-H

BDFE lowering is a function of the early transition metal redox couple, in which higher energy redox couples (for example, Zr^{IV}/Zr^{III} in **64** or Ti^{III}/Ti^{II} in **65**) lead to greater activation than lower energy couples (for example, Ti^{IV}/Ti^{III} in **63**). This allows for thermodynamically favourable PCET to occur from $(\eta^5-C_5Me_5)(py-Ph)RhH$ (**68**) — which has a weak Rh-H bond ($52.3 \text{ kcal mol}^{-1}$) but is nonetheless able to split molecular H_2 — to $(\eta^5-Me_4SiMe_3)_2TiCl(NR_2)$ derivatives ($R = H, Me, NH_2, NMe_2$) (BDFEs $\wedge 60 \text{ kcal mol}^{-1}$). Using this strategy, the catalytic hydrogenolysis of Ti^{IV} amide complex **66** to the Ti^{III} chloride complex **67** has been realized, opening the possibility of incorporating PCET into a catalytic N_2 reduction cycle with H_2 as the terminal reductant.

Similar to the effects seen with NH_3 , coordination of H_2O to $Cp_2Ti^{III}X$ results in dramatically reduced H-OH BDEs ($A_{bde} \approx 50 \text{ kcal mol}^{-1}$), enabling water to serve as a H-atom donor^{292–294}. This important effect can be used in the photochemical splitting of water. For example, ansa-titanocene dihydroxido complexes undergo Ti-O bond homolysis under visible light to form ansa-titanocene monohydroxido complexes and OH radicals^{295–297}. This reaction had been observed with non-tethered titanocenes before²⁹⁸, but the Cp-Ti BDFE is typically smaller than the Ti-O BDFE, leading primarily to loss of Cp in those systems. A catalytic model system for the photochemical splitting of water has been demonstrated using an ansa-titanocene aquo-hydroxo complex **70** (REF²⁹⁷) (FIG. 7b). Currently, this process requires TEMPO ((2,2,6,6-tetramethylpiperidin-1-yl) oxyl) to reoxidize the bis(aquo) complex **69** to **70**, which can liberate an OH radical upon irradiation. While this reaction may be catalytic, a turnover number could not be confirmed. In contrast to the above reaction, reaction of Cp^*_2TiOTf with water results in rapid oxidation to $Cp^*_2Ti(OH)(OTf)$ and hydrogen gas without TEMPO additive²⁹⁹.

Building upon this work showing that Cp_2MX_2 can undergo ligand-to-metal charge-transfer (LMCT) processes to liberate radicals, it has also been shown that early transition metals can be used as sensitizers for single-electron photoredox catalysis. For example, metal photosensitizers have been designed from π -donor ligands on electron-poor Zr centres (FIG. 7c). As Zr^{IV} is a d^0 metal, these complexes use LMCT processes as opposed to the typical metal-to-ligand charge-transfer (MLCT) processes seen in late transition metal photosensitizers such as $[Ru(bpy)_3]^{2+}$ (REF³⁰⁰). Zr is considerably less expensive and more earth abundant than the common late metals used in photosensitizers, making it an attractive alternative for catalysis. $Zr^{(Me)PDP}_2$ ($^{(Me)PDP} = 2,6\text{-bis}(5\text{-methyl-3-phenyl-1H-pyrrol-2-yl})\text{-pyridine}$) (**73**) has a visible excitation at 528 nm and a photoemission at 594 nm. Cyclic voltammetry (CV) experiments show a reversible reduction at -2.16 V , while Rehm-Weller analysis estimates an excited-state reduction potential of -0.07 V . This excited-state reduction potential is supported by excited-state quenching using weak organic reductants such as 1,3-dimethyl-2-phenyl-2,3-dihydro-1H-7-methylbenzo-[d]imidazole ($^{(Me)BIH}$). Complex **73** is able to catalyse the reductions of ethyl 2-bromo-2,2-difluoroacetate and diethyl maleate to ethyl 2,2-difluoroacetate and diethyl succinate, respectively (FIG. 7c, right). An in-depth study of the homocoupling of benzyl bromide helped support the photoredox mechanism³⁰¹.

Group-transfer catalysis

As is the case with small-molecule activation reactions, engendering catalytic reactivity in group-transfer reactions is particularly challenging owing to the strength of M-E multiple bonds. Classical examples such as methylene transfer in the Tebbe olefination result in the formation of strong Ti-O double bonds, precluding practical turnover. Nonetheless, there have been several notable advances toward catalytic oxidative group-transfer reactions, in particular nitrene transfer. Most of the catalytic early transition metal nitrene transfer reactions involve oxidative imination of unsaturated organic functional groups, which stands in contrast to late-transition-metal-catalysed nitrene transfers, which often perform C-H insertions^{302–305} or aziridinations^{306–308}.

Heyduk has developed several Zr and Ta complexes of aminoquinone-derived redox non-innocent ligands^{80,309}. In these complexes, multi-electron redox changes needed for group transfer take place on the ligand framework instead of at the metal centre, allowing facile, low-temperature access to imido complexes that can undergo further reactivity such as catalytic nitrene coupling^{310,311}, imide reduction³¹² and oxidative isocyanide imination (FIG. 8a, left). For example, in the (NNN) ZrCl (**74**)-catalysed isocyanide imination reaction, the trianionic (tris)amide (NNN)³⁻ ligand undergoes two-electron oxidation by RN₃ to yield the monoanionic (NNN)¹⁻ imido complex **75**, which can insert isocyanide to form an η²-carbodiimide **76**. Upon carbodiimide ligand loss, the NNN ligand framework is again reduced to the (NNN)³⁻ form in **74** to close the catalytic cycle³¹³. A related Ti system based on a diamide-dimine (dadi) framework has also been reported⁸². These (dadi) Ti complexes can access a large array of ligand oxidation states, thereby realizing a two-electron pathway for catalytic azide carbonylation by **77** (FIG. 8a, right). Productive catalysis hinges on weak CO binding to Ti resulting from the redox non-innocent ligand; one would expect Ti^{II} to exhibit strong CO binding due to backdonation into the C-O π* orbital, but because the redox events take place on the ligand, the electron density at Ti remains low throughout the catalytic cycle and thus Ti is incapable of strong backdonation.

In contrast to the above examples of overt ligand redox non-innocence in nitrene transfer, there are several recent examples of formally Ti^{II}/Ti^{IV}-catalysed oxidative nitrene transfer reactions. These reactions yield multi-substituted pyrroles from the coupling of alkynes with azobenzene^{314,315} or α,β-unsaturated imines and α-iminocyclopropanes from the coupling of alkynes and alkenes with azobenzene³¹⁶ (FIG. 8b). The crux of this system is that cooperative metal-ligand effects can also stabilize low-valent metals through backbonding (classical redox non-innocence) into the reagents (alkynes and diazenes) or products (pyrroles and imines), allowing for otherwise challenging reductive events, such as the reductive elimination from **82** to **83**, to occur with relative ease. Although it was initially speculated that azobenzene was a critical reagent for catalytic reoxidation of **83** to **80** through the hydrazido adduct **84**, recent advances have extended this chemistry into other nitrene sources such as azides³¹⁷.

Computational analysis has revealed several additional surprising facets of the reductive elimination events that are critical for catalytic turnover in these Ti^{II}/Ti^{IV} reactions^{318,319}. In general, it appears that direct reductive coupling of M-C and M-X σ-bonds from

intermediates **82** or **88** is unfavourable relative to other reductive pathways. For example, in the pyrrole synthesis, manifold intrinsic bond orbital (IBO) analysis indicates that reductive elimination from the six-membered metallacycle **82** is a π -type electrocyclic ring closure (FIG. 8b, left). By contrast, in the alkene-alkyne coupling reactions, the six-membered metallacycle **88** is semi-saturated (FIG. 8b, right) and thus lacks the requisite π -orbital for electrocyclization. This opens up alternative reductive elimination pathways: p-H elimination to ultimately form α,β -unsaturated imines (**86**) or a pericyclic Kulinkovich-like cyclopropanation to form α -iminocyclopropanes (**87**). It is proposed that the α,β -unsaturated imines are formed through N-H reductive elimination, which has not previously been observed on Ti. Computational studies have proposed a more complex insertion-type pathway for the formation of this product, although experimental evidence for either pathway is currently lacking³²⁰.

In contrast to group 4 metals, group 5 metal imidos tend to be considerably less reactive. To counteract this, efforts have been made to synthesize group 5 systems that electronically saturate the metal *d* orbitals (π -loading), thereby increasing the polarizability of the imido bonds³²¹. The re-loaded β -diketiminate (BDI) Nb bis(imido) complex **89** can thus perform a formally N^{III}/Nb^V-catalysed nitrene transfer reaction to generate asymmetric dialkylcarbodiimides from azides and isocyanides³²² (FIG. 8c).

Outside of catalytic nitrene transfer, there are also many examples of stoichiometric atom-transfer reactions (As₄ (REF.³²³), S₈ (REFS^{324, 325}) and P₄ (REF.³²⁶)). For example, reaction of either P₄ or As₄ with activated zirconocene derivatives results in pnictogen-silicon analogues of benzene³²⁷ and carbon-phosphorous cage compounds³²⁸. The Cummins group has also extensively studied P₄ transformations³²⁹ and synthesized phosphorous-rich organic clusters mediated by a Nb^{III}/Nb^V couple³³⁰.

O-atom transfer is an incredibly challenging target to achieve with oxophilic early transition metals because the strong metal-oxo bond is a thermodynamic sink. In order to circumvent this problem, Nb nitride **90** has been used to perform outer-sphere O-atom transfer from CO₂ (FIG. 8d). CO₂ binds to the terminal anionic nitride in **90**, and the resulting oxide ion in **91** can then be transferred to a RC(O)Z (Z = halide, OAc) acceptor to form **92**, an N-bound cyanate. Subsequent oxidation and reduction steps liberate CO from the bound cyanate in **93**, regenerating **90**. While this system is not catalytic, it remains a rare example of a closed synthetic cycle to achieve CO₂ reduction³³¹.

Summary and outlook

The field of low-valent early transition metal synthesis, reactivity and catalysis has undergone a resurgence of activity in recent years. Many of these new reactions and applications can be split into three categories: reactions providing a modern twist on classic low-valent organometallic transformations, such as catalytic ring-opening reactions and substrate-directed reductive couplings; reactions that push low-valent early transition metals into new, emerging fields, such as PCET, photoredox catalysis and catalyst-controlled radical reactions; and new modes of low-valent inorganic and organometallic reactivity, such as oxidative nitrene transfer catalysis and reversible end-on Sc^{II} N₂ binding.

Each of these categories illustrates that early transition metals play a rich and evolving role in a broad array of chemical applications, and that there likely remains substantially more to discover as the field of low-valent early metal chemistry is revisited through the lens of modern organometallic chemistry. For example, accessing low-valent intermediates under mild conditions (using, for example, organosilane reductants or photoredox chemistry) is only beginning to have an impact in the field and should help to overcome many of the chemical compatibility issues associated with the strong reductants previously used to access low-valent early transition metals.

While many of the reactions discussed here are catalytic in nature, it would be unwise to neglect the importance of fundamental and stoichiometric reactivity with the early transition metals — not only can these later provide the basis for catalytic reactions (as seen in many of the examples discussed above), but the early metals are incredibly earth abundant, environmentally friendly and inexpensive and often have practical utility even in stoichiometric quantities.

Acknowledgements

Financial support was provided by the National Institutes of Health (1R35GM119457) and the Alfred P. Sloan Foundation (I.A.T. is a 2017 Sloan Fellow).

References

1. Hunt AJ, Farmer TJ & Clark JH in *Element Recovery and Sustainability* (ed. Hunt AJ) 1–28 (Royal Society of Chemistry, 2013).
2. Greenwood NN & Earnshaw A *Chemistry of the Elements* (Butterworth-Heinemann, Oxford, 1997).
3. Kepp KP A quantitative scale of oxophilicity and thiophilicity. *Inorg. Chem* 55, 9461–9470 (2016). [PubMed: 27580183]
4. Gambarotta S & Scott J Multimetallic cooperative activation of N₂. *Angew. Chem. Int. Ed* 43, 5298–5308 (2004).
5. Gambarotta S Dinitrogen fixation and activation after 30 years: a puzzle still unsolved. *J. Organomet. Chem* 500, 117–126 (1995).
6. Hidai M & Mizobe Y Recent advances in the chemistry of dinitrogen complexes. *Chem. Rev* 95, 1115–1133 (1995).
7. MacKay BA & Fryzuk MD Dinitrogen coordination chemistry: on the biomimetic borderlands. *Chem. Rev* 104, 385–401 (2004). [PubMed: 14871129]
8. Gardiner MG & Stringer DN Dinitrogen and related chemistry of the lanthanides: a review of the reductive capture of dinitrogen, as well as mono- and di-aza containing ligand chemistry of relevance to known and postulated metal mediated dinitrogen derivatives. *Materials* 3, 841–862 (2010).
9. Studt F & Tuzcek F Theoretical, spectroscopic, and mechanistic studies on transition-metal dinitrogen complexes: implications to reactivity and relevance to the nitrogenase problem. *J. Comput. Chem* 27, 1278–1291 (2006). [PubMed: 16786542]
10. Bazhenova TA & Shilov AE Nitrogen fixation in solution. *Coord. Chem. Rev* 144, 69–145 (1995).
11. Mori M Activation of nitrogen for organic synthesis. *J. Organomet. Chem* 689, 4210–4227 (2004).
12. Fryzuk MD & Johnson SA The continuing story of dinitrogen activation. *Coord. Chem. Rev* 200–202, 379–409 (2000).
13. Burford RJ, Yeo A & Fryzuk MD Dinitrogen activation by group 4 and group 5 metal complexes supported by phosphine-amido containing ligand manifolds. *Coord. Chem. Rev* 334, 84–99 (2017).

14. Fryzuk MD Side-on end-on bound dinitrogen: an activated bonding mode that facilitates functionalizing molecular nitrogen. *Acc. Chem. Res* 42, 127–133 (2009). [PubMed: 18803409]
15. Chirik PJ Dinitrogen functionalization with bis(cyclopentadienyl) complexes of zirconium and hafnium. *Dalton Trans* 0, 16–25 (2007).
16. Burford RJ & Fryzuk MD Examining the relationship between coordination mode and reactivity of dinitrogen. *Nat. Rev. Chem* 1,0026 29(2017).
17. Jia H-P & Quadrelli EA Mechanistic aspects of dinitrogen cleavage and hydrogenation to produce ammonia in catalysis and organometallic chemistry: relevance of metal hydride bonds and dihydrogen. *Chem. Soc. Rev* 43, 547–564 (2014). [PubMed: 24108246]
18. Reichard HA & Micalizio GC Metallocycle-mediated cross-coupling with substituted and electronically unactivated alkenes. *Chem. Sci* 2,573–589 (2011).
19. Yan X & Xi C Conversion of zirconacyclopentadienes into metalloles: Fagan-Nugent reaction and beyond. *Acc. Chem. Res* 48, 935–946 (2015). [PubMed: 25831225]
20. Micalizio GC in *Comprehensive Organic Synthesis II* 2nd edn Vol. 5 (ed. Knochel P) 1660–1737 (Elsevier, 2014).
21. Xu S & Negishi E Zirconium-catalyzed asymmetric carboalumination of unactivated terminal alkenes. *Acc. Chem. Res* 49, 2158–2168 (2016). [PubMed: 27685327]
22. Negishi E Controlled carbometalation as a new tool for carbon-carbon bond formation and its application to cyclization. *Acc. Chem. Res* 20, 65–72 (1987).
23. Zweig JE, Kim DE & Newhouse TR Methods utilizing first-row transition metals in natural product total synthesis. *Chem. Rev* 117, 11680–11752.
24. Nugent WA & RajanBabu TV Transition-metalcentered radicals in organic synthesis. Titanium(III)-induced cyclization of epoxy olefins. *J. Am. Chem. Soc* 110, 8561–8562 (1988).
25. Rosales A et al. The Nugent-RajanBabu reagent: a formidable tool in contemporary radical and organometallic chemistry. *Eur. J. Org. Chem* 2015, 4567–4591 (2015).
26. Parkin G in *Comprehensive Organometallic Chemistry III* Vol. 1 (eds Mingos MP & Crabtree RH) 1–57 (Elsevier, 2007).
27. Grant LN, Miehlich ME, Meyer K & Mindiola DJ Arrested disproportionation in trivalent, mononuclear, and non-metallocene complexes of Zr(III) and Hf(III). *Chem. Commun* 54, 2052–2055 (2018).
28. Kurogi T, Miehlich ME, Halter D & Mindiola DJ, 1, 2-CH bond activation of pyridine across a transient titanium alkylidene radical and re-formation of the Ti=CHtBu moiety. *Organometallics* 37, 165–167 (2018).
29. Perera TH, Lord RL, Heeg MJ, Schlegel HB & Winter C Metallocene pyrimidines and metallocene pyrimidiniums from oxidative addition of pyrazolate N-N bonds to niobium(III), niobium(IV), and tantalum(IV) metal centers and assessment of their aromatic character. *Organometallics* 31,5971–5974 (2012).
30. Pun D, Leopold SM, Bradley CA, Lobkovsky E & Chirik PJ Bis(indenyl)hafnium chemistry: ligand-induced haptotropic rearrangement and fundamental reactivity studies at a reduced hafnium center. *Organometallics* 28, 2471–2484 (2009).
31. Chomitz WA, Sutton AD, Krinsky JL & Arnold J Synthesis and reactivity of titanium and zirconium complexes supported by a multidentate monoanionic [N2P2] ligand. *Organometallics* 28, 3338–3349 (2009).
32. Figueroa JS, Piro NA, Clough CR & Cummins CC A nitridoniobium(V) reagent that effects acid chloride to organic nitrile conversion: synthesis via heterodinuclear (Nb/Mo) dinitrogen cleavage, mechanistic insights, and recycling. *J. Am. Chem. Soc* 128, 940–950 (2006). [PubMed: 16417385]
33. Buschel S et al. Adduct formation of $[(\eta^7\text{-C}_7\text{H}_7)\text{Hf}(\eta^5\text{-C}_5\text{H}_5)]$ with isocyanides, phosphines and N-heterocyclic carbenes: an experimental and theoretical study. *J. Organomet. Chem* 694, 1244–1250 (2009).
34. Okamoto S Synthetic reactions using low-valent titanium reagents derived from Ti(OR)₄ or CpTiX₃ (X=O-i-Pr or Cl) in the presence of Me₃SiCl and Mg. *Chem. Rec* 16, 857–872 (2016). [PubMed: 26969349]
35. Gansauer A, Bluhm H & Pierobon M Emergence of a novel catalytic radical reaction: titanocene-catalyzed reductive opening of epoxides. *J. Am. Chem. Soc* 120, 12849–12859 (1998).

36. Galindo A et al. Alkyl alkyne mono((trimethylsilyl) cyclopentadienyl) niobium complexes. Synthesis and chemical behavior in insertion processes. X-ray crystal structures of $[\text{NbCp}^*(\text{CH}_2\text{SiMe}_3)_2(\text{Me}_3\text{SiCCSiMe}_3)]$ and $[\text{NbCp}^*(\text{NAr})\{\eta^4\text{-CH}(\text{SiMe}_3)\text{C}(\text{SiMe}_3)\text{C}(\text{CH}_2\text{SiMe}_3)=\text{CH}(\text{SiMe}_3)\}]$, ($\text{Cp}^*=\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3$, $\text{Ar} = 2,6\text{-Me}_2\text{C}_6\text{H}_3$). DFT studies of the model complexes $[\text{Nb}(\eta^5\text{-C}_5\text{H}_5)\text{R}_2(\text{HCCH})]$ ($\text{R} = \text{Cl}, \text{Me}$). *Organometallics* 21,293–304 (2002).
37. Calderazzo F, Pampaloni G, Rocchi L, Strahle J & Wurst K The reduction of the NbX₅/AIX₃ system with aluminum in the presence of aromatic hydrocarbons: an approach to niobium(II), niobium(I), and niobium(0) organometallics. *Angew. Chem. Int. Ed* 30, 102–103 (1991).
38. Oshiki T, Yamada A, Kawai K, Arimitsu H & Takai K Alkyne exchange reactions of silylalkyne complexes of tantalum: mechanistic investigation and its application in the preparation of new tantalum complexes having functional alkynes ($\text{PhC}=\text{CR}$ ($\text{R} = \text{COOMe}, \text{CONMe}_2$)). *Organometallics* 26, 173–182 (2007).
39. Kamitani M, Searles K, Chen C-H, Carroll PJ & Mindiola DJ p-hydrogen abstraction of an ethyl group provides entry to titanium and zirconium ethylene complexes. *Organometallics* 34,2558–2566 (2015).
40. Crestani MG et al. Room temperature dehydrogenation of ethane, propane, linear alkanes C₄–C₈, and some cyclic alkanes by titanium-carbon multiple bonds. *J. Am. Chem. Soc* 135, 14754–14767 (2013). [PubMed: 23981228]
41. Cavaliere VN et al. Room temperature dehydrogenation of ethane to ethylene. *J. Am. Chem. Soc* 133, 10700–10703 (2011). [PubMed: 21678955]
42. Fernandez FJ et al. β -hydrogen-containing zirconium alkyls with the doubly-bridged bis(dimethylsilylanediyl) dicyclopentadienyl ligand. X-ray molecular structures of $[\text{Zr}\{(\text{SiMe}_2)_2(\eta^5\text{-C}_5\text{H}_3)_2\}\text{ClEt}]$ and $[\text{Zr}\{(\text{SiMe}_2)_2(\eta^5\text{-C}_5\text{H}_3)_2\}\text{Et}]_2$ ($\text{h-Ch } 2=\text{CH } 2$). *Organometallics* 16, 1553–1561 (1997).
43. Epstein OL, Savchenko AI & Kulinkovich OG Titanium(IV) isopropoxide-catalysed reaction of alkylmagnesium halides with ethyl acetate in the presence of styrene. Non-hydride mechanism of ligand exchange in the titanacyclopropanes. *Tetrahedron Lett* 40, 5935–5938 (1999).
44. Negishi E-I & Takahashi T Alkene and alkyne complexes of zirconocene. Their preparation, structure, and novel transformations. *Bull. Chem. Soc. Jpn* 71, 755–769 (1998).
45. Negishi E-I, Cederbaum FE & Takahashi T Reaction of zirconocene dichloride with alkylolithiums or alkyl grignard reagents as a convenient method for generating a “zirconocene” equivalent and its use in zirconium-promoted cyclization of alkenes, alkynes, dienes, enynes, and diynes. *Tetrahedron Lett* 27, 2829–2832 (1986).
46. Negishi E-I & Takahashi T Patterns of stoichiometric and catalytic reactions of organozirconium and related complexes of synthetic interest. *Acc. Chem. Res* 27, 124–130 (1994).
47. Rosenthal U, Burlakov VV, Arndt P, Baumann W & Spannenberg A The titanocene complex of bis(trimethylsilyl)acetylene: synthesis, structure, and chemistry. *Organometallics* 22, 884–900 (2003).
48. Beweries T et al. Complexation of bis(trimethylsilyl) acetylene by decamethylhafnocene to give the hafnacyclopentene $\text{Cp}^*2\text{Hf}(\eta^2\text{-Me}_3\text{SiC}_2\text{SiMe}_3)$: an unusually strong metal-alkyne interaction. *Organometallics* 26, 247–249 (2007).
49. Beweries T et al. Synthesis of hafnacyclopentanes from hafnocene alkyne complexes: influence of styrene substituents on the C–C coupling regioselectivity. *Eur. J. Inorg. Chem* 2009, 1456–1459 (2009).
50. Fang M et al. Synthesis of the (N₂)₃- radical from Y²⁺ and its protonolysis reactivity to form (N₂H₂)₂- via the Y[N(SiMe₃)₂]₃/KC₈ reduction system. *J. Am. Chem. Soc* 133, 3784–3787 (2011). [PubMed: 21361288]
51. Fang M et al. Isolation of (CO)₁- and (CO)₂- radical complexes of rare earths via Ln(NR₂)₃/K reduction and [K₂(18-crown-6)]₂⁺ oligomerization. *J. Am. Chem. Soc* 6064–6067 (2012). [PubMed: 22435647]
52. Woen DH et al. Solution synthesis, structure, and CO₂ reduction reactivity of a scandium(II) complex, $\{\text{Sc}[\text{N}(\text{SiMe}_3)_2]_3\}^-$. *Angew. Chem. Int. Ed* 56, 2050–2053 (2017). This study presents the first crystallographically characterized Sc(II) compound with amide ligands that reversibly binds

N₂ in an end-on fashion. This example highlights the fact that with the right ligand design, even group 3 low-valent early transition metal complexes can be stabilized.

53. Woen DH et al. End-on bridging dinitrogen complex of scandium. *J. Am. Chem. Soc* 139, 14861–14864 (2017). [PubMed: 28957628]
54. Schmiede BM, Ziller JW & Evans WJ Reduction of dinitrogen with an yttrium metallocene hydride precursor, [(C₅Me₅)₂YH]₂. *Inorg. Chem* 49, 10506–10511 (2010). [PubMed: 20968284]
55. Demir S et al. Synthesis, structure, and density functional theory analysis of a scandium dinitrogen complex, [(C₅Me₄H)Sc]₂(μ-η²:η²-N₂). *J. Am. Chem. Soc* 132, 11151–11158 (2010). [PubMed: 20698681]
56. Clentsmith GKB et al. Stabilization of low-oxidation-state early transition-metal complexes bearing 1,2,4-triphosphacyclopentadienyl ligands: structure of [{Sc(P₃C₂tBu₂)₂]₂}; Sc(II) or mixed oxidation state? *Angew. Chem. Int. Ed* 42, 1038–1041 (2003).
57. Wijeratne GB et al. Electronic structure and reactivity of a well-defined mononuclear complex of Ti(II). *Inorg. Chem* 54, 10380–10397 (2015). [PubMed: 26451744]
58. Edema JH, Duchateau R, Gambarotta S, Hynes R & Gabe E Novel titanium(II) amine complexes L₄TiCl₂[L = 1/2 N,N,N',N'-tetramethylethylenediamine (TMEDA), 1/2 N,N,N'-trimethylethylenediamine, pyridine, 1/2,2',2''-bipyridine]: synthesis and crystal structure of monomeric trans-(TMEDA)₂TiCl₂. *Inorg. Chem* 30, 154–156 (1991).
59. Araya MA, Cotton FA, Matonic JH & Murillo CA An efficient reduction process leading to titanium(II) and niobium(II): preparation and structural characterization of trans-MCl₂(py)₄ compounds, M = Ti, Nb, and Mn. *Inorg. Chem* 34, 5424–5428 (1995).
60. Chakarawet K et al. Ta(CNDipp)₆: an isocyanide analogue of hexacarbonyltantalum(0). *Angew. Chem. Int. Ed* 56, 10577–10581 (2017).
61. Allen JM & Ellis JE Synthesis and characterization of titanium tetrakisocyanide complexes, [CpTi(CNXyl)₄E], E = I, SnPh₃, and SnMe₃. *J. Organomet. Chem* 693, 1536–1542 (2008).
62. Barybin MV et al. Homoleptic isocyanidometalates of 4d- and 5d-transition metals: [Nb(CNXyl)₆]-, [Ta(CNXyl)₆]-, and derivatives thereof. *J. Am. Chem. Soc* 129, 1141–1150 (2007). [PubMed: 17263395]
63. Jilek RE, Jang M, Smolensky ED, Britton JD & Ellis JE Structurally distinct homoleptic anthracene complexes, [M(C₁₄H₁₀)₃]₂-, M = titanium, zirconium, hafnium: tris(arene) complexes for a triad of transition metals. *Angew. Chem. Int. Ed* 47, 8692–8695 (2008).
64. Sussman VJ & Ellis JE From storable sources of atomic Nb- and Ta- ions to isolable anionic tris(1,3-butadiene)metal complexes: [M(η⁴-C₄H₅)₃]-, M = Nb, Ta. *Angew. Chem. Int. Ed* 47, 484–489 (2008).
65. MacDonald MR et al. Expanding rare-earth oxidation state chemistry to molecular complexes of holmium(II) and erbium(II). *J. Am. Chem. Soc* 134, 8420–8423 (2012). [PubMed: 22583320]
66. Arteaga-Muller R et al. New tantalum ligand-free catalyst system for highly selective trimerization of ethylene affording 1-hexene: new evidence of a metallacycle mechanism. *J. Am. Chem. Soc* 131, 5370–5371 (2009). [PubMed: 20560633]
67. Nishiyama H et al. Structural and electronic noninnocence of a-diimine ligands on niobium for reductive C-Cl bond activation and catalytic radical addition reactions. *J. Am. Chem. Soc* 139, 6494–6505 (2017). [PubMed: 28409626]
68. Tsurugi H, Saito T, Tanahashi H, Arnold J & Mashima K Carbon radical generation by d⁰ tantalum complexes with a-diimine ligands through ligand-centered redox processes. *J. Am. Chem. Soc* 133, 18673–18683 (2011). [PubMed: 21981364]
69. Saito T et al. Reduction of (tBuN=)NbCl₃(py)₂ in a salt-free manner for generating Nb(IV) dinuclear complexes and their reactivity toward benzo[c] cinnoline. *Inorg. Chem* 54, 6004–6009 (2015). [PubMed: 26017157]
70. Saito T et al. 1,4-bis(trimethylsilyl)-1,4-diaza-2,5-cyclohexadienes as strong salt-free reductants for generating low-valent early transition metals with electron-donating ligands. *J. Am. Chem. Soc* 136, 5161–5170 (2014). [PubMed: 24597916] This study shows that inner-sphere organosilicon reductants offer many advantages over traditional alkyl metal halide reductants to access low-valent early transition metals: they are less prone to over-reduction, and the organic by-products are easily removed.

71. Gomez M, Hernandez-Prieto C, Martin A, Mena M & Santamaria C An effective route to dinuclear niobium and tantalum imido complexes. *Inorg. Chem* 56, 11681–11687 (2017). [PubMed: 28901759]
72. Klesko JP, Thrush CM & Winter CH Thermal atomic layer deposition of titanium films using titanium tetrachloride and 2-methyl-1,4-bis(trimethylsilyl)-2,5-cyclohexadiene or 1,4-bis(trimethylsilyl)-1,4-dihydropyrazine. *Chem. Mater* 27, 4918–4921 (2015).
73. Frey G, Hausmann JN & Streuff J Titanium-catalyzed reductive umpolung reactions with a metal-free terminal reducing agent. *Chem. Eur. J* 21, 5693–5696 (2015). [PubMed: 25712472]
74. Satoh Y & Obora Y Active low-valent niobium catalysts from NbCl₅ and hydrosilanes for selective intermolecular cycloadditions. *J. Org. Chem* 76, 8569–8573 (2011). [PubMed: 21919436]
75. Paradies J et al. Photogeneration of titanium(III) from titanium(IV) citrate in aqueous solution. *J. Inorg. Biochem* 100, 1260–1264 (2006). [PubMed: 16603248]
76. Chong E, Xue W, Storr T, Kennepohl P & Schafer LL Pyridonate-supported titanium(III). Benzylamine as an easy-to-use reductant. *Organometallics* 34, 4941–4945 (2015).
77. Gianetti TL et al. Diniobium inverted sandwich complexes with [j-r]6:16-arene ligands: synthesis, kinetics of formation, and electronic structure. *J. Am. Chem. Soc* 135, 3224–3236 (2013). [PubMed: 23343247]
78. Plundrich GT, Wadepohl H, Clot E & Gade LH η^6 -arene-zirconium-PNP-pincer complexes: mechanism of their hydrogenolytic formation and their reactivity as zirconium(II) synthons. *Chem. Eur. J* 22, 9283–9292 (2016). [PubMed: 27258989]
79. Hananouchi S, Krull BT, Ziller JW, Furche F & Heyduk AF Metal effects on ligand non-innocence in group 5 complexes of the redox-active [ONO] pincer ligand. *Dalton Trans* 43, 17991–18000 (2014). [PubMed: 25352212]
80. Munha RF, Zarkesh RA & Heyduk AF Group transfer reactions of d⁰ transition metal complexes: redox-active ligands provide a mechanism for expanded reactivity. *Dalton Trans* 42, 3751–3766. (2013) [PubMed: 23334157] This perspective shows how redox non-innocent ligands can access multiple redox states and facilitate catalytic group-transfer reactions with early transition metals.
81. Duan L et al. Synthesis, characterization, and reversible multielectron redox properties of a biradical yttrium complex containing bis(2-isopropylaminophenyl) amide. *Eur. J. Inorg. Chem* 2017, 2231–2235 (2017).
82. Heins SP, Wolczanski PT, Cundari TR & MacMillan SN Redox non-innocence permits catalytic nitrene carbonylation by (dadi)Ti=NAd (Ad=adamantyl). *Chem. Sci* 8, 3410–3418 (2017). [PubMed: 28507712]
83. Fagan PJ, Nugent WA & Calabrese JC Metallacycle transfer from zirconium to main group elements: a versatile synthesis of heterocycles. *J. Am. Chem. Soc* 116, 1880–1889 (1994).
84. Fagan PJ, Burns EG & Calabrese JC Synthesis of boroles and their use in low-temperature Diels-Alder reactions with unactivated alkenes. *J. Am. Chem. Soc* 110, 2979–2981 (1988).
85. Fagan PJ & Nugent WA Synthesis of main group heterocycles by metallacycle transfer from zirconium. *J. Am. Chem. Soc* 110, 2310–2312 (1988).
86. Kulinkovich OG, Sviridov SV, Vasilevskii DA & Pritytskaya TS Reaction of ethylmagnesium bromide with esters of carboxylic-acid in the presence of tetraisopropoxytitanium. *Zh. Org. Khim* 25, 2244 (1989).
87. Epstein OL, Savchenko AI & Kulinkovich OG On the mechanism of titanium-catalyzed cyclopropanation of esters with aliphatic organomagnesium compounds. Deuterium distribution in the reaction products of (CD₃)₂CHMgBr with ethyl 3-chloropropionate in the presence of titanium tetraisopropoxide. *Russ. Chem. Bull* 49, 378–380 (2000).
88. Corey EJ, Rao SA & Noe MC Catalytic diastereoselective synthesis of cis-1,2-disubstituted cyclopropanols from esters using a vicinal dicarbanion equivalent. *J. Am. Chem. Soc* 116, 9345–9346 (1994).
89. Geis O & Schmalz H-G New developments in the Pauson-Khand reaction. *Angew. Chem. Int. Ed* 37, 911–914 (1998).
90. Hicks FA & Buchwald SL Highly enantioselective catalytic Pauson-Khand type formation of bicyclic cyclopentenones. *J. Am. Chem. Soc* 118, 11688–11689 (1996).

91. Hicks FA & Buchwald SL An intramolecular titanium-catalyzed asymmetric Pauson-Khand type reaction. *J. Am. Chem. Soc* 121, 7026–7033 (1999).
92. Hicks FA, Kablaoui NM & Buchwald SL Scope of the intramolecular titanocene-catalyzed Pauson-Khand type reaction. *J. Am. Chem. Soc* 121, 5881–5898 (1999).
93. Sturla SJ & Buchwald SL Catalytic asymmetric cyclocarbonylation of nitrogen-containing enynes. *J. Org. Chem* 64, 5547–5550 (1999). [PubMed: 11674619]
94. Kablaoui NM, Hicks FA & Buchwald SL Titanocene-catalyzed cyclocarbonylation of o-allyl aryl ketones to γ -butyrolactones. *J. Am. Chem. Soc* 119, 4424–4431 (1997).
95. Campbell AD, Taylor RJK & Raynham TM The total synthesis of (–)- α -kainic acid using titanium-mediated diene metallabicyclisation methodology. *Chem. Commun* 0, 245–246 (1999).
96. Cheng X & Micalizio GC Synthesis of neurotrophic seco-prezizaane sesquiterpenes (1R,10S)-2-oxo-3,4-dehydroneomajucin, (2S)-hydroxy-3,4-dehydroneomajucin, and (–)-jiadifenin. *J. Am. Chem. Soc* 138, 1150–1153 (2016). [PubMed: 26785051]
97. Meng Z et al. Total synthesis and antiviral activity of indolosesquiterpenoids from the xiamycin and oridamycin families. *Nat. Commun* 6, 6096 (2015) [PubMed: 25648883]
98. Chaplinski V & de Meijere A A versatile new preparation of cyclopropylamines from acid dialkylamides. *Angew. Chem. Int. Ed* 35, 413–414 (1996).
99. Quan LG, Kim S-H, Lee JC & Cha JK Diastereoselective synthesis of trans-1,2-dialkylcyclopropanols by the Kulinkovich hydroxycyclopropanation of homoallylic alcohols. *Angew. Chem. Int. Ed* 41, 2160–2162 (2002).
100. Kiel GR, Samkian AE, Nicolay A, Witzke RJ & Tilley TD Titanocene-mediated dinitrile coupling: a divergent route to nitrogen-containing polycyclic aromatic hydrocarbons. *J. Am. Chem. Soc* 140, 2450–2454 (2018). [PubMed: 29383934] This is an example of the emerging field of nitrile couplings, demonstrating their applications in the field of polycyclic aromatics and conducting materials.
101. Konik YA, Kananovich DG & Kulinkovich OG Enantioselective cyclopropanation of carboxylic esters with alkyl magnesium bromides in the presence of titanium(IV) (4 R,5 R)-TADDOLates. *Tetrahedron* 69, 6673–6678 (2013).
102. Kulinkovich OG, Kananovich DG, Lopp M & Snieckus V Insight into the mechanism and stereochemistry of the transformations of alkyltitanium ate-complexes. An enhanced enantioselectivity in the cyclopropanation of the carboxylic esters with titanacyclopropane reagents. *Adv. Synth. Catal* 356, 3615–3626 (2014).
103. Barysevich MV et al. Stereoselective synthesis of α -methyl and α -alkyl ketones from esters and alkenes via cyclopropanol intermediates. *Chem. Commun* 54, 2800–2803 (2018).
104. Finn PB, Derstine BP & Sieburth SM Carbocyclic amino ketones by Bredt's rule-arrested Kulinkovich-de Meijere reaction. *Angew. Chem. Int. Ed* 55, 2536–2539 (2016).
105. Fawcett FS Bredt's rule of double bonds in atomic-bridged-ring structures. *Chem. Rev* 47, 219–274 (1950). [PubMed: 24538877]
106. Kim S-H, Park Y, Choo H & Cha JK Regio- and stereochemistry of inter- and intramolecular titanium-mediated coupling of imides and mono-substituted olefins. *Tetrahedron Lett* 43, 6657–6660 (2002).
107. Santra S, Masalov N, Epstein OL & Cha JK Diastereoselective, titanium-mediated cyclization of w -vinyl tethered imides. *Org. Lett* 7, 5901–5904 (2005) [PubMed: 16354095]
108. Lee J, Ha JD & Cha JK New synthetic method for functionalized pyrrolizidine, indolizidine, and mitomycin alkaloids. *J. Am. Chem. Soc* 119, 8127–8128 (1997).
109. Takahashi M & Micalizio GC Regio- and stereoselective cross-coupling of substituted olefins and imines. A convergent stereoselective synthesis of saturated 1,5-aminoalcohols and substituted piperidines. *J. Am. Chem. Soc* 129, 7514–7516 (2007). [PubMed: 17530760] This study shows that highly regioselective and stereoselective transformations can be accomplished using alkoxide directing groups in combination with stoichiometric amounts of Ti complexes.
110. Hill JE, Balaich G, Fanwick PE & Rothwell IP The chemistry of titanacyclopentadiene rings supported by 2,6-diphenylphenoxide ligation: stoichiometric and catalytic reactivity. *Organometallics* 12, 2911–2924 (1993).

111. See XY et al. Generation of TiII alkyne trimerization catalysts in the absence of strong metal reductants. *Organometallics* 36, 1383–1390 (2017). [PubMed: 28690352]
112. Eisch JJ, Gitua JN, Otieno PO & Shi X Carbon-carbon bond formation via oxidative-addition processes of titanium(II) reagents with π -bonded organic substrates. Reactivity modifications by Lewis acids and Lewis bases: part 22. *Organic chemistry of subvalent transition metal complexes*. *J. Organomet. Chem* 624, 229–238 (2001).
113. Rassadin VA & Six Y A study of the reaction of *n*-BuLi with Ti(Oi-Pr)₄ as a method to generate titanacyclopropane and titanacyclopropene species. *Tetrahedron* 70, 787–794 (2014).
114. Tarselli MA & Micalizio GC Aliphatic imines in titanium-mediated reductive cross-coupling: unique reactivity of Ti(O-*i*-Pr)₄/*n*-BuLi. *Org. Lett* 11, 4596–4599 (2009). [PubMed: 19810765]
115. Chen MZ et al. Preparation of stereodefined homoallylic amines from the reductive cross-coupling of allylic alcohols with imines. *J. Org. Chem* 75, 8048–8059 (2010). [PubMed: 21069994]
116. Yang D & Micalizio GC Convergent and stereodivergent synthesis of complex 1-aza-7-oxabicyclo[2.2.1]heptanes. *J. Am. Chem. Soc* 133, 9216–9219 (2011). [PubMed: 21604673]
117. Chen MZ & Micalizio GC Three-component coupling sequence for the regiospecific synthesis of substituted pyridines. *J. Am. Chem. Soc* 134, 1352–1356 (2012). [PubMed: 22103772]
118. Greszler SN, Reichard HA & Micalizio GC Asymmetric synthesis of dihydroindanes by convergent alkoxide-directed metallacycle-mediated bond formation. *J. Am. Chem. Soc* 134, 2766–2774 (2012). [PubMed: 22235773]
119. Jeso V et al. Synthesis of angularly substituted trans-fused hydroindanes by convergent coupling of acyclic precursors. *J. Am. Chem. Soc* 136, 8209–8212 (2014). [PubMed: 24856045]
120. Cheng X & Micalizio GC An annulation reaction for the synthesis of cross-conjugated triene-containing hydroindanes from acyclic precursors. *Org. Lett* 16, 5144–5147 (2014). [PubMed: 25216381]
121. Mizoguchi H & Micalizio GC Synthesis of angularly substituted trans-fused decalins through a metallacycle-mediated annulative cross-coupling cascade. *Angew. Chem. Int. Ed* 55, 13099–13103 (2015).
122. Mizoguchi H & Micalizio GC Synthesis of highly functionalized decalins via metallacycle-mediated cross-coupling. *J. Am. Chem. Soc* 137, 6624–6628 (2015). [PubMed: 25930967]
123. Valadez TN, Norton JR & Neary MC Reaction of Cp*(Cl)M(diene) (M = Ti, Hf) with isonitriles. *J. Am. Chem. Soc* 137, 10152–10155 (2015). [PubMed: 26230451]
124. Hessen B, Blenkins J, Teuben JH, Helgesson G & Jagner S Carbon-carbon couplings in the reactions of unsaturated Group 4 metal σ -cis-butadiene complexes with 2,6-xylyl isocyanide. *Organometallics* 8, 830–835 (1989).
125. Valadez TN, Norton JR, Neary MC & Quinlivan PJ Reaction of Cp*₂Zr(2,3-dimethylbutadiene) with isonitriles and CO. *Organometallics* 35, 3163–3169 (2016).
126. Becker L, Haehnel M, Spannenberg A, Arndt P & Rosenthal U Reactions of group 4 metallocenes with monosubstituted acetonitriles: keteniminate formation versus C-C coupling. *Chem. Eur. J* 21, 3242–3248 (2015). [PubMed: 25641093]
127. Becker L, Arndt P, Jiao H, Spannenberg A & Rosenthal U Nitrile-nitrile C-C coupling at group 4 metallocenes to form 1-metalla-2,5-diaza-cyclopenta-2,4-dienes: synthesis and reactivity. *Angew. Chem. Int. Ed* 52, 11396–11400 (2013).
128. Rosenthal U Reactions of group 4 metallocene bis(trimethylsilyl)acetylene complexes with nitriles and isonitriles. *Angew. Chem. Int. Ed* 10.1002/anie.201805157 (2018).
129. Rehbaum F, Thiele K-H & Trojanov SI Darstellung und molekulstruktur des (μ -alkylidenamido) titanocenkomplexes [(C₅H₅)₂TiCl]₂[μ -{ N = C(CH₂C₅H₅) C(CH₂C₆H₅) = N. *J. Organomet. Chem* 410, 327–333 (1991).
130. Kiel GR, Patel SC, Smith PW, Levine DS & Tilley TD Expanded helicenes: a general synthetic strategy and remarkable supramolecular and solid-state behavior. *J. Am. Chem. Soc* 139, 18456–18459 (2017). [PubMed: 29215272]
131. Kiel GR, Ziegler MS & Tilley TD Zirconacyclopentadiene-annulated polycyclic aromatic hydrocarbons. *Angew. Chem. Int. Ed* 56, 4839–4844 (2016)

132. Nitschke JR & Tilley TD Novel templating effect in the macrocyclization of functionalized diynes by zirconocene coupling. *Angew. Chem. Int. Ed* 40, 2142–2145 (2001).
133. Nitschke JR, Zurcher S & Tilley TD New zirconocene-coupling route to large, functionalized macrocycles. *J. Am. Chem. Soc* 122, 10345–10352 (2000).
134. Altenburger K et al. Multiple and highly selective alkyne-isonitrile C-C and C-N couplings at group 4 metallocenes. *Chem. Eur. J* 22, 9169–9180 (2016). [PubMed: 27226122]
135. Rosenthal U, Burlakov VV, Bach MA & Beweries T Five-membered metallacycles of titanium and zirconium? Attractive compounds for organometallic chemistry and catalysis. *Chem. Soc. Rev* 36, 719 (2007). [PubMed: 17471397]
136. Beweries T et al. Reactions of decamethylhafnocene with 1,3-butadiynes: formation of hafnacyclocumulenes and C-H activation at pentamethylcyclopentadienyl ligands. *Organometallics* 26, 6827–6831 (2007).
137. Rosenthal U, Pellny P-M, Kirchbauer FG & Burlakov VV What do titano- and zirconocenes do with diynes and polyynes? *Acc. Chem. Res* 33, 119–129 (2000). [PubMed: 10673320]
138. Burlakov VV et al. The first titanacyclic five-membered cumulene. Synthesis, structure, and reactivity. *Chem. Ber* 128, 967–971 (1995).
139. Burlakov VV et al. Novel acetylene complexes of titanocene and permethyltitanocene without additional ligands. Synthesis spectral characteristics and X-ray diffraction study. *J. Organomet. Chem* 476, 197–206 (1994).
140. Rosenthal U, Burlakov VV, Arndt P, Baumann W & Spannenberg A Five-membered titana- and zirconacyclocumulenes: stable 1-metallacyclopenta- 2,3,4-trienes. *Organometallics* 24, 456–471. (2005)
141. Kaleta K et al. Reactions of group 4 metallocene alkyne complexes with carbodiimides: experimental and theoretical studies of the structure and bonding of five-membered hetero-metallacycloallenes. *J. Am. Chem. Soc* 133, 5463–5473 (2011). [PubMed: 21413717]
142. Tomaschun G et al. Group 4 metallocene complexes and cyanopyridines: coordination or coupling to metallacycles. *Eur. J. Inorg. Chem* 2016, 272–280 (2016)
143. Diekmann M et al. Chiral bis(η^5 : η^1 -pentafulvene) titanium complexes. *Organometallics* 25, 339–348 (2006)
144. Theilmann O, Saak W, Haase D & Beckhaus R Reactions of low-valent titanocene(II) fragments with trans-4,4'-azobispyridine (RN=NR, R=C₅H₄N): formation of tetranuclear molecular squares by trans-cis isomerization. *Organometallics* 28, 2799–2807 (2009).
145. Loose F, Schmidtman M, Saak W & Beckhaus R Imines in the titanium coordination sphere: highly reactive titanaaziridines and larger titanacycles formed by subsequent C-C coupling reactions. *Eur. J. Inorg. Chem* 2015, 5171–5187 (2015).
146. Jaroschik F et al. Synthesis, characterization and reactivity of formal 20 electron zirconocene-pentafulvene complexes. *Organometallics* 36, 2004–2013 (2017).
147. Manßen M, Lauterbach N, Woriescheck T, Schmidtman M & Beckhaus R Reactions of secondary amines with bis(η^5 : η^1 -pentafulvene)titanium complexes: formation of titanium amides and titanaaziridines. *Organometallics* 36, 867–876 (2017).
148. Fischer M, Schaper R, Jaugstetter M, Schmidtman M & Beckhaus R Electrophilic d⁰ cations of group 4 metals (M=Ti, Zr, Hf) derived from monopentafulvene complexes: direct formation of tridentate Cp, O, P-ligands. *Organometallics* 37, 1192–1205 (2018).
149. Eisch JJ, Owuor FA & Shi X Novel synthesis of unbridged, sterically substituted zirconocene dichlorides from fulvenes and dialkylzirconium dichlorides via zirconium(IV) hydride transfer. *Organometallics* 18, 1583–1585 (1999).
150. Eisch JJ, Shi X & Owuor FA Novel synthesis of ansa-metallocenes via the reductive dimerization of fulvenes with group 4 metal divalent halides. *Organometallics* 17, 5219–5221 (1998).
151. Parker KDJ & Fryzuk MD Carbon-carbon bond forming reactions with tantalum diamidophosphine complexes that incorporate alkyne ligands. *Organometallics* 33, 6122–6131 (2014).
152. Ziegler JA, Buckley HL & Arnold J Synthesis and reactivity of tantalum corrole complexes. *Dalton Trans* 46, 780–785 (2017). [PubMed: 27996068]

153. Yamamoto K, Tsurugi H & Mashima K Direct evidence for a [4 + 2] cycloaddition mechanism of alkynes to tantalacyclopentadiene on dinuclear tantalum complexes as a model of alkyne cyclotrimerization. *Chem. Eur. J* 21, 11369–11377 (2015). [PubMed: 26118787]
154. Yamamoto K, Nagae H, Tsurugi H & Mashima K Mechanistic understanding of alkyne cyclotrimerization on mononuclear and dinuclear scaffolds: [4 + 2] cycloaddition of the third alkyne onto metallacyclopentadienes and dimetallacyclopentadienes. *Dalton Trans* 45, 17072–17081 (2016). [PubMed: 27730228]
155. Obora Y, Takeshita K & Ishii Y NbCl₃-catalyzed [2 + 2 + 2] intermolecular cycloaddition of alkynes and alkenes to 1,3-cyclohexadiene derivatives. *Org. Biomol. Chem* 7, 428–431 (2009). [PubMed: 19156304]
156. Obora Y, Satoh Y & Ishii Y NbC⁺-catalyzed intermolecular [2 + 2 + 2] cycloaddition of alkynes and a, w-dienes: highly chemo- and regioselective formation of 5-w-alkenyl-1,4-substituted-1,3-cyclohexadiene derivatives. *J. Org. Chem* 75, 6046–6049 (2010). [PubMed: 20681534]
157. Satoh Y & Obora Y NbCl₃-catalyzed three-component [2 + 2 + 2] cycloaddition reaction of terminal alkynes, internal alkynes, and alkenes to 3,4,5-tetrasubstituted 1,3-cyclohexadienes. *Org. Lett* 13, 2568–2571 (2011). [PubMed: 21491900]
158. Satoh Y & Obora Y Low-valent niobium-catalyzed intermolecular [2 + 2 + 2] cycloaddition of tert-butylacetylene and aryl nitriles to form 2,3,6-trisubstituted pyridine derivatives. *J. Org. Chem* 78, 7771–7776 (2013). [PubMed: 23848456]
159. Satoh Y & Obora Y Niobium complexes in organic transformations: from stoichiometric reactions to catalytic [2 + 2 + 2] cycloaddition reactions. *Eur. J. Org. Chem* 2015, 5041–5054 (2015).
160. LaPointe RE, Wolczanski PT & Mitchell JF Carbon monoxide cleavage by (silox)₃ Ta (silox=t-Bu₃SiO-). *J. Am. Chem. Soc* 108, 6382–6384 (1986).
161. LaPointe RE & Wolczanski PT Silox hydrides (silox=t-Bu₃SiO-) of group 5: do [(silox)₂MH₂]₂ (M=Nb, Ta) complexes contain unbridged M-M bonds? *J. Am. Chem. Soc* 108, 3535–3537 (1986).
162. Steffey BD et al. Intramolecular arene hydrogenation by niobium aryloxide compounds: stereochemistry of cyclohexadiene formation. *J. Am. Chem. Soc* 111, 378–380 (1989).
163. Chesnut RW, Steffey BD & Rothwell IP The metallation and hydrogenation of aryl rings in early transition metal aryloxide systems. *Polyhedron* 8, 1607–1610 (1989).
164. Bell RA, Cohen SA, Doherty NM, Threlkel RS & Bercaw JE Borohydride, hydride, halide, and carbonyl derivatives of bis(pentamethylcyclopentadienyl)niobium. *Organometallics* 5, 972–975 (1986).
165. Parkin G et al. Alpha- and beta-migratory insertion and elimination processes for alkyl complexes of permethylscandocene and permethyltantalocene. *J. Mol. Catal* 41, 21–39 (1987).
166. Millar SP, Zubris DL, Bercaw JE & Eisenberg R On the mechanism of dihydrogen addition to tantalocene complexes. *J. Am. Chem. Soc* 120, 5329–5330 (1998).
167. Tebbe FN & Parshall GW Hydride derivatives of niobocene and tantalocene. *J. Am. Chem. Soc* 93, 3793–3795 (1971).
168. Ankianiec BC, Fanwick PE & Rothwell IP Isolation of a new series of seven-coordinate hydride compounds of tantalum(V) and their involvement in the catalytic hydrogenation of arene rings. *J. Am. Chem. Soc* 113, 4710–4712 (1991).
169. Parkin BC et al. Synthesis and characterization of a series of mononuclear tantalum(V) hydride compounds containing aryloxide ligation. *Organometallics* 14, 3002–3013 (1995).
170. Yu JS, Ankianiec BC, Nguyen MT & Rothwell IP All-cis catalytic hydrogenation of polynuclear aromatic hydrocarbons by group 5 metal aryloxide compounds. *J. Am. Chem. Soc* 114, 1927–1929 (1992).
171. Visciglio VM et al. Coordination and hydrogenation of 1,3-cyclohexadiene by niobium and tantalum aryl oxide compounds: relevance to catalytic arene hydrogenation. *J. Am. Chem. Soc* 119, 3490–3499 (1997).
172. Parker KG, Noll B, Pierpont CG & Dubois MR Syntheses of tantalum(V) complexes containing tetramethylpyrrolyl, pyrrolyl, and indolyl ligands. *Inorg. Chem* 35, 3228–3234 (1996). [PubMed: 11666522]

173. Fryzuk MD, Kozak CM, Bowdridge MR & Patrick BO Cyclohexadienyl niobium complexes and arene hydrogenation catalysis. *Organometallics* 21, 5047–5054 (2002).
174. Clark JR, Fanwick PE & Rothwell IP Aryloxi ligand dependent reactivity of tantalum dihydride compounds with alkenes. *J. Chem. Soc. Chem. Commun* 553–554 (1995).
175. Mulford DR, Clark JR, Schweiger SW, Fanwick PE & Rothwell IP Reactions of alkynes and olefins with tantalum hydrides containing aryloxi ancillary ligation: relevance to catalytic hydrogenation. *Organometallics* 18, 4448–4458 (1999).
176. Gianetti TL, Tomson NC, Arnold J & Bergman RG Z-selective, catalytic internal alkyne semihydrogenation under H₂/CO mixtures by a niobium(III) imido complex. *J. Am. Chem. Soc* 133, 14904–14907 (2011). [PubMed: 21854008]
177. Gianetti TL, Bergman RG & Arnold J Stoichiometric carbon-carbon bond formation mediated by well defined Nb(III) complexes. *Polyhedron* 84, 19–23 (2014).
178. Gianetti TL, La Pierre HS & Arnold J Group 5 imides and bis(imide)s as selective hydrogenation catalysts. *Eur. J. Inorg. Chem* 2013, 3771–3783 (2013).
179. Tomson NC, Arnold J & Bergman RG Synthesis, characterization, and reactions of isolable (beta-diketiminato)niobium(III) imido complexes. *Organometallics* 29, 5010–5025 (2010). [PubMed: 21116450]
180. Brintzinger HH & Bercaw JE The nature of so-called titanocene, (C₁₀H₁₀Ti)₂. *J. Am. Chem. Soc* 92, 6182–6185 (1970).
181. Figueroa JS, Piro NA, Mindiola DJ, Fickes MG & Cummins CC Niobaziridine hydrides. *Organometallics* 29, 5215–5229 (2010).
182. Rankin MA & Cummins CC Carbon dioxide reduction by terminal tantalum hydrides: formation and isolation of bridging methylene diolate complexes. *J. Am. Chem. Soc* 132, 10021–10023 (2010). [PubMed: 20593902]
183. Berno P & Gambarotta S Formation of a metallaaziridine ring and dinitrogen fixation promoted by a niobium amide complex. *Organometallics* 14, 2159–2161 (1995).
184. Curtis MD, Bell LG & Butler WM C-H activation. Synthesis of silyl derivatives of niobocene and tantalocene hydrides, their H/D exchange reactions with C₆D₆, and the structure of Cp₂Ta(H)₂SiMe₂Ph. *Organometallics* 4, 701–707 (1985).
185. Kriegel BM, Bergman RG & Arnold J Generation of low-valent tantalum species by reversible C-H activation in a cyclometallated tantalum hydride complex. *Dalton Trans* 43, 10046–10056 (2014). [PubMed: 24870398]
186. Ballard KR, Gardiner IM & Wigley DE Evidence for intramolecular C-H bond activation and the formation of a “tucked-in” complex of hexamethylbenzene at a tantalum(III) center. *J. Am. Chem. Soc* 111, 2159–2162 (1989).
187. Bercaw JE, Marvich RH, Bell LG & Brintzinger HH Titanocene as an intermediate in reactions involving molecular hydrogen and nitrogen. *J. Am. Chem. Soc* 94, 1219–1238 (1972).
188. Yu JS, Fanwick PE & Rothwell IP Intramolecular alkane dehydrogenation and functionalization at niobium metal centers. *J. Am. Chem. Soc* 11, 8171–8172 (1990).
189. Riley PN, Clark JR, Fanwick PE & Rothwell IP Synthesis and structure of niobium and tantalum derivatives of bis(dicyclohexylphosphino)methane (dcpm). *Inorg. Chim. Acta* 288, 35–39 (1999).
190. Yu JS et al. Intramolecular dehydrogenation of alkyl groups at niobium aryloxi centers: bonding and reactivity of the ensuing niobacyclopropane ring. *Organometallics* 15, 4443–4449 (1996).
191. Steffey BD et al. Intramolecular activation of aliphatic and aromatic carbon-hydrogen bonds by tantalum(III) metal centers: synthesis and structure of the bis-metalated compounds Ta(OC₆H₃ButCMe₂CH₂)₂Cl and Ta(OC₆H₃PhC₆H₄)₂ (OAr-2,6-Ph₂) (OAr-2,6-Ph₂=2,6-diphenylphenoxide). *Organometallics* 8, 1419–1423 (1989).
192. Mindiola DJ & Cummins CC Probing the niobium metallaaziridine functionality. *Organometallics* 20, 3626–3628 (2001).
193. Figueroa JS & Cummins CC The niobaziridine-hydride functional group: synthesis and divergent reactivity. *J. Am. Chem. Soc* 125, 4020–4021 (2003). [PubMed: 12670202]
194. Figueroa JS & Cummins CC A niobaziridine hydride system for white phosphorus or dinitrogen activation and N- or P-atom transfer. *Dalton Trans* 35, 2161–2168 (2006).

195. Aguilar-Calderon JR, Metta-Magana AJ, Noll B & Fortier S C(sp³)-H oxidative addition and transfer hydrogenation chemistry of a titanium(II) synthon: mimicry of late-metal type reactivity. *Angew. Chem. Int. Ed* 55, 14101–14105 (2016).
196. Erickson KA, Stelmach JPW, Mucha NT & Waterman R Zirconium-catalyzed amine borane dehydrocoupling and transfer hydrogenation. *Organometallics* 34, 4693–4699 (2015).
197. Tayebani M, Feghali K, Gambarotta S & Yap G Molecular rearrangements of a low-valent niobium amide: ligand C-H bond oxidative addition and reductive elimination. *Organometallics* 17, 4282–4290 (1998).
198. Shaver MP & Fryzuk MD Phosphine- induced ancillary ligand orthometalation at a tantalum-tantalum double bond. *Organometallics* 24, 2606–2601 (2005).
199. Wolczanski PT Activation of carbon-hydrogen bonds via 1,2-RH-addition/-elimination to early transition metal imides. *Organometallics* 37, 505–516 (2018).
200. Bailey BC, Fan H, Huffman JC, Baik M-H & Mindiola DJ Intermolecular C-H bond activation reactions promoted by transient titanium alkylidynes. Synthesis, reactivity, kinetic, and theoretical studies of the Ti=C linkage. *J. Am. Chem. Soc* 129, 8781–8793. [PubMed: 17592842]
201. Bailey BC et al. Intermolecular C-H bond activation promoted by a titanium alkylidyne. *J. Am. Chem. Soc* 127, 16016–16017 (2005). [PubMed: 16287275]
202. Solowey DP et al. A new and selective cycle for dehydrogenation of linear and cyclic alkanes under mild conditions using a base metal. *Nat. Chem* 9, 1126–1132 (2017). [PubMed: 29064500]
In this study, the mechanism for dehydrogenation is proposed to avoid metal hydride intermediates, resulting in high terminal selectivity, as the terminal alkenes cannot isomerize.
203. Hickey AK et al. Dehydrogenation of hydrocarbons with metal-carbon multiple bonds and trapping of a titanium(II) intermediate. *Dalton Trans* 43, 9834–9837 (2014). [PubMed: 24841924]
204. Kamitani M et al. Phosphinoalkylidene and -alkylidyne complexes of titanium: intermolecular C-H bond activation and dehydrogenation reactions. *J. Am. Chem. Soc* 137, 11872–11875 (2015). [PubMed: 26302157]
205. Clark TJ, Russell CA & Manners I Homogeneous, titanocene-catalyzed dehydrocoupling of amine-borane adducts. *J. Am. Chem. Soc* 128, 9582–9583 (2006). [PubMed: 16866483]
206. Pun D, Lobkovsky E & Chirik PJ Amineborane dehydrogenation promoted by isolable zirconium sandwich, titanium sandwich and N₂ complexes. *Chem. Commun* 43, 3297–3299 (2007).
207. Sloan ME et al. Homogeneous catalytic dehydrocoupling/dehydrogenation of amine-borane adducts by early transition metal, group 4 metallocene complexes. *J. Am. Chem. Soc* 132, 3831–3841 (2010). [PubMed: 20180565]
208. Beweries T, Hansen S, Kessler M, Klahn M & Rosenthal U Catalytic dehydrogenation of dimethylamine borane by group 4 metallocene alkyne complexes and homoleptic amido compounds. *Dalton Trans* 40, 7689–7692 (2011). [PubMed: 21512699]
209. Beweries T et al. Catalytic and kinetic studies of the dehydrogenation of dimethylamine borane with an iPr substituted titanocene catalyst. *ChemCatChem* 3, 1865–1868 (2011).
210. Thomas J, Klahn M, Spannenberg A & Beweries T Group 4 metallocene catalysed full dehydrogenation of hydrazine borane. *Dalton Trans* 42, 14668–14672 (2013). [PubMed: 23884155]
211. Jurca T et al. Step-growth titanium-catalysed dehydropolymerisation of amine-boranes. *Chem. Sci* 9, 3360–3366 (2018). [PubMed: 29780466]
212. Luo Y & Ohno K Computational study of titanocene-catalyzed dehydrocoupling of the adduct Me₂NH-BH₃: an intramolecular, stepwise mechanism. *Organometallics* 26, 3597–3600 (2007).
213. Lummis PA, McDonald R, Ferguson MJ & Rivard E Synthesis, characterisation, and dehydrocoupling ability of zirconium complexes bearing hindered bis(amido)silyl ligands. *Dalton Trans* 44, 7009–7020 (2015). [PubMed: 25781718]
214. Helten H et al. Paramagnetic titanium(III) and zirconium(III) metallocene complexes as precatalysts for the dehydrocoupling/dehydrogenation of amine-boranes. *Angew. Chem. Int. Ed* 52, 437–440 (2013).
215. Richmond TG in *Topics in Organometallic Chemistry Vol. 3* (eds Murai S et al.) 243–269 (1999).

216. Blanksby SJ & Ellison GB Bond dissociation energies of organic molecules. *Acc. Chem. Res* 36, 255–263 (2003). [PubMed: 12693923]
217. Giesy JP & Kannan K Global distribution of perfluorooctane sulfonate in wildlife. *Environ. Sci. Technol* 35, 1339–1342 (2001).
218. de Boer J, Wester PG, Klamer HJC, Lewis WE & Boon JP Do flame retardants threaten ocean life? *Nature* 394, 28–29 (1998). [PubMed: 9665124]
219. Schine KP & Sturges WT CO₂ is not the only gas. *Science* 315, 1804–1805 (2007). [PubMed: 17395819]
220. Perutz RN A catalytic foothold for fluorocarbon reactions. *Science* 321, 1168–1169 (2008). [PubMed: 18755964]
221. Kiplinger JL, Richmond TG & Osterberg CE Activation of carbon-fluorine bonds by metal complexes. *Chem. Rev* 94, 373–431 (1994).
222. Richmond TG Organometallic transformations demonstrate that fluorocarbons are reactive molecules. *Angew. Chem. Int. Ed* 39, 3241–3244 (2000).
223. Klahn M & Rosenthal U An update on recent stoichiometric and catalytic C-F bond cleavage reactions by lanthanide and group 4 transition-metal complexes. *Organometallics* 31, 1235–1244 (2012).
224. Driver TG Niobium-catalyzed activation of aryl trifluoromethyl groups and functionalization of C-H bonds: an efficient and convergent approach to the synthesis of N-heterocycles. *Angew. Chem. Int. Ed* 48, 7974–7976 (2009).
225. Amii H & Uneyama KC-F Bond activation in organic synthesis. *Chem. Rev* 109, 2119–2183 (2009). [PubMed: 19331346]
226. Clot E et al. C-F and C-H bond activation of fluorobenzenes and fluoropyridines at transition metal centers: how fluorine tips the scales. *Acc. Chem. Res* 44, 333–348 (2011). [PubMed: 21410234]
227. Kuehnel MF, Lentz D & Braun T Synthesis of fluorinated building blocks by transition-metal-mediated hydrodefluorination reactions. *Angew. Chem. Int. Ed* 52, 3328–3348 (2013).
228. Ahrens T, Kohlmann J, Ahrens M & Braun T Functionalization of fluorinated molecules by transition-metal-mediated C-F bond activation to access fluorinated building blocks. *Chem. Rev* 115, 931–972 (2015). [PubMed: 25347593]
229. Eisenstein O, Milani J & Perutz RN Selectivity of C-H activation and competition between C-H and C-F bond activation at fluorocarbons. *Chem. Rev* 117, 8710–8753 (2017). [PubMed: 28653537]
230. Bouwkamp MW et al. Structure of the decamethyl titanocene cation, a metallocene with two agostic C-H bonds, and its interaction with fluorocarbons. *J. Am. Chem. Soc* 124, 12956–12957 (2002). [PubMed: 12405816]
231. Piglosiewicz IM, Kraft S, Beckhaus R, Haase D & Saak W Selective C-H and C-F bond activation reactions of pyridine and fluoropyridines — formation of binuclear μ -X titanocene complexes (X = H, F) with α -functionalized N-heterocycles. *Eur. J. Inorg. Chem* 2005, 938–945 (2005).
232. Burk MJ, Staley DL & Tumas W Oxidatively induced reductive elimination. A novel titanium complex resulting from C-F bond activation. *J. Chem. Soc. Chem. Commun* 809–810 (1990).
233. Rosenthal U et al. Bis(trimethylsilyl)acetylene complexes of titanocenes and zirconocenes: their recent chemistry and reactions with Lewis acids. *Eur. J. Inorg. Chem* 2004, 4739–4749 (2004).
234. Kiplinger JL & Richmond TG Selective room temperature hydrogenolysis of aromatic fluorocarbons mediated by a low-valent zirconium complex. *Chem. Commun* 1115–1116 (1996).
235. Fujiwara M, Ichikawa J, Okauchi T & Minami T Vinylic C-F bond activation with low-valent zirconocene: the generation and cross-coupling reactions of 1-fluorovinylzirconocene. *Tetrahedron Lett* 40, 7261–7265 (1999).
236. O'Connor PE, Berg DJ & Barclay T Photolytic zirconium benzyl bond cleavage and subsequent aryl C-F activation in zirconium complexes of fluorinated aryl diamides. *Organometallics* 21, 3947–3954 (2002).
237. Jones WD Activation of C-F bonds using Cp*₂ZrH₂: a diversity of mechanisms. *Dalton Trans* 2003, 3991–3995 (2003).

238. Jager-Fiedler U et al. Reactions of zirconocene bis(trimethylsilyl)acetylene complexes with fluorinated pyridines: C-H versus C-F bond activation. *Eur. J. Inorg. Chem* 2005, 2842–2849 (2005).
239. Edelbach BL, Kraft BM & Jones WD Generation of perfluoropolyphenylene oligomers via carbon-fluorine bond activation by Cp₂Zr(C₆F₅)₂: a dual mechanism involving a radical chain and release of tetrafluorobenzene. *J. Am. Chem. Soc* 121, 10327–10331 (1999).
240. Rieth RD, Brennessel WW & Jones WD Activation of aromatic, aliphatic, and olefinic carbon-fluorine bonds using Cp*₂HfH₂. *Eur. J. Inorg. Chem* 2007, 2839–2847 (2007).
241. Fuchibe K & Akiyama T Low-valent niobium-mediated double activation of C-F/C-H bonds: fluorene synthesis from o-arylated a, a, a-trifluorotoluene derivatives. *J. Am. Chem. Soc* 128, 1434–1435 (2006). [PubMed: 16448098]
242. Gianetti TL, Bergman RG & Arnold J Disassembly of a benzylic CF₃ group mediated by a niobium(III) imido complex. *J. Am. Chem. Soc* 135, 8145–8148 (2013). [PubMed: 23663120]
243. Nechayev M, Gianetti TL, Bergman RG & Arnold J C-F sp² bond functionalization mediated by niobium complexes. *Dalton Trans* 44, 19494–19500 (2015). [PubMed: 26216334]
244. Huang W & Diaconescu PL Aromatic C-F bond activation by rare-earth-metal complexes. *Organometallics* 36, 89–96 (2017).
245. Simoes JAM & Beauchamp JL Transition metal-hydrogen and metal-carbon bond strengths: the keys to catalysis. *Chem. Rev* 90, 629–688 (1990).
246. Kiplinger JL & Richmond TG Group IV metallocene-mediated synthesis of fluoroaromatics via selective defluorination of saturated perfluorocarbons. *J. Am. Chem. Soc* 118, 1805–1806 (1996).
247. Kraft BM, Lachicotte RJ & Jones WD Aliphatic carbon-fluorine bond activation using (C₅Me₅)₂ZrH₂. *J. Am. Chem. Soc* 122, 8559–8560 (2000).
248. Kraft BM, Lachicotte RJ & Jones WD Aliphatic and aromatic carbon-fluorine bond activation with Cp*₂ZrH₂: mechanisms of hydrodefluorination. *J. Am. Chem. Soc* 123, 10973–10979 (2001). [PubMed: 11686701]
249. Fuchibe K, Ohshima Y, Mitomi K & Akiyama T Low-valent niobium-catalyzed reduction of a, a, a-trifluorotoluenes. *Org. Lett* 9, 1497–1499 (2007). [PubMed: 17352487]
250. Fuchibe K, Mitomi K, Suzuki R & Akiyama T C-C coupling reactions of superstrong CF₃ groups with C(sp²)-H bonds: reactivity and synthetic utility of zero-valent niobium catalyst. *Chem. Asian J* 3, 261–271 (2008). [PubMed: 18175305]
251. Gianetti TL, Bergman RG & Arnold J Carbon-fluorine bond cleavage in fluoroarenes via a niobium(III) imido complex: from stoichiometric to catalytic hydrodefluorination. *Chem. Sci* 5, 2517–2524 (2014). This study shows that the well-characterized (BDI)Nb platform allows for a more detailed understanding of the mechanism of Nb-catalysed hydrodefluorination.
252. Barrero AF et al. Couplings of benzylic halides mediated by titanocene chloride: synthesis of bibenzyl derivatives. *J. Org. Chem* 72, 2251–2254 (2007). [PubMed: 17309310]
253. Cha JY, Yeoman JTS & Reisman SE A concise total synthesis of (–)-maoecrystal Z. *J. Am. Chem. Soc* 133, 14964–14967 (2011). [PubMed: 21877709]
254. Plesniak MP, Huang H-M & Procter DJ Radical cascade reactions triggered by single electron transfer. *Nat. Rev. Chem* 1, 0077 (2017).
255. Castro Rodriguez M et al. Cp₂TiCl: an ideal reagent for green chemistry? *Org. Process Res. Dev* 21, 911–923 (2017).
256. Richrath RB et al. Cp₂TiX complexes for sustainable catalysis in single-electron steps. *Chem. Eur. J* 24, 6371–6379 (2018). [PubMed: 29327511]
257. Morcillo SP et al. Recent applications of Cp₂TiCl in natural product synthesis. *Org. Chem. Front* 1, 15–33 (2014).
258. Enemark RJ, Larsen J, Skrydstrup T & Daasbjerg K Revelation of the nature of the reducing species in titanocene halide-promoted reductions. *J. Am. Chem. Soc* 126, 7853–7864 (2004). [PubMed: 15212533]
259. Gansauer A et al. Cationic titanocene(III) complexes for catalysis in single-electron steps. *Angew. Chem. Int. Ed* 54, 7003–7006 (2015).

260. Friedrich J et al. Titanocene catalyzed 4-exo cyclizations: mechanism, experiment, catalyst design. *J. Am. Chem. Soc* 130, 1788–1796 (2008). [PubMed: 18189399]
261. Gansauer A, Shi L & Otte M Catalyticenantioselective radical cyclization via regiodivergent epoxide opening. *J. Am. Chem. Soc* 132, 11858–11859 (2010). [PubMed: 20687604]
262. Hildebrandt S & Gansauer A Synthesis of dihydropyrrolizine and tetrahydroindolizine scaffolds from pyrroles by titanocene(III) catalysis. *Angew.Chem. Int. Ed* 55, 9719–9722 (2016).
263. Gansauer A, Lauterbach T, Bluhm H & Noltemeyer M A catalytic enantioselective electron transfer reaction: titanocene-catalyzed enantioselective formation of radicals from meso-epoxides. *Angew. Chem. Int. Ed* 38, 2909–2910 (1999).
264. Streuff J A titanium(III)-catalyzed redox umpolung reaction for the reductive cross-coupling of enones with acrylonitriles. *Chem. Eur. J* 17, 5507–5510 (2011). [PubMed: 21488110]
265. Bichovski P, Haas TM, Keller M & Streuff J Direct conjugate alkylation of a, p-unsaturated carbonyls by TiIII-catalysed reductive umpolung of simple activated alkenes. *Org. Biomol. Chem* 14, 5673–5682 (2016). [PubMed: 26806535]
266. Leijendekker LH, Weweler J, Leuther TM & Streuff J Catalytic reductive synthesis and direct derivatization of unprotected aminoindoles, aminopyrroles, and iminoindolines. *Angew. Chem. Int. Ed* 56, 6103–6106 (2017).
267. Streuff J, Feurer M, Bichovski P, Frey G & Gellrich U Enantioselective titanium(III)-catalyzed reductive cyclization of ketonitriles. *Angew. Chem. Int. Ed* 51, 8661–8664 (2012).
268. Frey G, Luu H-T, Bichovski P, Feurer M & Streuff J Convenient titanium(III)-catalyzed synthesis of cyclic aminoketones and pyrrolidinones-development of a formal [4 + 1] cycloaddition. *Angew. Chem. Int. Ed* 52, 7131–7134 (2013).
269. Feurer M, Frey G, Luu H-T, Kratzert D & Streuff J The cross-selective titanium(III) catalysed acyloin reaction. *Chem. Commun* 50, 5370–5372 (2014)
270. Streuff J Reductive umpolung reactions with low-valent titanium catalysts. *Chem. Rec* 14, 1100–1113 (2014). [PubMed: 25234255]
271. Bichovski P, Haas TM, Kratzert D & Streuff J Synthesis of bridged benzazocines and benzoxocines by a titanium-catalyzed double-reductive umpolung strategy. *Chem. Eur. J* 21, 2339–2342 (2015). [PubMed: 25476744]
272. Luu H-T, Wiesler S, Frey G & Streuff J A titanium(III)- catalyzed reductive umpolung reaction for the synthesis of 1,1-disubstituted tetrahydroisoquinolines. *Org. Lett* 17, 2478–2481 (2015). [PubMed: 25928360]
273. Streuff J & Gansauer A Metal-catalyzed β -functionalization of Michael acceptors through reductive radical addition reactions. *Angew. Chem. Int. Ed* 54, 14232–14242 (2015). This review highlights strategies for accessing catalytic single-electron chemistry with early transition metal complexes and their subsequent applications in Michael-type chemistry.
274. Streuff J et al. Mechanism of the TiIII-catalyzed acyloin-type umpolung: a catalyst-controlled radical reaction. *J. Am. Chem. Soc* 137, 14396–14405 (2015) [PubMed: 26495878] This report details a bimolecular mechanism for the acyloin-type umpolung reaction that avoids free radical intermediates from open-shell catalysts.
275. Gansauer A & Hildebrandt S in *Topics in Heterocyclic Chemistry Vol. 54* (ed. Landais Y) 253–283 (Springer, 2018).
276. Zheng X et al. Umpolung of hemiaminals: titanocene-catalyzed dehydroxylative radical coupling reactions with activated alkenes. *Angew. Chem. Int. Ed* 52, 3494–3498 (2013).
277. Estevez RE et al. Stereocontrolled coupling between aldehydes and conjugated alkenals mediated by TiIII/H₂O. *Org. Lett* 8, 5433–5436 (2006). [PubMed: 17107040]
278. Kosal AD & Ashfeld BL Titanocene-catalyzed conjugate reduction of a, p-unsaturated carbonyl derivatives. *Org. Lett* 12, 44–47 (2010). [PubMed: 19968238]
279. Hao W et al. Radical redox-relay catalysis: formal [3+2] cycloaddition of N-acylaziridines and alkenes. *J. Am. Chem. Soc* 139, 12141–12144 (2016)
280. Zhang Y-Q, Vogelsang E, Qu Z-W, Grimme S & Gansauer A Titanocene-catalyzed radical opening of N-acylated aziridines. *Angew. Chem. Int. Ed* 56, 12654–12657 (2017).

281. Hao W, Harenberg JH, Wu X, MacMillan SN & Lin S Diastereo- and enantioselective formal [3 + 2] cycloaddition of cyclopropyl ketones and alkenes via Ti-catalyzed radical redox relay. *J. Am. Chem. Soc* 140, 3514–3517 (2018). [PubMed: 29465998]
282. Alvarado J, Herrmann AT & Zakarian A Stereoselective α -fluorination of N-acyloxazolidinones at room temperature within 1 h. *J. Org. Chem* 79, 6206–6220 (2014). [PubMed: 24926887]
283. Mabe PJ & Zakarian A Asymmetric radical addition of TEMPO to titanium enolates. *Org. Lett* 16, 516–519 (2014). [PubMed: 24377351]
284. Herrmann AT, Smith LL & Zakarian A A simple method for asymmetric trifluoromethylation of N-acyl oxazolidinones via Ru-catalyzed radical addition to zirconium enolates. *J. Am. Chem. Soc* 134, 6976–6979 (2012). [PubMed: 22486383]
285. Gu Z, Herrmann AT & Zakarian A Dual Ti-Ru catalysis in the direct radical haloalkylation of N-acyl oxazolidinones. *Angew. Chem. Int. Ed* 50, 7136–7139 (2011).
286. Ciez D, Patasz A & Trzewik B Titanium enolate chemistry at the beginning of the 21st century. *Eur. J. Org. Chem* 2016, 1476–1493 (2016).
287. Beaumont S, Ilardi EA, Monroe LR & Zakarian A Valence tautomerism in titanium enolates: catalytic radical haloalkylation and application in the total synthesis of heodysidenin. *J. Am. Chem. Soc* 132, 1482–1483 (2010). [PubMed: 20078122]
288. Moreira I & de PR et al. Unconventional biradical character of titanium enolates. *J. Am. Chem. Soc* 130, 3242–3243 (2008). [PubMed: 18288838]
289. Dieguez HR et al. Weakening C-O bonds: Ti(III), a new reagent for alcohol deoxygenation and carbonyl coupling olefination. *J. Am. Chem. Soc* 132, 254–259 (2010). [PubMed: 20000601]
290. Pappas I & Chirik PJ Ammonia synthesis by hydrogenolysis of titanium-nitrogen bonds using proton coupled electron transfer. *J. Am. Chem. Soc* 137, 3498–3501 (2015). [PubMed: 25719966]
291. Pappas I & Chirik PJ Catalytic proton coupled electron transfer from metal hydrides to titanocene amides, hydrazides and imides: determination of thermodynamic parameters relevant to nitrogen fixation. *J. Am. Chem. Soc* 138, 13379–13389 (2015) This report highlights the large impact that coordination to an early transition metal can have on the N-H BDFE in NH₃.
292. Cuerva JM et al. Water: the ideal hydrogen-atom source in free-radical chemistry mediated by Tl(III) and other single-electron-transfer metals? *Angew. Chem. Int. Ed* 45, 5522–5526 (2006).
293. Paradis M et al. Understanding the exceptional hydrogen-atom donor characteristics of water in Ti(III)-mediated free-radical chemistry. *J. Am. Chem. Soc* 132, 12748–12756 (2010). [PubMed: 20722395]
294. Gansauer A et al. H₂O activation for hydrogen-atom transfer: correct structures and revised mechanisms. *Angew. Chem. Int. Ed* 51, 3266–3270 (2012).
295. Kessler M et al. Photoassisted Ti-O activation in a decamethyltitanocene dihydroxido complex: insights into the elemental steps of water splitting. *Angew. Chem. Int. Ed* 51, 6272–6275 (2012).
296. Godemann C et al. Highly selective visible light-induced Ti-O bond splitting in an ansa-titanocene dihydroxido complex. *Chem. Commun* 51, 3065–3068 (2015).
297. Godemann C et al. A model of a closed cycle of water splitting using ansa-titanocene(III/IV) triflate complexes. *J. Am. Chem. Soc* 137, 16187–16195 (2015). [PubMed: 26641723]
298. Harrigan RW, Hammond GS & Gray HB Photochemistry of titanocene(IV) derivatives. *J. Organomet. Chem* 81, 79–85 (1974).
299. Kessler M et al. Synthesis of Cp*₂Ti(OTf) and its reaction with water. *Eur. J. Inorg. Chem* 2011, 627–631 (2011).
300. Zhang Y, Petersen JL & Milsmann C A luminescent zirconium(IV) complex as a molecular photosensitizer for visible light photoredox catalysis. *J. Am. Chem. Soc* 138, 13115–13118 (2016). [PubMed: 27643820]
301. Zhang Y, Lee TS, Petersen JL & Milsmann C A zirconium photosensitizer with a long-lived excited state: mechanistic insight into photoinduced single-electron transfer. *J. Am. Chem. Soc* 140, 5934–5947 (2018). [PubMed: 29671586] This study shows that early transition metals can act as photosensitizers, taking advantage of LMCT processes as opposed to the classic MLCT processes used for Ru(bpy)₃²⁺ and similar complexes.

302. Roizen JL, Harvey ME & Du Bois J Metal-catalyzed nitrogen-atom transfer methods for the oxidation of aliphatic C-H bonds. *Acc. Chem. Res* 45, 911–922 (2012). [PubMed: 22546004]
303. Qin C & Davies HML Role of sterically demanding chiral dirhodium catalysts in site-selective C-H functionalization of activated primary C-H bonds. *J. Am. Chem. Soc* 136, 9792–9796 (2014). [PubMed: 24933043]
304. Iovan DA, Wilding MJT, Baek Y, Hennessy ET & Betley TA Diastereoselective C-H bond amination for disubstituted pyrrolidines. *Angew. Chem. Int. Ed* 56, 15599–15602 (2017).
305. Basuli F, Wicker B, Huffman JC & Mindiola DJ Understanding the role of an easy-to-prepare aldimine-alkyne carboamination catalyst, [Ti(NMe₂)₃(NHMe₂)]B(C₅F₅)₄. *J. Organomet. Chem* 696, 235–243 (2011).
306. Fantauzzi S et al. Origin of the deactivation in styrene aziridination by aryl azides, catalyzed by ruthenium porphyrin complexes. Structural characterization of a A₂-1,2,3-triazoline RuII(TPP)CO complex. *Organometallics* 24, 4710–4713 (2005).
307. King ER, Hennessy ET & Betley TA Catalytic C-H bond amination from high-spin iron imido complexes. *J. Am. Chem. Soc* 133, 4917–4923 (2011). [PubMed: 21405138]
308. Cramer SA & Jenkins DM Synthesis of aziridines from alkenes and aryl azides with a reusable macrocyclic tetracarbene iron catalyst. *J. Am. Chem. Soc* 133, 19342–19345 (2011). [PubMed: 22081884]
309. Heyduk AF, Zarkesh RA & Nguyen AI Designing catalysts for nitrene transfer using early transition metals and redox-active ligands. *Inorg. Chem* 50, 9849–9863 (2011). [PubMed: 21774482]
310. Zarkesh RA, Ziller JW & Heyduk AF Four-electron oxidative formation of aryl diazenes using a tantalum redox-active ligand complex. *Angew. Chem. Int. Ed* 47, 4715–4718 (2008).
311. Zarkesh RA & Heyduk AF Reactivity of organometallic tantalum complexes containing a bis(phenoxy)amide (ONO)₃- ligand with aryl azides and 1,2-diphenylhydrazine. *Organometallics* 30, 4890–4898 (2011).
312. Blackmore KJ, Lal N, Ziller JW & Heyduk AF Catalytic reactivity of a zirconium(IV) redox-active ligand complex with 1,2-diphenylhydrazine. *J. Am. Chem. Soc* 130, 2728–2729 (2008). [PubMed: 18266370]
313. Nguyen AI, Zarkesh RA, Lacy DC, Thorson MK & Heyduk AF Catalytic nitrene transfer by a zirconium(IV) redox-active ligand complex. *Chem. Sci* 2, 166–169 (2011).
314. Gilbert ZW, Hue RJ & Tonks IA Catalytic formal [2 + 2 + 1] synthesis of pyrroles from alkynes and diazenes via TiII/TiIV redox catalysis. *Nat. Chem* 8, 63–68 (2016). [PubMed: 26673265]
This study provides an example of Ti redox catalysis and group transfer in the absence of redox non-innocent ligands, highlighting the role that π -accepting ligands can play in promoting catalysis with low-valent early transition metal complexes.
315. Chiu H & Tonks IA Trimethylsilyl-protected alkynes as selective cross-coupling partners in titanium-catalyzed [2 + 2 + 1] pyrrole synthesis. *Angew. Chem. Int. Ed* 57, 6090–6094 (2018).
316. Davis-Gilbert ZW, Yao LJ & Tonks IA Ti-catalyzed multicomponent oxidative carboamination of alkynes with alkenes and diazenes. *J. Am. Chem. Soc* 138, 14570–14573 (2016). [PubMed: 27790910]
317. Pearce AJ, See XY & Tonks IA Oxidative nitrene transfer from azides to alkynes via Ti(II)/Ti(IV) redox catalysis: formal [2 + 2 + 1] synthesis of pyrroles. *Chem. Commun* 54, 6891–6894 (2018).
318. Davis-Gilbert ZW, Wen X, Goodpaster JD & Tonks IA Mechanism of Ti-catalyzed oxidative nitrene transfer in [2 + 2 + 1] pyrrole synthesis from alkynes and azobenzene. *J. Am. Chem. Soc* 140, 7267–7281 (2018). [PubMed: 29763560]
319. Guo J et al. Differences between the elimination of early and late transition metals: DFT mechanistic insights into the titanium-catalyzed synthesis of pyrroles from alkynes and diazenes. *Chem. Sci* 8, 2413–2425 (2017). [PubMed: 28451348]
320. Guo J et al. Strong preference of the redox-neutral mechanism over the redox mechanism for the TiIV catalysis involved in the carboamination of alkyne with alkene and diazene. *Chem. Eur. J* 24, 7010–7025 (2018). [PubMed: 29709085]

321. Tomson NC, Arnold J & Bergman RG Halo, alkyl, aryl, and bis(imido) complexes of niobium supported by the (3-diketiminato) ligand. *Organometallics* 29, 2926–2942 (2010). [PubMed: 20671985]
322. Kriegel BM, Bergman RG & Arnold J Nitrene metathesis and catalytic nitrene transfer promoted by niobium bis(imido) complexes. *J. Am. Chem. Soc* 138, 52–55 (2016). [PubMed: 26698833]
323. Schmidt M et al. Transfer reagent for bonding isomers of iron complexes. *J. Am. Chem. Soc* 139, 13981–13984 (2017). [PubMed: 28933848]
324. Corbey JF, Fang M, Ziller JW & Evans WJ Cocrystallization of (η -S₂)²⁻ and (n-S)²⁻ and formation of an [r²-S₃N(SiMe₃)₂] ligand from chalcogen reduction by (N₂)²⁻ in a bimetallic yttrium amide complex. *Inorg. Chem* 54, 801–807 (2015). [PubMed: 25166587]
325. Fieser ME et al. Dinitrogen reduction, sulfur reduction, and isoprene polymerization via photochemical activation of trivalent bis(cyclopentadienyl) rare-earth-metal allyl complexes. *Organometallics* 34, 4387–4393 (2015).
326. Camp C, Maron L, Bergman RG & Arnold J Activation of white phosphorus by low-valent group 5 complexes: formation and reactivity of cyclo-P₄ inverted sandwich compounds. *J. Am. Chem. Soc* 136, 17652–17661 (2014). [PubMed: 25469924]
327. Seitz AE et al. Pnictogen-silicon analogues of benzene. *J. Am. Chem. Soc* 138, 10433–10436 (2016). [PubMed: 27513314]
328. Vogel U et al. Access to phosphorus-rich zirconium complexes. *Angew. Chem. Int. Ed* 50, 8982–8985 (2011).
329. Cossairt BM, Piro NA & Cummins CC Early transition metal mediated activation and transformation of white phosphorus. *Chem. Rev* 110, 4164–4177 (2010). [PubMed: 20175534]
330. Cossairt BM & Cummins CC A niobium-mediated cycle producing phosphorus-rich organic molecules from white phosphorus (P₄) through activation, functionalization, and transfer reactions. *Angew. Chem. Int. Ed* 47, 8863–8866 (2008).
331. Silvia JS & Cummins CC Ligand-based reduction of CO₂ to CO mediated by an anionic niobium nitride complex. *J. Am. Chem. Soc* 132, 2169–2171 (2010). [PubMed: 20121151]

Inner-sphere reductants

Reagents that act as reductants after first forming a bond to the complex of interest.

Umpolung reactions

Reactions that proceed through polarity inversion of the given functional group.

Redox non-innocent ligands

Ligands bound to a metal complex in which the oxidation state is ambiguous, wherein oxidation or reduction of the ligands may occur in tandem with or instead of metal oxidation or reduction.




Fig. 1 |. Relative abundance of the valence states of early transition metal organometallic complexes.

Reduced early transition metal complexes become increasingly rare as one proceeds from group 5 to earlier groups and also from period 3 to period 5. Data adapted from REF²⁶.




Fig. 2 | Examples of synthetic routes to low-valent early transition metal complexes.
a | Classical reduction routes involving strong reductants. **b** | Representative examples of modern low-valent metal synthesis. **c** | New organosilane reductants for low-valent metal synthesis and catalysis. These routes demonstrate the diverse array of reduction methods available and the common ability of π -accepting ligands to stabilize low-valent early transition metal complexes. Red-coloured metals are in their highest oxidation state, orange-coloured metals are one-electron reduced, and teal-coloured metals are two-electron

reduced. $h\nu$, irradiation; Dipp, 2,6-diisopropylphenyl; $E_{1/2}$, half-wave potential; Fc, ferrocene; THF, tetrahydrofuran; TMSCl, trimethylsilyl chloride.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig. 3 | Recent advances in two-electron reductive coupling reactions proceeding through titanacyclopentane and titanacyclopentene intermediates from $Ti(O^iPr)_4/RM$ and related species.

a | Catalytic Kulinkovich cyclopropanation reactions, in which ester C = O bond insertion into the Ti bond precedes pericyclic metalacyclopentane collapse. **b** | Stoichiometric arrested Kulinkovich-de Meijere-Like cyclopropanations, in which metalacyclopentane collapse is prevented through steric (top) or electronic (bottom) blocking. **c** | Alkoxide-directed regioselective and stereospecific reductive couplings that bias regioselective metalacyclopentane formation and subsequent insertion steps through covalent Ti-O bonding. DCE,


dichloroethane; DME, dimethoxyethane; equiv., equivalents; THF, tetrahydrofuran; TMS, trimethylsilyl.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



1,2- dichloroethane; DME, 1,2-dimethoxyethane; equiv., equivalents; THF, tetrahydrofuran; TM, transition metal; TMS, trimethylsilyl

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig. 6 | Single-electron processes catalysed by Ti^{III} complexes.

a | Catalytic epoxide ring opening using weak reductants and acids to drive turnover (Left) and examples in selective organic synthesis. **b** | Single-electron umpolung reactions can yield new C-C bonded products through a complex bimetallic radical coupling mechanism. **c** | SingLe-eLectron transfer (SET) from Ti^{III} into carbonyls adjacent to strained rings gives access to several new α lKene coupling reactions including cyclization, Michael addition and [3+2] cycLoadditions. COLL-HCL, 2,4,6-coLLidine hydrochloride; equiv, equivalents; e.r.,

enantiomeric ratio; EWG, eLectron-withdrawing group; TBDPS, tert-butyl-diphenylsilyl; THF, tetrahydrofuran; TMS, trimethylsilyl.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig. 7 | Single-electron strategies in small-molecule activation chemistry.

a | Hydrogenolysis of metal amides to ammonia via proton-coupled electron transfer (PCET), enabled by coordination-induced weakening of the N-H bond. **b** | Photochemical model of H₂O splitting with Ti^{III} catalysts, again enabled by coordination-induced weakening of O-H bonds. **c** | Photoredox catalysis via ligand-to-metal charge transfer (LMCT) with (MePDP)₂Zr complexes, in which charge transfer occurs in a direction opposite of state-of-the-art late transition metal photoredox catalysts, which undergo metal-to-ligand charge transfer (MLCT). *hν*, irradiation; BDFE, bond dissociation free energy;

Cp*, C₅Me₅⁻; E_{pc}, cathodic peak potential; equiv, equivalents; E_{red}, reduction potential; Fc, ferrocene; LED, light-emitting diode; MePDP, 2,6-bis (5-methyl-3-phenyl-1H-pyrrol-2-yl)-pyridine; TEMPO, (2,2,6,6-tetramethylpiperidin-1-yl)oxy; THF, tetrahydrofuran; TMS, trimethylsilyl; TON, turnover number.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig. 8 | Group-transfer catalysis mediated by early transition metal complexes.

a | Redox non-innocent ligand frameworks enable nitrene transfer reactions by undergoing redox changes at the ligand instead of the metal. **b** | Ti^{II}/Ti^{IV} redox catalytic nitrene transfer in oxidative amination catalysis — a new mechanism for pyrrole formation (left) — and examples in selective synthesis (right). **c** | Nitrene transfer promoted by a π -loaded bis(imido)Nb catalyst, which renders the imido fragment more labile than other group 5 imido complexes. **d** | A closed synthetic cycle for ligand-based CO₂ reduction to CO that avoids the formation of strong M-O multiple bonds. BDI, 2,6 diisopropylbenzene- β

diketimate; dadi, diamide-dimine; equiv, equivalents; THF, tetrahydrofuran; TMS, trimethylsilyl; TON, turnover number.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript