



**Cochrane**  
**Library**

Cochrane Database of Systematic Reviews

## Aspirin (single dose) for perineal pain in the early postpartum period (Review)

Molakatalla S, Shepherd E, Grivell RM

Molakatalla S, Shepherd E, Grivell RM.

Aspirin (single dose) for perineal pain in the early postpartum period.

*Cochrane Database of Systematic Reviews* 2017, Issue 2. Art. No.: CD012129.

DOI: 10.1002/14651858.CD012129.pub2.

[www.cochranelibrary.com](http://www.cochranelibrary.com)

## TABLE OF CONTENTS

HEADER . . . . .	1
ABSTRACT . . . . .	1
PLAIN LANGUAGE SUMMARY . . . . .	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON . . . . .	4
BACKGROUND . . . . .	6
OBJECTIVES . . . . .	7
METHODS . . . . .	7
Figure 1. . . . .	10
RESULTS . . . . .	13
Figure 2. . . . .	16
Figure 3. . . . .	17
Figure 4. . . . .	19
Figure 5. . . . .	20
Figure 6. . . . .	21
DISCUSSION . . . . .	23
AUTHORS' CONCLUSIONS . . . . .	25
ACKNOWLEDGEMENTS . . . . .	25
REFERENCES . . . . .	26
CHARACTERISTICS OF STUDIES . . . . .	29
DATA AND ANALYSES . . . . .	57

[Intervention Review]

# Aspirin (single dose) for perineal pain in the early postpartum period

Sujana Molakatalla<sup>1</sup>, Emily Shepherd<sup>2</sup>, Rosalie M Grivell<sup>3</sup>

<sup>1</sup>Department of Obstetrics and Gynaecology, Flinders Medical Centre, Adelaide, Australia. <sup>2</sup>ARCH: Australian Research Centre for Health of Women and Babies, Robinson Research Institute, Discipline of Obstetrics and Gynaecology, The University of Adelaide, Adelaide, Australia. <sup>3</sup>Department of Obstetrics and Gynaecology, Flinders University and Flinders Medical Centre, Bedford Park, Australia

Contact address: Sujana Molakatalla, Department of Obstetrics and Gynaecology, Flinders Medical Centre, Flinders Drive, Bedford Park, Adelaide, 5043, Australia. [drmsujana@gmail.com](mailto:drmsujana@gmail.com).

**Editorial group:** Cochrane Pregnancy and Childbirth Group.

**Publication status and date:** New, published in Issue 2, 2017.

**Citation:** Molakatalla S, Shepherd E, Grivell RM. Aspirin (single dose) for perineal pain in the early postpartum period. *Cochrane Database of Systematic Reviews* 2017, Issue 2. Art. No.: CD012129. DOI: 10.1002/14651858.CD012129.pub2.

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

## ABSTRACT

### Background

Perineal trauma (due to spontaneous tears, surgical incision (episiotomy) or in association with operative vaginal birth) is common after vaginal birth, and is often associated with postpartum perineal pain. Birth over an intact perineum may also lead to perineal pain. There are adverse health consequences associated with perineal pain for the women and their babies in the short- and long-term, and the pain may interfere with newborn care and the establishment of breastfeeding. Aspirin has been used in the management of postpartum perineal pain and its effectiveness and safety should be assessed.

### Objectives

To determine the efficacy of a single dose of aspirin (acetylsalicylic acid), including at different doses, in the relief of acute postpartum perineal pain.

### Search methods

We searched Cochrane Pregnancy and Childbirth's Trials Register (30 August 2016), [ClinicalTrials.gov](http://ClinicalTrials.gov), the WHO International Clinical Trials Registry Platform (ICTRP) (31 May 2016) and reference lists of retrieved studies.

### Selection criteria

Randomised controlled trials (RCTs) assessing single dose aspirin compared with placebo, no treatment, a different dose of aspirin, or single dose paracetamol/acetaminophen for women with perineal pain in the early postpartum period. We planned to include cluster-RCTs but none were identified. Quasi-RCTs and cross-over studies were not eligible for inclusion in this review.

### Data collection and analysis

Two review authors independently assessed study eligibility, extracted data and assessed the risk of bias of the included RCTs. Data were checked for accuracy. The quality of the evidence for the main comparison (aspirin versus placebo) was assessed using the GRADE approach.

## Main results

We included 17 RCTs, with 16 involving 1132 women randomised to aspirin or placebo (one RCT did not report numbers of women). Two RCTs (of 16) did not contribute data to review meta-analyses. All women had perineal pain post-episiotomy, and were not breastfeeding. Studies were published between 1967 and 1997, and the risk of bias was often unclear due to poor reporting.

We included four comparisons: aspirin versus placebo (data from 15 RCTs); 300 mg versus 600 mg aspirin (1 RCT); 600 mg versus 1200 mg aspirin (2 RCTs); and 300 mg versus 1200 mg aspirin (1 RCT).

## Primary outcomes

### *Aspirin versus placebo*

More women who received aspirin experienced adequate pain relief compared with women who received placebo over four to eight hours after administration (risk ratio (RR) 2.03, 95% confidence intervals (CI) 1.69 to 2.42; 13 RCTs, 1001 women; low-quality evidence). Women who received aspirin were less likely to need additional pain relief over four to eight hours after administration (RR 0.25, 95% CI 0.17 to 0.37; 10 RCTs, 744 women; very low-quality evidence). There was no difference in maternal adverse effects over four to eight hours post-administration (RR 1.08, 95% CI 0.57 to 2.06; 14 RCTs, 1067 women; very low-quality evidence). Subgroup analyses based on dose did not reveal any clear subgroup differences.

There was no clear difference over four hours after administration between 300 mg and 600 mg aspirin for adequate pain relief (RR 0.82, 95% CI 0.36 to 1.86; 1 RCT, 81 women) or need for additional pain relief (RR 0.68, 95% CI 0.12 to 3.88; 1 RCT, 81 women). There were no maternal adverse effects in either aspirin group.

There was no clear difference over four to eight hours after administration between 600 mg and 1200 mg aspirin for adequate pain relief (RR 0.85, 95% CI 0.52 to 1.39; 2 RCTs, 121 women), need for additional pain relief (RR 1.32, 95% CI 0.30 to 5.68; 2 RCTs, 121 women), or maternal adverse effects (RR 3.00, 95% CI 0.13 to 69.52; 2 RCTs, 121 women).

There was no clear difference over four hours after administration between 300 mg and 1200 mg aspirin for adequate pain relief (RR 0.62, 95% CI 0.29 to 1.32; 1 RCT, 80 women) or need for additional pain relief (RR 2.00, 95% CI 0.19 to 21.18; 1 RCT, 80 women). There were no maternal adverse effects in either aspirin group.

None of the included RCTs reported on neonatal adverse effects.

## Secondary outcomes

No studies reported on secondary review outcomes: prolonged hospitalisation due to perineal pain; re-hospitalisation due to perineal pain; fully breastfeeding at discharge; mixed feeding at discharge; fully breastfeeding at six weeks; mixed feeding at six weeks; perineal pain at six weeks; maternal views; maternal postpartum depression.

## Authors' conclusions

We found low-quality evidence to suggest that single dose aspirin compared with placebo can increase pain relief in women with perineal pain post-episiotomy. Very low-quality evidence also suggested that aspirin can reduce the need for additional analgesia, without increasing maternal adverse effects. Evidence was downgraded based on study limitations (risk of bias), imprecision, and publication bias or both. RCTs excluded breastfeeding women so there is no evidence to assess the effects of aspirin on neonatal adverse effects or breastfeeding.

With international guidance recommending mothers initiate breastfeeding within one hour of birth, and exclusively breastfeed for the first six months, the evidence from this review is not applicable to current recommended best practice. Aspirin may be considered for use in non-breastfeeding women with post-episiotomy perineal pain. Although formal assessment was beyond the remit of this review, current guidance suggests that other analgesic drugs (including paracetamol) should be considered first for postpartum perineal pain. Such agents are the focus of other reviews in this series on drugs for perineal pain in the early postpartum period. It is considered most likely that if RCTs are conducted in the future they could compare aspirin with other pain relievers. Future RCTs should be designed to ensure high methodological quality, and address gaps in the evidence, such as the secondary outcomes established for this review. Current research has focused on women with post-episiotomy pain, future RCTs could be extended to women with perineal pain associated with spontaneous tears or operative birth.

## PLAIN LANGUAGE SUMMARY

## **Aspirin (single dose) for relief of perineal pain after childbirth**

### **What is the issue?**

Can aspirin be given to women who experience perineal pain following childbirth to relieve the pain without causing side effects for either women or their babies?

### **Why is this important?**

Many women experience pain in the perineum (the area between the vagina and anus) following childbirth. The perineum may be bruised or torn during childbirth, or a cut made for the baby to be born (an episiotomy). After childbirth, perineal pain can interfere with women's ability to care for their newborns and establish breastfeeding. If perineal pain is not relieved effectively, longer-term problems for women may include painful sexual intercourse, pelvic floor problems resulting in incontinence, prolapse or chronic perineal pain. Aspirin may be given to women who have perineal pain after childbirth, but its effectiveness and safety had not been assessed in a systematic review. This is part of a series of reviews looking at drugs to help relieve perineal pain in first few weeks after childbirth.

### **What evidence did we find?**

We searched for evidence on 31 May 2016, and included 17 studies involving 1132 women published between 1967 and 1997. All women had perineal pain following an episiotomy (usually within 48 hours after birth) and were not breastfeeding. The women received either aspirin (doses ranging from 300 mg to 1200 mg) or fake pills (placebo), by mouth. The studies were assessed as low- or very low-quality evidence. Two trials did not contribute any data for analyses.

More women had adequate pain relief at four to eight hours after taking aspirin compared with women who received placebo (low-quality evidence). Women were less likely to need additional pain relief at four to eight hours after taking aspirin (very low-quality evidence). There was no difference in adverse effects for women in the four to eight hours after administration (very low-quality evidence).

We found no clear differences in effect for women who received 300 mg versus 600 mg aspirin (1 trial), 600 mg versus 1200 mg aspirin (2 trials), or 300 mg versus 1200 mg (1 trial) for adequate pain relief, the need for additional pain relief, or adverse effects for the mother.

No studies reported on adverse effects of aspirin for the baby, or other outcomes we planned to assess: prolonged hospital stay or readmission to hospital due to perineal pain; perineal pain six weeks after childbirth, women's views, or postpartum depression.

### **What does this mean?**

A single dose of aspirin may help with perineal pain following episiotomy for women who are not breastfeeding, when measured four to eight hours after administration. Breast milk is widely accepted as the best food for infants and it is recommended that where possible, mothers start breastfeeding within one hour of birth, and breastfeed for the first six months of their infant's life. We found no information to assess the effects of aspirin for women who are breastfeeding, but it is known that aspirin can be transferred into the breast milk.

## SUMMARY OF FINDINGS FOR THE MAIN COMPARISON *[Explanation]*

Aspirin compared with placebo for perineal pain in the early postpartum period						
<b>Patient or population:</b> women with perineal pain in the early postpartum period <b>Settings:</b> 17 RCTs published from 1967 to 1997 (11 RCTs conducted in USA, 3 in Venezuela, 1 each in Belgium, Canada and India) <b>Intervention:</b> aspirin (single dose) <b>Comparison:</b> placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Aspirin				
Adequate pain relief as reported by the woman (4 to 8 hours)	Study population		RR 2.03 (1.69, 2.42)	1001 (13 RCTs)	⊕⊕○○ low <sup>1</sup>	
	253 per 1000	513 per 1000 (427 to 612)				
Need for additional pain relief (4 to 8 hours)	Study population		RR 0.25 (0.17, 0.37)	744 (10 RCTs)	⊕○○○ very low <sup>1,2</sup>	
	267 per 1000	67 per 1000 (45 to 99)				
Maternal adverse effects (4 to 8 hours)	Study population		RR 1.08 (0.57, 2.06)	1067 (14 RCTs)	⊕○○○ very low <sup>1,3</sup>	
	27 per 1000	29 per 1000 (15 to 55)				
Neonatal adverse effects				(0 RCTs)		Not reported by any of the included RCTs
Perineal pain at six weeks postpartum				(0 RCTs)		Not reported by any of the included RCTs

\*The **corresponding risk** (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).  
**CI:** confidence interval; **RCT:** randomised controlled trial; **RR:** risk ratio

GRADE Working Group grades of evidence

**High quality:** Further research is very unlikely to change our confidence in the estimate of effect.

**Moderate quality:** Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

**Low quality:** Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

**Very low quality:** We are very uncertain about the estimate.

<sup>1</sup>Study limitations: downgraded two levels due to the serious risk of bias

<sup>2</sup>Publication bias: downgraded by one level based on visual inspection of funnel plot which indicates likely publication bias

<sup>3</sup>Imprecision: downgraded one level due to few events and wide 95% CI around the pooled estimate which includes no effect

## BACKGROUND

### Description of the condition

Perineal trauma may result from naturally occurring tears, surgical incisions such as episiotomy (cutting of the perineum to enlarge the vaginal opening during the second stage of labour), or in association with operative vaginal births (vacuum or forceps assisted births); and is frequently associated with acute perineal pain in the immediate postpartum period (Chou 2009). Birth over an intact perineum is also often associated with acute postpartum perineal pain. Perineal trauma is common, for example, in high-income countries such as Australia, where perineal trauma may occur in up to 68% of vaginal births, in association with episiotomy (15.6%), naturally occurring tears (first to fourth degree lacerations) (49.8%), and laceration plus episiotomy (3.0%) (Li 2013). Short-term morbidities for the mother arising from perineal trauma may include bleeding, infection, haematoma, and acute postpartum perineal pain, which may also interfere with newborn care and the establishment of breastfeeding (Chou 2009; East 2012a). In the longer-term, women are at an increased risk of dyspareunia (painful sexual intercourse), pelvic floor problems, and chronic perineal pain (Chou 2009; East 2012a).

Various practices can impact on the extent of perineal trauma sustained during birth, and so can influence the degree of perineal pain experienced by the woman in the immediate postpartum period. Cochrane systematic reviews have shown that antenatal digital perineal massage (Beckmann 2013) and the use of warm compresses on the perineum during the second stage of labour (Aasheim 2011) to be effective in preventing perineal trauma and associated pain.

A variety of practices and agents have also been assessed for the relief of perineal pain in the immediate postpartum period. Cochrane reviews have reported finding limited evidence to support routine use of local cooling (such as with ice packs or cold gel packs) of the perineum (East 2012b), or the application of topical local anaesthetics to the perineum for postpartum perineal pain relief (Hedayati 2005). A Cochrane review found some support for the use of paracetamol to reduce postpartum perineal pain, and decrease the need for additional pain relief. However, the overall quality of included evidence was assessed as unclear, and adverse effects were not assessed (Chou 2013). Other practices and agents that have been systematically reviewed and shown to have varied effectiveness in relieving postpartum perineal pain include: methods or materials or both for suturing perineal tears or episiotomies, therapeutic ultrasound, and rectal analgesia (East 2012a; Hay-Smith 1998; Hedayati 2003; Kettle 2012). For example, in regard to perineal suturing after childbirth, a Cochrane review showed that continuous suturing techniques for perineal closure, compared with interrupted methods, are associated with less short-term pain; if the continuous techniques are used for all

layers (vagina, perineal muscles and skin), the reduction of pain has been reported to be even greater (Kettle 2012).

### Description of the intervention

The history of aspirin began thousands of years ago, with early uses of extracts from plants and herbs containing salicylates (Vane 2003). In the 1870s, it was demonstrated that salicin and salicylic acid from white willow bark could reduce fever, pain and inflammation in people with rheumatic fever (Maclagan 1879). The success of salicylic acid prompted the German pharmaceutical manufacturer, Bayer, to search for a derivative that was equally or more effective. Felix Hoffman, a young chemist at Bayer, motivated by his father's inability to take salicylic acid for his arthritis due to its adverse effects (particularly vomiting), found a way to acetylate the hydroxyl group on the benzene ring of salicylic acid to form acetylated salicylic acid (Vane 2003).

In the first decades of the 1900s, acetylsalicylic acid, or 'aspirin' was considered the supreme analgesic (pain reliever); for three quarters of the 20th century, its use was solely as an analgesic and antipyretic (fever reducing) agent (Vane 2003). In the 1970s and 1980s, as part of his Nobel Prize-winning work, Sir John Vane demonstrated that aspirin could inhibit the formation of prostaglandins, associated with pain, fever, and inflammation, providing a physiological rationale for the effectiveness of one of the world's most widely used medications. As part of this work, Vane also discovered prostacyclin, an important prostaglandin that plays a vital role in the process of blood coagulation. The potential for aspirin to prevent a range of serious, life-threatening conditions, including heart attacks and stroke, was recognised following this discovery (Smith 2014).

Aspirin is now considered to be one of the most effective and versatile medications in the world. It is commonly recommended to be taken in the lowest effective dose to avoid adverse effects secondary to higher doses. For example, low-dose aspirin (75 mg to 150 mg daily) has been shown to provide substantial benefit for preventing serious cardiovascular events (heart attacks, stroke and vascular death) in people with pre-existing cardiovascular disease or with a history of events (secondary prevention) (ATT Collaboration 2002); in primary prevention, for people without histories of events or previous disease, the value of low and high-dose aspirin (75 mg to 500 mg daily) remains uncertain (ATT Collaboration 2009). Furthermore, there is increasing evidence that aspirin may reduce the risk of some cancers, and certain pregnancy complications. Long-term low-dose aspirin (at least 75 mg daily) has been shown to reduce colorectal cancer incidence and mortality (Rothwell 2010); and low-dose aspirin is reported to have moderate benefits in preventing pre-eclampsia and its consequences (Duley 2007), with 75 mg daily recommended for pregnant women a high risk of developing the condition (WHO 2011).



## How the intervention might work

Perineal pain is transmitted primarily through the pudendal nerve, a somatic sensory and motor nerve that innervates the external genitalia, as well as bladder and rectum sphincters (Cunningham 2005). Although a detailed description of the mechanism of action and pharmacology of aspirin is beyond the scope of this review, basic concepts are outlined.

The mechanisms by which aspirin exerts its analgesic, anti-inflammatory, and antipyretic effects were discovered in the 1970s. Aspirin inhibits the activity of the cyclo-oxygenase (COX) enzymes (irreversible inhibition of COX-1 and modification of COX-2), which play important roles in inflammation and nociceptive processes (the encoding and processing in the nervous system of noxious stimuli), such as through the formation of prostaglandins and thromboxanes (Vane 2003).

Through inhibiting these key enzymes, it has also been demonstrated that aspirin can prevent the production of physiologically important prostaglandins and thromboxanes, including those that protect the stomach mucosa from damage by hydrochloric acid, and those that aggregate platelets when required (Vane 2003). It is through these mechanisms that aspirin has been shown to cause adverse effects such as gastrointestinal irritation and occult (hidden) blood loss (Derry 2012). The availability of alternative agents with improved tolerability has reduced the use of aspirin for pain relief over recent years, however in many parts of the world, where alternatives are not available or are more expensive, aspirin is still the most commonly used analgesic for many different pain conditions (Derry 2012; Vane 2003).

A Cochrane systematic review that included 67 trials (involving 5743 adults) that were assessed at moderate to high quality overall, confirmed single-dose aspirin (300 mg to 1200 mg) to be an effective analgesic for acute postoperative moderate to severe intensity pain (Derry 2012). Higher doses (900 mg to 1000 mg) were shown to be more effective, however, these doses were associated with increased adverse effects, including gastric irritation, nausea, vomiting, drowsiness and dizziness. The pain relief achieved with aspirin was very similar to paracetamol given at the same dose (Derry 2012). Derry 2012 excluded trials where pain was due to trauma, such as is often the case for women with acute perineal pain in the immediate postpartum period. It is considered plausible that aspirin may also be effective in relieving acute perineal pain in the early postpartum period after birth, and so it was important to review the efficacy of single-dose aspirin when used for this indication.

## Why it is important to do this review

Perineal trauma is common after vaginal birth, and frequently associated with acute postpartum perineal pain; birth over an intact perineum is also often associated with perineal pain. Perineal pain may be associated with adverse health consequences for the mother

and her baby in the short and long term, such as dyspareunia, pelvic floor problems, and chronic perineal pain, and may also interfere with newborn care, including the establishment of breastfeeding (Chou 2009; East 2012a).

There is currently a dearth of evidence on effective interventions to reduce acute perineal pain in the immediate postpartum period. Previous Cochrane reviews have assessed practices and agents including therapeutic ultrasound (Hay-Smith 1998), rectal analgesia (Hedayati 2003), local cooling (East 2012b), topical anaesthetics (Hedayati 2005), paracetamol (Chou 2013), and most recently non-steroidal anti-inflammatory agents (Wuytack 2016), for the relief of perineal pain in the postpartum period. These reviews have reported mixed results. It is therefore important to establish if aspirin may be effective in relieving perineal pain and improving health outcomes for mother and their babies. Because it is known that salicylate and salicylate metabolites, including aspirin, are excreted in breast milk (NIH 2015), there is potential for effects on breastfeeding babies. Adverse effects or harms for both mothers and their babies therefore also need to be assessed.

We assessed the clinical effectiveness and adverse effects of aspirin given to relieve perineal pain in the early postpartum period after childbirth. This review is one of a series of reviews of drugs for perineal pain in the early postpartum period, all based on the same generic protocol (Chou 2009). This protocol is published in the *Cochrane Library* and describes the methods that shaped the production of all the reviews on drugs for perineal pain. It is available for consultation for prospective reviews undertaken on future drugs that may be introduced for this population and indication.

## OBJECTIVES

To determine the effects of a single dose of aspirin (acetylsalicylic acid), including at different doses, in the relief of acute postpartum perineal pain.

## METHODS

### Criteria for considering studies for this review

#### Types of studies

We included randomised controlled trials and planned to include cluster-randomised controlled trials. Quasi-randomised controlled trials and cross-over trials were excluded. We planned to include studies published as abstracts only, as well as studies published in full-text form.

## Types of participants

All women with acute perineal pain in the early postpartum period after childbirth; defined as the first four weeks after giving birth or as defined by the authors of the studies.

## Types of interventions

Single administration of aspirin used to treat perineal pain due to spontaneous lacerations, episiotomy or birth over an intact perineum in the early postpartum period. We included studies in which aspirin was compared with a placebo or no treatment, and where different doses of aspirin (e.g. 75 mg, 300 mg, etc) administered as a single dose, were compared. We also planned to include studies where aspirin was compared with a single dose of paracetamol/acetaminophen for perineal pain in the early postpartum period.

## Types of outcome measures

### Primary outcomes

1. Adequate pain relief as reported by the woman.\*
2. Need for additional pain relief in the first 48 hours for perineal pain.
3. Maternal adverse effects, composite of any of the following: nausea, vomiting, sedation, constipation, diarrhoea, drowsiness, sleepiness, gastric discomfort, psychological impact.
4. Neonatal adverse effects, composite of any of the following: vomiting, sedation, constipation, diarrhoea, drowsiness, sleepiness.

\* As determined by more than 50% relief of pain, as stated by the woman or calculated using a formula; see [Data collection and analysis](#) for details.

### Secondary outcomes

1. Prolonged hospitalisation due to perineal pain.
2. Rehospitalisation due to perineal pain.
3. Fully breastfeeding at discharge.
4. Mixed feeding at discharge.
5. Fully breastfeeding at six weeks.
6. Mixed feeding at six weeks.
7. Perineal pain at six weeks.
8. Maternal views (using a validated questionnaire).
9. Maternal postpartum depression.

## Search methods for identification of studies

The methods section of this review is based on a standard template used by Cochrane Pregnancy and Childbirth.

## Electronic searches

We searched Cochrane Pregnancy and Childbirth's Trials Register by contacting their Information Specialist (30 August 2016).

The Register is a database containing over 22,000 reports of controlled trials in the field of pregnancy and childbirth. For full search methods used to populate Pregnancy and Childbirth's Trials Register including the detailed search strategies for CENTRAL, MEDLINE, Embase and CINAHL; the list of handsearched journals and conference proceedings, and the list of journals reviewed via the current awareness service, please follow this link to the editorial information about [Cochrane Pregnancy and Childbirth](#) in the Cochrane Library and select the 'Specialized Register' section from the options on the left side of the screen.

Briefly, Cochrane Pregnancy and Childbirth's Trials Register is maintained by their Information Specialist and contains trials identified from:

1. monthly searches of the Cochrane Central Register of Controlled Trials (CENTRAL);
2. weekly searches of MEDLINE (Ovid);
3. weekly searches of Embase (Ovid);
4. monthly searches of CINAHL (EBSCO);
5. handsearches of 30 journals and the proceedings of major conferences; and
6. weekly current awareness alerts for a further 44 journals plus monthly BioMed Central email alerts.

Search results are screened by two people and the full text of all relevant trial reports identified through the searching activities described above is reviewed. Based on the intervention described, each trial report is assigned a number that corresponds to a specific Pregnancy and Childbirth review topic (or topics), and is then added to the Register. The Information Specialist searches the Register for each review using this topic number rather than keywords. This results in a more specific search set which has been fully accounted for in the relevant review sections ([Included studies](#); [Excluded studies](#); [Studies awaiting classification](#)).

We also searched [ClinicalTrials.gov](#) and the WHO International Clinical Trials Registry Platform ([ICTRP](#)) for unpublished, planned and ongoing trial reports using the terms given in [Appendix 1](#) on 31 May 2016.

## Searching other resources

We searched for further studies in the reference lists of the studies identified.

We did not apply language or date restrictions.

## Data collection and analysis

### Assessment of pain

The number of participants achieving adequate pain relief was defined as one of the following.

1. The number of participants reporting “good” or “excellent” pain relief when asked about their level of pain relief four to six hours after receiving their allocated treatment (the data were extracted as dichotomous data).

2. The number of women who reported 50% pain relief, or greater.

3. The number of participants who achieved 50% pain relief, or greater, as calculated by using derived pain relief scores (TOTPAR (total pain relief) or SPID (summed pain intensity differences)) over four to six hours.

It is common to use categorical or visual analogue scales for pain intensity and to calculate the results for each participant, over periods of four or six hours, as SPID or TOTPAR (Moore 1996). From these categorical scales, it was possible to convert results into dichotomous data (the proportion of participants achieving at the least 50%, or greater, max TOTPAR) using standard formulae (Moore 1996; Moore 1997b). Converting data in this way enabled these data to be used in a meta-analysis (Moore 1997a; Moore 1997b). The following equations were used to estimate the proportions of participants achieving at least 50% of maximum TOTPAR.

Proportion with greater than 50% maxTOTPAR =  $(1.33 \times \text{mean \%maxTOTPAR} - 11.5)$

With  $\% \text{maxTOTPAR} = \text{mean TOTPAR} \times 100 / (\text{maximum score} \times \text{number of hours})$

(Cooper 1991; Moore 1997b)

Proportion with greater than 50% maxTOTPAR =  $(1.36 \times \text{mean$

$\% \text{maxSPID} - 2.3)$

With  $\% \text{maxSPID} = \text{mean SPID} \times 100 / (\text{maximum score} \times \text{number of hours})$

(Cooper 1991; Moore 1997a)

The number of participants achieving at least 50% maxTOTPAR was then calculated by multiplying the proportions of participants with at least 50% maxTOTPAR by the total number of participants in the treatment groups. The number of participants with at least 50% maxTOTPAR was then used to calculate the relative benefit and number needed to treat to benefit.


Where studies used more than one method of calculating adequate pain relief, preference for analyses and reporting purposes, in order of decreasing preference, was: i) the proportion with at least 50% maxTOTPAR calculated using SPID; ii) the proportion with at least 50% maxTOTPAR calculated using TOTPAR; and iii) the number of participants reporting ‘good’ or ‘excellent’ pain relief/ number of participants reporting at least 50% pain relief. We also assessed the number of participants who re-medicated in the period of four to eight hours, as well as the median time to re-medication, if data were available.

### Selection of studies

Two review authors independently assessed for inclusion all the potential studies we identified as a result of the search strategy. We resolved any disagreement through discussion or, if required, we consulted a third review author.

We created a study flow diagram to illustrate numbers of records identified, included and excluded (Figure 1).

Figure 1. Study flow diagram



## Data extraction and management

We designed a form to extract data. At least two review authors extracted data using the agreed form for eligible studies. We resolved discrepancies through discussion or, if required, consultation with another member of the review author team. We entered data into Review Manager software (RevMan 2014) and checked for accuracy. When information regarding any steps was unclear, we attempted to contact authors of the original reports to provide further details.

## Assessment of risk of bias in included studies

Two review authors independently assessed risk of bias for each study using the criteria outlined in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). We resolved any disagreement by discussion or by involving a third assessor.

### (1) Random sequence generation (checking for possible selection bias)

We described for each included study the method used to generate the allocation sequence in sufficient detail to allow an assessment of whether it should produce comparable groups.

We assessed the method as:

- low risk of bias (any truly random process, e.g. random number table; computer random number generator);
- high risk of bias (any non-random process, e.g. odd or even date of birth; hospital or clinic record number);
- unclear risk of bias.

### (2) Allocation concealment (checking for possible selection bias)

We described for each included study the method used to conceal allocation to interventions prior to assignment and assessed whether intervention allocation could have been foreseen in advance of, or during recruitment, or changed after assignment.

We assessed the methods as:

- low risk of bias (e.g. telephone or central randomisation; consecutively numbered sealed opaque envelopes);
- high risk of bias (open random allocation; unsealed or non-opaque envelopes, alternation; date of birth);
- unclear risk of bias.

### (3.1) Blinding of participants and personnel (checking for possible performance bias)

We described for each included study the methods used, if any, to blind study participants and personnel from knowledge of which

intervention a participant received. We considered that studies were at low risk of bias if they were blinded, or if we judged that the lack of blinding would be unlikely to affect results.

We assessed the methods as:

- low, high or unclear risk of bias for participants;
- low, high or unclear risk of bias for personnel.

### (3.2) Blinding of outcome assessment (checking for possible detection bias)

We described for each included study the methods used, if any, to blind outcome assessors from knowledge of which intervention a participant received.

We assessed methods used to blind outcome assessment as:

- low, high or unclear risk of bias.

### (4) Incomplete outcome data (checking for possible attrition bias due to the amount, nature and handling of incomplete outcome data)

We described for each included study, and for each outcome or class of outcomes, the completeness of data including attrition and exclusions from the analysis. We stated whether attrition and exclusions were reported and the numbers included in the analysis at each stage (compared with the total randomised participants), reasons for attrition or exclusion where reported, and whether missing data were balanced across groups or were related to outcomes. Where sufficient information was reported, or could be supplied by the trial authors, we planned to re-include missing data in the analyses.

We assessed methods as:

- low risk of bias (e.g. no missing outcome data; missing outcome data balanced across groups);
- high risk of bias (e.g. numbers or reasons for missing data imbalanced across groups; 'as treated' analysis done with substantial departure of intervention received from that assigned at randomisation);
- unclear risk of bias.

### (5) Selective reporting (checking for reporting bias)

We described for each included study how we investigated the possibility of selective outcome reporting bias and what we found. We assessed the methods as:

- low risk of bias (where it was clear that all of the study's pre-specified outcomes and all expected outcomes of interest to the review were reported);
- high risk of bias (where not all the study's pre-specified outcomes were reported; one or more reported primary

outcomes were not pre-specified; outcomes of interest were reported incompletely and so could not be used; study failed to include results of a key outcome that would have been expected to have been reported);

- unclear risk of bias.

#### **(6) Other bias (checking for bias due to problems not covered by points (1) to (5))**

We described for each included study any important concerns we had about other possible sources of bias including: was the trial stopped early due to some data-dependent process? Was there extreme baseline imbalance? Has the study been claimed to be fraudulent?

We assessed whether each study was free of other problems that could put it at risk of bias:

- low risk of other bias;
- high risk of other bias;
- unclear whether there was risk of other bias.

#### **(7) Overall risk of bias**

We made explicit judgements about whether studies were at high risk of bias, according to the criteria given in the *Handbook* (Higgins 2011). With reference to points (1) to (6), we assessed the likely magnitude and direction of the bias and whether we considered it is likely to impact on the findings. We planned to assess the impact of the level of bias through undertaking sensitivity analyses - see [Sensitivity analysis](#).

### **Assessing the quality of the evidence using the GRADE approach**

We planned to assess the quality of the evidence using the GRADE approach as outlined in the [GRADE handbook](#) in order to assess the quality of the body of evidence relating to the following outcomes for the main comparison: aspirin versus placebo.

1. Adequate pain relief as reported by the woman.
2. Need for additional pain relief in the first 48 hours for perineal pain.
3. Maternal adverse effects, composite of any of the following: nausea, vomiting, sedation, constipation, diarrhoea, drowsiness, sleepiness, gastric discomfort, psychological impact.
4. Neonatal adverse effects, composite of any of the following: vomiting, sedation, constipation, diarrhoea, drowsiness, sleepiness.
5. Perineal pain at six weeks postpartum.

We could however only assess the quality of the evidence for the first three outcomes, as for outcomes 4 and 5, we had no data from the included trials.

We created a 'Summary of findings' table for our main comparison in Review Manager 5.3 (RevMan 2014). A summary of the

intervention effect and a measure of quality for each of the outcomes was produced using the GRADE approach. The GRADE approach uses five considerations (study limitations, consistency of effect, imprecision, indirectness and publication bias) to assess the quality of the body of evidence for each outcome. The evidence can be downgraded from 'high quality' by one level for serious (or by two levels for very serious) limitations, depending on assessments for risk of bias, indirectness of evidence, serious inconsistency, imprecision of effect estimates or potential publication bias.

### **Measures of treatment effect**

#### **Dichotomous data**

We presented results as summary risk ratio with 95% confidence intervals for dichotomous data.

#### **Continuous data**

We planned to use the mean difference if outcomes were measured in the same way between trials. We planned to use the standardised mean difference to combine trials that measured the same outcome, but used different methods.

### **Unit of analysis issues**

#### **Cluster-randomised trials**

We planned to include cluster-randomised trials in the analyses along with individually-randomised trials. If we include cluster-randomised trials in future updates, we will adjust their sample sizes using the methods described in the *Handbook* (Higgins 2011) using an estimate of the intra cluster correlation co-efficient (ICC) derived from the trial (if possible), from a similar trial or from a study of a similar population. If we use ICCs from other sources, we will report this and conduct sensitivity analyses to investigate the effect of variation in the ICC. If we identify both cluster-randomised trials and individually-randomised trials, we plan to synthesise the relevant information. We will consider it reasonable to combine the results from both if there is little heterogeneity between the study designs and the interaction between the effect of intervention and the choice of randomisation unit is considered to be unlikely.

We will also acknowledge heterogeneity in the randomisation unit and perform a sensitivity analysis to investigate the effects of the randomisation unit.

### Cross-over trials

We considered cross-over trials to be inappropriate for this research question and these were not included.

### Multi-armed trials

We included all the relevant intervention groups (aspirin) and control groups (placebo) from multi-arm trials. We excluded other arms that were not relevant to this review.

### Dealing with missing data

We noted levels of attrition for the included studies. We planned to explore the impact of including studies with high levels of missing data in the overall assessment of treatment effect by conducting sensitivity analyses.

We carried out analyses, as far as possible, on an intention-to-treat basis for all outcomes. That is, we attempted to include all participants randomised to each group in the analyses, and all participants were analysed in the group to which they were allocated, regardless of whether or not they received the allocated intervention. The denominator for each outcome in each trial was the number randomised minus any participants whose outcomes were known to be missing.

### Assessment of heterogeneity

We assessed statistical heterogeneity in meta-analyses using the  $T^2$ ,  $I^2$  and  $\text{Chi}^2$  statistics. We regarded heterogeneity as substantial if an  $I^2$  was greater than 30% and either a  $T^2$  was greater than zero, or there was a low P value ( $P < 0.10$ ) in the  $\text{Chi}^2$  test for heterogeneity.

### Assessment of reporting biases

Because there were 10 or more studies included in the meta-analyses for 'Adequate pain relief as reported by the woman', 'Need for additional pain relief' and 'Maternal adverse effects', we investigated reporting biases (such as publication bias) using funnel plots. We assessed funnel plot asymmetry visually.

### Data synthesis

We carried out statistical analysis using Review Manager software (RevMan 2014). We used fixed-effect meta-analysis for combining data because it was considered reasonable to assume that studies were estimating the same underlying treatment effect.

In future updates of this review, if there is clinical heterogeneity sufficient to expect that the underlying treatment effects differed between trials, or where substantial statistical heterogeneity is detected, we will use random-effects meta-analysis to produce an overall summary. We will treat the random-effects summary as the

average of the range of possible treatment effects and discuss the clinical implications of treatment effects differing among trials. If the average treatment effect is not clinically meaningful, we will not combine trials. If we use random-effects analyses in future updates, the results will be presented as the average treatment effect with 95% confidence intervals, and the estimates of  $\text{Tau}^2$  and  $I^2$ .

### Subgroup analysis and investigation of heterogeneity

If we had identified substantial heterogeneity, we planned to investigate possible sources using subgroup analyses and sensitivity analyses. We planned to consider whether an overall summary was meaningful, and if it was, use random-effects analysis to produce the effect.

We planned to carry out the following subgroup analyses:

1. primiparous versus multiparous women;
2. women with perineal trauma versus women who gave birth over intact perineum; and
3. dose of aspirin (i.e. low-dose versus high-dose).

However due to the absence of relevant data in the included trials, we were able to conduct analyses based on dose only.

Subgroup analysis were restricted to the review's primary outcomes with reported data.

We assessed subgroup differences by interaction tests available in RevMan 2014. We reported the results of subgroup analyses quoting the  $\text{Chi}^2$  statistic and P value, and the interaction test  $I^2$  value.

### Sensitivity analysis

We planned to conduct sensitivity analyses to explore the effects of trial quality on the outcomes. We planned to explore the effects of trial quality assessed by allocation concealment and random sequence generation (considering selection bias), by omitting studies rated as high or unclear risk of bias for these components. However, because all included trials were assessed with an unclear rating for at least one of these two components, we did not conduct sensitivity analyses.

We also planned to investigate the effects of the randomisation unit (individual versus cluster) on the outcomes, and the impact of including studies with high levels of missing data. We planned to explore the effects of fixed-effect or random-effects analyses for outcomes with statistical heterogeneity, and the effects of any assumptions made such as the value of the ICC used for cluster-randomised trials. However, because we did not include any cluster-randomised trials, trials with high levels of missing data, or identify outcomes with substantial statistical heterogeneity, we did not conduct sensitivity analyses.

We planned to use only primary outcomes in sensitivity analyses.

## RESULTS

## Description of studies

### Results of the search

The search of the Cochrane Pregnancy and Childbirth's Trials Register retrieved 35 reports. The search of additional sources yielded seven records (ClinicalTrials.gov = five, ICTRP = two records), all of which were screened and considered to be ineligible based on title, abstract or both. This resulted in screening 35 full text reports. Of these, 17 studies (22 records) were included, and 10 studies (11 records) were excluded.

Two studies (two records) await classification: the method of allocation was not clearly reported in [Bhounsule 1990](#); and we were unable to locate a copy of [Sunshine 1989](#).

See [Characteristics of included studies](#); [Characteristics of excluded studies](#); [Characteristics of studies awaiting classification](#); [Figure 1](#).

### Included studies

#### Design and setting

We included 17 studies (22 reports) in this review. All were reported to be randomised controlled trials (RCTs).

The included studies were published between 1967 and 1997; one trial was published in the 1960s; six in the 1970s; nine in the 1980s; and one in the 1990s. Most (11 trials) were conducted in the USA ([Bloomfield 1967](#); [Bloomfield 1970a](#); [Bloomfield 1970b](#); [Bloomfield 1974](#); [Friedrich 1983](#); [Jain 1978a](#); [Jain 1978b](#); [Jain 1985](#); [London 1983a](#); [London 1983b](#); [Okun 1982](#)), three in Venezuela ([Olson 1997](#); [Sunshine 1983a](#); [Sunshine 1983b](#)), and one each in Belgium ([De Vroey 1978](#)), Canada ([Trop 1983](#)) and India ([Mukherjee 1980](#)).

Only one trial ([Bloomfield 1970b](#)) had two trial arms, comparing aspirin and placebo. Another trial compared only aspirin and placebo, but had four trial arms ([London 1983b](#)). The remaining 15 trials had between three and five trial arms, and in addition to aspirin, assessed a number of other agents for perineal pain in the early postpartum period. These agents included chlorphenesin 400 mg, 800 mg and combination aspirin 300 mg and chlorphenesin 400 mg ([Bloomfield 1967](#)); flufenisal 300 mg and 600 mg ([Bloomfield 1970a](#)); ibuprofen 300 mg and 900 mg ([Bloomfield 1974](#)); diflunisal 125 mg, 250 mg and 500 mg ([De Vroey 1978](#)), etodolac 25 mg and 100 mg ([Friedrich 1983](#)), piroxicam 20 mg and 40 mg ([Jain 1978a](#)); combination aspirin 800 mg and caffeine 64 mg ([Jain 1978b](#)); indoprofen 50 and 100 mg ([Jain 1985](#)); fluproquazone 100 mg and 200 mg ([London 1983a](#)); dipyron 500 mg ([Mukherjee 1980](#)); fendosal 100 mg, 200 mg and 400 mg ([Okun 1982](#)); potassium 25 mg, 50 mg and 100 mg ([Olson 1997](#)); zomepirac and ibuprofen ([Sunshine 1983a](#)); flurbiprofen 25 mg 50 mg and 100 mg ([Sunshine 1983b](#)); and tiaprofenic acid 200 and 400 mg ([Trop 1983](#)). For the purposes of the review,

we analysed only the aspirin and placebo arms from the included trials.

### Participants and sample sizes

In total, there were 1132 women in the aspirin and placebo arms of 16 of the 17 included trials, with 617 women randomised to receive aspirin, and 515 to a placebo. In three trials, only the numbers analysed (not randomised) were reported ([De Vroey 1978](#); [London 1983a](#); [London 1983b](#)), and in one trial ([Trop 1983](#)), the numbers of women were not reported at all. The sample sizes of the trials (including only the relevant arms) ranged from 26 ([Bloomfield 1970b](#)) to 178 ([Mukherjee 1980](#)). We reported the number of women in other arms of the trials not included in analyses in the [Characteristics of included studies](#) tables.

All included trials included women with perineal pain in the early postpartum period post-episiotomy. One trial recruited and randomised women on the first post-operative morning ([Mukherjee 1980](#)), two within 24 hours of birth ([Bloomfield 1967](#); [Bloomfield 1970b](#)), one from 16 to 48 hours following induction of anaesthesia ([Friedrich 1983](#)), six within 48 hours of birth ([Bloomfield 1970a](#); [Bloomfield 1974](#); [De Vroey 1978](#); [Jain 1978a](#); [London 1983a](#); [Okun 1982](#)); and seven trials did not specify a time period following birth ([Jain 1978a](#); [Jain 1985](#); [London 1983b](#); [Olson 1997](#); [Sunshine 1983a](#); [Sunshine 1983b](#); [Trop 1983](#)). No trials were identified that assessed perineal pain associated with naturally occurring tears or birth over an intact perineum. The intensity of women's pain following episiotomy varied among the included trials; eight trials included women with moderate or severe pain ([Bloomfield 1967](#); [De Vroey 1978](#); [Friedrich 1983](#); [Jain 1978a](#); [London 1983a](#); [London 1983b](#); [Mukherjee 1980](#); [Sunshine 1983b](#)); three included women with moderate to very severe pain ([Bloomfield 1970a](#); [Bloomfield 1974](#); [Okun 1982](#)); one included women with mild to severe pain ([Bloomfield 1970b](#)); one included women with at least moderate pain ([Jain 1985](#)); and three included women with severe pain ([Jain 1978b](#); [Olson 1997](#); [Sunshine 1983a](#)). Pain intensity was not specified in one trial ([Trop 1983](#)). Most trials clearly specified that breastfeeding was an exclusion criterion, and excluded women with known sensitivity or allergy to aspirin, and women who had recently received analgesia.

### Interventions and comparisons

Of the 17 included trials, 15 included additional arms that assessed other agents. However, we included data from aspirin and placebo arms only in this review. Fifteen trials included comparisons of aspirin and placebo only; the single, oral doses of aspirin in these were 500 mg ([Mukherjee 1980](#)), 600 mg ([Bloomfield 1967](#); [De Vroey 1978](#); [Jain 1985](#); [Sunshine 1983a](#); [Sunshine 1983b](#)), 648 mg ([Jain 1978a](#)), 650 mg ([Friedrich 1983](#); [Jain 1978b](#); [London 1983a](#); [Okun 1982](#); [Olson 1997](#)), 900 mg ([Bloomfield 1974](#)), and 1200 mg ([Bloomfield 1970b](#)). Three trials included two or more



aspirin arms (in addition to a placebo arm); Bloomfield 1970a and Trop 1983 compared 600 mg and 1200 mg aspirin, and London 1983b compared 300 mg, 600 mg and 1200 mg aspirin. The number of aspirin and placebo tablets (and dose of the tablets) taken varied across the trials.

## Outcomes

Some measure of 'Adequate pain relief as reported by the woman' could be extracted four to eight hours after drug administration from 13 trials and meta-analysed. Data were not presented in a way that enabled inclusion in the meta-analysis in four trials (Jain 1978a; Jain 1978b; Okun 1982; Trop 1983).

Three trials in the meta-analysis provided data on adequate pain relief four hours after taking the medication (London 1983b; Olson 1997; Sunshine 1983a); two trials reported this outcome after five hours (Bloomfield 1970b; Jain 1985), seven trials after six hours (Bloomfield 1967; Bloomfield 1974; De Vroey 1978; Friedrich 1983; London 1983a; Mukherjee 1980; Sunshine 1983b) and one trial after eight hours (Bloomfield 1970a). SPID scores were used to calculate the number of women with adequate pain relief for the meta-analysis in 11 trials; (Bloomfield 1967; Bloomfield 1970a; Bloomfield 1970b; Bloomfield 1974; De Vroey 1978; Friedrich 1983; Jain 1985; London 1983b; Olson 1997; Sunshine 1983a; Sunshine 1983b), one used TOTPAR scores (Mukherjee 1980), and one used the number of women reporting pain relief to be good or excellent (London 1983a).

Five trials provided both SPID and TOTPAR scores (Friedrich 1983; Jain 1985; Olson 1997; Sunshine 1983a; Sunshine 1983b); two trials provided both SPID scores, and the number of women reporting pain relief to be good or excellent (Friedrich 1983; Jain 1985); and five reported SPID scores and the number of women with at least 50% pain relief (or similar) (Bloomfield 1970a; Bloomfield 1970b; Bloomfield 1974; De Vroey 1978; Mukherjee 1980). In these cases, we used SPID data to calculate the number of women with adequate pain relief for inclusion in the meta-analysis. In some cases, the number of women with adequate pain relief according to these different measures did not match, and the reasons for discrepancies were not entirely clear, particularly in numbers of women with adequate pain relief when calculated using the SPID versus TOTPAR scores.

Data on the need for additional analgesia which could be included in a meta-analysis were available from 10 trials (Bloomfield 1970a; Bloomfield 1974; De Vroey 1978; Jain 1978a; Jain 1985; London 1983a; London 1983b; Olson 1997; Sunshine 1983a; Sunshine 1983b) and 14 trials reported data on any maternal adverse effects suitable for meta-analysis (Bloomfield 1967; Bloomfield 1970a; Bloomfield 1974; De Vroey 1978; Friedrich 1983; Jain 1978a; Jain 1978b; Jain 1985; London 1983a; London 1983b; Mukherjee

1980; Olson 1997; Sunshine 1983a; Sunshine 1983b).

Two trials (Okun 1982; Trop 1983) did not provide any data that could be meta-analysed.

None of the 17 included trials reported on any of the prespecified secondary outcomes.

## Funding sources

Two of the trials reported support or funding solely from the National Institutes of Health (Bloomfield 1967), and the United States Public Health Service and National Heart Institute (Bloomfield 1970b). Ten of the trials reported at least partial support from pharmaceutical companies/commercial medical research organisations: Merck Sharp and Dohme Research Laboratories (Bloomfield 1970a; De Vroey 1978), the Upjohn Company (Bloomfield 1974; Sunshine 1983b), American Home Products, Ives Laboratories and Wyeth Laboratories (Jain 1978b), Adria Laboratories (Jain 1985), Sandoz Inc. (London 1983a), the Ciba-Geigy Corporation (Olson 1997), Boots Pharmaceuticals (Sunshine 1983b) and Roussel Canada Inc. (Trop 1983). Five of the trials (Friedrich 1983; Jain 1978a; London 1983b; Mukherjee 1980; Okun 1982) did not report any sources of support or funding.

## Declarations of interests

None of the 17 included trials provided specific declarations of interest for the manuscript authors. It was noted that three of the trials had author(s) with affiliations to pharmaceutical companies/commercial medical research organisations: Merck Sharp and Dohme Research Laboratories (De Vroey 1978); Analgesic Development Ltd. (Olson 1997), and Roussel Canada Inc. (Trop 1983).


## Excluded studies

We excluded 10 studies (11 records) for the following reasons: five trials (Bruni 1965; Gruber 1979; Moggian 1972; Sunshine 1983c; Sunshine 1985) included mixed populations of women with postpartum pain (such as uterine cramping), and no results were reported separately for women with perineal pain; one was not a randomised trial (Santiago 1959); three assessed combination agents (not aspirin alone) (Gindhart 1971; Prockop 1960; Rubin 1984); and one assessed twice daily aspirin (not single dose aspirin) (Van der Pas 1984).

## Risk of bias in included studies

See Figure 2 and Figure 3.

**Figure 2. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies**



**Figure 3. Risk of bias summary: review authors' judgements about each risk of bias item for each included study**

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Bloomfield 1967	?	?	+	+	-	?	+
Bloomfield 1970a	?	?	+	+	-	?	?
Bloomfield 1970b	?	?	+	+	+	?	?
Bloomfield 1974	?	?	+	+	-	?	+
De Vroey 1978	?	?	+	+	-	-	?
Friedrich 1983	?	?	?	?	?	?	?
Jain 1978a	?	?	?	?	?	-	+
Jain 1978b	?	?	?	?	?	-	?
Jain 1985	?	?	+	+	+	?	+
London 1983a	?	?	+	+	?	?	?
London 1983b	?	?	+	+	-	?	?
Mukherjee 1980	?	?	+	+	+	?	+
Okun 1982	?	?	+	+	-	?	+
Olson 1997	+	?	+	+	-	?	+
Sunshine 1983a	?	?	+	+	-	-	+
Sunshine 1983b	+	?	+	+	-	?	+
Trop 1983	?	?	+	+	?	-	?

## Allocation

Only two trials were assessed to have applied adequate sequence generation methods, both used computer-generated random sequences (Olson 1997; Sunshine 1983b). The remaining 15 trials did not report on methods used for random sequence generation, and simply reported that the women were 'randomised' (Bloomfield 1967; Bloomfield 1970a; Bloomfield 1970b; Bloomfield 1974; De Vroey 1978; Friedrich 1983; Jain 1978a; Jain 1978b; Jain 1985; London 1983a; London 1983b; Mukherjee 1980; Okun 1982; Sunshine 1983a; Trop 1983). All included trials were judged at unclear risk of selection bias, with none reporting methods of allocation concealment.

## Blinding

Of the 17 trials, 14 were judged at low risk of both performance and detection bias, with blinding of women and study personnel (who were also the outcome assessors) by using identical placebos (Bloomfield 1967; Bloomfield 1970a; Bloomfield 1970b; Bloomfield 1974; De Vroey 1978; Jain 1985; London 1983a; London 1983b; Mukherjee 1980; Okun 1982; Olson 1997; Sunshine 1983a; Sunshine 1983b; Trop 1983). Three trials (Friedrich 1983; Jain 1978a; Jain 1978b) were at unclear risk of performance and detection bias, because although the trials were reported to be 'double-blind', no information was provided on the nature of the placebos used to determine if blinding could have been successfully achieved.

## Incomplete outcome data

Only three trials (Bloomfield 1970b; Jain 1985; Mukherjee 1980) were judged to be at low risk of attrition bias, with no losses or exclusions.

Five trials were assessed at unclear risk of attrition bias (Friedrich 1983; Jain 1978a; Jain 1978b; London 1983a; Trop 1983), largely due to unclear reporting regarding any losses and exclusions, and reasons for missing data, or both.

Nine trials (Bloomfield 1967; Bloomfield 1970a; Bloomfield 1974; De Vroey 1978; London 1983b; Okun 1982; Olson 1997; Sunshine 1983a; Sunshine 1983b) were assessed at high risk of attrition bias. In eight, the trial authors imputed data (i.e. for women requesting additional analgesia, trial authors either used women's pre-treatment pain intensity/relief scores for all subsequent hours, or used the last observation carried forward method for subsequent hours), which may have introduced bias; in one trial (Bloomfield 1967), women who requested additional analgesia were excluded from the analyses, which may have similarly introduced bias.

## Selective reporting

We assessed 12 trials (Bloomfield 1967; Bloomfield 1970a; Bloomfield 1970b; Bloomfield 1974; Friedrich 1983; Jain 1985; London 1983a; London 1983b; Mukherjee 1980; Okun 1982; Olson 1997; Sunshine 1983b) at unclear risk of reporting bias, with no access to trial protocols or registrations to confidently assess the risk of selective reporting.

Five trials (De Vroey 1978; Jain 1978a; Jain 1978b; Sunshine 1983a; Trop 1983) were judged to be at high risk of reporting bias. In all five, some of outcome data and results were reported incompletely in the text, which meant these data could not have been extracted, such as for inclusion in a meta-analysis. For example: "The three drugs were much the same for mean onset, duration and time to peak values. The hypothesis that there is no difference among treatments was rejected at the 0.05 level or better for all variables" (Sunshine 1983a).

Very few outcome data were reported in all 17 trials.

## Other potential sources of bias

Nine trials (Bloomfield 1967; Bloomfield 1970a; Jain 1978a; Jain 1985; Mukherjee 1980; Okun 1982; Olson 1997; Sunshine 1983a; Sunshine 1983b) were assessed at low risk of other potential sources of bias. Eight trials (Bloomfield 1970b; Bloomfield 1974; De Vroey 1978; Friedrich 1983; Jain 1978b; London 1983a; London 1983b; Trop 1983) were judged to be at unclear risk of other bias. These trials did not report baseline characteristics in a way that enabled assessment of comparability among groups (with no baseline characteristics reported, or lack of detail reported); one trial (Bloomfield 1970a) reported that most baseline characteristics were similar between groups "However, body weight was not similar in all treatment groups".

## Effects of interventions

See: [Summary of findings for the main comparison Aspirin compared with placebo for perineal pain in the early postpartum period](#)

## Comparison 1: Aspirin versus placebo for perineal pain


Fifteen of the 17 included trials contributed data to meta-analyses in this comparison (Bloomfield 1967; Bloomfield 1970a; Bloomfield 1970b; Bloomfield 1974; De Vroey 1978; Friedrich 1983; Jain 1978a; Jain 1978b; Jain 1985; London 1983a; London 1983b; Mukherjee 1980; Olson 1997; Sunshine 1983a; Sunshine 1983b). Two trials (Okun 1982; Trop 1983) did not provide any data that could be meta-analysed.

## Primary outcomes

### Adequate pain relief as reported by the woman

Over four to eight hours after drug administration, more women who had received aspirin experienced adequate pain relief compared with women who received placebo (risk ratio (RR) 2.03, 95% confidence intervals (CI) 1.69 to 2.42; 13 trials, 1001 women; [Analysis 1.1](#)). The quality of the evidence (GRADE) for this outcome was judged to be low, with downgrading based on study limitations (risk of bias) ([Summary of findings for the main comparison](#)). Visual inspection of the funnel plot for this outcome suggested no clear evidence of reporting bias ([Figure 4](#)).

**Figure 4. Funnel plot of comparison: I Aspirin versus placebo for perineal pain, outcome: 1.1 Adequate pain relief as reported by the woman**



### Data from the trials not included in the meta-analysis

- [Jain 1978a](#): “By all measurements of drug effect... aspirin 648 mg [was] significantly ( $P < 0.01$ ) superior to placebo in [its]

overall analgesic effect and also at second, third and fourth hours after dosing”.

- [Jain 1978b](#): “In comparing 650 mg aspirin with placebo,


we detected no significant difference at 1, 2, or 3 hr, but at the fourth hour we noted trends toward significant in favour of aspirin ( $P < 0.10$ ) by each Kruskal-Wallis analysis for pain analogue, pain intensity, and pain relief scores. The corresponding analysis of covariance at hour 4 showed differences in favour of aspirin for both pain analogue and pain intensity scores ( $P < 0.02$ )”.

- **Okun 1982**: “In patients with either uterine cramp or episiotomy pain, aspirin... provided greater pain relief (lower mean pain intensity scores) than did placebo from the 2nd through the 8th study hour”.
- **Trop 1983**: “When compared to placebo both patient’s self-rating scale and nurse’s impression scale have shown a significant reduction in pain following treatment... with ASA”.

### Need for additional pain relief in the first 48 hours for perineal pain

Women who received aspirin were less likely to require additional analgesia over four to eight hours (RR 0.25, 95% CI 0.17 to 0.37; 10 trials, 744 women; **Analysis 1.2**) after drug administration. The quality of the evidence (GRADE) for this outcome was judged to be very low, with downgrading based on study limitations (risk of bias), and possible publication bias (**Summary of findings for the main comparison**). Visual inspection of the funnel plot for this outcome indicated possible evidence of reporting bias, which could be due to some smaller trials producing exaggerated intervention effect estimates (**Figure 5**).

**Figure 5. Funnel plot of comparison: I Aspirin versus placebo for perineal pain, outcome: I.2 Need for additional pain relief**



#### Data from the trials not included in the meta-analysis

- **Bloomfield 1967**: not reported; although one woman in the aspirin group was reported to have been “withdrawn owing to

distressing pain unrelieved by the study drugs” compared with no women in the placebo group.


- Okun 1982: “The proportion of patients requiring additional analgesic was significantly different... Approximately 71% of patients in the placebo group needed additional analgesic as compared with... 48% in the aspirin group” (these data related to women with uterine cramp or episiotomy pain).

- Trop 1983 “None of the patients on... ASA required any additional analgesic during the 4-hour observation period, but four patients in the placebo group required supplementary medication for pain” (the denominators for each group were not reported).

### Maternal adverse effects

There was no clear difference in overall maternal adverse effects over four to eight hours post-administration (RR 1.08, 95% CI 0.57 to 2.06; 14 trials, 1067 women; Analysis 1.3). The quality of the evidence (GRADE) for this outcome was judged to be very low due to study limitations (risk of bias) and imprecision (Summary of findings for the main comparison). Visual inspection of the funnel plot for this outcome suggested no clear evidence of reporting bias (Figure 6).

**Figure 6. Funnel plot of comparison: I Aspirin versus placebo for perineal pain, outcome: I.3 Maternal adverse effects**



### Data from the trials not included in the meta-analysis

- Okun 1982: “The incidence of side effects was not significantly different among the treatment groups... Three patients each in the placebo group... and 5 patients in the aspirin group reported side effects” (these data related to women with uterine cramp or episiotomy pain).

- Trop 1983: one woman receiving 1200 mg aspirin (dizziness), no women receiving 600 mg aspirin and two women receiving placebo (nausea) experienced side effects (the denominators for each group were not reported).

### Neonatal adverse effects

None of the included trials reported on the primary outcome: neonatal adverse effects.

### Subgroup analyses based on dose

We integrated subgroup analyses based on dose into the main analyses, comparing trials using 300 mg, 500 to 650 mg, 900 mg and 1200 mg aspirin. No clear subgroup differences were observed based on dose of aspirin for 'Adequate pain relief as reported by the woman' (Test for subgroup differences:  $\text{Chi}^2 = 0.75$ ,  $P = 0.86$ ,  $I^2 = 0\%$ ; [Analysis 1.1](#)); 'Need for additional pain relief' (Test for subgroup differences:  $\text{Chi}^2 = 0.63$ ,  $P = 0.89$ ,  $I^2 = 0\%$ ; [Analysis 1.2](#)); or 'Maternal adverse effects' (Test for subgroup differences:  $\text{Chi}^2 = 3.76$ ,  $df = 2$  ( $P = 0.15$ ),  $I^2 = 46.8\%$ ; [Analysis 1.3](#)).

### Secondary outcomes

None of the included trials reported on any of the secondary review outcomes: prolonged hospitalisation due to perineal pain; re-hospitalisation due to perineal pain; fully breastfeeding at discharge; mixed feeding at discharge; fully breastfeeding at six weeks; mixed feeding at six weeks; perineal pain at six weeks; maternal views (using a validated questionnaire); maternal postpartum depression.

### Comparison 2: 300 mg aspirin versus 600 mg aspirin for perineal pain

[London 1983b](#) contributed data to this comparison.

#### Primary outcomes

##### Adequate pain relief as reported by the woman

There was no clear difference over four hours after drug administration between the 300 mg and 600 mg aspirin groups in the proportion of women who experienced adequate pain relief (RR 0.82, 95% CI 0.36 to 1.86; 1 trial, 81 women; [Analysis 2.1](#)).

##### Need for additional pain relief in the first 48 hours for perineal pain

There was no clear difference over four hours after drug administration between the 300 mg and 600 mg aspirin groups in the proportion of women who required additional pain relief (RR 0.68, 95% CI 0.12 to 3.88; 1 trial, 81 women; [Analysis 2.2](#)).

##### Maternal adverse effects

There were no adverse effects among women in the 300 mg and 600 mg aspirin groups ([Analysis 2.3](#)).

### Neonatal adverse effects

[London 1983b](#) did not report on the primary outcome: neonatal adverse effects.

### Secondary outcomes

[London 1983b](#) did not report on any of the secondary review outcomes.

### Comparison 3: 600 mg aspirin versus 1200 mg aspirin for perineal pain

[Bloomfield 1970a](#) and [London 1983b](#) contributed data to this comparison.

#### Primary outcomes

##### Adequate pain relief as reported by the woman

There was no clear difference over four to eight hours after drug administration between the 600 mg and 1200 mg aspirin groups in the proportion of women who experienced adequate pain relief (RR 0.85, 95% CI 0.52 to 1.39; 2 trials, 121 women; [Analysis 3.1](#)).

##### Need for additional pain relief in the first 48 hours for perineal pain

There was no clear difference over four to eight hours after drug administration between the 600 mg and 1200 mg aspirin groups in the proportion of women who required additional pain relief (RR 1.32, 95% CI 0.30 to 5.68; 2 trials, 121 women; [Analysis 3.2](#)).

##### Maternal adverse effects

There was no clear difference over four to eight hours after drug administration between the 300 mg and 1200 mg aspirin groups in the proportion of women who experienced adverse effects (RR 3.00, 95% CI 0.13 to 69.52; 2 trials, 121 women; [Analysis 3.3](#)).

### Neonatal adverse effects

[Bloomfield 1970a](#) and [London 1983b](#) did not report on the primary outcome: neonatal adverse effects.

### Secondary outcomes

[Bloomfield 1970a](#) and [London 1983b](#) did not report on any of the secondary review outcomes.



#### **Comparison 4: 300 mg aspirin versus 1200 mg aspirin for perineal pain**

[London 1983b](#) contributed data to this comparison.

#### **Primary outcomes**

##### **Adequate pain relief as reported by the woman**

There was no clear difference over four hours after drug administration between the 300 mg and 1200 mg aspirin groups in the proportion of women who experienced adequate pain relief (RR 0.62, 95% CI 0.29 to 1.32; 1 trial, 80 women; [Analysis 4.1](#)).

##### **Need for additional pain relief in the first 48 hours for perineal pain**

There was no clear difference over four hours after drug administration between the 300 mg and 1200 mg aspirin groups in the proportion of women who required additional pain relief (RR 2.00, 95% CI 0.19 to 21.18; 1 trial, 80 women; [Analysis 4.2](#)).

##### **Maternal adverse effects**

There were no adverse effects among women in the 300 mg and 1200 mg aspirin groups ([Analysis 4.3](#)).

##### **Neonatal adverse effects**

[London 1983b](#) did not report on the primary outcome: neonatal adverse effects.

#### **Secondary outcomes**

[London 1983b](#) did not report on any of the secondary review outcomes.

from four trials ([Jain 1978a](#); [Jain 1978b](#); [Okun 1982](#); [Trop 1983](#)) could not be included in the meta-analysis for this outcome. However, results indicated a benefit from aspirin when compared with placebo. In a subgroup analysis, no clear differences were observed based on dose of aspirin. Neither were clear differences observed between groups for adequate pain relief as reported by the women in comparisons of 300 mg and 600 mg aspirin (1 trial, 81 women), 600 mg and 1200 mg aspirin (2 trials, 121 women), and 300 mg and 1200 mg aspirin (1 trial, 80 women).

Very low-quality evidence also showed that women receiving aspirin (doses ranging from 300 mg to 1200 mg) had a 75% relative reduction in the need for additional analgesia for perineal pain over four to eight hours after administration (from 27% in the placebo group to 6% in the aspirin group) across 10 trials involving 744 women. Subgroup analysis did not reveal any clear differences based on dose of aspirin. No clear differences were observed between groups for need of additional pain relief in comparisons of 300 mg and 600 mg aspirin (1 trial, 81 women), 600 mg and 1200 mg aspirin (2 trials, 121 women), and 300 mg and 1200 mg aspirin (1 trial, 80 women).

Very low-quality evidence also showed no clear difference between women receiving aspirin (doses ranging from 300 mg to 1200 mg) versus placebo for maternal adverse effects over four to eight hours after administration (3% in both the aspirin and placebo groups) across 14 trials involving 1067 women. Similarly, there were no subgroup differences based on dose. No clear difference was observed in a comparison of 300 mg and 1200 mg aspirin (2 trials, 121 women).

None of the included trials reported on the review primary outcome - neonatal adverse effects, nor the secondary review outcomes: prolonged hospitalisation due to perineal pain; re-hospitalisation due to perineal pain; fully breastfeeding at discharge; mixed feeding at discharge; fully breastfeeding at six weeks; mixed feeding at six weeks; perineal pain at six weeks; maternal views (using a validated questionnaire); and maternal postpartum depression.

## **DISCUSSION**

### **Summary of main results**

We included 17 trials, of these, 16 randomised 1132 women to single dose aspirin or placebo for perineal pain in the early postpartum period. Fifteen trials contributed data to four comparisons (aspirin versus placebo; 300 mg versus 600 mg aspirin; 600 mg versus 1200 mg aspirin; 300 mg versus 1200 mg aspirin).

Low-quality evidence showed that women receiving aspirin (doses ranging from 300 mg to 1200 mg) had a 103% relative increase in adequate pain relief at four to eight hours after administration compared with placebo (from 25% in the placebo group to 47% in the aspirin group) across 13 trials involving 1001 women. Data

### **Overall completeness and applicability of evidence**

The included trials enrolled women with perineal pain in the early postpartum period post-episiotomy. Accordingly, results may not be applicable to other women with perineal pain, such as those with pain following naturally occurring tears or birth over an intact perineum. All included trials compared aspirin with placebo; we were unable to assess the comparative effects of aspirin versus paracetamol as proposed in the protocol for this review.

Most trials recruited women from the USA (11 trials); three trials were conducted in Venezuela, and one in each Belgium, Canada and India. Sixteen trials were published before the 1990s (one in the 1960s, six in the 1970s, and nine in the 1980s). Results may

not be applicable to all settings or countries worldwide, nor to current clinical practice.

Although there were more than 1000 women and their babies in the included trials, individually, sample sizes were small, ranging from 26 to 178 women. Most trials reported on the review primary outcomes adequate pain relief as reported by the woman (N = 13), need for additional analgesia (N = 10) and maternal adverse effects (N = 14). Only three (of 13) pre-specified outcomes were examined in the included trials; there were no data reported for the primary outcome - neonatal adverse effects - or for any of the secondary review outcomes.

Breastfeeding was clearly stated as an exclusion criterion in most trials, and as a result, no data were available to determine any neonatal adverse effects or effects on breastfeeding. Guidance for the management of perineal pain, including in breastfeeding women, recommends that if oral analgesia is required then paracetamol/acetaminophen should be used first unless contraindicated; if paracetamol is not effective, an oral or rectal non-steroidal anti-inflammatory (NSAID) agent such as ibuprofen should be considered in the absence of contraindications (Montgomery 2012; NICE 2015; NIH 2015). Although some guidance indicates that low-dose aspirin may be considered as an antiplatelet drug for use in breastfeeding women (Bell 2011), it is generally recommended to be used cautiously or avoided during breastfeeding because salicylate and salicylate metabolites are excreted in breast milk. Therefore, there is potential for adverse effects in infants. Longer-term, high-dose maternal aspirin administration has been associated with a report of infant metabolic acidosis, and aspirin administration to infants with viral infections has been associated with Reye's syndrome (NIH 2015).

It is recognised that breastfeeding is an unequalled way of providing the ideal food for infants. International guidance, including from the World Health Organization, recommends (where possible) initiation of breastfeeding within the first hour after birth, and exclusive breastfeeding for the first six months of life for optimal growth, development and health, followed by age-appropriate complementary feeding alongside breastfeeding for two years or more (WHO 2001; WHO 2003). The evidence in this review is thus not applicable to current globally recommended best practice.

## Quality of the evidence

Many aspects of methodological quality were unclear for several of the included trials (Figure 2; Figure 3). Except for one included trial, all studies were published before the 1990s. We found a lack of methodological detail provided in published reports. Attempts to contact trial authors to obtain additional information have been unsuccessful. Of the 17 included trials, 15 were assessed at unclear risk of selection bias; this judgement was based on study reports not detailing methods for sequence generation. All trials had unclear risk of selection bias; assessment was based on study reports not

detailing methods for concealment of allocation. We assessed 14 trials at low risk of performance and detection bias; the risk was judged as unclear for three trials. Most trials were judged to be at unclear or high risk of attrition bias (a number imputing data); and all were at unclear or high risk of reporting bias, with many reporting very limited outcome data, and none had available trial registration or protocols.

We assessed the quality of the evidence using the GRADE approach as outlined in the GRADE Handbook for pre-specified outcomes analysed in the main comparison (aspirin versus placebo). Our assessment was that evidence quality was low (adequate pain relief as reported by the women) or very low (need for additional pain relief; maternal adverse effects). Our judgements were based on design limitations in the included trials (all outcomes), possible publication bias (need for additional pain relief) and imprecision (maternal adverse effects). See [Summary of findings for the main comparison](#).

## Potential biases in the review process

We took steps to minimise the introduction of bias during the review process. At least two review authors independently assessed trials for inclusion, performed data extraction, and assessed risk of bias for each of the included trials.

A detailed, systematic search process was conducted by the Information Specialist of Cochrane Pregnancy and Childbirth, without language or publication status restrictions to reduce the risk for potential publication bias. We also searched trial registries for unpublished, planned or ongoing trials. It is possible that additional trials assessing aspirin for perineal pain in the early postpartum period have been published but not identified; and that further trials have been conducted but are not yet published; or both. Should any such studies be identified in the future, we will assess these for inclusion in future updates of this review.

We investigated reporting biases (such as publication bias) using funnel plots for our primary outcomes. Although no clear evidence of reporting bias was observed for adequate pain relief as reported by the woman and maternal adverse effects, the funnel plot for need for additional pain relief demonstrated some asymmetry. This could indicate possible reporting bias, with the smaller published trials reporting exaggerated intervention effect estimates, and the possibility of additional small trials (including those reporting smaller effect estimates) remaining unpublished.

## Agreements and disagreements with other studies or reviews

Previous Cochrane Reviews have assessed therapeutic ultrasound (Hay-Smith 1998), rectal analgesia (Hedayati 2003), local cooling (East 2012b) and topical anaesthetics (Hedayati 2005) for the relief of perineal pain in the postpartum period, revealing mixed

results. More recently, following publication of a generic protocol (Chou 2009) for a series of reviews of drugs for perineal pain in the early postpartum period, two reviews have assessed paracetamol (Chou 2013), and NSAIDs (Wuytack 2016). Both Chou 2013 (including 10 trials involving 2307 women) and Wuytack 2016 (including 28 trials involving 4181 women) showed benefits for paracetamol and NSAIDs compared with placebo, in terms of an increase in adequate pain relief and reduced need for additional pain relief for women with perineal pain in the early postpartum period. However, like our review, Chou 2013 and Wuytack 2016 found that the risk of bias was unclear for many of the included studies (most of which were also conducted from the 1960s to 1990s), and adverse effects were often not assessed for women, and were not assessed for infants. Breastfeeding women, and thus breastfeeding outcomes, were not included.

Another Cochrane Review (including 37 trials involving 5743 adults) assessed single dose aspirin (doses ranging from 300 mg to 1200 mg) for acute postoperative pain in adults (Derry 2012). Like our review, Derry 2012 found that compared with placebo, aspirin was shown to increase adequate pain relief, and reduced the need for rescue medication. Although Derry 2012 reported that benefits were seen for 600 mg to 650 mg aspirin, 900 mg to 1000 mg aspirin and 1200 mg aspirin, it was reported that lower doses of aspirin (500 mg) were not significantly different from placebo; however, no formal subgroup interaction tests were performed or reported. Derry 2012 reported no difference in adverse effects when 600 mg to 650 mg aspirin was compared with placebo, but reported an increase in adverse effects with 900 mg to 1000 mg aspirin compared with placebo. No formal subgroup interaction test was performed.

No other reviews were identified that assessed single dose aspirin for perineal pain in the early postpartum period.

## AUTHORS' CONCLUSIONS

### Implications for practice

Low-quality evidence suggests that single dose aspirin (at doses ranging from 300 mg to 1200 mg) compared with placebo, can increase adequate pain relief in women with perineal pain post-episiotomy in the early postpartum period. Very low-quality evidence also suggests that aspirin compared with placebo can reduce the need for additional pain relief, and does not appear to increase the occurrence of maternal adverse effects. Evidence was considered to be low to very low quality, and downgrading was based on study limitations (risk of bias), imprecision and publication bias, or both. Current evidence does not suggest clear differences in pain relief or maternal adverse effects based on the dose of aspirin administered. All trials to date have excluded women who were breastfeeding. Therefore, there was no evidence to formally assess the effects of single dose aspirin on neonatal adverse effects

or breastfeeding outcomes. There was no evidence to assess any of the secondary review outcomes.

With international guidance recommending mothers initiate breastfeeding within one hour of birth, and exclusively breastfeeding for the first six months, the evidence from this review is not applicable to current recommended best practice. Aspirin may be considered for use in non-breastfeeding women with post-episiotomy perineal pain. Although formal assessment was beyond the remit of this review, current guidance suggests that other analgesic drugs (such as paracetamol and NSAIDs including ibuprofen) should first be considered for postpartum perineal pain, particularly for breastfeeding women. These agents are the focus of further reviews in this series on drugs for perineal pain in the early postpartum period.

### Implications for research

Due to current guidance suggesting other analgesics be first considered (particularly for breastfeeding women), and possible ethical concerns regarding withholding pain relief, it is considered unlikely that future trials will be conducted to determine the effects of aspirin compared with placebo; it is also considered unlikely that trials will be conducted in breastfeeding women. It is considered most likely, that if conducted, future trials would compare single dose aspirin with other pain relievers. Such trials should be designed to ensure robust methodological quality, and address gaps in the evidence, such as maternal views, postpartum depression, and prolonged hospitalisation or re-hospitalisation. Because research to date has focused on women post-episiotomy, future trials could be extended to include women with perineal trauma associated with naturally occurring tears or birth over an intact perineum.

## ACKNOWLEDGEMENTS

As part of the pre-publication editorial process, this review has been commented on by three peers (an editor and two referees who are external to the editorial team), a member of the Pregnancy and Childbirth Group's international panel of consumers and the Group's Statistical Adviser.

This project was supported by the National Institute for Health Research, via Cochrane Infrastructure funding to Cochrane Pregnancy and Childbirth. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, NHS or the Department of Health.

We acknowledge the support from the Cochrane Pregnancy and Childbirth editorial team in Liverpool, and the Australian and New Zealand Satellite of Cochrane Pregnancy and Childbirth

(funded by the Australian National Health and Medical Research Council (NHMRC)).

## REFERENCES

### References to studies included in this review

#### **Bloomfield 1967** {published data only}

Bloomfield SS, Gaffney TE, Howett M. Comparative analgesic efficacy of chlorphenesin carbamate and acetylsalicylic acid after episiotomy. *Anesthesia and Analgesia* 1967;**46**(5):515–20.

#### **Bloomfield 1970a** {published data only}

\* Bloomfield SS, Barden TP, Hille R. Clinical evaluation of flufenisal, a long-acting analgesic. *Clinical Pharmacology and Therapeutics* 1970;**11**(5):747–54.

Bloomfield SS, Barden TP, Mitchell J. Aspirin and codeine in two postpartum pain models. *Clinical Pharmacology and Therapeutics* 1976;**20**(4):499–503.

#### **Bloomfield 1970b** {published data only}

Bloomfield SS, Hurwitz HN. Tourniquet and episiotomy pain as test models for aspirin-like analgesics. *Journal of Clinical Pharmacology* 1970;**10**(6):361–9.

#### **Bloomfield 1974** {published data only}

Bloomfield SS, Barden TP, Mitchell J. Comparative efficacy of ibuprofen and aspirin in episiotomy pain. *Clinical Pharmacology and Therapeutics* 1974;**15**(6):565–70.

#### **De Vroey 1978** {published data only}

\* De Vroey P. A double-blind comparison of diflunisal and aspirin in the treatment of post-operative pain after episiotomy. *Current Medical Research and Opinion* 1978;**5**(7):544–7.

De Vroey P. The treatment of postoperative pain with a single dose of diflunisal. *Clinical Therapeutics* 1977;**1**(Suppl A):30–3.

Devroey P, Steelman SL, Caudron J. A double blind, placebo controlled, single dose study comparing three dose levels of diflunisal with aspirin and placebo in patients with pain due to episiotomy. *Acta Therapeutica* 1977;**3**(3):205–16.

#### **Friedrich 1983** {published data only}

Friedrich E. A comparison of etodolac (Ultradol) with aspirin and placebo in patients with episiotomy pain. *Current Therapeutic Research, Clinical and Experimental* 1983;**33**(1):100–7.

#### **Jain 1978a** {published data only}

Jain AK, McMahon FG, Ryan JR, Raphan H, Richard W. Piroxicam, a novel analgesic in postpartum pain. *European Journal of Rheumatology and Inflammation* 1978;**1**(3):356–9.

#### **Jain 1978b** {published data only}

Jain AK, McMahon FG, Ryan JR, Unger D, Richard W. A comparison of aspirin-caffeine versus aspirin: results of two double-blind placebo controlled studies in postpartum pain

[abstract]. *Clinical Pharmacology and Therapeutics* 1978;**23**(1):116.

\* Jain AK, McMahon FG, Ryan JR, Unger D, Richard W. Aspirin and aspirin-caffeine in postpartum pain relief. *Clinical Pharmacology and Therapeutics* 1978;**24**(1):69–75.

#### **Jain 1985** {published data only}

Jain AK, McMahon FG, Ryan JR, Smith GB. Analgesic efficacy of indoprofen in postpartum episiotomy pain. *Current Therapeutic Research, Clinical and Experimental* 1985;**38**(5):677–81.

#### **London 1983a** {published data only}

London R, Sundaram GS, Feldman S, Goldstein P. Evaluation of analgesic efficacy and safety of fluproquazone compared to aspirin in the treatment of episiotomy pain. *Federation Proceedings* 1982;**41**(4):No. 5573.

\* London R, Sundaram GS, Feldman S, Goldstein PJ. Episiotomy pain: efficacy and safety of fluproquazone compared to aspirin and placebo. *International Journal of Gynecology & Obstetrics* 1983;**21**(3):251–5.

#### **London 1983b** {published data only}

London RS, Sundaram GS, Feldman S, Goldstein PJ. Aspirin in the treatment of episiotomy pain. *Southern Medical Journal* 1983;**76**(7):844–5.

#### **Mukherjee 1980** {published data only}

Mukherjee S, Sood S. A controlled evaluation of orally administered aspirin dipyron and placebo in patients with post-operative pain. *Current Medical Research and Opinion* 1980;**6**(9):619–23.

#### **Okun 1982** {published data only}

Okun R. Evaluation of the analgesic effect of fendosal in patients with postpartum uterine cramp or episiotomy pain. *Current Therapeutic Research, Clinical and Experimental* 1982;**32**(1):65–73.

#### **Olson 1997** {published data only}

Olson NZ, Sunshine A, Zigelboim I, DeCastro A. Onset and duration of analgesia of diclofenac potassium in the treatment of postepisiotomy pain. *American Journal of Therapeutics* 1997;**4**(7-8):239–46.

#### **Sunshine 1983a** {published data only}

Sunshine A, Olson NZ, Laska EM, Zigelboim I, De Castro A, De Sarrazin C. Analgesic effect of graded doses of flurbiprofen in post-episiotomy pain. *Pharmacotherapy* 1983;**3**(3):177–81.

#### **Sunshine 1983b** {published data only}

Sunshine A, Olson NZ, Laska EM, Zigelboim I, De Castro A, De Sarrazin C. Ibuprofen, zomepirac, aspirin, and placebo in the relief of postepisiotomy pain. *Clinical Pharmacology and Therapeutics* 1983;**34**(2):254–8.

**Trop 1983** *{published data only}*

Trop D, Nucci C, Elie R, Gareau J. Double-blind comparative evaluation of tiaprofenic acid (Surgam) versus acetylsalicylic acid (ASA) in relieving pain following episiotomy. *Current Therapeutic Research, Clinical and Experimental* 1983;**34**(21):274–9.

**References to studies excluded from this review****Bruni 1965** *{published data only}*

Bruni JR, Holt RE. Controlled double-blind evaluation of three analgesic medications for postpartum discomfort. *Obstetrics and Gynecology* 1965;**25**:76–81.

**Gindhart 1971** *{published data only}*

Gindhart JD. A rationale for studying analgesia. A double blind study in postpartum patients. *Current Therapeutic Research, Clinical and Experimental* 1971;**13**(4):240–50.

**Gruber 1979** *{published data only}*

Gruber CM, Bauer RO, Bettigole JB, Lash AF, McDonald JS. A multicenter study for analgesia involving fenoprofen, propoxyphene (alone or in combination) with placebo and aspirin controls in postpartum pain. *Journal of Medicine* 1979;**10**(1-2):65–98.

**Moggian 1972** *{published data only}*

Moggian G, Cervellati I, Tamburini E. Critical study of the post-partum pain by means of a clinical pharmacology experiment. *Bollettino Chimico Farmaceutico* 1972;**111**(8):497–9.

**Prockop 1960** *{published data only}*

Prockop LD, Eckenhoff JE, McElroy RC. Evaluation of dextropropoxyphene, codeine and acetylsalicylic compound. *Obstetrics and Gynecology* 1960;**16**:113–8.

**Rubin 1984** *{published data only}*

Rubin A, Winter LJ. A double-blind randomized study of an aspirin/caffeine combination versus acetaminophen/ aspirin combination versus acetaminophen versus placebo in patients with moderate to severe post-partum pain. *Journal of International Medical Research* 1984;**12**(6):338–45.

**Santiago 1959** *{published data only}*

Santiago FS, Danforth DN. Non-narcotic analgesia to simplify postpartum care. *Obstetrics and Gynecology* 1959;**14**(1):22–3.

**Sunshine 1983c** *{published data only}*

Sunshine A, Olson JZ, Siegel C, Laska EM. Oral analgesic study of ketoprofen, aspirin and placebo in post-partum pain. *Clinical Pharmacology and Therapeutics* 1983;**33**:154. Sunshine A, Zigelboim I, Laska E, Siegel C, Olson NZ, De Castro A. A double-blind, parallel comparison of ketoprofen, aspirin, and placebo in patients with postpartum pain. *Journal of Clinical Pharmacology* 1986;**26**(8):706–11.

**Sunshine 1985** *{published data only}*

Sunshine A, Zigelboim I, Olson NZ, De Sarrazin C, Laska E. A comparative oral analgesic study of indoprofen, aspirin, and placebo in postpartum pain. *Journal of Clinical Pharmacology* 1985;**25**(5):374–80.

**Van der Pas 1984** *{published data only}*

Van der Pas HFM, Vertommen F. A double-blind study of naproxen-suppositories for post episiotomal pain compared with asa, oxyfenbutazon and placebo. *Ars Medici Revue Internationale de Therapie Pratique* 1984;**39**(5):79–81.

**References to studies awaiting assessment****Bhounsule 1990** *{published data only}*

Bhounsule SA, Nevreker PR, Agshikar NV, Pal MN, Dhume VG. A comparison of four analgesics in post-episiotomy pain. *Indian Journal of Physiology and Pharmacology* 1990;**34**(1):34–8.

**Sunshine 1989** *{published data only}*

Sunshine A, Zigelboim I, Hoburg A, Byrd W, Olson NZ, Laska E, et al. Ibuprofen, aspirin and placebo in postpartum pain (abstract). *International Journal of Clinical Pharmacology and Therapeutics* 1989;**45**(2):174.

**Additional references****Aasheim 2011**

Aasheim V, Nilsen ABV, Lukasse M, Reinart LM. Perineal techniques during the second stage of labour for reducing perineal trauma. *Cochrane Database of Systematic Reviews* 2011, Issue 12. [DOI: 10.1002/14651858.CD006672.pub2]

**ATT Collaboration 2002**

Antithrombotic Trialists' Collaboration. Collaborative meta-analysis of randomised trials of antiplatelet therapy for prevention of death, myocardial infarction, and stroke in high risk patients. *BMJ* 2002;**324**(7329):71–86.

**ATT Collaboration 2009**

Antithrombotic Trialists' (ATT) Collaboration, Baigent C, Blackwell L, Collins R, Emberson J, Godwin J, et al. Aspirin in the primary and secondary prevention of vascular disease: collaborative meta-analysis of individual participant data from randomised trials. *Lancet* 2009;**373**(9678):1849–60.

**Beckmann 2013**

Beckmann MM, Stock OM. Antenatal perineal massage for reducing perineal trauma. *Cochrane Database of Systematic Reviews* 2013, Issue 4. [DOI: 10.1002/14651858.CD005123.pub3]

**Bell 2011**

Bell AD, Roussin A, Cartier R, Chan WS, Douketis JD, Gupta A, et al. The use of antiplatelet therapy in the outpatient setting: Canadian Cardiovascular Society Guidelines Executive Summary. *Canadian Journal of Cardiology* 2011;**27**(2):208–21.

**Chou 2009**

Chou D, Abalos E, Gyte Gillian ML, Gülmezoglu AM. Drugs for perineal pain in the early postpartum period: generic protocol. *Cochrane Database of Systematic Reviews* 2009, Issue 3. [DOI: 10.1002/14651858.CD007734.pub2]

**Chou 2013**

Chou D, Abalos E, Gyte GM, Gulmezoglu AM. Paracetamol/acetaminophen (single administration) for perineal pain in the early postpartum period. *Cochrane Database of Systematic Reviews* 2013, Issue 1. [DOI: 10.1002/14651858.CD008407.pub2]

**Cooper 1991**

Cooper SA. Commentary: Single-dose analgesic studies: the upside and downside of assay sensitivity. In: Max M, Portenov R, Laska E editor(s). *Advances in Pain Research and Therapy*. Vol. 18, New York: Raven Press, 1991:117–24.

**Cunningham 2005**

Cunningham G. Maternal anatomy. In: Cunningham G, Leveno K, Bloom S, Hauth J, Gilstrap L, Wenstrom K editor(s). *Williams Obstetrics*. 22nd Edition. New York: McGraw-Hill, 2005:21.

**Derry 2012**

Derry S, Moore RA. Single dose oral aspirin for acute postoperative pain in adults. *Cochrane Database of Systematic Reviews* 2012, Issue 4. [DOI: 10.1002/14651858.CD002067.pub2]

**Duley 2007**

Duley L, Henderson-Smart DJ, Meher S, King JF. Antiplatelet agents for preventing pre-eclampsia and its complications. *Cochrane Database of Systematic Reviews* 2007, Issue 2. [DOI: 10.1002/14651858.CD004659.pub2]

**East 2012a**

East CE, Sherburn M, Nagle C, Said J, Forster D. Perineal pain following childbirth: prevalence, effects on postnatal recovery and analgesia usage. *Midwifery* 2012;**28**(1):93–7.

**East 2012b**

East CE, Begg L, Henshall NE, Marchant PR, Wallace K. Local cooling for relieving pain from perineal trauma sustained during childbirth. *Cochrane Database of Systematic Reviews* 2012, Issue 5. [DOI: 10.1002/14651858.CD006304.pub3]

**Hay-Smith 1998**

Hay-Smith J. Therapeutic ultrasound for postpartum perineal pain and dyspareunia. *Cochrane Database of Systematic Reviews* 1998, Issue 3. [DOI: 10.1002/14651858.CD000495]

**Hedayati 2003**

Hedayati H, Parsons J, Crowther CA. Rectal analgesia for pain from perineal trauma following childbirth. *Cochrane Database of Systematic Reviews* 2003, Issue 3. [DOI: 10.1002/14651858.CD003931]

**Hedayati 2005**

Hedayati H, Parsons J, Crowther CA. Topically applied anaesthetics for treating perineal pain after childbirth. *Cochrane Database of Systematic Reviews* 2005, Issue 2. [DOI: 10.1002/14651858.CD004223.pub2]

**Higgins 2011**

Higgins JPT, Green S, editors. *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated

March 2011]. The Cochrane Collaboration, 2011. Available from [www.cochrane-handbook.org](http://www.cochrane-handbook.org).

**Kettle 2012**

Kettle C, Dowswell T, Ismail KMK. Continuous and interrupted suturing techniques for repair of episiotomy or second-degree tears. *Cochrane Database of Systematic Reviews* 2012, Issue 11. [DOI: 10.1002/14651858.CD000947.pub3]

**Li 2013**

Li Z, Zeki R, Hilder L, Sullivan EA. *Australia's Mothers and Babies 2011. Perinatal Statistics Series No. 28*. Vol. Cat no. PER 59, Canberra: AIHW National Perinatal Epidemiology and Statistics Unit, 2013.

**Maclagan 1879**

Maclagan TJ. The treatment of acute rheumatism by salicin and salicylic acid. *Lancet* 1879;**113**(2912):875–7.

**Montgomery 2012**

Montgomery A, Hale TW, Academy of Breastfeeding Medicine. ABM clinical protocol #15: analgesia and anesthesia for the breastfeeding mother, revised 2012. *Breastfeeding Medicine* 2012;**7**(6):547–53.

**Moore 1996**

Moore A, McQuay H, Gavaghan D. Deriving dichotomous outcome measures from continuous data in randomised controlled trials of analgesics. *Pain* 1996;**66**(2-3):229–37.

**Moore 1997a**

Moore A, Moore O, McQuay H, Gavaghan D. Deriving dichotomous outcome measures from continuous data in randomised controlled trials of analgesia: use of pain intensity and visual analogue scales. *Pain* 1997;**69**(3):311–5.

**Moore 1997b**

Moore A, McQuay H, Gavaghan D. Deriving dichotomous outcome measures from continuous data in randomised controlled trials of analgesia: verification from independent data. *Pain* 1997;**69**(1-2):127–30.

**NICE 2015**

National Institute for Health and Care Excellence (NICE). Postnatal care up to 8 weeks after birth, NICE Clinical Guideline CG37. [nice.org.uk/guidance/cg37](http://nice.org.uk/guidance/cg37) (accessed 20 August 2016).

**NIH 2015**

US National Library of Medicine, National Institutes of Health, Health and Human Services. LacMed, a ToxNet database. [www.toxnet.nlm.nih.gov](http://www.toxnet.nlm.nih.gov) (accessed 20 June 2016).

**RevMan 2014 [Computer program]**

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

**Rothwell 2010**

Rothwell PM, Wilson M, Elwin CE, Norrving B, Algra A, Warlow CP, et al. Long-term effect of aspirin on colorectal cancer incidence and mortality: 20-year follow-up of five randomised trials. *Lancet* 2010;**376**(9754):1741–50.

**Smith 2014**

Smith T, Elwood P, Keating C, Rothwell P, Detering E, Freedman A, et al. The Aspirin Foundation Scientific Conference: the history, the present state and the future of aspirin prophylaxis. *Ecancermedalscience* 2014;**8**:388. [PUBMED: 24678343]

**Vane 2003**

Vane JR, Botting RM. The mechanism of action of aspirin. *Thrombosis Research* 2003;**110**(5-6):255–8.

**WHO 2001**

World Health Organization. *The Optimal Duration of Exclusive Breastfeeding. WHO Report of an Expert Consultation*. Geneva, Switzerland: World Health Organization, 2001.

**WHO 2003**

World Health Organization, UNICEF. *Global Strategy for Infant and Young Child Feeding*. Geneva, Switzerland: World Health Organization, 2003.

**WHO 2011**

World Health Organization. WHO recommendations for prevention and treatment of pre-eclampsia and eclampsia. *WHO recommendations for prevention and treatment of pre-Eclampsia and eclampsia*. Geneva: World Health Organization, 2011.

**Wuytack 2016**

Wuytack F, Smith V, Cleary BJ. Oral non-steroidal anti-inflammatory drugs (single dose) for perineal pain in the early postpartum period. *Cochrane Database of Systematic Reviews* 2016, Issue 7. [DOI: 10.1002/14651858.CD011352.pub2]

**References to other published versions of this review****Molakatalla 2016**

Molakatalla S, Shepherd E, Grivell RM. Aspirin (single dose) for perineal pain in early postpartum period. *Cochrane Database of Systematic Reviews* 2016, Issue 3. [DOI: 10.1002/14651858.CD012129]

\* Indicates the major publication for the study

## CHARACTERISTICS OF STUDIES

### Characteristics of included studies [ordered by study ID]

#### Bloomfield 1967

Methods	RCT
Participants	<p><b>Setting:</b> Cincinnati General Hospital, Ohio, USA</p> <p><b>Date:</b> December 1965 to April 1966</p> <p><b>Inclusion criteria:</b> women with a painful ('moderate' or 'severe') mediolateral episiotomy, within 24 hours following an uncomplicated labour and birth</p> <p><b>Exclusion criteria:</b> breastfeeding; aged &lt; 18 years; known aspirin sensitivity; 'mild' pain at interview within 24 hours of birth</p>
Interventions	<p><b>Aspirin (N = 17 randomised)</b> 600 mg aspirin; women received a single oral dose in a black capsule</p> <p><b>Placebo (N = 18 randomised)</b> Women received a single oral dose in a black capsule</p> <p><b>All women:</b> women did not receive other analgesics during the 6 hours of study or during the 6 hours before entering the study</p>
Outcomes	<p><b>Adequate pain relief as reported by the woman:</b> pain intensity evaluated by 1 research nurse hourly for 6 hours; women were asked "How much do your stitches hurt you?", and answers were transposed into an ordinal scale from 0 to 3 (0 = no pain; 1 = slight pain; 2 = moderate pain; 3 = severe pain). The difference between a woman's pre-treatment pain intensity score and each hourly post-treatment score gave an hourly pain relief score; a total 6 hour pain relief score was calculated for each woman by adding these scores. Mean Pain Relief scores (equivalent to SPID scores) were used to calculate 'Adequate pain relief as reported by the woman' (taken over 6 hours)</p> <p><b>Maternal adverse effects:</b> women were asked on the day following treatment whether they noticed any other effects of the treatment; if they answered 'yes' they were asked 'What were they'; no leading questions were asked</p>
Notes	<p><b>Funding:</b> the study was supported in part by USPHS grants HE 05622 and HE 07392 from the National Institutes of Health</p> <p><b>Declarations of interest:</b> not reported (short 'About the authors' section describing affiliations)</p> <p><b>Additional trial arms:</b> this was a 5-arm trial, also assessing chlorphenesin 400 mg (N = 18), 800 mg (N = 17) and combination aspirin 300 mg and chlorphenesin 400 mg (N = 18); we included only the relevant arms in this review</p>

#### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "were randomly assigned... according to a predetermined schedule"
Allocation concealment (selection bias)	Unclear risk	Not detailed



**Bloomfield 1967** (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quotes: “double-blind”; and “All patients received a single dose of coded medication by mouth in identical black capsules”. Assumed that blinding of women and personnel was successful with the use of an identical placebo
Blinding of outcome assessment (detection bias) All outcomes	Low risk	1 research nurse evaluated pain intensity and side effects by interviewing women. Assumed that blinding of the research nurse and women was successful with the use of an identical placebo
Incomplete outcome data (attrition bias) All outcomes	High risk	88 women had moderate or severe episiotomy pain; 84 completed the 6 hours of study and “form the basis of this report”... “Four of the 88 patients entering the trial were withdrawn owing to distressing pain unrelieved by the study drugs” (1/17 from the aspirin group; 0/18 from the placebo group)
Selective reporting (reporting bias)	Unclear risk	Very few outcomes reported (pain relief and side effects only); no access to trial registration or protocol to further assess selective reporting
Other bias	Low risk	Baseline characteristics were comparable between groups; no other obvious risk of bias identified

**Bloomfield 1970a**

Methods	RCT
Participants	<p><b>Setting:</b> Cincinnati General Hospital, University of Cincinnati College of Medicine, Ohio, USA (assumed from affiliation)</p> <p><b>Inclusion criteria:</b> healthy, consenting, ward patients with moderate to very severe episiotomy pain (mediolateral or midline) within 48 hours of an otherwise uncomplicated birth</p> <p><b>Exclusion criteria:</b> mild pain; under the age of 18; history of aspirin allergy; breastfeeding; given analgesics within the previous 6 hours</p>
Interventions	<p><b>Aspirin Group 1* (N = 20 randomised)</b> 1200 mg aspirin; women received a single oral dose in capsules</p> <p><b>Aspirin Group 2* (N = 20 randomised)</b> 600 mg aspirin; women received a single oral dose in a capsules</p> <p><b>Placebo (N = 19 randomised)</b> Lactose placebo; women received a single oral dose in a capsule</p> <p><b>All women:</b> lactose capsules were included with medication where necessary to provide a total of 4 capsules per dose. All drugs were administered before breakfast with a full glass of water, and women were instructed to lie on their right side for 2 hours after administration. Stilbestrol and ferrous sulphate were given routinely in the postpartum period, but all other drugs except for the study drugs were avoided, and except for “cleansing: all perineal care was suspended for the 8-hour study period</p>

<p>Outcomes</p>	<p><b>Adequate pain relief as reported by the woman:</b> the same trained nurse observer interviewed women hourly for 8 hours; they were asked “How much do the stitches hurt you”; answers were transposed to an ordinal score on a scale of 0 to 4 (0 = no pain; 1 = mild pain; 2 = moderate pain; 3 = severe pain; 4 = very severe pain):</p> <ul style="list-style-type: none"> <li>• Pain intensity difference scores were calculated by the difference between a woman’s pre-treatment pain intensity score and early hourly post-treatment score; these scores (equivalent to SPID scores) were used to calculate ‘Adequate pain relief as reported by the woman’ (taken over 8 hours)</li> <li>• Percentage of women with pain reduction &gt; 50% (a fall of &gt; 50% in the pre-treatment pain intensity) was also reported</li> </ul> <p><b>Need for additional pain relief in the first 48 hours for perineal pain:</b> requirement for additional known analgesic medication (codeine or propoxyphene) for inadequate response to study drugs</p> <p><b>Maternal adverse effects:</b> side effects were evaluated at the last interview by the question, “Did you notice any other effects from today’s medicine?” If the answer was “yes”, the woman was asked, “What are they?” No other leading questions were asked</p>	
<p>Notes</p>	<p><b>Funding:</b> “Supported in part by United States Public Health Service Grant HE-05622 and by Merck Sharp &amp; Dohme Research Laboratories. Supplies of flufenisal and other coded medications were provided by Dr. A. W. Vogel, Merck Sharp &amp; Dohme Research Laboratories”</p> <p><b>Declarations of interest:</b> not reported</p> <p><b>Additional trial arms:</b> this was a 5-arm trial, also assessing flufenisal 300 mg (N = 20) and flufenisal 600 mg (N = 21); we included only the relevant arms in this review</p> <p><b>Note:</b> we combined the 2 aspirin groups for the main analysis</p>	
<p><i>Risk of bias</i></p>		
<p><b>Bias</b></p>	<p><b>Authors’ judgement</b></p>	<p><b>Support for judgement</b></p>
<p>Random sequence generation (selection bias)</p>	<p>Unclear risk</p>	<p>Quote: “Within each of the 3 strata of pain intensity, patients were randomly assigned under double-blind conditions to one of the 5 treatment groups according to a predetermined balanced allocation schedule”</p>
<p>Allocation concealment (selection bias)</p>	<p>Unclear risk</p>	<p>Not detailed</p>
<p>Blinding of participants and personnel (performance bias) All outcomes</p>	<p>Low risk</p>	<p>Quote: “All test medications were prepackaged in individual patient-coded vials containing a single oral dose in identical capsules. Lactose capsules were included with medication where necessary to uniformly provide a total of 4 capsules per dose” and “double-blind conditions”</p>
<p>Blinding of outcome assessment (detection bias) All outcomes</p>	<p>Low risk</p>	<p>Subjective evaluation of pain relief and side effects; the same trained nurse observer interviewed women; considered reasonable to assume nurse and women were blinded</p>

**Bloomfield 1970a** (Continued)

Incomplete outcome data (attrition bias) All outcomes	High risk	Imputation of data likely to have influenced results. Quote: “Pain relief data collected before additional analgesic was given to each of these 14 patients was included in the analysis without qualification, but interviews were discontinued. By convention each patient’s pain intensity score for each of the residual hours was adjusted to the value of her pretreatment score, and these adjusted scores were used for calculations of pain relief which were then analysed together with the earlier recorded data. Although such an adjustment was arbitrary and tended to underestimate in these 14 patients the analgesic response to the study treatments, bias in the opposite direction, i.e. tending to exaggerate analgesic response to treatments, would have occurred if all or part of the hourly data for these 14 patients would have been excluded from the analysis or no adjustment made”
Selective reporting (reporting bias)	Unclear risk	Limited number of outcomes reported; no access to trial registration or protocol to further assess selective reporting
Other bias	Unclear risk	Most baseline characteristics were similar among groups; quote “However, body weight was not similar in all treatment groups”

**Bloomfield 1970b**

Methods	RCT
Participants	<p><b>Setting:</b> Cincinnati General Hospital, University of Cincinnati College of Medicine, Ohio, USA (assumed from affiliation)</p> <p><b>Inclusion criteria:</b> healthy, consenting women with mild to severe episiotomy pain within 24 hours of an otherwise uncomplicated birth</p> <p><b>Exclusion criteria:</b> allergy to aspirin; receipt of medication during the 6 hours before treatment</p>
Interventions	<p><b>Aspirin (N = 13 randomised)</b> 1200 mg aspirin given in coded single oral dose of 4 capsules</p> <p><b>Placebo (N = 13 randomised)</b> Identical lactose placebo given in coded single oral dose of 4 capsules</p> <p><b>All women:</b> no additional medications were received in the 5-hour period of pain evaluation. Medications were given after a non-fatty breakfast, and all other foods except water were withheld until after the pain evaluations were completed</p>
Outcomes	<p><b>Adequate pain relief as reported by the woman:</b> the same trained nurse observer interviewed women hourly for 5 hours; they were asked “How much do the stitches hurt you”; answers were transposed to an ordinal score on a scale of 0 to 3 (0 = no pain; 1 = mild pain; 2 = moderate pain; 3 = severe pain):</p> <ul style="list-style-type: none"> <li>• Pain relief scores were calculated by the difference between a woman’s pre-treatment score and early hourly post-treatment score; mean pain relief scores at 0-5 hours were presented in Figure 2 of manuscript, and were used to calculate ‘Adequate pain relief as reported by the woman’</li> </ul>

**Bloomfield 1970b** (Continued)

	<ul style="list-style-type: none"> <li>Percentage of women with pain reduction &gt; 50% (a fall of more than 50% in the pre-treatment pain intensity) was also reported</li> </ul>	
Notes	<p><b>Funding:</b> "This investigation was supported in part by USPHS training grant HE-05622 and by the Special Research Fellowship HE-34688 of the National Heart Institute"</p> <p><b>Declarations of interest:</b> not reported</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Quote: "Within each of the three strata of pain intensity... patients were randomly assigned to one of the two treatment groups"
Allocation concealment (selection bias)	Unclear risk	Not detailed
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "identical lactose placebo, given in a coded single oral dose of four capsules under double-blind conditions"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Subjective evaluation of episiotomy pain; the same trained nurse observed interviewed women; reasonable to assume women and the nurse were blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	No losses/exclusions for pain relief
Selective reporting (reporting bias)	Unclear risk	Pain relief was the only outcome reported; no access to trial registration or protocol to further assess selective reporting
Other bias	Unclear risk	Very limited methodological detail provided; no detail of baseline characteristics

**Bloomfield 1974**

Methods	RCT
Participants	<p><b>Setting:</b> University of Cincinnati Medical Center, Ohio, USA (assumed from affiliation)</p> <p><b>Inclusion criteria:</b> healthy postpartum women with moderate to very severe episiotomy pain (mediolateral or midline) within 48 hours of an otherwise uncomplicated birth</p> <p><b>Exclusion criteria:</b> mild pain; unmarried, aged &lt; 18 years; history of aspirin allergy; given analgesics, sedatives or other psychotropic drugs within the previous 6 hours; breastfeeding; known drug dependence</p>
Interventions	<p><b>Aspirin (N = 20 randomised)</b> 900 mg aspirin; single oral dose of 3 tablets of 300 mg</p> <p><b>Placebo (N = 20 randomised)</b> Lactose placebo; single oral dose of 3 tablets</p>

	<p><b>All women:</b> stilbestrol and ferrous sulphate were given routinely during the postpartum period, however all other drugs were avoided unless necessary, and except for “cleansing”, all perineal care was suspended for the 6-hour study period; women were confined to bed for the first 2 hours and were intermittently out of bed during the last 4 hours. Tablets were administered on demand with a full glass of water at approximately the same time of the day throughout the study (2 hours before breakfast) and women were instructed to lie on their right sides for 2 hours afterwards</p>	
Outcomes	<p><b>Adequate pain relief as reported by the woman:</b> the same trained nurse observer interviewed women hourly for 6 hours; women estimated the severity of ‘stitch’ pain on an ordinal score on a scale of 0 to 4 (0 = no pain; 1 = mild pain; 2 = medium pain; 3 = severe pain; 4 = very severe pain):</p> <ul style="list-style-type: none"> <li>• Pain intensity difference scores were calculated by the difference between a woman’s pre-treatment pain intensity score and early hourly post-treatment score; these scores (equivalent to SPID scores), presented in Figure 4 in the manuscript, were used to calculate ‘Adequate pain relief as reported by the woman’ (taken over 6 hours)</li> <li>• Number of women with pain reduction &gt; 50% at any time during the 6 hours was also reported</li> </ul> <p><b>Need for additional pain relief in the first 48 hours for perineal pain:</b> request for additional analgesic medication (codeine or propoxyphene) before the end of the 6-hour study period</p> <p><b>Maternal adverse effects:</b> side effects were elicited spontaneously at final interview “with a minimum use of leading questions and without invoking a checklist of possible side effects”</p>	
Notes	<p><b>Funding:</b> “Supported in part by United States Public Health Service Grant No. HL-05622 and by the Upjohn Company. Supplies of ibuprofen and other coded medications were provided by Carter D. Brooks, M.D., The Upjohn Company”</p> <p><b>Declarations of interests:</b> not reported</p> <p><b>Additional trial arms:</b> this was a 4-arm trial also assessing ibuprofen 300 mg (N = 20) and 900 mg (N = 20); we have only included the relevant arms in this review</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors’ judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Quote: “On entering the study patients were randomly allocated to one of the 4 groups according to a predetermined schedule. The randomization provided for stratification of patients on the basis of initial intensity of pain”
Allocation concealment (selection bias)	Unclear risk	Not detailed
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quotes: “double-blind conditions”... “All were in the form of film-coated tablets identical in appearance and taste, and were prepackaged in code-numbered individual dose vials”

**Bloomfield 1974** (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: “Changes in pain intensity and side effects associated with the treatments were evaluated subjectively in uniformly conducted interviews”; reasonable to assume blinding of women and interviewer
Incomplete outcome data (attrition bias) All outcomes	High risk	No losses/exclusions reported; assumed only 80 women randomised, and all included in analyses; 4 women requested additional analgesic (3 in the placebo group); their pain relief data before additional analgesics were given were included in the analysis, but interviews were then discontinued, and for the remaining hours, each woman’s pain intensity score was adjusted to the value of her pre-treatment score. Imputation of data likely to have influenced results
Selective reporting (reporting bias)	Unclear risk	Very limited outcome data; no access to trial registration or protocol to further assess selective reporting
Other bias	Low risk	Baseline characteristics comparable for relevant groups (“Two exceptions were a preponderance of unmarried patients in the ibuprofen 300 mg group compared with the 3 other groups, and body weight, which in the group of patients receiving ibuprofen 900 mg was distinctly higher than in patients in the other 3 groups. These were chance occurrences with an uncertain influence on the results”); no other obvious risk of bias identified

**De Vroey 1978**

Methods	RCT
Participants	<p><b>Setting:</b> Department of Obstetrics and Gynaecology, St Christiana Clinic, Dendermonde, Belgium</p> <p><b>Inclusion criteria:</b> primiparae who had undergone mediolateral episiotomy (3 cm to 5 cm) during the course of an otherwise uncomplicated birth within the previous 48 hours, with moderate to severe pain</p> <p><b>Exclusion criteria:</b> a more extensive episiotomy (because of forceps birth or other procedures); multigravida women; known allergy to aspirin; breastfeeding; other analgesic therapy within the previous 6 hours; mild pain</p>
Interventions	<p><b>Aspirin (N randomised was unclear; N = 32 analysed)</b> 600 mg; single oral dose in 2 identical capsules</p> <p><b>Placebo (N randomised was unclear; N = 31 analysed)</b> Placebo; single oral dose in 2 identical capsules</p>
Outcomes	<p><b>Adequate pain relief as reported by the woman:</b> the same trained nurse observer questioned women hourly for 6 hours; women estimated the severity of pain on a scale of 0 to 3 (0 = no pain; 1 = mild pain; 2 = moderate pain; 3 = severe pain):</p> <ul style="list-style-type: none"> <li>• Mean pain scores were reported for each hour; the sum of the difference between these scores and pre-treatment scores (SPID scores) were used to calculate 'Adequate</li> </ul>

	<p>pain relief as reported by the woman' (taken over 6 hours)</p> <ul style="list-style-type: none"> <li>The percentage of women who showed an improvement of at least 2 rating score points (i.e. from severe to mild; or from moderate to no pain) was also reported</li> </ul> <p><b>Need for additional pain relief in the first 48 hours for perineal pain:</b> request for additional analgesic medication 4 hours after administration of study drugs</p> <p><b>Maternal adverse effects:</b> close observation was made for any "adverse reactions"</p>	
Notes	<p><b>Funding:</b> "The statistical assistance of T. COOK, B. RODDA and C. DAURIO of the Merck Sharp &amp; Dohme Research Laboratories is gratefully acknowledged"</p> <p><b>Declarations of interests:</b> not reported; though author affiliations include "Merck Sharp &amp; Dohme Research Laboratories"</p> <p><b>Additional arms:</b> this was a 5-arm trial also assessing diflunisal 125 mg (N = 33 analysed), 250 mg (N = 30 analysed) and 500 mg (N = 30 analysed); we have only included the relevant arms in this review</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Quote: "Patients were allocated at random"
Allocation concealment (selection bias)	Unclear risk	Not detailed
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quotes: "double-blind"... "All test medications were prepackaged in individual patient-coded vials containing a single oral dose in two identically appearing capsules"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Efficacy parameters and side-effects were recorded by the investigator or by the same trained nurse observer, who questioned the patient at hourly intervals"
Incomplete outcome data (attrition bias) All outcomes	High risk	5/161 women admitted to the trial were excluded from the analysis: 2 as their initial pain was not considered severe enough to meet protocol; 2 due to incomplete data; 1 due to lack of cooperation (unclear from which groups; leaving 156 in total - 32 in the aspirin group and 31 in the placebo group). 3 women in the placebo group were withdrawn at 4 hours because of severe pain, and 1 woman in the aspirin group at 3 hours for reasons unrelated to pain or the drug; women who dropped out of the study were included in the analysis; they were assigned a pain score of 4 = worse than the scores of all women who remained in the study
Selective reporting (reporting bias)	High risk	Very limited outcome data reported; no access to trial registration/protocol to further assess selective reporting Quote: "As pain relief was still very marked in the 500 mg diflunisal group at 6 hours, it was decided to extend the period of observation to 8 hours in 42 patients, who were approximately

De Vroey 1978 (Continued)

		evenly distributed between the three groups”
Other bias	Unclear risk	Few baseline characteristics reported (initial pain score rating; age); limited methodological data reported

Friedrich 1983

Methods	RCT
Participants	<p><b>Setting:</b> Department of Obstetrics and Gynaecology, Washington University, St Louis, Missouri, USA (assumed from affiliation)</p> <p><b>Inclusion criteria:</b> women, suffering moderate or severe pain following episiotomy</p> <p><b>Exclusion criteria:</b> current or recent history of gastrointestinal bleeding; peptic ulcer; other GI disorders; alcohol or drug abuse; disorders of the nervous system, kidney, heart or blood; known allergies to aspirin or aspirin-like analgesics; conditions likely to interfere with absorption, distribution, metabolism, or excretion of drugs; other pain requiring narcotic analgesics; acute dermatitis or other skin lesions; past or present malignancies; taking corticosteroids or other NSAIDs, anticoagulants or other drugs that may interfere with study medication; experiencing pain due to other causes; breastfeeding</p>
Interventions	<p><b>Aspirin (N = 39 randomised)</b> 650 mg aspirin</p> <p><b>Placebo (N = 40 randomised)</b></p> <p><b>All women:</b> women received the study medication at the onset or recurrence of moderate or severe pain at least 16 but not more than 48 hours following induction of anaesthesia</p>
Outcomes	<p><b>Adequate pain relief as reported by the woman:</b> pain intensity and relief at 0.5 hours, then hourly for 8 hours was measured</p> <ul style="list-style-type: none"> <li>• Pain intensity was rated hourly on a scale of 1 to 5 (1 = no pain; 2 = mild pain; 3 = moderate pain; 4 = severe pain; 5 = very severe pain); reported SPID scores were used to calculate 'Adequate pain relief as reported by the woman' (taken over 6 hours)</li> <li>• Pain relief was rated hourly on a scale of 1 to 5 (1 = complete; 2 = a lot; 3 = some; 4 = little; 5 = no relief); reported TOTPAR scores were reported</li> <li>• Women provided their opinion of the medication on a scale 1 to 4 (1 = excellent; 2 = good; 3 = fair; 4 = poor); ratings of excellent and good were also reported</li> </ul> <p><b>Maternal adverse effects:</b> patient complaints were reported</p>
Notes	<p><b>Funding:</b> not reported</p> <p><b>Declarations of interests:</b> not reported</p> <p><b>Additional arms:</b> this was a 4-arm trial also assessed etodolac 25 mg (N = 40) and 100 mg (N = 40); we have only included the relevant arms in this review</p>

*Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: “were randomly assigned”



**Friedrich 1983** (Continued)

Allocation concealment (selection bias)	Unclear risk	Not detailed
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Described as “double-blind” but no description of whether the study medications were identical in appearance, taste, etc
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not detailed
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No losses/exclusions reported, but %/no of women ‘remaining in study’ at 4, 6 and 8 hours reported
Selective reporting (reporting bias)	Unclear risk	No access to trial registration or protocol to further assess selective reporting
Other bias	Unclear risk	Results report patients were “well matched” for a variety of baseline characteristics, but no table of these characteristics presented

**Jain 1978a**

Methods	RCT
Participants	<p><b>Setting:</b> Department of Medicine, Tulane University School of Medicine, New Orleans, Louisiana, USA (assumed from affiliation)</p> <p><b>Inclusion criteria:</b> women with moderate to severe episiotomy pain within 48 hours of a normal vaginal birth</p> <p><b>Exclusion criteria:</b> nursing; systemic diseases; allergic to aspirin</p>
Interventions	<p><b>Aspirin (N = 30 randomised)</b> 648 mg aspirin; single oral dose</p> <p><b>Placebo (N = 30 randomised)</b> Placebo; single oral dose</p> <p><b>All women:</b> the time between the test drug and previous analgesic, tranquillisers or sedatives was at least 5 hours</p>
Outcomes	<p><b>Adequate pain relief as reported by the woman:</b> a trained nurse observer rated pain intensity and relief hourly for 4 hours</p> <ul style="list-style-type: none"> <li>• Observer rating of pain intensity of 4 point scale (0 = none; 1 = slight; 2 = moderate; 3 = severe)</li> <li>• Observer rating of pain relief on 5 point scale (0 = none; 1 = slight; 2 = moderate; 3 = marked; 4 = complete)</li> <li>• Patient self-rating of pain from 0 to 1 on a continuous scale (no pain to severe pain)</li> </ul> <p>The results were not reported in such a way to calculate ‘Adequate pain relief as reported by the woman.’ Figure 2 in manuscript provides patient self-rating of pain (continuous, analogue scale)</p> <p><b>Need for additional pain relief in the first 48 hours for perineal pain:</b> need for extra</p>

**Jain 1978a** (Continued)

	analgesia during the 4-hour study period <b>Maternal adverse effects:</b> volunteered or observed side effects	
Notes	<b>Funding:</b> not reported <b>Declarations of interests:</b> not reported <b>Additional arms:</b> this was a 4-arm trial also assessed piroxicam 20 mg (N = 31) and 40 mg (N = 29); we have only included the relevant arms in this review	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Quote: "By random assignment", no other detail described
Allocation concealment (selection bias)	Unclear risk	No detail provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Quote: "double blind"; no further detail provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No detail provided
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Did not report on losses to follow up or exclusions
Selective reporting (reporting bias)	High risk	No access to trial protocol; however results reported incompletely in text
Other bias	Low risk	Baseline characteristics were reported to be comparable across groups; no other obvious sources of bias identified

**Jain 1978b**

Methods	RCT
Participants	<b>Setting:</b> Department of Medicine, Tulane University School of Medicine, USA (assumed from affiliation) <b>Inclusion criteria:</b> women with severe episiotomy pain or severe uterine cramping pain, following an uncomplicated vaginal birth <b>Exclusion criteria:</b> mild or moderate pain or baseline pain < 60% on a pain analogue; dependent on analgesics or tranquillisers; hypersensitive to salicylates or caffeine; gastrointestinal, hepatic, or renal history or a history of psychiatric illness; emotionally unstable or overtly anxious

Interventions	<p><b>Aspirin (N = 16 randomised)</b> 650 mg aspirin; single oral dose</p> <p><b>Placebo (N = 16 randomised)</b> Placebo; single oral dose</p> <p><b>All women:</b> duration between previous analgesic and test medication was at least 6 hours</p>	
Outcomes	<p><b>Adequate pain relief as reported by the woman:</b> a trained nurse observer rated pain intensity and relief hourly for 4 hours</p> <ul style="list-style-type: none"> <li>• Pain intensity: rated from 0 to 8 (0 = no pain; 2 = slight pain; 4 = moderate pain; 6 = severe pain; 8 = very severe pain)</li> <li>• Pain relief: rated from 0 to 8 (0 = worse; 2 = unchanged; 4 = less than half gone; 6 = more than half gone; 8 = complete relief)</li> <li>• Pain analogue: rated on visual analogue scale 0 = no pain; 100 = worst pain I have ever experienced</li> <li>• Subjective measure of global performance at last interview: rated from 2 to 8 (2 = poor; 4 = fair; 6 = good; 8 = very good)</li> </ul> <p>The results were not reported in such a way to calculate 'Adequate pain relief as reported by the woman'</p> <p><b>Maternal adverse effects:</b> women questioned about adverse effects at the last interview</p>	
Notes	<p><b>Funding:</b> "We wish to thank Mr. Garrett Swenson of American Home Products for the double-blind supplies of test drug, and Dr. Ilbok lee (Ives Laboratories), Dr. Bruce Schneider (Wyeth Laboratories), and Dr. Syliva Wassertheil-Smoller (Albert Einstein College of Medicine) for their assistance in the statistical analysis of data"</p> <p><b>Declarations of interest:</b> not reported</p> <p><b>Additional arms:</b> manuscript reports results of 2 randomised controlled trials; we have excluded the first, as it combined women with uterine and episiotomy pain, and did not report any results separately for the subset of women with episiotomy pain. The included trial was a 3-arm trial also assessed 800 mg aspirin and 64 mg caffeine (N = 15); we have only included the relevant arms in this review</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Quote: "Patients were divided at random", no further detail provided
Allocation concealment (selection bias)	Unclear risk	No detail provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Quote: "double blind", no further detail provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not detailed

**Jain 1978b** (Continued)

Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Complete data has not been reported and published
Selective reporting (reporting bias)	High risk	No access to trial protocol; limited data presented, results reported incompletely in text
Other bias	Unclear risk	No baseline characteristics reported

**Jain 1985**

Methods	RCT
Participants	<p><b>Setting:</b> Tulane University School of Medicine, New Orleans, Louisiana, USA (assumed from affiliation)</p> <p><b>Inclusion criteria:</b> postpartum women who had undergone episiotomy and requested analgesic medication for pain of at least moderate intensity, aged <math>\geq 18</math> years</p> <p><b>Exclusion criteria:</b> receipt of analgesics or tranquillisers within 4 hours of stud entry; planned to breast feed; history of convulsive disorders, known peptic ulcer, renal, hepatic or haematological disease; known allergic reactions to salicylates or other NSAIDs</p>
Interventions	<p><b>Aspirin (N = 30 randomised)</b> 600 mg aspirin; single dose of 2 matching capsules</p> <p><b>Placebo (N = 30 randomised)</b> Placebo; single dose of 2 matching capsules</p> <p><b>All women:</b> the test drug was given in a single dose in the form of 2 matching capsules</p>
Outcomes	<p><b>Adequate pain relief as reported by the woman:</b> a trained nurse observed recorded pain intensity and relief at 0.5 hours and hourly to 5 hours</p> <ul style="list-style-type: none"> <li>• Pain intensity was rated from 0 to 3 (0 = no pain; 1 = mild pain; 2 = moderate pain; 3 = severe pain); SPID scores were reported and used to calculate 'Adequate pain relief as reported by the woman' (taken over 5 hours)</li> <li>• Pain relief was rated from 0 to 4 (none; 1 = a little; 2 = some; 3 = a lot; 4 = complete); TOTPAR scores were also reported</li> <li>• Women's overall rating of the medication's efficacy was also reported, rated from 0 to 3 (0 = poor; 1 = fair; 2 = good; 3 = excellent)</li> </ul> <p><b>Need for additional pain relief in the first 48 hours for perineal pain:</b> need for supplemental analgesia in 5-hour study period</p> <p><b>Maternal adverse effects:</b> adverse effects reported by women or observed by the nurse were recorded</p>
Notes	<p><b>Funding sources:</b> "Supported in part by a grant from Adria Laboratories, inc., Columbus, Ohio"</p> <p><b>Declarations of interests:</b> not reported</p> <p><b>Additional arms:</b> this was a 4-arm trial also assessed indoprofen 50 (N = 30) and 100 mg (N = 30); we included only the relevant arms in this review</p>

*Risk of bias*

**Jain 1985** (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "randomly assigned", no further detail provided
Allocation concealment (selection bias)	Unclear risk	No detail provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quotes: "double blind" "the test drug was given in a single dose in the form of two matching capsules"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Not specifically stated; reasonable to assume women and the nurse were blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "Any patients who experienced inadequate pain relief were permitted to remedicate with an alternate analgesic. In such cases pain evaluations were discontinued and for the balance of the study, patients were assigned a pain intensity score equal to that at the time of remedication and pain relief scores of zero" ... There were however, no women who required re-medication
Selective reporting (reporting bias)	Unclear risk	No access to trial registration or protocol to further assess selective reporting
Other bias	Low risk	Baseline characteristics comparable between groups; no other obvious sources of bias identified

**London 1983a**

Methods	RCT
Participants	<p><b>Setting:</b> Sinai Hospital of Baltimore, MD, USA (assumed from affiliation)</p> <p><b>Inclusion criteria:</b> women with no systemic medical illness, experiencing moderate to severe episiotomy pain within 48 hours following an otherwise uncomplicated vaginal birth</p> <p><b>Exclusion criteria:</b> "those used by Hermann et al"</p>
Interventions	<p><b>Aspirin (N randomised was unclear; N = 40 analysed)</b> 650 mg aspirin; single dose in identical capsules</p> <p><b>Placebo (N randomised was unclear; N = 40 analysed)</b> Placebo; single dose in identical capsules</p>
Outcomes	<p><b>Adequate pain relief as reported by the woman:</b> one investigator assessed pain intensity and relief hourly for 6 hours;</p> <ul style="list-style-type: none"> <li>Though SPID and TOTPAR scores were reported, the scales used to measure pain intensity and relief were not reported, and thus these data could not be used to calculate 'Adequate pain relief as reported by the woman'</li> </ul>

London 1983a (Continued)

	<ul style="list-style-type: none"> <li>Women's "overall impression" was reported (excellent, very good, good, fair and poor); excellent, very good and good ratings were used to calculate 'Adequate pain relief as reported by the woman' (taken at 6 hours)</li> </ul> <p><b>Need for additional pain relief in the first 48 hours for perineal pain:</b> frequency of re-medication</p> <p><b>Maternal adverse effects:</b> women were observed hourly for adverse reactions</p>
Notes	<p><b>Funding:</b> "We would like to acknowledge the support of Sandoz, Inc. in this study"</p> <p><b>Declarations of interests:</b> not reported</p> <p><b>Additional arms:</b> this was a 4-arm trial also assessed fluproquazone 100 mg (N = 41 analysed) and 200 mg (N = 39 analysed); we included only the relevant arms in this review</p>

*Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "randomly assigned"
Allocation concealment (selection bias)	Unclear risk	No detail provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quotes: "double-blind"... "All medication was supplied in identical capsules... packaged in individually sealed envelopes"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Considered reasonable to assume blinding
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	In the whole trial, there were 166 who entered, and 160 provided "valid data for analyses"; other losses/exclusions not clearly reported
Selective reporting (reporting bias)	Unclear risk	No access to trial protocol to further assess; scales used to assess pain intensity and relief (needed to use SPID and TOTPAR scores to calculate adequate pain relief) were not reported
Other bias	Unclear risk	Baseline characteristics not reported

London 1983b

Methods	RCT
Participants	<p><b>Setting:</b> Sinai Hospital, Baltimore, MD, USA (assumed from affiliation); between July and December 1980</p> <p><b>Inclusion criteria:</b> postpartum women with moderate to severe episiotomy pain</p> <p><b>Exclusion criteria:</b> allergy to salicylates; asthma; history of chronic use of analgesics, alcohol, tranquillisers, or other drugs; blood dyscrasia; gastrointestinal disorders; hepatic</p>

London 1983b (Continued)

	and/or renal disease; psychiatric illness
Interventions	<p><b>Aspirin group 1 (N randomised not reported; N = 40 analysed)</b> 5 grains (300 mg) aspirin; 4 tablet single dose</p> <p><b>Aspirin group 2 (N randomised not reported; N = 41 analysed)</b> 10 grains (600 mg) aspirin; 4 tablet single dose</p> <p><b>Aspirin group 3 (N randomised not reported; N = 40 analysed)</b> 20 grains (1200 mg) aspirin; 4 tablet single dose</p> <p><b>Placebo (N randomised not reported; N = 39 analysed)</b> Placebo; 4 tablet single dose</p> <p><b>All women:</b> if women had been medicated previously for pain, the experimental protocol was not initiated for 4 hours</p>
Outcomes	<p><b>Adequate pain relief as reported by the woman:</b> trained research nurse investigator questioned women at 0.5 hours and hourly for 4 hours regarding pain intensity which was recorded on a 4 point scale (0 = none; 1 = slight pain; 2 = moderate pain; 3 = severe pain). Pain intensity scores were provided in Table 1 from 0 to 4 hours and were thus used to calculate SPID scores and 'Adequate pain relief as reported by the woman' (taken over 4 hours)</p> <p><b>Need for additional pain relief in the first 48 hours for perineal pain:</b> women re-medicated for episiotomy pain within 4-hour study period</p> <p><b>Maternal adverse effects:</b> side effects observed or reported</p>
Notes	<p><b>Funding:</b> not reported</p> <p><b>Declarations of interests:</b> not reported</p> <p><b>Note:</b> we combined the 3 aspirin groups for the main analysis</p>

*Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "randomly assigned"
Allocation concealment (selection bias)	Unclear risk	No detail provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quotes: "prospective blind study"... "Each single four-tablet dose was individually packaged and identified only by study and patient number, and all tablets appeared identical"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	As above; considered reasonable to assume blinding
Incomplete outcome data (attrition bias) All outcomes	High risk	"Because of protocol ineffectiveness, participation in the study was terminated if the patient requested additional analgesic medication, topical analgesics, or Sitz baths. In such cases pain intensity was measured at all intervals up to the time of termination"... 6/121 women in the aspirin groups and 10/39 in the

London 1983b (Continued)

		placebo groups required re-medication. No clear reporting of other losses/exclusions
Selective reporting (reporting bias)	Unclear risk	No access to trial protocol to further assess risk of selective reporting
Other bias	Unclear risk	Baseline characteristics not reported

Mukherjee 1980

Methods	RCT
Participants	<p><b>Setting:</b> LNJP Hospital, New Delhi, India (assumed from affiliation)</p> <p><b>Inclusion criteria:</b> women from an otherwise healthy population whose chief complaint was moderate to severe pain following episiotomy on the first postoperative morning</p> <p><b>Exclusion criteria:</b> known hypersensitivity to dipyrone and aspirin; receipt of any analgesics 8 hours before entry to the study</p>
Interventions	<p><b>Aspirin (N = 90 randomised)</b> 500 mg aspirin; single oral dose in identical tablet form</p> <p><b>Placebo (N = 88 randomised)</b> Placebo; single oral dose in identical tablet form</p> <p><b>All women:</b> nothing was permitted to be taken orally for the first hour after treatment administration</p>
Outcomes	<p><b>Adequate pain relief as reported by the woman:</b></p> <ul style="list-style-type: none"> <li>A research worker interviewed women at 0.5 hours, and hourly for 6 hours. Women were asked “By how many paise in the rupee is your pain less?”; pain relief was arbitrarily equated as 25% = slight (given a score of 1), 50% = moderate (given a score of 2), 75% = marked (given a score of 3, and 100% = complete (given a score of 4). Mean pain relief scores from 0 to 6 hours were provided in Figure 2, and were thus used to calculate TOTPAR scores and to calculate ‘Adequate pain relief as reported by the woman’ (taken over 6 hours)</li> <li>More than 50% pain relief was also reported</li> </ul> <p><b>Maternal adverse effects:</b> adverse drug reactions</p>
Notes	<p><b>Funding:</b> not reported</p> <p><b>Declarations of interests:</b> not reported</p> <p><b>Additional arms:</b> this was a 3 arm trial also assessed dipyrone 500 mg (N = 89); we have only included the relevant arms in this review</p>

*Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: “allocated at random”



**Mukherjee 1980** (Continued)

Allocation concealment (selection bias)	Unclear risk	No detail provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: “approved double-blind approach, which was strictly adhered to” “identical tablet form”
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Reasonable to assume blinding
Incomplete outcome data (attrition bias) All outcomes	Low risk	No losses/exclusions
Selective reporting (reporting bias)	Unclear risk	No access to trial protocol to further assess risk selective reporting
Other bias	Low risk	Figures reported for baseline characteristics (such as pain severity, age, weight and height at baseline), and reported “all three groups were also comparable...”

**Okun 1982**

Methods	RCT
Participants	<p><b>Setting:</b> Cedars-Sinai Medical Center, Los Angeles, California, USA (assumed from affiliation)</p> <p><b>Inclusion criteria:</b> hospitalised women with moderate, severe or very severe pain due to uterine cramps or episiotomy within 48 hours of delivery (94 women in total with episiotomy pain)</p> <p><b>Exclusion criteria:</b> breastfeeding; receipt of any analgesic, sedative or psychotropic medication within 6 hours before administration of the study drug</p>
Interventions	<p><b>Aspirin (N = 20 randomised)</b> 650 mg aspirin; single oral dose of 2 x 325 mg identical looking capsules</p> <p><b>Placebo (N = 18 randomised)</b> Placebo; single oral dose of 2 identical looking capsules</p>
Outcomes	<p><b>Adequate pain relief as reported by the woman:</b> 1 nurse observer recorded pain intensity hourly for 8 hours; intensity was rated from 1 to 5 (1 = no pain; 2 = mild pain; 3 = moderate pain; 4 = severe pain; 5 = very severe pain); pain intensity differences were calculated, as were SPID scores at 4, 6, and 8 hours; the time of maximum pain relief; duration of pain relief; the proportion of women with at least 50% pain relief 1 and 2 hours after treatment</p> <p><b>Need for additional pain relief in the first 48 hours for perineal pain:</b> proportion requiring additional analgesics</p> <p><b>Adverse effects:</b> adverse effects mentioned by women were recorded</p> <p>No data included in the meta-analyses as results were not reported separately for women with post-episiotomy pain</p>

Okun 1982 (Continued)

Notes	<p><b>Funding:</b> not reported</p> <p><b>Declarations of interests:</b> not reported</p> <p><b>Additional arms:</b> this was a 5-arm trial also assessed fendosal 100 (N = 19), 200 (N = 19) and 400 mg (N = 18); we included only the relevant arms in this review</p> <p><b>Note:</b> trial also included women with postpartum uterine cramps pain; not included in review (157/250); no data were able to be included in the meta-analyses as results were not reported separately for women with post-episiotomy pain</p>
-------	---

*Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Assignment to treatment was randomised"
Allocation concealment (selection bias)	Unclear risk	No detail provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quotes: "double-blind" and "identical looking capsules"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	As above; 1 nurse observer recorded the pain intensity scores prior and hourly after administration of medication, reasonable to assume that outcome assessment is blinded
Incomplete outcome data (attrition bias) All outcomes	High risk	A total of 250 eligible women were "admitted to the study"... "The PI scores from patients taking another analgesic were handled as treatment failures by substituting the initial PI score for all hours after the analgesic was taken"
Selective reporting (reporting bias)	Unclear risk	Results (such as SPID scores) were reported incompletely in text
Other bias	Low risk	Baseline characteristics (such as initial intensity and type of pain; age and weight) were comparable between groups

Olson 1997

Methods	RCT
Participants	<p><b>Setting:</b> Hospital Maternidad Concepcion Palacios, Caracas, Venezuela</p> <p><b>Inclusion criteria:</b> women of legal age (aged <math>\geq 18</math> years), who were able to communicate meaningfully with the nurse-observer, who were hospitalised and had severe post episiotomy pain after an uncomplicated birth and could tolerate oral medications</p> <p><b>Exclusion criteria:</b> planning to breast feed within 24 hours after administration of the study medications; serious complicating illness or abnormal postpartum bleeding, with active peptic ulcer disease or other gastrointestinal disease associated with blood loss; receipt of any other investigational drug within the 1 month prior; history of drug or alcohol abuse; known allergic sensitivities to aspirin, diclofenac, or other NSAIDs</p>

Interventions	<p><b>Aspirin (N = 50 randomised)</b> 650 mg aspirin; single oral dose of 2 x 325 mg capsules; and 3 placebo tablets</p> <p><b>Placebo (N = 52 randomised)</b> Placebo; single oral dose of 2 placebo capsule and 3 placebo tablets</p> <p><b>All women:</b> each woman received a single unit dose consisting 2 capsules and 3 tablets, with at least 8 ounces of water; women were asked to sit up or lie on their right side for 2 hours after administration. No medications (analgesics, sedatives, hypnotics, tranquillisers) were permitted concomitantly or during the 4 hours prior to taking the medication</p>
Outcomes	<p><b>Adequate pain relief as reported by the woman:</b> the same nurse-observer interviewed the women at 0.5 hours, and hourly for 8 hours</p> <ul style="list-style-type: none"> <li>• Women assessed their episiotomy pain on a scale of 0 to 3 (none = 0; slight pain = 1; moderate pain = 2; severe pain = 3); SPID scores were reported, and used to calculate 'Adequate pain relief as reported by the woman' (taken at 4 hours; also reported at 8 hours)</li> <li>• Women were asked to classify pain relief on a scale of 0 to 4 (none = 0; a little = 25% (1); some = 50% (2); a lot = 75% (3); complete = 100% (4); TOTPAR scores were also reported</li> <li>• Women were asked to rate the study medication and assess their overall improvement; study medication: 0 = poor; 1 = fair; 2 = good; 3 = excellent; overall improvement: 1 = very much worse; 7 = very much better</li> </ul> <p><b>Need for additional pain relief in the first 48 hours for perineal pain:</b> re-medication within 8-hour study period</p> <p><b>Maternal adverse effects:</b> adverse effects were recorded if they were observed or volunteered</p>
Notes	<p><b>Funding:</b> "This work was supported in part by a grant from the Ciba-Geigy Corporation, Summit, NJ"</p> <p><b>Declarations of interests:</b> not reported; though first and second authors affiliated to "Analgesic Development Ltd."</p> <p><b>Additional arms:</b> this was a 5-arm trial also assessed diclofenac potassium 25 mg (N = 52), 50 mg (N = 50) and 100 mg (N = 51); we have only included the relevant arms in this review</p>

*Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "randomly assigned... computer program generated a random permutation such that two patients received each treatment"
Allocation concealment (selection bias)	Unclear risk	No detail provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quotes: "double-blind" and "Each patient received a single-unit dose consisting of 3 tablets and 2 capsules... all unit doses were identical in appearance and packaging"

Olson 1997 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	1 nurse observer involved in outcome assessment
Incomplete outcome data (attrition bias) All outcomes	High risk	If a woman wished to withdraw before the first hour because of inadequate relief, a non-study analgesic was administered, and she was discontinued from the study (none were re-medicated in this first hour; therefore no discontinuations); if a woman required additional analgesic after the first hour, she was included, and relief scores of 0 and intensity scores equal to the pain at time of re-medication were assumed for the duration (5/50 in the aspirin group 19/52 in the placebo group)
Selective reporting (reporting bias)	Unclear risk	No access to trial protocol to further assess risk of selective reporting
Other bias	Low risk	Baseline characteristics reported (age, weight, height, parity, days post-delivery) were comparable between groups; no other obvious sources of bias identified

Sunshine 1983a

Methods	RCT
Participants	<p><b>Setting:</b> Hospital Maternidad Concepcion Palacios, Caracas, Venezuela</p> <p><b>Inclusion criteria:</b> women with severe post-episiotomy pain after an uncomplicated birth; aged <math>\geq 18</math> years, who could tolerate oral medications</p> <p><b>Exclusion criteria:</b> known allergic sensitivities to study medication; abnormal postpartum bleeding, or complicating illnesses; breastfeeding; history of drug dependence; receipt of other investigational drugs prior to enrolment</p>
Interventions	<p><b>Aspirin (N = 30 randomised)</b> 600 mg aspirin; single oral dose of 1 aspirin capsule and 1 placebo tablet</p> <p><b>Placebo (N = 30 randomised)</b> Placebo; single oral dose of 1 capsule and 1 tablet</p> <p><b>All women:</b> as a single dose; women were given the study medication by the nurse observer when their pain was severe; no medications that might alter the response to the study analgesics were permitted concomitantly or during the 4 hours before the test medication was taken</p>
Outcomes	<p><b>Adequate pain relief as reported by the woman:</b> the same nurse observer interviewed at the time of medication, 0.5 hours, and hourly for 4 hours</p> <ul style="list-style-type: none"> <li>• Women were asked to classify the intensity of their pain on a scale of 0 to 3 (0 = none; 1 = slight pain; 2 = moderate pain; 3 = severe pain); SPID scores were reported and used to calculate 'Adequate pain relief as reported by the woman' (taken over 4 hours)</li> <li>• Women were asked to estimate their percentage of pain relief from 0 to 4 (0 = none; 25% = 1; 50% = 2; 75% = 3; 100% = 4); total scores were also reported</li> </ul>

	<ul style="list-style-type: none"> <li>Women also were asked to estimate the time to onset of effect; to rate their overall improvement on a 7 point scale (1 = very much worse; 2 = much worse; 3 = a little worse; 4 = no change; 5 = a little better; 6 = much better; 7 = very much better); and to rate the study medication on a 4 point scale (0 = poor; 1 = fair; 2 = good; 3 = excellent)</li> </ul> <p><b>Need for additional pain relief in the first 48 hours for perineal pain:</b> re-medication within 4 hours</p> <p><b>Maternal adverse effects:</b> adverse reactions were noted if observed or volunteered</p>	
Notes	<p><b>Funding:</b> “A grant-in-aid and test medication from Upjohn Company made this research possible”</p> <p><b>Declarations of interests:</b> not reported</p> <p><b>Additional arms:</b> this was a 4-arm trial also assessed zomepirac 100 mg (N = 30) and ibuprofen 400 mg (N = 30); we have only included the relevant arms in this review</p>	
<i>Risk of bias</i>		
Bias	Authors’ judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: “In a randomised study”
Allocation concealment (selection bias)	Unclear risk	No detail provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: “in a double-blind fashion. Because the study medications were not identical in appearance, a double-dummy technique was used”; each woman received one tablet and one capsule as appropriate
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The same nurse observer interviewed the patients
Incomplete outcome data (attrition bias) All outcomes	High risk	If before the first hour a woman reported inadequate pain relief, a conventional analgesic was given and she was removed from the study; if a woman requested ‘rescue’ medication after the first hour she was given it and was included in the evaluation; responses at the time of re-medication were assumed for the duration of the study; all 120 women who participated in the study were included in the analysis; 5 women who received placebo required rescue medication during the study; no women receiving aspirin required re-medication
Selective reporting (reporting bias)	High risk	Some incomplete reporting “The three drugs were much the same for mean onset, duration and time to peak values. The hypothesis that there is no difference among treatments was rejected at the 0.05 level or better for all variables”; patients rating of overall improvement and of study medication mentioned in methods and not reported

Sunshine 1983a (Continued)

Other bias	Low risk	Baseline characteristics presented were comparable between groups; no other obvious sources of bias identified
------------	----------	--

Sunshine 1983b

Methods	RCT
Participants	<p><b>Setting:</b> Hospital Maternidad Concepcion Palacios, Caracas, Venezuela</p> <p><b>Inclusion criteria:</b> women with moderate or severe post-episiotomy pain after an uncomplicated delivery, who could tolerate oral medication, aged <math>\geq 18</math> years</p> <p><b>Exclusion criteria:</b> breastfeeding; any complicating illness or abnormal postpartum bleeding; receipt of any other investigational drug within 1 month prior to enrolment; history of drug dependence or known allergic sensitivities to prolonic acid derivatives or aspirin</p>
Interventions	<p><b>Aspirin (N = 29 randomised)</b> 600 mg aspirin; single oral dose of 5 identical tablets</p> <p><b>Placebo (N = 31 randomised)</b> Placebo; single oral dose of 5 identical tablets</p> <p><b>All women:</b> no medications that might confound the interpretation of the efficacy and/or adverse effect liability of the study analgesics were permitted concomitantly or during the 4 hours before taking the study medication</p>
Outcomes	<p><b>Adequate pain relief as reported by the woman:</b> the same nurse-observer interviewed the women at the time medication was administered and hourly for 6 hours:</p> <ul style="list-style-type: none"> <li>• Women were asked to assess the intensity of their pain from 0 to 3 (0 = none; 1 = light pain; 2 = moderate pain; 3 = severe pain); SPID scores were reported and used to calculate 'Adequate pain relief as reported by the woman' (taken over 6 hours)</li> <li>• Women were asked to classify their degree of pain relief from 0 to 4 (0 = none; 25% (1) = a little; 50% (2) = some; 75% (3) = a lot; 100% (4) = complete); total scores were also reported</li> </ul> <p><b>Need for additional pain relief in the first 48 hours for perineal pain:</b> re-medication within 6-hour study period</p> <p><b>Maternal adverse effects:</b> adverse reactions were noted if they were observed or volunteered</p>
Notes	<p><b>Funding:</b> "Supported by a grant from Boots Pharmaceuticals, Inc"</p> <p><b>Declarations of interests:</b> not reported</p> <p><b>Additional arms:</b> this was a 5-arm trial also assessed flurbiprofen 25 (N = 32), 50 (N = 29), 100 mg (N = 31); we included only the relevant arms in this review</p>

*Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "In each successive block of ten patients a computer program generated a random permutation such that, two patient received each treatment"

Sunshine 1983b (Continued)

Allocation concealment (selection bias)	Unclear risk	No detail provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: “double-blind... Each patient was given one dose of five tablets that were identical in appearance and packaging”
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Same nurse interviewer interviewed patients at administration and hourly afterwards; reasonable to assume blinding of outcome assessment
Incomplete outcome data (attrition bias) All outcomes	High risk	If women reported inadequate pain relief before the first hour, a conventional analgesic agent was given and they were removed from the study; if women requested rescue medication after the first hour, they were given the conventional analgesic and included in the analyses - baseline pain intensity and zero pain relief were assumed for the duration of scheduled observations; 168 women were enrolled in the study; 16 were “dropped from the analysis because they received concomitant oxytocic medication that the sponsor felt might confound the interpretation of the efficacy of the study drug” 1/32 in the placebo group; 4/33 in the aspirin group; no women re-medicated in the first hour; 14 re-medicated in placebo group; 1 re-medicated in the aspirin group
Selective reporting (reporting bias)	Unclear risk	No access to trial protocol to further assess selective reporting
Other bias	Low risk	Baseline characteristics reported were balanced between groups; no other obvious risk of bias identified

Trop 1983

Methods	RCT
Participants	<p><b>Setting:</b> “in an obstetric and gynecology unit”; Montreal, Quebec, Canada (assumed from affiliation)</p> <p><b>Inclusion criteria:</b> women who had an episiotomy</p> <p><b>Exclusion criteria:</b> receipt of tranquillisers, sedatives, hypnotics or other analgesics during the 4 hours preceding the study; breastfeeding</p>
Interventions	<p><b>Aspirin (N = not reported)</b> 1200 mg aspirin; single dose of 4 x 300 mg tablets</p> <p><b>Aspirin (N = not reported)</b> 600 mg aspirin; single dose of 2 x 300 mg tablets and 2 placebo tablets</p> <p><b>Placebo (N = not reported)</b> Placebo; single dose of 4 placebo tablets</p> <p><b>All women:</b> the medication was administered 10.5 to 14.4 hours after episiotomy, upon request by the woman or when pain was judged moderate to severe by the nurse</p>

Outcomes	<p><b>Adequate pain relief as reported by the woman:</b> severity of pain was judged using a 30 cm visual analogue scale (no pain, slight, moderate, severe, unbearable); the women registered the intensity of pain by putting a stroke on the place on the scale before drug administration and every hour for 4 hours (women were not allowed to see the result of their previous assessment); the research nurse independently recorded her own evaluation of the analgesic effect of the medication (scale of 0 to 4: no pain, to worse than before)</p> <p><b>Need for additional pain relief in the first 48 hours for perineal pain:</b> additional analgesic during 4-hour period</p> <p><b>Maternal adverse effects:</b> side effects reported by the women were noted on case report forms</p>
Notes	<p><b>Funding:</b> “This study was supported by a grant from Roussel Canada Inc., Montreal, Quebec, Canada”</p> <p><b>Declarations of interests:</b> not reported; though last author affiliated to “Roussel Canada Inc.”</p> <p><b>Additional arms:</b> this was a 5-arm trial also assessed tiaprofenic acid 200 mg (N = not reported) and 400 mg (N = not reported); we included only relevant arms in this review</p> <p><b>Note:</b> no data could be included in the meta-analyses as numbers for each group were not reported</p>

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: “randomised”
Allocation concealment (selection bias)	Unclear risk	No detail provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quotes: “double-blind”... “The drugs were prepared as tablets... all were of identical appearance”
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quotes: “double-blind”... “The drugs were prepared as tablets... all were of identical appearance”
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Not clearly reported (nor were the numbers of women in each group)
Selective reporting (reporting bias)	High risk	Results incompletely reported within text (including nurse's evaluation of analgesic effect); numbers in each group also not reported
Other bias	Unclear risk	Baseline characteristics incompletely reported in text “There were no significant differences among the 5 groups with respect to age, height, weight or vital signs”



GI: gastro-intestinal  
 NSAIDs: non-steroidal anti-inflammatory drugs  
 RCT: randomised controlled trial  
 SPID: summed pain intensity differences  
 TOTPAR: total pain relief

### Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
<a href="#">Bruni 1965</a>	RCT. Included women with “postpartum pain”, not exclusively women with perineal pain; results not reported separately for women with perineal pain
<a href="#">Gindhart 1971</a>	RCT. Did not assess aspirin for perineal pain, and rather assessed 2 agents: 1 (Darvon), containing propoxyphene with aspirin, phenacetin, and caffeine; and 1 (Fiorinal) which combined aspirin, phenacetin, and caffeine with a mild sedative, butalbital
<a href="#">Gruber 1979</a>	RCT. Included women with “postpartum pain”, not exclusively women with perineal pain; results not reported separately for women with perineal pain
<a href="#">Moggian 1972</a>	RCT. Included women with “postpartum pain”, not exclusively women with perineal pain; results not reported separately for women with perineal pain
<a href="#">Prockop 1960</a>	RCT. Did not assess aspirin for perineal pain, and rather assessed an aspirin compound (acetophenetidin acetylsalicylic acid and caffeine); codeine and an aspirin compound; dextropropoxyphene; dextropropoxyphene and an aspirin compound; and a starch placebo
<a href="#">Rubin 1984</a>	RCT. Did not assess aspirin for perineal pain, and rather assessed an aspirin/caffeine combination; an acetaminophen/aspirin combination; acetaminophen alone; and placebo
<a href="#">Santiago 1959</a>	Not RCT: “In consecutive cases of vaginal delivery with episiotomy, orders in alternating patients for analgesic 1 or analgesic 2... were written”. This study assessed Darvon Compound (dextro propoxyphene and acetylsalicylic acid compound); and a preparation containing acetylsalicylic acid, acetophenetidin, caffeine and codeine phosphate. Aspirin not assessed
<a href="#">Sunshine 1983c</a>	RCT. Included women with “postpartum pain”, not exclusively women with perineal pain; results not reported separately for women with perineal pain
<a href="#">Sunshine 1985</a>	RCT. Included women with “postpartum pain”, not exclusively women with perineal pain; results not reported separately for women with perineal pain
<a href="#">Van der Pas 1984</a>	Did not assess single dose aspirin, and rather assessed twice daily acetylsalicylic acid; and naproxen; it was not clear whether this was a randomised controlled trial

RCT: randomised controlled trial

## Characteristics of studies awaiting assessment *[ordered by study ID]*

### Bhounsule 1990

Methods	Unclear
Participants	100 women with post-episiotomy pain
Interventions	Single oral doses of 400 mg ibuprofen, 500 mg analgin, 500 mg paracetamol, 600 mg aspirin, and placebo
Outcomes	Pain intensity; pain relief; side effects
Notes	Awaiting classification pending further details regarding allocation

### Sunshine 1989

Methods	Unclear
Participants	Unclear
Interventions	Unclear
Outcomes	Unclear
Notes	No access to trial publication; awaiting classification pending availability of manuscript

## DATA AND ANALYSES

### Comparison 1. Aspirin versus placebo for perineal pain

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Adequate pain relief as reported by the woman	13	1001	Risk Ratio (M-H, Fixed, 95% CI)	2.03 [1.69, 2.42]
1.1 300 mg aspirin	1	53	Risk Ratio (M-H, Fixed, 95% CI)	2.6 [0.36, 18.88]
1.2 500 to 650 mg aspirin	11	800	Risk Ratio (M-H, Fixed, 95% CI)	1.98 [1.64, 2.39]
1.3 900 mg aspirin	1	40	Risk Ratio (M-H, Fixed, 95% CI)	1.83 [0.84, 3.99]
1.4 1200 mg aspirin	3	108	Risk Ratio (M-H, Fixed, 95% CI)	2.75 [1.25, 6.06]
2 Need for additional pain relief	10	744	Risk Ratio (M-H, Fixed, 95% CI)	0.25 [0.17, 0.37]
2.1 300 mg aspirin	1	53	Risk Ratio (M-H, Fixed, 95% CI)	0.16 [0.03, 0.79]
2.2 500 to 650 mg aspirin	9	569	Risk Ratio (M-H, Fixed, 95% CI)	0.27 [0.17, 0.41]
2.3 900 mg aspirin	1	40	Risk Ratio (M-H, Fixed, 95% CI)	0.14 [0.01, 2.60]
2.4 1200 mg aspirin	2	82	Risk Ratio (M-H, Fixed, 95% CI)	0.20 [0.06, 0.70]
3 Maternal adverse effects	14	1067	Risk Ratio (M-H, Fixed, 95% CI)	1.08 [0.57, 2.06]
3.1 300 mg aspirin	1	53	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.2 500 to 650 mg aspirin	13	892	Risk Ratio (M-H, Fixed, 95% CI)	1.13 [0.51, 2.53]
3.3 900 mg aspirin	1	40	Risk Ratio (M-H, Fixed, 95% CI)	2.5 [0.55, 11.41]
3.4 1200 mg aspirin	2	82	Risk Ratio (M-H, Fixed, 95% CI)	0.10 [0.01, 1.80]

### Comparison 2. 300 mg aspirin versus 600 mg aspirin for perineal pain

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Adequate pain relief as reported by the woman	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
2 Need for additional pain relief	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
3 Maternal adverse effects	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only

### Comparison 3. 600 mg aspirin versus 1200 mg aspirin for perineal pain

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Adequate pain relief as reported by the woman	2	121	Risk Ratio (M-H, Fixed, 95% CI)	0.85 [0.52, 1.39]
2 Need for additional pain relief	2	121	Risk Ratio (M-H, Fixed, 95% CI)	1.32 [0.30, 5.68]
3 Maternal adverse effects	2	121	Risk Ratio (M-H, Fixed, 95% CI)	3.0 [0.13, 69.52]