



RESEARCH ARTICLE

**REVISED** Presence and diversity of *Salmonella* isolated from layer farms in central Ecuador [version 2; peer review: 2 approved]

Previously titled: Prevalence and diversity of *Salmonella* isolated from layer farms in central Ecuador

Gabriela A. Salazar<sup>1</sup>, Ricardo Guerrero-López<sup>1</sup>, Liliana Lalaleo<sup>1</sup>,  
 Diana Avilés-Esquivel<sup>1</sup>, Christian Vinueza-Burgos<sup>2</sup>, William Calero-Cáceres <sup>1</sup>

<sup>1</sup>UTA RAM OneHealth Group, Faculty of Agricultural Sciences, Universidad Técnica de Ambato, Cevallos, Ecuador

<sup>2</sup>Facultad de Medicina Veterinaria y Zootecnia, Universidad Central del Ecuador, Quito, Ecuador

**v2** First published: 28 Feb 2019, 8:235 (<https://doi.org/10.12688/f1000research.18233.1>)  
 Latest published: 09 Apr 2019, 8:235 (<https://doi.org/10.12688/f1000research.18233.2>)

**Abstract**

**Background:** Given the considerable role played by *Salmonella* in the incidence of food contamination, around the world, surveillance of this infection is prioritized by both food producers and health care authorities. Data remains insufficient concerning the prevalence of *Salmonella* in poultry systems in Ecuador and in Latin America in general.

**Methods:** In this study, we evaluated the presence and diversity of *Salmonella* serovars in samples taken from 21 layer farms and backyard layers in central Ecuador during August-November 2017. *Salmonella* was isolated following standardized methods (ISO 6579) and the serovar determination was carried out by PCR.

**Results:** A significant presence of *Salmonella* was detected in the 21 farms evaluated, with a frequency of 76% (95% confidence interval (CI): 53-92) in environmental surfaces, 33% (95%CI: 15-57) in pooled cloacal swabs from layer hens, 33% (95% CI: 13-59) on feed samples, and 10% (95%CI: 1-30) in backyard layer feces from traditional local markets. The dominant serovars detected were *S. Infantis* and *S. Typhimurium*.

**Conclusions:** This study forms a basis for further surveillance of *Salmonella* serovars in layer farms in central Ecuador.

**Keywords**

Salmonella, Layer Poultry, Ecuador, Serovars.

**Open Peer Review**

Referee Status:

	Invited Referees	
	1	2
<b>REVISED</b>		
<b>version 2</b>		report
published 09 Apr 2019		
<b>version 1</b>		
published 28 Feb 2019	report	report

- 1 **Tania Gaviria Cantín**, Universidad de Santiago de Cali, Colombia
- 2 **Eloy Gonzalez-Gustavson** , Universidad Nacional Mayor de San Marcos, Peru

Any reports and responses or comments on the article can be found at the end of the article.

**Corresponding author:** William Calero-Cáceres ([wr.calero@uta.edu.ec](mailto:wr.calero@uta.edu.ec))

**Author roles:** **Salazar GA:** Formal Analysis, Investigation, Writing – Original Draft Preparation; **Guerrero-López R:** Conceptualization, Supervision, Validation; **Lalaleo L:** Formal Analysis, Investigation, Writing – Review & Editing; **Avilés-Esquivel D:** Methodology, Validation; **Vinueza-Burgos C:** Formal Analysis, Methodology, Writing – Original Draft Preparation; **Calero-Cáceres W:** Conceptualization, Formal Analysis, Funding Acquisition, Investigation, Methodology, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

**Competing interests:** No competing interests were disclosed.

**Grant information:** This study was supported by the Dirección de Investigación y Desarrollo DIDE-Universidad Técnica de Ambato (Project 1568-CU-P-2017).

*The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

**Copyright:** © 2019 Salazar GA *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**How to cite this article:** Salazar GA, Guerrero-López R, Lalaleo L *et al.* **Presence and diversity of *Salmonella* isolated from layer farms in central Ecuador [version 2; peer review: 2 approved]** F1000Research 2019, 8:235 (<https://doi.org/10.12688/f1000research.18233.2>)

**First published:** 28 Feb 2019, 8:235 (<https://doi.org/10.12688/f1000research.18233.1>)

**REVISED** Amendments from Version 1

- Considering the sampling procedures, the term "presence" is more accurate in relation to "prevalence".
- Additional information is provided about the selected genes for PCR serovar identification.
- The general conditions for the PCR procedures were added.
- The confidence intervals were calculated using the binomial distribution.

See referee reports

## Introduction

The genus *Salmonella* is considered a leading cause of foodborne illnesses around the world (WHO, 2017). These bacteria are among the most significant agents of food and water poisoning in the United States and Europe (Bäumler *et al.*, 2000; Callejón *et al.*, 2015; Varma *et al.*, 2005). Globally, it is estimated that 93.8 million cases of *Salmonella*-related gastroenteritis occur annually, resulting in 155,000 deaths (Majowicz *et al.*, 2010). In Ecuador, typhoid, together with paratyphoid fever, causes around 1,500 hospitalizations per year, while non-typhoidal salmonellosis leads to more than 2,000 hospitalizations over the same period. These infections have accounted for approximately 25% of total reported gastrointestinal illnesses in recent years (MSP, 2018). *Salmonella* represents a complex and diverse genus, but only a small number of serovars are involved in human infections (Issenhuth-Jeanjean *et al.*, 2014; Thiennimitr *et al.*, 2012).

*Salmonella* detection and the investigation of foodborne outbreaks is of maximum importance to public health, which has resulted in the establishment of epidemiological surveillance programs in many developed countries (EFSA, 2013). These measures provide information about endemic *Salmonella*-serovar patterns, outbreaks, temporal trends and the monitoring of control actions (CDC, 2018). The principal reservoir of *S. enterica* is the intestinal tract of livestock, representing one of the main sources of infection for humans (Antunes *et al.*, 2016). Despite this epidemiological and economic importance, in South America there is a paucity of information concerning *Salmonella* in poultry systems (Alexandre *et al.*, 2000; Donado-Godoy *et al.*, 2012). Research conducted in 2016 pertaining to broiler chicken farming in Ecuador indicated the presence of *Salmonella* serotypes *S. Infantis*, *S. Enteritidis* and *S. Corvallis* (Vinueza-Burgos *et al.*, 2016). However, among layer hens, no information about the prevalence or diversity of *Salmonella* has been reported in this country nowadays. The purpose of this study was to estimate the presence and diversity of *Salmonella* bacteria present in 21 layer farms in central Ecuador.

## Methods

### Samples

Samples were collected between August–November 2017 from different layer farms in central Ecuador (Tungurahua and

Cotopaxi provinces), which accounts for around 60% of egg production in the country. A total of 21 farms (>1,000 birds) in Latacunga, Cevallos, Quero and Ambato (all Ecuador) were sampled, based on their willingness to provide verbal consent for this study (verbal consent was obtained over written consent owing to the farmers' reluctance to sign their names, as they perceived this could be used to identify them). Further details for each site can be found in Dataset 1 (Calero-Cáceres, 2019a)). One laying hen house per farm was selected. The following samples were collected in each house: 21 pooled cloacal swabs (10 cloacal swabs per pool); 21 manure drag swabs (environmental swabs, 1 per business); 21 caecum content samples (1 layer per farm). To evaluate the potential risk from contaminated feed, 18 composite samples were taken from farmyards (18 of the 21 farms consented verbally to having these samples taken). Additionally, 21 fecal samples from backyard layers were sampled in traditional local markets. All samples were transported in an icebox at 3–5°C within 2 hours of collection for bacterial isolation. The experiment was performed under supervision of the ethical committee of the Faculty of Agricultural Sciences, Universidad Técnica de Ambato.

### Detection of *Salmonella*

*Salmonella* was isolated following standardized methods (ISO 6579) (ISO, 2017). The samples were pre-enriched in buffered peptone water (Oxoid, Basingstoke, England) and then incubated at 37±1°C for 18 h ± 2 h. Next, Rappaport Vassiliadis Soy Broth (RVS Broth) (Merck Millipore, Darmstadt, Germany) was used as a selective medium, being inoculated with the pre-enriched culture and incubated at 41.5±1°C for 24±3 h. One loopful of the selective enrichment medium was streaked onto xylose lysine deoxycholate agar (XLD agar) (Becton Dickinson GmbH, Heidelberg, Germany) and incubated at 37±1°C for 24±3 h. Presumptive *Salmonella* isolates (identified as red/yellow colonies with a black center) were purified in MacConkey agar (Merck, Darmstadt, Germany) and incubated at 37±1°C for 24 h. Isolates were Gram stained and the following biochemical tests were performed for confirmation of the genus *Salmonella*: a catalase test using 30% hydrogen peroxide (Merck Millipore, Darmstadt, Germany); triple sugar iron agar test (TSI) (Becton Dickinson GmbH, Heidelberg, Germany), Simmons citrate agar test (Merck, Darmstadt, Germany), Christensen urea agar test (Britania Lab., Buenos Aires, Argentina), and indole reaction using tryptone water (Merck, Darmstadt, Germany) and Kovac's reagent (Sigma Aldrich, St. Louis, USA). One isolate per positive sample was selected and cryopreserved using overnight growth in LB broth (Sigma Aldrich, St. Louis, USA) supplemented with 30% glycerol (Merck Millipore, Darmstadt, Germany) and maintained at -80°C until analysis.

### Serovar determination by PCR

PCR assays were performed to identify the genes under specific conditions (Table 1). One virulence gene related with fimbrial cluster (*bcfC*) was evaluated as target of *S. enterica* species (Zhu *et al.*, 2015). For serovars and biotypes: A modification methylase gene that is specific of *S. Infantis* (M.SinI) (Ranjbar *et al.*, 2017). 23S rRNA gene associated to *S. Typhi*

**Table 1. Oligonucleotides used in this study.**

Organism	Name and direction	5' to 3' sequence	Gene	Annealing temperature (°C)	Amplicon size (bp)	References
Enteritidis	sdf-F	TGT GTT TTA TCT GAT GCA AGA G	sdf locus	56	293	(Agron <i>et al.</i> , 2001)
	sdf-R	CGT TCT TCT GGT ACT TCA GAT GAC				
Enterica	bcfC-F	GGG TGG GCG GAA AAC TAT TTC	bcfC	56	993	(Zhu <i>et al.</i> , 2015)
	bcfC-R	CGG CAC GGC GGA ATA GAG CAC				
Infantis	M. SinI F	CAC AAT GAA CGT GGT GAA GG	M.Sin I	56	184	(Ranjbar <i>et al.</i> , 2017)
	M. SinI R	TGA ACT ACG TTC GTT CTT CTGG				
Gallinarum biotype Gallinarum	steB-F	TGT CGA CTG GGA CCC GCC CGC CCG C	steB	56	636	(Pugliese <i>et al.</i> , 2011)
	steB-R	CCA TCT TGT AGC GCA CCA T				
Gallinarum biotype Pullorum	rhs-F	TCG TTT ACG GCA TTA CAC AAG TA	rhs locus	56	402	(Zhu <i>et al.</i> , 2015)
	rhs-R	CAA ACC CAG AGC CAA TCT TAT CT				
Typhi	sty-1	TGC CGG AAA CGA ATC T	23S rRNA gene	53	300	(Zhu <i>et al.</i> , 1996)
	sty-2	GGT TGT CAT GCC AAT GCA CT				
Typhimurium	Fli15	CGG TGT TGC CCA GGT TGG TAA T	fliC gene	53	620	(Pui <i>et al.</i> , 2011)
	Typ04	ACT GGT AAA GAT GGC T				

(*sty*) (Pui *et al.*, 2011). A flagellin gene that show specificity to *S. Typhimurium* (*fliC*) (Pui *et al.*, 2011). Gallinarum biotypes were identified by *steB* fimbrial and *rhs* locus (Zhu *et al.*, 2015). Specific DNA difference fragment (*SdfI*) served as target to distinguish *S. Enteritidis* from other serovars (Agron *et al.*, 2001). And a putative membrane protein (*gly*) to identify *S. Kentucky* (Zhu *et al.*, 2015). DNA was extracted from overnight cultures in Casein-Peptide Soy meal-Peptide Broth (Merck, Darmstadt, Germany) as described by Muniesa *et al.* (2004). Approximately  $\approx 50$  ng/reaction resulting from the thermal shock of bacterial suspensions ( $>10^7$  UFC/ml) diluted in sterile ddH<sub>2</sub>O was used as template for the PCR reactions. Amplifications were carried out as follows: initial denaturation at 95°C for 1 min; 35 cycles of 95°C for 30 s, annealing temperature according to Table 1 for 30 s, extension at 72°C for 1 min; and a final elongation step at 72°C for 7 min. The following reference *Salmonella* strains were used as positive controls for the six serovars under investigation: *S. Enteritidis* UNIETAR 1, *S. Gallinarum* b. Gallinarum NCTC 13346, *S. Gallinarum* b. Pullorum ATCC 19945®, *S. Typhi* ATCC® 19430, *S. Typhimurium* ATCC® 26930 and *S. Infantis* UNIETAR 3CT7.

The reaction mixture contained: 12.5 µl of DreamTaq Green PCR Master Mix (Thermo Fisher Scientific, Massachusetts, USA), 0.5 µl of each primer (30 µM stock), 9 µl of nuclease-free water (Thermo Fisher Scientific, Massachusetts, USA), and 2.5 µl of crude DNA were used. PCRs were performed with an Applied Biosystems SimplyAmp Thermal Cycler (Thermo Fisher Scientific, Massachusetts, USA). A total of 10 µl of each PCR product were analyzed by agarose gel electrophoresis and stained using Sybr® Safe DNA Gel Stain (Invitrogen, Carlsbad, USA).

## Results

### Assessing the presence of *Salmonella*


Overall, 31 out of 34 isolates which showed phenotypic characteristics in accordance with *Salmonella* were confirmed by PCR as *S. enterica* (Table 2). Of the 21 farms, 16 (76%, 95%CI: 53-92) showed the presence of *Salmonella* on environmental surfaces, and *Salmonella* bacteria were isolated in 7 pooled cloacal swabs (33%, 95%CI: 15-57) and in 6 of 18 composite feed samples (33%, 95%CI: 13-59). Feces from backyard layers showed the presence of *Salmonella* in 2 of 21 samples (10%, 95%CI: 1-30). Dataset 1 shows the location of each sample and the presence or absence of *Salmonella* (Calero-Cáceres, 2019a)

### Determining presence of serovars

Regarding to the presence of individual serovars (Figure 1), the highest occurrence was of *S. Infantis*, detected in 58% (18/31) of isolates, followed by *S. Typhimurium*, present in 32% of samples (10/31) (Supplementary Figure 1 (Calero-Cáceres, 2019c)). The serovar of 10% of *Salmonella* isolates (3/31) were unable to be serotyped by the panel of tests. In poultry feed samples, *S. Infantis* was present in 100% (6/6), while in *S. enterica* isolated from manure drag swabs, *S. Infantis* was detected in 50% (8/8) and *S. Typhimurium* in 50% (8/8). In backyard poultry feces from local markets, *S. Infantis* was detected in 100% of positive samples (2/2). The most heterogeneous diversity of *Salmonella* serotypes was observed in pooled cloacal swabs, with 2 *S. Infantis*, 2 *S. Typhimurium* and 3 *S. enterica* consisting of serovars not covered within the panel. Dataset 1 lists the identity of the serovar taken from each sampling location (Calero-Cáceres, 2019a). Dataset 2 shows PCR gels for confirmation of *Salmonella* serovars (Calero-Cáceres, 2019b).

**Table 2. Salmonella-positive samples in relation to evaluated matrices.**

Location	Sample	Matrix	Samples, n	Positive samples, n (%)
Farms	Feed	Composite feed	18	6 (33%)
	Environment	Environmental swabs	21	16 (76%)
	Animal	Laying hen cloacal swab (composite)	21	7 (33%)
Local markets	Environment	Backyard poultry feces	21	2 (10%)

**Figure 1. Distribution of Salmonella serovars according to the evaluated matrices.**

## Discussion

The overall results showed a significant presence of *Salmonella* in layer farms in the central Ecuador region, one of the main sources of egg production in the country, with a detection of at least one sample positive per farm in the 76% of the evaluated sites. This result is similar to that reported in Colombia (65%) (Donado-Godoy *et al.*, 2012), and considerably higher than the presence of *Salmonella* in broiler chicken farms in northern Ecuador (15.9%) (Vinueza-Burgos *et al.*, 2016).

The notable presence of *Salmonella* in poultry feed suggests that this may constitute a potential reservoir of this bacteria for poultry systems in the evaluated region, and consequently, a potential route of infection and colonization of poultry and subsequent entry into the food chain (Fink-Gremmels, 2012; Jones, 2011). In recent research conducted in an integrated poultry farm in Ecuador, 4.1% (8/194) of samples were positive for *Salmonella*, particularly in animal-based feed (Villagómez Estrada *et al.*, 2017). Therefore, the accurate detection of this pathogen in feed is necessary in order to identify the critical

points of contamination (facilities, raw materials, transport, storage, producers), that one may apply effective measures to reduce the risk of transmission.

The finding of only *S. Infantis* in poultry feed may be attributed to the high level of persistence of this serovar over time in poultry feed, which in turn results in its considerable presence in Ecuadorian broiler chicken (Andino *et al.*, 2014; Vinueza-Burgos *et al.*, 2016). Although the origin of this serovar in feed it is not well defined, studies have pointed to cross-contamination with feces, persistent contamination of storage bins and surfaces, and poor ingredient selection as main causes of feed contamination with *Salmonella* (Davies & Wray, 1997; Huss *et al.*, 2018; Laban *et al.*, 2014). Genomic tools, such as multilocus sequence typing, BOX-PCR, (GTG)<sub>5</sub>-PCR or whole-genome sequencing may help to identify the origin of feed contamination in a larger study.

Drag swabs revealed the presence of two different serovars in the sampled poultry farms: *Infantis* and *Typhimurium*. These

non-typhoidal *S. enterica* serovars are commonly associated with poultry systems and are linked to outbreaks of foodborne illness (Anderson *et al.*, 2016; Pui *et al.*, 2011). Serovars vary in their persistence over time and geographic distribution around the world (Hendriksen *et al.*, 2011), making further studies desirable in order to evaluate the variations in serovar persistence in the locations sampled in this research.

In backyard layer feces sampled at local markets, *S. Infantis* alone was detected (2/2), but further evaluation of *Salmonella* serovars in backyard flocks is recommended in order to improve surveillance of this potential source of salmonellosis (Behraves *et al.*, 2014). In pooled cloacal swabs, the serovars detected were *Infantis* (2/7), *Typhimurium* (2/7) and unidentified serovars (3/7). *Infantis* and *Typhimurium* serovars are commonly detected in poultry around the world (Foley & Lynne, 2008; Foley *et al.*, 2011). Complementary analysis by serotyping the unclassified serovars is necessary in order to identify *Salmonella* bacteria not covered by the panel.

Data remains insufficient concerning the prevalence of *Salmonella* in poultry systems in Ecuador and in Latin America in general. The coordination of similar future studies may provide a starting point for surveillance of zoonotic bacteria within a defined public health area, leading to an improvement in policies and safe practices in the food industry. Such studies may reduce the risk of infection and establish protocols for corrective measures to be implemented in key upstream points of the chain as indicated by the data.

At the same time, the intervention of public health authorities may be required in order to ensure the participation of a fully representative range of poultry businesses in future research. This study depended on the voluntary consent of farm owners to providing samples, which may have led to selection bias. In order to establish a thorough program of surveillance, all potential sources of *Salmonella* infection in the region need to be made accessible to researchers.

These findings show a significant presence of *Salmonella* in layer farms in the central zone of Ecuador. The predominant serovars are *S. Infantis* and *S. Typhimurium*, typified by PCR. The principal source of infection could be related with poultry feed. From a public health perspective, it is necessary to establish adequate surveillance of *Salmonella*, including protocols covering biosecurity practices, antibiotic usage and random sampling programs.

## Data availability

### Underlying data

Figshare: Dataset 1. Origin, biochemical tests, PCR results and serovar of *Salmonella* isolates. <https://doi.org/10.6084/m9.figshare.7726163.v2> (Calero-Cáceres, 2019a)

Figshare: Dataset 2. PCR results of *bcfC*, *fliC* and *M.SinI* genes. <https://doi.org/10.6084/m9.figshare.7726217.v2> (Calero-Cáceres, 2019b)

### Extended data

Figshare: Supplementary figure 1. Multiplex PCR amplification of serovar-specific genes. a) *bcfC* and *mSinI* fragments (*S. Infantis*), b) *bcfC* and *fliC* fragments (*S. Typhimurium*).

<https://doi.org/10.6084/m9.figshare.7732934.v1> (Calero-Cáceres, 2019c.)

Data are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

## Grant information

This study was supported by the Dirección de Investigación y Desarrollo DIDE-Universidad Técnica de Ambato (Project 1568-CU-P-2017).

*The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

## References

- Agron PG, Walker RL, Kinde H, *et al.*: Identification by subtractive hybridization of sequences specific for *Salmonella enterica* serovar enteritidis. *Appl Environ Microbiol.* 2001; **67**(11): 4984–4991.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Alexandre SM, Pozo MC, González GV, *et al.*: Detección de *Salmonella enteritidis* en muestras de productos avícolas de consumo humano en la Región Metropolitana. *Revista Médica de Chile.* 2000; **128**(10): 1075–1083.  
[Publisher Full Text](#)
- Anderson TC, Nguyen TA, Adams JK, *et al.*: Multistate outbreak of human *Salmonella* Typhimurium infections linked to live poultry from agricultural feed stores and mail-order hatcheries, United States 2013. *One Health.* 2016; **2**: 144–149.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Andino A, Pendleton S, Zhang N, *et al.*: Survival of *Salmonella enterica* in poultry feed is strain dependent. *Poult Sci.* 2014; **93**(2): 441–447.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Antunes P, Mourão J, Campos J, *et al.*: Salmonellosis: the role of poultry meat. *Clin Microbiol Infect.* 2016; **22**(2): 110–121.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bäumler AJ, Hargis BM, Tsois RM: Tracing the origins of *Salmonella* outbreaks. *Science.* 2000; **287**(5450): 50–52.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Behraves CB, Brinson D, Hopkins BA, *et al.*: Backyard poultry flocks and salmonellosis: a recurring, yet preventable public health challenge. *Clin Infect Dis.* 2014; **58**(10): 1432–1438.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Callejón RM, Rodríguez-Naranjo MI, Ubeda C, *et al.*: Reported foodborne outbreaks due to fresh produce in the United States and European Union: trends and causes. *Foodborne Pathog Dis.* 2015; **12**(1): 32–38.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Calero-Cáceres W: Dataset 1. Origin, biochemical tests, PCR results and serovar of *Salmonella* isolates. *figshare.* Dataset. 2019a.  
<http://www.doi.org/10.6084/m9.figshare.7726163.v2>
- Calero-Cáceres W: Dataset 2. PCR results of *bcfC*, *fliC* and *M.SinI* genes. *figshare.* Figure. 2019b.  
<http://www.doi.org/10.6084/m9.figshare.7726217.v2>
- Calero-Cáceres W: Supplementary figure 1. *figshare.* Figure. 2019c.  
<http://www.doi.org/10.6084/m9.figshare.7732934.v1>

CDC: **Salmonella Atlas**. Reports and Publications. *Salmonella*. CDC [Online]. [Accessed: 7 February 2019]. 2018.

[Reference Source](#)

Davies RH, Wray C: **Distribution of *Salmonella* contamination in ten animal feedmills**. *Vet Microbiol*. 1997; **57**(2–3): 159–169.

[PubMed Abstract](#) | [Publisher Full Text](#)

Donado-Godoy P, Gardner I, Byrne BA, *et al.*: **Prevalence, risk factors, and antimicrobial resistance profiles of *Salmonella* from commercial broiler farms in two important poultry-producing regions of Colombia**. *J Food Prot*. 2012; **75**(5): 874–883.

[PubMed Abstract](#) | [Publisher Full Text](#)

EFSA: **EU Summary Report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2013**. [Online]. [Accessed: 7 February 2019]. 2013.

[Reference Source](#)

Fink-Gremmels J: **Animal feed contamination: effects on livestock and food safety**. 2012.

[Reference Source](#)

Foley SL, Lynne AM: **Food animal-associated *Salmonella* challenges: pathogenicity and antimicrobial resistance**. *J Anim Sci*. 2008; **86**(14 Suppl): E173–87.

[PubMed Abstract](#) | [Publisher Full Text](#)

Foley SL, Nayak R, Hanning IB, *et al.*: **Population dynamics of *Salmonella enterica* serotypes in commercial egg and poultry production**. *Appl Environ Microbiol*. 2011; **77**(13): 4273–4279.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Hendriksen RS, Vieira AR, Karlsmose S, *et al.*: **Global monitoring of *Salmonella* serovar distribution from the World Health Organization Global Foodborne Infections Network Country Data Bank: results of quality assured laboratories from 2001 to 2007**. *Foodborne Pathog Dis*. 2011; **8**(8): 887–900.

[PubMed Abstract](#) | [Publisher Full Text](#)

Huss A, Cochrane R, Muckey M, *et al.*: **Animal Feed Mill Biosecurity: Prevention of Biological Hazards**. In *Food and feed safety systems and analysis*. Elsevier, 2018; 63–81.

[Publisher Full Text](#)

ISO: **ISO 6579-1:2017 - Microbiology of the food chain -- Horizontal method for the detection, enumeration and serotyping of *Salmonella* -- Part 1: Detection of *Salmonella* spp.** International Organization for Standardization. 2017.

[Reference Source](#)

Issenhuth-Jeanjean S, Roggentin P, Mikoleit M, *et al.*: **Supplement 2008-2010 (no. 48) to the White-Kauffmann-Le Minor scheme**. *Res Microbiol*. 2014; **165**(7): 526–530.

[PubMed Abstract](#) | [Publisher Full Text](#)

Jones FT: **A review of practical *Salmonella* control measures in animal feed**. *J Appl Poult Res*. 2011; **20**(1): 102–113.

[Publisher Full Text](#)

Laban SE, Moustafa GZ, Anwer W, *et al.*: **Microbial Load of Poultry By-Products Following Rendering Process**. *Global Veterinaria*. 2014; **12**: 756–759.

[Reference Source](#)

Majowicz SE, Musto J, Scallan E, *et al.*: **The global burden of nontyphoidal *Salmonella* gastroenteritis**. *Clin Infect Dis*. 2010; **50**(6): 882–889.

[PubMed Abstract](#) | [Publisher Full Text](#)

MSP: **Gaceta Epidemiológica Ecuador SIVE-ALERTA – Ministerio de Salud Pública**. [Online]. [Accessed: 6 February 2019]. 2018.

[Reference Source](#)

Muniesa M, Blanco JE, De Simón M, *et al.*: **Diversity of *stx*<sub>2</sub> converting bacteriophages induced from Shiga-toxin-producing *Escherichia coli* strains isolated from cattle**. *Microbiology*. 2004; **150**(Pt 9): 2959–2971.

[PubMed Abstract](#) | [Publisher Full Text](#)

Pugliese N, Circella E, Pazzani C, *et al.*: **Validation of a seminested PCR approach for rapid detection of *Salmonella enterica* subsp. *enterica* serovar Gallinarum**. *J Microbiol Methods*. 2011; **85**(1): 22–27.

[PubMed Abstract](#) | [Publisher Full Text](#)

Pui CF, Wong WC, Chai LC, *et al.*: **Simultaneous detection of *Salmonella* spp., *Salmonella* Typhi and *Salmonella* Typhimurium in sliced fruits using multiplex PCR**. *Food Control*. 2011; **22**(2): 337–342.

[Publisher Full Text](#)

Ranjbar R, Mortazavi SM, Mehrabi Tavana A, *et al.*: **Simultaneous Molecular Detection of *Salmonella enterica* Serovars Typhi, Enteritidis, Infantis, and Typhimurium**. *Iran J Public Health*. 2017; **46**(1): 103–111.

[PubMed Abstract](#) | [Free Full Text](#)

Thiennimitr P, Winter SE, Bäumlér AJ: ***Salmonella*, the host and its microbiota**. *Curr Opin Microbiol*. 2012; **15**(1): 108–114.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Varma JK, Greene KD, Ovitt J, *et al.*: **Hospitalization and antimicrobial resistance in *Salmonella* outbreaks, 1984-2002**. *Emerg Infect Dis*. 2005; **11**(6): 943–946.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Villagómez Estrada S, Logacho Pilataxi M, Vinueza Burgos C: **Presencia y Resistencia a los Antimicrobianos de serovariedades de *Salmonella enterica* aisladas en una empresa avícola integrada del Ecuador**. *Rev Ecuat Med Cienc Biol*. 2017; **38**(1).

[Publisher Full Text](#)

Vinueza-Burgos C, Cevallos M, Ron-Garrido L, *et al.*: **Prevalence and Diversity of *Salmonella* Serotypes in Ecuadorian Broilers at Slaughter Age**. *PLoS One*. 2016; **11**(7): e0159567.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

WHO: **Salmonella**. WHO. [Online]. [Accessed: 7 February 2019]. 2017.

[Reference Source](#)

Zhu Q, Lim CK, Chan YN: **Detection of *Salmonella typhi* by polymerase chain reaction**. *J Appl Bacteriol*. 1996; **80**(3): 244–251.

[PubMed Abstract](#) | [Publisher Full Text](#)

Zhu C, Yue M, Rankin S, *et al.*: **One-Step Identification of Five Prominent Chicken *Salmonella* Serovars and Biotypes**. *J Clin Microbiol*. 2015; **53**(12): 3881–3883.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

# Open Peer Review

Current Referee Status:  

---

## Version 2

Referee Report 17 April 2019

<https://doi.org/10.5256/f1000research.20651.r46971>



**Tania Gaviria Cantín**

Departamento de Ciencias Naturales, Exactas y Estadística. Facultad de Ciencias Básicas, Universidad de Santiago de Cali, Cali, Colombia

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Molecular microbiology, bacterial genetics, molecular biology, genetic engineering

**I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

Referee Report 09 April 2019

<https://doi.org/10.5256/f1000research.20651.r46970>



**Eloy Gonzalez-Gustavson** 

Facultad de Medicina Veterinaria, Universidad Nacional Mayor de San Marcos, Lima, Peru

I agree with the changes.

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Epidemiology, biostatistics, environmental microbiology, parasitology

**I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

---

## Version 1

Referee Report 21 March 2019

<https://doi.org/10.5256/f1000research.19945.r45085>





**Eloy Gonzalez-Gustavson** 

Facultad de Medicina Veterinaria, Universidad Nacional Mayor de San Marcos, Lima, Peru

The scientific names in the manuscript need to be reviewed.

The author use incorrectly the terms prevalence and incidence as synonyms.

Incidence is defined as the presence of new cases of an illness over a period of time. The manuscript does not correspond to an incidence study.

I would like to see more information in the manuscript about the design to consider the results as prevalence. I have not found details about the representativeness of the sampling (sample size to get a prevalence) or if they were obtained randomly. If the method does not satisfy the requirement to be considered as prevalence, I recommend replacing the term prevalence by frequency, presence or proportion.

The confidence interval in one case was incorrectly estimated.

"Feces from backyard layers showed the presence of *Salmonella* in 2 of 21 samples (10%, 95%CI: 0–22)"  
This values have to be calculated based on binomial distribution. Actually in the manuscript, the confidence interval was estimated based on normal or t distribution and 0 was included as lower interval.

**Is the work clearly and accurately presented and does it cite the current literature?**

Partly

**Is the work clearly and accurately presented and does it cite the current literature?**

Partly

**Is the study design appropriate and is the work technically sound?**

Yes

**Is the study design appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and analysis provided to allow replication by others?**

Yes

**Are sufficient details of methods and analysis provided to allow replication by others?**

Yes

**If applicable, is the statistical analysis and its interpretation appropriate?**

Partly

**If applicable, is the statistical analysis and its interpretation appropriate?**

Partly

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Epidemiology, biostatistics, environmental microbiology, parasitology

**I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.**

Referee Report 11 March 2019

<https://doi.org/10.5256/f1000research.19945.r45084>



**Tania Gaviria Cantín**

Departamento de Ciencias Naturales, Exactas y Estadística. Facultad de Ciencias Básicas, Universidad de Santiago de Cali, Cali, Colombia

#### **Title**

I consider that the title is not totally in agreement with the study. This could be modified at "*Samonella* diversity isolated from ... " without the word "Prevalence" because the study is not a follow-up of previously reported cases in those same conditions

#### **Introduction**

In the fourth line you would change the word "poisoning" by contamination

At the end correct one "and" that is in italics

When you say "among the layer hens, no information about prevalence or diversity of *Salmonella* has been reported", you could specify or support if that lack of information is only in Ecuador or at a South American or global level. In the same way, it could important to include some epidemiological data of salmonellosis in Ecuador.

#### **Methodology**

I think that is not necessary to do DNA extraction to perform PCR, because *Salmonella* an is gram negative bacteria and during the start of the PCR where the temperature rises to 90 ° C it is possible to destroy cellular wall and release the DNA. However, it does not affect the results, taking into account that the procedure has been carried out correctly avoiding cross contamination.

On the other hand, in order to reproduce the methodology of this study, the program used for PCR should be indicated. The concentration of primers (30 uM) is the final or initial concentration?

It would be useful to indicate why those particular genes were used as molecular markers for the identification of the species and the serovars

#### **Results**

You can ignore the phrase "amplification of the bcfC gene" and leave the table 1 referenced because it is

a bit redundant

Regarding statistical analysis and sample number in representation or not, I prefer not to give many opinions because I'm not qualified enough for that. However, it would be important that the methodology explain in detail how the statistical analysis was performed and the calculation of the incidence.

In order of reference, table 1 of results should be table 2 and vice versa

Regarding table 1 (positive *Salmonella* samples ...), it could be a little more consistent with what they explain in the methodology because it took me a bit to understand it, taking into account the classification of columns in terms of location, samples, etc.

### **Discussion**

I believe that at some point it can be said that the regulations of Ecuador in relation to the health and control of layer farms and local markets and explain how the work is contributing to improve local control, taking into account that this layer farms produce around 60% of egg production in the country.

**Is the work clearly and accurately presented and does it cite the current literature?**

Yes

**Is the work clearly and accurately presented and does it cite the current literature?**

Yes

**Is the study design appropriate and is the work technically sound?**

Partly

**Is the study design appropriate and is the work technically sound?**

Partly

**Are sufficient details of methods and analysis provided to allow replication by others?**

Partly

**Are sufficient details of methods and analysis provided to allow replication by others?**

Partly

**If applicable, is the statistical analysis and its interpretation appropriate?**

I cannot comment. A qualified statistician is required.

**If applicable, is the statistical analysis and its interpretation appropriate?**

I cannot comment. A qualified statistician is required.

**Are all the source data underlying the results available to ensure full reproducibility?**

Partly

**Are all the source data underlying the results available to ensure full reproducibility?**

Partly

**Are the conclusions drawn adequately supported by the results?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Molecular microbiology, bacterial genetics, molecular biology, genetic engineering

**I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.**

---

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact [research@f1000.com](mailto:research@f1000.com)

**F1000Research**