

## A *mecC* allotype, *mecC3*, in the CoNS *Staphylococcus caeli*, encoded within a variant SCC*mecC*

A. C. MacFadyen<sup>1</sup>, E. M. Harrison<sup>2</sup>, I. Drigo<sup>3</sup>, J. Parkhill<sup>2</sup>, M. A. Holmes<sup>4</sup> and G. K. Paterson<sup>1\*</sup>

<sup>1</sup>Royal (Dick) School of Veterinary Studies and The Roslin Institute, University of Edinburgh, Easter Bush Campus, Midlothian EH25 9RG, UK; <sup>2</sup>The Wellcome Trust Sanger Institute, Wellcome Trust, Genome Campus, Hinxton CB10 1SA, UK; <sup>3</sup>Istituto Zooprofilattico Sperimentale delle Venezie, via dell'Università 10, 35020 Legnaro, PD, Italy; <sup>4</sup>Department of Veterinary Medicine, University of Cambridge, Madingley Road, Cambridge CB3 0ES, UK

\*Corresponding author. E-mail: gavin.paterson@ed.ac.uk

Received 20 July 2018; returned 10 October 2018; revised 7 November 2018; accepted 7 November 2018

**Background:** Methicillin resistance in staphylococci is conferred by an alternative PBP (PBP2a/2') with low affinity for most  $\beta$ -lactam antibiotics. PBP2a is encoded by *mecA*, which is carried on a mobile genetic element known as SCC*mec*. A variant of *mecA*, *mecC*, was described in 2011 and has been found in *Staphylococcus aureus* from humans and a wide range of animal species as well as a small number of other staphylococcal species from animals.

**Objectives:** We characterized a novel *mecC* allotype, *mecC3*, encoded by an environmental isolate of *Staphylococcus caeli* cultured from air sampling of a commercial rabbit holding.

**Methods:** The *S. caeli* isolate 82B<sup>T</sup> was collected in Italy in 2013 and genome sequenced using MiSeq technology. This allowed the assembly and comparative genomic study of the novel SCC*mec* region encoding *mecC3*.

**Results:** The study isolate encodes a novel *mecA* allotype, *mecC3*, with 92% nucleotide identity to *mecC*. *mecC3* is encoded within a novel SCC*mec* element distinct from those previously associated with *mecC*, including a *ccrAB* pairing (*ccrA5B3*) not previously linked to *mecC*.

**Conclusions:** This is the first description of the novel *mecC* allotype *mecC3*, the first isolation of a *mecC*-positive *Staphylococcus* in Italy and the first report of *mecC* in *S. caeli*. Furthermore, the SCC*mec* element described here is highly dissimilar to the archetypal SCC*mec* XI encoding *mecC* in *S. aureus* and to elements encoding *mecC* in other staphylococci. Our report highlights the diversity of *mecC* allotypes and the diverse staphylococcal species, ecological settings and genomic context in which *mecC* may be found.

### Introduction

Methicillin resistance in *Staphylococcus aureus* is typically conferred by the gene *mecA* along with two variants, *mecB* and *mecC*.<sup>1–4</sup> *mec* gene resistance is mediated by an alternative PBP with reduced affinity for almost all  $\beta$ -lactam antimicrobials.<sup>5,6</sup> Since its first discovery in bulk tank milk on an English dairy farm,<sup>1</sup> *mecC* has been found in *S. aureus* isolates from a wide range of host species, including human carriage and infection and various wildlife, companion and livestock species<sup>7,8</sup> with genomic analysis indicating zoonotic transmission from livestock to humans.<sup>9,10</sup> *mecC*-MRSA have been reported from a range of countries and while this geographical distribution has centred on Europe,<sup>8</sup> *mecC*-MRSA have also been reported in Australia.<sup>11</sup> In addition to *S. aureus*, *mecC* has been described in other species of staphylococci, albeit in only a limited number of species and a small number of isolates, which have all come from animals: *Staphylococcus*

*xylosus*,<sup>12</sup> *Staphylococcus sciuri*,<sup>13</sup> *Staphylococcus stepanovici*<sup>14</sup> and *Staphylococcus saprophyticus*<sup>15,16</sup> or in the case of *Staphylococcus edaphicus* from environmental sampling.<sup>17</sup> *mecC* in MRSA is found within a staphylococcal cassette chromosome *mec* (SCC*mec*) type XI while in other staphylococci it is found in a range of genomic contexts, although always within the *orfX* region and with features common to SCC*mec* elements. Divergent *mecC* allotypes *mecC1* and *mecC2* have been described in *S. xylosus*<sup>12</sup> and *S. saprophyticus*,<sup>15</sup> respectively, but have not been reported in *S. aureus*, suggesting a greater diversity and ancestral association of *mecC* with non-*aureus* staphylococci.

Here we describe a novel *mecC* allotype, *mecC3*, encoded within a distinct and novel SCC*mec* element in a newly described species, *Staphylococcus caeli*,<sup>18</sup> which furthers our understanding of the diversity of *mecC* genes and the diverse species and genetic elements that carry it.

## Materials and methods

### Isolate collection and WGS

The *S. caeli* isolate 82B<sup>T</sup> (=NCTC 14063<sup>T</sup>=CCUG 71912<sup>T</sup>) was collected from air sampling in a commercial rabbit holding in Italy in 2013 as part of a previous study<sup>19</sup> and has been described elsewhere, including its genome sequencing.<sup>18</sup> MRSA was also present on the sampled farm.<sup>19</sup> Six months prior to the isolation of 82B<sup>T</sup>, ST398 MRSA belonging to t034 and t5210 were isolated from farm workers, their relatives and rabbits on the farms.<sup>19</sup> At the time of 82B<sup>T</sup> isolation MRSA belonging to t034, t5210, t1190 and t2970 were isolated from rabbits and humans with non-typed MRSA isolates found in air samples and surface swabs.<sup>19</sup>

### Sequence assembly and identification of SCCmec

Sequencing reads were *de novo* assembled using Velvet<sup>20</sup> and annotated using the NCBI Prokaryotic Genome Annotation Pipeline.<sup>21</sup> Contiguous sequences (contigs) containing the *orfX* and *mecC* genes were identified using BLAST, using the respective genes from *S. aureus* LGA251 (accession number FR821779) as query sequences. Contigs NZ\_FMPG01000005.1 and NZ\_FMPG01000008.1 were identified as containing the *orfX* and *mecC* genes, respectively. Primers were designed for the end of contig NZ\_FMPG01000005.1 and start of contig NZ\_FMPG01000008.1, followed by PCR, with the resulting amplicon being ABI sequenced (Source BioScience, Cambridge, UK), to close the gap between the contigs. Primers were designed using Primer3 (<http://primer3.ut.ee/>).

### Phylogenetic analysis of *mecA* homologues

Phylogenetic analyses were carried out in MEGA7.<sup>22</sup> All nucleotide sequences were obtained from NCBI databases, using the following accession numbers: *S. aureus* LGA251, FR821779, *mecC*; *S. xylosum* S04009, HE993884, *mecC1*; *S. saprophyticus* 210, KF955540, *mecC2*; *S. caeli* 82B, *mecC3*, FMPG01000008; *S. sciuri* K11, Y13094, *mecA1*; *S. aureus* N315, NC\_002745, *mecA*; *Staphylococcus vitulinus* CSB08, AM048810, *mecA2*; *Macrococcus caseolyticus* IMD0819, KY013611, *mecD*; and *M. caseolyticus* JCS5402, NC\_011996, *mecB*. The sequences were aligned using MUSCLE and a maximum likelihood tree was generated, using a general time reversible model. Site substitution rates were calculated using a discrete gamma distribution model.

### Nucleotide accession numbers

The whole genome nucleotide sequences for *S. caeli* 82B<sup>T</sup> have been deposited previously<sup>18</sup> in the NCBI database under accession numbers NZ\_FMPG00000000 (assembly) and ERR473447 (sequence reads). The assembled SCCmec region of 82B<sup>T</sup> generated in this work has been deposited under accession number MH155596.

### Antimicrobial susceptibility testing

Antimicrobial susceptibility testing was performed by Vitek2 using AST P260 cards following the manufacturer's instructions. Disc diffusion was performed following the EUCAST disc diffusion method Version 6.0 January 2017. Growth on MRSA Brilliance (Oxoid, Basingstoke, UK) was tested by spreading a 1 µL loop taken from a 0.5 McFarland suspension (~1.5 × 10<sup>5</sup> cfu) on to the Brilliance plate and incubating at 30°C, 35°C and 37°C for 24 h. Interpretation was done based on the criteria for CoNS with *S. aureus* NCTC 12497 and NCTC 12493 used as control strains for susceptibility testing. PBP2a detection was performed using the PBP2<sup>a</sup> Latex Agglutination Test Kit (Oxoid) following the manufacturer's instructions.

## Results and discussion


### *S. caeli* 82B<sup>T</sup> contains a novel *mecC* allotype, *mecC3*

Isolate 82B<sup>T</sup> has been described as the type strain of a novel staphylococcal species, *S. caeli*, most closely related to *S. xylosum*, *S. saprophyticus* and *S. edaphicus*.<sup>18</sup> As part of the characterization of 82B<sup>T</sup> it underwent WGS. Further investigation of the 82B<sup>T</sup> genome by BLAST analysis using the *mecC* gene from *S. aureus* LGA251 as the query sequence revealed that 82B<sup>T</sup> encodes a *mecC* homologue on contig NZ\_FMPG01000008.1. Alignments of the 82B<sup>T</sup> *mecC* gene with known *mecC* allotypes showed that it shares 92% nucleotide identity with *mecC* from LGA251, 93% with *mecC1* from *S. xylosum* S04009 and 94% with *mecC2* from *S. saprophyticus* 210. Based on these results and in line with guidelines for the classification for *mecA* homologues,<sup>23</sup> the *mecC* gene for *S. caeli* 82B<sup>T</sup> is a new allotype of *mecC* and therefore designated *mecC3*. The phylogenetic relationship of *mecC3* to the other *mecA* homologues is illustrated in Figure 1; this suggests that the more common *mecC* has evolved from the more ancestral forms (*mecC1*, *mecC2*, *mecC3*) and that, of the three, *mecC3* is closest to the ancestral form. The *mecC* complex of *S. caeli* 82B<sup>T</sup> shares 93% and 92% nucleotide identity with the complete *mecC* complex of LGA251 and S04009, respectively, with the partial *mecC* complex of *S. saprophyticus* 210 sharing 94%. 82B<sup>T</sup> gave a negative reaction using the commercial PBP2a latex agglutination test kit from Oxoid. This reaction was not altered after cefoxitin induction. *mecC3* in 82B<sup>T</sup> was, however, detected by the published *mecC* primer pairs 1A and 1B<sup>24</sup> and *mecC*-Uni-F and *mecC*-Uni-R.<sup>13</sup>

### *S. caeli* 82B<sup>T</sup> *mecC3* is encoded within a distinct SCCmec-like element

*mecC* genes are typically found within SCCmec elements,<sup>13,14</sup> which insert into the genome at the 3'-end of a 23S rRNA methyltransferase gene, often referred to as *orfX*.<sup>25,26</sup> BLASTn analysis using *orfX* from LGA251 as the query sequence identified the *orfX* region of 82B<sup>T</sup> within contig NZ\_FMPG01000005.1. PCR and amplicon sequencing confirmed that the two contigs (NZ\_FMPG01000005.1 containing *orfX* and NZ\_FMPG01000008.1 containing *mecC3*) are contiguous in the 82B<sup>T</sup> genome with a gap of 140 bp between the ends of the two contigs. The assembled sequence generated in this study from these two contigs, encoding the *orfX* region of 82B<sup>T</sup> and *mecC3* including a SCCmec-like element, has been deposited in NCBI with accession number MH155596.

The integration site on the genome for SCCmec elements within *orfX*, known as the attachment (*attB*) site, is identified by the recombinase CcrAB/CcrC via a 14 bp sequence.<sup>27</sup> Insertion into the *attB* generally results in the SCCmec being flanked by direct repeats, *attR* and *attL*.<sup>13,27,28</sup> We therefore searched for these repeat sequences within the *orfX* region of 82B<sup>T</sup> to define the limits of its putative SCCmec-like element. The search sequences; gc[ag]-tatca[tc]aatgatgctggttt and aacc[tg]catca[tc][tc][at][ac][tc]gataag[ct], produced from previously identified *attR* and *attL* sequences, respectively, identified an *attR* site 51 kbp downstream of *orfX* and an *attL* site 3.8 kbp downstream of *mecC3* (Figure 2a). This indicates an SCCmec in 82B<sup>T</sup>, which is 127 kbp in size, with the *orfX* and *mecC3* regions at opposite ends of the element (Figure 2a). However, it is not clear if this represents a single entity or a composite element generated by consecutive insertions.



**Figure 1.** Phylogenetic relationship of representative *mec* allotypes. Nucleotide sequences were aligned using MUSCLE, with the maximum likelihood method, based on the general time reversible model, being used to build the tree. The highest log likelihood (−9785.4010) tree is shown. The percentage of trees in which the associated genes clustered together, based on bootstrapping with 500 replicates, is shown at the branches. A discrete gamma distribution was used to model site substitution rates, with branch lengths measured in the number of substitutions per site.

Within this SCCmec region, 82B<sup>T</sup> contains two *ccr* complexes and three joining (J) regions (Figure 2a). The *ccrA* and *ccrB* genes of 82B<sup>T</sup> share 90% and 91% nucleotide identity with *ccrA5* and *ccrB3*, respectively, from *Staphylococcus pseudintermedius* KM241 (accession number AM904731). This represents a type 6 *ccr* allotype,<sup>25,28</sup> which is novel for an SCCmec encoding a *mecC*. Indeed this pairing is rare among SCCmec in staphylococci, with a search of NCBI finding only two other staphylococcal isolates that contained this *ccrAB* pairing [*Staphylococcus cohnii* WC28 (accession number GU370073)<sup>29</sup> and *S. cohnii* SNUDS-2 (CP019597; locus tags BZ166\_04625 and BZ166\_04620)]. In addition to the *ccrAB* genes described above, the SCCmec of 82B<sup>T</sup> also contains a *ccrC* gene, sharing 90% nucleotide identity with *S. aureus* PM1 (SCCmec VII, accession number AB462393, gene *ccrC8*). The *attR* site was identified 51 kbp away from the *orfX* gene (Figure 2a), which deviates from typical SCCmec elements and therefore the *orfX* region was examined to find any additional *attR*-like site.

Within the 3' region of *orfX* an *att* site with sequence homology to the *attR* SCCmec type VII of PM1 was identified. As there are two *ccr* regions within the 82B<sup>T</sup> SCCmec, it was proposed that the *attR* within the 3'-end of *orfX* (Figure 2, *attR2*) might be linked to the CcrC recombinase. Therefore, a search sequence: [at][at][at][ct][ct][ga][cg][atc][ta][ca][at][ta][ct][ac]act[ga][ga][tc]a, based on the *attL* sequence of SCCmec elements containing only the *ccrC* gene, was used to identify the corresponding *attL* site linked to *attR2*, *attL2*. This revealed only one potential *attL2* site, located 16.8 kbp upstream of the *mec* region (Figure 2, *attL2*). Sequence alignment of the *attR/L* sites of 82B<sup>T</sup> with *attR/L* sites of *ccrAB*-containing SCCmec, compared with the alignment of the *attR/L2* sites of 82B<sup>T</sup> versus those *attR/L* sites of *ccrC*-containing SCCmec, suggest a varied *att* site for CcrAB/CcrC (Figure 3). Though it appears that the


highly conserved central 8 bp sequence is consistent, there is notable variation between the *ccrAB*- and *ccrC*-associated *att* sites within those bases that flank the 8 bp region (Figure 3). Although the location of *attL2* was identified with the predicted region, the only other potential *attL* site identified was *attL'*, recently described for a highly conserved type XI SCCmec found in *S. xylosus* 47–83.<sup>30</sup> As with the *attL'* site identified for *S. xylosus* 47–83, the *attL'* of 82B<sup>T</sup> lies downstream of the *lip* gene; however, when compared with the *attL* of the *ccrC*-associated SCCmec *attL* sequences, it shares little similarity and lacks the central cysteine required for *attB/attSCC* recombination (Figure 3b).<sup>27</sup> Indeed, the *attL'* of 82B<sup>T</sup> shows more sequence similarity to that of *attL2* from *S. scuiri* GVGS2 and the other *attL* sites related to CcrAB (Figure 3a). This suggests that *attL2* is most likely linked with *attR2*, though its unusual genomic location within the SCCmec suggests various DNA recombination events may have occurred.

SCCmec include J regions, defined by the areas between the *orfX*, *ccr* and *mec* genes, with the SCCmec of 82B<sup>T</sup> having three such J regions. The J1 region contains primarily hypothetical genes and the *ccrC* gene and shares 86% identity with the J1 region of *S. aureus* PM1 SCCmec. The J2 region upstream of *attL2*, contains genes similar to those of SCCmec type V, with the presence of type 1 restriction modification genes, although there is little similarity between the restriction modification genes of 82B<sup>T</sup> and those present on SCCmec type V.


The J2 region downstream of *attL'* shows the greatest similarity to the SCCmec XI of LGA251, sharing 89% identity, with ABC transporters, genes associated with arsenic resistance and a lipase gene. Unlike the lipase gene in LGA251 the version in 82B<sup>T</sup> appears to be intact. The J3 region is divergent from the rest of the SCCmec of LGA251, with the exception of a putative membrane protein and a putative DNA helicase protein. A notable feature of the J3 region is the presence of a lantibiotic biosynthetic cluster that shares 90% nucleotide identity with one present on the plasmid pETB797 of *S. aureus* NRL 08/797 (accession number KY436025). The cluster encodes two peptide homologues of Lacticin 3147, produced by *Lactococcus lactis*, which has been shown to be active against Gram-positive bacteria.<sup>31</sup> Within the cluster is also a gene encoding a homologue of LtnT, which is required for the transport of the Lacticin 3147 peptides as well as unrelated peptides.<sup>32</sup> The J3 region also contains two ISL3-like transposases, with the majority of the other genes found within SCCmec being from different staphylococci.

### Antimicrobial susceptibility of *S. caeli* 82B<sup>T</sup>

Using Vitek 2, isolate 82B<sup>T</sup> was found to be susceptible to ciprofloxacin, daptomycin, gentamicin, linezolid, mupirocin, rifampicin, teicoplanin, tigecycline and vancomycin. The isolate was resistant to clindamycin, erythromycin, fusidic acid, tetracycline and trimethoprim. Genomic analysis revealed that 82B<sup>T</sup> has the resistance genes *erm(B)*, *erm(C)*, *fusD*, *tet(L)* and *dfcK*, which is consistent with the resistance profile of the isolate. With regards to  $\beta$ -lactam antibiotics, it was resistant to benzylpenicillin (MIC  $\geq 0.5$  mg/L) and oxacillin (MIC 1–2 mg/L) but negative in the cefoxitin resistance screen. However, by disc diffusion it was resistant to cefoxitin and displayed an MIC of 3 mg/L when assessed by Etest. The isolate failed to grow on the MRSA screening agar MRSA Brilliance despite a high inoculum and the use of different incubation temperatures.



**Figure 2.** Overview of *S. caeli* 82B<sup>T</sup> SCCmec. SCCmec Type XI-LGA251 corresponds to *S. aureus* LGA251, accession number FR821779, with SCCmec-82B corresponding to *S. caeli* 82B<sup>T</sup>. Regions of homology are represented by bands connecting the two sequences, with the percentage identity key shown on the right. Blue denotes normal sequence alignment (N); red denotes inverted sequence alignment (I). Key features associated with SCCmec elements are labelled. *att* sites are highlighted by filled circles and labelled above. This figure appears in colour in the online version of JAC and in black and white in the print version of JAC.



**Figure 3.** Comparison of *attR/L* sites from *ccrAB*- or *ccrC*-containing SCCmec. Conserved nucleotide bases within the core 8 bp region, represented in black, bold font, are indicated by an asterisk. The black triangle indicates the position of the central cytosine, thought to be essential for recombination between *attB* and *attSCC*.<sup>27</sup> Inverted repeats are marked by the underlined bases. (a) Sequences of known *attR* and *attL* sites associated with *ccrAB* [from: *S. aureus* N315 (N315), NC\_002745; *S. aureus* LGA251 (LGA251), FR821779; *S. xylosus* S04009 (S04009), HE993884; *S. sciuri* GVGS2 (GVGS2), HG515014; and *S. aureus* ZTA09 (ZTA09), LK024544] were aligned and compared with those identified in *S. caeli* 82B<sup>T</sup>. (b) Sequences of known *attR* and *attL* sites associated with *ccrC* [from: *S. aureus* 85/2082 (85\_2082), AB037671; *S. aureus* WIS (WIS), AB121219; *S. aureus* PM1 (PM1), AB462393; *S. aureus* TSGH17 (TSGH17), AB512767; and *S. aureus* JCSC6082 (JCSC6082), AB373032] were aligned and compared with *attR2* and *attL2* from *S. caeli* 82B<sup>T</sup>.

This study extends the known distribution and diversity of *mecC* genes in terms of country of isolation, sample type, staphylococcal host species, genomic context and allotypes, all of which are novel to the best of our knowledge. These highlight the complex epidemiology of this resistance determinant, particularly among CoNS. It is interesting to note that the species most closely related to *S. caeli* (*S. xylosus*, *S. saprophyticus* and *S. edaphicus*) have all been reported to carry *mecC* or *mecC* allotypes, which may indicate a prominent role for this group in the origins, evolution and epidemiology of *mecC* and SCC*mecC*.

## Acknowledgements

The help of the core sequencing and informatics teams and the Pathogens Informatics team at the Wellcome Trust Sanger Institute (WTSI) is gratefully acknowledged.

## Funding

This project was supported by internal funding from the Royal (Dick) School of Veterinary Studies, University of Edinburgh, a Medical Research Council (MRC) partnership grant (G1001787/1) and the Wellcome Trust, grant number 098051. E. M. H. is supported by a UK Research and Innovation (UKRI) Fellowship: MR/S00291X/1. The funders had no role in the study design, data collection, analysis, decision to publish or preparation of the manuscript.

## Transparency declarations

None to declare.

## References

- García-Álvarez L, Holden MTG, Lindsay H et al. Methicillin-resistant *Staphylococcus aureus* with a novel *mecA* homologue in human and bovine populations in the UK and Denmark: a descriptive study. *Lancet Infect Dis* 2011; **11**: 595–603.
- Shore AC, Deasy EC, Slickers P et al. Detection of staphylococcal cassette chromosome *mec* type XI carrying highly divergent *mecA*, *mecI*, *mecR1*, *blaZ*, and *ccr* genes in human clinical isolates of clonal complex 130 methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* 2011; **55**: 3765–73.
- Peacock SJ, Paterson GK. Mechanisms of methicillin resistance in *Staphylococcus aureus*. *Annu Rev Biochem* 2015; **84**: 577–601.
- Becker K, van Alen S, Idelevich EA et al. Plasmid-encoded transferable *mecB*-mediated methicillin resistance in *Staphylococcus aureus*. *Emerg Infect Dis* 2018; **24**: 242–8.
- Hartman BJ, Tomasz A. Low-affinity penicillin-binding protein associated with  $\beta$ -lactam resistance in *Staphylococcus aureus*. *J Bacteriol* 1984; **158**: 513–6.
- Kim C, Milheiro C, Gardete S et al. Properties of a novel PBP2A protein homolog from *Staphylococcus aureus* strain LGA251 and its contribution to the  $\beta$ -lactam-resistant phenotype. *J Biol Chem* 2012; **287**: 36854–63.
- Becker K, Ballhausen B, Köck R et al. Methicillin resistance in *Staphylococcus* isolates: the “*mec* alphabet” with specific consideration of *mecC*, a *mec* homolog associated with zoonotic *S. aureus* lineages. *Int J Med Microbiol* 2014; **304**: 794–804.
- Paterson GK, Harrison EM, Holmes MA. The emergence of *mecC* methicillin-resistant *Staphylococcus aureus*. *Trends Microbiol* 2014; **22**: 42–7.
- Harrison EM, Coll F, Toleman MS et al. Genomic surveillance reveals low prevalence of livestock-associated methicillin-resistant *Staphylococcus aureus* in the East of England. *Sci Rep* 2017; **7**: 7406.
- Harrison EM, Paterson GK, Holden MTG et al. Whole genome sequencing identifies zoonotic transmission of MRSA isolates with the novel *mecA* homologue *mecC*. *EMBO Mol Med* 2013; **5**: 509–15.
- Worthing KA, Coombs GW, Pang S et al. Isolation of *mecC* MRSA in Australia. *J Antimicrob Chemother* 2016; **71**: 2348–9.
- Harrison EM, Paterson GK, Holden MTG et al. A *Staphylococcus xylosus* isolate with a new *mecC* allotype. *Antimicrob Agents Chemother* 2013; **57**: 1524–8.
- Harrison EM, Paterson GK, Holden MTG et al. A novel hybrid SCC*mecC* region in *Staphylococcus sciuri*. *J Antimicrob Chemother* 2014; **69**: 911–8.
- Semmler T, Harrison EM, Lübke-Becker A et al. A look into the melting pot: the *mecC*-harboring region is a recombination hot spot in *Staphylococcus stepanovicii*. *PLoS One* 2016; **11**: e0147150.
- Małyszko I, Schwarz S, Hauschild T. Detection of a new *mecC* allotype, *mecC2*, in methicillin-resistant *Staphylococcus saprophyticus*. *J Antimicrob Chemother* 2014; **69**: 2003–5.
- Loncaric I, Kubber-Heiss A, Posautz A et al. Characterization of methicillin-resistant *Staphylococcus* spp. carrying the *mecC* gene, isolated from wildlife. *J Antimicrob Chemother* 2013; **14**: 2222–5.
- Pantucek R, Sedlacek I, Indrakova A et al. *Staphylococcus edaphicus* sp. nov., isolated in Antarctica, harbours *mecC* gene and genomic islands with suspected role in adaptation to extreme environment. *Appl Environ Microbiol* 2018; **84**: e01746-17.
- MacFadyen AC, Drigo I, Harrison EM et al. *Staphylococcus caeli* sp. nov., isolated from air sampling in an industrial rabbit holding. *Int J Syst Evol Microbiol* 2018; doi:10.1099/ijsem.0.003098.
- Agnoletti F, Mazzolini E, Bacchin C et al. First reporting of methicillin-resistant *Staphylococcus aureus* (MRSA) ST398 in an industrial rabbit holding and in farm-related people. *Vet Microbiol* 2014; **170**: 172–7.
- Zerbino DR, Birney E. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res* 2008; **18**: 821–9.
- Angiuoli SV, Gussman A, Klimke W et al. Toward an online repository of standard operating procedures (SOPs) for (meta)genomic annotation. *OMICS* 2008; **12**: 137–41.
- Kumar S, Stecher G, Tamura K. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol* 2016; **33**: 1870–4.
- Ito T, Hiramatsu K, Tomasz A et al. Guidelines for reporting novel *mecA* gene homologues. *Antimicrob Agents Chemother* 2012; **56**: 4997–9.
- Paterson GK, Larsen AR, Robb A et al. The newly described *mecA* homologue, *mecA*<sub>LGA251</sub>, is present in methicillin-resistant *Staphylococcus aureus* isolates from a diverse range of host species. *J Antimicrob Chemother* 2012; **67**: 2809–13.
- Liu J, Chen D, Peters BM et al. Staphylococcal chromosomal cassettes *mec* (SCC*mec*): a mobile genetic element in methicillin-resistant *Staphylococcus aureus*. *Microb Pathog* 2016; **101**: 56–67.
- Monecke S, Jatzwauk L, Müller E et al. Diversity of SCC*mec* elements in *Staphylococcus aureus* as observed in south-eastern Germany. *PLoS One* 2016; **11**: e0162654.
- Wang L, Safa M, Archer GL. Characterization of DNA sequences required for the CcrAB-mediated integration of staphylococcal cassette chromosome *mec*, a *Staphylococcus aureus* genomic island. *J Bacteriol* 2012; **194**: 486–98.
- International Working Group on the Classification of Staphylococcal Cassette Chromosome. Classification of staphylococcal cassette chromosome *mec* (SCC*mec*): guidelines for reporting novel SCC*mec* elements. *Antimicrob Agents Chemother* 2009; **53**: 4961–7.

- 29** Zong Z, Lü X. Characterization of a new SCCmec element in *Staphylococcus cohnii*. *PLoS One* 2010; **5**: e14016.
- 30** MacFadyen AC, Harrison EM, Ellington MJ *et al.* A highly conserved *mecC*-encoding SCCmec type XI in a bovine isolate of methicillin-resistant *Staphylococcus xylosus*. *J Antimicrob Chemother* 2018; **73**: 3516–8.
- 31** Galvin M, Hill C, Ross RP. Lacticin 3147 displays activity in buffer against Gram-positive bacterial pathogens which appear insensitive in standard plate assays. *Lett Appl Microbiol* 1999; **28**: 355–8.
- 32** Kuipers A, Meijer-Wierenga J, Rink R *et al.* Mechanistic dissection of the enzyme complexes involved in biosynthesis of lacticin 3147 and nisin. *Appl Environ Microbiol* 2008; **74**: 6591–7.