



Cochrane
Library

Cochrane Database of Systematic Reviews

Genomics-based non-invasive prenatal testing for detection of fetal chromosomal aneuploidy in pregnant women (Review)

Badeau M, Lindsay C, Blais J, Nshimyumukiza L, Takwoingi Y, Langlois S, Légaré F, Giguère Y, Turgeon AF, Witteman W, Rousseau F

Badeau M, Lindsay C, Blais J, Nshimyumukiza L, Takwoingi Y, Langlois S, Légaré F, Giguère Y, Turgeon AF, Witteman W, Rousseau F.

Genomics-based non-invasive prenatal testing for detection of fetal chromosomal aneuploidy in pregnant women.

Cochrane Database of Systematic Reviews 2017, Issue 11. Art. No.: CD011767.

DOI: [10.1002/14651858.CD011767.pub2](https://doi.org/10.1002/14651858.CD011767.pub2).

www.cochranelibrary.com

Genomics-based non-invasive prenatal testing for detection of fetal chromosomal aneuploidy in pregnant women (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	12
Figure 1.	13
Figure 2.	15
OBJECTIVES	16
METHODS	16
RESULTS	19
Figure 3.	20
Figure 4.	23
Figure 5.	25
Figure 6.	26
Figure 7.	27
Figure 8.	28
Figure 9.	30
Figure 10.	31
Figure 11.	32
Figure 12.	34
Figure 13.	35
Figure 14.	36
Figure 15.	37
Figure 16.	38
Figure 17.	38
Figure 18.	39
Figure 19.	39
Figure 20.	40
Figure 21.	41
Figure 22.	42
Figure 23.	43
DISCUSSION	44
AUTHORS' CONCLUSIONS	46
ACKNOWLEDGEMENTS	46
REFERENCES	47
CHARACTERISTICS OF STUDIES	71
DATA	230
Test 1. MPSS T21.	231
Test 2. MPSS T18.	231
Test 3. MPSS T13.	231
Test 4. MPSS 45,X.	232
Test 5. MPSS 47, XXX.	232
Test 6. MPSS 47,XXY.	232
Test 7. MPSS 47,XXY.	232
Test 8. MPSS all 7 aneuploidies.	232
Test 9. MPSS, autosomes.	232
Test 10. MPSS, SCA.	232
Test 11. TMPS T21.	232
Test 12. TMPS T18.	232
Test 13. TMPS T13.	232
Test 14. TMPS 45,X.	232

Test 15. TMPS 47,XXX.	232
Test 16. TMPS 47,XXY.	232
Test 17. TMPS 47,XYY.	233
Test 18. TMPS all 7 aneuploidies.	233
Test 19. TMPS, autosomes.	233
Test 20. TMPS, SCA.	233
Test 21. Traditional screening tests, autosomes.	233
Test 22. Traditional screening tests T21.	233
Test 23. Traditional screening tests T18.	233
Test 24. Traditional screening tests T13.	233
ADDITIONAL TABLES	233
APPENDICES	264
HISTORY	281
CONTRIBUTIONS OF AUTHORS	281
DECLARATIONS OF INTEREST	281
SOURCES OF SUPPORT	282
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	282
INDEX TERMS	283

[Diagnostic Test Accuracy Review]

Genomics-based non-invasive prenatal testing for detection of fetal chromosomal aneuploidy in pregnant women

Mylène Badeau¹, Carmen Lindsay¹, Jonatan Blais^{2,3}, Leon Nshimyumukiza⁴, Yemisi Takwoingi⁵, Sylvie Langlois⁶, France Légaré¹, Yves Giguère^{7,8}, Alexis F Turgeon^{9,10}, William Witteman¹, François Rousseau^{8,10}

¹Population Health and Optimal Health Practices Research Axis, CHU de Québec - Université Laval, Québec City, Canada. ²Department of Medical Biology, CHAU-Hôtel-Dieu de Lévis, Lévis, Canada. ³Department of Molecular Biology, Medical Biochemistry and Pathology, Faculty of Medicine, Université Laval, Québec City, Canada. ⁴School of Public Health, University of Alberta, Edmonton, Canada. ⁵Institute of Applied Health Research, University of Birmingham, Birmingham, UK. ⁶Department of Medical Genetics, Faculty of Medicine, University of British Columbia, Vancouver, Canada. ⁷Reproductive, Mother and Child Health Research Axis, CHU de Québec - Université Laval, Québec City, Canada. ⁸Department of Molecular Biology, Medical Biochemistry and Pathology, Faculty of Medicine, Université Laval, Québec City, Canada. ⁹Department of Anesthesiology and Critical Care Medicine, Division of Critical Care Medicine, and Population Health and Optimal Health Practices Research Unit, CHU de Québec - Université Laval Research Center, CHU de Québec - Université Laval, Université Laval, Québec City, Canada. ¹⁰Population Health and Optimal Health Practices Research Axis, CHU de Québec Research Center, Université Laval, Québec City, Canada

Contact address: François Rousseau, Population Health and Optimal Health Practices Research Axis, CHU de Québec Research Center, Université Laval, 1401, 18^{eme} rue, Québec City, Québec, G1J 1Z4, Canada. francois.rousseau@mac.com, francois.rousseau@fmed.ulaval.ca.

Editorial group: Cochrane Pregnancy and Childbirth Group.

Publication status and date: New, published in Issue 11, 2017.

Citation: Badeau M, Lindsay C, Blais J, Nshimyumukiza L, Takwoingi Y, Langlois S, Légaré F, Giguère Y, Turgeon AF, Witteman W, Rousseau F. Genomics-based non-invasive prenatal testing for detection of fetal chromosomal aneuploidy in pregnant women. *Cochrane Database of Systematic Reviews* 2017, Issue 11. Art. No.: CD011767. DOI: [10.1002/14651858.CD011767.pub2](https://doi.org/10.1002/14651858.CD011767.pub2).

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Common fetal aneuploidies include Down syndrome (trisomy 21 or T21), Edward syndrome (trisomy 18 or T18), Patau syndrome (trisomy 13 or T13), Turner syndrome (45,X), Klinefelter syndrome (47,XXY), Triple X syndrome (47,XXX) and 47,XYY syndrome (47,XYY). Prenatal screening for fetal aneuploidies is standard care in many countries, but current biochemical and ultrasound tests have high false negative and false positive rates. The discovery of fetal circulating cell-free DNA (ccfDNA) in maternal blood offers the potential for genomics-based non-invasive prenatal testing (gNIPT) as a more accurate screening method. Two approaches used for gNIPT are massively parallel shotgun sequencing (MPSS) and targeted massively parallel sequencing (TMPS).

Objectives

To evaluate and compare the diagnostic accuracy of MPSS and TMPS for gNIPT as a first-tier test in unselected populations of pregnant women undergoing aneuploidy screening or as a second-tier test in pregnant women considered to be high risk after first-tier screening for common fetal aneuploidies. The gNIPT results were confirmed by a reference standard such as fetal karyotype or neonatal clinical examination.

Search methods

We searched 13 databases (including MEDLINE, Embase and Web of Science) from 1 January 2007 to 12 July 2016 without any language, search filter or publication type restrictions. We also screened reference lists of relevant full-text articles, websites of private prenatal diagnosis companies and conference abstracts.

Selection criteria

Studies could include pregnant women of any age, ethnicity and gestational age with singleton or multifetal pregnancy. The women must have had a screening test for fetal aneuploidy by MPSS or TMPS and a reference standard such as fetal karyotype or medical records from birth.

Data collection and analysis

Two review authors independently carried out study selection, data extraction and quality assessment (using the QUADAS-2 tool). Where possible, hierarchical models or simpler alternatives were used for meta-analysis.

Main results

Sixty-five studies of 86,139 pregnant women (3141 aneuploids and 82,998 euploids) were included. No study was judged to be at low risk of bias across the four domains of the QUADAS-2 tool but applicability concerns were generally low. Of the 65 studies, 42 enrolled pregnant women at high risk, five recruited an unselected population and 18 recruited cohorts with a mix of prior risk of fetal aneuploidy. Among the 65 studies, 44 evaluated MPSS and 21 evaluated TMPS; of these, five studies also compared gNIPT with a traditional screening test (biochemical, ultrasound or both). Forty-six out of 65 studies (71%) reported gNIPT assay failure rate, which ranged between 0% and 25% for MPSS, and between 0.8% and 7.5% for TMPS.

In the population of unselected pregnant women, MPSS was evaluated by only one study; the study assessed T21, T18 and T13. TMPS was assessed for T21 in four studies involving unselected cohorts; three of the studies also assessed T18 and T13. In pooled analyses (88 T21 cases, 22 T18 cases, eight T13 cases and 20,649 unaffected pregnancies (non T21, T18 and T13)), the clinical sensitivity (95% confidence interval (CI)) of TMPS was 99.2% (78.2% to 100%), 90.9% (70.0% to 97.7%) and 65.1% (9.16% to 97.2%) for T21, T18 and T13, respectively. The corresponding clinical specificity was above 99.9% for T21, T18 and T13.

In high-risk populations, MPSS was assessed for T21, T18, T13 and 45,X in 30, 28, 20 and 12 studies, respectively. In pooled analyses (1048 T21 cases, 332 T18 cases, 128 T13 cases and 15,797 unaffected pregnancies), the clinical sensitivity (95% confidence interval (CI)) of MPSS was 99.7% (98.0% to 100%), 97.8% (92.5% to 99.4%), 95.8% (86.1% to 98.9%) and 91.7% (78.3% to 97.1%) for T21, T18, T13 and 45,X, respectively. The corresponding clinical specificities (95% CI) were 99.9% (99.8% to 100%), 99.9% (99.8% to 100%), 99.8% (99.8% to 99.9%) and 99.6% (98.9% to 99.8%). In this risk group, TMPS was assessed for T21, T18, T13 and 45,X in six, five, two and four studies. In pooled analyses (246 T21 cases, 112 T18 cases, 20 T13 cases and 4282 unaffected pregnancies), the clinical sensitivity (95% CI) of TMPS was 99.2% (96.8% to 99.8%), 98.2% (93.1% to 99.6%), 100% (83.9% to 100%) and 92.4% (84.1% to 96.5%) for T21, T18, T13 and 45,X respectively. The clinical specificities were above 100% for T21, T18 and T13 and 99.8% (98.3% to 100%) for 45,X. Indirect comparisons of MPSS and TMPS for T21, T18 and 45,X showed no statistical difference in clinical sensitivity, clinical specificity or both. Due to limited data, comparative meta-analysis of MPSS and TMPS was not possible for T13.

We were unable to perform meta-analyses of gNIPT for 47,XXX, 47,XXY and 47,XYY because there were very few or no studies in one or more risk groups.

Authors' conclusions

These results show that MPSS and TMPS perform similarly in terms of clinical sensitivity and specificity for the detection of fetal T21, T18, T13 and sex chromosome aneuploidy (SCA). However, no study compared the two approaches head-to-head in the same cohort of patients. The accuracy of gNIPT as a prenatal screening test has been mainly evaluated as a second-tier screening test to identify pregnancies at very low risk of fetal aneuploidies (T21, T18 and T13), thus avoiding invasive procedures. Genomics-based non-invasive prenatal testing methods appear to be sensitive and highly specific for detection of fetal trisomies 21, 18 and 13 in high-risk populations. There is paucity of data on the accuracy of gNIPT as a first-tier aneuploidy screening test in a population of unselected pregnant women. With respect to the replacement of invasive tests, the performance of gNIPT observed in this review is not sufficient to replace current invasive diagnostic tests.

We conclude that given the current data on the performance of gNIPT, invasive fetal karyotyping is still the required diagnostic approach to confirm the presence of a chromosomal abnormality prior to making irreversible decisions relative to the pregnancy outcome. However, most of the gNIPT studies were prone to bias, especially in terms of the selection of participants.

PLAIN LANGUAGE SUMMARY

Accuracy of gNIPT for identifying genetic abnormalities in unborn babies

What is the issue?

How accurate is the new test (genomics-based non-invasive prenatal testing (gNIPT)) for detecting abnormal chromosome number in an unborn baby's genetic material (DNA) found in the mother's blood? We assessed the accuracy for the screening of Down syndrome (trisomy 21), Edward syndrome (trisomy 18), Patau syndrome (trisomy 13), Turner syndrome (45,X), Klinefelter syndrome (47,XXY), Triple X syndrome (47,XXX) and 47,XYY syndrome. There are different methods in use for gNIPT. We assessed MPSS (massively parallel shotgun sequencing) that tests whole DNA and TMPS (targeted massively parallel sequencing) that tests targeted DNA.

Background

There are 46 chromosomes (23 pairs) in humans. Abnormal numbers of chromosomes can cause genetic disorders for which there are no cures. Having an extra chromosome is called trisomy and an excess (or less) of sexual chromosome is called sex chromosome abnormality (SCA). The most common trisomy is Down syndrome which occurs in about one in 1000 babies. Children with Downs have slow growth, characteristic facial features and mild to moderate intellectual disability, with some requiring specialist education later in life. However, the symptoms vary from mild to severe so that some infants lead relatively normal lives. The other trisomy or SCA conditions have varying degrees of disability but the chance of a baby being affected is much less.

Current screening tests for these conditions require confirmation if the baby has the condition or not and for this an invasive test like amniocentesis is used. Amniocentesis is where fetal cells that float in the fluid surrounding the unborn baby are collected by putting a fine needle through the mother's abdomen and collecting the fluid. Alternatively, tissue can be collected from the placenta (chorionic villus sampling (CVS)). With these invasive tests, pregnant women are exposed to a higher chance of losing their baby even if the baby is unaffected by Down syndrome. So, this invasive test is only offered to women who are thought to have a higher chance of having an affected unborn baby

What we did

We looked for studies that included women of any age, ethnicity and gestational age who were carrying either a single baby or more than one. We searched for studies (up to July 2016) that assessed the accuracy of the new test.

What we found

We found 65 studies with a total of 86,139 pregnant women, including 3141 affected pregnancies. Forty-two studies (65%) enrolled pregnant women with a high chance of having babies with abnormal chromosome number. Forty-eight (74%) studies included only women with a singleton pregnancy. Forty-four studies (68%) used MPSS and 21 studies (32%) used TMPS.

gNIPT seems to be accurate for screening unborn babies (either singletons or twins), especially for detecting Down syndrome, trisomy 18 and trisomy 13. However, there were some problems with how the studies were conducted which makes us cautious about our findings. This may result in gNIPT appearing to perform better than it really does.

Other important information to consider

gNIPT method appears to perform well in identifying unborn babies with abnormal number of chromosomes. However, when a gNIPT detects an abnormal chromosome number, then a confirmation using invasive tests (like amniocentesis or CVS) is still needed before pregnancy-related decisions can be made.

It is important that pregnant women are given full information on the possible health problems that might arise for babies affected by an additional chromosome. For example, with Down syndrome though some children have considerable disability, others can lead relatively normal lives. In addition, in this review most studies enrolled pregnant women with increased chance of having babies with abnormal chromosome number, so our findings do not directly apply to general populations of pregnant women.

SUMMARY OF FINDINGS

Summary of findings 1. Summary characteristics of included studies

Summary characteristics of included studies	
Review question	What is the diagnostic accuracy of massively parallel shotgun sequencing (MPSS) and targeted massively parallel sequencing (TMPS) using circulating cell-free DNA (ccfDNA) in maternal blood for the detection of common fetal aneuploidies (T21, T18, T13, 45,X, 47,XXY, 47,XXX and 47,XYY) in pregnant women according to their prior risk of fetal aneuploidy?
Importance (rationale)	These new genomics-based non-invasive prenatal testing (gNIPT) approach report higher sensitivity and lower false positive rate than traditional screening tests. gNIPT is already advertised and marketed. How gNIPT should be used in clinical practice should be assessed in order to provide a framework for its use.
Study design	There were 40 prospective cohort studies, 8 retrospective cohort studies, 16 case-control studies and 1 prospective and retrospective cohort study.
Population	Pregnant women of any age, ethnicity and gestational age, with singleton or multifetal pregnancy who had a screening test for fetal aneuploidy using gNIPT and received a reference standard. 42 studies enrolled pregnant women selected at high risk of fetal aneuploidy, 5 enrolled unselected pregnant women undergoing aneuploidy screening and 18 enrolled pregnant women from a mixed-risk population of fetal aneuploidy. 48 studies included only women with singleton pregnancy, 5 included only multifetal pregnancies, 4 included either type of pregnancy and 8 did not report type of pregnancy. 10 studies included only women in the first trimester (15 weeks or less), 21 studies included women in the first 2 trimesters (29 weeks or less), 24 studies included women in the 3 trimesters (42 weeks or less) and 10 studies (15%) did not report gestational age.
Index tests	gNIPT by MPSS (44 studies) or TMPS (21 studies), including 5 studies that compared a gNIPT with a traditional screening test. 37 studies were industry-funded or were written by 1 or more authors affiliated with a company who sells gNIPT. 22 studies were not reported to be funded by industry but samples were sequenced and analysed by a commercial laboratory. 3 studies had no links with industry.
Target conditions	36 studies reported results for only autosomes (T21, T18, T13), 4 for only SCA (45,X, 47,XXY, 47,XXX and 47,XYY), and 25 for both autosomes and SCA.
Reference standard	Fetal karyotyping performed on cells obtained from chorionic villi sampling, amniotic fluid, placental tissue, a fetus lost by miscarriage or other equivalent and recognised methods on the same materials for autosomes and SCA. If fetal karyotyping was not performed, we used neonatal clinical examination or medical records from birth (for autosomes only). Only 1 reference standard was used for all pregnant women included in 36 studies while multiple reference standards were used in 29 studies.
Risk of bias	<p>The QUality Assessment of Diagnostic Accuracy Studies (QUADAS-2) tool was used to assess the methodological quality of included studies.</p> <p>No study was assessed as being at low risk of bias across all domains. For the patient selection domain, no study was assessed as being at low risk of bias. For the index test, reference standard and flow and timing domains, the risk of bias was low for 94%, 77% and 23% of studies, respectively.</p>

Applicability concerns

Applicability was of low concern for all studies in the index test and reference standard domains because the studies matched the review question. In the patient selection domain, 47 (71%) studies were judged to be of low applicability concern because they included pregnant women matching the review question.

45,X: Turner syndrome, 47,XXX: triple X syndrome, 47,XXY: Klinefelter syndrome, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, T21: trisomy 21, T18: trisomy 18, T13: trisomy 13.

Summary of findings 2. Performance of gNIPT for detection of T21

Performance of gNIPT for detection of T21

Test strategy	Number of studies	Number of affected pregnancies (Number of unaffected pregnancies) ^a	Sensitivity % (95% CI)	Specificity % (95% CI)	Median prevalence ^b % (range)	Missed cases (FN) ^c	False positives (FP) ^d
Unselected pregnant women							
MPSS	1	8 (1733)	100 (67.6 to 100)	100 (99.8 to 100)	0.46 (0.24 to 5.21)	0	0
TMPS	4	88 (20,679)	99.2 (78.2 to 100)	100 (> 99.9 to 100)		4	0
Traditional screening test ^e	1	38 (15,803)	78.9 (63.7 to 88.9)	94.6 (94.2 to 94.9)		97	5375
Implications	<ul style="list-style-type: none"> • 460 of 100,000 pregnancies expected to be affected by T21; • MPSS will detect all cases and no pregnant woman will undergo an unnecessary invasive test; • with TMPS, 4 cases will be missed and no pregnant woman will undergo unnecessary invasive test; and • with traditional screening tests, 363 cases will be detected and 5375 unaffected pregnant women will undergo unnecessary invasive test. 						
Selected high-risk pregnant women							
MPSS	30	1048 (15,937)	99.7 (98.0 to 100)	99.9 (99.8 to 100)	4.95 (0.44 to 27.66)	15	95
TMPS	6	246 (4380)	99.2 (96.8 to 99.8)	100 (99.8 to 100)		40	0
Difference between MPSS and TMPS			0.53 (-0.73 to 1.78)	-0.03 (-0.11 to 0.04)	NA		

- Implications
- 4950 of 100,000 pregnancies expected to be affected by T21;
 - 4936 and 4911 cases will be detected while 15 and 40 cases will be missed by MPSS and TMPS, respectively; and
 - of 95,050 expected pregnancies unaffected by T21, 95 and 0 pregnant women will undergo unnecessary invasive tests with MPSS and TMPS, respectively.

MPSS: massively parallel shotgun sequencing, NA; not applicable, TMPS: targeted massively parallel sequencing, T21: trisomy 21.

^aUnaffected pregnancies: we included patients with any other aneuploidy than the one under analysis with all euploid cases as "unaffected" pregnancies.

^bThe median prevalence and range were calculated by using all prospective or retrospective studies for each category considered.

^cMissed cases per 100,000 tested. FN: false negatives.

^dFalse positives per 100,000 tested. A false positive result may lead to unnecessary invasive tests depending on choices by the pregnant woman.

^eTraditional screening tests are first-trimester combined test, second-trimester quadruple test, second-trimester fully integrated test, second-trimester sequential test or second-trimester triple test.

Summary of findings 3. Performance of gNIPT for detection of T18

Performance of gNIPT for detection of T18

Test strategy	Number of studies	Number of affected pregnancies (Number of unaffected pregnancies) ^a	Sensitivity % (95% CI)	Specificity % (95% CI)	Median prevalence ^b % (range)	Missed cases (FN) ^c	False positives (FP) ^d
Unselected pregnant women							
MPSS	1	2 (1739)	100 (34.3 to 100)	99.9 (99.7 to 100)	0.11 (0.06 to 0.36)	0	100
TMPS	3	22 (20,553)	90.9 (70.0 to 97.7)	100 (99.9 to 100)		10	0
Traditional screening test ^e	1	10 (15,831)	80.0 (49.0 to 94.3)	99.7 (99.6 to 99.8)		22	300

- Implications
- 109 of 100,000 pregnancies expected to be affected by T18;
 - MPSS will detect all cases and 100 unaffected pregnant women will undergo an unnecessary invasive test;
 - with TMPS, 10 cases will be missed and no unaffected pregnant woman will undergo unnecessary invasive test; and
 - with traditional screening tests, 87 cases will be detected, 22 will be missed and 300 unaffected pregnant women will undergo unnecessary invasive test.

Selected high-risk pregnant women

MPSS	28	332 (16,180)	97.8 (92.5 to 99.4)	99.9 (99.8 to 100)	1.46 (0.22 to 17.02)	32	99
TMPS	5	112 (4010)	98.2 (93.1 to 99.6)	100 (99.8 to 100)		26	0
Difference between MPSS and TMPS			-0.41 (-4.11 to 3.28)	-0.06 (-0.14 to 0.03)	NA		
Implications	<ul style="list-style-type: none"> • 1463 of 100,000 pregnancies expected to be affected by T18; • 1431 and 1437 cases will be detected while 32 and 26 cases will be missed by MPSS and TMPS, respectively; and • of 98,537 expected unaffected by T18, 99 and 0 pregnant women will undergo unnecessary invasive test with MPSS and TMPS, respectively. 						

MPSS: massively parallel shotgun sequencing, NA: not applicable, TMPS: targeted massively parallel sequencing, T18: trisomy 18.

^aUnaffected pregnancies: we included patients with any other aneuploidy than the one under analysis with all euploid cases as "unaffected" pregnancies.

^bThe median prevalence and range were calculated by using all prospective or retrospective studies for each category considered.

^cMissed cases per 100,000 tested. FN: false negatives.

^dFalse positives per 100,000 tested. A false positive result may lead to unnecessary invasive tests depending on choices by the pregnant woman.

^eTraditional screening tests are first-trimester combined test, second-trimester quadruple test, second-trimester fully integrated test, second-trimester sequential test or second-trimester triple test.

Summary of findings 4. Performance of gNIPT for detection of T13

Performance of gNIPT for detection of T13

Test strategy	Number of studies	Number of affected pregnancies (Number of unaffected pregnancies) ^a	Sensitivity % (95% CI)	Specificity % (95% CI)	Median prevalence ^b % (range)	Missed cases (FN) ^c	False positives (FP) ^d
Unselected pregnant women							
MPSS	1	1 (1740)	100 (20.7 to 100)	100 (99.8 to 100)	0.12 (0.01 to 0.52)	0	0
TMPS	3	8 (14,154)	65.1 (9.16 to 97.2)	100 (99.9 to 100)		41	0
Traditional screening test ^e	1	2 (11,183)	50.0 (9.45 to 90.5)	99.7 (99.6 to 99.8)		59	300

- Implications
- 118 of 100,000 pregnancies expected to be affected by T13;
 - MPSS will detect all cases and no unaffected pregnant woman will undergo an unnecessary invasive test;
 - with TMPS, 41 cases will be missed and no unaffected pregnant woman will undergo unnecessary invasive test; and
 - with traditional screening tests, 59 cases will be missed and 300 unaffected pregnant women will undergo unnecessary invasive test.

Selected high-risk pregnant women

MPSS	20	128 (13,810)	95.8 (86.1 to 98.9)	99.8 (99.8 to 99.9)	1.09 (0.04 to 3.54)	46	198
TMPS	2	20 (293)	100 (83.9 to 100) ^f	100 (98.7 to 100) ^f		0	0

- Implications
- 1087 of 100,000 pregnancies expected to be affected by T13;
 - 1041 and 1087 cases will be detected while 46 and 0 cases will be missed by MPSS and TMPS, respectively; and
 - of 98,913 expected unaffected by T13, 198 and 0 pregnant women will undergo unnecessary invasive test with MPSS and TMPS, respectively.

MPSS: massively parallel shotgun sequencing, NA: not applicable, TMPS: targeted massively parallel sequencing, T13: trisomy 13.

^aUnaffected pregnancies: we included patients with any other aneuploidy than the one under analysis with all euploid cases as "unaffected" pregnancies.

^bThe median prevalence and range were calculated by using all prospective or retrospective studies for each category considered.

^cMissed cases per 100,000 tested. FN: false negatives.

^dFalse positives per 100,000 tested. A false positive result may lead to unnecessary invasive tests depending on choices by the pregnant woman.

^eTraditional screening tests are first-trimester combined test, second-trimester quadruple test, second-trimester fully integrated test, second-trimester sequential test or second-trimester triple test.

^fSimple pooling used to obtain summary estimates of sensitivity, specificity or both.

Summary of findings 5. Performance of gNIPT for detection of 45,X

Performance of gNIPT for detection of 45,X

Test strategy	Number of studies	Number of affected pregnancies (Number of unaffected pregnancies) ^a	Sensitivity % (95% CI)	Specificity % (95% CI)	Median prevalence ^b % (range)	Missed cases (FN) ^c	False positives (FP) ^d
Selected high-risk pregnant women							
MPSS	12	119 (7440)	91.7 (78.3 to 97.1)	99.6 (98.9 to 99.8)	1.04 (0.27 to 18.58)	86	396
TMPS	4	79 (985)	92.4 (84.1 to 96.5)	99.8 (98.3 to 100)		79	198
Difference between MPSS and TMPS			-0.74 (-11.1 to 9.60)	-0.23 (-0.82 to 0.36)	NA		

- Implications
- 1039 of 100,000 pregnancies expected to be affected by 45X;
 - 953 and 960 cases will be detected while 86 and 79 cases will be missed by MPSS and TMPS, respectively; and
 - of 98,961 expected unaffected by 45X, 396 and 198 pregnant women will undergo unnecessary invasive test with MPSS and TMPS, respectively.

45,X: Turner syndrome, MPSS: massively parallel shotgun sequencing, NA: not applicable, TMPS: targeted massively parallel sequencing.

^aUnaffected pregnancies: we included patients with any other aneuploidy than the one under analysis with all euploid cases as "unaffected" pregnancies.

^bThe median prevalence and range were calculated by using all prospective or retrospective studies for each category considered.

^cMissed cases per 100,000 tested. FN: false negatives.

^dFalse positives per 100,000 tested. A false positive result may lead to unnecessary invasive tests depending on choices by the pregnant woman.

Summary of findings 6. Performance of gNIPT for detection of autosomes aneuploidies (T21, T18 and T13 combined)

Performance of gNIPT for detection of autosomes aneuploidies (T21, T18 and T13 combined)

Test strategy	Number of studies	Number of affected pregnancies (Number of unaffected pregnancies) ^a	Sensitivity % (95% CI)	Specificity % (95% CI)	Median prevalence ^b % (range)	Missed cases (FN) ^c	False positives (FP) ^d
Unselected pregnant women							
MPSS	1	11 (1730)	100 (74.1 to 100)	99.9 (99.7 to 100)	0,63	0	99
TMPS	4	118 (20,649)	94.9 (89.1 to 97.7)	99.9 (99.8 to 99.9)	(0.32 to 5.73)	32	99
Traditional screening test ^e	4	120 (22,247)	ND ^f			ND	
Implications	<ul style="list-style-type: none"> • 632 of 100,000 pregnancies expected to be affected by T21, T18 or T13; • 632 and 600 cases will be detected whereas 0 and 32 cases will be missed by MPSS and TMPS, respectively; and • of 99,368 unaffected, 99 pregnant women will undergo unnecessary invasive test with MPSS or TMPS. 						
Selected high-risk pregnant women							
MPSS	32	1508 (15,797)	98.8 (97.2 to 99.5)	99.9 (99.7 to 100)	5.85	70	94
TMPS	7	378 (4282)	98.9 (97.2 to 99.6)	99.9 (99.8 to 100)	(0.67 to 46.81)	64	94
Difference between MPSS and TMPS			-0.11	-0.08	NA		

(-1.58 to 1.35) (-0.22 to 0.07)

- Implications
- 5851 of 100,000 pregnancies expected to be affected by T21, T18 or T3;
 - 5781 and 5787 cases will be detected, whereas 70 and 64 cases will be missed by MPSS and TMPS, respectively; and
 - of 94,149 unaffected, 94 pregnant women will undergo unnecessary invasive test with MPSS or TMPS.

MPSS: massively parallel shotgun sequencing, NA: not applicable, ND: no data available, TMPS: targeted massively parallel sequencing, T13: trisomy 13, T18: trisomy 18, T21: trisomy 21.

^aUnaffected pregnancies: we included patients with any other aneuploidy than the one under analysis with all euploid cases as "unaffected" pregnancies.

^bThe median prevalence and range were calculated by using all prospective or retrospective studies for each category considered.

^cMissed cases per 100,000 tested. FN: false negatives.

^dFalse positives per 100,000 tested. A false positive result may lead to unnecessary invasive tests depending on choices by the pregnant woman.

^eTraditional screening tests are first-trimester combined test, second-trimester quadruple test, second-trimester fully integrated test, second-trimester sequential test or second-trimester triple test.

^fSummary sensitivity and specificity were not obtained for traditional screening tests because the four studies used different cut-offs to determine test positivity. Three of the four studies compared TMPS and traditional screening tests in the same population (direct comparison).

Summary of findings 7. Performance of gNIPT for detection of sex chromosome aneuploidies (45,X, 47,XXX, 47,XXY and 47,XYY combined)^a

Performance of gNIPT for detection of sex chromosome aneuploidies (45,X, 47,XXX, 47,XXY and 47,XYY combined)

Test strategy	Number of studies	Number of affected pregnancies (Number of unaffected pregnancies) ^b	Sensitivity % (95% CI)	Specificity % (95% CI)	Median prevalence ^c % (range)	Missed cases (FN) ^d	False positives (FP) ^e
Selected high-risk pregnant women							
MPSS	12	151 (7452)	91.9 (73.8 to 97.9)	99.5 (98.8 to 99.8)	1.53 (0.45 to 18.58)	124	492
TMPS	4	96 (968)	93.8 (86.8 to 97.2)	99.6 (98.1 to 99.9)		95	394
Difference between MPSS and TMPS			-1.85 (-13.3 to 9.60)	-0.06 (-0.82 to 0.71)	NA		

- Implications
- 1535 of 100,000 pregnancies expected to be affected by SCA;
 - 1411 and 1440 cases will be detected while 124 and 95 cases will be missed by MPSS and TMPS, respectively;
 - of 98,465 unaffected by SCA, 492 and 394 pregnant women will undergo unnecessary invasive test with MPSS and TMPS, respectively.

45,X: Turner syndrome, 47,XXX: triple X syndrome, 47,XXY: Klinefelter syndrome, MPSS: massively parallel shotgun sequencing, NA: not applicable, ND: no data available, TMPS: targeted massively parallel sequencing

^aWe did not assess the accuracy of gNIPT individually for 47,XXX, 47,XXY and 47,XYY due to paucity data.

- ^bUnaffected pregnancies: we included patients with any other aneuploidy than the one under analysis with all euploid cases as "unaffected" pregnancies.
- ^cThe median prevalence and range were calculated by using all prospective or retrospective studies for each category considered.
- ^dMissed cases per 100,000 tested. FN: false negatives.
- ^eFalse positives per 100,000 tested. A false positive result may lead to unnecessary invasive tests depending on choices by the pregnant woman.

BACKGROUND

Aneuploidies^[1] are chromosomal abnormalities characterised by a different (additional or missing) number of chromosomes than the 23 pairs normally present in humans. These chromosomal anomalies are among the most common types of genetic disorders and they represent a significant cause of both childhood and adulthood morbidity or death. In addition, they may lead to perinatal complications (Wellesley 2012; Wu 2013a). The severity of associated symptoms is often variable and typically less severe in mosaic cases (not all cells affected) (Fishler 1991; Modi 2003; Zhu 2013). Although offering prenatal screening for fetal aneuploidies such as Down syndrome is now considered standard of care in routine antenatal care in most upper-middle and high-income countries, prenatal screening methods and strategies are evolving. Prenatal screening consists of blood-based biochemical testing or ultrasound measurements or a combination of both, in addition to maternal age (Aldred 2012). Because of the serious health consequences of various aneuploidies and given their incurable nature, prenatal screening is an option available to pregnant women. An invasive diagnostic test (e.g. amniocentesis) is offered to pregnant women found to be at high risk of fetal aneuploidy after prenatal screening, but there is a procedure-related risk of miscarriage. The discovery of circulating cell-free DNA (ccfDNA) in maternal blood has enabled the development of genomics-based non-invasive prenatal testing (gNIPT) to analyse the fetal genome. Prenatal screening, and ultimately prenatal diagnosis, provides couples with the information necessary for taking informed decisions (the optimisation of medical intervention and psychological counselling for managing the identified condition or pregnancy termination). The decision to terminate pregnancy among women who received a positive diagnosis of fetal aneuploidy during the prenatal period varies between 86% and 97% (Choi 2012; Irving 2011). Many factors, such as religion, maternal age, gestational age at the time of diagnosis, number of existing children, past history of induced abortion and psychosocial factors (perceived parenting burden/reward, quality of life of a child with a chromosomal abnormality, attitudes toward, and comfort with individuals with disabilities, and support from others) influence women's decision making following prenatal anomaly detection (Choi 2012).

In this systematic review, we assessed the accuracy of gNIPT for the detection of common fetal aneuploidies in pregnant women according to their prior risk of fetal aneuploidy. More specifically, we evaluated and compared the diagnostic performance of two new next-generation sequencing approaches (i.e. massively parallel shotgun sequencing (MPSS) and targeted massively parallel sequencing (TMPS)) that have recently been proposed as methods of choice to detect fetal aneuploidies by analysing ccfDNA in maternal plasma. We also made comparisons between MPSS and TMPS or between gNIPT and their combination with other first-tier screening approaches. gNIPT could be used as a first-tier test in pregnant women without prior risk (i.e. in unselected pregnant women or the general population) or as a second-tier test after a positive result for traditional first-tier screening tests such as biochemical, ultrasound or both markers (with maternal age

included in risk assessment) and previous maternal history when possible.

[1] For a glossary of terms, see [Appendix 1](#). For a list of acronyms and abbreviations, see [Appendix 2](#).

Target condition being diagnosed

The target conditions are fetal chromosomal abnormalities diagnosed in pregnant women. The seven target conditions assessed were Down syndrome (trisomy 21 or T21), Edward syndrome (trisomy 18 or T18), Patau syndrome (trisomy 13 or T13), Turner syndrome (45,X), Klinefelter syndrome (47,XXY), Triple X syndrome (47,XXX) and 47,XYY syndrome (47,XYY) ([Table 1](#)). The majority of aneuploidies are associated with an extra copy (trisomy) of one chromosome (e.g. three copies of chromosome 21 for T21 instead of two) or a loss of one chromosome (e.g. female 45,X). Chromosomal abnormality is usually caused by a chromosome division failure or a chromosomal translocation. For example, most cases (76.2%) of 45,X karyotype (all cells affected) are caused by paternal chromosome division failure ([Uematsu 2002](#)). The most common chromosomal abnormalities are T21 and 45,X, respectively. For T21, the prevalences reported for pregnant women are 0.11% and 0.44% at 25 and 35 years old, respectively at diagnosis procedure ([Snijders 1999](#)).


Clinical characteristics and spectrum of severity are variable among aneuploidies. It has been reported that 50% of 45,X cases are mosaic ([Sybert 2004](#)). During the past few decades, caring for children with T21 or sex chromosomal abnormalities and provision of counselling to their family has changed fundamentally. These changes, including medical and surgical advances, specific interventions in the classroom for those with learning disabilities, interventions and support for parents and family members, have helped individuals with T21 live longer and enjoy an improved quality of life ([Van Riper 2001](#)). Many health problems associated with T21, 45,X, 47,XXY, 47,XXX and 47,XYY aneuploidies can be treated but fetuses with T18 and T13 are most affected and usually die in utero. The age at diagnosis varies widely depending on the condition. T21, T18 and T13 are generally detected during the perinatal period, while detection of 45,X, 47,XXX and 47,XYY is often delayed, sometimes up to 60 years old ([Stochholm 2006](#); [Stochholm 2010a](#); [Tartaglia 2010](#)). Around 10% of fetuses with 47,XXY are diagnosed prenatally and the mean age at diagnosis is in the mid-30s. Most 47,XXY cases are never diagnosed ([Groth 2013](#); [Tyler 2004](#)). The incidence, clinical features and prognosis of the target conditions are summarised in [Table 1](#).

Index test(s)

Genomics-based non-invasive prenatal tests are based on the finding that placental cells continuously release detectable amounts of fetal ccfDNA into maternal blood. This fetal ccfDNA originates from normal placental cell death and consists mainly of relatively short fragments of < 300 base pairs ([Bianchi 2004](#); [Fan 2010](#)). Proof-of-concept studies showed the feasibility of such tests to detect fetal aneuploidy in 2008 ([Chiu 2008](#); [Fan 2008](#)).

We assessed these two gNIPT approaches ([Figure 1](#)):

Figure 1. Difference between massively parallel shotgun sequencing (MPSS) and targeted massively parallel sequencing (TMPS). Genomics-based non-invasive prenatal testing (gNIPT) aims to count the number of copies of DNA fragments from the chromosomes of interest (chromosome 21 (Chrom. 21) in this example) present in circulating cell-free DNA (ccfDNA) from a pregnant woman, relative to a reference set of chromosomes (Ref. Chrom.). DNA fragments circulating in maternal blood in the case of a euploid (left) and aneuploid (right) pregnancy are illustrated (top). MPSS produces a large number of sequence reads from all chromosomes while TMPS generates a larger proportion of reads from the chromosomes of interest (bottom). In both methods, sequence reads can be used to detect a slight excess of fetal genomic material coming from the chromosome of interest. Figure was created by FR.



- massively parallel shotgun sequencing (MPSS) which randomly analyses all DNA fragments of a sample; and
- targeted massively parallel sequencing (TMPS) which targets specific DNA fragments from the chromosomal regions of interest.

The fraction of the total ccfDNA in maternal circulation that is of fetal origin (the fetal fraction) is an important parameter for correctly identifying an aneuploid fetus by gNIPT (Canick 2013). Although the fetal ccfDNA fraction is a relatively small fraction (about 2% to 20%) of all ccfDNA in maternal blood, it can be detected from five weeks of gestation (Birch 2005; Canick 2013; Lo 1997; Lun 2008). Invasive procedures such as amniocentesis, may (Samura 2003) or may not be (Bussani 2011; Vora 2010) associated with a statistically significant increase of ccfDNA in maternal blood, which could affect fetal DNA concentration and affect gNIPT results. Therefore, in the context of clinical studies, maternal blood for

gNIPT is usually collected either before or after waiting for a minimum of 24 hours following an invasive test. Indeed, the half-life of ccfDNA has been estimated to be less than one day (Lo 1999; Yu 2013). On average, euploid multifetal pregnancies have a higher fetal ccfDNA fraction than euploid singleton pregnancies (Attilakos 2011; Canick 2012). There is no reported difference in ccfDNA concentration between monozygotic and dizygotic multifetal pregnancies (Attilakos 2011). However, dizygotic pregnancies complicate gNIPT analysis by the presence of an additional genome (or more in the presence of more than two fetuses) as opposed to the two genomes of mother and fetus present in singleton or monozygotic twin pregnancies.

Next generation sequencing (NGS) applied on DNA extracted from the plasma of pregnant women generates millions of DNA sequences from both maternal and fetal genomes in relative proportion to their original abundance (for technical details see Appendix 3). The data thus produced can be used to detect a slight

excess (or loss) of fetal genomic material associated with cases of fetal aneuploidy (Papageorgiou 2012). These NGS technologies have paved the way for the development of gNIPT by alleviating the need for fetal-specific genetic markers and with potentially better test accuracy than current fetal aneuploidy screening methods.

Currently, gNIPT for the detection of common aneuploidies has been developed by companies in America, Asia and Europe and are commercially available. As part of their marketing material, these companies have published the diagnostic performance of their respective tests on their websites (Table 2). In addition, several research and clinical laboratories have developed in-house gNIPT.

Before taking a personal decision to accept or decline gNIPT, pregnant women should be given information on the screening process, which must include a discussion with a health professional (Gagnon 2010; Legare 2010; Legare 2011; St-Jacques 2008). Following screening, the results should be explained in the context of the harms and benefits of definitive diagnosis through non directive counselling (Benn 2013b). In their recent guideline, the American College of Obstetricians and Gynecologists (ACOG) recommends that gNIPT should not be used to replace diagnostic testing and that all pregnant women with a positive gNIPT result should have a diagnostic procedure before undertaking any irreversible action such as pregnancy termination. Guidelines also recommend that pregnant women with an unreported, indeterminate or uninterpretable gNIPT result should receive further genetic counselling and be offered comprehensive ultrasound evaluation and diagnostic testing (ACOG #163 2016).

Clinical pathway

Prior test(s)

Prenatal screening for fetal aneuploidy (mostly T21) is part of public health programs in most upper-middle and high-income countries and is typically offered to all pregnant women (Benn 2013b; Chitayat 2011). Up to now, screening tests for aneuploidies have relied on blood-based biochemical testing of placental markers with or without ultrasound imaging to assess for nuchal translucency thickness and other markers of fetal aneuploidy in the first trimester. The age of the pregnant woman is combined with levels of biomarkers and nuchal translucency as predictive markers for T21 in the first or second trimester (Benn 2011; Chitayat 2011; Summers 2007). Table 3 presents the various testing combinations (e.g. sequential, integrated or contingent algorithms) that have been described and are currently in use in prenatal clinics (Allred 2017b). The screening performance of these algorithms is mostly related to the detection rates of different marker combinations and the accepted level of false positive rates. A large prospective Canadian study of 32,227 pregnant women showed that the detection rate of existing screening strategies for T21 can reach about 88.4%, with a screen-positive rate of 3.3% when applying the integrated prenatal screening procedure (Okun 2008).

A woman is classified as screen-positive if her risk is equal to or exceeds a predetermined threshold following prenatal screening result or due to some other factors such as personal or familial history of aneuploidies or translocations. Although these factors are considered to significantly increase the risk of fetal aneuploidy, the indications for invasive testing may vary between countries. To confirm the presence or absence of fetal aneuploidy in these high-risk pregnant women, a diagnostic


test involving karyotyping by an invasive procedure such as amniocentesis or chorionic villi sampling (CVS) is offered (ACOG #88 2007; Benn 2011; Chitayat 2011). Karyotyping by traditional banding techniques of fetal cells obtained from amniotic fluid or placental tissue has been considered the standard of care for prenatal diagnosis of aneuploidies (ACOG #545 2012; Benn 2013a; ICFMM 2013). Fluorescence in situ hybridisation (FISH) and quantitative fluorescence polymerase chain reaction (QF-PCR) are appropriate standards of care for pregnant women at increased risk of common fetal aneuploidies based on screening results (Duncan 2011; Langlois 2011; South 2013). Microarray analysis by array comparative genomic hybridisation (aCGH) is recommended in pregnancies with fetal anomalies and it is increasingly replacing karyotyping (ACOG #682 2016).

Five reviews published in the Cochrane Library examined serum, urine, ultrasound or a combination of these tests for T21 screening. For first-trimester serum tests (Allred 2015a), the authors concluded that two markers in combination with maternal age, specifically pregnancy associated plasma protein A (PAPP-A) and free human chorionic gonadotropin (hCG) are significantly better than those involving single markers combined with or without maternal age. For second-trimester serum tests (Allred 2012), the authors concluded that two or more markers, with or without inhibin A, in combination with maternal age are significantly more sensitive than one marker alone. Their review also showed that no test combination was superior to the others and therefore it was not possible to recommend a specific test combination. For first-trimester ultrasound tests alone of in combination with first-trimester serum tests (Allred 2017a), the authors concluded that test strategies that combine ultrasound markers with serum markers, especially PAPP-A and free β hCG, and maternal age were significantly better than those involving only ultrasound markers (with or without maternal age) except nasal bone. For first- and second-trimester serum tests with and without first-trimester ultrasound tests (Allred 2017b), the authors concluded that tests involving first-trimester ultrasound with first- and second-trimester serum markers in combination with maternal age are significantly better than those without ultrasound, or those evaluating first-trimester ultrasound in combination with second-trimester serum markers, without first-trimester serum markers. For first- and second-trimester urine tests (Allred 2015b), the authors concluded that second-trimester β -core fragment and oestriol with maternal age are significantly more sensitive than the single marker second-trimester β -core fragment and maternal age. However, there were few studies and the evidence does not support the use of urine tests for T21 screening for the first 24 weeks of pregnancy.

Role of index test(s)

Genomics-based non-invasive prenatal testing such as MPSS or TMPS could be offered to pregnant women after a first-tier screening and before a diagnostic test in order to better identify which pregnant women at increased risk of fetal aneuploidy should be offered further testing (triage) (Figure 2). The use of such NGS-based approaches has also been suggested as a replacement for current first-tier screening tests (biochemical, ultrasound or both) or as potential diagnostic tests to replace current diagnostic test (karyotyping of fetal cells from amniocentesis or CVS) (Bianchi 2012).

Figure 2. Current clinical pathway and three proposed uses of genomics-based non-invasive prenatal testing (gNIPT). Currently (on the left), pregnant women can have a prenatal screening test consisting of biomarkers or ultrasound, or both. For high-risk pregnant women, an invasive diagnostic test (karyotyping) is offered. In the present review, we propose 3 different clinical pathways. First, gNIPT could be used as a triage test, to decide which pregnant women should receive further testing. Second, gNIPT could be used to replace current prenatal screening tests. Finally, gNIPT could be used to replace current invasive diagnostic tests (if diagnostic performance permits). At any point in a clinical pathway, a pregnant woman may decide not to proceed with other tests (not shown in the figure). Figure was designed by CL, JB, MB and YT.



^aPrenatal screening test consists of serum biomarkers or ultrasound measurements or both offer to all pregnant women.

^bKaryotyping from amniocentesis or chorionic villus sampling.

Rationale

Current screening tests (biochemical, ultrasound or both) have relatively high false positive rates, which may result in undue anxiety for many pregnant women who will be offered an invasive diagnostic procedure. For example, at a prenatal screening risk cut-off of 1:300, fetal aneuploidy is confirmed by karyotyping in only about 1/34 to 1/14 (3% to 7%) screen-positive cases (Renshaw 2013; Wald 2005). As a result, many more women will undergo invasive diagnostic testing following positive screening tests than the number carrying a fetus with aneuploidy. In France, each year, about 800,000 pregnant women opt for prenatal T21 biochemical screening, ultrasound measurements or both, and about 24,000 of them (3%) will have karyotype testing (Basset 2013). Invasive testing methods for prenatal diagnosis of aneuploidy identify pregnancies with fetal chromosomal abnormalities, but contribute to an additional procedure-related fetal loss rate (Wilson 2007).

A recent meta-analysis showed that weighted pooled procedure-related risks of miscarriage of invasive testing methods before 24 weeks' gestation were 0.11% for amniocentesis and 0.22% for CVS (Akolekar 2015). The risk of miscarriage of normal fetuses associated with such invasive procedures has fostered the development of alternative screening and diagnostic approaches.

The discovery of fetal circulating cells and fetal cfDNA in maternal blood during pregnancy has enabled the development of non-invasive methods to analyse the fetal genome (Birch 2005; Lo 1997; Wright 2009). Fetal DNA offers advantages over circulating fetal cells because it is more easily extracted from maternal plasma samples and it disappears within hours after birth (undetectable about one to two days postpartum), as compared to the paucity and persistence of fetal cells in maternal blood over several consecutive pregnancies (up to 27 years) (Wright 2009; Yu 2013). At present, the analysis of cfDNA by NGS technologies seems to be the most

promising alternative gNIPT approach for the detection of fetal aneuploidies from maternal blood. This allows sequencing of tens of millions of these DNA fragments simultaneously, paving the way for the development of a non-invasive, less psychologically stressful method potentially able of detecting fetal aneuploidies earlier and with better accuracy than current screening programs. As such, NGS technologies have the potential to radically change prenatal screening for fetal aneuploidy. Indeed, a study exploring the impact of gNIPT on prenatal care showed that more pregnant women with positive first-trimester screening opt for further testing (from 47.2% to 78.8%) than before the introduction of gNIPT, while the rate of invasive diagnostic testing has decreased significantly (from 47.2% to 39.2%). Additionally, fewer pregnant women declined follow-up testing when gNIPT was an option (from 52.8% to 21.2%) (Chetty 2013). Another study suggested that gNIPT could reduce procedure-related fetal losses in high-risk women by up to 88% (O'Leary 2013).

For instance, the new gNIPT approach is reported to detect aneuploidy with high sensitivity to select a subset of pregnant women for an invasive diagnostic procedure and could be performed in high-risk pregnant women (as a second-tier test) following a positive screening result (Benn 2013a). The major expected advantage of gNIPT by NGS over current (biochemical, ultrasound or both) screening tests is the significant decrease in false positive results and thus the reduction of invasive procedures and their associated normal fetus losses. Also, it was reported that a reduction of invasive prenatal procedures with the introduction gNIPT has indeed been documented (Chetty 2013; Larion 2014; Tiller 2014). Assessment of how NGS should be used in clinical practice for aneuploidy detection is currently being studied. NGS approaches could also be performed in general obstetrical population (as first-tier test), in place of current screening algorithms (biochemical, ultrasound or both) (Figure 2). However, the field is moving rapidly. From January to July 2014, around 60 NIPT studies were published in PubMed compared to 70 studies in 2013 and 40 studies in 2012.

Up to now, no comprehensive systematic review including meta-analyses has analysed and compared the diagnostic accuracy of MPSS and TMPS methods for the detection of fetal aneuploidies, either as a second-tier test (i.e. in women at increased risk of fetal aneuploidy after current screening procedures) or as a first-tier test (i.e. in all pregnant women). Benn 2013b published a review on gNIPT focused on providing the information needed by clinicians and public health providers before implementation of this technology in routine clinical practice. However, their review included only T21 and T18. Mersy 2013 published a systematic review on quality and outcome of diagnostic test accuracy studies on non-invasive detection of fetal T21 only. One updated meta-analysis (Gil 2015a) pooled all gNIPT methods but did not assess the relative performance of MPSS and TMPS technologies separately. More recently, Taylor-Phillips 2016 published a meta-analysis on gNIPT accuracy for major autosomal anomalies (T21, T18 and T13) without sex chromosome aneuploidies (SCAs) assessment and using restrictive inclusion criteria for included publications (e.g. limited to the English language, cohorts of more than 50 pregnant women) and including studies with incomplete follow-up (pregnant women without reference standard). In the meta-analysis of Mackie 2017, multifetal pregnancies and case-control study design were excluded. In the meta-analysis published by the Haute Autorité de Santé in France (HAS 2015), the accuracy of gNIPT was evaluated

for T21 only and included studies with pregnant women selected at high risk of fetal aneuploidy as well as studies with pregnant women unselected for their risk (general population). Only studies published in English were included. The review of Agarwal 2013 described the properties of commercial tests available (e.g. type of gNIPT method, costs, turnaround times and reimbursement), intellectual property, commercialisation, patenting, patenting litigation and licensing landscape of technologies underlying these tests.

Genomics-based non-invasive prenatal tests are already advertised and marketed to North-American, European and Asian healthcare providers. Leading companies are summarised in Table 2. Other entities are trying to make their way into the market (Birmingham Women's NHS; Counsyl; GENDIA; Genesis Genetics; Integrated Genetics; NIPD Genetics; Progenity; Quest Diagnostics; RAVGEN; Xcelom). Some of these assays have yet to be approved by the US Food and Drug Administration. There is significant pressure for increasing their use in clinical practice, but comparative effectiveness and cost-effectiveness studies, as well as studies of the ethical, legal and social issues are scarce. Furthermore, tools needed for their patient value-based implementation are not available or have not been validated.

OBJECTIVES

To evaluate and compare the diagnostic accuracy of massively parallel shotgun sequencing (MPSS) and targeted massively parallel sequencing (TMPS) using circulating cell-free DNA (ccfDNA) in maternal blood for the detection of common fetal aneuploidies (T21, T18, T13, 45,X, 47,XXY, 47,XXX and 47,XYY) according to their prior risk of fetal aneuploidy. The genomics-based non-invasive prenatal testing (gNIPT) results were confirmed by a reference standard such as fetal karyotype or neonatal clinical examination.

To evaluate the screening performance of MPSS and TMPS as triage tests (a second-tier screening test) for identifying which pregnant women at increased risk of fetal aneuploidy should be offered further testing, that is, after a first-tier screening, but before a diagnostic test.

To assess the screening performance of MPSS and TMPS as a first-tier test in pregnant women without prior risk (i.e. in unselected pregnant women or general population) as a replacement for current offered first-tier tests (biochemical, ultrasound or both).

To assess the diagnostic performance of MPSS and TMPS as a second-tier test as potential diagnostic tests to replace current invasive diagnostic tests.

Secondary objectives

To investigate potential sources of heterogeneity that may influence the diagnostic accuracy of MPSS and TMPS such as gestational age at the time of blood collection and type of reference standard used.

METHODS

Criteria for considering studies for this review

Types of studies

We included studies that met the following inclusion criteria:

- randomised studies where pregnant women were randomised to receive one gNIPT (MPSS or TMPS) as well as the reference standard;
- retrospective and prospective cohort studies where all pregnant women were tested with one or more gNIPT methods and the reference standard (including head-to-head studies); and
- retrospective and prospective case-control studies comparing one or more of the gNIPT methods with the reference standard.

Although studies with a retrospective or case-control design are prone to biases, we included such studies because we anticipated a paucity of other study designs. When data were sufficient, we explored the effect of excluding case-control studies in sensitivity analyses.

We excluded studies for which it was not possible to extract or derive the number of true positives, false positives, false negatives and true negatives.

Participants

We included women of any age, ethnicity and gestational age with a singleton or multifetal (monochorionic and dichorionic) pregnancy.

Index tests

Genomics-based non-invasive prenatal tests based on plasma ccfDNA in maternal blood, analysis by either MPSS or TMPS methods.

Target conditions

We considered seven fetal aneuploidies, namely T21, T18, T13, 45,X, 47,XXY, 47,XXX and 47,XYY.

Reference standards

We considered the following test as reference standard: fetal karyotyping performed on cells obtained from chorionic villi sampling (CVS), amniotic fluid, placental tissue, a fetus lost by miscarriage or other equivalent and recognised methods on the same materials. By "fetal karyotyping" we mean traditional banding techniques, spectral karyotyping, fluorescence in situ hybridisation (FISH), array comparative genomic hybridisation (aCGH) or quantitative fluorescence polymerase chain reaction (QF-PCR). If fetal karyotyping was not performed, we used neonatal clinical examination or medical records from birth as a secondary reference standard for T21, T18 or T13. For sex chromosome aneuploidies (SCA), only fetal karyotype was an appropriate reference standard because newborns usually have a normal phenotype.

Search methods for identification of studies

Electronic searches

We used a sensitive search strategy that included the following three sets of search terms and synonyms:

- index test (e.g. cell-free DNA, sequencing, non-invasive and genetic diagnosis);
- participants' description (e.g. pregnant women, fetus and prenatal); and
- target condition (e.g. aneuploidy and chromosome anomalies).

We combined free-text words and subject headings used within each set with the Boolean operator OR and then combined the three sets using AND. We reviewed publications from 1st January 2007 because MPSS and TMPS were introduced in the literature in 2008 (Chiu 2008; Fan 2008). We did not limit our search by language, search filter or publication type (e.g. journal article, clinical trial, validation study, review and comment).

We applied a comparable search strategy (Appendix 4) with adaptations for each of the following databases:

- MEDLINE (Ovid) (January 2007 to July 2016);
- Embase (January 2007 to July 2016);
- Web of Science (ISI) (January 2007 to July 2016);
- Cochrane Register of Diagnostic Test Accuracy Studies, *Cochrane Library* (January 2007 to October 2016);
- ClinicalTrials.gov (January 2007 to September 2016);
- European Clinical Trials Register (January 2007 to September 2016);
- WHO ICTRP (January 2007 to September 2016);
- The National Technical Information Service (NTIS) (January 2007 to September 2016);
- OpenGrey (January 2007 to October 2016); and
- National Guideline Clearing House (January 2007 to September 2016).

Searching other resources

We examined references cited in potentially relevant full-text papers and those cited in previous reviews by cross-checking bibliographies. We examined grey literature by searching data available on the websites of private prenatal diagnosis companies (Ariosa Diagnostics 2016; BGI 2016; Berry Genomics 2016; Genoma 2016; Genome Care 2016; Illumina 2016; LabGenomics 2016; LifeCodexx 2016; Natera 2016; Genesupport 2016; Premaithe Health plc 2016; Sequenom 2016) using gNIPT technologies (January 2007 to December 2016). We also searched for conference abstracts and theses in appropriate sources (e.g. TheseNet, Theses Canada Portal) (January 2007 to October 2016).

Data collection and analysis

We used the methods suggested by the Cochrane Diagnostic Test Accuracy Working Group (Deeks 2013). For selection of studies, data extraction and assessment of methodological quality, we conducted a pilot using 20 randomly selected articles to trial our forms in order to ensure criteria were applied consistently.

None of the review authors involved in conducting a gNIPT primary study (FL, FR, SL and YG) took part in the selection of studies, nor in any decisions/analyses related to their own studies. Furthermore, by the final date of data collection, these authors had not published a primary gNIPT study.

Selection of studies

Two review authors (MB and CL) independently identified relevant studies by screening the titles and abstracts of all studies identified by the search strategy. We obtained the full-text version of all potentially relevant studies and assessed them for inclusion by using a study eligibility table based on prespecified inclusion criteria. The data collection form (Excel[®] format) for classifying studies during the full-text assessment is presented in Appendix

5. We considered all comments, statements or errata related to included studies. We excluded studies that did not match the inclusion criteria and we recorded the reason(s) for exclusion. If results from the same study cohort were reported in multiple publications, we considered all the publications and included results from the most relevant and comprehensive publications. We excluded papers with preliminary results whose full published results were available. We resolved any disagreement between assessors (MB and CL) by iteration, discussion and consensus. If required, we consulted a third review author (JB or LN).

Data extraction and management

Two review authors (MB and CL, JB or LN) independently extracted information and data from each included study by using a data extraction form that we developed in Excel® format. We included the following items:

- study characteristics (e.g. reference details allowing identification of the publication, language and study design);
- population characteristics (e.g. gestational age, maternal age, ethnicity, total number of pregnant women, number of aneuploid cases, number of euploid cases, recruitment location (country, geographic locations or regions), recruitment period and other relevant tests carried out prior to index test (e.g. ultrasonography, biochemical screening));
- features of the reference standard (e.g. fetal karyotyping, chromosome analysis or clinical examination);
- features of the index test (e.g. technical details, commercial or in-house gNIPT, cutpoint, failure rate, blood sample collection time (before or after reference standard) and first-tier test or second-tier test); and
- data for constructing two-by-two tables (number of true positives, false positives, false negatives and true negatives) or summary statistics from which the data were derived. In the two-by-two tables, the true negative cases were patients with any other aneuploidy than the one under analysis and all euploid cases were considered unaffected. When data were presented in three-by-two tables due to unclassified index test results (defined as grey zone between positive and negative test results), we constructed two-by-two tables by considering all unclassified gNIPT results as test positives. This is because in practice such results will lead to further testing and investigation to ensure a case of fetal aneuploidy is not missed.

We cross-checked all extracted and recorded data and we resolved any disagreement by iteration, discussion and consensus between two review authors (MB and CL, JB or LN). If required, we consulted a third author (JB, LN or CL). We wrote to the study contact author if information was missing or unclear or to clarify potential overlap between publications based on the same dataset to avoid including the same women more than once. If an article presented results including other aneuploidies than the ones under review, we considered only the subset of the cohort with the aneuploidies of interest.

Assessment of methodological quality

We used the revised QUality Assessment of Diagnostic Accuracy Studies (QUADAS-2) tool for assessment of methodological quality of included studies (Whiting 2011). We tailored the tool to this review question using the operational criteria detailed in Appendix 6 to answer signalling questions and make the overall judgment

of risk of bias and applicability concerns for each domain of the tool. We answered each signalling question with a 'yes', 'no' or 'unclear' response for each included study and we recorded the reason for the judgment made. If a study was recorded as 'yes' on all signalling questions related to risk of bias, then it was deemed appropriate to have an overall judgment of 'low risk of bias'. If a study is recorded 'no' or 'unclear' on one or more signalling questions in a domain, then it was judged as having 'high or unclear risk of bias'. Judgments about applicability concern were rated as 'low', 'high' or 'unclear' in relation to our review question. 'Unclear concern' was used only if insufficient information was available. Two review authors (MB and CL, JB or LN) independently applied the QUADAS-2 tool to each included study and we resolved any disagreement by iteration, discussion and consensus. If required, we consulted a third review author (JB, LN or CL).

Statistical analysis and data synthesis

The unit of analysis was the pregnant woman irrespective of the type of pregnancy (multifetal or singleton pregnancy). We evaluated the performance of MPSS and TMPS for the detection of each type of aneuploidy under study both individually and globally for any type of aneuploidy (all autosomal aneuploidies combined and all sex chromosomal aneuploidies combined). We distinguished between each of the following groups of pregnant women and performed separate analyses for each subgroup:

- unselected pregnant women undergoing aneuploidy screening (first-tier gNIPT, i.e. offered to all pregnant women) and women selected at high risk of fetal aneuploidy (second-tier gNIPT);
- women with singleton and multifetal pregnancy because ccfDNA's fetal fraction in multifetal pregnancy is higher than in singleton pregnancy (Attilakos 2011; Canick 2012); and
- pregnant women who underwent gNIPT during the first trimester (15 weeks or less), the first or second trimester (29 weeks or less) or at any time during pregnancy (42 weeks or less).

For each gNIPT method, we used Review Manager® to produce coupled forest plots of sensitivity and specificity, together with their 95% confidence intervals (CIs). We also plotted study-specific estimates of sensitivity and specificity in receiver operating characteristic (ROC) space. All gNIPTs are laboratory-developed tests based on differently calibrated assays with specific cutpoints to classify samples as euploid or aneuploid. There is no consensus on the cutpoints to use in practice. For this reason, we had planned to use a modelling strategy that focuses on the estimation of summary ROC curves (Macaskill 2010; Rutter 2001) and to estimate summary points (summary sensitivity and specificity) if a sufficient number of studies reported common cutpoints. However, given the qualitative nature of the cutpoints, which is highly dependent on each laboratory's developed gNIPT and study populations, it was not possible to identify a common cutpoint. Therefore, we reasoned that this was a special case where we can assume gNIPT results were binary (positive or negative). The rationale was further strengthened by the lack of apparent threshold effect when we examined the studies in ROC space. If a study reported more than one cutpoint, we considered all cutpoints and chose one cutpoint, the most commonly reported across all studies, such that only one pair of sensitivity and specificity from a study was included in meta-analysis.

Due to limited or absence of threshold effect, there was no requirement to account for correlation between sensitivity

and specificity across studies in meta-analysis. Therefore, we removed the correlation parameter from the bivariate model (Chu 2006), thus simplifying the model to two univariate random-effects logistic regression models for separate meta-analyses of sensitivities and specificities (Takwoingi 2015). In cases where there were few studies in the meta-analysis or a random-effects analysis failed to converge, we used fixed-effect logistic regression models. Where all studies in the meta-analysis reported 100% sensitivity or 100% specificity, these fixed-effect models fail as the prediction is perfect. Therefore, in such situations we used simple pooling by summing up the numbers of true positives and total cases to compute sensitivity, and the numbers of true negatives and unaffected pregnancies to compute specificity. CIs were obtained using the Wilson method (Newcombe 1998).

We compared the diagnostic accuracy of MPSS and TMPS by first using all available data (indirect comparison). If studies that compared MPSS and TMPS in the same population (head-to-head or direct comparison) were available, we had planned a second set of analyses restricted to direct comparisons. Comparative meta-analyses were done by adding a covariate for test type to random-effects or fixed-effect models. We used likelihood ratio tests to assess the statistical significance of differences between tests by comparing models that included covariate terms for test type with models that did not include the terms. If data were available, comparisons between gNIPTs and traditional screening approaches were planned using a similar strategy to that described above. Meta-analyses were performed using the `xmlogit` and `blogit` functions in the Stata software package (version 13; StataCorp, College Station, Texas 77845, USA). When meta-analyses of direct comparisons were not possible, we examined individual study results. For each comparative study, we computed differences in sensitivity and specificity, and 95% CIs were calculated for the differences using the Newcombe-Wilson method without continuity correction (Newcombe 1998).

Investigations of heterogeneity

We examined forest plots of sensitivity and specificity and summary ROC plots for each gNIPT method to visually assess heterogeneity. If sufficient data were available for meta-regression (by adding a covariate to a logistic regression model to explore its effect on sensitivity and specificity), we had planned to investigate the effect of the following:

- study population (e.g. ethnicity, gestational age at blood collection); and
- type of reference standard (i.e. karyotype or mixed reference standard).

However, formal investigations using meta-regression were not possible due to limited data and little or no heterogeneity in test accuracy.

Sensitivity analyses

We performed sensitivity analyses to assess the effect of excluding case-control studies and studies with a small number of cases of aneuploidy (less than 10 cases) on the summary estimates of test accuracy.

We had planned to also assess the effect of:

- studies where pregnant women received an invasive diagnostic test less than one day before blood collection for gNIPT;
- third trimester gestational age at the moment of blood collection for gNIPT;
- studies available only as abstracts; and
- studies at 'high or unclear risk of bias' according to the QUADAS-2 assessment tool.

However, due to lack of data or lack of variability in estimates of sensitivity and specificity, only assessments of the impact of study design and number of cases were performed.

RESULTS

Results of the search

We found a total of 11,912 articles through our electronic searches from January 2007 to October 2016 (see PRISMA study flow diagram in Figure 3). A total of 11,700 articles were identified through databases (941 through MEDLINE, 8381 through Embase, 1986 through Web of Science, 18 through Cochrane Diagnostic Test Accuracy register of studies, 245 through ClinicalTrials.gov, 43 through European Clinical Trials Register, 21 through WHO ICTRP, 34 through NTIS, 19 through OpenGrey and 12 through the National Guideline Clearing House). We found 212 publications through other sources (two articles received from the author, 175 from gNIPT company's website, 27 from TheseNet and eight from These Canada Portal). After removing 2354 duplicates, two review authors independently screened the titles and abstracts of 9558 publications. Of the 9558 publications, 9209 were deemed irrelevant to our review question. We retrieved the full texts of the remaining 349 articles to assess their eligibility. After resolving disagreement between two or three review authors, 261 articles were excluded (see [Characteristics of excluded studies](#) for details) and 63 articles fulfilled our inclusion criteria (see [Characteristics of included studies](#) for details). Among these 63 articles, 62 were journal articles and one was a letter to the editor with sufficient information to be included (Jackson 2014). From the 63 articles, two articles presented two studies (two different cohort, two 2x2 tables). At all, we included 65 studies of 86,139 pregnant women (3141 aneuploids and 82,998 euploids). No studies are awaiting classification. We identified 25 ongoing trials through clinical trials databases (see [Characteristics of ongoing studies](#) for details). We will consider these trials in future updates.

Figure 3. PRISMA flow diagram for selection of studies from January 2007 to October 2016. #: number, DTA: diagnostic test accuracy, NTIS: The National Technical Information Service and WHO ICTRP: World Health Organization International Clinical Trials Registry Platform.





Figure 3. (Continued)



Basic features of the included studies

The clinical characteristics of pregnant women and sequencing method were generally well described or referenced. Some studies did not clearly report how patient selection was done and which inclusion and exclusion criteria were applied. Patients’ enrolment flow-charts, pregnancy outcome flow-chart and 2 x 2 tables were unclear in many studies. We therefore contacted study authors to clarify unclear information, to obtain missing data or to clarify potential overlap of patients between publications.

We described the characteristics of included studies in [Characteristics of included studies](#) table and provided a summary in [Table 4](#). Forty-two studies (65%) enrolled pregnant women selected at high risk of fetal aneuploidy ([Alberti 2015](#); [Ashoor 2012](#); [Benachi 2015](#); [Bianchi 2012](#); [Bianchi 2013](#); [Bijok 2014](#); [Canick 2012](#); [Chen 2011](#); [Ehrich 2011](#); [Hall 2014](#); [Hooks 2014](#); [Hou 2012](#); [Huang 2014](#); [Jeon 2014](#); [Jiang 2012](#); [Johansen 2016](#); [Ke 2015](#); [Kim 2016](#); [Lee 2015](#); [Lefkowitz 2016](#); [Liang 2013](#); [Liu 2012](#); [Mazloom 2013](#); [Nicolaidis 2013](#); [Nicolaidis 2014a](#); [Norton 2012](#); [Palomaki 2012](#); [Papageorghiou 2016a](#); [Papageorghiou 2016b](#); [Persico 2016](#); [Poon 2016](#); [Porreco 2014](#); [Sehnert 2011](#); [Song 2015](#); [Sparks 2012a](#); [Stumm 2014](#); [Sukhikh 2015](#); [Sung-Hee 2015](#); [Verweij 2013](#); [Wang 2014](#); [Wang 2015a](#); [Zhang 2016](#)); five studies (8%) enrolled pregnant women without prior risk of fetal aneuploidy ([del Mar Gil 2014](#); [Nicolaidis 2012](#); [Norton 2015](#); [Quezada 2015](#); [Song 2013](#)); and 18 studies (28%) enrolled pregnant women from a mixed risk cohort of fetal aneuploidy ([Ashoor 2013](#); [Bevilacqua 2015](#); [Bianchi 2014a](#); [Chiu 2011](#); [Comas 2015](#); [Fiorentino 2016](#); [Gil 2016](#); [Jackson 2014](#); [Korostelev 2014](#); [Lau 2012](#); [Ma 2016](#); [Pergament 2014](#); [Samango-Sprouse 2013](#); [Shaw 2014](#); [Tynan 2016](#); [Yao 2014](#); [Zhou 2014a](#); [Zhou 2014b](#)). Mixed-risk samples included a mixture of selected pregnant women with low, high or no prior risk of fetal aneuploidy. Such samples do not represent the real-life situation (i.e. using gNIPT as a first-tier screening test or as a second-tier test) and so such studies were not used for addressing our research objectives. Nevertheless, as we did not pre-specify exclusion of such studies, we analysed the data and the results are presented in [Appendix 7](#).

The studies assessed MPSS and TMPS using various algorithms and cutpoints. [Table 4](#) describes the specific gNIPT assay used in the included studies. Each assay was developed and validated by the testing laboratory. Among the 65 studies, 44 studies (68%) used a whole genome sequencing method (MPSS) ([Alberti 2015](#); [Benachi 2015](#); [Bianchi 2012](#); [Bianchi 2013](#); [Bianchi 2014a](#); [Bijok 2014](#); [Canick 2012](#); [Chen 2011](#); [Chiu 2011](#); [Ehrich 2011](#); [Fiorentino 2016](#); [Hou 2012](#); [Huang 2014](#); [Jeon 2014](#); [Jiang 2012](#); [Johansen 2016](#); [Ke 2015](#); [Kim 2016](#); [Lau 2012](#); [Lee 2015](#); [Lefkowitz 2016](#); [Liang 2013](#); [Liu 2012](#); [Ma 2016](#); [Mazloom 2013](#); [Palomaki 2012](#); [Papageorghiou 2016a](#); [Papageorghiou 2016b](#); [Poon 2016](#); [Porreco 2014](#); [Sehnert 2011](#); [Shaw 2014](#); [Song 2013](#); [Song 2015](#); [Stumm 2014](#); [Sukhikh 2015](#); [Sung-Hee 2015](#); [Tynan 2016](#); [Wang 2014](#); [Wang 2015a](#); [Yao](#)

[2014](#); [Zhang 2016](#); [Zhou 2014a](#); [Zhou 2014b](#)), and 21 (32%) used a targeted method (TMPS) ([Ashoor 2012](#); [Ashoor 2013](#); [Bevilacqua 2015](#); [Comas 2015](#); [del Mar Gil 2014](#); [Gil 2016](#); [Hall 2014](#); [Hooks 2014](#); [Jackson 2014](#); [Korostelev 2014](#); [Nicolaidis 2012](#); [Nicolaidis 2013](#); [Nicolaidis 2014a](#); [Norton 2012](#); [Norton 2015](#); [Pergament 2014](#); [Persico 2016](#); [Quezada 2015](#); [Samango-Sprouse 2013](#); [Sparks 2012a](#); [Verweij 2013](#)). Of the 65 studies, five studies compared gNIPT with traditional screening tests ([Bianchi 2014a](#); [Nicolaidis 2012](#); [Norton 2015](#); [Quezada 2015](#); [Song 2013](#)). MPSS studies involved 50,864 pregnant women, TMPS studies involved 35,275 pregnant women and traditional screening tests involved 24,279 pregnant women. The most commonly (15 studies) used cutpoint for gNIPT assays was a chromosomal ratio Z score of 3. Thirteen studies used the FORTE risk score, eight studies used a normalised chromosome value (NCV) and 13 studies did not report their cutpoint. The remaining studies used other cutpoints ([Table 4](#)). Timing of blood sampling for gNIPT was before invasive procedure in 55 studies, before or more than 24 hours after invasive sampling in four studies ([Ashoor 2013](#); [Lefkowitz 2016](#); [Pergament 2014](#); [Samango-Sprouse 2013](#)), and was not reported in six studies ([Bevilacqua 2015](#); [Jiang 2012](#); [Song 2013](#); [Sparks 2012a](#); [Wang 2014](#); [Zhang 2016](#)).

Among all aneuploidies considered, 36 studies (55%) reported analyses only for autosomes, four (6%) for only sex chromosome aneuploidies (SCA) and 25 studies (39%) for both autosomes and SCA. Fifty-seven studies (82,620 pregnant women) evaluated T21, 50 studies (79,322 pregnant women) evaluated T18, 39 studies (68,958 pregnant women) evaluated T13, 20 studies (10,081 pregnant women) evaluated 45,X, seven studies (6035 pregnant women) evaluated 47,XXX, 12 studies (7609 pregnant women) evaluated 47,XXY and 10 studies (6987 pregnant women) evaluated 47,XYY ([Table 4](#)). Among all 65 included studies, there are a total of 2004 T21 cases, 634 T18 cases, 215 T13 cases, 232 45,X cases, 14 47,XXX cases, 25 47,XXY cases and 16 47,XYY cases. All 65 studies used an appropriate reference standard such as fetal or neonatal karyotype, genetic testing, neonatal clinical examination or medical records from birth. In 36 studies (55%), only one reference standard was used while 29 studies (45%) used more than one reference standard ([Table 4](#)).

Among the 65 studies, 40 (62%) studies were prospective cohort studies ([Ashoor 2013](#); [Bevilacqua 2015](#); [Bianchi 2014a](#); [Bijok 2014](#); [Comas 2015](#); [Fiorentino 2016](#); [Gil 2016](#); [Hou 2012](#); [Huang 2014](#); [Jackson 2014](#); [Jeon 2014](#); [Jiang 2012](#); [Johansen 2016](#); [Ke 2015](#); [Kim 2016](#); [Korostelev 2014](#); [Lau 2012](#); [Lee 2015](#); [Liang 2013](#); [Liu 2012](#); [Mazloom 2013](#); [Nicolaidis 2013](#); [Norton 2012](#); [Norton 2015](#); [Pergament 2014](#); [Persico 2016](#); [Porreco 2014](#); [Quezada 2015](#); [Samango-Sprouse 2013](#); [Shaw 2014](#); [Song 2013](#); [Song 2015](#); [Stumm 2014](#); [Sukhikh 2015](#); [Verweij 2013](#); [Wang 2014](#); [Wang 2015a](#); [Zhang 2016](#); [Zhou 2014a](#); [Zhou 2014b](#)), eight (12%) studies were retrospective cohort studies ([Benachi 2015](#); [Bianchi 2013](#); [del Mar Gil 2014](#); [Nicolaidis 2012](#); [Sehnert 2011](#); [Sung-Hee 2015](#); [Tynan](#)

2016; Yao 2014), one (1%) study was a prospective and retrospective cohort study (Ma 2016) and 16 (25%) studies used a case-control design (Alberti 2015; Ashoor 2012; Bianchi 2012; Canick 2012; Chen 2011; Chiu 2011; Ehrich 2011; Hall 2014; Hooks 2014; Lefkowitz 2016; Nicolaidis 2014a; Palomaki 2012; Papageorghiou 2016a; Papageorghiou 2016b; Poon 2016; Sparks 2012a) (Table 4).

Forty-eight (74%) studies included only singleton pregnancies, while five (8%) studies included only multifetal pregnancies. Four (6%) studies included women with either type of pregnancy and eight (12%) studies did not report the type of pregnancy. Ten (15%) studies included only pregnant women in the first trimester (15 weeks or less), 21 (33%) studies included pregnant women in the first two trimesters (29 weeks or less), 24 studies (37%) included pregnant women in the three trimesters (42 weeks or less) and 10 studies (15%) did not report gestational age. Eighteen studies (28%) had more than 50% Caucasian women in their cohort, 21 studies (32%) had more than 50% Asian women and 26 studies (40%) did not report ethnicity.

Thirty-seven studies (57%) were industry-funded or were written by one or more author affiliated with a company who sells gNIPT (Benachi 2015; Bianchi 2012; Bianchi 2013; Bianchi 2014a; Canick 2012; Chen 2011; Chiu 2011; Ehrich 2011; Hall 2014; Hooks 2014; Huang 2014; Jackson 2014; Jiang 2012; Kim 2016; Lau 2012; Lee 2015; Lefkowitz 2016; Ma 2016; Mazloom 2013; Nicolaidis 2012; Nicolaidis 2013; Norton 2012; Norton 2015; Palomaki 2012; Papageorghiou 2016a; Papageorghiou 2016b; Pergament 2014; Persico 2016; Porreco 2014; Samango-Sprouse 2013; Sehnert 2011; Shaw 2014; Sparks 2012a; Stumm 2014; Tynan 2016; Verweij 2013; Yao 2014); 22 studies (34%) were not reported to be funded by industry but samples were sequenced and analysed by a commercial laboratory (Ashoor 2012; Ashoor 2013; Bevilacqua 2015; Bijok 2014; Comas 2015; del Mar Gil 2014; Fiorentino 2016; Gil 2016; Hou 2012; Jeon 2014; Ke 2015; Korostelev 2014; Liang 2013; Poon 2016; Quezada 2015; Song 2013; Song 2015; Sung-Hee 2015; Wang 2014; Wang 2015a; Zhou 2014a; Zhou 2014b); three studies (4.5%) had no link with industry (Alberti 2015; Johansen 2016; Sukhikh 2015); and the funding source was not reported for three studies (4.5%) (Liu 2012; Nicolaidis 2014a; Zhang 2016). Table 5 describes the specific gNIPT assay used in the included studies. Of the 65 studies, 61 (94%) used a commercial gNIPT (15 from Ariosa Diagnostics, Inc., 12 from Beijing Genomics Institute, four from Illumina (or Verinata Health), six from Natera, nine from Sequenom and 15 from other companies) (Table 5). It appears

that, for three of the commercially available assays, there are nine studies or more adding up to a large number of cases and unaffected cases analysed. Further, only two assays (one TMPS and one MPSS) were used in one of the five studies involving unselected pregnant women and one assay (Ariosa's Harmony™ test) was used in four of them. Twelve studies (19%) included their entire cohort in the analyses, 36 studies (55%) included between 80% to 99.9%, and 17 studies (26%) included less than 80%. We found 54 (83%) studies where patient exclusions and failed samples were reported (Table 6; Table 7).

Summary of excluded studies

We described the excluded studies in the PRISMA flow diagram (Figure 3) as well as in [Characteristics of excluded studies](#). After full-text assessment, we excluded 261 articles.

Of these 261:

- 93 (36%) studies were not diagnostic test accuracy studies (e.g. implementation study, simulation model, method development, proof-of-concept, method without sequencing approach);
- 55 (21%) studies had overlapping samples and were excluded to avoid double counting;
- 54 (21%) studies had incomplete 2 X 2 data or insufficient information to derive a 2 X 2 table;
- 22 (8%) studies had either an inappropriate or no reference standard;
- 8 (3%) studies were identified as reviews or Health Technology Assessment reports;
- 11 (4%) studies had target conditions, methods or sampling schemes other than those specified in our review; and
- 18 (7%) studies were news, letters, comments, notes, replies or editorials without new data.

The 25 ongoing studies are described in [Characteristics of ongoing studies](#).

Methodological quality of included studies

Figure 4 and Figure 5 show the risk of bias and applicability concerns for each included study for MPSS and TMPS, respectively. In Figure 6, the quality assessment results are summarised across all studies.

Figure 4. Risk of bias and applicability concerns summary: review authors' judgements about each domain for each of the studies included for massively parallel shotgun sequencing (MPSS).

	Risk of Bias				Applicability Concerns		
	Patient Selection	Index Test: MPSS	Reference Standard	Flow and Timing	Patient Selection	Index Test: MPSS	Reference Standard
Alberti 2015	-	+	+	-	+	+	+
Benachi 2015	-	+	+	-	+	+	+
Bianchi 2012	-	+	+	-	+	+	+
Bianchi 2013	-	+	+	+	+	+	+
Bianchi 2014a	-	+	+	-	-	+	+
Bijok 2014	?	+	+	-	+	+	+
Canick 2012	-	+	+	+	+	+	+
Chen 2011	-	?	+	+	+	+	+
Chiu 2011	-	+	?	-	-	+	+
Ehrich 2011	-	+	+	-	+	+	+
Fiorentino 2016	-	+	?	-	-	+	+
Hou 2012	-	+	+	+	+	+	+
Huang 2014	-	+	+	+	+	+	+
Jeon 2014	-	-	+	+	+	+	+
Jiang 2012	-	+	+	?	+	+	+
Johansen 2016	-	+	+	-	+	+	+
Ke 2015	-	+	?	+	+	+	+
Kim 2016	-	-	+	+	+	+	+
Lau 2012	-	+	+	+	-	+	+
Lee 2015	-	+	+	-	+	+	+
Lefkowitz 2016	-	+	+	-	+	+	+
Liang 2013	-	+	+	-	+	+	+
Liu 2012	?	+	+	+	+	+	+
Ma 2016	-	+	+	-	-	+	+
Mazloom 2013	-	+	+	-	+	+	+
Palomaki 2012	-	+	+	-	+	+	+

Figure 4. (Continued)

Palomaki 2012	⊖	⊕	⊕	⊖	⊕	⊕	⊕
Papageorghiou 2016a	⊖	⊕	⊕	⊖	⊕	⊕	⊕
Papageorghiou 2016b	⊖	⊕	⊕	⊖	⊕	⊕	⊕
Poon 2016	⊖	⊕	⊕	⊖	⊕	⊕	⊕
Porreco 2014	⊖	⊕	⊕	⊖	⊕	⊕	⊕
Sehnert 2011	⊖	⊕	⊕	⊕	⊕	⊕	⊕
Shaw 2014	⊖	⊕	⊕	⊕	⊖	⊕	⊕
Song 2013	⊖	⊕	⊕	⊖	⊕	⊕	⊕
Song 2015	⊖	⊕	⊕	⊖	⊕	⊕	⊕
Stumm 2014	⊖	⊖	⊕	⊖	⊕	⊕	⊕
Sukhikh 2015	⊖	?	?	⊕	⊕	⊕	⊕
Sung-Hee 2015	⊖	⊕	⊖	⊖	⊕	⊕	⊕
Tynan 2016	⊖	⊕	⊕	⊖	⊖	⊕	⊕
Wang 2014	⊖	⊕	?	?	⊕	⊕	⊕
Wang 2015a	⊖	⊕	?	⊕	⊕	⊕	⊕
Yao 2014	⊖	⊕	?	⊖	⊖	⊕	⊕
Zhang 2016	⊖	⊕	⊕	?	⊕	⊕	⊕
Zhou 2014a	⊖	⊕	?	⊖	⊖	⊕	⊕
Zhou 2014b	⊖	?	?	⊖	⊖	⊕	⊕


⊖ **High**
? **Unclear**
⊕ **Low**

Figure 5. Risk of bias and applicability concerns summary: review authors' judgements about each domain for each study included for targeted massively parallel sequencing (TMPS).

	Risk of Bias				Applicability Concerns		
	Patient Selection	Index Test: TMPS	Reference Standard	Flow and Timing	Patient Selection	Index Test: TMPS	Reference Standard
Ashoor 2012	-	+	+	-	+	+	+
Ashoor 2013	-	+	+	-	-	+	+
Bevilacqua 2015	-	+	?	-	-	+	+
Comas 2015	-	+	+	-	-	+	+
del Mar Gil 2014	?	+	+	-	+	+	+
Gil 2016	-	+	-	-	-	+	+
Hall 2014	-	?	?	-	+	+	+
Hooks 2014	-	?	+	-	+	+	+
Jackson 2014	?	+	+	-	-	+	+
Korostelev 2014	-	+	?	+	-	+	+
Nicolaides 2012	-	+	+	-	+	+	+
Nicolaides 2013	-	+	+	-	+	+	+
Nicolaides 2014a	-	+	+	-	+	+	+
Norton 2012	-	+	+	-	+	+	+
Norton 2015	-	+	+	-	+	+	+
Pergament 2014	-	+	+	-	-	+	+
Persico 2016	-	+	+	-	+	+	+
Quezada 2015	-	+	?	-	+	+	+
Samango-Sprouse 2013	-	+	+	-	-	+	+
Sparks 2012a	-	?	+	?	+	+	+
Verweij 2013	-	+	+	-	+	+	+

- High
 ? Unclear
 + Low

Figure 6. Risk of bias and applicability concerns (all tests included): review authors' judgements about each domain presented as percentages across included studies. MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing.



Risk of bias

No study was assessed as being at low risk of bias across all domains (Figure 4). For the patient selection domain, the 'Risk of bias' judgement was influenced mainly by inappropriate exclusions than the other signalling questions in this domain. Of the 61 studies judged to be at high risk of bias, 57 (93%) had inappropriate exclusions. The exclusions were mainly due to multifetal pregnancy, gestational age limits, and the prior risk of fetal aneuploidy. The remaining four (7%) studies were judged to be at unclear risk of bias (Figure 6).

In the index test domain, the risk of bias was considered to be low in 38 (58%) of the 44 MPSS studies and unclear in three (5%) studies. The remaining three (5%) MPSS studies were judged to be at high risk of bias because the index test was performed knowing the results of the reference standard or the threshold was not pre-specified. The risk of bias was low in 18 (27%) of the 21 TMPS studies. The remaining three (5%) TMPS studies were judged to be at unclear risk of bias. All five studies that assessed traditional screening approaches were judged to be at low risk of bias for the index test domain (Figure 6).

In the reference standard domain, all studies used a reference standard likely to correctly classify the target condition. We considered 50 (77%) studies to be at low risk of bias because the studies stated that the reference standard results were interpreted without knowledge of the results of the index test. Of the remaining 15 studies, two (3%) studies were at high risk of bias because the reference standard was performed knowing the results of the index test while it was unclear what was done in the other 13 (20%) studies (Figure 6).

For the flow and timing domain, 46 (71%) studies were considered to be at high risk of bias because some pregnant women were excluded from 2 x 2 tables because gNIPT failed during the sequencing process. Fifteen (23%) studies were judged to be at low risk of bias. For the remaining four (6%) studies, information about the appropriate interval between the index test and reference standard was not provided (Figure 6).

Applicability concerns

We judged all studies to be of low applicability concern in the index test and reference standard domains because the studies matched the review question (Figure 4; Figure 6). All studies used a gNIPT method with cfDNA in maternal blood and appropriate reference standard for the detection of common fetal aneuploidies. In the patient selection domain, 47 (72%) studies included cohort of pregnant women selected at high risk of fetal aneuploidy or cohort of unselected pregnant women and were judged to be of low applicability concern. In the other 18 (28%) studies, the cohorts comprised pregnant women with different prior risk of fetal aneuploidy (mixed risk cohorts). This population did not represent the real-life situation and those cohorts were judged to be of high applicability concern.

Findings

The characteristics of the studies are summarised in Table 4 and Summary of findings 1. Results are presented separately for each of the main fetal aneuploidies (T21, T18, T13 and 45,X) and globally for all autosomes or all sex chromosome aneuploidies (SCA) combined (Summary of findings 2; Summary of findings 3; Summary of findings 4; Summary of findings 5; Summary of findings 6; Summary of findings 7). For each aneuploidy, results are presented according to the prior risk of chromosomal abnormality as high risk or unselected population and according to MPSS and TMPS methods. Results from mixed-risk populations are summarised in Appendix 7. No study directly compared the accuracy of MPSS and TMPS. There were insufficient data to separately consider monochorionic and dichorionic pregnancies and four of the nine studies did not report chorionicity.

1. Trisomy 21 (T21 or Down syndrome)

A total of 57 studies assessed gNIPT for T21 in 2004 affected and 80,616 non T21 pregnancies. Five studies enrolled an unselected population of pregnant women undergoing aneuploidy screening, 36 studies enrolled pregnant women selected at high risk of fetal aneuploidy and 16 studies enrolled pregnant women with various prior risk and no a priori risk of fetal aneuploidy (mixed risk). Of the 57 studies, 41 assessed MPSS and 16 assessed TMPS. The results are summarised in Summary of findings 2.

a. Unselected population of pregnant women undergoing aneuploidy screening


Five cohort studies evaluated gNIPT in an unselected population of pregnant women undergoing aneuploidy screening. The studies

included 22,412 non T21 pregnancies and 96 (0.43%) T21 cases. MPSS was assessed in one study and TMPS was assessed in four studies (Figure 7).

Figure 7. Forest plot of MPSS and TMPS for T21 in unselected pregnant women undergoing aneuploidy screening. FN: false negative, FP: false positive, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, TN: true negative and TP: true positive.


MPSS T21

Study	TP	FP	FN	TN	Case-control	Sensitivity (95% CI)	Specificity (95% CI)
Song 2013	8	0	0	1733	No	1.00 [0.63, 1.00]	1.00 [1.00, 1.00]




MPSS T18

Study	TP	FP	FN	TN	Case-control	Sensitivity (95% CI)	Specificity (95% CI)
Song 2013	2	1	0	1738	No	1.00 [0.16, 1.00]	1.00 [1.00, 1.00]




MPSS T13

Study	TP	FP	FN	TN	Case-control	Sensitivity (95% CI)	Specificity (95% CI)
Song 2013	1	0	0	1740	No	1.00 [0.03, 1.00]	1.00 [1.00, 1.00]




TMPS T21

Study	TP	FP	FN	TN	Case-control	Sensitivity (95% CI)	Specificity (95% CI)
Norton 2015	38	9	0	15794	No	1.00 [0.91, 1.00]	1.00 [1.00, 1.00]
Quezada 2015	32	1	0	2752	No	1.00 [0.89, 1.00]	1.00 [1.00, 1.00]
del Mar Gil 2014	9	0	1	182	No	0.90 [0.55, 1.00]	1.00 [0.98, 1.00]
Nicolaidis 2012	8	0	0	1941	No	1.00 [0.63, 1.00]	1.00 [1.00, 1.00]




TMPS T18

Study	TP	FP	FN	TN	Case-control	Sensitivity (95% CI)	Specificity (95% CI)
Norton 2015	9	1	1	15830	No	0.90 [0.55, 1.00]	1.00 [1.00, 1.00]
Quezada 2015	9	5	1	2770	No	0.90 [0.55, 1.00]	1.00 [1.00, 1.00]
Nicolaidis 2012	2	2	0	1945	No	1.00 [0.16, 1.00]	1.00 [1.00, 1.00]



TMPS T13

Study	TP	FP	FN	TN	Case-control	Sensitivity (95% CI)	Specificity (95% CI)
Norton 2015	2	2	0	11181	No	1.00 [0.16, 1.00]	1.00 [1.00, 1.00]
Quezada 2015	2	2	3	2778	No	0.40 [0.05, 0.85]	1.00 [1.00, 1.00]
del Mar Gil 2014	1	0	0	191	No	1.00 [0.03, 1.00]	1.00 [0.98, 1.00]



i. MPSS

One prospective cohort study included eight T21 cases and 1733 non T21 pregnancies (Song 2013). The sensitivity (95% confidence interval (CI)) of MPSS was 100% (67.6% to 100%) and the specificity (95% CI) was 100% (99.8% to 100%).

ii. TMPS

TMPS was evaluated in four studies comprising 20,679 non T21 pregnancies and 88 T21 cases (del Mar Gil 2014; Nicolaidis 2012; Norton 2015; Quezada 2015). The summary sensitivity (95% CI) was 99.2% (78.2% to 100%) and the summary specificity (95% CI) was 100% (> 99.9% to 100%).

iii. Comparative accuracy of MPSS and TMPS


It was not possible to compare the accuracy of MPSS and TMPS in a meta-analysis because of limited data.

b. Selected population of pregnant women at high risk of fetal aneuploidy


Overall, 36 studies included pregnant women selected at high risk of fetal aneuploidy involving 20,317 non T21 pregnancies and 1294 (6.37%) T21 cases. MPSS was assessed in 30 studies and TMPS in six studies (Figure 8).

Figure 8. Forest plot of MPSS and TMPS for T21 in pregnant women selected at high risk of fetal aneuploidy. FN: false negative, FP: false positive, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, TN: true negative and TP: true positive.

MPSS T21



TMPS T21



i. MPSS

The 30 MPSS studies included 15,937 non T21 pregnancies and 1048 T21 cases (Alberti 2015; Benachi 2015; Bianchi 2012; Bianchi 2013; Canick 2012; Ehrich 2011; Hou 2012; Huang 2014; Jeon 2014; Jiang 2012; Johansen 2016; Ke 2015; Kim 2016; Lee 2015; Lefkowitz 2016; Liang 2013; Liu 2012; Palomaki 2012; Papageorghiou 2016a; Papageorghiou 2016b; Poon 2016; Porreco 2014; Sehnert 2011; Song 2015; Stumm 2014; Sukhikh 2015; Sung-Hee 2015; Wang 2014; Wang 2015a; Zhang 2016). The summary sensitivity (95% CI) was 99.7% (98.0% to 100%) and the summary specificity (95% CI) was 99.9% (99.8% to 100%).

ii. TMPS

Six studies evaluated TMPS in 4380 non T21 pregnancies and 246 T21 cases (Ashoor 2012; Nicolaides 2013; Norton 2012; Persico 2016; Sparks 2012a; Verweij 2013). The summary sensitivity (95% CI) was 99.2% (96.8% to 99.8%) and the summary specificity (95% CI) was 100% (99.8% to 100%).

iii. Comparative accuracy of MPSS and TMPS

An indirect comparison of the 30 MPSS and six TMPS studies showed no statistical evidence of a difference in sensitivity or specificity or both (P value = 0.52). The differences in sensitivity and specificity were negligible (Summary of findings 2).

2. Trisomy 18 (T18)

Fifty studies assessed T18 in 634 cases and 78,688 non T18 pregnancies. Four studies enrolled unselected population of pregnant women undergoing aneuploidy screening, 33 studies enrolled pregnant women selected at high risk of fetal aneuploidy and 13 studies enrolled a cohort with mixed prior risk. Of the 50 studies, 38 evaluated MPSS and 12 evaluated TMPS. The results are summarised in [Summary of findings 3](#).

a. Unselected population of pregnant women undergoing aneuploidy screening

Four studies, comprising 22,292 non T18 pregnancies and 24 (0.11%) T18 cases, assessed gNIPT for fetal aneuploidy in unselected pregnant women. One study assessed MPSS and three studies assessed TMPS ([Figure 7](#)).

i. MPSS

One MPSS study evaluated two T18 cases and 1739 non T18 pregnancies ([Song 2013](#)). The sensitivity (95% CI) was 100% (34.3% to 100%) and the specificity (95% CI) was 99.9% (99.7% to 100%).

ii. TMPS

Three studies evaluated TMPS in 20,553 non T18 pregnancies and 22 T18 cases ([Nicolaidis 2012](#); [Norton 2015](#); [Quezada 2015](#)). The summary sensitivity (95% CI) was 90.9% (70.0% to 97.7%) and the summary specificity (95% CI) was 100% (99.9% to 100%).

iii. Comparative accuracy of MPSS and TMPS


It was not possible to compare the accuracy of MPSS and TMPS in a meta-analysis because data were sparse.

b. Selected population of pregnant women at high risk of fetal aneuploidy


A total of 33 studies included pregnant women selected at high risk of fetal aneuploidy involving 444 (2.20%) T18 cases and 20,190 non T18 pregnancies. Of these, 28 studies assessed MPSS and five studies assessed TMPS ([Figure 9](#)).

Figure 9. Forest plot of MPSS and TMPS for T18 in pregnant women selected at high risk of fetal aneuploidy. FN: false negative, FP: false positive, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, TN: true negative and TP: true positive.

MPSS T18



TMPS T18



i. MPSS

Twenty-eight studies evaluated MPSS in 16,180 non T18 pregnancies and 332 T18 cases (Benachi 2015; Bianchi 2012; Bianchi 2013; Bijok 2014; Chen 2011; Hou 2012; Huang 2014; Jeon 2014; Jiang 2012; Johansen 2016; Ke 2015; Lee 2015; Lefkowitz 2016; Liang 2013; Liu 2012; Palomaki 2012; Papageorghiou 2016a; Papageorghiou 2016b; Poon 2016; Porreco 2014; Sehnert 2011; Song 2015; Stumm 2014; Sukhikh 2015; Sung-Hee 2015; Wang 2014; Wang 2015a; Zhang 2016). The summary sensitivity (95% CI) was 97.8% (92.5% to 99.4%) and the summary specificity (95% CI) was 99.9% (99.8% to 100%).

ii. TMPS

Five studies evaluated TMPS in 4010 non T18 pregnancies and 112 T18 cases (Ashoor 2012; Nicolaides 2013; Norton 2012; Persico 2016; Sparks 2012a). The summary sensitivity (95% CI) was 98.2% (93.1% to 99.6%) and the summary specificity (95% CI) was 100% (99.8% to 100%).

iii. Comparative accuracy of MPSS and TMPS

An indirect comparison of the 28 MPSS and five TMPS studies showed no statistical evidence of a difference in sensitivity, specificity or both (P value = 0.47). The differences in sensitivity and specificity were negligible (Summary of findings 3).

3. Trisomy 13 (T13)

T13 was assessed in 39 studies comprising 215 affected and 68,743 non T13 pregnancies. Four studies evaluated unselected population of pregnant women undergoing fetal aneuploidy screening, while 22 studies evaluated women at high risk of fetal aneuploidy and 13 studies evaluated mixed prior risk cohorts. Of the 39 studies, 29 assessed MPSS and 10 assessed TMPS. The results are summarised in [Summary of findings 4](#).

a. Unselected population of pregnant women undergoing aneuploidy screening

Four studies assessed gNIPT for T13 in unselected pregnant women. The studies included 15,894 non T13 pregnancies and nine (0.06%) T13 cases. Three studies evaluated TMPS and one study evaluated MPSS ([Figure 7](#)).

i. MPSS

One study evaluated MPSS in one T13 case and 1740 non T13 pregnancies ([Song 2013](#)). The sensitivity (95% CI) was 100% (20.7% to 100%) and the specificity (95% CI) was 100% (99.8% to 100%).

ii. TMPS

Three studies evaluated TMPS in 14,154 non T13 pregnancies and eight T13 cases ([del Mar Gil 2014](#); [Norton 2015](#); [Quezada 2015](#)). The summary sensitivity (95% CI) was 65.1% (9.2% to 97.2%) and the summary specificity (95% CI) was 100% (99.9% to 100%).

iii. Comparative accuracy of MPSS and TMPS


It was not possible to compare the accuracy of MPSS and TMPS in a meta-analysis because data were sparse.

b. Selected population of pregnant women at high risk of fetal aneuploidy


A total of 22 studies evaluated pregnant women selected at high risk of fetal aneuploidy. The studies included 14,103 non T13 pregnancies and 148 (1.05%) T13 cases. Twenty studies assessed MPSS and two studies assessed TMPS ([Figure 10](#)).

Figure 10. Forest plot of MPSS and TMPS for T13 in pregnant women selected at high risk of fetal aneuploidy. FN: false negative, FP: false positive, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, TN: true negative and TP: true positive.

MPSS T13



TMPS T13



i. MPSS

Twenty studies evaluated MPSS in 13,810 non T13 pregnancies and 128 T13 cases ([Benachi 2015](#); [Bianchi 2012](#); [Bianchi 2013](#);

[Canick 2012](#); [Chen 2011](#); [Jiang 2012](#); [Johansen 2016](#); [Ke 2015](#); [Lee 2015](#); [Lefkowitz 2016](#); [Liang 2013](#); [Liu 2012](#); [Palomaki 2012](#); [Papageorghiou 2016a](#); [Poon 2016](#); [Porreco 2014](#); [Sehnert 2011](#);

Song 2015; Stumm 2014; Sukhikh 2015). The summary sensitivity (95% CI) was 95.8% (86.1% to 98.9%) and the summary specificity (95% CI) was 99.8% (99.8% to 99.9%).

ii. TMPS

Two studies evaluated TMPS in 293 non T13 pregnancies and 20 T13 cases (Hall 2014; Persico 2016). The summary sensitivity (95% CI) was 100% (83.9% to 100%) and the summary specificity (95% CI) was 100% (98.7% to 100%).

iii. Comparative accuracy of MPSS and TMPS

It was not possible to compare the accuracy of MPSS and TMPS in a meta-analysis because data were sparse.

4. Turner syndrome (45,X)

Turner syndrome (45,X) was assessed in 20 studies, comprising 232 affected and 9849 non 45,X pregnancies. Among these studies, 16

enrolled pregnant women selected at high risk of fetal aneuploidy and four enrolled a cohort of pregnant women with mixed prior risk. Of the 20 studies, 14 evaluated MPSS and six evaluated TMPS. The results are summarised in Summary of findings 5.

a. Unselected population of pregnant women undergoing aneuploidy screening

No study assessed 45,X in this population.


b. Selected population of pregnant women at high risk of fetal aneuploidy

Sixteen studies included 198 (2.35%) affected and 8421 non 45,X pregnancies. MPSS and TMPS were assessed by 12 and four studies respectively (Figure 11).

Figure 11. Forest plot of MPSS and TMPS for 45,X in pregnant women selected at high risk of fetal aneuploidy. FN: false negative, FP: false positive, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, TN: true negative and TP: true positive.


MPSS 45,X

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Lefkowitz 2016	19	7	0	1118	1.00 [0.82, 1.00]	0.99 [0.99, 1.00]
Song 2015	0	1	1	202	0.00 [0.00, 0.97]	1.00 [0.97, 1.00]
Sukhikh 2015	4	1	0	195	1.00 [0.40, 1.00]	0.99 [0.97, 1.00]
Porreco 2014	9	11	0	3258	1.00 [0.66, 1.00]	1.00 [0.99, 1.00]
Bianchi 2013	20	0	1	92	0.95 [0.76, 1.00]	1.00 [0.96, 1.00]
Liang 2013	5	1	3	403	0.63 [0.24, 0.91]	1.00 [0.99, 1.00]
Mazloom 2013	20	1	1	389	0.95 [0.76, 1.00]	1.00 [0.99, 1.00]
Bianchi 2012	21	46	6	416	0.78 [0.58, 0.91]	0.90 [0.87, 0.93]
Hou 2012	1	1	0	203	1.00 [0.03, 1.00]	1.00 [0.97, 1.00]
Jiang 2012	3	1	1	898	0.75 [0.19, 0.99]	1.00 [0.99, 1.00]
Liu 2012	1	0	0	152	1.00 [0.03, 1.00]	1.00 [0.98, 1.00]
Sehnert 2011	3	1	0	43	1.00 [0.29, 1.00]	0.98 [0.88, 1.00]



TMPS 45,X

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Persico 2016	2	0	1	246	0.67 [0.09, 0.99]	1.00 [0.99, 1.00]
Hooks 2014	26	2	1	385	0.96 [0.81, 1.00]	0.99 [0.98, 1.00]
Nicolaides 2014a	43	0	4	125	0.91 [0.80, 0.98]	1.00 [0.97, 1.00]
Nicolaides 2013	2	0	0	227	1.00 [0.16, 1.00]	1.00 [0.98, 1.00]



i. MPSS

Twelve studies evaluated MPSS in 119 affected and 7440 non 45,X pregnancies (Bianchi 2012; Bianchi 2013; Hou 2012; Jiang 2012; Lefkowitz 2016; Liang 2013; Liu 2012; Mazloom 2013; Porreco 2014; Sehnert 2011; Song 2015; Sukhikh 2015). The summary sensitivity (95% CI) was 91.7% (78.3% to 97.1%) and the summary specificity (95% CI) was 99.6% (98.9% to 99.8%).

ii. TMPS

Four studies evaluated TMPS in 79 affected and 985 non 45,X pregnancies (Hooks 2014; Nicolaides 2013; Nicolaides 2014a; Persico 2016). The summary sensitivity (95% CI) was 92.4% (84.1% to 96.5%) and the summary specificity (95% CI) was 99.8% (98.3% to 100%).

iii. Comparative accuracy of MPSS and TMPS

An indirect comparison of the 12 MPSS and four TMPS studies showed no statistical evidence of a difference in sensitivity,

specificity or both (P value = 0.40). The differences in sensitivity and specificity were negligible ([Summary of findings 5](#)).


5. Triple X syndrome (47,XXX)

Seven studies assessed 47,XXX, comprising 14 (0.23%) affected and 6021 non 47,XXX pregnancies ([Hooks 2014](#); [Lefkowitz 2016](#); [Liang](#)


[2013](#); [Mazloom 2013](#); [Nicolaidis 2014a](#); [Porreco 2014](#); [Song 2015](#)). The studies enrolled pregnant women selected at high risk of fetal aneuploidy. Five studies evaluated MPSS and two studies evaluated TMPS. ([Figure 12](#); [Table 8](#)). We did not perform a separate meta-analysis for 47,XXX due to sparse data (very few cases or studies, or one or more subgroups had no study).

Figure 12. Forest plot of MPSS and TMPS for 47,XXX, 47,XXY and 47,YYY in pregnant women selected at high risk of fetal aneuploidy. FN: false negative, FP: false positive, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, TN: true negative and TP: true positive.


MPSS 47,XXX




MPSS 47,XXY




MPSS 47,YYY




TMPS 47,XXX



TMPS 47,XXY



TMPS 47,YYY



6. Klinefelter syndrome (47,XXY)

Twelve studies assessed 47,XXY in 25 (0.33%) affected and 7584 non 47,XXY pregnancies (Hooks 2014; Hou 2012; Jiang 2012; Lau 2012; Lefkowitz 2016; Liang 2013; Mazloom 2013; Nicolaidis 2014a; Persico 2016; Porreco 2014; Samango-Sprouse 2013; Song 2015).

Ten studies enrolled pregnant women selected at high risk of fetal aneuploidy (Figure 12; Table 8) and two studies enrolled pregnant women with mixed risk (See Finding section 11). No study assessed 47,XXY in an unselected population of pregnant women undergoing aneuploidy screening. Eight studies assessed MPSS

and four studies assessed TMPS. We did not perform a separate meta-analysis for 47,XXY due to sparse data (very few cases or studies, or one or more subgroups had no study).

7. 47,XXY

Ten studies assessed 47,XXY in 16 (0.23%) affected and 6971 non 47,XXY pregnancies (Hou 2012; Jiang 2012; Lefkowitz 2016; Liang 2013; Liu 2012; Mazloom 2013; Nicolaidis 2014a; Porreco 2014; Samango-Sprouse 2013; Shaw 2014). Eight studies enrolled pregnant women selected at high risk of fetal aneuploidy (Figure 12; Table 8) and two studies enrolled pregnant women with mixed risk (See Finding section 11). Eight studies used MPSS and two studies used TMPS. We did not perform a separate meta-analysis for 47,XXX due to sparse data (very few cases or studies, or one or more subgroups had no study).

8. All autosomes combined

Autosomal aneuploidies were assessed in 61 studies. The studies included 84,954 pregnant women of which 2853 were T21, T18 or T13 pregnancies and 82,073 were unaffected. Among these 61 studies, 43 assessed MPSS and 18 assessed TMPS. Of the 61 studies, five enrolled unselected pregnant women, 39 enrolled high-risk pregnant women and 17 enrolled a cohort of mixed prior risk. The results are summarised in Summary of findings 6. The results for mixed risk cohorts are summarised in Appendix 7.


a. Unselected population of pregnant women undergoing aneuploidy screening

Five studies assessed 129 (0.58%) affected and 22,379 unaffected (non T21, T18 and T13) pregnancies. Of the five studies, one study assessed MPSS and four studies assessed TMPS (Figure 13).

Figure 13. Forest plot of MPSS and TMPS for autosomes (T21, T18 and T13 combined) in unselected pregnant women undergoing aneuploidy screening. FN: false negative, FP: false positive, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, TN: true negative and TP: true positive.


MPSS, autosomes

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Song 2013	11	1	0	1729	1.00 [0.72, 1.00]	1.00 [1.00, 1.00]



TMPS, autosomes

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Norton 2015	49	12	1	15779	0.98 [0.89, 1.00]	1.00 [1.00, 1.00]
Quezada 2015	43	8	4	2730	0.91 [0.80, 0.98]	1.00 [0.99, 1.00]
del Mar Gil 2014	10	0	1	181	0.91 [0.59, 1.00]	1.00 [0.98, 1.00]
Nicolaidis 2012	10	2	0	1937	1.00 [0.69, 1.00]	1.00 [1.00, 1.00]



i. MPSS

Only one study assessed MPSS (Song 2013). The study evaluated 1730 unaffected (non T21, T18 and T13) pregnancies and 11 cases in women with singleton pregnancy. The sensitivity (95% CI) was 100% (74.1% to 100%) and the specificity (95% CI) was 99.9% (99.7% to 100%).

ii. TMPS

Four studies assessed TMPS in 20,649 unaffected (non T21, T18 and T13) pregnancies and 118 cases (del Mar Gil 2014; Nicolaidis 2012; Norton 2015; Quezada 2015). Of the four studies, three studies included only women with singleton pregnancy and the remaining study included only women with multifetal pregnancy (Table 9).

Based on the four studies, the summary sensitivity (95% CI) was 94.9% (89.1% to 97.7%) and the summary specificity (95% CI) was 99.9% (99.8% to 99.9%).

iii. Comparative accuracy of MPSS and TMPS


It was not possible to compare the accuracy of MPSS and TMPS in a meta-analysis due to limited data.

b. Selected population of pregnant women at high risk of fetal aneuploidy


A total of 39 studies included 1886 (9.39%) affected and 20,079 unaffected (non T21, T18 and T13) pregnancies. Of the 39 studies, 32 assessed MPSS and seven assessed TMPS (Figure 14).

Figure 14. Forest plot of MPSS and TMPS for autosomes (T21, T18 and T13) in pregnant women selected at high risk of fetal aneuploidy. FN: false negative, FP: false positive, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, TN: true negative and TP: true positive.

MPSS, autosomes



TMPS, autosomes



i. MPSS

Thirty-two MPSS studies evaluated 15,797 unaffected (non T21, T18 and T13) pregnancies and 1508 cases (Alberti 2015; Benachi 2015; Bianchi 2012; Bianchi 2013; Bijok 2014; Canick 2012; Chen 2011; Ehrich 2011; Hou 2012; Huang 2014; Jeon 2014; Jiang 2012; Johansen 2016; Ke 2015; Kim 2016; Lee 2015; Lefkowitz 2016; Liang 2013; Liu 2012; Palomaki 2012; Papageorghiou 2016a; Papageorghiou 2016b; Poon 2016; Porreco 2014; Sehnert 2011; Song 2015; Stumm 2014; Sukhikh 2015; Sung-Hee 2015;

Wang 2014; Wang 2015a; Zhang 2016). Of the 32 studies, 19 evaluated only singleton pregnancies, three evaluated only multifetal pregnancies, three evaluated singleton and multifetal pregnancies, and the remaining seven studies did not report type of pregnancy. Based on the 32 studies, the summary sensitivity (95% CI) was 98.8% (97.2% to 99.5%) and the summary specificity (95% CI) was 99.9% (99.7% to 100%). Results are presented separately for singleton and multifetal pregnancy studies in Table 9. The sensitivity tends to be lower in multifetal pregnancies but there are

no enough studies in this subgroup to compare MPSS performance according to pregnancy type.

ii. TMPS

Seven TMPS studies evaluated 378 cases and 4282 unaffected (non T21, T18 and T13) pregnancies in women with singleton pregnancy (Ashoor 2012; Hall 2014; Nicolaides 2013; Norton 2012; Persico 2016; Sparks 2012a; Verweij 2013). The summary sensitivity (95% CI) was 98.9 (97.2% to 99.6%) and the summary specificity (95% CI) was 99.9% (99.8% to 100%) (Table 9).

iii. Comparative accuracy of MPSS and TMPS

An indirect comparison of the 32 MPSS and seven TMPS studies showed no statistical evidence of a difference in sensitivity, specificity or both (P value = 0.11). The differences in sensitivity and specificity were negligible (Summary of findings 6).

9. All sex chromosome aneuploidies (SCA) combined

The sex chromosome aneuploidies (45,X, 47,XXX, 47,XXY and 47,YYY) were considered together as one target condition. SCA was

assessed in 20 studies, comprising 286 affected cases and 9839 non SCA pregnancies. MPSS and TMPS were assessed by 14 and six studies, respectively. Among the 20 studies, 16 enrolled pregnant women selected at high risk of fetal aneuploidy and four enrolled a cohort of pregnant women with mixed prior risk. The results are summarised in Summary of findings 7. The results for mixed risk cohorts are summarised in Appendix 7.

a. Unselected population of pregnant women undergoing aneuploidy screening

No study assessed SCA in an unselected population of pregnant women.


b. Selected population of pregnant women at high risk of fetal aneuploidy

Sixteen studies involving 247 (2.93%) affected and 8420 non SCA pregnancies were included. MPSS and TMPS were assessed by 12 and four studies respectively (Figure 15).

Figure 15. Forest plot of MPSS and TMPS for SCA (45,X, 47,XXX, 47,XXY and 47,YYY combined) in pregnant women selected at high risk of fetal aneuploidy. FN: false negative, FP: false positive, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, TN: true negative and TP: true positive.


MPSS, SCA

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Lefkowitz 2016	26	7	0	1111	1.00 [0.87, 1.00]	0.99 [0.99, 1.00]
Song 2015	0	1	2	201	0.00 [0.00, 0.84]	1.00 [0.97, 1.00]
Sukhikh 2015	4	1	0	195	1.00 [0.40, 1.00]	0.99 [0.97, 1.00]
Porreco 2014	15	16	0	3291	1.00 [0.78, 1.00]	1.00 [0.99, 1.00]
Bianchi 2013	20	0	1	92	0.95 [0.76, 1.00]	1.00 [0.96, 1.00]
Liang 2013	8	1	3	400	0.73 [0.39, 0.94]	1.00 [0.99, 1.00]
Mazloom 2013	29	1	1	380	0.97 [0.83, 1.00]	1.00 [0.99, 1.00]
Bianchi 2012	21	46	6	416	0.78 [0.58, 0.91]	0.90 [0.87, 0.93]
Hou 2012	1	1	2	201	0.33 [0.01, 0.91]	1.00 [0.97, 1.00]
Jiang 2012	6	1	1	895	0.86 [0.42, 1.00]	1.00 [0.99, 1.00]
Liu 2012	2	0	0	151	1.00 [0.16, 1.00]	1.00 [0.98, 1.00]
Sehnert 2011	3	1	0	43	1.00 [0.29, 1.00]	0.98 [0.88, 1.00]



TMPS, SCA

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Persico 2016	3	0	1	245	0.75 [0.19, 0.99]	1.00 [0.99, 1.00]
Hooks 2014	33	4	1	376	0.97 [0.85, 1.00]	0.99 [0.97, 1.00]
Nicolaides 2014a	52	1	4	115	0.93 [0.83, 0.98]	0.99 [0.95, 1.00]
Nicolaides 2013	2	0	0	227	1.00 [0.16, 1.00]	1.00 [0.98, 1.00]



i. MPSS

Twelve MPSS studies evaluated 151 affected and 7452 non SCA pregnancies (Bianchi 2012; Bianchi 2013; Hou 2012; Jiang 2012; Lefkowitz 2016; Liang 2013; Liu 2012; Mazloom 2013; Porreco 2014; Sehnert 2011; Song 2015; Sukhikh 2015). Of the 12 studies, seven included only women with singleton pregnancy, one evaluated singleton and multifetal pregnancies, and the remaining four studies did not report type of pregnancy. Results are presented

separately for singleton and multifetal pregnancy studies in Table 9. Based on all 12 studies, the summary sensitivity (95% CI) was 91.9% (73.8% to 97.9%) and the summary specificity (95% CI) was 99.5% (98.8% to 99.8%).

ii. TMPS

Four TMPS studies evaluated 96 affected and 968 non SCA pregnancies in women with singleton pregnancy (Hooks 2014;

Nicolaides 2013; Nicolaides 2014a; Persico 2016). The summary sensitivity (95% CI) was 93.8% (86.8% to 97.2%) and the summary specificity (95% CI) was 99.6% (98.1% to 99.9%).

iii. Comparative accuracy of MPSS and TMPS

An indirect comparison of the 12 MPSS and four TMPS studies showed no statistical evidence of a difference in sensitivity, specificity or both (P value = 0.41). The differences in sensitivity and specificity were negligible (Summary of findings 7).

10. gNIPT approach (MPSS or TMPS) against traditional screening tests

Five studies directly compared a gNIPT approach (MPSS or TMPS) and traditional screening tests for autosomal aneuploidies by using cohorts of pregnant women who were tested by both methods. Three studies compared TMPS and traditional screening tests, and

two studies compared MPSS and traditional screening tests. The results are summarised in Summary of findings 2, Summary of findings 3, Summary of findings 4 and Summary of findings 6.

a. Unselected population of pregnant women undergoing aneuploidy screening

Only one study that compared TMPS and a traditional screening test evaluated T21, T18 and T13 individually in an unselected population of pregnant women undergoing aneuploidy screening (Norton 2015). This study evaluated 38, 10 and two cases of T21, T18 and T13, respectively and 15,803, 15,831 and 11,183 non T21, T18 and T13, respectively (Figure 16). Direct comparisons between gNIPT and traditional screening tests were not possible because there was only one study but authors observed eight, two and one cases of T21, T18 and T13 respectively missed by traditional screening test and only one T18 case missed by TMPS.

Figure 16. Forest plot of traditional screening tests for T21, T18 and T13 in unselected pregnant women undergoing aneuploidy screening. FN: false negative, FP: false positive, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, TN: true negative and TP: true positive.

Traditional screening tests T21

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Norton 2015	30	854	8	14949	0.79 [0.63, 0.90]	0.95 [0.94, 0.95]		

Traditional screening tests T18

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Norton 2015	8	49	2	15782	0.80 [0.44, 0.97]	1.00 [1.00, 1.00]		

Traditional screening tests T13

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Norton 2015	1	28	1	11155	0.50 [0.01, 0.99]	1.00 [1.00, 1.00]		

Four studies compared a gNIPT approach with a traditional screening test for autosomal aneuploidies (T21, T18 and T13 combined) in 22,367 unselected pregnant women (Figure 17). Three studies (Nicolaides 2012; Norton 2015; Quezada 2015) compared TMPS and first-trimester combined test (Figure 18), and one study (Song 2013) compared MPSS and a second-trimester triple test. The three TMPS studies had similar characteristics. Meta-analyses of direct comparisons between gNIPT and traditional screening tests

were not possible because traditional screening tests used different cutpoints and there were very few studies to enable estimation of summary sensitivity and specificity at specific cutpoints. Individual study results are presented in Table 10. Overall, 16 aneuploid cases were missed by traditional screening test and only five cases were missed by gNIPT approach. While specificity was consistently higher for TMPS than traditional screening tests, sensitivity was not consistently higher as shown in Figure 18.


Figure 17. Forest plot of traditional screening tests for autosomes (T21, T18 and T13 combined) in unselected pregnant women undergoing aneuploidy screening. FN: false negative, FP: false positive, TN: true negative and TP: true positive.

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Norton 2015	39	931	11	14860	0.78 [0.64, 0.88]	0.94 [0.94, 0.94]		
Quezada 2015	49	124	0	2663	1.00 [0.93, 1.00]	0.96 [0.95, 0.96]		
Song 2013	6	243	5	1487	0.55 [0.23, 0.83]	0.86 [0.84, 0.88]		
Nicolaides 2012	10	87	0	1852	1.00 [0.69, 1.00]	0.96 [0.94, 0.96]		

Figure 18. Forest plot of comparative studies of TMPS and traditional screening tests for autosomes (T21, T18 and T13 combined) in unselected pregnant women undergoing aneuploidy screening. FN: false negative, FP: false positive, TN: true negative and TP: true positive.


TMPS, autosomes

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Norton 2015	49	12	1	15779	0.98 [0.89, 1.00]	1.00 [1.00, 1.00]
Quezada 2015	43	8	4	2730	0.91 [0.80, 0.98]	1.00 [0.99, 1.00]
Nicolaides 2012	10	2	0	1937	1.00 [0.69, 1.00]	1.00 [1.00, 1.00]



Traditional screening tests, autosomes

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Norton 2015	39	931	11	14860	0.78 [0.64, 0.88]	0.94 [0.94, 0.94]
Quezada 2015	49	124	0	2663	1.00 [0.93, 1.00]	0.96 [0.95, 0.96]
Nicolaides 2012	10	87	0	1852	1.00 [0.69, 1.00]	0.96 [0.94, 0.96]



b. Mixed-risk cohort of fetal aneuploidy


One study compared MPSS and traditional screening test for autosomal aneuploidies (T21, T18 and T13 combined) in a cohort with mixed prior risk of fetal aneuploidy including 1908 non T21, T18 and T31 pregnancies and four cases of autosomal aneuploidy

(Bianchi 2014a). Traditional screening tests included first-trimester combined test or a second-trimester result (quadruple, serum integrated, fully integrated or sequential) (Figure 19). Overall, 80 unaffected pregnancies were detected as affected by traditional screening test against 12 for TMPS.

Figure 19. Forest plot of traditional screening tests for autosomes (T21, T18 and T13 combined) in pregnant women with mixed prior risk of fetal aneuploidy. FN: false negative, FP: false positive, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, TN: true negative and TP: true positive.


Traditional screening tests, autosomes

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Bianchi 2014a	4	80	0	1828	1.00 [0.40, 1.00]	0.96 [0.95, 0.97]




Traditional screening tests T21

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Bianchi 2014a	3	69	0	1840	1.00 [0.29, 1.00]	0.96 [0.95, 0.97]



Traditional screening tests T18

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Bianchi 2014a	1	11	0	1894	1.00 [0.03, 1.00]	0.99 [0.99, 1.00]



11. Pregnant women with mixed prior risk of fetal aneuploidy

Summary sensitivities and specificities for cohorts of pregnant women with mixed prior risk of fetal aneuploidy are presented in Appendix 7. For autosomal aneuploidies, 17 studies included 838 cases and 39,615 unaffected (non T21, T18 and T13) pregnancies. Of the 17 studies, 10 assessed MPSS and seven assessed TMPS (Figure 20). For T21, 16 studies included 614 cases (1.6%) and 37,887 non T21 pregnancies. Of the 16 studies, 10 assessed MPSS and


six assessed TMPS. For T18, 13 studies included 166 cases (0.5%) and 36,206 non T18 pregnancies. Of the 13 studies, nine assessed MPSS and four assessed TMPS. For T13, 13 studies included 58 cases (0.1%) and 38,746 non T13 pregnancies. Eight of the 13 studies assessed MPSS and the other five assessed TMPS (Figure 21). For SCA, four studies included 39 cases and 1419 non SCA pregnancies; two of the studies assessed MPSS and the other two assessed TMPS (Figure 22). For 45,X, four studies included 34 cases (2.4%) and 1424 non 45,X pregnancies. Of the four studies, two studies assessed

MPSS and two studies assessed TMPS. For 47,XXY, two studies (one of MPSS and one of TMPS) included three cases (1%) and 291 non 47,XXY pregnancies. For 47,XYY, two studies included two cases (0.5%) and 384 non 47,XYY pregnancies; one study assessed MPSS

and the other study assessed TMPS. No study assessed gNIPT for 47,XXX in cohorts of pregnant women with mixed prior risk of fetal aneuploidy (Figure 23).

Figure 20. Forest plot of MPSS and TMPS for autosomes (T21, T18 and T13 combined) in pregnant women with mixed prior risk of fetal aneuploidy. FN: false negative, FP: false positive, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, TN: true negative and TP: true positive.

MPSS, autosomes



TMPS, autosomes






Figure 21. Forest plot of MPSS and TMPS for T21, T18 or T13 in pregnant women with mixed prior risk of fetal aneuploidy. FN: false negative, FP: false positive, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, TN: true negative and TP: true positive.


MPSS T21




MPSS T18




MPSS T13



TMPS T21



TMPS T18



TMPS T13

Figure 21. (Continued)

TMPS T13





Figure 22. Forest plot of MPSS and TMPS for SCA (45,X, 47,XXX, 47,XXY and 47,XYY combined) in pregnant women with mixed prior risk of fetal aneuploidy. FN: false negative, FP: false positive, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, TN: true negative and TP: true positive.

MPSS, SCA



TMPS, SCA





Figure 23. Forest plot of MPSS and TMPS for 45,X, 47,XXY or 47,XYY in pregnant women with mixed prior risk of fetal aneuploidy. FN: false negative, FP: false positive, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, TN: true negative and TP: true positive.


MPSS 45,X

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Shaw 2014	3	0	1	196	0.75 [0.19, 0.99]	1.00 [0.98, 1.00]
Lau 2012	8	0	0	100	1.00 [0.63, 1.00]	1.00 [0.96, 1.00]




MPSS 47,XXY

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Lau 2012	1	0	0	107	1.00 [0.03, 1.00]	1.00 [0.97, 1.00]




MPSS 47,XYY

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Shaw 2014	1	0	0	199	1.00 [0.03, 1.00]	1.00 [0.98, 1.00]




TMPS 45,X

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	
Pergament 2014		9	1	1	953	0.90 [0.55, 1.00]	1.00 [0.99, 1.00]
Samango-Sprouse 2013		11	0	1	174	0.92 [0.62, 1.00]	1.00 [0.98, 1.00]




TMPS 47,XXY

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Samango-Sprouse 2013	2	0	0	184	1.00 [0.16, 1.00]	1.00 [0.98, 1.00]



TMPS 47,XYY

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Samango-Sprouse 2013	1	0	0	185	1.00 [0.03, 1.00]	1.00 [0.98, 1.00]



12. Failure rates

Table 7 shows the non-negligible failure rate of gNIPT reported in the studies. gNIPT assay failure rate was reported in 46 out of 65 (71%) studies. The largest failure rate (25%) was observed in a study that used its own developed MPSS assay (Alberti 2015). The main reasons for assay failure included low amount of ccfDNA, low fetal fraction DNA and failure of sample to pass quality control. The failure rate ranged between 0% and 25% for MPSS and between 0.8% and 7.5% for TMPS. The number of aneuploid and euploid cases in failed samples was reported in 23 of 46 (50%) studies. Among these 23 studies, there were 1064 euploid cases and 79 aneuploid cases among 1143 failed samples. The failure rate among aneuploid cases, ranged between 0% and 50% for MPSS and between 0% and 23% for TMPS. The failure rate among euploid cases ranged between 0% and 6.7% for MPSS and between 1% and 7.6% for TMPS.

Investigation of heterogeneity

We planned to evaluate the effect of potential sources of heterogeneity such as type of reference standard and ethnicity. However, formal investigations using meta-regression were not possible due to limited data and little or no heterogeneity in the sensitivities and specificities. Most studies (55%) used karyotyping

while the remaining 29 studies (45%) used multiple reference standards. Ethnicity was not reported by 26 (40%) studies while the population in 21 (32%) studies was more than 50% Asian and in 18 (28%) studies the population was more than 50% Caucasian. In Appendix 8, the number of studies, affected and unaffected pregnancies are shown according to the gNIPT approach and prior risk of fetal aneuploidy. We also planned to assess gNIPT performance according to gestational age and gNIPT approach for autosomes and SCA aneuploidies. The accuracy of gNIPT appears to be high in all gestational age groups.

Sensitivity analyses

We did not perform sensitivity analyses to assess the effect of the interval between blood collection for gNIPT and fluid collection for reference standard because most studies had an acceptable interval between sample collection for index test and reference standard. Due to lack of data or lack of variability in estimates of sensitivity and specificity, analyses of the effect of high or unclear risk of bias according to the QUADAS-2 domains were not done. We performed sensitivity analyses using data from all autosomes combined and all SCA combined in order to have enough studies to assess the impact of study design and number of cases. The results are presented in Table 11. Excluding case-control studies or studies

with less than 10 aneuploid cases had little or no impact on our findings.

DISCUSSION

Summary of main results

This review included data from 65 studies of 86,139 pregnant women (including 3141 aneuploids) tested by genomics-based non-invasive prenatal testing (gNIPT) and a reference standard. The gNIPT method used circulating cell-free DNA (ccfDNA) in maternal blood for the detection of common fetal aneuploidies (T21, T18, T13, 45,X, 47,XXY, 47,XXX and 47,XYY). The number of gNIPT studies in unselected populations was limited (five studies), but 42 studies in high-risk cohorts provided data for various meta-analyses. Few (14%) studies included more than 100 aneuploid cases. Importantly, in almost all studies, the risk of bias was generally high with respect to patient selection as well as flow and timing. Some women can spontaneously lose their pregnancy after enrolment into a study. However, none of the studies reported such events. Since women with spontaneous abortions are likely to be lost to follow-up, we believe that any risk of bias has been captured in the quality assessment of studies. Blood samples for gNIPT were mainly taken just before the invasive test (reference standard) and so pregnancies were unlikely to terminate naturally between the gNIPT and the reference standard. Across all studies, applicability concerns were low in the index test and reference standard domains.

These results show that massively parallel shotgun sequencing (MPSS) and targeted massively parallel sequencing (TMPS) perform similarly in terms of clinical sensitivity and specificity for the detection of fetal T21, T18, T13 and sex chromosome aneuploidy (SCA). However, no study compared the two approaches head-to-head in the same cohort of patients.

In high-risk pregnancies, gNIPT methods (MPSS and TMPS) were highly accurate for detection of any of the three major trisomies (T21, T18 and T13) with sensitivities from 95.8% to 99.7% depending on specific trisomies and specificities above 99%. There were no statistically significant differences in accuracy between MPSS and TMPS.

In unselected cohorts of pregnant women, only one study evaluated MPSS. Based on meta-analytic findings for each trisomy, TMPS appeared to be accurate for the detection of T21, with lower accuracy for T18 and T13. When compared to traditional prenatal screening tests, only four studies were identified (three for TMPS and one for MPSS). Genomics-based non-invasive prenatal testing showed greater specificity for T21 and T18 than traditional screening tests, while inconsistent results were observed for sensitivity. The inconsistency may be due to different cutpoints for traditional screening tests though one would expect that to also affect specificity. Given the small number of studies, the differences may be due to chance or there may be other differences between the studies that were not apparent.

With respect to the replacement of invasive tests, the performance of gNIPT observed in this review is not sufficient to replace current invasive diagnostic tests.

We also compared the diagnostic test accuracy of MPSS and TMPS for all three autosomes combined because gNIPT is being clinically

proposed as one test during prenatal follow-up to detect any of the three conditions. Under this scenario, in high-risk pregnancies of fetal aneuploidy, there was no statistically significant difference in diagnostic accuracy between MPSS and TMPS. In unselected cohorts of pregnant women, a test comparison was not possible due to limited data.

There was paucity of data for each SCA. In high-risk cohorts, all SCAs combined gave a pooled sensitivity (95% CI) and specificity (95% CI) of 91.9% (73.8% to 97.9%) and 99.5% (98.8% to 99.8%) from 12 MPSS studies. The pooled sensitivity (95% CI) and specificity (95% CI) were 93.8% (86.8% to 97.2%) and 99.6% (98.1% to 99.9%) from four TMPS studies. SCAs are considered “incidental” findings of current aneuploidy screening programs. It should be noted that SCAs are not of interest for prenatal screening since they do not lead to any intervention prior to birth.

The failure rate associated with gNIPT, which is higher than the current failure rate of traditional screening tests which is close to zero, is worrying and may be a source of bias. Furthermore, the large heterogeneity between laboratory-developed assays in their protocol details and observed failure rates highlight the fact that each laboratory providing gNIPT services should determine its own failure rate and inform healthcare professionals ordering the test about this important test characteristic. Failed samples were excluded from the analyses in the studies. This systematic review found a slightly larger failure rate for TMPS than the MPSS approach. This was also reported by [Yaron 2016](#). We also found that the proportion of failed samples for aneuploid samples was higher than the proportion of failed samples for euploid samples. If these failed samples were included in the summary statistics, the diagnostic performance of gNIPT would be lower.

Comparison with other systematic reviews with meta analysis

At the time of writing, there are four published systematic reviews with meta-analyses of gNIPT ([Gil 2015a](#); [HAS 2015](#); [Mackie 2017](#); [Taylor-Phillips 2016](#)). Although these meta-analyses had different criteria for including studies and analyses, they reported similar sensitivities and specificities to our findings.

As reported by [Gil 2015a](#), the detection rate of gNIPT for autosomes was between 91.0% to 99.2% and specificity above 99.9% in singleton pregnancies. The detection rate for 45,X and SCA other than 45,X was 90.3% and 93.0%, respectively with specificity above 99.8% in singleton pregnancies. The results from [HAS 2015](#) group for T21 were respectively 98.0% and 99.9% for sensitivity and specificity. Regarding [Mackie 2017](#), the sensitivity was between 90.6% to 99.4% and specificity above 99.9% for autosomes. For 45,X, the sensitivity and specificity was 92.9% and 99.9%, respectively. They also pointed out that failed results were poorly reported across studies. Finally, [Taylor-Phillips 2016](#) reported sensitivity between 97.4% to 99.3% for autosomes and specificity of 99.9%.

This is the first Cochrane diagnostic test accuracy (DTA) review on gNIPT. There are five published Cochrane DTA reviews on prenatal screening tests ([Alldred 2012](#); [Alldred 2015](#); [Alldred 2015a](#); [Alldred 2017a](#); [Alldred 2017b](#)). The suite of reviews addressed traditional biochemical, ultrasound and urine markers for Down syndrome screening ([Alldred 2010](#)) and none of the other fetal aneuploidies considered in this review were evaluated in this suite. In the first of the three reviews, Alldred and colleagues

evaluated second-trimester serum markers and found that double and triple test combinations (involving alpha-fetoprotein, human chorionic gonadotropin (hCG) (free and total) or unconjugated estriol) significantly outperformed individual markers, detecting six to seven out of every 10 Down syndrome pregnancies at a 5% false positive rate (Allred 2012). The second review evaluated first-trimester serum markers and found that a test strategy involving maternal age, PAPP-A and free β hCG significantly outperformed individual markers, detecting about seven out of every 10 Down's syndrome pregnancies at a 5% false positive rate (Allred 2015a). The third review evaluated urine markers and concluded there was a paucity of evidence to support the use of urine testing for Down syndrome screening (Allred 2015b). The fourth review evaluated first-trimester ultrasound tests alone or in combination with first-trimester serum tests and found that a combination of ultrasound and serum markers (especially PAPP-A and free β hCG) and maternal age can detect about nine of 10 T21 affected pregnancies for a fixed 5% false positive rate (Allred 2017a). The fifth review evaluated first- and second-trimester serum tests with and without first-trimester ultrasound tests and found that a combination of first-trimester ultrasound with first- and second-trimester serum markers with maternal age are significantly better than those without ultrasound or those evaluating first-trimester ultrasound in combination with second-trimester serum markers, without first-trimester serum markers (the authors cannot make recommendations about a specific strategy) (Allred 2017b).

Strengths and weaknesses of the review

Strengths

The review methodology was transparent with the full protocol published in the Cochrane Library (1 July 2015) and in PROSPERO (11 November 2015). The review evaluated the screening and diagnostic accuracy of gNIPT by MPSS and TMPS for seven common aneuploidies with no restriction imposed on population characteristics such as maternal age, gestational age, aneuploidy risk, number of fetuses and ethnicity. We performed a comprehensive search with no language restriction and we included studies in the languages used by various authors in the field, including Chinese, Bulgarian, Russian, Polish, Korean and Spanish. Study selection, data extraction and quality assessment were independently performed by two review authors. We contacted authors to clarify data and to avoid duplication of data as a result of overlapping populations.

We evaluated the performance of the two major gNIPT methods (MPSS and TMPS which included digital analysis of selected regions (DANSR) and single nucleotide polymorphism (SNP)-based method) and included data on traditional screening tests when compared to gNIPT.

We collected and reported data on excluded and failed samples and presented the failure rate at first attempt, the number of repeated tests and the final failure rate for each study. When it was possible, we also reported separate failure rates among aneuploid and euploid cases. Where possible, we performed subgroups analyses to investigate heterogeneity, and also performed sensitivity analyses to assess the robustness of these findings.

Weaknesses

Fetal karyotyping is the reference standard for establishing a diagnosis of fetal aneuploidy. This is an invasive procedure

with some risk for the fetus and the pregnant woman. Many pregnant women included in the studies, especially those involving unselected cohorts, were not tested by karyotyping. Rather, clinical examination of the newborn or medical records from birth were used as a secondary reference standard. We are aware that these secondary reference standards are not as accurate as fetal karyotype and some cases may have been missed.

Studies rarely reported the qualification of the person conducting the neonatal clinical examination at birth. Such examination is expected to be more reliable if it was made by a paediatrician or a geneticist. Ideally, this examination should be done a few months after birth because the phenotypic characteristics of aneuploidies are more apparent than at birth (Devlin 2004).

Genomics-based non-invasive prenatal testing assays are laboratory-developed tests that are not standardised in their methods, sequencing platforms, sequencing data manipulation, measures used or cut-offs for interpretation. Each assay was developed and validated by the testing laboratory and each laboratory has a different method. Usually detailed information about the assays were not available. As shown in Table 5, 15, different gNIPT assays were used in the studies included in this review. Thus, they may differ in various aspects and show different analytical and clinical validity. We have grouped them accordingly to the type of assay used (targeted versus shotgun), but there are also differences within each of these two subgroups that we were not able to account for, given the small number of studies published on most of these different assays. Thirteen of the assays were used only in studies of high-risk pregnancies or mixed cohorts. Only a few gNIPT assays were used in a significant number of studies. Thus, caution should be used before generalising the diagnostic accuracy observed in this category of patients to all gNIPT assays. This limits the generalisability of these findings and we cannot infer that all gNIPT assays will show the same performance.

Applicability of findings to the review question

These findings suggest that gNIPT has high sensitivity and specificity for detection of fetal aneuploidies in high-risk pregnancies. Performance varied depending on the type of aneuploidy. There was limited evidence of the performance of gNIPT in unselected cohorts of pregnant women. Most studies involved either high-risk pregnancies or mixed populations where it was not possible to differentiate between high-risk pregnancies and unselected pregnant women. Thus, more studies are needed in the general population of pregnant women before firm conclusions can be made about the sensitivity of gNIPT as a first-tier screening test. The two major types of gNIPT method (MPSS and TMPS) appear to have comparable performance, but there are many different gNIPT assays for each approach. For many of these assays, very little data have been published about their diagnostic accuracy. Additionally, performance in the cohorts studied may not reflect performance in other populations owing to differences in fetal fraction distribution because of, for example, differences in mean body mass index or gestational age. Importantly, summary sensitivities and specificities derived from cohort data can be very different from the probability associated with any particular patient sample to be positive or negative depending on the sample's specific fetal fraction. Thus, summary sensitivity, specificity and associated predictive values of an assay cannot be used as a straightforward measure of the probability of a specific patient's sample to be affected given a positive or negative result. This

underscores the importance, before clinically offering a laboratory developed gNIPT assay, that it is fully validated according to recognised best practice clinical laboratory molecular diagnostics guidelines. Finally, the methodological quality of studies was generally poor with high risk of bias, especially in terms of patient selection and flow and timing.

AUTHORS' CONCLUSIONS

Implications for practice

Genomics-based non-invasive prenatal testing (gNIPT) appears to be an accurate prenatal screening test, its accuracy having been evaluated as a second-tier screening test to identify pregnancies at very low risk of fetal aneuploidies (T21, T18 and T13) and thus to decrease the false positive rate of traditional screening approaches and avoid invasive procedures in those pregnant women. As a first-tier aneuploidy screening test, based on limited data from comparative studies, gNIPT appears to have significantly better specificity than current screening approaches using maternal serum biochemical markers, ultrasound or both, but evidence about sensitivity is inconsistent. At current gNIPT pricing levels, gNIPT as a second-tier screening test provides the best value for money, especially for publicly-funded screening programs while gNIPT as a first-tier screening test was found not to be cost-effective (Nshimyumukiza 2017). The failure rate of gNIPT is a concern as it is substantially larger than the current failure rate of traditional prenatal screening approaches.

It is worth noting that gNIPT shows good performance for the detection of sex chromosome aneuploidies though data are sparse. The number of studies for sex chromosome aneuploidy (SCA) was small and confidence intervals on sensitivity and specificity estimates are therefore wide. Thus, sex chromosome aneuploidies appear to be more difficult to detect since performances of gNIPT are not as good as for detecting autosomal aneuploidies. SCAs are considered "incidental" findings of current aneuploidy screening programs and they do not lead to any intervention prior to birth.

Maternal serum screening, ultrasound fetal examination, gNIPT and invasive diagnostic tests are thus complementary approaches because in its current state, gNIPT cannot detect all chromosomal abnormalities or adverse obstetrical outcomes. About 44% to 64% of all chromosomal abnormalities found during prenatal diagnostic are common aneuploidies which gNIPT can detect (Kazerouni 2011; Shani 2016). Counselling expectant mothers and their partners is essential for explaining the advantages, limitations and risks of these procedures.

We conclude that given the current data on the performance of gNIPT, invasive fetal karyotyping is still the required diagnostic approach to confirm the presence of a chromosomal abnormality

prior to making irreversible decisions relative to the pregnancy outcome.

Implications for research

This systematic review has highlighted the fact that most published studies on gNIPT have high risk of bias in the patient selection and flow and timing domains. Many different gNIPT assays are in use and for the majority of them, there is insufficient published data to individually assess their clinical performance. Therefore, the results in this systematic review may not be generalisable to all gNIPT assays. Studies are needed that directly compare the accuracy of gNIPT with that of current traditional prenatal screening methods for fetal aneuploidy, especially in unselected populations of pregnant women. Such studies can provide valid evidence of the incremental accuracy of gNIPT if gNIPT is being considered as a first-tier test. Particular attention should be paid to study design in order to minimise patient selection biases as well as biases in flow and timing domain. Further well-designed, independent large-scale studies on real life gNIPT's implementation into prenatal care should be performed. Large scale randomised clinical trials of tests and patient outcomes are needed to validate the clinical utility of gNIPT in the various clinical settings. Given the rapid evolution of gNIPT and its capacity to detect other fetal chromosomal anomalies (Benn 2016), future systematic reviews may have to widen the scope of target conditions.

ACKNOWLEDGEMENTS

We thank the Editorial Staff of Cochrane Pregnancy and Childbirth and the Editorial Team for Cochrane Diagnostic Test Accuracy Reviews for their support. We also thank Lynn Hampson for her help with development of the search strategy and members of "Cochrane Canada Francophone" for their support.

As part of the pre-publication editorial process, this review has undergone dual peer review with Cochrane Pregnancy and Childbirth and the Cochrane Diagnostic Test Accuracy Reviews Unit and has been commented by peer reviewers.

This project was supported by the National Institute for Health Research, via Cochrane Infrastructure funding to Cochrane Pregnancy and Childbirth. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, NHS or the Department of Health.

We wish to thank our translators. Thanks to Dr Ze Zhang for his help with Chinese publications, Roumiana Gulemetova for her help with Bulgarian and Russian publications, Sylvie Giroux for her help with Spanish publications, Richard Szostak for his help with Polish publication and Hyun Jin Park for her help with Korean publication.

REFERENCES

References to studies included in this review

Alberti 2015 {published and unpublished data}

* Alberti A, Salomon LJ, Le Lorc'h M, Couloux A, Bussi eres L, Goupil S, et al. Non-invasive prenatal testing for trisomy 21 based on analysis of cell-free fetal DNA circulating in the maternal plasma. *Prenatal Diagnosis* 2015;**35**(5):471-6. [PUBMED: 25643828]

NCT01118507. Trisomy of chromosome 21 diagnosis by high output sequencing. clinicaltrials.gov/show/NCT01118507 Date first received: 28 April 2010.

Ashoor 2012 {published data only}

Ashoor G, Syngelaki A, Wagner M, Birdir C, Nicolaides KH. Chromosome-selective sequencing of maternal plasma cell-free DNA for first-trimester detection of trisomy 21 and trisomy 18. *American Journal of Obstetrics and Gynecology* 2012;**206**(4):322.e1-5. [PUBMED: 22464073]

Ashoor 2013 {published data only}

Ashoor G, Syngelaki A, Wang E, Struble C, Oliphant A, Song K, et al. Trisomy 13 detection in the first trimester of pregnancy using a chromosome-selective cell-free DNA analysis method. *Ultrasound in Obstetrics & Gynecology* 2013;**41**(1):21-5. [PUBMED: 22996646]

Benachi 2015 {published data only}

Benachi A, Letourneau A, Kleinfinger P, Senat MV, Gautier E, Favre R, et al. Cell-free DNA analysis in maternal plasma in cases of fetal abnormalities detected on ultrasound examination. *Obstetrics and Gynecology* 2015;**125**(6):1330-7. [PUBMED: 26000504]

Bevilacqua 2015 {published data only}

Bevilacqua E, Gil MM, Nicolaides KH, Ordonez E, Cirigliano V, Dierickx H, et al. Performance of screening for aneuploidies by cell-free DNA analysis of maternal blood in twin pregnancies. *Ultrasound in Obstetrics & Gynecology* 2015;**45**(1):61-6. [PUBMED: 25297464]

Bianchi 2012 {published data only}

Bianchi DW, Platt LD, Goldberg JD, Abuhamad AZ, Sehnert AJ, Rava RP, et al. Genome-wide fetal aneuploidy detection by maternal plasma DNA sequencing. *Obstetrics and Gynecology* 2012;**119**(5):890-901. [PUBMED: 22362253]

Bianchi 2013 {published and unpublished data}

* Bianchi DW, Prosen T, Platt LD, Goldberg JD, Abuhamad AZ, Rava RP, et al. Massively parallel sequencing of maternal plasma DNA in 113 cases of fetal nuchal cystichygroma. *Obstetrics and Gynecology* 2013;**121**(5):1057-62.

NCT01122524. MatErnal blood IS Source to accurately diagnose fetal Aneuploidy (MELISSA). clinicaltrials.gov/show/NCT01122524 Date first received: 11 May 2010.

Bianchi 2014a {published and unpublished data}

* Bianchi DW, Parker RL, Wentworth J, Madankumar R, Saffer C, Das AF, et al. DNA sequencing versus standard prenatal

aneuploidy screening. *New England Journal of Medicine* 2014;**370**(9):799-808. [PUBMED: 24571752]

NCT01663350. Comparison of aneuploidy risk evaluations. clinicaltrials.gov/show/NCT01663350 Date first received: 31 July 2012.

Bijok 2014 {published data only}

Bijok J, Gorzelnik K, Massalska D, Ilnicka A, Pawlowska B, Zimowski JG, et al. Non-invasive prenatal diagnosis of the most common aneuploidies with cell-free fetal DNA in maternal serum--preliminary results [Nieinwazyjna diagnostyka prenatalna najczestszych aneuploidii na podstawie plodowego DNA we krwi matki --doniesienie wstepne]. *Ginekologia Polska* 2014;**85**(3):208-13. [PUBMED: 24783433]

Canick 2012 {published data only}

Canick JA, Kloza EM, Lambert-Messerlian GM, Haddow JE, Ehrich M, van den Boom D, et al. DNA sequencing of maternal plasma to identify Down syndrome and other trisomies in multiple gestations. *Prenatal Diagnosis* 2012;**32**(8):730-4. [PUBMED: 22585317]

Chen 2011 {published data only}

Chen EZ, Chiu RW, Sun H, Akolekar R, Chan KC, Leung TY, et al. Noninvasive prenatal diagnosis of fetal trisomy 18 and trisomy 13 by maternal plasma DNA sequencing. *PLOS One* 2011;**6**(7):e21791. [PUBMED: 21755002]

Chiu 2011 {published data only}

Chiu RW, Akolekar R, Zheng YW, Leung TY, Sun H, Chan KC, et al. Non-invasive prenatal assessment of trisomy 21 by multiplexed maternal plasma DNA sequencing: large scale validity study. *BMJ (Clinical Research Ed.)* 2011;**342**:c7401. [PUBMED: 21224326]

Comas 2015 {published data only}

Comas C, Echevarria M, Rodriguez MA, Prats P, Rodriguez I, Serra B. Initial experience with non-invasive prenatal testing of cell-free DNA for major chromosomal anomalies in a clinical setting. *Journal of Maternal-fetal & Neonatal Medicine* 2015;**28**(10):1196-201. [PUBMED: 25048745]

del Mar Gil 2014 {published data only}

del Mar Gil M, Quezada MS, Bregant B, Syngelaki A, Nicolaides KH. Cell-free DNA analysis for trisomy risk assessment in first-trimester twin pregnancies. *Fetal Diagnosis and Therapy* 2014;**35**(3):204-11. [PUBMED: 24247435]

Ehrich 2011 {published data only}

Ehrich M, Deciu C, Zwiefelhofer T, Tynan JA, Cagasan L, Tim R, et al. Noninvasive detection of fetal trisomy 21 by sequencing of DNA in maternal blood: a study in a clinical setting. *American Journal of Obstetrics and Gynecology* 2011;**204**(3):205.e1-11. [PUBMED: 21310373]

Fiorentino 2016 {published data only}

Fiorentino F, Bono S, Pizzuti F, Mariano M, Polverari A, Duca S, et al. The importance of determining the limit of detection of

- non-invasive prenatal testing methods. *Prenatal Diagnosis* 2016;**36**(4):304-11. [PUBMED: 26815144]
- Gil 2016** {published data only}
 Gil MM, Revello R, Poon LC, Akolekar R, Nicolaides KH. Clinical implementation of routine screening for fetal trisomies in the UK NHS: cell-free DNA test contingent on results from first-trimester combined test. *Ultrasound in Obstetrics & Gynecology* 2016;**47**(1):45-52. [PUBMED: 26498918]
- Hall 2014** {published data only}
 Hall MP, Hill M, Zimmermann B, Sigurjonsson S, Westemeyer M, Saucier J, et al. Non-invasive prenatal detection of trisomy 13 using a single nucleotide polymorphism- and informatics-based approach. *PLOS One* 2014;**9**(5):e96677. [PUBMED: 24805989]
- Hooks 2014** {published data only}
 Hooks J, Wolfberg AJ, Wang ET, Struble CA, Zahn J, Juneau K, et al. Non-invasive risk assessment of fetal sex chromosome aneuploidy through directed analysis and incorporation of fetal fraction. *Prenatal Diagnosis* 2014;**34**(5):496-9. [PUBMED: 24510887]
- Hou 2012** {published data only}
 Hou QF, Wu D, Chu Y, Kang B, Liao SX, Yang YL, et al. Clinical application of noninvasive prenatal diagnosis using cell free fetal DNA in maternal plasma. *Zhonghua Fu Chan Ke za Zhi* 2012;**47**(11):813-7. [PUBMED: 23302120]
- Huang 2014** {published data only}
 Huang X, Zheng J, Chen M, Zhao Y, Zhang C, Liu L, et al. Noninvasive prenatal testing of trisomies 21 and 18 by massively parallel sequencing of maternal plasma DNA in twin pregnancies. *Prenatal Diagnosis* 2014;**34**(4):335-40. [PUBMED: 24357023]
- Jackson 2014** {published data only}
 Jackson J, Hamar B, Lazar E, Lim K, Rodriguez D, Stock K, et al. Nuchal translucency measurement plus non-invasive prenatal testing to screen for aneuploidy in a community-based average-risk population. *Ultrasound in Obstetrics & Gynecology* 2014; Vol. 44, issue 4:491. [PUBMED: 24890031]
- Jeon 2014** {published data only}
 Jeon YJ, Zhou Y, Li Y, Guo Q, Chen J, Quan S, et al. The feasibility study of non-invasive fetal trisomy 18 and 21 detection with semiconductor sequencing platform. *PLOS One* 2014;**9**(10):e110240.
- Jiang 2012** {published data only}
 Jiang F, Ren J, Chen F, Zhou Y, Xie J, Dan S, et al. Noninvasive Fetal Trisomy (NIFTY™) test: an advanced noninvasive prenatal diagnosis methodology for fetal autosomal and sex chromosomal aneuploidies. *BMC Medical Genomics* 2012;**5**:57. [PUBMED: 23198897]
- Johansen 2016** {published data only}
 Johansen P, Richter SR, Balslev-Harder M, Miltoft CB, Tabor A, Duno M, et al. Open source non-invasive prenatal testing platform and its performance in a public health laboratory. *Prenatal Diagnosis* 2016;**36**(6):530-6. [PUBMED: 27027563]
- Ke 2015** {published data only}
 Ke WL, Zhao WH, Wang XY. Detection of fetal cell-free DNA in maternal plasma for Down syndrome, Edward syndrome and Patau syndrome of high risk fetus. *International Journal of Clinical and Experimental Medicine* 2015;**8**(6):9525-30. [PUBMED: 26309618]
- Kim 2016** {published data only}
 Kim S, Jung H, Han SH, Lee S, Kwon J, Kim MG, et al. Comparison of two high-throughput semiconductor chip sequencing platforms in noninvasive prenatal testing for Down syndrome in early pregnancy. *BMC Medical Genomics* 2016;**9**(1):22. [PUBMED: 27129388]
- Korostelev 2014** {published data only}
 Korostelev S, Totchiev G, Kanivets I, Gnetetskaya V. Association of non-invasive prenatal testing and chromosomal microarray analysis for prenatal diagnostics. *Gynecological Endocrinology* 2014;**30**:13-6.
- Lau 2012** {published data only}
 Lau TK, Chen F, Pan X, Pooh RK, Jiang F, Li Y, et al. Noninvasive prenatal diagnosis of common fetal chromosomal aneuploidies by maternal plasma DNA sequencing. *Journal of Maternal-Fetal & Neonatal Medicine* 2012;**25**(8):1370-4. [PUBMED: 22070770]
- Lee 2015** {published data only}
 Lee MY, Cho DY, Won HS, Hwang AR, Jeong B, Kim J, et al. Performance of Momguard, a new non-invasive prenatal testing protocol developed in Korea. *Obstetrics & Gynecology Science* 2015;**58**(5):340-5. [PUBMED: 26430657]
- Lefkowitz 2016** {published data only}
 Lefkowitz RB, Tynan JA, Liu T, Wu Y, Mazloom AR, Almasri E, et al. Clinical validation of a noninvasive prenatal test for genomewide detection of fetal copy number variants. *American Journal of Obstetrics and Gynecology* 2016;**215**(2):227.e1-227.e16. [PUBMED: 26899906]
- Liang 2013** {published data only}
 Liang D, Lv W, Wang H, Xu L, Liu J, Li H, et al. Non-invasive prenatal testing of fetal whole chromosome aneuploidy by massively parallel sequencing. *Prenatal Diagnosis* 2013;**33**(5):409-15. [PUBMED: 23299662]
- Liu 2012** {published data only}
 Liu HY, Wu D, Li H, Guo SK, Zhang CY, Liao SX, et al. Significance of detecting free DNA from maternal plasma for the diagnosis of fetal chromosomal aneuploidies. *Zhonghua yi xue yi chuan xue za zhi = Zhonghua yixue yichuanxue zazhi [Chinese journal of medical genetics]* 2012;**29**(4):435-8. [PUBMED: 22875501]
- Ma 2016** {published data only}
 Ma J, Wang Y, Wang W, Dong Y, Xu C, Zhou A, et al. Validation of combinatorial probe-anchor ligation (cPAL) based sequencing method for non-invasive prenatal testing in trisomy detection by a central laboratory. *Ultrasound in Obstetrics & Gynecology* 2016;**50**(1):49-57. [PUBMED: 27363706]

Mazloom 2013 {published data only}

Mazloom AR, Dzakula Z, Oeth P, Wang H, Jensen T, Tynan J, et al. Noninvasive prenatal detection of sex chromosomal aneuploidies by sequencing circulating cell-free DNA from maternal plasma. *Prenatal Diagnosis* 2013;**33**(6):591-7. [PUBMED: 23592550]

Nicolaidis 2012 {published data only}

Nicolaidis KH, Syngelaki A, Ashoor G, Birdir C, Touzet G. Noninvasive prenatal testing for fetal trisomies in a routinely screened first-trimester population. *American Journal of Obstetrics and Gynecology* 2012;**207**(5):374.e1-6. [PUBMED: 23107079]

Nicolaidis 2013 {published data only}

Nicolaidis KH, Syngelaki A, Gil M, Atanasova V, Markova D. Validation of targeted sequencing of single-nucleotide polymorphisms for non-invasive prenatal detection of aneuploidy of chromosomes 13, 18, 21, X, and Y. *Prenatal Diagnosis* 2013;**33**(6):575-9.

Nicolaidis 2014a {published data only}

Nicolaidis KH, Musci TJ, Struble CA, Syngelaki A, Gil MM. Assessment of fetal sex chromosome aneuploidy using directed cell-free DNA analysis. *Fetal Diagnosis and Therapy* 2014;**35**(1):1-6. [PUBMED: 24335155]

Norton 2012 {published data only}

Norton ME, Brar H, Weiss J, Karimi A, Laurent LC, Caughey AB, et al. Non-Invasive Chromosomal Evaluation (NICE) Study: results of a multicenter prospective cohort study for detection of fetal trisomy 21 and trisomy 18. *American Journal of Obstetrics and Gynecology* 2012;**207**(2):137.e1-8. [PUBMED: 22742782]

Norton 2015 {published and unpublished data}

NCT01511458. Non-invasive chromosomal examination of Trisomy Study. clinicaltrials.gov/show/NCT01511458 Date first received: 13 January 2012.

* Norton ME, Jacobsson B, Swamy GK, Laurent LC, Ranzini AC, Brar H, et al. Cell-free DNA analysis for noninvasive examination of trisomy. *New England Journal of Medicine* 2015;**372**(17):1589-97. [PUBMED: 25830321]

Palomaki 2012 {published data only}

Palomaki GE, Decui C, Kloza EM, Lambert-Messerlian GM, Haddow JE, Neveux LM, et al. DNA sequencing of maternal plasma reliably identifies trisomy 18 and trisomy 13 as well as Down syndrome: an international collaborative study. *Genetics in Medicine* 2012;**14**(3):296-305. [PUBMED: 22281937]

Papageorghiou 2016a {published data only}

Papageorghiou AT, Khalil A, Forman M, Hulme R, Mazey R, Mousa HA, et al. Clinical evaluation of the IONA[®] test: a non-invasive prenatal screening test for trisomies 21, 18 and 13. *Ultrasound in Obstetrics & Gynecology* 2016;**47**(2):188-93. [PUBMED: 26493543]

Papageorghiou 2016b {published data only}

Papageorghiou AT, Khalil A, Forman M, Hulme R, Mazey R, Mousa HA, et al. Clinical evaluation of the IONA[®] test: a non-

invasive prenatal screening test for trisomies 21, 18 and 13. *Ultrasound in Obstetrics & Gynecology* 2016;**47**(2):188-93. [PUBMED: 26493543]

Pergament 2014 {published data only}

Pergament E, Cuckle H, Zimmermann B, Banjevic M, Sigurjonsson S, Ryan A, et al. Single-nucleotide polymorphism-based noninvasive prenatal screening in a high-risk and low-risk cohort. *Obstetrics and Gynecology* 2014;**124**(2 Pt 1):210-8. [PUBMED: 25004354]

Persico 2016 {published data only}

Persico N, Boito S, Ischia B, Cordisco A, De Robertis V, Fabietti I, et al. Cell-free DNA testing in the maternal blood in high-risk pregnancies after first-trimester combined screening. *Prenatal Diagnosis* 2016;**36**(3):232-6. [PUBMED: 26749576]

Poon 2016 {published data only}

Poon LC, Dumidrascu-Diris D, Francisco C, Fantasia I, Nicolaidis KH. IONA[®] test for first-trimester detection of trisomies 21, 18 and 13. *Ultrasound in Obstetrics & Gynecology* 2016;**47**(2):184-7. [PUBMED: 26387684]

Porreco 2014 {published and unpublished data}

NCT00847990. Non-Invasive screening for fetal aneuploidy. clinicaltrials.gov/show/NCT00847990 Date first received: 18 February 2009.

* Porreco RP, Garite TJ, Maurel K, Marusiak B, Ehrich M, van den Boom D, et al. Noninvasive prenatal screening for fetal trisomies 21, 18, 13 and the common sex chromosome aneuploidies from maternal blood using massively parallel genomic sequencing of DNA. *American Journal of Obstetrics and Gynecology* 2014;**211**(4):365.e1-12. [PUBMED: 24657131]

Quezada 2015 {published data only}

Quezada MS, Gil MM, Francisco C, Orosz G, Nicolaidis KH. Screening for trisomies 21, 18 and 13 by cell-free DNA analysis of maternal blood at 10-11 weeks' gestation and the combined test at 11-13 weeks. *Ultrasound in Obstetrics & Gynecology* 2015;**45**(1):36-41. [PUBMED: 25251385]

Samango-Sproue 2013 {published data only}

Samango-Sproue C, Banjevic M, Ryan A, Sigurjonsson S, Zimmermann B, Hill M, et al. SNP-based non-invasive prenatal testing detects sex chromosome aneuploidies with high accuracy. *Prenatal Diagnosis* 2013;**33**(7):643-9.

Sehnert 2011 {published data only}

Sehnert AJ, Rhee B, Comstock D, de Feo E, Heilek G, Burke J, et al. Optimal detection of fetal chromosomal abnormalities by massively parallel DNA sequencing of cell-free fetal DNA from maternal blood. *Clinical Chemistry* 2011;**57**(7):1042-9. [PUBMED: 21519036]

Shaw 2014 {published data only}

Shaw SW, Hsiao CH, Chen CY, Ren Y, Tian F, Tsai C, et al. Noninvasive prenatal testing for whole fetal chromosomal aneuploidies: a multicenter prospective cohort trial in Taiwan. *Fetal Diagnosis and Therapy* 2014;**35**(1):13-7. [PUBMED: 24296685]

Song 2013 {published data only}

Song Y, Liu C, Qi H, Zhang Y, Bian X, Liu J. Noninvasive prenatal testing of fetal aneuploidies by massively parallel sequencing in a prospective Chinese population. *Prenatal Diagnosis* 2013;**33**(7):700-6.

Song 2015 {published data only}

Song Y, Huang S, Zhou X, Jiang Y, Qi Q, Bian X, et al. Non-invasive prenatal testing for fetal aneuploidies in the first trimester of pregnancy. *Ultrasound in Obstetrics & Gynecology* 2015;**45**(1):55-60.

Sparks 2012a {published data only}

Sparks AB, Struble CA, Wang ET, Song K, Oliphant A. Noninvasive prenatal detection and selective analysis of cell-free DNA obtained from maternal blood: evaluation for trisomy 21 and trisomy 18. *American Journal of Obstetrics and Gynecology* 2012;**206**(4):319.e1-9. [PUBMED: 22464072]

Stumm 2014 {published data only}

Stumm M, Entezami M, Haug K, Blank C, Wustemann M, Schulze B, et al. Diagnostic accuracy of random massively parallel sequencing for non-invasive prenatal detection of common autosomal aneuploidies: a collaborative study in Europe. *Prenatal Diagnosis* 2014;**34**(2):185-91. [PUBMED: 24222400]

Sukhikh 2015 {published data only}

Sukhikh GT, Karetnikova NA, Shubina ES, Baranova EE, Korostin DO, Ekimov AN, et al. Noninvasive prenatal diagnosis of aneuploidies by next-generation sequencing (NGS) in a group of high-risk women. *Akusherstvo i Ginekologiya* 2015;**4**:5-10.

Sung-Hee 2015 {published data only}

Sung-Hee H, Young-Ho Y, Jae-Song R, Myung-Soo K, Young-Jin K, Kyoung-Ryul L. Noninvasive prenatal test for fetal chromosomal aneuploidies by massively parallel sequencing of cell-free fetal DNA in maternal plasma: The first clinical experience in Korea. *Journal of Genetic Medicine* 2015;**12**(2):85-91.

Tynan 2016 {published data only}

Tynan JA, Kim SK, Mazloom AR, Zhao C, McLennan G, Tim R, et al. Application of risk score analysis to low-coverage whole genome sequencing data for the noninvasive detection of trisomy 21, trisomy 18, and trisomy 13. *Prenatal Diagnosis* 2016;**36**(1):56-62. [PUBMED: 26505614]

Verweij 2013 {published data only}

Verweij EJ, Jacobsson B, van Scheltema PA, de Boer MA, Hoffer MJ, Hollemon D, et al. European non-invasive trisomy evaluation (EU-NITE) study: a multicenter prospective cohort study for non-invasive fetal trisomy 21 testing. *Prenatal Diagnosis* 2013;**33**(10):996-1001. [PUBMED: 23794121]

Wang 2014 {published data only}

Wang S, Gao Z, Lu Y, Li Y, Jiang S, Wang L, et al. Detection of fetal chromosomal aneuploidy in pregnant women at advanced maternal age during the first trimester. *Nan Fang Yi Ke da Xue Xue Bao [Journal of Southern Medical University]* 2014;**34**(5):655-8. [PUBMED: 24849430]

Wang 2015a {published data only}

Wang L, Meng Q, Tang X, Yin T, Zhang J, Yang S, et al. Maternal mosaicism of sex chromosome causes discordant sex chromosomal aneuploidies associated with noninvasive prenatal testing. *Taiwanese Journal of Obstetrics & Gynecology* 2015;**54**(5):527-31. [PUBMED: 26522104]

Yao 2014 {published data only}

Yao H, Jiang F, Hu H, Gao Y, Zhu Z, Zhang H, et al. Detection of fetal sex chromosome aneuploidy by massively parallel sequencing of maternal plasma DNA: initial experience in a Chinese hospital. *Ultrasound in Obstetrics & Gynecology* 2014;**44**(1):17-24. [PUBMED: 24616044]

Zhang 2016 {published data only}

Zhang J, Zhang B. Second-generation non-invasive high-throughput DNA sequencing technology in the screening of Down's syndrome in advanced maternal age women. *Biomedical Reports* 2016;**4**(6):715-8. [PUBMED: 27313855]

Zhou 2014a {published data only}

Zhou Q, Pan L, Chen S, Chen F, Hwang R, Yang X, et al. Clinical application of noninvasive prenatal testing for the detection of trisomies 21, 18, and 13: a hospital experience. *Prenatal Diagnosis* 2014;**34**(11):1061-5. [PUBMED: 24899146]

Zhou 2014b {published data only}

Zhou Q, Pan L, Chen S, Chen F, Hwang R, Yang X, et al. Clinical application of noninvasive prenatal testing for the detection of trisomies 21, 18, and 13: a hospital experience. *Prenatal Diagnosis* 2014;**34**(11):1061-5. [PUBMED: 24899146]

References to studies excluded from this review
Anderson 2015 {published data only}

Anderson B, Zhang K, Nguyen Q, Tsao D, Liu Y, Livingston K, et al. An automated, non-invasive prenatal screening assay (NIPS) for trisomy 21,18,13 in singleton and twin gestations. *International Journal of Gynecology and Obstetrics* 2015;**131**:E264.

Anselem 2016 {published data only}

Anselem O, Keroui S, Deput-Rampon C, Chartier M, Costa JM, Goffinet F, et al. Analysis of cell-free DNA in maternal blood for detection of fetal trisomy 21 in high-risk population: Couples acceptance and grounds for refusal [Étude de l'ADN foetal dans le sang maternel pour la détection de la trisomie 21 en population à risque accru : adhésion des couples et motifs de refus]. *Journal de Gynécologie, Obstétrique et Biologie de la Reproduction* 2016;**45**(8):918-23. [PUBMED: 26780844]

Bayindir 2015 {published data only}

Bayindir B, Dehaspe L, Brison N, Brady P, Ardui S, Kammoun M, et al. Noninvasive prenatal testing using a novel analysis pipeline to screen for all autosomal fetal aneuploidies improves pregnancy management. *European Journal of Human Genetics* 2015;**23**(10):1286-93. [PUBMED: 25585704]

Beamon 2013 {published data only}

Beamon C, Hardisty E, Harris S, Vora N. Promises and pitfalls of a new technology: a single center experience with noninvasive prenatal testing (NIPT). *American Journal of Obstetrics and Gynecology* 2013;**208**(1):S244-S5.

Beamon 2014 {published data only}

Beamon CJ, Hardisty EE, Harris SC, Vora NL. A single center's experience with noninvasive prenatal testing. *Genetics in Medicine* 2014;**16**(9):681-7. [PUBMED: 24675675]

Belloin 2016 {published data only}

Belloin C, Jacquemard F, Bernabe-Dupont C, Viot G, Lohmann L, Grange G. The noninvasive prenatal testing for Down's Syndrome. Retrospective study of 8821 patients [Le dépistage prénatal non invasif de la trisomie 21. Étude rétrospective à propos de 8821 patientes]. *Journal de Gynécologie, Obstétrique et Biologie de la Reproduction* 2016;**45**(9):1127-32. [PUBMED: 27091545]

Benachi 2015b {published data only}

Benachi A, Costa JM. Non invasive prenatal diagnosis of trisomy 21. *La Revue du Praticien* 2015;**65**(2):160-2. [PUBMED: 25939212]

Benachi 2016 {published data only}

Benachi A, Letourneau A, Kleinfinger P, Senat MV, Gautier E, Favre R, et al. Performance and indications of noninvasive prenatal testing using cell free circulating fetal DNA (cffDNA) for the detection of fetal trisomy 21, 18 and 13 in France [Performance et indication du dépistage des trisomies 21, 18 et 13 en France par l'analyse de l'ADN foetal dans le sang maternel.]. *Journal de Gynécologie, Obstétrique et Biologie de la Reproduction* 2016;**45**(6):633-40. [PUBMED: 26518155]

Benn 2015 {published data only}

Benn P. Re: Non-invasive prenatal testing for trisomies 21, 18 and 13: clinical experience from 146 958 pregnancies. Zhang H, Gao Y, Jiang F, Fu M, Yuan Y, Guo Y, et al. *Ultrasound in Obstetrics & Gynecology* 2015; **45**(5): 530-8.. *Ultrasound in Obstetrics & Gynecology* 2015;**45**(5):512-3. [PUBMED: 25914392]

Bhatt 2014 {published data only}

Bhatt S, Parsa S, Snyder H, Taneja P, Halks-Miller M, Seltzer W, et al. Clinical laboratory experience with noninvasive prenatal testing: Update on clinically relevant metrics. *Prenatal Diagnosis* 2014;**34**(Suppl 1):48.

Bianchi 2012a {published data only}

Bianchi D, Platt L, Goldberg J, Abuhamad A, Sehnert A, Rava R. Whole genome maternal plasma DNA sequencing accurately detects autosomal and sex chromosome aneuploidies. *Prenatal Diagnosis* 2012;**32**((Bianchi D.) Tufts Medical Center, Mother Infant Research Institute, Boston, United States):3-4.

Bianchi 2014b {published data only}

Bianchi DW, Lamar Parker R, Wentworth J, Madankumar R, Saffer C, Das AF, et al. DNA sequencing versus standard prenatal aneuploidy screening. *Obstetrical and Gynecological Survey* 2014;**69**(6):319-21. [EMBASE: 2014389701]

Bianchi 2014c {published data only}

Bianchi D, Swanson A, Parsa S, Bhatt S, Halks-Miller M, Sehnert A, et al. NIPT for sex chromosome aneuploidy: Initial clinical laboratory experience and biologic reasons for discordant results. *American Journal of Obstetrics and Gynecology* 2014;**210**(1):S87-S8.

Bianchi 2015a {published data only}

Bianchi DW, Chudova D, Sehnert AJ, Bhatt S, Murray K, Prosen TL, et al. Noninvasive prenatal testing and incidental detection of occult maternal malignancies. *JAMA* 2015;**314**(2):162-9. [PUBMED: 26168314]

Bianchi 2015b {published data only}

Bianchi DW, Parsa S, Bhatt S, Halks-Miller M, Kurtzman K, Sehnert AJ, et al. Fetal sex chromosome testing by maternal plasma DNA sequencing: clinical laboratory experience and biology. *Obstetrics and Gynecology* 2015;**125**(2):375-82.

Bianchi 2015c {published data only}

Bianchi DW, Chudova D, Sehnert AJ, Bhatt S, Murray K, Prosen TL, et al. Noninvasive prenatal testing and incidental detection of occult maternal malignancies. *Obstetrical and Gynecological Survey* 2015;**70**(12):744-6.

Bianchi 2015d {published data only}

Bianchi D, Chudova D, Sehnert A, Bhatt S, Murray K, Prosen T, et al. Incidental detection of occult maternal malignancies by noninvasive prenatal testing. *Prenatal Diagnosis* 2015;**35**:10-1.

BlueCross BlueShield Association 2014 {published data only}

BlueCross BlueShield Association. Noninvasive prenatal cell-free fetal DNA-based screening for aneuploidies other than trisomy 21. *Technology Evaluation Center Assessment Program Executive summary* 2014;**29**(7):1-7. [PUBMED: 25577816]

Brady 2016 {published data only}

Brady P, Brison N, Van Den Bogaert K, de Ravel T, Peeters H, Van Esch H, et al. Clinical implementation of NIPT - technical and biological challenges. *Clinical Genetics* 2016;**89**(5):523-30. [PUBMED: 25867715]

Chen 2013 {published data only}

Chen F, Zheng J, Chen M, Fang Q, Wei Y, Zhang C, et al. Noninvasive prenatal testing of Trisomy 21 and 18 by maternal plasma sequencing in twin pregnancies. *Prenatal Diagnosis* 2013;**33**:14.

Chen 2014 {published data only}

Chen M, Ma GC, Yeang CH. Genome-wide normalized score: A novel counting algorithm to detect fetal trisomies in non-invasive prenatal testing. *Prenatal Diagnosis* 2014;**34**:49.

Cherry 2014 {published data only}

Cherry AM, Williams JM Jr, Manning MA. Accuracy of non-invasive prenatal screening for trisomies 13, 18 and 21 and sex chromosome aneuploidy. *Cytogenetic and Genome Research* 2014;**142**(3):240-1.

Cheung 2015 {published data only}

Cheung SW, Patel A, Leung TY. Accurate description of DNA-based noninvasive prenatal screening. *New England Journal of Medicine* 2015; Vol. 372, issue 17:1675-7. [PUBMED: 25830325]

Chiu 2008 {published data only}

Chiu RW, Chan KC, Gao Y, Lau VY, Zheng W, Leung TY, et al. Noninvasive prenatal diagnosis of fetal chromosomal aneuploidy by massively parallel genomic sequencing of DNA in maternal plasma. *Proceedings of the National Academy of Sciences of the United States of America* 2008;**105**(51):20458-63. [PUBMED: 19073917]

Chiu 2010 {published data only}

Chiu RW, Sun H, Akolekar R, Clouser C, Lee C, McKernan K, et al. Maternal plasma DNA analysis with massively parallel sequencing by ligation for noninvasive prenatal diagnosis of trisomy 21. *Clinical Chemistry* 2010;**56**(3):459-63. [PUBMED: 20026875]

Christina 2012 {published data only}

Christina Fan H, Gu W, Wang J, Blumenfeld YJ, El-Sayed YY, Quake SR. Non-invasive prenatal measurement of the fetal genome. *Nature* 2012;**487**(7407):320-4. [EMBASE: 2012416711]

Cinnioglu 2012 {published data only}

Cinnioglu C. The application of bioinformatics to genetic testing for the detection of human aneuploidy and genotyping. *Reproductive BioMedicine Online* 2012;**24**:S37.

Cirigliano 2013 {published data only}

Cirigliano V, Ordonez E, Rueda L, Moreno M, Palao B, De La Paz Canadas M. Introduction of cfDNA based screening for common trisomies in Spain. *Prenatal Diagnosis* 2013;**33**:85-6.

Cirigliano 2014 {published data only}

Cirigliano V, Ordoñez E, Rueda L, Cañadas P, Moreno M, Palao B. cfDNA based aneuploidy screening in Spain: Results of one year clinical application. *Prenatal Diagnosis* 2014;**34**:51.

Cuckle 2015 {published data only}

Cuckle H, Benn P, Pergament E. Cell-free DNA screening for fetal aneuploidy as a clinical service. *Clinical Biochemistry* 2015;**48**(15):932-41. [PUBMED: 25732593]

Curnow 2014 {published data only}

Curnow K, Ryan A, Banjevic M, Shchegrova S, Babiarz J, Constantin T, et al. Observations following clinical implementation of a cfDNA and single-nucleotide polymorphism-based non-invasive prenatal aneuploidy test (NIPT). *Prenatal Diagnosis* 2014;**34**:51-2.

Dan 2012 {published data only}

Dan S, Wang W, Ren J, Li Y, Hu H, Xu Z, et al. Clinical application of massively parallel sequencing-based prenatal noninvasive fetal trisomy test for trisomies 21 and 18 in 11,105 pregnancies with mixed risk factors. *Prenatal Diagnosis* 2012;**32**(13):1225-32. [PUBMED: 23138752]

Dar 2014 {published data only}

Dar P, Curnow KJ, Gross SJ, Hall MP, Stosic M, Demko Z, et al. Clinical experience and follow-up with large scale single-nucleotide polymorphism-based noninvasive prenatal aneuploidy testing. *American Journal of Obstetrics and Gynecology* 2014;**211**(5):527.e1-527.e17. [PUBMED: 25111587]

De Ligt 2013 {published data only}

De Ligt J, Janssen I, Van Bon B, Buysse K, Gomes I, Eggink A, et al. Detecting partial fetal aneuploidies by MPS: An unexpected discrepancy between amniotic fluid and cffDNA. *Prenatal Diagnosis* 2013;**33**(Suppl 1):72-3. [ISSN 01973851]

Denona 2016 {published data only}

Denona B, Mone F, Cathcart B, Mahony R, Carroll S, Higgins S, et al. Changing rates of noninvasive prenatal testing and prenatal invasive testing in a tertiary centre. *BJOG: an international journal of obstetrics and gynaecology* 2016;**123**:75-6.

Discenza 2015 {published data only}

Discenza M, Ahern D, Coles K, Dobson L, Reiss R. Sex chromosome aneuploidy detection by NIPT: helpful or hazardous?. *Prenatal Diagnosis* 2015;**35**:105-6.

Dobson 2015 {published data only}

Dobson L, Reiff E, Little S, Bromley B, Wilkins-Haug L. Clinical course and patient outcomes following positive noninvasive prenatal testing (NIPT). *Prenatal Diagnosis* 2015;**35**:3.

Dobson 2016 {published data only}

Dobson LJ, Reiff ES, Little SE, Wilkins-Haug L, Bromley B. Patient choice and clinical outcomes following positive noninvasive prenatal screening for aneuploidy with cell-free DNA (cfDNA). *Prenatal Diagnosis* 2016;**36**(5):456-62. [PUBMED: 26938930]

Dong 2016 {published data only}

Dong Z, Zhang J, Hu P, Chen H, Xu J, Tian Q, et al. Low-pass whole-genome sequencing in clinical cytogenetics: a validated approach. *Genetics in Medicine* 2016;**18**(9):940-8. [PUBMED: 26820068]

Duenwald 2016 {published data only}

Duenwald S, Barbacioru C, Deciu C, Chen G, Comstock D, Skvortsov D, et al. Development of a novel paired-end sequencing-based noninvasive prenatal test. *American Journal of Obstetrics and Gynecology* 2016;**214**(1):S254-5.

Ehrich 2011a {published data only}

Ehrich M, Deciu C, Zwiefelhofer T, Tynan JA, Cagasan L, Tim R, et al. Noninvasive detection of fetal trisomy 21 by sequencing of DNA in maternal blood: a study in a clinical setting. *Obstetrical & Gynecological Survey* 2011;**66**(6):342-4.

Eiben 2014 {published data only}

Eiben B, Glaubitz R, Kagan KO. Non-invasive prenatal diagnostics: ETS and NGS-based tests. *Medizinische Genetik* 2014;**26**(4):382-90. [EMBASE: 2014948590]

Ellison 2015 {published data only}

Ellison C, Sun Y, Hogg G, Fox J, Tao H, McCarthy E, et al. Leveraging targeted sequencing of paired homologs for noninvasive detection of fetal aneuploidies. *Prenatal Diagnosis* 2015;**35**:54.

Faas 2011 {published data only}

Faas B, Vissers L, Janssen I, De Ligt J, Eggink A, Veltman J, et al. Multiplex massively parallel sequencing for noninvasive prenatal diagnosis. *Chromosome Research* 2011;**19**:S35-S6.

Faas 2012 {published data only}

Faas BH, de Ligt J, Janssen I, Eggink AJ, Wijnberger LD, van Vugt JM, et al. Non-invasive prenatal diagnosis of fetal aneuploidies using massively parallel sequencing-by-ligation and evidence that cell-free fetal DNA in the maternal plasma originates from cytotrophoblastic cells. *Expert Opinion on Biological Therapy* 2012;**12 Suppl 1**:S19-26. [PUBMED: 22500971]

Fairbrother 2013a {published data only}

Fairbrother G, Johnson S, Musci TJ, Song K. Clinical experience of noninvasive prenatal testing with cell-free DNA for fetal trisomies 21, 18, and 13, in a general screening population. *Prenatal Diagnosis* 2013;**33**(6):580-3. [PUBMED: 23494956]

Fairbrother 2013b {published data only}

Fairbrother G, Johnson S, Musci T, Song K. Clinical experience of Harmony™ Prenatal Test, a noninvasive prenatal test, in a general screening population. *Prenatal Diagnosis* 2013;**33**:78.

Fan 2008 {published data only}

Fan HC, Blumenfeld YJ, Chitkara U, Hudgins L, Quake SR. Noninvasive diagnosis of fetal aneuploidy by shotgun sequencing DNA from maternal blood. *Proceedings of the National Academy of Sciences of the United States of America* 2008;**105**(42):16266-71. [PUBMED: 18838674]

Fang 2015 {published data only}

Fang Y, Wang G, Wang C, Suo F, Gu M, Xia Y. The diagnosis pattern of mid-trimester fetal chromosomal aneuploidy in Xuzhou and the clinical applications. *Cell Biochemistry and Biophysics* 2015;**73**(2):267-70. [PUBMED: 25733332]

Ferres 2013 {published data only}

Ferres MA, Lichten L, Sachs A, Lau KM, Bianchi DW. Early experience with noninvasive DNA testing for aneuploidy in prenatal care. *Prenatal Diagnosis* 2013;**33**:73.

Fiorentino 2015 {published data only}

Fiorentino F, Spinella F, Bono S, Pizzuti F, Mariano M, Polverari A, et al. Feasibility of noninvasive prenatal testing for common fetal aneuploidies in maternal serum with low levels circulating fetal cell-free DNA fraction. *Prenatal Diagnosis* 2015;**35**:1.

Fosler 2015 {published data only}

Fosler L, Winters P, Jones K, Platt L. Clinical laboratory experience with noninvasive prenatal testing in twin gestations. *American Journal of Obstetrics and Gynecology* 2015;**212**(1):S330-S1.

Futch 2013 {published data only}

Futch T, Spinosa J, Bhatt S, de Feo E, Rava RP, Sehnert AJ. Initial clinical laboratory experience in noninvasive prenatal testing for fetal aneuploidy from maternal plasma DNA samples. *Prenatal Diagnosis* 2013;**33**(6):569-74.

Gabriel 2014 {published data only}

Gabriel H, Biskup S. Parallel prenatal testing for trisomies (-13,-18 and-21), turner syndrome, 22q11.2 del/dup-syndromes (e.g. DiGeorge), Smith-Lemli-Opitz syndrome and Noonan syndrome in fetuses with increased nuchal translucency using the AmpliSeq™ technology (Life Tech). *Medizinische Genetik* 2014;**26**(1):192-3. [ISSN 09365931]

Galea 2014 {published data only}

Galea M, Grehan S, Ives D, Evans K, Harraway J. A one-year experience of non-invasive prenatal testing in an Australian private pathology setting. *Prenatal Diagnosis* 2014;**34**:54.

Gao 2014 {published data only}

Gao Y, Xie B, Liu R. Delivering noninvasive prenatal testing in a clinical setting using semiconductor sequencing platform. *Science China Life Sciences* 2014;**57**(7):737-8. [PUBMED: 24969704]

Gao 2015 {published data only}

Gao Y, Zeng F, Fu M, Zhang H, Jiang F, Chen F, et al. Comparison of NIPT clinical performance in 72,382 high risk pregnant women and 40,287 low-risk pregnant women. *Prenatal Diagnosis* 2015;**35**:2.

Geifman-Holtzman 2013 {published data only}

Geifman-Holtzman O, Berman J, Ness AL, Cohen A, Fischer R, Carre A, et al. NIPDT registry-a critical utilization of non-invasive prenatal diagnosis test recently introduced to clinical practice. *American Journal of Obstetrics and Gynecology* 2013;**208**(1):S254.

Geifman-Holtzman 2014 {published data only}

Geifman-Holtzman O, Berman J, Xiong Y, Carre A, Ness A, Fang YMV, et al. Noninvasive prenatal testing (NIPT) registry patients' results and providers' perspective. *American Journal of Obstetrics and Gynecology* 2014;**210**(1):S96.

Gerundino 2017 {published data only}

Gerundino F, Giachini C, Contini E, Benelli M, Marseglia G, Giuliani C, et al. Validation of a method for noninvasive prenatal testing for fetal aneuploidies risk and considerations for its introduction in the Public Health System. *Journal of Maternal-Fetal & Neonatal Medicine* 2017;**30**(6):710-6. [PUBMED: 27226231]

Gil 2013 {published data only}

Gil MM, Quezada MS, Bregant B, Ferraro M, Nicolaidis KH. Implementation of maternal blood cell-free DNA testing in early screening for aneuploidies. *Ultrasound in Obstetrics & Gynecology* 2013;**42**(1):34-40. [PUBMED: 23744609]

Gil 2015 {published data only}

Gil MM, Giunta G, Macalli EA, Poon LC, Nicolaidis KH. UK NHS pilot study on cell-free DNA testing in screening for fetal

trisomies: factors affecting uptake. *Ultrasound in Obstetrics & Gynecology* 2015;**45**(1):67-73. [PUBMED: 25302655]

Gnetetskaya 2015 {published data only}

Gnetetskaya V, Kurtser M. Diagnosis of chromosomal aneuploidies using non-invasive prenatal test in Moscow. *Journal of Perinatal Medicine* 2015;**43**(s1):1300-64.

Grati 2014 {published data only}

Grati FR, Malvestiti F, Ferreira JC, Bajaj K, Gaetani E, Agrati C, et al. Fetoplacental mosaicism: potential implications for false-positive and false-negative noninvasive prenatal screening results. *Genetics in Medicine* 2014;**16**(8):620-4. [PUBMED: 24525917]

Gray 2013 {published data only}

Gray KJ, Wilkins-Haug L, Gerrol P, Reiss R. Introduction of cfDNA testing for aneuploidy screening: Indications, changes in invasive testing rates, and caveats. *Prenatal Diagnosis* 2013;**33**:74.

Gromminger 2014 {published data only}

Gromminger S, Yagmur E, Erkan S, Nagy S, Schock U, Bonnet J, et al. Fetal aneuploidy detection by cell-free DNA sequencing for multiple pregnancies and quality issues with vanishing twins. *Journal of Clinical Medicine* 2014;**3**(3):679-92. [PUBMED: 26237471]

Guex 2013 {published data only}

Guex N, Iseli C, Syngelaki A, Deluen C, Pescia G, Nicolaides KH, et al. A robust second-generation genome-wide test for fetal aneuploidy based on shotgun sequencing cell-free DNA in maternal blood. *Prenatal Diagnosis* 2013;**33**(7):707-10.

Halks-Miller 2015 {published data only}

Halks-Miller M, Chudova D, Bianchi DW. Maternal malignancies detected with noninvasive prenatal testing--reply. *JAMA* 2015; Vol. 314, issue 20:2192-3. [PUBMED: 26599192]

Harasim 2016 {published data only}

Harasim T, Wagner A, Rath S, Behaqi P, Rost I, Klein H-G. Prenatalis® NIPT: Accredited high resolution non-invasive prenatal testing by using massive parallel ultra-deep sequencing. *Medizinische Genetik* 2016;**28**(1):152-3.

Hernandez-Gomez 2015 {published data only}

Hernandez-Gomez M, Ramirez-Arroyo E, Melendez-Hernandez R, Garduno-Zaraza LM, Mayen-Molina DG. Non invasive prenatal test (NIPT) in maternal blood by parallel massive sequencing. Initial experience in Mexican women and literature review [Prueba prenatal no invasiva (NIPT) en sangre materna a traves de secuenciacion masiva paralela (MPS). Experiencia inicial en mujeres mexicanas y revision de la bibliografia]. *Ginecologia y Obstetricia de Mexico* 2015;**83**(5):277-88. [PUBMED: 26233973]

Hofmann 2013 {published data only}

Hofmann W, Entezami M, Haug K, Blank C, Wustemann M, Schulze B, et al. Diagnostic accuracy for the noninvasive prenatal detection of common autosomal aneuploidies. *Prenatal Diagnosis* 2013;**33**:75.

Hofmann 2014 {published data only}

Hofmann W, Grömminger S, Schöck U, Bonnet J, Smerdka P. Non-invasive prenatal testing (NIPT): Laboratory experiences of PrenaTest®. *Medizinische Genetik* 2014;**26**(1):194.

Hofmann 2015 {published data only}

Hofmann W, Grömminger S, Sachse M, Bonnet J, Schöck U. Recent bioinformatic advances of non-invasive prenatal detection to enhance diagnostic accuracy and aneuploidy discovery. *Prenatal Diagnosis* 2015;**35**:57.

Hu 2014 {published data only}

Hu Y, Duan H, Li J, Zhu R. Study of clinical indications for non-invasive prenatal testing based on distributions of fetal chromosomal abnormalities. *Prenatal Diagnosis* 2014;**34**:59.

Hu 2015 {published data only}

Hu HJ, Kwon YJ, Oh M, Kim J, Cho DY, Seo DH. Evaluating the results of the Momguard™ noninvasive prenatal test. *Journal of Genetic Medicine* 2015;**12**(2):96-9. [DOI: [10.5734/JGM.2015.12.2.96](https://doi.org/10.5734/JGM.2015.12.2.96)]

Hui 2015a {published data only}

Hui L, Teoh M, Da Silva Costa F, Ramsay P, Palma-Dias R, Richmond Z, et al. Clinical implementation of noninvasive prenatal testing by Australian sonologists. *BJOG: a international journal of obstetrics and gynaecology* 2015;**122**:52.

Hui 2015b {published and unpublished data}

Hui L, Teoh M, da Silva C, Ramsay P, Palma-Dias R, Richmond Z, et al. Clinical implementation of cell-free DNA-based aneuploidy screening: perspectives from a national audit. *Ultrasound in Obstetrics & Gynecology* 2015;**45**(1):10-5.

Jackson 2013 {published data only}

Jackson J, Dever K, Hamar B, Lazar E, Lim KH, Rodriguez D, et al. Nuchal translucency plus noninvasive prenatal testing screening for aneuploidy in a population of low-and high-risk patients. *Prenatal Diagnosis* 2013;**33**:22.

Jensen 2013 {published data only}

Jensen TJ, Zwiefelhofer T, Tim RC, Dzakula Z, Kim SK, Mazloom AR, et al. High-throughput massively parallel sequencing for fetal aneuploidy detection from maternal plasma. *PLOS One* 2013;**8**(3):e57381. [PUBMED: 23483908]

Jensen 2015 {published data only}

Jensen TJ, Deciu C, Ehrich M, Geis J, Kim SK, Tao H. Selective enrichment of genomic loci for the noninvasive detection of fetal aneuploidies. *Obstetrics and Gynecology* 2015;**125**:92S.

Jin 2014 {published data only}

Jin Y, Miao Z, Ge J, Zhang W, Li S, Liu X. Prenatal diagnosis of fetal chromosome aneuploidy by massively parallel genomic sequencing. *National Medical Journal of China* 2014;**94**(23):1788-90.

Johnson 2013 {published data only}

Johnson J, Pastuck M, Metcalfe A, Connors G, Krause R, Wilson D, et al. First-trimester Down syndrome screening using additional serum markers with and without nuchal

translucency and cell-free DNA. *Prenatal Diagnosis* 2013;**33**(11):1044-9.

Juneau 2014 {published data only}

Juneau K, Bogard PE, Huang S, Mohseni M, Wang ET, Ryvkin P, et al. Microarray-based cell-free DNA analysis improves noninvasive prenatal testing. *Fetal Diagnosis and Therapy* 2014;**36**(4):282-6.

Kagan 2015 {published data only}

Kagan KO, Schmid M, Hoopmann M, Wagner P, Abele H. Screening performance and costs of different strategies in prenatal screening for trisomy 21. *Geburtshilfe und Frauenheilkunde* 2015;**75**(3):244-50. [PUBMED: 25914417]

Kalantar 2014 {published data only}

Kalantar SM, Sitar G, Carabresy J. Is time for using non-invasive pre-natal diagnosis for genetic disorders?. *Iranian Journal of Reproductive Medicine* 2014;**12**(6):15.

Karlsson 2015 {published data only}

Karlsson K, Sahlin E, Iwarsson E, Westgren M, Nordenskjold M, Linnarsson S. Amplification-free sequencing of cell-free DNA for prenatal non-invasive diagnosis of chromosomal aberrations. *Genomics* 2015;**105**(3):150-8.

Kershberg 2015 {published data only}

Kershberg H, Greenberg J, Liou A, Logg A, Alvarado M, Natoli J, et al. Positive and negative predictive values of NIPT in an integrated health care system: summary of NIPT outcomes in Kaiser Permanente Southern California. *Prenatal Diagnosis* 2015;**35**:58.

Kinde 2012 {published data only}

Kinde I, Papadopoulos N, Kinzler KW, Vogelstein B. FAST-SeqS: a simple and efficient method for the detection of aneuploidy by massively parallel sequencing. *PLoS One* 2012;**7**(7):e41162. [PUBMED: 22815955]

Korabecna 2012 {published data only}

Korabecna M. Non invasive prenatal diagnostics of most frequent chromosomal aneuploidies in the language of numbers. *Aktualni Gynekologije a Porodnictvi* 2012;**4**(1):114-5.

Koumbaris 2016 {published data only}

Koumbaris G, Kypri E, Tsangaras K, Achilleos A, Mina P, Neofytou M, et al. Cell-free DNA analysis of targeted genomic regions in maternal plasma for non-invasive prenatal testing of trisomy 21, trisomy 18, trisomy 13, and fetal sex. *Clinical Chemistry* 2016;**62**(6):848-55. [PUBMED: 27117469]

Kurtser 2015 {published data only}

Kurtser MA, Gnetetskaya VA. A noninvasive prenatal test in the diagnosis of chromosome aneuploidies. *Akushersivo i Ginekologiya* 2015;**8**:65-9.

Lambert-Messerlian 2014 {published data only}

Lambert-Messerlian G, Kloza EM, Williams J 3rd, Loucky J, O'Brien B, Wilkins-Haug L, et al. Maternal plasma DNA testing for aneuploidy in pregnancies achieved by

assisted reproductive technologies. *Genetics in Medicine* 2014;**16**(5):419-22. [PUBMED: 24091801]

Larion 2015 {published data only}

Larion S, Warsof S, Romary L, Mlynarczyk M, Abuhamad A. Three year clinical experience with noninvasive prenatal testing in 3000 high risk cases in the United States. *Prenatal Diagnosis* 2015;**35**:59.

Lau 2012a {published data only}

Lau TK, Chan MK, Lo PS, Chan HY, Chan WS, Koo TY, et al. Clinical utility of noninvasive fetal trisomy (NIFTY™) test-early experience. *Journal of Maternal-fetal & Neonatal Medicine* 2012;**25**(10):1856-9. [PUBMED: 22471583]

Lau 2013 {published data only}

Lau TK, Jiang F, Chan MK, Zhang H, Lo PS, Wang W. Non-invasive prenatal screening of fetal Down syndrome by maternal plasma DNA sequencing in twin pregnancies. *Journal of Maternal-Fetal & Neonatal Medicine* 2013;**26**(4):434-7. [PUBMED: 23035860]

Lau 2014 {published data only}

Lau TK, Cheung SW, Lo PS, Pursley AN, Chan MK, Jiang F, et al. Non-invasive prenatal testing for fetal chromosomal abnormalities by low-coverage whole-genome sequencing of maternal plasma DNA: review of 1982 consecutive cases in a single center. *Ultrasound in Obstetrics & Gynecology* 2014;**43**(3):254-64. [PUBMED: 24339153]

Lebo 2015 {published data only}

Lebo RV, Novak RW, Wolfe K, Michelson M, Robinson H, Mancuso MS. Discordant circulating fetal DNA and subsequent cytogenetics reveal false negative, placental mosaic, and fetal mosaic cfDNA genotypes. *Journal of Translational Medicine* 2015;**13**:260. [PUBMED: 26260800]

Leung 2013 {published data only}

Leung TY, Qu JZZ, Liao GJW, Jiang P, Cheng YKY, Chan KCA, et al. Noninvasive twin zygoty assessment and aneuploidy detection by maternal plasma DNA sequencing. *Prenatal Diagnosis* 2013;**33**(7):675-81.

Levandoski 2015 {published data only}

Levandoski K, Marchand K, Briere L, Ferres M. One clinic's experience with discordant NIPT results for trisomies 18 and 13: Practical and psychosocial implications of non-invasive prenatal testing. *Prenatal Diagnosis* 2015;**35**:106-7.

Levy 2013 {published data only}

Levy B, Banjevic M, Hill M, Zimmermann B, Ryan A, Sigurjonsson S, et al. Targeted sequencing of SNPs results in highly accurate noninvasive detection of fetal aneuploidy of chromosomes 13, 18, 21, X, and Y: A validation study. *Human Reproduction* 2013;**28**:i26-i7.

Levy 2013a {published data only}

Levy B, Banjevic M, Hill M, Zimmermann B, Ryan A, Sigurjonsson S, et al. Targeted sequencing of single-nucleotide polymorphisms for accurate, noninvasive fetal aneuploidy detection at autosomal and sex chromosomes. *Prenatal Diagnosis* 2013;**33**:76.

Levy 2013b {published data only}

Levy, Ryan B, Banjevic A, Hill M, Zimmermann M, Sigurjonsson B, et al. Highly multiplexed single-nucleotide polymorphism amplification and sequencing to identify fetal aneuploidy from maternal cell-free DNA. *Journal of Perinatal Medicine* 2013;**41**(Suppl 1):34-218. [ISSN 03005577]

Li 2012 {published data only}

Li X, Sehnert A, Rava R. Percentage of fetal cell-free DNA in maternal plasma: Dependence on multiple clinical factors. *Prenatal Diagnosis* 2012;**32**:6.

Li 2015 {published data only}

Li WH, Wang PH, Chuang CM, Chang YW, Yang MJ, Chen CY, et al. Noninvasive prenatal testing for fetal trisomy in a mixed risk factors pregnancy population. *Taiwanese Journal of Obstetrics & Gynecology* 2015;**54**(2):122-5. [PUBMED: 25951714]

Liao 2011 {published data only}

Liao GJ, Lun FM, Zheng YW, Chan KC, Leung TY, Lau TK, et al. Targeted massively parallel sequencing of maternal plasma DNA permits efficient and unbiased detection of fetal alleles. *Clinical Chemistry* 2011;**57**(1):92-101. [PUBMED: 21078840]

Liao 2012 {published data only}

Liao GJ, Chan KC, Jiang P, Sun H, Leung TY, Chiu RW, et al. Noninvasive prenatal diagnosis of fetal trisomy 21 by allelic ratio analysis using targeted massively parallel sequencing of maternal plasma DNA. *PLOS One* 2012;**7**(5):e38154. [PUBMED: 22666469]

Liao 2013 {published data only}

Liao C, Fu YG, Huang SY, Fu F, Xie GE. Rapid noninvasive prenatal diagnosis of Down syndrome with Ion Proton™. *Prenatal Diagnosis* 2013;**33**:76-7.

Liao 2014 {published data only}

Liao C, Yin AH, Peng CF, Fu F, Yang JX, Li R, et al. Noninvasive prenatal diagnosis of common aneuploidies by semiconductor sequencing. *Proceedings of the National Academy of Sciences of the United States of America* 2014;**111**(20):7415-20. [PUBMED: 24799683]

Liao 2014a {published data only}

Liao C, Zhengfeng X, Zhang K. DNA sequencing versus standard prenatal aneuploidy screening. *New England Journal of Medicine* 2014; Vol. 371, issue 6:577-8. [PUBMED: 25099588]

Liu 2015 {published data only}

Liu J, Wang H, Xi H, Jia Z, Zhou Y, Wu L. Application of next-generation DNA sequencing for prenatal testing of fetal chromosomal aneuploidies. *Zhonghua Yi Xue Yi Chuan Xue za Zhi [Chinese Journal of Medical Genetics]* 2015;**32**(4):533-7. [PUBMED: 26252102]

Lo 2014 {published data only}

Lo KK, Boustred C, Chitty LS, Plagnol V. RAPIDR: an analysis package for non-invasive prenatal testing of aneuploidy. *Bioinformatics (Oxford, England)* 2014;**30**(20):2965-7. [PUBMED: 24990604]

Lo 2014a {published data only}

Lo K, Boustred C, McKay F, Fielding S, Plagnol V, Chitty L. A comparison of statistical methods for NIPT for aneuploidies and development of the RAPID-R package. *Prenatal Diagnosis* 2014;**34**:59-60.

Loucký 2013 {published data only}

Loucký J, Zemánek M. Non invasive prenatal testing of most frequent chromosomal aneuploidies - Some other aspects [Neinvazivní prenatalní testování nejčastějších chromozomálních aneuploidií - Některé další aspekty]. *Actual Gynecology and Obstetrics* 2013;**5**(1):6-7.

Louis-Jacques 2014 {published data only}

Louis-Jacques A, Burans C, Robinson S, Schofield E, Smulian J, Rochon M. Use of commercial tests for aneuploidy screening using cell-free fetal DNA in clinical practice. *Obstetrics and Gynecology* 2014;**123**(Suppl 1):154S.

Ma 2015 {published data only}

Ma J, Pan H, Fu J, Yu L, Yang H. Perspective study of non-invasive prenatal testing using cell-free fetal DNA in high-risk population. *Zhonghua Yi Xue za Zhi* 2015;**95**(11):849-52. [PUBMED: 26080919]

Ma 2015a {published data only}

Ma J, Yang H, Chen F, Wang Y. Nationwide evaluation on noninvasive prenatal testing through maternal plasma DNA sequencing under governmental regulation in China. *Prenatal Diagnosis* 2015;**35**:67.

Manotaya 2016 {published data only}

Manotaya S, Xu H, Uerpairojkit B, Chen F, Charoenvidhya D, Liu H, et al. Clinical experience from Thailand: noninvasive prenatal testing as screening tests for trisomies 21, 18 and 13 in 4736 pregnancies. *Prenatal Diagnosis* 2016;**36**(3):224-31. [PUBMED: 26748603]

Marchili 2015 {published data only}

Marchili JP, Chaulet A, Guetmonovitch O, Margulies ND, Solarz V. One clinic's experience with cell free fetal DNA testing in Argentina. *Prenatal Diagnosis* 2015;**35**:60.

Mayen 2015 {published data only}

Mayen DG, Hernandez-Gomez M, Ramirez E, Meléndez R, Garduño LM. Noninvasive prenatal test, one year of experience in the genetics clinic at the Hospital Angeles Lomas in Mexico City. *Prenatal Diagnosis* 2015;**35**:60-1.

Mazloom 2013a {published data only}

Mazloom A, Dzakula Z, Wang H, Oeth P, Jensen T, Tynan J, et al. Detection of fetal sex chromosomal aneuploidies by sequencing circulating cell-free DNA from maternal plasma. *Prenatal Diagnosis* 2013;**33**:77.

McCullough 2014 {published data only}

McCullough RM, Almasri EA, Guan X, Geis JA, Hicks SC, Mazloom AR, et al. Non-invasive prenatal chromosomal aneuploidy testing--clinical experience: 100,000 clinical samples. *PLOS One* 2014;**9**(10):e109173.

McCullough 2014a {published data only}

McCullough R, Almasri E, Guan X, Oeth P, Bombard A, Saldivar JS. Noninvasive prenatal testing: 100,000 patients clinical impact. *American Journal of Obstetrics and Gynecology* 2014;**210**(1):S58.

McCullough 2015 {published data only}

McCullough R, Almasri E, Boomer T, Wardrop J, Oeth P, Paxton W, et al. High volume clinical laboratory noninvasive prenatal testing: A synopsis (or summary) of results from >375,000 patients. *Prenatal Diagnosis* 2015;**35**:51-2.

McLennan 2016 {published data only}

McLennan A, Palma-Dias R, da Silva Costa F, Meagher S, Nisbet DL, Scott F. Noninvasive prenatal testing in routine clinical practice--an audit of NIPT and combined first-trimester screening in an unselected Australian population. *Australian & New Zealand Journal of Obstetrics & Gynaecology* 2016;**56**(1):22-8. [PUBMED: 26817523]

Meck 2014 {published data only}

Meck JM, Kramer DE, Aviram A, Trunca C, Riethmaier D, Pineda AD, et al. Non-invasive prenatal screening from viewpoint of the cytogenetics laboratory. *Cytogenetic and Genome Research* 2014;**142**(3):241.

Meck 2015 {published data only}

Meck JM, Kramer Dugan E, Matyakhina L, Aviram A, Trunca C, Pineda-Alvarez D, et al. Noninvasive prenatal screening for aneuploidy: positive predictive values based on cytogenetic findings. *American Journal of Obstetrics and Gynecology* 2015;**213**(2):214.e1-5. [PUBMED: 25843063]

Meck 2015a {published data only}

Meck J, Dugan EK, Pineda-Alvarez D, Wray A, Richard G, Matyakhina L. Comparison of results from noninvasive prenatal screening and diagnostic cytogenetic testing. *Prenatal Diagnosis* 2015;**35**:61.

Mennuti 2015 {published data only}

Mennuti MT, Chandrasekaran S, Khalek N, Dugoff L. Cell-free DNA screening and sex chromosome aneuploidies. *Prenatal Diagnosis* 2015;**35**(10):980-5. [PUBMED: 26088741]

Minarik 2015 {published data only}

Minarik G, Repiska G, Hyblova M, Nagyova E, Soltys K, Budis J, et al. Utilization of Benchtop Next Generation Sequencing Platforms Ion Torrent PGM and MiSeq in noninvasive prenatal testing for chromosome 21 trisomy and testing of impact of in silico and physical size selection on its analytical performance. *PLOS One* 2015;**10**(12):e0144811. [PUBMED: 26669558]

Miron 2011 {published data only}

Miron P. Prenatal screening for trisomy 21 and other aneuploidies at first trimester [Dépistage prénatal de la trisomie 21 et autres aneuploïdies au premier trimestre]. Université de Montréal (Faculté de Médecine) 2011:244 pages. [No AMICUS: 39273224]

Mundy 2008 {published data only}

Mundy L, Hiller JE. Non-invasive prenatal diagnostic test for Down's Syndrome (Structured abstract). Health Technology Assessment Database. Adelaide; Adelaide Health Technology Assessment (AHTA), 2008, issue 2. [HTA-32012000229]

Mundy 2009 {published data only}

Mundy L, Hiller JE. Non-invasive prenatal diagnostic test for trisomy-21 (Down's Syndrome) (Structured abstract). Health Technology Assessment Database. Adelaide Health Technology Assessment (AHTA) on behalf of National Horizon Scanning Unit (HealthPACT and MSAC), 2009, issue 2. [HTA-32010000766]

Musci 2014 {published data only}

Musci T. The largest study on NIPT thus far: The next study. *Journal of Maternal-fetal & Neonatal Medicine* 2014;**27**:16.

Musci 2014a {published data only}

Musci T, Struble C, Wang E, Hooks J, Syngelaki A, Del Mar Gil M, et al. Risk assessment for fetal sex chromosome aneuploidies using digital analysis of selected regions (DANSRTM) assays. *Prenatal Diagnosis* 2014;**34**:1-2.

NCT00770458 {unpublished data only}

NCT00770458. Non-invasive screening for fetal aneuploidy: a new maternal plasma marker. ClinicalTrials.gov/show/NCT00770458 Date first received: 8 October 2008.

NCT00877292 {unpublished data only}

NCT00877292. A new prenatal blood test for Down syndrome. ClinicalTrials.gov/show/NCT00877292 Date first received: 6 April 2009.

NCT00891852 {unpublished data only}

NCT00891852. Non-invasive determination of fetal chromosome abnormalities. ClinicalTrials.gov/show/NCT00891852 Date first received: 29 April 2009.

NCT00971334 {unpublished data only}

NCT00971334. Noninvasive screening for fetal aneuploidy: a new maternal plasma marker. ClinicalTrials.gov/show/NCT00971334 Date first received: 1 September 2009.

NCT01052688 {unpublished data only}

NCT01052688. Noninvasive screening for affected pregnancies: assay development and optimization in affected pregnancies. ClinicalTrials.gov/show/NCT01052688 Date first received: 15 January 2010.

NCT01256606 {unpublished data only}

NCT01256606. Prenatal test for fetal aneuploidy detection. ClinicalTrials.gov/show/NCT01256606 Date first received: 22 November 2010.

NCT01451671 {unpublished data only}

NCT01451671. Development of a prenatal test for fetal aneuploidy detection. ClinicalTrials.gov/show/NCT01451671 Date first received: 10 October 2011.

NCT01451684 {unpublished data only}

NCT01451684. Development of a non-invasive prenatal test. ClinicalTrials.gov/show/NCT01451684 Date first received: 10 October 2011.

NCT01555346 {unpublished data only}

NCT01555346. Clinical evaluation of the SEQureDx T21 test in high risk pregnancies. ClinicalTrials.gov/show/NCT01555346 Date first received: 15 February 2012.

NCT01574781 {unpublished data only}

NCT01574781. Non-invasive prenatal diagnostic validation study. ClinicalTrials.gov/show/NCT01574781 Date first received: 2 April 2012.

NCT01597063 {unpublished data only}

NCT01597063. Clinical evaluation of the SEQureDx Trisomy test in low risk pregnancies. ClinicalTrials.gov/show/NCT01597063 Date first received: 9 May 2012.

NCT01661010 {unpublished data only}

NCT01661010. The clinical study of sex chromosome variants. ClinicalTrials.gov/show/NCT01661010 Date first received: 7 August 2012.

NCT01663675 {unpublished data only}

NCT01663675. Trisomy 21 in adulthood. ClinicalTrials.gov/show/NCT01663675 Date first received: 8 August 2012.

NCT01668251 {unpublished data only}

NCT01668251. Turner syndrome prenatal diagnosis study. ClinicalTrials.gov/show/NCT01668251 Date first received: 3 November 2011.

NCT01725438 {unpublished data only}

NCT01725438. Non invasive prenatal diagnosis of trisomy 21 by genetic analysis of circulating fetal cells. ClinicalTrials.gov/show/NCT01725438 Date first received: 8 November 2012.

NCT01837979 {unpublished data only}

NCT01837979. Down syndrome screening based on dried blood spots and cell-free fetal DNA. ClinicalTrials.gov/show/NCT01837979 Date first received: 14 April 2013.

NCT01966991 {unpublished data only}

NCT01966991. Prenatal screening for Down syndrome with DNAFirst. ClinicalTrials.gov/show/NCT01966991 Date first received: 17 October 2013.

NCT02127515 {unpublished data only}

NCT02127515. Non invasive prenatal testing of Down syndrome. ClinicalTrials.gov/show/NCT02127515 Date first received: 2 April 2014.

NCT02226315 {unpublished data only}

NCT02226315. Clinical performance of the MaterniT21™ PLUS LDT in multiple gestation pregnancies. ClinicalTrials.gov/show/NCT02226315 Date first received: 20 August 2014.

NCT02872948 {unpublished data only}

NCT02872948. Diagnosis accuracy of noninvasive screening by PCR Digital for Down syndrome. ClinicalTrials.gov/show/NCT02872948 Date first received: 16 August 2016.

Neufeld-Kaiser 2015 {published data only}

Neufeld-Kaiser WA, Cheng EY, Liu YJ. Positive predictive value of non-invasive prenatal screening for fetal chromosome disorders using cell-free DNA in maternal serum: independent clinical experience of a tertiary referral center. *BMC Medicine* 2015;**13**:129. [PUBMED: 26033224]

Neveling 2015 {published data only}

Neveling K, Thung DT, Beulen L, Buijsman W, Gomes I, Van Den Heuvel S, et al. Clinical utility of NextSeq 500 sequencing for noninvasive prenatal testing. *Prenatal Diagnosis* 2015;**35**:54-5.

Nickolich 2016 {published data only}

Nickolich S, Farahi N, Jones K, Mounsey A. PURLs: Aneuploidy screening: Newer noninvasive test gains traction. *Journal of Family Practice* 2016;**65**(1):49-52. [PUBMED: 26845165]

Nicolaides 2013a {published data only}

Nicolaides K, Syngelaki A, Ashoor G, Musci T, Wang E, Song K. Clinical performance comparison of Harmony™ Prenatal Test and first-trimester combined screening in general pregnancy population. *Prenatal Diagnosis* 2013;**33**:3.

Nicolaides 2014 {published data only}

Nicolaides KH, Syngelaki A, Poon LC, Gil MM, Wright D. First-trimester contingent screening for trisomies 21, 18 and 13 by biomarkers and maternal blood cell-free DNA testing. *Fetal Diagnosis and Therapy* 2014;**35**(3):185-92. [PUBMED: 24192489]

Nicolaides 2014b {published data only}

Nicolaides KH, Musci TJ, Struble CA, Syngelaki A, Del Mar Gil M. Assessment of fetal sex chromosome aneuploidy using directed cell-free DNA analysis. *Obstetrical and Gynecological Survey* 2014;**69**(5):249-50.

Nicolaides 2014c {published data only}

Nicolaides KH, Syngelaki A, del Mar Gil M, Quezada MS, Zinevich Y. Prenatal detection of fetal triploidy from cell-free DNA testing in maternal blood. *Fetal Diagnosis and Therapy* 2014;**35**(3):212-7. [PUBMED: 24135152]

Norem 2015 {published data only}

Norem C, Obolensky E, Bijesse E, Turocy J, Blumberg B, Fehlen-Quizon P, et al. Non-invasive prenatal screening for trisomies-2 years experience in a large Health Maintenance Organization (HMO). *Prenatal Diagnosis* 2015;**35**:62.

Norton 2014 {published data only}

Norton ME, Jelliffe-Pawlowski LL, Currier RJ. Chromosome abnormalities detected by current prenatal screening and noninvasive prenatal testing. *Obstetrics and Gynecology* 2014;**124**(5):979-86.

Norton 2014a {published data only}

Norton ME, Jacobsson B, Swamy G, Laurent L, Ranzini A, Brar H, et al. Non-invasive EXamination of Trisomy using directed cell-free DNA analysis: The NEXT study. *Prenatal Diagnosis* 2014;**34**:e2.

Norton 2015a {published data only}

Norton M, Wapner R, Kuppermann M, Jelliffe-Pawłowski L, Currier R. Cell free DNA analysis vs sequential screening as primary testing considering all fetal chromosomal abnormalities. *American Journal of Obstetrics and Gynecology* 2015;**212**(1):S2.

Norton 2015b {published data only}

Norton ME, Jacobsson B, Swamy GK, Laurent LC, Ranzini AC, Brar H, et al. Cell-Free DNA analysis for noninvasive examination of trisomy. *Obstetrical & Gynecological Survey* 2015;**70**(8):483-4. [DOI: [10.1097/01.ogx.0000470657.58577.f2](https://doi.org/10.1097/01.ogx.0000470657.58577.f2)]

Norton 2015c {published data only}

Norton ME, Wapner RJ. Cell-free DNA analysis for noninvasive examination of trisomy. *New England Journal of Medicine* 2015; Vol. 373, issue 26:2582. [PUBMED: 26699179]

Norton 2016 {published data only}

Norton ME, Baer RJ, Wapner RJ, Kuppermann M, Jelliffe-Pawłowski LL, Currier RJ. Cell-free DNA vs sequential screening for the detection of fetal chromosomal abnormalities. *American Journal of Obstetrics and Gynecology* 2016;**214**(6):727.e1-6. [PUBMED: 26709085]

O'Leary 2014 {published data only}

O'Leary P, Maxwell S, Murch A, Dickinson J. What could non-invasive prenatal testing miss in highrisk pregnancies and would it change outcomes?. *Prenatal Diagnosis* 2014;**34**:52-3.

Oepkes 2015 {published data only}

Oepkes D, Schuring-Blom H, Pajkrt E, Faas B, Bax C, Coumans A, et al. TRIDENT: Or monitored NIPT implementation in the Netherlands. *Prenatal Diagnosis* 2015;**35**:10.

Oneda 2016 {published data only}

Oneda B, Steindl K, Masood R, Reshetnikova I, Krejci P, Baldinger R, et al. Noninvasive prenatal testing: More caution in counseling is needed in high risk pregnancies with ultrasound abnormalities. *European Journal of Obstetrics Gynecology and Reproductive Biology* 2016;**200**:72-5.

Ordoñez 2015 {published data only}

Ordoñez E, Rueda LR, Cañadas MP, Moreno M, Palao B, Cirigliano V. cfDNA-Based aneuploidy screening, the more the better?. *Prenatal Diagnosis* 2015;**35**:107.

Palomaki 2011 {published data only}

Palomaki GE, Kloza EM, Lambert-Messerlian GM, Haddow JE, Neveux LM, Ehrich M, et al. DNA sequencing of maternal plasma to detect Down syndrome: an international clinical validation study. *Genetics in Medicine* 2011;**13**(11):913-20. [PUBMED: 22005709]

Palomaki 2012a {published data only}

Palomaki GE, Kloza EM, Canick JA. Evidence that next-generation sequencing of maternal plasma will identify aneuploidies in a low risk population. *Prenatal Diagnosis* 2012;**32**:4.

Palomaki 2012b {published data only}

Palomaki GE, Kloza EM, Lambert-Messerlian GM, Haddow JE, Neveux LM, Ehrich M, et al. Editorial comment on DNA sequencing of maternal plasma to detect Down syndrome: an international clinical validation study. *Obstetrical and Gynecological Survey* 2012;**67**(2):86-8. [ISSN 0029-7828]

Palomaki 2015 {published data only}

Palomaki GE, Kloza EM, Lambert-Messerlian GM, van den Boom D, Ehrich M, Deciu C, et al. Circulating cell free DNA testing: Are some test failures informative?. *Prenatal Diagnosis* 2015;**35**(3):289-93.

Palomaki 2015a {published data only}

Palomaki GE, Kloza EM, Lambert-Messerlian GM, Van Den Boom D, Ehrich M, Deciu C, et al. Circulating cell free DNA testing: are some test failures informative?. *Obstetrical and Gynecological Survey* 2015;**70**(8):492-4.

Perez-Pedregosa 2015 {published data only}

Pérez-Pedregosa J, Paredes Ros B, Calles Hernandez LC, Izquierdo López L, Cabrillo Rodríguez E, Hurtado Caballero IV, et al. Non-invasive prenatal screening for aneuploidy through analysis of cell-free fetal DNA from maternal blood [Cribado prenatal no invasivo de aneuploidías mediante análisis de ADN fetal en sangre materna]. *Progresos de Obstetricia y Ginecología* 2015;**58**(3):113-7.

Pescia 2017 {published data only}

Pescia G, Guex N, Iseli C, Brennan L, Osteras M, Xenarios I, et al. Cell-free DNA testing of an extended range of chromosomal anomalies: clinical experience with 6,388 consecutive cases. *Genetics in Medicine* 2017;**19**(2):169-75. [PUBMED: 27362910]

Petersen 2014 {published data only}

Petersen OB, Vogel I, Ekelund C, Hyett J, Tabor A. Potential diagnostic consequences of applying non-invasive prenatal testing: population-based study from a country with existing first-trimester screening. *Ultrasound in Obstetrics & Gynecology* 2014;**43**(3):265-71.

Pettit 2014 {published data only}

Pettit KE, Hull AD, Korty L, Jones MC, Pretorius DH. The utilization of circulating cell-free fetal DNA testing and decrease in invasive diagnostic procedures: an institutional experience. *Journal of Perinatology* 2014;**34**(10):750-3. [PUBMED: 24875410]

Porreco 2014a {published data only}

Porreco RP. Reply. *American Journal of Obstetrics and Gynecology* 2014;**211**(6):712.

Rabinowitz 2012 {published data only}

Rabinowitz M, Gemelos G, Hill M, Demko Z, McAdoo S, Levy B. Noninvasive prenatal aneuploidy testing of chromosomes 13, 18, 21, X, and y. *Fertility and Sterility* 2012;**98**(3):S28-S9.

Rabinowitz 2012a {published data only}

Rabinowitz M, Gemelos G, Banjevic M, Zimmermann B, Baner J, Levy B, et al. Non-invasive detection of fetal aneuploidy across five chromosomes in the first trimester by targeted sequencing. *Prenatal Diagnosis* 2012;**32**:67.

Rabinowitz 2012b {published data only}

Rabinowitz M, Gemelos G, Banjevic M, Zimmermann B, Baner J, Levy B, et al. Highly accurate first trimester non-invasive detection of fetal aneuploidy of 13, 18, 21 and sex chromosomes by targeted sequencing. *Human Reproduction* 2012;**27**(Suppl 2):ii106-ii108. [ISSN 0268-1161]

Rabinowitz 2013 {published data only}

Rabinowitz M, Hill M, Demko Z, McAdoo S, Zimmermann B, Levy B. Using targeted sequencing of SNPs to achieve a highly accurate non-invasive detection of fetal aneuploidy of 13,18, 21 and sex chromosomes. *American Journal of Obstetrics and Gynecology* 2013;**208**(1):S253.

Rabinowitz 2014 {published data only}

Rabinowitz M, Valenti E, Pettersen B, Sigurjonsson S, Hill M, Zimmermann B. Noninvasive aneuploidy detection by multiplexed amplification and sequencing of polymorphic loci. *Obstetrics and Gynecology* 2014;**123** Suppl 1(5):167S-S.

Rad 2014 {published data only}

Rad S, Turner AL, Beauchamp S, Aghajanian P, Williams J 3rd, Esakoff TF. Noninvasive prenatal testing compared with invasive diagnostic testing in the setting of an abnormal state aneuploidy screen. *Obstetrics and Gynecology* 2014;**123** Suppl 1:100s.

Radoi 2015 {published data only}

Radoi VE, Bohiltea CL, Bohiltea RE, Albu DN. Cell free fetal DNA testing in maternal blood of Romanian pregnant women. *Iranian Journal of Reproductive Medicine* 2015;**13**(10):623-6. [PUBMED: 26644790]

Rava 2012 {published data only}

Rava R, Bianchi D, Platt L, Goldberg J, Abuhamad A, Sehnert A. Genome wide fetal aneuploidy detection by sequencing of maternal plasma DNA: diagnostic accuracy in a prospective, blinded, multicenter study. *American Journal of Obstetrics and Gynecology* 2012;**206**(1):S367-S.

Rava 2014 {published data only}

Rava RP, Srinivasan A, Sehnert AJ, Bianchi DW. Circulating fetal cell-free DNA fractions differ in autosomal aneuploidies and monosomy X. *Clinical Chemistry* 2014;**60**(1):243-50.

Reiff 2015 {published data only}

Reiff E, Little S, Dobson L, Agarwala V, Wilkins-Haug L, Bromley B. The role of first trimester ultrasound in prenatal aneuploidy screening for women with a negative cell free fetal DNA. *Prenatal Diagnosis* 2015;**35**:16-7.

Reiff 2016 {published data only}

Reiff ES, Little SE, Dobson L, Wilkins-Haug L, Bromley B. What is the role of the 11- to 14-week ultrasound in women with

negative cell-free DNA screening for aneuploidy?. *Prenatal Diagnosis* 2016;**36**(3):260-5. [PUBMED: 26748490]

Reimers 2015 {published data only}

Reimers R, Mason-Suares H, Little S, Reiff E, Bromley B, Dobson L, et al. What will be missed if Noninvasive Prenatal Testing (NIPT) is the only assessment for "Ultrasound Indicated" genetic studies?. *Prenatal Diagnosis* 2015;**35**:63-4.

Revello 2016 {published data only}

Revello R, Sarno L, Ispas A, Akolekar R, Nicolaides KH. Screening for trisomies by cell-free DNA testing of maternal blood: consequences of a failed result. *Ultrasound in Obstetrics & Gynecology* 2016;**47**(6):698-704. [PUBMED: 26743020]

Ryan 2016 {published data only}

Ryan A, Hunkapiller N, Banjevic M, Vankayalapati N, Fong N, Jinnett KN, et al. Validation of an enhanced version of a single-nucleotide polymorphism-based noninvasive prenatal test for detection of fetal aneuploidies. *Fetal Diagnosis and Therapy* 2016;**40**:219-23. [PUBMED: 27028530]

Sachse 2015 {published data only}

Sachse M, Grömminger S, Schöck U, Bonnet J, Hofmann W. Development of a non-invasive prenatal test (NIPT) assay for trisomy 21 (T21) based on comparative quantification of chromosome 21 and a reference chromosome via quantitative real-time PCR. *Medizinische Genetik* 2015;**27**(1):177.

Samura 2015 {published data only}

Samura O, Sago H, Sekizawa A. Nationwide project using next-generation sequencing of cellfree DNA in maternal plasma in Japan: One and a half years of experience. *Prenatal Diagnosis* 2015;**35**:64.

Sarno 2016 {published data only}

Sarno L, Revello R, Hanson E, Akolekar R, Nicolaides KH. Prospective first-trimester screening for trisomies by cell-free DNA testing of maternal blood in twin pregnancy. *Ultrasound in Obstetrics & Gynecology* 2016;**47**(6):705-11. [PUBMED: 26970114]

Schöck 2015 {published data only}

Schöck U, Grömminger S, Bonnet J, Sachse M, Hofmann W. Recent bioinformatic advances of non-invasive prenatal detection to enhance diagnostic accuracy and aneuploidy discovery. *Medizinische Genetik* 2015;**27**(1):177.

Sehnert 2013 {published data only}

Sehnert A, Futch T, Seltzer W, Bianchi D, Rava R. Clinical laboratory experience using noninvasive prenatal testing for fetal aneuploidy in late gestation. *Prenatal Diagnosis* 2013;**33**:80-1.

Sehnert 2014 {published data only}

Sehnert A, Bianchi D, Rava R. Maternal cell-free DNA (cfDNA) sequencing versus standard prenatal aneuploidy screening in a general obstetrical population. *Prenatal Diagnosis* 2014;**34**:8.

Sentilhes 2015 {published data only}

Sentilhes L, Salomon LJ, Vayssiere C. Cell-free DNA analysis for noninvasive examination of trisomy. *New England Journal of Medicine* 2015; Vol. 373, issue 26:2581-2. [PUBMED: 26699181]

Seo 2015 {published data only}

Seo DH, Cho DY, Kim J, Kim SY, Cho SE, Oh M. Clinical study of non-invasive prenatal testing using next-generation sequencing. *Journal of Laboratory Medicine and Quality Assurance* 2015;**37**:214-8. [DOI: [10.15263/jlmqa.2015.37.4.214](https://doi.org/10.15263/jlmqa.2015.37.4.214)]

Settler 2015 {published data only}

Settler C, Dohany L. Clinical laboratory experience in a general obstetrical population of noninvasive prenatal screening for fetal aneuploidy from cell free DNA. *Prenatal Diagnosis* 2015;**35**:64-5.

Shani 2016 {published data only}

Shani H, Goldwaser T, Keating J, Klugman S. Chromosomal abnormalities not currently detected by cell-free fetal DNA: a retrospective analysis at a single center. *American Journal of Obstetrics and Gynecology* 2016;**214**(6):729.e1-729.e11. [PUBMED: 26721783]

Shaohua 2012 {published data only}

Shaohua T, Huanzheng L, Erle Z, Xueqing X. Noninvasive prenatal diagnosis for aneuploidy by massively parallel genomic sequencing of DNA in maternal plasma. *Prenatal Diagnosis* 2012;**32**:79.

Sharma 2015 {published data only}

Sharma P, Metcalfe A, Pastuck M, Laberge A-M, Haidar H, Ravitsky V, et al. Women's understanding of non-invasive prenatal testing based on cell free DNA versus first trimester combined screening. *Prenatal Diagnosis* 2015;**35**:106.

Shaw 2013 {published data only}

Shaw SWS, Chen CY, Hsiao CH, Ren Y, Tian F, Tsai C, et al. Non-invasive prenatal testing for whole fetal chromosomal aneuploidies: A multi-center prospective cohort trial in Taiwan. *Prenatal Diagnosis* 2013;**33**:81.

Shen 2016 {published data only}

Shen J, Wen Z, Qin X, Shi Y. Noninvasive fetal trisomy detection by multiplexed semiconductor sequencing: a barcoding analysis strategy. *Journal of Human Genetics* 2016;**61**(3):247-52. [PUBMED: 26657936]

Shi 2015 {published data only}

Shi X, Zhang Z, Cram DS, Liu C. Feasibility of noninvasive prenatal testing for common fetal aneuploidies in an early gestational window. *Clinica Chimica Acta* 2015;**439**:24-8.

Shulman 2014 {published data only}

Shulman L, Dungan J, Ginsberg N. The use of noninvasive prenatal screening (NIPS) in the assessment of an abnormal fetal ultrasound. *Prenatal Diagnosis* 2014;**34**:62.

Sistermans 2015a {published data only}

Sistermans E, Straver R, Faas BH. Maternal malignancies detected with noninvasive prenatal testing. *JAMA* 2015; Vol. 314, issue 20:2192. [PUBMED: 26599191]

Smith-Bindman 2015 {published data only}

Smith-Bindman R, Miglioretti D. Cell-free DNA analysis for noninvasive examination of trisomy. *New England Journal of Medicine* 2015; Vol. 373, issue 26:2581. [PUBMED: 26699180]

Song 2012 {published data only}

Song K, Ashoor G, Syngelaki A, Wagner M, Birdir C, Struble C, et al. Clinical evaluation of a directed cfDNA analysis method for non-invasive prenatal fetal trisomy detection. *Prenatal Diagnosis* 2012;**32**:16-7.

Sparks 2012 {published data only}

Sparks AB, Wang ET, Struble CA, Barrett W, Stokowski R, McBride C, et al. Selective analysis of cell-free DNA in maternal blood for evaluation of fetal trisomy. *Prenatal Diagnosis* 2012;**32**(1):3-9. [PUBMED: 2223233]

Srinivasan 2013 {published data only}

Srinivasan A, Bianchi D, Liao W, Sehnert A, Rava R. Maternal plasma DNA sequencing: Effects of multiple gestation on aneuploidy detection and the relative cell-free fetal DNA (cffDNA) per fetus. *American Journal of Obstetrics and Gynecology* 2013;**208**(1):S31.

Stokowski 2015 {published data only}

Stokowski R, Wang E, White K, Batey A, Jacobsson B, Brar H, et al. Clinical performance of non-invasive prenatal testing (NIPT) using targeted cell-free DNA analysis in maternal plasma with microarrays or next generation sequencing (NGS) is consistent across multiple controlled clinical studies. *Prenatal Diagnosis* 2015;**35**(12):1243-6. [PUBMED: 26332378]

Strah 2015 {published data only}

Strah D, Ovniček P, Bernik J. Non-invasive prenatal cell-free fetal DNA testing for down syndrome and other chromosomal abnormalities [Neinvazivno predrojstveno testiranje prostih plodov DNA za downov sindrom in ostale kromosomske nepravilnosti]. *Zdravniški Vestnik* 2015;**84**(11):727-33.

Straver 2014 {published data only}

Straver R, Sistermans EA, Holstege H, Visser A, Oudejans CBM, Reinders MJT. WISECONDOR: Detection of fetal aberrations from shallow sequencing maternal plasma based on a within-sample comparison scheme. *Nucleic Acids Research* 2014;**42**(5):e31.

Strom 2015 {published data only}

Strom C. The positive predictive value of noninvasive prenatal screening (NIPS). *Chromosome Research* 2015;**23**(1):S23.

Stumm 2011 {published data only}

Stumm M, Trunk N, Beck M, Entezami M, Becker R, Locherbach J, et al. Non-invasive prenatal detection of chromosome aneuploidies using next-generation sequencing: First steps towards clinical application. *Medizinische Genetik* 2011;**23**(1):86.

Stumm 2012 {published data only}

Stumm M, Entezami M, Trunk N, Beck M, Locherbach J, Wegner RD, et al. Noninvasive prenatal detection of chromosomal aneuploidies using different next generation sequencing strategies and algorithms. *Prenatal Diagnosis* 2012;**32**(6):569-77. [PUBMED: 22573401]

Stumm 2012a {published data only}

Stumm M, Entezami M, Haug K, Blank C, Wustemann M, Schulze B, et al. Non-invasive prenatal detection of trisomy 21 using massively parallel sequencing: A collaborative study in Europe. *Prenatal Diagnosis* 2012;**32**:63-4.

Stumm 2013 {published data only}

Stumm M, Entezami M, Haug K, Blank C, Wustemann C, Schulze B, et al. Diagnostic accuracy of PrenaTest® for non-invasive prenatal detection of common autosomal aneuploidies. *Medizinische Genetik* 2013;**25**(1):171-2.

Stumm 2016 {published data only}

Stumm M. Noninvasive prenatal tests from the viewpoint of human geneticists: Possibilities and limits in daily routine [Nichtinvasive pränatale Tests aus Sicht des Humangenetikers: Möglichkeiten und Grenzen in der täglichen Routine]. *Der Gynäkologe* 2016;**49**(6):429-36.

Swanson 2012 {published data only}

Swanson A, Coffeen C, Sehnert AJ. Non-invasive prenatal testing for fetal aneuploidy by massively parallel DNA sequencing of maternal plasma: The future has arrived today. *Laboratoriums Medizin* 2012;**36**(5):269-75.

Syngelaki 2014 {published data only}

Syngelaki A, Pergament E, Homfray T, Akolekar R, Nicolaidis KH. Replacing the combined test by cell-free DNA testing in screening for trisomies 21, 18 and 13: impact on the diagnosis of other chromosomal abnormalities. *Fetal Diagnosis and therapy* 2014;**35**(3):174-84. [PUBMED: 24525399]

Tan 2016 {published data only}

Tan Y, Gao Y, Lin G, Fu M, Li X, Yin X, et al. Noninvasive prenatal testing (NIPT) in twin pregnancies with treatment of assisted reproductive techniques (ART) in a single center. *Prenatal Diagnosis* 2016;**36**(7):672-9. [PUBMED: 27150972]

Taneja 2016 {published data only}

Taneja PA, Snyder HL, de Feo E, Kruglyak KM, Halks-Miller M, Curnow KJ, et al. Noninvasive prenatal testing in the general obstetric population: clinical performance and counseling considerations in over 85 000 cases. *Prenatal Diagnosis* 2016;**36**(3):237-43. [PUBMED: 26715197]

Taneja 2017 {published data only}

Taneja PA, Prosen TL, de Feo E, Kruglyak KM, Halks-Miller M, Curnow KJ, et al. Fetal aneuploidy screening with cell-free DNA in late gestation. *Journal of Maternal-Fetal & Neonatal Medicine* 2017;**30**(3):338-42. [PUBMED: 27124739]

Tarrier 2015 {published data only}

Tarrier B, Sekedat M, Mann T, Stoerker J, Bombard A. Whole genome maternal plasma DNA sequencing for fetal

aneuploidy accurately detects autosomal and sex chromosome abnormalities: A validation study with independent fetal fraction observations. *Prenatal Diagnosis* 2015;**35**:65-6.

Taylor 2014 {published data only}

Taylor JB, Chock VY, Hudgins L. NIPT in a clinical setting: an analysis of uptake in the first months of clinical availability. *Journal of Genetic Counseling* 2014;**23**(1):72-8.

Togneri 2016 {published data only}

Togneri F, Court S, Parks M, Clokie S, Hamilton S, Bibb N, et al. Noninvasive prenatal testing for fetal aneuploidy: The experience of an NHS regional genetics laboratory. *BJOG: an international journal of obstetrics and gynaecology* 2016;**123**:70.

Tong 2016 {published data only}

Tong H, Jin Y, Xu Y, Zou B, Ye H, Wu H, et al. Prenatal diagnosis of trisomy 21, 18 and 13 by quantitative pyrosequencing of segmental duplications. *Clinical Genetics* 2016;**90**(5):451-5. [PUBMED: 26948280]

Valderramos 2016a {published data only}

Valderramos SG, Rao R, Scibetta EW, Clark K, Krakow D, Silverman NS, et al. Clinical accuracy of abnormal cell-free fetal DNA (cfDNA) results for the sex chromosomes. *American Journal of Obstetrics and Gynecology* 2016;**214**(1):S402-3.

Valderramos 2016b {published data only}

Valderramos SG, Rao R, Scibetta EW, Clark K, Krakow D, Silverman NS, et al. Clinical accuracy of abnormal autosomal cell-free fetal DNA (cfDNA) screening. *American Journal of Obstetrics and Gynecology* 2016;**214**(1):S330.

Valderramos 2016c {published data only}

Valderramos SG, Rao RR, Scibetta EW, Silverman NS, Han CS, Platt LD. Cell-free DNA screening in clinical practice: abnormal autosomal aneuploidy and microdeletion results. *American Journal of Obstetrics and Gynecology* 2016;**215**(5):626.e1-626.e10. [PUBMED: 27371353]

van den Oever 2012a {published data only}

van den Oever JM, Balkassmi S, Verweij EJ, van Iterson M, Adama van Scheltema PN, Oepkes D, et al. Single molecule sequencing of free DNA from maternal plasma for noninvasive trisomy 21 detection. *Clinical Chemistry* 2012;**58**(4):699-706. [PUBMED: 22278607]

van den Oever 2012b {published data only}

Van Den Oever JME, Balkassmi S, Verweij EJ, Van Iterson M, Van Scheltema PNA, Oepkes D, et al. Single molecule sequencing of free DNA from maternal plasma for non-invasive trisomy 21 detection. *Prenatal Diagnosis* 2012;**32**:60-1.

van den Oever 2013 {published data only}

van den Oever JM, Balkassmi S, Johansson LF, Adama van Scheltema PN, Suijkerbuijk RF, Hoffer MJ, et al. Successful noninvasive trisomy 18 detection using single molecule sequencing. *Clinical Chemistry* 2013;**59**(4):705-9. [PUBMED: 23315481]

Van Opstal 2016 {published data only}

Van Opstal D, Srebnik MI, Polak J, de Vries F, Govaerts LC, Joosten M, et al. False negative NIPT results: risk figures for chromosomes 13, 18 and 21 based on chorionic villi results in 5967 cases and literature review. *PLOS One* 2016;**11**(1):e0146794. [PUBMED: 26771677]

Verweij 2013a {published data only}

Verweij EJ, Jacobsson B, Van Scheltema PNA, De Boer MA, Hoffer MJV, Hollemon D, et al. European Noninvasive Trisomy Evaluation (EU-NITE) study: Multicenter prospective study for noninvasive fetal trisomy 21 testing. *Prenatal Diagnosis* 2013;**33**:83-4.

Wald 2015a {published data only}

Wald NJ, Bestwick JP. Performance of antenatal reflex DNA screening for Down's syndrome. *Journal of Medical Screening* 2015;**22**(4):168-74. [PUBMED: 25882667]

Wald 2015b {published data only}

Wald NJ, Huttly WJ, Bestwick JP, Aquilina J, Peregrine E. Reflex antenatal DNA screening for Down syndrome. *Prenatal Diagnosis* 2015; Vol. 35, issue 11:1154. [PUBMED: 26211510]

Wang 2012 {published data only}

Wang SJ, Gao ZY, Lu YP, Li YL, You YQ, Zhang LW, et al. Value of detection of cell-free fetal DNA in maternal plasma in the prenatal diagnosis of chromosomal abnormalities. *Zhonghua Fu Chan Ke za Zhi* 2012;**47**(11):808-12. [PUBMED: 23302119]

Wang 2015b {published data only}

Wang JC, Sahoo T, Schonberg S, Kopita KA, Ross L, Patek K, et al. Discordant noninvasive prenatal testing and cytogenetic results: a study of 109 consecutive cases. *Genetics in Medicine* 2015;**17**(3):234-6. [PUBMED: 25101914]

Wang 2015c {published data only}

Wang S, Huang S, Ma L, Liang L, Zhang J, Zhang J, et al. Maternal X chromosome copy number variations are associated with discordant fetal sex chromosome aneuploidies detected by noninvasive prenatal testing. *Clinica Chimica Acta; International Journal of Clinical Chemistry* 2015;**444**:113-6. [PUBMED: 25689220]

Wang 2015d {published data only}

Wang JC, Sahoo T, Schonberg S, Kopita KA, Ross L, Patek K, et al. Discordant noninvasive prenatal testing and cytogenetic results: A study of 109 consecutive cases. *Obstetrical and Gynecological Survey* 2015;**70**(7):434-6.

Wang 2015e {published data only}

Wang E, Struble C, Kingsley C, Steeke R, Batey A, Hollemon D, et al. Importance of fetal fraction analysis for CFDNA testing in the general pregnancy population. *Reproductive Sciences* 2015;**22**:176A.

Xiong 2015 {published data only}

Xiong Y, Berman J, Weiner S, Carre A, Seligman NS, Steinbach G, et al. Noninvasive prenatal testing (NIPT) registry-patients' results and providers' perspective. *Reproductive Sciences* 2015;**22**:176A-7A.

Yankova 2015 {published data only}

Yankova M, Chaveeva P, Stratieva V. Models of clinical implementation of cell free fetal DNA in the maternal serum screening test-analysis. *Akusherstvo i Ginekologija* 2015;**54**(7):15-21. [PUBMED: 27025103]

Yaron 2015 {published data only}

Yaron Y, Jani J, Schmid M, Oepkes D. Current status of testing for microdeletion syndromes and rare autosomal trisomies using cell-free DNA technology. *Obstetrics and Gynecology* 2015;**126**(5):1095-9. [PUBMED: 26444108]

Yeang 2014 {published data only}

Yeang CH, Ma GC, Hsu HW, Lin YS, Chang SM, Cheng PJ, et al. Genome-wide normalized score: a novel algorithm to detect fetal trisomy 21 during non-invasive prenatal testing. *Ultrasound in Obstetrics & Gynecology* 2014;**44**(1):25-30. [PUBMED: 24700679]

Yu 2014 {published data only}

Yu SC, Chan KC, Zheng YW, Jiang P, Liao GJ, Sun H, et al. Size-based molecular diagnostics using plasma DNA for noninvasive prenatal testing. *Proceedings of the National Academy of Sciences of the United States of America* 2014;**111**(23):8583-8. [PUBMED: 24843150]

Yuan 2013 {published data only}

Yuan Y, Jiang F, Hua S, Du B, Hao Y, Ye L, et al. Feasibility study of semiconductor sequencing for noninvasive prenatal detection of fetal aneuploidy. *Clinical Chemistry* 2013;**59**(5):846-9. [PUBMED: 23364181]

Zhang 2015 {published data only}

Zhang H, Gao Y, Jiang F, Fu M, Yuan Y, Guo Y, et al. Non-invasive prenatal testing for trisomies 21, 18 and 13: clinical experience from 146,958 pregnancies. *Ultrasound in Obstetrics & Gynecology* 2015;**45**(5):530-8. [PUBMED: 25598039]

Zhou 2013 {published data only}

Zhou D, Liang D, Lv W, Tian F, Song Z, Zhang J, et al. Noninvasive prenatal testing in China: Results and learning from a clinical study of more than 50,000 pregnancies. *Prenatal Diagnosis* 2013;**33**:1.

Zimmermann 2012 {published data only}

Zimmermann B, Hill M, Gemelos G, Demko Z, Banjevic M, Baner J, et al. Noninvasive prenatal aneuploidy testing of chromosomes 13, 18, 21, X, and Y, using targeted sequencing of polymorphic loci. *Prenatal Diagnosis* 2012;**32**(13):1233-41. [PUBMED: 23108718]

Zimmermann 2013 {published data only}

Zimmermann B, Banjevic M, Hill M, Lacroute P, Dodd M, Sigurjonsson S, et al. Highly multiplexed single-nucleotide polymorphism amplification and sequencing to identify fetal aneuploidy from maternal cell-free DNA. *Prenatal Diagnosis* 2013;**33**:68.

Zwiefelhofer 2013 {published data only}

Zwiefelhofer T, Whitley P, Roy K, Jean-Jacques R, Ehrich M. Prenatal detection of fetal aneuploidy using bench top sequencing. *Prenatal Diagnosis* 2013;**33**:85.

References to ongoing studies
Basaran 2015 {published and unpublished data}

Basaran S, Yuksel A, Has R, Kirgiz M, Dehgan T, Karaman B. False positive and false negative results of cell free DNA testing. *Chromosome Research* 2015;**23**(1):S124-S5.

Buresch 2016 {published data only}

Buresch A, Rosner M, Suskin B, Einstein F, Bircaj E, Lister R, et al. Actual rates of recommended diagnostic testing after first trimester screening vs. same-day screening by cell free DNA. *American Journal of Obstetrics and Gynecology* 2016;**214**(1):S326.

Chen 2011a {published data only}

Chen F, Wang W, Shan D, Ren J, Xie J, Huang Y, et al. Noninvasive prenatal diagnosis of fetal aneuploidy by massively parallel sequencing of maternal plasma DNA. *Journal of Perinatal Medicine* 2011;**39**:299.

Da Fonseca 2015 {published data only}

Da Fonseca JP, El Khattabi L, Brun S, Chatron N, Gueguen P, Nectoux J, et al. Non-Invasive prenatal testing for the most common aneuploidies (trisomies 21, 18, and 13) using a semiconductor-sequencing platform: a French multicenter pilot study. *Chromosome Research* 2015;**23**(1):S110-S1.

ISRCTN11174071 {unpublished data only}

ISRCTN11174071. Comparison of false positive rates in prenatal combined screening and cell free DNA screening for trisomy 21. isrctn.com/ISRCTN11174071 Date first received: 26 July 2016. [DOI: [10.1186/ISRCTN11174071](https://doi.org/10.1186/ISRCTN11174071)]

Lin 2014 {published data only}

Lin G, Gao Y, Yin X, Tan Y, Zhang H, Lu G, et al. Clinical implementation of noninvasive prenatal testing in twin pregnancies with assisted reproductive technique treatment. *Prenatal Diagnosis* 2014;**34**:14.

Mu 2014 {published data only}

Mu Y, Sun F. Maternal non-invasive fetal DNA test used in prenatal diagnosis. *International Journal of Laboratory Hematology* 2014;**36**:106-7.

NCT01429389 {unpublished data only}

NCT01429389. Specimen collection from pregnant women at increased risk for fetal aneuploidy. [ClinicalTrials.gov/show/NCT01429389](https://clinicaltrials.gov/show/NCT01429389) Date first received: 2 September 2011.

NCT01472523 {unpublished data only}

NCT01472523. A safer pre-natal diagnosis using free DNA in maternal blood. [ClinicalTrials.gov/show/NCT01472523](https://clinicaltrials.gov/show/NCT01472523) Date first received: 11 October 2011.

NCT01545674 {unpublished data only}

NCT01545674. Prenatal non-invasive aneuploidy test utilizing SNPs trial. [ClinicalTrials.gov/show/NCT01545674](https://clinicaltrials.gov/show/NCT01545674) Date first received: 1 March 2012.

NCT01925742 {unpublished data only}

NCT01925742. Study of the efficacy of new non-invasive prenatal tests for screening for fetal trisomies using maternal blood. [ClinicalTrials.gov/show/NCT01925742](https://clinicaltrials.gov/show/NCT01925742) Date first received: 16 August 2013.

NCT02201862 {unpublished data only}

NCT02201862. Non-invasive chromosomal evaluation of trisomy study. [ClinicalTrials.gov/show/NCT02201862](https://clinicaltrials.gov/show/NCT02201862) Date first received: 22 July 2014.

NCT02278536 {unpublished data only}

NCT02278536. Multiple gestation study. [ClinicalTrials.gov/show/NCT02278536](https://clinicaltrials.gov/show/NCT02278536) Date first received: 28 October 2014.

NCT02278874 {unpublished data only}

NCT02278874. High risk multiple gestation study. [ClinicalTrials.gov/show/NCT02278874](https://clinicaltrials.gov/show/NCT02278874) Date first received: 27 August 2014.

NCT02317965 {unpublished data only}

NCT02317965. Non-invasive screening for fetal aneuploidy. [ClinicalTrials.gov/show/NCT02317965](https://clinicaltrials.gov/show/NCT02317965) Date first received: 9 December 2014.

NCT02424474 {unpublished data only}

NCT02424474. T21,18 and 13 screening by cell free fetal DNA in low risk patients. [ClinicalTrials.gov/show/NCT02424474](https://clinicaltrials.gov/show/NCT02424474) Date first received: 8 April 2015.

NCT02787486 {unpublished data only}

NCT02787486. Expanded noninvasive genomic medical assessment: the Enigma Study. [ClinicalTrials.gov/show/NCT02787486](https://clinicaltrials.gov/show/NCT02787486) Date first received: 26 May 2016.

Sago 2015 {published data only}

Sago H, Sekizawa A, and Japan NIPT consortium. Nationwide demonstration project of next-generation sequencing of cell-free DNA in maternal plasma in Japan: 1-year experience. *Prenatal Diagnosis* 2015;**35**(4):331-6.

Sanchez-Usabiaga 2015 {published data only}

Sanchez-Usabiaga RA, Aguinaga-Rios M, Batista-Espinoza A, Hurtado-Amador R, Romero-Tovar S. [Clinical implementation of non-invasive prenatal study for detecting aneuploidies by fetal DNA based on single nucleotide polymorphisms: two years in Mexico] [Implementacion clinica del estudio prenatal no invasivo para la deteccion de aneuploidias mediante ADN fetal con base en polimorfismos de nucleotido unico: dos anos en Mexico]. *Ginecologia y Obstetricia de Mexico* 2015;**83**(4):220-31. [PUBMED: 26727755]

Sistermans 2015 {published data only}

Sistermans EA, Schuring-Blom GH, Faas BHW, Boon EMJ, Bax CJ, Coumans ABC, et al. TRIDENT: or monitored NIPT

implementation in the Netherlands. *European Journal of Human Genetics* 2015;**23**(Suppl 1):C01.2.

Torres 2015 {published data only}

Torres Y, Suela J, Cigudosa JC. Genetic non invasive prenatal testing: A clinical and technical experience of 3.000 cases with follow-up. *Chromosome Research* 2015;**23**(1):S140-S141. [ISSN 09673849]

Van Wymersch 2015 {published data only}

Van Wymersch D, Gilson G. Introduction of noninvasive prenatal testing for fetal trisomies: preliminary results and consequences on invasive samplings [Introduction du dépistage sanguin des trisomies foetales par recherche d'ADN foetal circulant: Resultats preliminaires d'une annee et evolution des prelevements invasifs]. *Bulletin de la Societe des Sciences Medicales du Grand-Duche de Luxembourg* 2015;**1**:65-72. [PUBMED: 26946853]

Willems 2014 {published data only}

Willems PJ, Dierickx H, Vandenaeker E, Bekedam D, Segers N, Deboulle K, et al. The first 3,000 Non-Invasive Prenatal Tests (NIPT) with the Harmony™ test in Belgium and the Netherlands. *Facts, Views & Vision in ObGyn* 2014;**6**(1):7-12. [PUBMED: 25009720]

Yu 2014a {published data only}

Yu M, Fei S. Maternal non-invasive fetal DNA test used in prenatal diagnosis. *Clinical Chemistry and Laboratory Medicine* 2014;**52**:S1570.

Zwiefelhofer 2014 {published data only}

Zwiefelhofer T, Whitley P, Roy K, Saha M, Burcham T, Van Den Boom D, et al. Prenatal detection of fetal aneuploidy on the Ion Torrent Proton™ platform. *Prenatal Diagnosis* 2014;**34**:64-5.

Additional references

ACOG #163 2016

American College of Obstetricians and Gynecologists. Practice Bulletin No. 163 Summary: Screening for fetal aneuploidy. *Obstetrics and Gynecology* 2016;**127**(5):979-81. [PUBMED: 27101120]

ACOG #545 2012

American College of Obstetricians and Gynecologists Committee on Genetics. Committee Opinion No. 545: Noninvasive prenatal testing for fetal aneuploidy. *Obstetrics and Gynecology* 2012;**120**(6):1532-4. [PUBMED: 23168792]

ACOG #682 2016

American College of Obstetricians and Gynecologists and Society for Maternal-Fetal Medicine. Committee Opinion No.682: Microarrays and next-generation sequencing technology: The use of advanced genetic diagnostic tools in obstetrics and gynecology. *Obstetrics and Gynecology* 2016;**128**(6):e262-8. [PUBMED: 27875474]

ACOG #88 2007

American College of Obstetricians and Gynecologists. ACOG Practice Bulletin No. 88, December 2007. Invasive

prenatal testing for aneuploidy. *Obstetrics and Gynecology* 2007;**110**(6):1459-67. [PUBMED: 18055749]

Agarwal 2013

Agarwal A, Sayres LC, Cho MK, Cook-Deegan R, Chandrasekharan S. Commercial landscape of noninvasive prenatal testing in the United States. *Prenatal Diagnosis* 2013;**33**(6):521-31. [PUBMED: 23686656]

Akolekar 2015

Akolekar R, Beta J, Picciarelli G, Ogilvie C, D'Antonio F. Procedure-related risk of miscarriage following amniocentesis and chorionic villus sampling: a systematic review and meta-analysis. *Ultrasound in Obstetrics & Gynecology* 2015;**45**(1):16-26. [PUBMED: 25042845]

Allred 2010

Allred SK, Deeks JJ, Neilson JP, Alfirevic Z. Antenatal screening for Down's syndrome: generic protocol. *Cochrane Database of Systematic Reviews* 2010, Issue 4. [DOI: [10.1002/14651858.CD007384.pub2](https://doi.org/10.1002/14651858.CD007384.pub2); CD007384]

Allred 2012

Allred SK, Deeks JJ, Guo B, Neilson JP, Alfirevic Z. Second trimester serum tests for Down's Syndrome screening. *Cochrane Database of Systematic Reviews* 2012, Issue 6. [DOI: [10.1002/14651858.CD009925](https://doi.org/10.1002/14651858.CD009925)]

Allred 2015a

Allred SK, Takwoingi Y, Guo B, Pennant M, Deeks JJ, Neilson JP, et al. First trimester serum tests for Down's syndrome screening. *Cochrane Database of Systematic Reviews* 2015, Issue 11. [DOI: [10.1002/14651858.CD011975](https://doi.org/10.1002/14651858.CD011975); PUBMED: 26617074]

Allred 2015b

Allred SK, Guo B, Takwoingi Y, Pennant M, Wisniewski S, Deeks JJ, et al. Urine tests for Down's syndrome screening. *Cochrane Database of Systematic Reviews* 2015, Issue 12. [DOI: [10.1002/14651858.CD011984](https://doi.org/10.1002/14651858.CD011984); PUBMED: 26662198]

Allred 2017a

Allred SK, Takwoingi Y, Guo B, Pennant M, Deeks JJ, Neilson JP, et al. First trimester ultrasound tests alone or in combination with first trimester serum tests for Down's syndrome screening. *Cochrane Database of Systematic Reviews* 2017, Issue 3. [DOI: [10.1002/14651858.CD012600](https://doi.org/10.1002/14651858.CD012600); PUBMED: 28295158]

Allred 2017b

Allred SK, Takwoingi Y, Guo B, Pennant M, Deeks JJ, Neilson JP, et al. First and second trimester serum tests with and without first trimester ultrasound tests for Down's syndrome screening. *Cochrane Database of Systematic Reviews* 2017, Issue 3. [DOI: [10.1002/14651858.CD012599](https://doi.org/10.1002/14651858.CD012599); PUBMED: 28295159]

APA 2013

American Psychiatric Association. Neurodevelopmental disorders. Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DMS-5®). Arlington: American Psychiatric Publishing, 2013. [DOI: [10.1176/appi.books.9780890425596.514988](https://doi.org/10.1176/appi.books.9780890425596.514988)]

Ariosa Diagnostics 2016

Ariosa Diagnostics, Inc, USA. www.ariosadx.fr/healthcare-professionals/ (accessed 25 October 2016).

Attilakos 2011

Attilakos G, Maddocks DG, Davies T, Hunt LP, Avent ND, Soothill PW, et al. Quantification of free fetal DNA in multiple pregnancies and relationship with chorionicity. *Prenatal Diagnosis* 2011;**31**(10):967-72. [PUBMED: 21769896]

Bardsley 2013

Bardsley MZ, Kowal K, Levy C, Gosek A, Ayari N, Tartaglia N, et al. 47,XXY syndrome: clinical phenotype and timing of ascertainment. *Journal of Pediatrics* 2013;**163**(4):1085-94. [PUBMED: 23810129]

Basset 2013

Basset C, Beauvils F, Dreifuss-Netter F, Droit RP, Gaudray P, Matuchansky C, et al. Avis N° 120: Questions éthiques associées au développement des tests génétiques foetaux sur sang maternel. Comité Consultatif National d'Ethique (CCNE) pour les sciences de la vie et de la santé, 2013:50 pages.

Benn 2011

Benn P, Borrell A, Crossley J, Cuckle H, Dugoff L, Gross S, et al. Aneuploidy screening: a position statement from a committee on behalf of the Board of the International Society for Prenatal Diagnosis, January 2011. *Prenatal Diagnosis* 2011;**31**(6):519-22. [PUBMED: 21604286]

Benn 2013a

Benn P, Borell A, Chiu R, Cuckle H, Dugoff L, Faas B, et al. Position statement from the Aneuploidy Screening Committee on behalf of the Board of the International Society for Prenatal Diagnosis. *Prenatal Diagnosis* 2013;**33**(7):622-9. [PUBMED: 23616385]

Benn 2013b

Benn P, Cuckle H, Pergament E. Non-invasive prenatal testing for aneuploidy: current status and future prospects. *Ultrasound in Obstetrics & Gynecology* 2013;**42**(1):15-33. [PUBMED: 23765643]

Benn 2016

Benn P. Expanding non-invasive prenatal testing beyond chromosomes 21, 18, 13, X and Y. *Clinical Genetics* 2016;**90**(6):477-85. [PUBMED: 27283893]

Berry Genomics 2016

Berry Genomics Co. Ltd, China. www.berrygenomics.com/bambnittest/ (accessed 5 December 2016).

BGI 2014

Bejing Genomics Institute, China. www.niftytest.com/plus/list.php?tid=14 (accessed 20 June 2014).

BGI 2016

Bejing Genomics Institute, China. www.niftytest.com/healthcare-providers/clinical-data/ (accessed 25 October 2016).

Bianchi 2004

Bianchi DW. Circulating fetal DNA: its origin and diagnostic potential-a review. *Placenta* 2004;**25 Suppl A**:S93-S101. [PUBMED: 15033315]

Birch 2005

Birch L, English CA, O'Donoghue K, Barigye O, Fisk NM, Keer JT. Accurate and robust quantification of circulating fetal and total DNA in maternal plasma from 5 to 41 weeks of gestation. *Clinical Chemistry* 2005;**51**(2):312-20. [PUBMED: 15608152]

Bussani 2011

Bussani C, Di Tommaso M, Cioni R, Pasquini L, Quitadamo L, Scarselli G. Quantitative variation analysis of fetal DNA in maternal plasma samples collected before and after amniocentesis. *Journal of Obstetrics and Gynaecology Research* 2011;**37**(6):571-4. [PUBMED: 21375672]

Canick 2013

Canick JA, Palomaki GE, Kloza EM, Lambert-Messerlian GM, Haddow JE. The impact of maternal plasma DNA fetal fraction on next generation sequencing tests for common fetal aneuploidies. *Prenatal Diagnosis* 2013;**33**(7):667-74. [PUBMED: 23592541]

Cereda 2012

Cereda A, Carey JC. The trisomy 18 syndrome. *Orphanet Journal of Rare Diseases* 2012;**7**:81. [PUBMED: 23088440]

Chen 2009

Chen CP. Prenatal sonographic features of fetuses in trisomy 13 pregnancies (I). *Taiwanese Journal of Obstetrics & Gynecology* 2009;**48**(3):210-7. [PUBMED: 19797008]

Chetty 2013

Chetty S, Garabedian MJ, Norton ME. Uptake of noninvasive prenatal testing (NIPT) in women following positive aneuploidy screening. *Prenatal Diagnosis* 2013;**33**(6):542-6. [PUBMED: 23592525]

Chitayat 2011

Chitayat D, Langlois S, Wilson RD, Genetics Committee of the Society of Obstetricians and Gynaecologists of Canada, Prenatal Diagnosis Committee of the Canadian College of Medical Geneticists. Prenatal screening for fetal aneuploidy in singleton pregnancies. *Journal of Obstetrics and Gynaecology Canada : JOGC* 2011;**33**(7):736-50. [PUBMED: 21749752]

Choi 2012

Choi H, Van Riper M, Thoyre S. Decision making following a prenatal diagnosis of Down syndrome: an integrative review. *Journal of Midwifery & Women's Health* 2012;**57**(2):156-64. [PUBMED: 22432488]

Christianson 2006

Christianson A, Howson CP, Modell B. March of dimes. Global report on birth defects. The hidden toll of dying and disabled children. March of Dimes Birth Defects Foundation 2006:1-76. [<http://www.marchofdimes.com/glue/files/global-report-on-birth-defects-the-hidden-toll-of-dying-and-disabled-children-full-report.pdf>]

Chu 2006

Chu H, Cole SR. Bivariate meta-analysis of sensitivity and specificity with sparse data: a generalized linear mixed model approach. *Journal of Clinical Epidemiology* 2006; Vol. 59, issue 12:1331-2; author reply 1332-3. [PUBMED: 17098577]

Cochrane Glossary 2014

The Cochrane Collaboration's Glossary of terms. community-archive.cochrane.org/glossary (accessed 17 July 2014).

Deeks 2013

Deeks JJ, Wisniewski S, Davenport C. Chapter 4: Guide to the contents of a Cochrane Diagnostic Test Accuracy Protocol. In: Deeks JJ, Bossuyt PM, Gatsonis C (editors), *Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy* Version 1.0.0. The Cochrane Collaboration, 2013. Available from: methods.cochrane.org/sdt/handbook-dta-reviews.

Devlin 2004

Devlin L, Morrison PJ. Accuracy of the clinical diagnosis of Down syndrome. *Ulster Medical Journal* 2004;**73**(1):4-12. [PUBMED: 15244118]

Driscoll 2009

Driscoll DA, Gross S. Clinical practice. Prenatal screening for aneuploidy. *New England Journal of Medicine* 2009;**360**(24):2556-62. [PUBMED: 19516035]

Duncan 2011

Duncan A, Langlois S. Use of array genomic hybridization technology in prenatal diagnosis in Canada. *Journal of Obstetrics and Gynaecology Canada* 2011;**33**(12):1256-9. [PUBMED: 22166281]

Fan 2010

Fan HC, Blumenfeld YJ, Chitkara U, Hudgins L, Quake SR. Analysis of the size distributions of fetal and maternal cell-free DNA by paired-end sequencing. *Clinical Chemistry* 2010;**56**(8):1279-86. [PUBMED: 20558635]

Fishler 1991

Fishler K, Koch R. Mental development in Down syndrome mosaicism. *American Journal of Mental Retardation* 1991;**96**(3):345-51. [PUBMED: 1836734]

Gagnon 2010

Gagnon S, Labrecque M, Njoya M, Rousseau F, St-Jacques S, Legare F. How much do family physicians involve pregnant women in decisions about prenatal screening for Down syndrome?. *Prenatal Diagnosis* 2010;**30**(2):115-21. [PUBMED: 20013876]

Gekas 2009

Gekas J, Gagne G, Bujold E, Douillard D, Forest JC, Reinharz D, et al. Comparison of different strategies in prenatal screening for Down's syndrome: cost effectiveness analysis of computer simulation. *BMJ (Clinical Research Ed.)* 2009;**338**:b138. [PUBMED: 19218323]

Genesupport 2016

Genesupport, Switzerland. www.prendia.ch/fileadmin/user_upload/pdf/Tableau_Anomalies-Numeriques.pdf (accessed 25 October 2016).

Genoma 2016

Genoma, Switzerland. genoma.com/product_page.php?id=2 (accessed 10 November 2016).

Genome Care 2016

Genome Care, Korea. http://genomecare.net/en/ (accessed 5 December 2016).

Gil 2015a

Gil MM, Quezada MS, Revello R, Akolekar R, Nicolaides KH. Analysis of cell-free DNA in maternal blood in screening for fetal aneuploidies: updated meta-analysis. *Ultrasound in Obstetrics & Gynecology* 2015;**45**(3):249-66. [PUBMED: 25639627]

Groth 2013

Groth KA, Skakkebaek A, Host C, Gravholt CH, Bojesen A. Clinical review: Klinefelter syndrome--a clinical update. *Journal of Clinical Endocrinology and Metabolism* 2013;**98**(1):20-30. [PUBMED: 23118429]

HAS 2015

Haute Autorité de Santé. Screening test accuracy for fetal trisomy 21 with circulating cell-free DNA. [Les performances des tests de dépistage de la trisomie 21 foetale par analyse de l'ADN libre circulant]. Collège de la Haute Autorité de Santé 2015.

ICFMM 2013

Anonymous. Position Statement from the Italian College of Fetal Maternal Medicine: Non-invasive prenatal testing (NIPT) by maternal plasma DNA sequencing. *Journal of Prenatal Medicine* 2013;**7**(2):19-20. [PUBMED: 23991274]

illumina 2014

Verinata Health, Inc. www.verifitest.com/clinical-data/ (accessed 3 March 2014).

illumina 2016

illumina, Inc, USA. www.illumina.com/clinical/reproductive-genetic-health/nip.html (accessed 25 October 2016).

Irving 2011

Irving C, Richmond S, Wren C, Longster C, Embleton ND. Changes in fetal prevalence and outcome for trisomies 13 and 18: a population-based study over 23 years. *Journal of Maternal-Fetal & Neonatal Medicine* 2011;**24**(1):137-41. [PUBMED: 20384468]

Irving 2012

Irving CA, Chaudhari MP. Cardiovascular abnormalities in Down's syndrome: spectrum, management and survival over 22 years. *Archives of Disease in Childhood* 2012;**97**(4):326-30. [PUBMED: 21835834]

Karnis 2012

Karnis MF. Fertility, pregnancy, and medical management of Turner syndrome in the reproductive years. *Fertility and Sterility* 2012;**98**(4):787-91. [PUBMED: 23020910]

Kazerouni 2011

Kazerouni NN, Currier RJ, Flessel M, Goldman S, Hennigan C, Hodgkinson C, et al. Detection rate of quadruple-marker screening determined by clinical follow-up and registry data in the statewide California program, July 2007 to February 2009. *Prenatal Diagnosis* 2011;**31**(9):901-6. [PUBMED: 21706514]

LabGenomics 2016

LabGenomics Clinical Research Institute, Korea. www.momguard.com/eng/home.php?go=main (accessed 5 December 2016).

Langlois 2011

Langlois S, Duncan A. Use of a DNA method, QF-PCR, in the prenatal diagnosis of fetal aneuploidies. *Journal of Obstetrics and Gynaecology Canada* 2011;**33**(9):955-60. [PUBMED: 21923994]

Larion 2014

Larion S, Warsof SL, Romary L, Mlynarczyk M, Peleg D, Abuhamad AZ. Association of combined first-trimester screen and noninvasive prenatal testing on diagnostic procedures. *Obstetrics and Gynecology* 2014;**123**(6):1303-10. [PUBMED: 24807333]

Legare 2010

Legare F, Kearing S, Clay K, Gagnon S, D'Amours D, Rousseau M, et al. Are you SURE?: Assessing patient decisional conflict with a 4-item screening test. *Canadian Family Physician Medecin de Famille Canadien* 2010;**56**(8):e308-14. [PUBMED: 20705870]

Legare 2011

Legare F, St-Jacques S, Gagnon S, Njoya M, Brisson M, Fremont P, et al. Prenatal screening for Down syndrome: a survey of willingness in women and family physicians to engage in shared decision-making. *Prenatal Diagnosis* 2011;**31**(4):319-26. [PUBMED: 21268046]

Leggett 2010

Leggett V, Jacobs P, Nation K, Scerif G, Bishop DV. Neurocognitive outcomes of individuals with a sex chromosome trisomy: XXX, XYY, or XXY: a systematic review. *Developmental Medicine and Child Neurology* 2010;**52**(2):119-29. [PUBMED: 20059514]

LifeCodexx 2016

LifeCodexx AG, Germany. lifecodexx.com/en/for-physicians/methods-technology/clinical-validation/ (accessed 10 November 2016).

Lo 1997

Lo YM, Corbetta N, Chamberlain PF, Rai V, Sargent IL, Redman CW, et al. Presence of fetal DNA in maternal plasma and serum. *Lancet* 1997;**350**(9076):485-7. [PUBMED: 9274585]

Lo 1999

Lo YM, Zhang J, Leung TN, Lau TK, Chang AM, Hjelm NM. Rapid clearance of fetal DNA from maternal plasma. *American Journal of Human Genetics* 1999;**64**(1):218-24. [PUBMED: 9915961]

Lun 2008

Lun FM, Chiu RW, Allen Chan KC, Yeung Leung T, Kin Lau T, Dennis Lo YM. Microfluidics digital PCR reveals a higher than expected fraction of fetal DNA in maternal plasma. *Clinical Chemistry* 2008;**54**(10):1664-72. [PUBMED: 18703764]

Macaskill 2010

Macaskill P, Gatsonis C, Deeks JJ, Harbord RM, Takwoingi Y. Chapter 10: Analysing and presenting results. In: Deeks JJ, Bossuyt PM, Gatsonis C, (editors), *Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy* Version 1.0. The Cochrane Collaboration 2010. Available from: methods.cochrane.org/sdt/handbook-dta-reviews.

Mackie 2017

Mackie FL, Hemming K, Allen S, Morris RK, Kilby MD. The accuracy of cell-free fetal DNA-based non-invasive prenatal testing in singleton pregnancies: a systematic review and bivariate meta-analysis. *BJOG: an international journal of obstetrics and gynaecology* 2017;**124**(1):32-46. [PUBMED: 27245374]

Mazzanti 1998

Mazzanti L, Cacciari E. Congenital heart disease in patients with Turner's syndrome. Italian Study Group for Turner Syndrome (ISGTS). *Journal of Pediatrics* 1998;**133**(5):688-92. [PUBMED: 9821430]

MedlinePlus 2014

Medical Encyclopedia, A service of the U.S. National Library of Medicine, National Institutes of Health. medlineplus.gov/mpplusdictionary.html (updated daily, accessed 17 July 2014).

Mersy 2013

Mersy E, Smits LJ, van Winden LA, de Die-Smulders CE, Paulussen AD, Macville MV, et al. Noninvasive detection of fetal trisomy 21: systematic review and report of quality and outcomes of diagnostic accuracy studies performed between 1997 and 2012. *Human Reproduction Update* 2013;**19**(4):318-29. [PUBMED: 23396607]

Modi 2003

Modi D, Berde P, Bhartiya D. Down syndrome: a study of chromosomal mosaicism. *Reproductive Biomedicine Online* 2003;**6**(4):499-503. [PUBMED: 12831601]

Natera 2016

Natera, Inc, USA. www.natera.com/panorama-test/clinical-information (accessed 25 October 2016).

Newcombe 1998

Newcombe RG. Interval estimation for the difference between independent proportions: comparison of eleven methods. *Statistics in Medicine* 1998;**17**(8):873-90. [PUBMED: 9595617]

Nshimyumukiza 2017

Nshimyumukiza L, Menon S, Hina H, Rousseau F, Reinharz D. Cell-free DNA noninvasive prenatal screening for aneuploidy versus conventional screening: a systematic review of economic evaluations. *Clinical Genetics in press*.

O'Leary 2013

O'Leary P, Maxwell S, Murch A, Hendrie D. Prenatal screening for Down syndrome in Australia: costs and benefits of current and novel screening strategies. *Australian & New Zealand Journal of Obstetrics & Gynaecology* 2013;**53**(5):425-33. [PUBMED: 24090461]

Okun 2008

Okun N, Summers AM, Hoffman B, Huang T, Winsor E, Chitayat D, et al. Prospective experience with integrated prenatal screening and first trimester combined screening for trisomy 21 in a large Canadian urban center. *Prenatal Diagnosis* 2008;**28**(11):987-92. [PUBMED: 18925623]

Papageorgiou 2012

Papageorgiou EA, Patsalis PC. Non-invasive prenatal diagnosis of aneuploidies: new technologies and clinical applications. *Genome Medicine* 2012;**4**(5):46. [PUBMED: 22640877]

Parker 2010

Parker SE, Mai CT, Canfield MA, Rickard R, Wang Y, Meyer RE, et al. Updated National Birth Prevalence estimates for selected birth defects in the United States, 2004-2006. *Birth Defects Research. Part A, Clinical and Molecular Teratology* 2010;**88**(12):1008-16. [PUBMED: 20878909]

Premaitha Health plc 2016

Premaitha Health public limited company, UK. (accessed 15 November 2016).

Renshaw 2013

Renshaw R, Ellis K, Jacobs P, Morris J. Antenatal screening for Down syndrome: a quantitative demonstration of the improvements over the past 20 years. *Journal of Health Services Research & Policy* 2013;**18**(4):195-201. [PUBMED: 23864125]

Rothberg 2011

Rothberg JM, Hinz W, Rearick TM, Schultz J, Mileski W, Davey M, et al. An integrated semiconductor device enabling non-optical genome sequencing. *Nature* 2011;**475**(7356):348-52. [PUBMED: 21776081]

Rutter 2001

Rutter CM, Gatsonis CA. A hierarchical regression approach to meta-analysis of diagnostic test accuracy evaluations. *Statistics in Medicine* 2001;**20**(19):2865-84. [PUBMED: 11568945]

Saenger 1996

Saenger P. Turner's syndrome. *New England Journal of Medicine* 1996;**335**(23):1749-54. [PUBMED: 8929268]

Samura 2003

Samura O, Miharu N, Hyodo M, Honda H, Ohashi Y, Honda N, et al. Cell-free fetal DNA in maternal circulation after

amniocentesis. *Clinical Chemistry* 2003;**49**(7):1193-5. [PUBMED: 12816922]

Schoemaker 2008

Schoemaker MJ, Swerdlow AJ, Higgins CD, Wright AF, Jacobs PA. Mortality in women with turner syndrome in Great Britain: a national cohort study. *Journal of Clinical Endocrinology and Metabolism* 2008;**93**(12):4735-42. [PUBMED: 18812477]

Sequenom 2016

Sequenom, Inc. www.sequenom.com/tests/reproductive-health/maternit21-plus#provider-references and www.sequenom.com/tests/reproductive-health/visibilit#provider-test-details (accessed 25 October 2016).

Snijders 1999

Snijders RJ, Sundberg K, Holzgreve W, Henry G, Nicolaides KH. Maternal age- and gestation-specific risk for trisomy 21. *Ultrasound in Obstetrics & Gynecology* 1999;**13**(3):167-70. [PUBMED: 10204206]

South 2013

South ST, Lee C, Lamb AN, Higgins AW, Kearney HM. ACMG Standards and Guidelines for constitutional cytogenomic microarray analysis, including postnatal and prenatal applications: revision 2013. *Genetics in Medicine* 2013;**15**(11):901-9. [PUBMED: 24071793]

St-Jacques 2008

St-Jacques S, Grenier S, Charland M, Forest JC, Rousseau F, Legare F. Decisional needs assessment regarding Down syndrome prenatal testing: a systematic review of the perceptions of women, their partners and health professionals. *Prenatal Diagnosis* 2008;**28**(13):1183-203. [PUBMED: 19097031]

Stochholm 2006

Stochholm K, Juul S, Juel K, Naeraa RW, Gravholt CH. Prevalence, incidence, diagnostic delay, and mortality in Turner syndrome. *Journal of Clinical Endocrinology and Metabolism* 2006;**91**(10):3897-902. [PUBMED: 16849410]

Stochholm 2010a

Stochholm K, Juul S, Gravholt CH. Diagnosis and mortality in 47,XXX persons: a registry study. *Orphanet Journal of Rare Diseases* 2010;**5**:15. [PUBMED: 20509956]

Stochholm 2010b

Stochholm K, Juul S, Gravholt CH. Mortality and incidence in women with 47,XXX and variants. *American Journal of Medical Genetics. Part a* 2010;**152A**(2):367-72. [PUBMED: 20101696]

Summers 2007

Summers AM, Langlois S, Wyatt P, Wilson RD, Society of Obstetricians and Gynaecologists of Canada. Prenatal screening for fetal aneuploidy. *Journal of Obstetrics and Gynaecology Canada : JOGC* 2007;**29**(2):146-79. [PUBMED: 17346485]

Sybert 2004

Sybert VP, McCauley E. Turner's syndrome. *New England Journal of Medicine* 2004;**351**(12):1227-38. [PUBMED: 15371580]

Takwoingi 2015

Takwoingi Y, Guo B, Riley RD, Deeks JJ. Performance of methods for meta-analysis of diagnostic test accuracy with few studies or sparse data. *Statistical Methods in Medical Research* 2015;**26**(4):1896-911. [PUBMED: 26116616]

Tartaglia 2010

Tartaglia NR, Howell S, Sutherland A, Wilson R, Wilson L. A review of trisomy X (47,XXX). *Orphanet Journal of Rare Diseases* 2010;**5**:8. [PUBMED: 20459843]

Taylor-Phillips 2016

Taylor-Phillips S, Freeman K, Geppert J, Agbebiyi A, Uthman OA, Madan J, et al. Accuracy of non-invasive prenatal testing using cell-free DNA for detection of Down, Edwards and Patau syndromes: a systematic review and meta-analysis. *BMJ Open* 2016;**6**(1):e010002. [PUBMED: 26781507]

Tiller 2014

Tiller GE, Kershberg HB, Goff J, Coffeen C, Liao W, Sehnert AJ. Women's views and the impact of noninvasive prenatal testing on procedures in a managed care setting. *Prenatal Diagnosis* 2014;**34**:1-6. [PUBMED: 25201151]

Tyler 2004

Tyler C, Edman JC. Down syndrome, Turner syndrome, and Klinefelter syndrome: primary care throughout the life span. *Primary Care* 2004;**31**(3):627-48, x-xi. [PUBMED: 15331252]

Uematsu 2002

Uematsu A, Yorifuji T, Muroi J, Kawai M, Mamada M, Kaji M, et al. Parental origin of normal X chromosomes in Turner syndrome patients with various karyotypes: implications for the mechanism leading to generation of a 45,X karyotype. *American Journal of Medical Genetics* 2002;**111**(2):134-9. [PUBMED: 12210339]

UK Screening Glossary 2012

United Kingdom National Screening Committee Glossary. webarchive.nationalarchives.gov.uk/20150408175925/http://www.screening.nhs.uk/glossary (Last reviewed June 2012, accessed 18 November 2013).

Van Riper 2001

Van Riper M, Cohen WI. Caring for children with Down syndrome and their families. *Journal of Pediatric Health Care* 2001;**15**(3):123-31. [PUBMED: 11353361]

Vora 2010

Vora NL, Johnson KL, Peter I, Tighiouart H, Ralston SJ, Craigo SD, et al. Circulating cell-free DNA levels increase variably following chorionic villus sampling. *Prenatal Diagnosis* 2010;**30**(4):325-8. [PUBMED: 20127897]

Wald 2005

Wald NJ, Rodeck C, Hackshaw AK, Rudnicka A. SURUSS in perspective. *Seminars in Perinatology* 2005;**29**(4):225-35. [PUBMED: 16104673]

Weijerman 2010

Weijerman ME, de Winter JP. Clinical practice. The care of children with Down syndrome. *European Journal of Pediatrics* 2010;**169**(12):1445-52. [PUBMED: 20632187]

Wellesley 2012

Wellesley D, Dolk H, Boyd PA, Greenlees R, Haeusler M, Nelen V, et al. Rare chromosome abnormalities, prevalence and prenatal diagnosis rates from population-based congenital anomaly registers in Europe. *European Journal of Human Genetics* 2012;**20**(5):521-6. [PUBMED: 22234154]

Whiting 2011

Whiting PF, Rutjes AW, Westwood ME, Mallett S, Deeks JJ, Reitsma JB, et al. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Annals of Internal Medicine* 2011;**155**(8):529-36. [PUBMED: 22007046]

Wilson 2007

Wilson RD, Langlois S, Johnson JA. Mid-trimester amniocentesis fetal loss rate. *Journal of Obstetrics and Gynaecology Canada : JOGC* 2007;**29**(7):586-95. [PUBMED: 17623573]

Wright 2009

Wright C. Cell-free fetal nucleic acids for non-invasive prenatal diagnosis, Report of the UK expert working group. PHG foundation (2009).

Wu 2013a

Wu J, Springett A, Morris JK. Survival of trisomy 18 (Edwards syndrome) and trisomy 13 (Patau Syndrome) in England and Wales: 2004-2011. *American Journal of Medical Genetics. Part A* 2013;**161**(10):2512-8. [PUBMED: 23949924]

Wu 2013b

Wu J, Morris JK. The population prevalence of Down's syndrome in England and Wales in 2011. *European Journal of Human Genetics* 2013;**21**(9):1016-9. [PUBMED: 23321618]

Yaron 2016

Yaron Y. The implications of non-invasive prenatal testing failures: a review of an under-discussed phenomenon. *Prenatal Diagnosis* 2016;**36**(5):391-6. [PUBMED: 26941176]

Yu 2013

Yu SC, Lee SW, Jiang P, Leung TY, Chan KC, Chiu RW, et al. High-resolution profiling of fetal DNA clearance from maternal plasma by massively parallel sequencing. *Clinical Chemistry* 2013;**59**(8):1228-37. [PUBMED: 23603797]

Zhu 2013

Zhu JL, Hasle H, Correa A, Schendel D, Friedman JM, Olsen J, et al. Survival among people with Down syndrome: a nationwide population-based study in Denmark. *Genetics in Medicine* 2013;**15**(1):64-9. [PUBMED: 22878506]

References to other published versions of this review

 Cochrane Database of Systematic Reviews 2015, Issue 7. [DOI: [10.1002/14651858.CD011767](https://doi.org/10.1002/14651858.CD011767)]

Badeau 2015

Badeau M, Lindsay C, Blais J, Takwoingi Y, Langlois S, Légaré F, et al. Genomics-based non-invasive prenatal testing for detection of fetal chromosomal aneuploidy in pregnant women.

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES
Characteristics of included studies [ordered by study ID]

Alberti 2015

Study characteristics	
Patient sampling	Study design: case-control study (1:2) from a prospective cohort. Participants: pregnant women selected from a high-risk population of fetal aneuploidy. Inclusion criteria: pregnant women who had a risk of fetal trisomy 21 (> 1 in 250), based on the combination of maternal age with ultrasound and maternal serum markers during the first or second trimester and prior invasive testing. Exclusion criteria: multifetal pregnancies, absence of medical coverage by the National Health System and women declining an invasive procedure.
Patient characteristics and setting	Number enrolled: 976 pregnant women. Number available for 2 x 2 table: 183 pregnant women (subgroup of 19%). 23 euploid samples were used as reference set and 8 samples randomly chosen for pretesting phase. Setting: 3 centres in France. Recruitment period: March 2010 to April 2013. Ethnicity: not reported. Mean gestational age (\pm SD): 14 (\pm 2) weeks. Mean maternal age (\pm SD): 35.2 (\pm 6.7) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening. Language of the study: English.
Index tests	gNIPT by MPSS on Illumina HiSeq 2000 without multiplexing. Each library was sequenced using 50 bases-length reads chemistry in a single end-flow cell. Mean fetal fraction DNA: (male only) euploid: 20.11% and T21: 16.86%. Blood samples for gNIPT were collected before reference standard. Cutpoint: positive if Z score > 3. In-house gNIPT.
Target condition and reference standard(s)	Target condition: T21. Reference standard: fetal karyotype of chorionic villi or amniotic fluid.
Flow and timing	Blood samples were obtained prior to the invasive procedure (reference standard). gNIPT was a second-tier test. 701/976 samples were not selected for the case-control study. 50/275 samples were excluded during DNA extraction (47 for low amount of DNA and 3 for haemolysis) (no gNIPT results). 31/225 samples were excluded from analysis (8 for pretesting phase and 23 for reference set). 11/194 samples were excluded from analysis for insufficient fetal fraction DNA (no gNIPT results). No repeated test reported.
Comparative	

Alberti 2015 (Continued)

Aim to study	To evaluate the implementation of gNIPT for trisomy 21 into a cytogenetics laboratory in a university teaching hospital as well as validate gNIPT's clinical use on samples collected prospectively.
--------------	--

Funding source or sponsor of the study	Study not funded by industry.
--	-------------------------------

Informations about the authors contacted	Authors were contacted on: 23 March and 4 May 2016. Last reply received on: 16 May 2016.
--	---

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	No		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			

Alberti 2015 (Continued)

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Ashoor 2012
Study characteristics

Patient sampling	<p>Study design: nested case-control study (1:3) from a prospective cohort. Participants: pregnant women selected from a high-risk population (archived maternal plasma samples) of fetal aneuploidy.</p> <p>Inclusion criteria: singleton pregnancies between 11 to 13 weeks' gestation. Exclusion criteria: pregnancies that were conceived by in vitro fertilization.</p>
Patient characteristics and setting	<p>Number enrolled: 400 pregnant women. Number available for 2 x 2 table: 397 archived plasma samples (subgroup of 99%). Setting: 1 centre. Tertiary Referral Centre, King's College Hospital, London, United Kingdom. Recruitment period: March 2006 to August 2011. Ethnicity: Caucasian (88.5%), Afro-Caribbean (5%), South Asian (4%), East Asian (2%) and multiracial (0.5%). Mean gestational age (range): 13.3 (12.1 to 13.7) weeks. Mean maternal age (range): 36.2 (29.9 to 41.2) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening. Language of the study: English.</p>
Index tests	<p>gNIPT by TMPS (DANSR assay) on Illumina HiSeq 2000 in 96-plex.</p> <p>Fetal fraction DNA: not reported. Blood samples for gNIPT were collected before reference standard. Cutpoint: not reported. Usually, Harmony™ prenatal test uses FORTE algorithm; positive if FORTE risk score ≥ 1%. Commercial test: Harmony™ prenatal test by Ariosa Diagnostics, Inc.</p>
Target condition and reference standard(s)	<p>Target conditions: T21 and T18. Reference standard: fetal karyotype of chorionic villi.</p>
Flow and timing	<p>Blood samples were obtained prior to the invasive procedure (reference standard).</p> <p>gNIPT was a second-tier test. 3/400 samples failed amplification and sequencing (no gNIPT result).</p> <p>25 samples did not meet Ariosa Diagnostics, Inc acceptance criteria but they were replaced with the next available cases.</p> <p>No repeated test reported.</p>
Comparative	

Ashoor 2012 (Continued)

Aim to study	To assess the prenatal detection rate of T21 and T18 and the false-positive rate by chromosome-selective sequencing of maternal plasma cfDNA.
Funding source or sponsor of the study	Study not funded by industry but samples were analysed at Ariosa Diagnostics, Inc.
Informations about the authors contacted	No need for further contact.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	No		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test TMPS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		

Ashoor 2012 (Continued)

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Ashoor 2013
Study characteristics

Patient sampling Study design: blinded prospective cohort (second phase). First phase (case-control study) not shown in the present review.
 Participants: euploid pregnancies underwent routine first-trimester combined screening and confirmed T13 cases were selected.
 Inclusion criteria: singleton pregnancies.
 Exclusion criteria: multifetal pregnancies.

Patient characteristics and setting Number enrolled: 2167 pregnant women.
 Number available for 2 x 2 table: 1949 pregnant women (subgroup of 90%).
 Setting: several centres. Euploid pregnancies were from King's College Hospital, London, UK and T13 cases were from the USA.
 Recruitment period: October 2010 to January 2011 for euploid pregnancies. Not reported for T13 cases.
 Ethnicity: Caucasian (70.8%), African (20%), Asian (6.8%), mixed (2.6%).
 Mean gestational age (\pm SD; range): 12.7 (\pm 0.62; 13 to 26) weeks.
 Mean maternal age (\pm SD): 31.8 (\pm 5.6) years.
 Relevant tests carried out prior to index test: not reported.
 Language of the study: English.

Index tests gNIPT by TMPS (DANSR assay) on Illumina HiSeq 2000 in 96-plex.
 Median fetal fraction DNA (range): euploids: 10.0% (4.1% to 31.0%) and T21: 14.0% (6.1% to 24.0%).
 Blood samples for gNIPT were collected before reference standard for euploid pregnancies. T13 samples were collected post-confirmation of trisomy by karyotyping (reference standard).
 Cutpoint: positive if FORTE algorithm risk score \geq 1%.
 Commercial test: Harmony™ Prenatal Test by Ariosa Diagnostics, Inc.

Target condition and reference standard(s) Target condition: T13.
 Reference standards: fetal karyotype of chorionic villi or amniotic fluid or neonatal clinical examination.

Flow and timing Blood samples were obtained at the time of screening for euploid pregnancies (before reference standard).
 Blood samples were obtained after T13 confirmation following invasive procedure (reference standard).
 gNIPT was a first- or a second-tier test.
 165/2167 samples were excluded because they were used in the first phase.
 53/2002 samples failed during amplification or sequencing (no gNIPT result).
 No repeated test reported.

Ashoor 2013 (Continued)

Comparative

Aim to study	To assess the performance of chromosome-selective sequencing of maternal plasma cell-free DNA (cfDNA) in non-invasive prenatal testing for trisomy 13.
Funding source or sponsor of the study	Study not funded by industry but samples were analysed at Ariosa Diagnostics, Inc.
Informations about the authors contacted	No need for further contact.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	High
DOMAIN 2: Index Test TMPS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			

Ashoor 2013 (Continued)

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Benachi 2015
Study characteristics

Patient sampling Study design: blinded, retrospective analysis from a prospective cohort.
 Participants: all pregnant women considered at high risk of fetal aneuploidies who were willing to undergo invasive procedure.
 Inclusion criteria: at least 18 years old, more than 10 weeks of gestation and singleton or twin pregnancies.
 Exclusion criteria: vanishing twin or < 18 years old.

Patient characteristics and setting Number enrolled: 900 pregnant women.
 Number available for 2 x 2 table: 886 pregnant women (subgroup of 98%).
 Setting: 29 centres. French Fetal Medicine Centres in France.
 Recruitment period: December 2012 to October 2013.
 Ethnicity: Caucasian (84.2%), Black or Caribbean (4.6%), Asian (2.0%), mixed (5.7%) and unknown (3.5%).
 Median gestational age (range): 15.1 (10.2 to 34.6) weeks.
 Median maternal age (range): 35 (30 to 39) years.
 Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening.
 Language of the study: English.

Index tests gNIPT by MPSS with Illumina v3 flow-cell on a HiSeq 1500 sequencer in 12-plex.
 Mean fetal fraction DNA: group 1 (patients without abnormal fetal ultrasound findings, but at high risk of fetal aneuploidy): 10.9% and group 2 (high risk of fetal aneuploidy after ultrasound finding): 11.2%.
 Blood samples for gNIPT were collected just before reference standard.
 Cutpoint: positive if Z score > 3 (T21) or > 3.95 (T18 and T13).
 Commercial test: Laboratoire CERBA's prenatal test.

Target condition and reference standard(s) Target conditions: T21, T18 and T13.
 Reference standards: fetal karyotype of chorionic villi or amniotic fluid or neonatal clinical examination.

Flow and timing Blood samples were obtained prior to the invasive procedure (reference standard).
 gNIPT was a second-tier test.
 8/900 samples without karyotype result were excluded.
 42 samples failed the initial MPSS testing for technical issues.
 42/42 repeated tests using a second aliquot and 36/42 samples obtained gNIPT results.

Benachi 2015 (Continued)

6/892 samples failed during gNIPT process (low fetal fraction DNA or result appeared atypical) (no gNIPT result).

Comparative

Aim to study To evaluate the performance of the gNIPT (using fetal ccfDNA) for detection of the 3 main autosomal fetal trisomies in a very high-risk population of patients whose fetuses display ultrasonographically identified anomalies by comparing the results with those obtained by conventional fetal karyotyping.

Funding source or sponsor of the study Funding source not reported. 1 author is an employee of Laboratoire CERBA and also a shareholder.

Informations about the authors contacted Authors were contacted on: 25 May 2016.
Reply received on: 26 May 2016.

Notes Authors are from de Collaborative SEquençage a Haut Debit et Aneuploidies (SEHDA) Study Group.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	Yes		
--	-----	--	--

Was a case-control design avoided?	Yes		
------------------------------------	-----	--	--

Did the study avoid inappropriate exclusions?	No		
---	----	--	--

High

Low

DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
---	-----	--	--

If a threshold was used, was it pre-specified?	Yes		
--	-----	--	--

Low

Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes		
---	-----	--	--

Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
--	-----	--	--

Benachi 2015 (Continued)

	Low	Low
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Yes	
Did all analysed patients receive the reference standard?	Yes	
Were all patients included in the analysis?	No	
High		

Bevilacqua 2015
Study characteristics

Patient sampling	<p>Study design: prospective cohort study.</p> <p>Participants: pregnant women between 10 to 28 weeks' gestation selected at high risk of fetal trisomy or women who wanted to have the new test as a primary method of screening (unselected population).</p> <p>Inclusion criteria: singleton (not reported in the present review) or twin pregnancies between 10 to 28 weeks' gestation.</p> <p>Exclusion criteria: not reported.</p>
Patient characteristics and setting	<p>Number enrolled: 2362 pregnant women including 1847 singleton pregnancies (data not reported in the present review) and 515 twin pregnancies.</p> <p>Number available for 2 x 2 table: 340 twin pregnancies (subgroup of 66%).</p> <p>Setting: multicentre.</p> <p>Recruitment period: May 2013 to September 2014 (twin).</p> <p>Ethnicity: not reported.</p> <p>Median gestational age (range): 13.0 (10 to 28) weeks.</p> <p>Median maternal age (range): 36.8 (19 to 50.3) years.</p> <p>Chorionicity (368/515): 13% monochorionic and 58.4% dichorionic.</p> <p>Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening for some women.</p> <p>Language of the study: English.</p>
Index tests	<p>gNIPT by TMPS (DANSR assay) on Illumina HiSeq 2000 in 96-plex.</p> <p>Mean fetal fraction DNA (range): twins: 8.7% (4.1% to 30.0%) and singleton: 11.7% (4.0% to 38.9%).</p> <p>Blood samples for gNIPT were collected before reference standard.</p> <p>Cutpoint: not reported. Usually, Harmony™ prenatal test uses FORTE algorithm; positive if FORTE risk score \geq 1%.</p> <p>Commercial test: Harmony™ Prenatal Test by Ariosa Diagnostics, Inc.</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18 and T13.</p> <p>Reference standards: fetal karyotype of chorionic villi or amniotic fluid or neonatal karyotype.</p>
Flow and timing	<p>Blood samples were obtained prior to the invasive procedure (reference standard).</p> <p>gNIPT was a first- or second-tier test.</p>

Bevilacqua 2015 (Continued)

164/515 samples without follow-up were excluded including 138 lost to follow-up, 19 still pregnant and 7 miscarriages or stillbirths without karyotype of fetal tissue.

29/515 samples failed the initial TMPS testing.

26/29 samples resequenced with a second aliquot of the first sampling and 13/26 samples obtained a gNIPT result.

16/515 samples failed during sequencing process (no gNIPT result).

Comparative
Aim to study

To report the clinical implementation of cfDNA analysis of maternal blood in screening for T21, T18 and T13 in a large series of twin pregnancies and examine variables that could influence the failure rate of the test.

Funding source or sponsor of the study

Study not funded by industry but Ariosa Diagnostics, Inc made blinded sequencing and analysis.

Informations about the authors contacted

Author was contacted on: 1 June and 27 September 2016.

No replies received from the author.

Notes

gNIPT results from singleton pregnancies were not reported in the present review for incomplete 2 x 2 tables.

Methodological quality
Item
Authors' judgement
Risk of bias
Applicability concerns
DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?

No

Was a case-control design avoided?

Yes

Did the study avoid inappropriate exclusions?

No

High

High

DOMAIN 2: Index Test TMPS

Were the index test results interpreted without knowledge of the results of the reference standard?

Yes

If a threshold was used, was it pre-specified?

Yes

Low

Low

DOMAIN 3: Reference Standard

Bevilacqua 2015 (Continued)

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Unclear

Unclear

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Bianchi 2012
Study characteristics

Patient sampling Study design: nested case-control (1:4) study from the MELISSA prospective cohort. Participants: pregnant women randomly selected from a high-risk population (archived maternal plasma samples). Inclusion criteria: singleton pregnancies at high risk of fetal aneuploidy between 8 and 22 weeks of gestation. Exclusion criteria: multifetal pregnancies.

Patient characteristics and setting Number enrolled: 2882 pregnant women. Number available for 2 x 2 table: 503 pregnant women for T21, 502 for T18, 501 for T13 and 489 for 45,X (subgroup of 17%). Setting: 60 centres. Medical centre in 25 states in USA. Samples from 53 centres were analysed. Recruitment period: June 2010 to August 2011. Ethnicity: Caucasian (72.7%), Afro American (10.9%), Asian (9.9%), Native American or Alaska Native (0.9%) and multiracial (5.6%). Mean gestational age (\pm SD; range): 15.1 (\pm 3.16; 10 to 23) weeks. Mean maternal age (\pm SD; range): 35.2 (\pm 6.40; 18 to 46) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening. Language of the study: English.

Index tests gNIPT by MPSS on Illumina HiSeq 2000 sequencer in 6-plex. Fetal fraction DNA: amount measured but not reported. Blood samples for gNIPT were collected before reference standard. Cutpoint: 1) for T21, T18, and T13: positive if NCV > 4 (aneuploidy suspected if NCV is between 2.5 and 4).

Bianchi 2012 (Continued)

- 2) for 45,X: positive if NCV for Chrom. X < -4 and NCV for Chrom. Y < 2.5.
- 3) for 47,XXX: positive if NCV for Chrom. X > 4 and NCV for Chrom. Y < 2.5.
- 4) for 47,XXY: positive if NCV for Chrom. X between -2.5 and 2.5 and NCV for Chrom. Y > 33.
- 5) for 47,XYY: positive if NCV for Chrom. X < -4 and NCV for Chrom. Y > 4 with NCV for Chrom. Y is 2 times greater than expected NCV Chrom. X.
- Commercial test: Verinata's prenatal test.

Target condition and reference standard(s)	Target conditions: T21, T18, T13, 45,X, 47,XXX, 47,XXY and 47,XYY. Reference standard: fetal karyotype of chorionic villi (42.7%), amniotic fluid (56.4%) or products of conception (0.9%).
Flow and timing	Blood samples were obtained prior to the invasive procedure (reference standard). gNIPT was a second-tier test. 257/2882 samples were excluded (127 were ineligible, 45 without karyotype and 85 for multifetal pregnancies). 2091/2625 samples were not selected for this case-control study. 2/534 samples were excluded for tracking issue. 16/532 samples without fetal DNA detected were excluded during process (no gNIPT result). 13/516 samples were excluded of T21 2 x 2 table for censored complex karyotype. 14/516 samples were excluded of T18 2 x 2 table for censored complex karyotype. 15/516 samples were excluded of T13 2 x 2 table for censored complex karyotype. 27/516 samples were excluded of 45,X 2 x 2 table for censored complex karyotype. No repeated test reported.
Comparative	
Aim to study	To prospectively determine the diagnostic accuracy of massively parallel sequencing to detect whole chromosome fetal aneuploidy from maternal plasma.
Funding source or sponsor of the study	Study funded by Verinata Health, Inc. (a wholly owned subsidiary of Illumina, Inc.).
Informations about the authors contacted	Authors were contacted on: 1 March and 30 November 2016. Replies received on: 1 March and 8 December 2016.
Notes	This study is a clinical trial. MELISSA study. Clinicaltrials.gov NCT01122524. Data for 47,XXY, 47,XYY and 47,XXX were incomplete in the publication (data not shown in the present review).

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		

Bianchi 2012 *(Continued)*

Was a case-control design avoided? No

Did the study avoid inappropriate exclusions? No

High
Low
DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High
Bianchi 2013
Study characteristics

Bianchi 2013 (Continued)

Patient sampling	<p>Study design: retrospective study (archived maternal plasma samples) from a prospective cohort.</p> <p>Participants: pregnant women selected from a high-risk population (archived maternal plasma samples).</p> <p>Inclusion criteria: eligible blood samples, singleton pregnancies with karyotype result and nuchal cystic hygroma on fetal ultrasound.</p> <p>Exclusion criteria: multifetal pregnancies.</p>
Patient characteristics and setting	<p>Number enrolled: 2882 pregnant women.</p> <p>Number available for 2 x 2 table: 113 pregnant women (subgroup of 4%).</p> <p>Setting: 60 centres in USA.</p> <p>Recruitment period: June 2010 to August 2011.</p> <p>Ethnicity: Caucasian (73%), Afro-American (10%), Asian (9%) and multiracial (8%).</p> <p>Mean gestational age (\pm SD): 13.2 (\pm 2.0) weeks.</p> <p>Median gestational age (range): 12.6 (10 to 21) weeks.</p> <p>Mean maternal age (\pm SD): 32.2 (\pm 5.8) years.</p> <p>Median maternal age (range): 32.9 (18 to 44) years.</p> <p>Relevant test carried out prior to index test: ultrasonography (nuchal translucency measurement).</p> <p>Language of the study: English.</p>
Index tests	<p>gNIPT by MPSS with the sequencing chemistry Illumina TrueSeq 3.0.</p> <p>Fetal fraction DNA: not reported.</p> <p>Blood samples for gNIPT were collected before reference standard.</p> <p>Cutpoint:</p> <ol style="list-style-type: none"> 1) for T21, T18 and T13: positive if NCV > 4 (aneuploidy suspected zone between 3 and 4). 2) for 45,X: positive if NCV Chrom. X < -3 and NCV Chrom. Y < 3. <p>Commercial test: Verinata's prenatal test.</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18, T13 and 45,X.</p> <p>Reference standard: fetal karyotype of chorionic villi (78%), amniotic fluid (20%) or products of conception (2%).</p>
Flow and timing	<p>Blood samples were obtained prior to the invasive procedure (reference standard).</p> <p>gNIPT was a second-tier test.</p> <p>2769/2882 samples were not selected for this study.</p> <p>No failed sample reported.</p> <p>No repeated test reported.</p>
Comparative	
Aim to study	<p>To estimate the accuracy and potential clinical effect of using massively parallel sequencing of maternal plasma DNA to detect fetal aneuploidy in a population of pregnant women carrying fetuses with nuchal cystic hygroma.</p>
Funding source or sponsor of the study	<p>Study funded by Verinata Health, Inc. (a wholly owned subsidiary of Illumina, Inc.).</p>
Informations about the authors contacted	<p>No need for further contact.</p>
Notes	<p>74/113 samples were previously sequenced during the MELISSA trial. In this study, all 113 samples were newly resequenced (no overlap) with MELISSA study.</p>

Bianchi 2013 (Continued)

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all analysed patients receive the reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Bianchi 2014a

Study characteristics

Patient sampling	<p>Study design: blinded, prospective cohort study.</p> <p>Participants: pregnant women who planned to undergo (without prior risk) or had completed (high- or low-risk) standard prenatal serum screening for fetal aneuploidy.</p> <p>Inclusion criteria: pregnant women of 18 years or older, gestational age ≥ 8 weeks, able to provide consent and pregnancy records accessible and available for data collection.</p> <p>Exclusion criteria: invasive procedure (amniocentesis or CVS) performed within 2 weeks prior enrolment or prenatal screening determination by nuchal translucency measurement only.</p>
Patient characteristics and setting	<p>Number enrolled: 2052 pregnant women.</p> <p>Number available for 2 x 2 table: 1952 for T21 and T18 (subgroup of 95%) and 1914 for T13 (subgroup of 93%).</p> <p>Setting: 21 centres. In 14 states (USA).</p> <p>Recruitment period: 2 July 2012 to 4 January 2013.</p> <p>Ethnicity: Caucasian (65.4%), Afro-American (22.3%), Asian (7.3%) and other (5%).</p> <p>Mean gestational age (\pm SD; range): 20.3 (\pm 8.6; 8 to 39.4) weeks.</p> <p>Mean maternal age (\pm SD; range): 29.6 (\pm 5.54; 18 to 48.6) years.</p> <p>Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening.</p> <p>Language of the study: English.</p>
Index tests	<p>gNIPT by MPSS on Illumina HiSeq 2000 in 8-plex.</p> <p>Mean fetal fraction DNA: more than 35 years old: 11.3%, less than 35 years old: 11.6%, and at third trimester only: 24.6%.</p> <p>Blood samples for gNIPT were collected before or after reference standard.</p> <p>Cutpoint: positive if NCV ≥ 4. Resequenced if NCV is between 3 and 4.</p> <p>Commercial test: verifi[®] prenatal test by Verinata Health.</p> <p>The traditional screening tests (first-trimester combined test or a second-trimester result (quadruple, serum integrated, fully integrated or sequential)) were also assessed. Mixed cutpoints used.</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18 and T13.</p> <p>Reference standards: fetal karyotype of chorionic villi, amniotic fluid or products of conception, neonatal clinical examination or medical record from birth.</p>
Flow and timing	<p>Blood samples were obtained prior or after the invasive procedure (reference standard).</p> <p>gNIPT was a first- or second-tier test.</p> <p>10/2052 samples failed blood quality control before sequencing process.</p> <p>72/2042 samples without clinical outcome.</p> <p>38/2042 samples without standard screening result.</p> <p>17/2042 samples without gNIPT result.</p> <p>1/2042 samples without standard screening result and without gNIPT result.</p> <p>12 resequenced samples were in the grey zone (between affected and unaffected) and were successfully resequenced in uniplex.</p>
Comparative	
Aim to study	<p>To compare the results of gNIPT with ccfDNA for fetal autosomal aneuploidy with the results of conventional screening for T21 and T18 in a general obstetrical population. To compare false positive rates with the use of each method. To compare false positive rates for T13 in a subset of pregnant women in whom standard screening results included a risk assessment for trisomy 13. To compare fe-</p>

Bianchi 2014a (Continued)

tal ccfDNA fractions in low-risk patients and those in high-risk patients in the CARE study population to assess the potential effects of demographic differences on test performance.

Funding source or sponsor of the study Study funded by Illumina, Inc.

Informations about the authors contacted Author was contacted on: 10 February, 1 June and 28 June 2016.
 No replies received from the author.

Notes This study is a clinical trial (Comparison of Aneuploidy Risk Evaluations; CARE study). ClinicalTrials.gov number: NCT0166335.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	Unclear		
--	---------	--	--

Was a case-control design avoided?	Yes		
------------------------------------	-----	--	--

Did the study avoid inappropriate exclusions?	No		
---	----	--	--

High

High

DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
---	-----	--	--

If a threshold was used, was it pre-specified?	Yes		
--	-----	--	--

Low

Low

DOMAIN 2: Index Test Traditional screening tests

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
---	-----	--	--

If a threshold was used, was it pre-specified?	Yes		
--	-----	--	--

Low

Low

DOMAIN 3: Reference Standard

Bianchi 2014a (Continued)

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Bijok 2014
Study characteristics

Patient sampling Study design: prospective cohort study.
 Participants: pregnant women selected at high risk of fetal aneuploidy.
 Inclusion criteria: pregnant women at high risk of fetal aneuploidy with invasive test result.
 Exclusion criteria: not reported.

Patient characteristics and setting Number enrolled: 10 pregnant women.
 Number available for 2 x 2 table: 9 pregnant women (subgroup of 90%).
 Setting: obstetric and gynaecology clinic in Warsaw, Poland.
 Recruitment period: not reported.
 Ethnicity: not reported.
 Median gestational age (range): 16 (13 to 23) weeks.
 Median maternal age (range): 31 (26 to 36) years.
 Relevant test carried out prior to index test: ultrasonography (nuchal translucency measurement).
 Language of the study: Polish.

Index tests gNIPT by MPSS on Illumina Genome Analyzer Ix or HiSeq 2000 sequencer in multiplex with BGI's algorithm.
 Fetal fraction DNA: amount measured but not reported.
 Blood samples for gNIPT were collected before reference standard.
 Cutpoint: not reported.
 Commercial test: NIFTY™ test by Beijing Genomics Institute.

Target condition and reference standard(s) Target conditions: T21, T18, and T13.
 Reference standard: fetal karyotype of chorionic villi (30%) or amniotic fluid (70%).

Bijok 2014 (Continued)

Flow and timing	Blood samples were obtained prior to the invasive procedure (reference standard). gNIPT was a second-tier test. 1/10 sample failed during sequencing process (no gNIPT result) for low fetal fraction DNA. No repeated test reported.
-----------------	--

Comparative

Aim to study	To present initial results of non-invasive prenatal diagnosis of common aneuploidies (T21, T18, and T13) based on ccfDNA in maternal plasma in high-risk pregnant women, and to compare the results with routine karyotyping.
--------------	---

Funding source or sponsor of the study	Study not funded by industry but NIFTY™ tests were provided by Beijing Genomics Institute, Shenzhen, China.
--	---

Informations about the authors contacted	Authors were contacted on: 2 May and 4 July 2016. Replies received on: 4 and 16 May 2016.
--	--

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	Unclear		
--	---------	--	--

Was a case-control design avoided?	Yes		
------------------------------------	-----	--	--

Did the study avoid inappropriate exclusions?	Unclear		
---	---------	--	--

Unclear

Low

DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
---	-----	--	--

If a threshold was used, was it pre-specified?	Yes		
--	-----	--	--

Low

Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes		
---	-----	--	--

Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
--	-----	--	--

Bijok 2014 (Continued)

Low
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High
Canick 2012
Study characteristics

Patient sampling Study design: case-control study.
 Participants: all multifetal pregnant women with T21, T18 or T13 fetus were selected along with all euploid triplet pregnancies and a random selection of euploid twin pregnancies.
 Inclusion criteria: multifetal pregnant women, at least 18 years old, between about 10 weeks and 21 weeks 6 days of gestation, at high risk of aneuploidies and who undergo an invasive procedure.
 Exclusion criteria: singleton pregnancies or low risk of aneuploidy.

Patient characteristics and setting Number enrolled: 4664 pregnant women.
 Number available for 2 x 2 table: 27 multifetal pregnancies (25 twin and 2 triplet pregnancies) (subgroup of 0.6%).
 Setting: 27 centres. Prenatal diagnostic centres (Canada, Italy, Spain, Czech Republic, Argentina, Ireland, Hungary, USA, Israel and Australia).
 Recruitment period: April 2009 to February 2011.
 Ethnicity: not reported.
 Mean gestational age (range): 15.0 (10.9 to 19) years.
 Maternal age: not reported.
 Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening.
 Language of the study: English.

Index tests gNIPT by MPSS on Illumina HiSeq 2000 sequencer in 4-plex.
 Fetal fraction DNA range: 7% to 55%.
 Blood samples for gNIPT were collected before reference standard.
 Cutpoint: positive if Z score ≥ 3 .
 Commercial test: Sequenom's test.

Target condition and reference standard(s) Target conditions: T21 and T13. T18 was also assessed but no case was found.
 Reference standard: fetal karyotype of chorionic villi or amniotic fluid.

Flow and timing Blood samples were obtained immediately prior the invasive procedure (reference standard).
 gNIPT was a second-tier test.
 4637/4664 samples were not selected for this case-control study.
 No failed sample was reported in multifetal pregnancies.

Canick 2012 (Continued)

No repeated test reported.

Comparative

Aim to study	To study prenatal testing for T21, T18, and T13 by MPSS of fetal ccfDNA in high-risk multifetal pregnant women.
Funding source or sponsor of the study	Study funded by Sequenom, Inc. Some authors are employees and shareholders of Sequenom, Inc. or of Sequenom Center for Molecular Medicine.
Informations about the authors contacted	Author was contacted on: 10 March 2016. Reply received on: 16 March 2016.
Notes	This study is a clinical trial "A New Prenatal Blood Test for Down Syndrome" ClinicalTrials.gov number: NCT00877292.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	No		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low

Canick 2012 (Continued)

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? Yes

Low

Chen 2011
Study characteristics

Patient sampling

Study design: nested case-control study from a prospective cohort and archived plasma.

Participants: pregnant women with clinical indications of fetal aneuploidy (high risk of fetal aneuploidy) for invasive procedure.

Inclusion criteria: singleton pregnancies with and without trisomy 13, 18 or 21, matched for gestational ages.

Exclusion criteria: twin pregnancies.

Patient characteristics and setting

Number enrolled: 392 pregnant women (252 from the prospective cohort and 140 were archived plasma).

Number available for 2 x 2 table: 289 pregnant women (subgroup of 74%).

Setting: 10 centres in Hong Kong, the Netherlands, and UK.

Recruitment period for the prospective cohort: October 2008 to May 2009.

Recruitment period for the archived plasma samples collection: October 2003 to September 2008.

Ethnicity: not reported.

Gestational age: not reported.

Maternal age: not reported.

Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening.

Language of the study: English.

Index tests

gNIPT by MPSS on Illumina Genome Analyzer Iix in 2-plex.

Feta fraction DNA: not reported.

Blood samples for gNIPT were collected before reference standard.

Cutpoint: positive if Z score > 3.

Commercial test: Sequenom's test.

Chen 2011 (Continued)

Target condition and reference standard(s)	Target conditions: T18 and T13. Reference standard: fetal karyotype of chorionic villi or amniotic fluid.
Flow and timing	Blood samples were obtained prior to the invasive procedure (reference standard). gNIPT was a second-tier test. 103/392 samples were selected as reference control. No failed sample reported. No repeated test reported.
Comparative	
Aim to study	To assess the prenatal diagnostic performance by MPSS of maternal plasma DNA on a cohort of pregnant women with T13 and T18 fetuses.
Funding source or sponsor of the study	Study co-sponsored by Sequenom, Inc and Life Technologies. Some authors have filed patent on gNIPT (part of this patent has been licensed to Sequenom, Inc).
Informations about the authors contacted	Author was contacted on: 14 December 2015 and 10 May 2016. Reply received on: 12 May 2016.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	No		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear		
If a threshold was used, was it pre-specified?	Yes		
		Unclear	Low
DOMAIN 3: Reference Standard			

Chen 2011 (Continued)

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? Yes

Low

Chiu 2011
Study characteristics

Patient sampling	<p>Study design: blinded, case-control study (1:5) from a prospective cohort and archived plasma.</p> <p>Participants: pregnant women with clinical indications for invasive procedure, mixed risk (mostly high risk (> 1/300 at traditional screening test), intermediate risk (between 1/300 and 1/1000) or other risk factors).</p> <p>T21 and non T21 pregnancies matched for gestational ages.</p> <p>Inclusion criteria: singleton pregnancies.</p> <p>Exclusion criteria: multifetal pregnancies.</p>
Patient characteristics and setting	<p>Number enrolled: 824 pregnant women.</p> <p>Number available for 2 x 2 table: 753 (8-plex) (subgroup of 91%).</p> <p>Setting: 10 centres in Hong Kong, the Netherlands, and UK.</p> <p>Recruitment period for the prospective cohort: October 2008 to May 2009.</p> <p>Recruitment period for the archived plasma samples collection: October 2003 to September 2008.</p> <p>Ethnicity: not reported.</p> <p>Median gestational age: 13.1 weeks.</p> <p>Median maternal age: 35.4 years.</p> <p>Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening.</p>

Chiu 2011 (Continued)

	Language of the study: English.
Index tests	<p>gNIPT by MPSS on Illumina Genome Analyzer II in 8-plex and 2-plex (not reported in the present review).</p> <p>Median fetal fraction DNA (interquartile 1 and 3): male euploid: 15.2% (10.6% and 19.1%), archived samples: 14.7%, and prospective samples: 15.4%.</p> <p>Blood samples for gNIPT were collected before reference standard.</p> <p>Cutpoint: positive if Z score > 3.</p> <p>Commercial test: Sequenom's test.</p>
Target condition and reference standard(s)	<p>Target condition: T21.</p> <p>Reference standard: fetal karyotype of chorionic villi or amniotic fluid.</p>
Flow and timing	<p>Blood samples were obtained prior to the invasive procedure (reference standard).</p> <p>gNIPT was a second-tier test.</p> <p>60/824 samples were excluded before sequencing process (2 twin pregnancies, 12 without karyotype and 46 failed quality control for blood sampling).</p> <p>11/764 samples failed quality control during sequencing process (no gNIPT result).</p> <p>96/753 samples were also used for reference controls (8-plex).</p> <p>No repeated test reported.</p>
Comparative	
Aim to study	To validate the diagnostic performance and practical feasibility of massively parallel genomic sequencing for the non-invasive prenatal assessment of trisomy 21 in pregnant women who had undergone conventional screening and were clinically indicated for definitive testing.
Funding source or sponsor of the study	Study sponsored by Sequenom, Inc. Some authors have filed patent applications on gNIPT (part of this patent has been licensed to Sequenom, Inc).
Informations about the authors contacted	No need for further contact.
Notes	Data from 2-plex sequencing were excluded from the present review to avoid double counting. We kept data from 8-plex because it is the method most likely to be used for routine testing.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	No		
Did the study avoid inappropriate exclusions?	No		

Chiu 2011 (Continued)

		High	High
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all analysed patients receive the reference standard?	Yes		
Were all patients included in the analysis?	No		
		High	

Comas 2015
Study characteristics

Patient sampling	Study design: blinded, observational prospective cohort study. Participants: all pregnant women who underwent conventional first-trimester combined screening for fetal aneuploidies (without prior risk of fetal aneuploidy). Some pregnant women were referred after their combined test (high risk of fetal aneuploidy). Inclusion criteria: singleton pregnancies. Exclusion criteria: multifetal pregnancies, cases of ultrasound anomalies, nuchal translucency > 99 centile, combined risk at first-trimester screening > 1/10, or women at high risk of other genetic conditions.
Patient characteristics and setting	Number enrolled: 333 pregnant women (85.5% without prior risk and 16.5% were at high risk of fetal aneuploidy).

Comas 2015 (Continued)

Number available for 2 x 2 table: 312 pregnant women (subgroup of 95%).
 Setting: 1 private prenatal diagnostics centre in Barcelona, Spain (Hospital Universitari Quiron Dexeus).
 Recruitment period: January to December 2013.
 Ethnicity: not reported.
 Mean gestational age (range): 14.6 (9.5 to 23.5) weeks.
 Mean maternal age (range): 37 (21 to 46) years.
 Relevant test carried out prior to index test: biochemical screening for a part of the cohort.
 Language of the study: English

Index tests	<p>gNIPT by TMPS (DANSR assay or SNP-based method).</p> <p>Mean fetal fraction DNA (range): 12.7% (4.2% to 27.9%), Harmony™ prenatal test: 13.1%, and Panorama™ prenatal test: 12.7%. Blood samples for gNIPT were collected before reference standard. DANSR assay cutpoint: not reported. Usually, Harmony™ prenatal test uses FORTE algorithm; positive if FORTE risk score ≥ 1%. SNP-based method cutpoint: not reported. Commercial test: Panorama™ prenatal test by Natera, Inc. or Harmony™ prenatal test by Ariosa Diagnostics, Inc.</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18, T13, 45,X, 47,XXX, 47,XXY, 47,XYY. SCA data were not reported in the present review. T18 and T13 were also assessed but no case was found.</p> <p>Reference standards: fetal karyotype of chorionic villi or amniotic fluid or neonatal clinical examination.</p>
Flow and timing	<p>Blood samples were obtained prior to the invasive procedure (reference standard).</p> <p>gNIPT was a first- or second-tier test.</p> <p>17/333 samples excluded because still pregnant at the time of publication (no follow-up). 9/333 samples failed the initial TMPS testing. 6/9 repeated sampling was performed and results were obtained in 5/6. 3/333 samples without gNIPT result were excluded (unrepeated samples). 1/333 samples without gNIPT result and follow-up were excluded (still pregnant).</p>
Comparative	
Aim to study	To evaluate gNIPT of ccfDNA as a screening method for major chromosomal anomalies in a clinical setting.
Funding source or sponsor of the study	Study not funded by industry but Ariosa Diagnostics, Inc and Natera, Inc. made sequencing and analysis.
Informations about the authors contacted	Author was contacted on: 27 May 2016 and 31 May 2016. Reply received on: 31 May 2016.
Notes	gNIPT is offered to pregnant women at their own cost. 45,X, 47,XXY, 47,XYY and 47,XXX were also screened but inappropriate reference standard for the present review was used. gNIPT data from SCA were not shown in this review.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

Comas 2015 (Continued)

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled? Yes

Was a case-control design avoided? Yes

Did the study avoid inappropriate exclusions? No

High
High
DOMAIN 2: Index Test TMPS

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

del Mar Gil 2014
Study characteristics

Patient sampling	<p>Study design: retrospective cohort study. Data from prospective cohort were not shown in the present review.</p> <p>Participants: pregnant women without a priori risk who undergo first-trimester screening for trisomies (archived maternal plasma samples).</p> <p>Inclusion criteria: multifetal pregnancies between 11 to 13 weeks' gestation.</p> <p>Exclusion criteria: singleton pregnancies.</p>
Patient characteristics and setting	<p>Number enrolled: 207 pregnant women from the retrospective cohort.</p> <p>Number available for 2 x 2 table: 192 pregnant women (subgroup of 93%).</p> <p>Setting: 1 centre at Kings' College Hospital in London, UK.</p> <p>Recruitment period: not reported.</p> <p>Ethnicity: not reported.</p> <p>Median gestational age (range): 13.0 (12.4 to 13.9) weeks.</p> <p>Median maternal age (range): 33.7 (26.7 to 37.9) years.</p> <p>Chorionicity: 41% of pregnancies were monochorionic (85/207) and 59% of pregnancies were dichorionic (122/207).</p> <p>Relevant tests carried out prior to index test: none.</p> <p>Language of the study: English.</p>
Index tests	<p>gNIPT by TMPS (DANSR assay) on Illumina HiSeq 2000 in 96-plex.</p> <p>Mean fetal fraction DNA (range): euploids: 9.8% (7.4% to 12.1%), T21: 10.8% (6.8% to 12.1%), and T13: 7%.</p> <p>Blood samples for gNIPT were collected before reference standard.</p> <p>Cutpoint: not reported. Usually, Harmony™ prenatal test uses FORTE algorithm; positive if FORTE risk score $\geq 1\%$.</p> <p>Commercial test: Harmony™ prenatal test by Ariosa Diagnostics, Inc.</p>
Target condition and reference standard(s)	<p>Target conditions: T21 and T13. T18 was also assessed but no case was found.</p> <p>Reference standard: fetal karyotype of chorionic villi or amniotic fluid.</p>
Flow and timing	<p>Blood samples were obtained prior to the invasive procedure (reference standard).</p> <p>gNIPT was a first-tier test.</p> <p>15/207 samples failed during sequencing process (11 for low fetal fraction DNA and 4 for laboratory processing failures) (no gNIPT result).</p> <p>No repeated test reported.</p>
Comparative	
Aim to study	<p>To examine the clinical implementation of TMPS of ccfDNA in maternal blood and an algorithm that relies on the lower fetal fraction DNA contribution of the 2 fetuses in the assessment of risk for trisomies in twin pregnancies.</p>
Funding source or sponsor of the study	<p>Study not funded by industry but Ariosa Diagnostics, Inc have performed gNIPT at their own expense. Study funded by a grant from The Fetal Medicine Foundation, UK.</p>
Informations about the authors contacted	<p>Author was contacted on: 27 May and 27 September 2016.</p> <p>No reply received from the author.</p>
Notes	<p>Data from prospective cohort study were not shown in the present review because patients with gNIPT negative result were without follow-up to confirm gNIPT result.</p>

del Mar Gil 2014 (Continued)

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Unclear		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Unclear		
		Unclear	Low
DOMAIN 2: Index Test TMPS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all analysed patients receive the reference standard?	Yes		
Were all patients included in the analysis?	No		
		High	

Ehrich 2011

Study characteristics

Patient sampling	Study design: blinded, case-control study (1:11) from a prospective cohort. Participants: pregnant women selected from a high-risk population. Inclusion criteria: not reported. Exclusion criteria: not reported.
Patient characteristics and setting	Number enrolled: 480 pregnant women. Number available for 2 x 2 table: 449 pregnant women (subgroup of 94%). Setting: in clinical practice and pregnancy termination centres. Recruitment period: May 2009 to unknown date. Ethnicity: not reported. Median gestational age (range): 16 (8 to 36) weeks. Mean maternal age (range): 37 (18 to 47) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening. Language of the study: English.
Index tests	gNIPT by MPSS on Illumina Genome Analyzer Iix in 4-plex. Minimum fetal fraction DNA as estimated with the fetal quantifier assay: 3.9%. Blood samples for gNIPT were collected before reference standard. Cutpoint: positive if Z score > 2.5. Commercial test: Sequenom's test.
Target condition and reference standard(s)	Target condition: T21. Reference standard: fetal karyotype of chorionic villi (19%) or amniotic fluid (81%).
Flow and timing	Blood samples were obtained prior or after the invasive procedure (reference standard). gNIPT was a second-tier test. 13/480 samples excluded before sequencing process (9 for plasma volume < 3.5 mL and 4 for processing errors). 20/467 samples failed the initial MPSS testing. 20/20 samples were resequenced using the same library (10 samples in 4-plex and 10 in monoplex) and 2/20 samples obtained a gNIPT results. 18/467 samples failed quality control during sequencing process, including 7 samples for low fetal fraction DNA (no gNIPT result).
Comparative	
Aim to study	To evaluate a multiplexed massively parallel shotgun sequencing assay for noninvasive trisomy 21 detection using circulating cell-free fetal DNA.
Funding source or sponsor of the study	Study funded by Sequenom, Inc.
Informations about the authors contacted	Author was been contacted on: 5 May and 28 September 2016. No reply received from the author.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

Ehrich 2011 (Continued)

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	No
--	----

Was a case-control design avoided?	No
------------------------------------	----

Did the study avoid inappropriate exclusions?	Unclear
---	---------

High
Low
DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes
---	-----

If a threshold was used, was it pre-specified?	Yes
--	-----

Low
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes
---	-----

Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes
--	-----

Low
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes
--	-----

Did all analysed patients receive the reference standard?	Yes
---	-----

Were all patients included in the analysis?	No
---	----

High
Fiorentino 2016
Study characteristics

Patient sampling	Study design: blinded, prospective cohort study. Retrospective cohort (training set) not reported in the present review.
------------------	--

Fiorentino 2016 (Continued)

Participants: mostly pregnant women selected from a high-risk population and pregnant women without prior risk (14%).
Inclusion criteria: singleton pregnancies.
Exclusion criteria: multifetal pregnancies.

Patient characteristics and setting	Number enrolled: 7103 pregnant women. Number available for 2 x 2 table: 7082 pregnant women (subgroup of 99.7%). Setting: in Italy. Recruitment period: September to December 2014. Ethnicity: not reported. Mean gestational age (\pm SD; range): 12.8 (\pm 2.3; 10 to 30) weeks. Mean maternal age (\pm SD; range): 36.4 (\pm 4.7; 24 to 54) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both. Language of the study: English.
Index tests	gNIPT by MPSS on Illumina HiSeq 2500 sequencer in 15-plex with SAFer™ algorithm. Fetal fraction DNA: the limit of detection (the lowest fetal fraction DNA with a detectable aneuploidy) for T21 was determined at 2% fetal fraction level. Blood samples for gNIPT were collected before reference standard. Cutpoint: positive if NCV > 4 (aneuploidy suspected if NCV was between 3 and 4). Commercial test: Genoma's prenatal test.
Target condition and reference standard(s)	Target conditions: T21, T18 and T13. Reference standards: fetal karyotype of chorionic villi or amniotic fluid or neonatal clinical examination.
Flow and timing	Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard). gNIPT was a first- or a second-tier test. 100/7103 samples failed the initial MPSS testing. 79/100 repeated samples with a second blood draw and all obtained a gNIPT result. 21/100 unrepeated samples failed quality control metrics (no gNIPT result).
Comparative	
Aim to study	To determine the limit of detection of a gNIPT method, in order to define the actual lower fetal fraction DNA required to detect common fetal autosomal trisomies, using a model system to simulate samples at different proportions of fetal ccfDNA. Secondly, to assess the impact of low fetal fraction DNA on the performance of ccfDNA-based maternal plasma testing for aneuploidies.
Funding source or sponsor of the study	Study not funded by industry but the samples were analysed in the GENOMA laboratory (Rome, Italy).
Informations about the authors contacted	Authors were contacted on: 30 August and 6 September 2016. Reply received on: 6 September.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			

Fiorentino 2016 (Continued)

Was a consecutive or random sample of patients enrolled? Unclear

Was a case-control design avoided? Yes

Did the study avoid inappropriate exclusions? No

High

High

DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low

Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Unclear

Unclear

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Gil 2016
Study characteristics

Patient sampling Study design: prospective cohort study.

Gil 2016 (Continued)

Participants: pregnant women with a first-trimester combined test selected for their risk of fetal aneuploidy (cut-off of 1 in 100 for high risk and 1 in 101 to 1 in 2500 for intermediate risk).
Inclusion criteria: singleton pregnancies.
Exclusion criteria: multifetal pregnancies, terminations of pregnancy, miscarriages or stillbirths without follow-up.

Patient characteristics and setting

Number enrolled: 11,692 pregnant women.
Number available for 2 x 2 table: 3633 pregnant women (subgroup of 31%).
Setting: 2 centres. King's College Hospital, London, and Medway Maritime Hospital, Gillingham, Kent in UK.
Recruitment period: October 2013 to February 2015.
Ethnicity: Caucasian (70%), Afro-Caribbean (20%), Asian (7%) and mixed (3%).

Gestational age: not reported.
Median maternal age (range): 31.6 (25.8 to 39.5) years.
Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both.
Language of the study: English.

Index tests

gNIPT by TMPS (DANSR assay).

Fetal fraction DNA: not reported.
Blood samples for gNIPT were collected before reference standard.
Cutpoint: not reported. Usually, Harmony™ prenatal test uses FORTE algorithm; positive if FORTE risk score \geq 1%.
Commercial test: Harmony™ prenatal test by Ariosa Diagnostics, Inc.

Traditional screening test was also assessed but 2 x 2 tables were incomplete.

Target condition and reference standard(s)

Target conditions: T21, T18 and T13.
Reference standards: fetal karyotype of chorionic villi, postnatal karyotype or neonatal clinical examination.

Flow and timing

Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard).
gNIPT was a second-tier test.

7994/11,692 samples did not undergo a gNIPT (no gNIPT result).

99/3698 samples failed the initial TMPS testing.

54/99 repeated sampling were processed and 34/54 gNIPT results were obtained.

65/3698 samples without gNIPT result.

Comparative

Aim to study

To report the feasibility of implementing gNIPT. To examine the factors affecting patient decisions concerning their options for screening and decisions on the management of affected pregnancies. To report the prenatal diagnosis of fetal trisomies and outcome of affected pregnancies following the introduction of contingent screening.

Funding source or sponsor of the study

Study not funded by industry but the cost of collection and analysis of the blood samples for the cell-free DNA test was covered by Ariosa Diagnostics, Inc. These organisations had no role in study design, data collection, data analysis, data interpretation or writing of the report. Study was funded by a grant from The Fetal Medicine Foundation, UK.

Informations about the authors contacted

No need for further contact.

Notes

Gil 2016 (Continued)

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Unclear		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	High
DOMAIN 2: Index Test TMPS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	No		
		High	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all analysed patients receive the reference standard?	Yes		
Were all patients included in the analysis?	No		
		High	

Hall 2014

Study characteristics

Patient sampling	<p>Study design: case-control study (1:3), age-matched randomly selected from a larger cohort. Participants: pregnant women with an affected fetus or considered to be at high risk of fetal aneuploidy were recruited.</p> <p>Inclusion criteria: pregnant women at least 18 years of age who had signed an informed consent, and with singleton pregnancy. Exclusion criteria: fetal mosaicism.</p>
Patient characteristics and setting	<p>Number enrolled: more than 1000 pregnant women. Number available for 2 x 2 table: 64 pregnant women (subgroup of 6%). Setting: 6 centres. Western Institutional (WA, USA), Einstein Institutional (CA and MO, USA), Polish Mother's Memorial Hospital Institutional (Polish), Bio Medical Research Institute of America (CA, USA), and the Mt. Sinai School of Medicine (NY, USA). Recruitment period: March to December of 2012. Ethnicity: not reported. Median gestational age (range): 16.0 (12.1 to 22.7) weeks. Maternal age: not reported. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening. Language of the study: English.</p>
Index tests	<p>gNIPT by TMPS (SNP-based method) on Illumina Genome Analyzer IIx or HiSeq sequencer. Samples were amplified using 11,000-plex or 19,488-plex targeted polymerase chain reaction (targets included SNPs from chromosomes 13, 18, 21, X, and Y).</p> <p>Mean fetal fraction DNA (median; range): 12.1% (11.1%; 2.2% to 30.4%). Blood samples for gNIPT were collected before reference standard. Cutpoint: not reported. Commercial test: Natera's prenatal test.</p>
Target condition and reference standard(s)	<p>Target condition: T13. Reference standards: fetal karyotype of chorionic villi or amniotic fluid or genetic testing of the cord blood, buccal, saliva or products of conception.</p>
Flow and timing	<p>Blood samples were obtained prior to the invasive procedure (reference standard).</p> <p>gNIPT was a second-tier test.</p> <p>About 932 samples were not selected for this case-control study. 4/68 samples failed DNA quality threshold for low fetal fraction DNA (no gNIPT result).</p> <p>No repeated test reported.</p>
Comparative	
Aim to study	<p>To determine how a single nucleotide polymorphism (SNP)- and informatics-based non-invasive prenatal aneuploidy test performs in detecting trisomy 13.</p>
Funding source or sponsor of the study	<p>Study funded by Natera, Inc. (involved in study design, data collection and analysis, decision to publish, and preparation of the manuscript).</p>
Informations about the authors contacted	<p>Authors were contacted on: 21 April 2016, and 27 May 2016. No reply received from the authors.</p>
Notes	

Hall 2014 (Continued)

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	No		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test TMPS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear		
If a threshold was used, was it pre-specified?	Yes		
		Unclear	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all analysed patients receive the reference standard?	Yes		
Were all patients included in the analysis?	No		
		High	

Hooks 2014
Study characteristics

Patient sampling	<p>Study design: case-control study from archived plasma samples from a prospective cohort.</p> <p>Participants: pregnant women selected at high risk of fetal aneuploidy presenting for invasive testing.</p> <p>Inclusion criteria: pregnant women 18 years and older, with a singleton pregnancy at gestational age 10 weeks or greater, and who were planning to undergo invasive prenatal diagnosis.</p> <p>Exclusion criteria: multifetal pregnancies, pregnant women with a known maternal aneuploidy, active malignancy or a history of metastatic cancer, or those who had already undergone chorionic villus sampling or amniocentesis during the current pregnancy.</p>
Patient characteristics and setting	<p>Number enrolled: not reported. 432 maternal plasma samples were retrieved from the prospective cohort.</p> <p>Number available for 2 x 2 table: 414 samples (subgroup of 96%).</p> <p>Setting: 16 centres. Selected prenatal care centres in the USA, the Netherlands and Sweden.</p> <p>Recruitment period: not reported.</p> <p>Ethnicity: not reported.</p> <p>Mean gestational age (\pm SD; range): 15.4 (\pm 3.7; 10 to 34.1) weeks.</p> <p>Mean maternal age (\pm SD; range): 35.6 (\pm 5.7; 18.5 to 45.5) years.</p> <p>Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both.</p> <p>Language of the study: English.</p>
Index tests	<p>gNIPT by TMPS (DANSR assay) on Illumina HiSeq 2000 in 96-plex.</p> <p>Fetal fraction DNA: amount measured but not reported.</p> <p>Blood samples for gNIPT were collected before reference standard.</p> <p>Cutpoint: not reported. Usually, Harmony™ prenatal test uses FORTE algorithm; positive if FORTE risk score \geq 1%.</p> <p>Commercial test: Harmony™ Prenatal Test by Ariosa Diagnostics, Inc.</p>
Target condition and reference standard(s)	<p>Target conditions: 45,X, 47,XXY and 47,XXX. 47,XYY was also assessed but no case was found.</p> <p>Reference standard: fetal karyotype of chorionic villi or amniotic fluid.</p>
Flow and timing	<p>Blood samples were obtained prior the invasive procedure (reference standard).</p> <p>gNIPT was a second-tier test.</p> <p>18/432 samples failed during sequencing process (no gNIPT result) for low fetal fraction DNA, unusually high variation in ccfDNA counts or failure to pass the quality control measures of the DANSR assay.</p> <p>No repeated test reported.</p>
Comparative	
Aim to study	To assess the performance of a directed chromosomal analysis approach in the prenatal evaluation of fetal sex chromosome aneuploidy.
Funding source or sponsor of the study	Study funded by Ariosa Diagnostics, Inc.
Informations about the authors contacted	<p>BGI-Shenzhen were contacted on: 19 May 2016.</p> <p>Author was contacted on: 16 June 2016.</p> <p>No reply received from the author.</p>
Notes	

Hooks 2014 (Continued)

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	No		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test TMPS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear		
If a threshold was used, was it pre-specified?	Yes		
		Unclear	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all analysed patients receive the reference standard?	Yes		
Were all patients included in the analysis?	No		
		High	

Hou 2012
Study characteristics

Patient sampling	Study design: prospective cohort study. Participants: pregnant women selected at high risk of fetal aneuploidy presenting for invasive testing. Inclusion criteria: singleton pregnancies. Exclusion criteria: multifetal pregnancies.
Patient characteristics and setting	Number enrolled: 308 pregnant women. Number available for 2 x 2 table: 205 pregnant women (subgroup of 67%). Setting: 1 centre. Henan Province People's Hospital in China. Recruitment period: October 2010 to January 2012. Ethnicity: Asian. Gestational age range: 14 to 24 weeks. Mean maternal age (range): 31 (21 to 44) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both. Language of the study: Chinese.
Index tests	gNIPT by MPSS on Illumina HiSeq 2000 sequencer with BGI's algorithm. Fetal fraction DNA: not reported. Blood samples for gNIPT were collected just before reference standard. Cutpoint: not reported. Commercial test: BGI-Shenzhen's prenatal test.
Target condition and reference standard(s)	Target conditions: T21, T18, 45,X, 47,XXY and 47,XYY. T13 and 47,XXX were also assessed but no cases were found. Reference standard: fetal karyotype of amniotic fluid.
Flow and timing	Blood samples were obtained just prior the invasive procedure (reference standard). gNIPT was a second-tier test. 103/308 patients did not undergo gNIPT (no gNIPT result). No failed sample reported. No repeated test reported.
Comparative	
Aim to study	To investigate the clinical value of gNIPT using ccfDNA in maternal blood.
Funding source or sponsor of the study	Study not funded by industry but BGI-Shenzhen provided the test.
Informations about the authors contacted	Author was contacted on: 11 April 2016 (author) and 19 May 2016 (BGI's contact). No reply received from the author.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		

Hou 2012 (Continued)

Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all analysed patients receive the reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Huang 2014
Study characteristics

Patient sampling	Study design: blinded, prospective cohort study. Participants: pregnant women selected at high risk of fetal aneuploidy (as real clinical samples). Inclusion criteria: twin pregnancies with live fetuses and karyotype result. Exclusion criteria: singleton pregnancies, twins with intrauterine fetal demise at the time of sampling or without fetal karyotype result.
Patient characteristics and setting	Number enrolled: 189 pregnant women. Number available for 2 x 2 table: 189 pregnant women (whole cohort included in analyses). Setting: 7 centres. Hospitals in China. Recruitment period: April 2012 to April 2013.

Huang 2014 (Continued)

Ethnicity: most Asian.
 Median gestational age (range): 19 (11 to 39) weeks.
 Median maternal age (range): 31 (22 to 44) years.
 Chorionicity: 17% monochorionics (33/189), 80% dichorionics (152/189) and 2% unknown (4/189).

Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening.
 Language of the study: English.

Index tests	gNIPT by MPSS on Illumina Genome Analyzer IIX or HiSeq 2000 platform. Fetal fraction DNA: not reported. Blood samples for gNIPT were collected 30 minutes before reference standard. Cutpoint: positive if t score > 2.5 and L score risk > 1 (warning zone if t score > 2.5 or L score > 1). Commercial test: BGI's prenatal test.
Target condition and reference standard(s)	Target conditions: T21 and T18. Reference standard: fetal karyotype of chorionic villi (2.1%), amniotic fluid (94.2%) or cord blood (3.7%).
Flow and timing	Blood samples were obtained prior to the invasive procedure (reference standard). gNIPT was a second-tier test. No failed sample reported. No repeated test reported.
Comparative	
Aim to study	To assess the performance of noninvasive prenatal testing for trisomies 21 and 18 on the basis of MPSS of ccfDNA from maternal plasma in twin pregnancies.
Funding source or sponsor of the study	Funded by the Shenzhen Engineering Laboratory for Clinical Molecular Diagnostic, the China National GeneBank-Shenzhen, the Medical Centre for Critical Pregnant Women in Guangzhou and Prenatal monitoring, In utero therapy and Follow-up after birth in the complexity of Twin Pregnancy. Some authors worked for BGI-Shenzhen.
Informations about the authors contacted	Author was contacted on: 10 February 2016. BGI-Shenzhen were contacted on: 19 May 2016. No reply received from author.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		

Huang 2014 (Continued)

	High	Low
DOMAIN 2: Index Test MPSS		
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes	
If a threshold was used, was it pre-specified?	Yes	
	Low	Low
DOMAIN 3: Reference Standard		
Is the reference standards likely to correctly classify the target condition?	Yes	
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes	
	Low	Low
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Yes	
Did all analysed patients receive the reference standard?	Yes	
Were all patients included in the analysis?	Yes	
	Low	

Jackson 2014

Study characteristics	
Patient sampling	Study design: prospective cohort study. Participants: pregnant women selected at high risk and low risk of fetal aneuploidy presenting for screening. Inclusion criteria: not reported. Exclusion criteria: not reported.
Patient characteristics and setting	Number enrolled: 1228 pregnant women screened at first-trimester, including 1184 pregnant women with normal first-trimester ultrasound and 44 with abnormal ultrasound. Number available for 2 x 2 table: 1161 pregnant women (subgroup of 95%). Setting: 1 centre. South Shore Hospital in USA. Recruitment period: June 2012 to January 2013.

Jackson 2014 (Continued)

Ethnicity: not reported.
Gestational age: not reported.
Median maternal age: 31.5 years.
Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both.
Language of the study: English.

Index tests

gNIPT by TMPS (DANSR assay).

Fetal fraction DNA: not reported.
Blood samples for gNIPT were collected before reference standard.
Cutpoint: not reported. Usually, Harmony™ prenatal test uses FORTE algorithm; positive if FORTE risk score $\geq 1\%$.
Commercial test: Harmony™ Prenatal Test.

Target condition and reference standard(s)

Target conditions: T21, T18 and T13.
Reference standards: fetal karyotype of chorionic villi or amniotic fluid, or medical record from birth.

Flow and timing

Blood samples were obtained prior to the invasive procedure (reference standard).

gNIPT was a second-tier test.
67/1228 samples excluded of 2 x 2 tables, including 7 women with other abnormal ultrasound, 14 women opted for CVS only without gNIPT, 32 women declined all testing and 14 samples failed after 2 attempts during sequencing process (no gNIPT result).

Comparative

Aim to study

To assess the performance of nuchal translucency measurement followed by gNIPT in the first-trimester to screen for aneuploidy in a community-based average-risk population.

Funding source or sponsor of the study

Funding source not reported but 1 author is employed by Ariosa Diagnostics, Inc.

Informations about the authors contacted

Author was contacted on: 22 February 2016 and 15 March 2016.
No reply received from the author.

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Unclear		
Was a case-control design avoided?	Unclear		
Did the study avoid inappropriate exclusions?	Unclear		
		Unclear	High

DOMAIN 2: Index Test TMPS

Jackson 2014 (Continued)

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes	
If a threshold was used, was it pre-specified?	Yes	
		Low
DOMAIN 3: Reference Standard		
Is the reference standards likely to correctly classify the target condition?	Yes	
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes	
		Low
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Yes	
Did all analysed patients receive the reference standard?	Yes	
Were all patients included in the analysis?	No	
		High

Jeon 2014
Study characteristics

Patient sampling	Study design: prospective cohort study. Participants: pregnant women selected at high risk of fetal aneuploidy presenting for invasive testing. Inclusion criteria: women who gave written informed consent participated in the study if they were ≥ 19 years old and had a singleton pregnancy with a gestational age of at least 12 weeks. Exclusion criteria: multifetal pregnancies.
Patient characteristics and setting	Number enrolled: 155 pregnant women. Number available for 2 x 2 table: 155 pregnant women (whole cohort included in analyses). Setting: 1 centre. Xiamen Maternal & Child Health Care Hospital, Xiamen, Fujian, China. Recruitment period: March 2012 to October 2013. Ethnicity: Asian. Gestational age ranges: 12 to 16 weeks (18.1%), 17 to 21 weeks (55.5%), ≥ 22 weeks (26.5%). All between 12 to 24 weeks. Mean maternal age (\pm SD; range): 30.73 (\pm 4.99; 19 to 43) years.

Jeon 2014 (Continued)

Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both.
 Language of the study: English.

Index tests	gNIPT by MPSS on Ion Torrent PGM sequencer with 10 samples per chip. Fetal fraction DNA: not reported. Blood samples for gNIPT were collected just before reference standard. Cutpoint: positive if Z score > 2.566 (T21) or > 2.459 (T18). Commercial test: Genome Care's prenatal test.
Target condition and reference standard(s)	Target conditions: T21 and T18. Reference standard: fetal karyotype of amniotic fluid.
Flow and timing	Blood samples were obtained prior to the invasive procedure (reference standard). gNIPT was a second-tier test. No failed sample reported. No repeated test reported.
Comparative	
Aim to study	To investigate whether fetal T18 and T21 were sensitively and specifically detectable by semiconductor sequencer: Ion Proton™.
Funding source or sponsor of the study	Study funded by the Industrial Strategic Technology Development Program, "Bioinformatics platform development for next generation bioinformation analysis" funded by the Ministry of Knowledge Economy (MKE, Korea).
Informations about the authors contacted	Author was contacted on: 6 and 11 April 2016. Reply received on: 11 April 2016.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		

Jeon 2014 (Continued)

If a threshold was used, was it pre-specified? No

High
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? Yes

Low
Jiang 2012
Study characteristics

Patient sampling Study design: prospective cohort study.
 Participants: pregnant women at high risk of fetal aneuploidy presenting for invasive testing selected from the cohort.
 Inclusion criteria: not reported.
 Exclusion criteria: not reported.

Patient characteristics and setting Number enrolled: 903 pregnant women.
 Number available for 2 x 2 table: 903 pregnant women (whole cohort included in analyses).
 Setting: 3 centres in Shenzhen, China.
 Recruitment period: June 2009 to August 2010.
 Ethnicity: Asian.
 Gestational age range: 10 to 39 weeks.
 Maternal age range: 20 to 45 years.
 Relevant tests carried out prior to index test: not reported.
 Language of the study: English.

Index tests gNIPT by MPSS on platforms Illumina Genome Analyzer IIx or Illumina HiSeq 2000 by multiplex sequencing.

Fetal fraction DNA (range): quality control criteria > 3.5% (1% to 33%).

Jiang 2012 (Continued)

It is not reported if the blood samples for gNIPT were collected before or after reference standard.

Cutpoint:

1) Positive if binary hypothesis t score (first hypothesis) > 3 and t score (second hypothesis) < 3 and if logarithmic LR > 1 (autosomal aneuploidy).

2) Positive if t score < -2.5 (45,X and 47,XXX) without Chrom. Y representation.

3) Positive if t score > 2.5 combined with estimation of fetal ccfDNA concentration by Chrom. X and Y independently (47,XXY and 47,XYY) for male fetus.

Commercial test: NIFTY™ prenatal test by Beijing Genomics Institute (BGI).

Target condition and reference standard(s)	Target conditions: T21, T18, T13, 45,X, 47,XXY, 47,XYY and 47,XXX. Reference standard: fetal karyotype of amniotic fluid.
Flow and timing	It is not reported if the blood samples were obtained prior or after the invasive procedure (reference standard). gNIPT was a second-tier test. No failed sample reported. No repeated test reported.
Comparative	
Aim to study	To develop an advanced gNIPT method based on MPSS.
Funding source or sponsor of the study	Study funded by industry. BGI was involved in the study design, conduct of the study, analysis and interpretation of results.
Informations about the authors contacted	Author was contacted on: 19 May 2016. No reply received from the author.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Unclear		
		High	Low
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		

Jiang 2012 (Continued)

If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Unclear		
Did all analysed patients receive the reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Unclear	

Johansen 2016
Study characteristics

Patient sampling	Study design: prospective cohort study. Participants: pregnant women selected at high risk of fetal aneuploidy presenting for invasive testing. Inclusion criteria: singleton pregnancies. Exclusion criteria: multifetal pregnancies.
Patient characteristics and setting	Number enrolled: 375 pregnant women (184 for the validation set). Number available for 2 x 2 table: 173 pregnant women (subgroup of 94%). Setting: Danish public health setting. Recruitment period: not reported. Ethnicity: not reported. Median gestational age (range): 13.4 (10.6 to 31) weeks. Maternal age: not reported. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both. Language of the study: English.
Index tests	gNIPT by MPSS on Ion Proton™ sequencer in 5-plex. Fetal fraction DNA: amount measured but not reported. Blood samples for gNIPT were collected just before reference standard.

Johansen 2016 (Continued)

Cutpoint: positive if Z score ≥ 4 and WISECONDOR $\geq 1\%$ (unclassified if Z score between 3 and 4).
 In-house test.

Target condition and reference standard(s)

Target conditions: T21, T18 and T13.
 Reference standard: fetal karyotype of chorionic villi or amniotic fluid.

Flow and timing

Blood samples for gNIPT were obtained just prior the invasive procedure (reference standard).
 gNIPT was a second-tier test.

191/375 not selected, samples for the validation set were excluded.
 11/184 samples failed during sequencing process for low fetal fraction DNA (no gNIPT result).

2/173 samples were resequenced because gNIPT results were in the inconclusive zone and 2 results were obtained.

Comparative

Aim to study

To introduce gNIPT for fetal autosomal trisomies and gender in a Danish public health setting, using semi-conductor sequencing and published open source scripts for analysis.

Funding source or sponsor of the study

No funding source was reported.

Informations about the authors contacted

No need for further contact.

Notes

Methodological quality

Item

Authors' judgement

Risk of bias

Applicability concerns

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?

No

Was a case-control design avoided?

Yes

Did the study avoid inappropriate exclusions?

No

High

Low

DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard?

Yes

If a threshold was used, was it pre-specified?

Yes

Low

Low

DOMAIN 3: Reference Standard

Johansen 2016 (Continued)

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Ke 2015
Study characteristics

Patient sampling Study design: prospective cohort study.
 Participants: pregnant women considered at high risk of fetal aneuploidy.
 Inclusion criteria: singleton pregnancies. Pregnant women at high risk of fetal aneuploidy describe as follows: over age 35, the histories of abnormal pregnancy including children with T21 and repeated spontaneous abortion, stillbirth in pregnancy periods, abnormal serological screening for T21 at early and mid pregnancy, abnormal screening for fetal nuchal translucency using colour duplex ultrasonography between 11-14 weeks of gestation.
 Exclusion criteria: not reported.

Patient characteristics and setting Number enrolled: 2340 pregnant women.
 Number available for 2 x 2 table: 2340 pregnant women (whole cohort included in analyses).
 Setting: 1 centre. Clinical setting at Shenzhen Second People's Hospital in China.
 Recruitment period: March 2012 to May 2013.
 Ethnicity: Asian.
 Gestational age: positive cases were between 16 to 24 weeks. All cohort: 95% were between 15 to 20 weeks, 3% were between 12 to 14 weeks and 0.9% were ≥ 24 weeks.
 Maternal age: 88% were less than 35 years old and 12% were 35 years old or more.
 Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both.
 Language of the study: English.

Index tests gNIPT by MPSS.
 Fetal fraction DNA: not reported.
 Blood samples for gNIPT were collected before reference standard.
 Cutpoint: positive if t score > 3.
 Commercial test: BGI-Shenzhen's prenatal test.

Ke 2015 (Continued)

Target condition and reference standard(s)	Target conditions: T21, T18, and T13. Reference standards: fetal karyotype or newborn outcome.
Flow and timing	Blood samples were obtained prior to the invasive procedure (reference standard). gNIPT was a second-tier test. No failed sample reported. No repeated test reported.
Comparative	
Aim to study	To validate the efficacy of detection of fetal cell-free DNA in maternal plasma of trisomy 21, 18 and 13 in a clinical setting.
Funding source or sponsor of the study	Study not funded by industry but patients had obtained insurance plans on behalf of Shenzhen Huada Genomics Institute.
Informations about the authors contacted	Author was contacted on: 22 April 2016. No reply received from the author.

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		

Ke 2015 (Continued)

Were the reference standard results interpreted without knowledge of the results of the index tests?

Unclear

Unclear

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?

Yes

Did all analysed patients receive the reference standard?

Yes

Were all patients included in the analysis?

Yes

Low

Kim 2016
Study characteristics

Patient sampling

Study design: blinded, prospective cohort study.
 Participants: pregnant women selected at high risk of fetal aneuploidy presenting for invasive testing.
 Inclusion criteria: not reported.
 Exclusion criteria: not reported.

Patient characteristics and setting

Number enrolled: 101 pregnant women.
 Number available for 2 x 2 table: 101 pregnant women (whole cohort included in analyses).
 Setting: 3 centres (Mirae & Heemang, Namujungwon and GN hospitals) in Korea.
 Recruitment period: December 2014 to April 2015.
 Ethnicity: Asian.
 Gestational age range: 11 to 18 weeks.
 Mean maternal age (\pm SD; range): 35.45 (\pm 3.64; 25 to 42) years.
 Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening (quadruple test screening).
 Language of the study: English.

Index tests

gNIPT by MPSS on Ion Torrent PGM (data not shown in the present review) and Ion Proton™ sequencer in multiplex.

Fetal fraction DNA: not reported.
 Blood samples for gNIPT were collected before reference standard.
 Cutpoint: positive if Z score > 2.10 for Ion Proton™.
 Commercial test: Genome Care's prenatal test.

Target condition and reference standard(s)

Target condition: T21.
 Reference standard: fetal karyotype of amniotic fluid.

Flow and timing

Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard).

Kim 2016 (Continued)

gNIPT was a second-tier test.

No failed sample reported.

No repeated test reported.

Comparative

Aim to study	To compare the Ion Torrent PGM and Ion Proton™ platforms for gNIPT for fetal T21 directly using PGM and Ion Proton™ simultaneously for the same set of samples.
--------------	---

Funding source or sponsor of the study	Study funded by Genome Care internal research funding. The first author is employee of Genome Care.
--	---

Informations about the authors contacted	No need for further contact.
--	------------------------------

Notes	Data from PGM sequencer are not shown in the present review to avoid patients overlap.
-------	--

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	No		
--	----	--	--

Was a case-control design avoided?	Yes		
------------------------------------	-----	--	--

Did the study avoid inappropriate exclusions?	No		
---	----	--	--

High
Low
DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
---	-----	--	--

If a threshold was used, was it pre-specified?	No		
--	----	--	--

High
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes		
---	-----	--	--

Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
--	-----	--	--

Low
Low
DOMAIN 4: Flow and Timing

Kim 2016 (Continued)

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? Yes

Low

Korostelev 2014
Study characteristics

Patient sampling Study design: prospective cohort study.
 Participants: pregnant women selected from a population at high risk or without prior risk of fetal aneuploidy.
 Inclusion criteria: women who had a singleton pregnancy and more than 9 weeks of gestation.
 Exclusion criteria: multifetal pregnancies.

Patient characteristics and setting Number enrolled: 1968 pregnant women.
 Number available for 2 x 2 table: 685 pregnant women (subgroup of 35%).
 Setting: private clinics in Moscow, Russia.
 Recruitment period: 2012 to 2014.
 Ethnicity: not reported.
 Median gestational age (range): 14 (9 to 33) weeks.
 Mean maternal age (range): 34.4 (26 to 45) years.
 Relevant tests carried out prior to index test: biochemical screening or ultrasonography (nuchal translucency measurement) or both.
 Language of the study: English.

Index tests gNIPT by TMPS (SNP-based method) on Illumina Genome Analyzer Ix or HiSeq sequencers with NATUS algorithm.
 Fetal fraction DNA: not reported (usually NATERA used quality control criteria > 4%).
 Blood samples for gNIPT were collected before reference standard.
 Cutpoint: not reported.
 Commercial test: Natera's prenatal test.

Target condition and reference standard(s) Target conditions: T21, T18 and T13. 45,X, 47,XXY, 47,XYY and 47,XXX were also screened but inappropriate reference standard for the present review was used (data not shown in this review).
 Reference standards: fetal karyotype of chorionic villi or amniotic fluid or medical record from birth.

Flow and timing Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard).
 gNIPT was a second-tier test.
 240/1968 samples did not undergo gNIPT (no gNIPT result).
 1043/1728 samples without follow-up were excluded.
 No repeated test reported.

Comparative

Korostelev 2014 (Continued)

Aim to study	To examine possibility to use combination of gNIPT and chromosomal microarray analysis for prenatal diagnostics and their advantages between combined first-trimester screen with confirmation by karyotyping of CVS or amniocytes.
--------------	---

Funding source or sponsor of the study	Study not funded by industry but gNIPT was carried out by Natera, Inc.
--	--

Informations about the authors contacted	Author was contacted on: 21 June 2016. No reply received from the author.
--	--

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	High
DOMAIN 2: Index Test TMPS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			

Korostelev 2014 (Continued)

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? Yes

Low

Lau 2012
Study characteristics

Patient sampling

Study design: blinded, prospective cohort study.

Participants: pregnant women mostly at high risk of fetal aneuploidy presenting for invasive testing.

Inclusion criteria: not reported.

Exclusion criteria: not reported.

Patient characteristics and setting

Number enrolled: 108 pregnant women.

Number available for 2 x 2 table: 108 pregnant women (whole cohort included in analyses).

Setting: 1 centre in Japan.

Recruitment period: not reported.

Ethnicity: Asian.

Median gestational age (range): 12.7 (11.6 to 28) weeks, 89.8% < 14 weeks.

Mean maternal age (\pm SD): 37 (\pm 4.3) years.

Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening.

Language of the study: English.

Index tests

gNIPT by MPSS on Illumina HiSeq 2000 sequencer in 12-plex.

Fetal fraction DNA: not reported.

Blood samples for gNIPT were collected immediately before reference standard.

Cutpoint:

- 1) positive if Z score \geq 3 (T21, T18 and T13).
- 2) for female fetus, positive if Chrom. X Z score \leq -3 (45,X).
- 3) for female fetus, positive if Chrom. X Z score \geq 3 (47,XXX).
- 4) for male fetus, positive if Chrom. Y Z score \geq 3 (47,XXY).

Lau 2012 (Continued)

Commercial test: NIFTY™ prenatal test by BGI-Shenzhen.

Target condition and reference standard(s)	Target conditions: T21, T18, T13, 45,X and 47,XXY. 47,XYY and 47,XXX were also assessed but no case was found. Reference standard: fetal karyotype of chorionic villi or amniotic fluid.
Flow and timing	Blood samples for gNIPT were collected immediately before invasive procedure (reference standard). gNIPT was a second-tier test. No failed sample reported. No repeated test reported.
Comparative	
Aim to study	To investigate the inclusion of an internal reference in the noninvasive prenatal identification of common fetal aneuploidies using massively parallel sequencing on maternal plasma.
Funding source or sponsor of the study	Study funded by BGI-Shenzhen.
Informations about the authors contacted	BGI-Shenzhen contacted on: 19 May 2016. No reply received from the author.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Unclear		
		High	High
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			

Lau 2012 (Continued)

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? Yes

Low

Lee 2015
Study characteristics

Patient sampling Study design: blinded, prospective cohort study.
 Participants: pregnant women selected at high risk of fetal aneuploidy presenting for invasive testing.
 Inclusion criteria: pregnant women who were > 18 years old and gestational age > 8 weeks, multifetal and singleton pregnancies.
 Exclusion criteria: not reported.

Patient characteristics and setting Number enrolled: 93 pregnant women.
 Number available for 2 x 2 table: 92 pregnant women (subgroup of 99%).
 Setting: 1 centre at Asan Medical Centre, Seoul, Korea.
 Recruitment period: August 2014 to February 2015.
 Ethnicity: Asian.
 Median gestational age (range): 21.1 (8.2 to 31.1) weeks.
 Median maternal age (range): 32 (21 to 43) years.
 Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both.
 Language of the study: English.

Index tests gNIPT by MPSS on Illumina MiSeq sequencer in 12-plex or on NextSeq 500 sequencer in 96-plex.
 Median fetal fraction DNA (range): male fetus only: 10.2% (3.85% to 25.0%).
 Blood samples for gNIPT were collected before reference standard.
 Cutpoint:
 1) positive if Z score > 4 (intermediate risk if Z score between 2.5 and 4) for T21 and T18.
 2) positive if Z score > 2.8 (intermediate risk if Z score between 1.9 and 2.8) for T13.

Lee 2015 (Continued)

Commercial test: MomGuard™ by LabGenomics.

Target condition and reference standard(s)	Target conditions: T21, T18 and T13. SCA were also assessed but no case was found. Reference standards: fetal karyotype of chorionic villi, amniotic fluid, cord blood or products of conception or neonatal karyotype from peripheral blood.
Flow and timing	Blood samples for gNIPT were obtained just prior to the invasive procedure (reference standard). gNIPT was a second-tier test. 1/93 samples failed during sequencing process for low fetal fraction DNA (no gNIPT result). No repeated test reported.
Comparative	
Aim to study	To evaluate the performance of MomGuard™, a gNIPT, for detecting T21, T18, T13, and SCA abnormalities recently developed in Korea.
Funding source or sponsor of the study	Study funded by a grant from the LabGenomics Clinical Research Institute.
Informations about the authors contacted	No need for further contact.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		

Lee 2015 (Continued)

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Lefkowitz 2016
Study characteristics

Patient sampling Study design: Retrospective cohort, blinded case-control study.
 Participants: pregnant women selected at high risk of fetal aneuploidy from 4 cohorts (archived maternal plasma samples).
 Inclusion criteria: not reported.
 Exclusion criteria: cases of fetal mosaicism or incomplete karyotype or microarray information.

Patient characteristics and setting Number enrolled: 5321 pregnant women in all 4 cohorts. 1222 pregnant women selected for this study.
 Number available for 2 x 2 table: 1166 pregnant women (subgroup of 95%) for autosomes and 1144 pregnant women (subgroup of 94%) for SCA.
 Setting: multicentre.
 Recruitment period: not reported.
 Ethnicity: not reported.
 Median gestational age (range): 17 (8 to 38) weeks.
 Median maternal age (range): 36.0 (17.8 to 47) years.
 Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both.
 Language of the study: English.

Index tests gNIPT by MPSS on Illumina HiSeq 2000 in 6-plex or uniplex.
 Fetal fraction DNA: amount measured but not reported.
 Blood samples for gNIPT were collected before (for 1189 pregnant women) or after (for 24 pregnant women) reference standard.
 Cutpoint:
 1) positive for T21 if Z score ≥ 3 .
 2) positive for T18 or T13 if Z score ≥ 3.95 .
 3) positive for 45,X if Z score < -3.5 (non-reportable regions between -2.5 and -3.5).
 4) positive for 47,XXX if Z score > 3.5 (non-reportable regions between 2.5 and 3.5).

Lefkowitz 2016 (Continued)

- 5) positive for 47,XYY if Z score < -3.5 with Chrom. Y representation.
- 6) positive for 47,XXY if Z score is between -3.5 and 3.5 with Chrom. Y representation.
- Commercial test: Sequenom's test.

Target condition and reference standard(s)	Target conditions: T21, T18, T13, 45,X, 47,XXY, 47,XYY and 47,XXX. copy number variants ≥ 7 Mb were also assessed but data not shown in the present review. Reference standard: fetal karyotype of chorionic villi or amniotic fluid.
Flow and timing	Blood samples for gNIPT were obtained prior or after the invasive procedure (reference standard). gNIPT was a second-tier test. 4099/5321 samples not selected for this study. 14/1222 samples were excluded before sequencing process (11/14 samples excluded for incomplete diagnostic information and 3/14 samples excluded for confirmed mosaicism). 42/1208 samples failed during autosome sequencing process (no gNIPT result) including 11/42 failed samples for low fetal fraction DNA, 29/42 failed samples for technical reasons and 2/42 failed samples for other biological reasons (maternal event). 22/1166 samples failed SCA sequencing process (no gNIPT result). No repeated test reported.
Comparative	
Aim to study	To provide a clinical validation of the sensitivity and specificity of a novel NIPT for detection of genome wide abnormalities.
Funding source or sponsor of the study	Study funded by Sequenom, Inc.
Informations about the authors contacted	No need for further contact.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	No		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge	Yes		

Lefkowitz 2016 *(Continued)*
 of the results of the reference
 standard?

If a threshold was used, was it pre-specified?	Yes	
		Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes	
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes	
		Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes	
Did all analysed patients receive the reference standard?	Yes	
Were all patients included in the analysis?	No	
		High

Liang 2013
Study characteristics

Patient sampling	Study design: blinded, prospective cohort study. Participants: pregnant women considered at high risk for fetal T21. Inclusion criteria: singleton and twin pregnancies underwent conventional serum screening and ultrasound scanning, and who invasive prenatal diagnostics were offered. Exclusion criteria: not reported.
Patient characteristics and setting	Number enrolled: 435 pregnant women. Number available for 2 x 2 table: 412 pregnant women (subgroup of 94.7%). Setting: 3 hospitals in China. Recruitment period: March 2009 to June 2011. Ethnicity: Asian. Median gestational age (range): 21.4 (11.4 to 39.4) weeks. Most pregnant women (60%) are between 21 to 40 weeks. Only 1 case is in the first trimester (0.23%). Mean maternal age (\pm SD): 31 (\pm 5.9) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or multiple screening tests. Language of the study: English.

Liang 2013 (Continued)

Index tests	<p>gNIPT by MPSS on Illumina HiSeq 2000 in 8-plex or 12-plex.</p> <p>Fetal fraction DNA: for a Z score cutoff value of 3 for chromosome 21, fetal DNA was estimated to 5.52%.</p> <p>Blood samples for gNIPT were collected before reference standard.</p> <p>Cutpoint:</p> <ol style="list-style-type: none"> 1) positive if Z score > 3 (T21). 2) positive if Z score > 5.91 (T18). 3) positive if Z score > 5.72 (T13). 4) positive if Z score Chrom. X < -2.91 and Z score Chrom. Y < 3 (45,X). 5) positive if Z score Chrom. X range from -2.91 to +2.91 and Z score Chrom. Y > 3 (47,XXY). 6) positive if Z score Chrom. X > 2.91 and Z score Chrom. Y < 3 (47,XXX). 7) positive if Z score Chrom. X < -2.91 and Z score Chrom. Y > 3 (47,XYY). <p>Commercial test: Berry Genomics's prenatal test.</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18, T13, 45,X, 47,XXY, 47,XYY, and 47,XXX.</p> <p>Reference standard: fetal karyotype of chorionic villi (1%) or amniotic fluid (77%) or cord blood (22%).</p>
Flow and timing	<p>Blood samples were obtained prior to the invasive procedure (reference standard).</p> <p>gNIPT was a second-tier test.</p> <p>12/435 samples failed sequencing process quality control (no gNIPT result).</p> <p>11/423 samples without karyotype were excluded (no reference standard result).</p>
Comparative	
Aim to study	To determine whether gNIPT by maternal plasma DNA sequencing can uncover all fetal chromosome aneuploidies in 1 simple sequencing event.
Funding source or sponsor of the study	Study not funded by industry but Berry Genomics Co. Ltd performed the sequencing analysis for free. This study was supported by the grants from the National High Technology Research and Development Program of China (863 Program) (No.2011AA02A112), the National Key Basic Research Program of China (2012CB944600) and the National Key Technology R&D Program of China (2012BAI09B05).
Informations about the authors contacted	No need for further contact.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		

Liang 2013 (Continued)

Did the study avoid inappropriate exclusions? Yes

High
Low
DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High
Liu 2012
Study characteristics

Patient sampling	Study design: prospective cohort study. Participants: pregnant women selected at high risk of fetal aneuploidy presenting for invasive testing. Inclusion criteria: women who planned an invasive testing for 1 or more of the following reasons: abnormality in plasma test, older than 35 years old, infant deformity (ultrasound), taken drugs (teratogen) during early pregnancy or history of malformation caused by virus infection, history of birth defect caused by abnormal chromosome, history of fetus stopping growth or re-
------------------	---

Liu 2012 (Continued)

peated spontaneous abortion or dead fetus or dead birth for unknown reason, history of chromosome abnormality in family or either of the couple, too much or little amniotic fluid.
 Exclusion criteria: not reported.

Patient characteristics and setting	<p>Number enrolled: 153 pregnant women. Number available for 2 x 2 table: 153 pregnant women (whole cohort included in analyses).</p> <p>Setting: Henan Province People Hospital Medical. Recruitment period: October to November 2011. Ethnicity: Asian. Gestational age: more than 14 weeks. Mean maternal age (\pm SD; range): 32.3 (\pm 1.2; 20 to 44) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both. Language of the study: Chinese.</p>
Index tests	<p>gNIPT by MPSS on Illumina HiSeq sequencer in multiplex.</p> <p>Fetal fraction DNA: not reported. Blood samples for gNIPT were collected 30 minutes before reference standard. Cutpoint: positive if Z score \geq 3. It is not reported if gNIPT was a commercial or an in-house test.</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18, T13, 45,X and 47,XXY. 47,XXY and 47,XXX were also assess but no case were found. Reference standard: fetal karyotype of amniotic fluid.</p>
Flow and timing	<p>Blood samples for gNIPT were obtained 30 minutes prior to the invasive procedure (reference standard). gNIPT was a second-tier test.</p> <p>No failed sample reported. No repeated test reported.</p>
Comparative	
Aim to study	To determine the feasibility and accuracy of detecting numerical chromosomal abnormalities by high-flux sequencing analysis of ccfDNA from maternal plasma.
Funding source or sponsor of the study	Study funded by by the National Natural Science Foundation of China and a Medical Science and Technology Research Project of Henan Province.
Informations about the authors contacted	Author was contacted on 11 April 2016 but contact author's email is no longer valid.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		

Liu 2012 (Continued)

Did the study avoid inappropriate exclusions? Unclear

Unclear

Low

DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low

Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? Yes

Low

Ma 2016
Study characteristics

Patient sampling	Study design: blinded, prospective and retrospective cohort study. Participants: pregnant women selected from a high-risk population presenting for invasive testing (prospective cohort) and archived maternal plasma from mixed-risk (high and low risk of fetal aneuploidy) pregnant women (retrospective cohort). Inclusion criteria: singleton pregnancies with gestational age of 12 weeks or above at the time of sampling. Exclusion criteria: women with twin pregnancy or organ donation history or maternal chromosome abnormality.
------------------	---

Ma 2016 (Continued)

Patient characteristics and setting	<p>Number enrolled: 10,598 pregnant women. 2439 from prospective cohort and 8159 from retrospective cohort.</p> <p>Number available for 2 x 2 table: 10,579 pregnant women (subgroup of 99.8%).</p> <p>Setting: 20 centres. Prenatal diagnosis clinics in China.</p> <p>Recruitment period: January 2012 to January 2014 (retrospective) and February to May 2014 (prospective).</p> <p>Ethnicity: Asian.</p> <p>Median gestational age: 19 weeks.</p> <p>Median maternal age (range): 32 (16 to 53) years.</p> <p>Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both.</p> <p>Language of the study: English.</p>
Index tests	<p>gNIPT by MPSS on BGISEQ-1000 platform in 16 or 24-plex.</p> <p>Fetal fraction DNA: not reported.</p> <p>Blood samples for gNIPT were collected before reference standard.</p> <p>Cutpoint: positive if Z score > 3.</p> <p>Commercial test: BGI-Shenzhen's prenatal test.</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18 and T13.</p> <p>Reference standards: fetal karyotype of chorionic villi, amniotic fluid or cord blood, or postnatal follow-up.</p>
Flow and timing	<p>Blood samples were obtained prior to the invasive procedure (reference standard).</p> <p>gNIPT was a second- or a first-tier test.</p> <p>19/10,598 samples were excluded from the analysis including 5 from retrospective cohort (4 samples had incomplete clinical information and 1 sample failed quality control during sequencing) and 14 from prospective cohort (10 samples had incomplete clinical information and 4 samples failed quality control during sequencing).</p> <p>No repeated test reported.</p>
Comparative	
Aim to study	<p>To report the established gNIPT screening system and the clinical performance of a new ultrahigh throughput gNIPT method based on combinatorial probe-anchor ligation sequencing (cPAL) of ccfDNA in detecting T21, T18 and T13 in the multicentre network using a centralised testing mode.</p>
Funding source or sponsor of the study	<p>Some authors are employees of BGI-Shenzhen, BGI-Manufacture or BGI-DX. Study funded by Shenzhen Birth Defect Screening Project Lab, Key Laboratory of Cooperation Project in Guangdong Province, Shenzhen Municipal Government of China, Pilot projects of regional strategic emerging industry cluster development by Hubei provincial development and Reform Commission and Action plan for the development of high-tech industry in biotechnology and new medicine in 2012 by Wuhan Science and Technology Bureau.</p>
Informations about the authors contacted	<p>No need for further contact.</p>

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			

Ma 2016 (Continued)

Was a consecutive or random sample of patients enrolled? Unclear

Was a case-control design avoided? Yes

Did the study avoid inappropriate exclusions? No

High

High

DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low

Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Mazloom 2013
Study characteristics

Mazloom 2013 (Continued)

Patient sampling	<p>Study design: blinded, prospective cohort study (validation set). Participants: pregnant women selected at high risk of fetal aneuploidy. Inclusion criteria: ≥ 18 years old and singleton pregnancies between 10.5 and 20 weeks of gestation. Exclusion criteria: multifetal pregnancies, mosaic cases for sex chromosomes, or samples without documented karyotype report available.</p>
Patient characteristics and setting	<p>Number enrolled: 1975 pregnant women including 1564 in the training set (data not shown in the present review) and 411 in the validation set. Number available for 2 x 2 table: 411 pregnant women (subgroup of 95% of validation set). Setting: not reported. Recruitment period: not reported. Ethnicity: Caucasian (58.4%), Asian (18.5%), Afro-American (7.5%), other and not specified (15.6%). Median gestational age (range): 17 (8 to 29) weeks. Median maternal age (range): 36 (19 to 47) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening. Language of the study: English.</p>
Index tests	<p>gNIPT by MPSS on Illumina v3 flow cells on HiSeq 2000 sequencer in 12-plex.</p> <p>Fetal fraction DNA: amount measured but not reported. Blood samples for gNIPT were collected before reference standard. Cutpoint:</p> <ol style="list-style-type: none"> 1) positive for 45,X if Z score < -3.5 (non-reportable regions between -2.5 and -3.5). 2) positive for 47,XXX if Z score > 3.5 (non-reportable regions between 2.5 and 3.5). 3) positive for 47,XXY if Z score < -3.5 with Chrom. Y representation. 4) positive for 47,XXY if Z score is between -3.5 and 3.5 with Chrom. Y representation. <p>Commercial test: Sequenom's prenatal test.</p>
Target condition and reference standard(s)	<p>Target conditions: 45,X, 47,XXY, 47,XYY and 47,XXX. Reference standard: fetal karyotype of chorionic villi or amniotic fluid.</p>
Flow and timing	<p>Blood samples were obtained prior to the invasive procedure (reference standard). gNIPT was a second-tier test. 1564/1975 excluded samples were used for the training set. 21/411 failed samples were in the non reportable region and were considered positive gNIPT result by authors. No repeated test reported.</p>
Comparative	
Aim to study	<p>To extend the detection of autosomal aneuploidies by MPSS of ccfDNA from maternal plasma to include common sex chromosome aneuploidies.</p>
Funding source or sponsor of the study	<p>Study funded by Sequenom, Inc. and Sequenom Center for Molecular Medicine (SCMM).</p>
Informations about the authors contacted	<p>Author was contacted on: 26 May 2016. No reply received from the author.</p>
Notes	<p>Data from the training set were not shown in the present review.</p>

Mazloom 2013 (Continued)

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Unclear		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all analysed patients receive the reference standard?	Yes		
Were all patients included in the analysis?	No		
		High	

Nicolaides 2012
Study characteristics

Patient sampling	<p>Study design: retrospective study from a prospective cohort.</p> <p>Participants: selected archived plasma samples from pregnant women without prior risk of fetal aneuploidy (general population) attending for their routine first-trimester combined screening for aneuploidies.</p> <p>Inclusion criteria: singleton pregnancies between 11 to 13.9 weeks' gestation. Archived samples of at least 2 mL.</p> <p>Exclusion criteria: multifetal pregnancies.</p>
Patient characteristics and setting	<p>Number enrolled: 2230 pregnant women.</p> <p>Number available for 2 x 2 table: 1949 pregnant women (subgroup of 87%).</p> <p>Setting: not reported.</p> <p>Recruitment period: October 2010 to January 2011.</p> <p>Ethnicity: Caucasian (69.8%), African (20.6%), South Asian (4%), East Asian (2.8%) and mixed (2.8%).</p> <p>Gestational age range: 11 to 13.9 weeks.</p> <p>Median maternal age (range): 31.8 (27.7 to 35.4) years.</p> <p>Relevant tests carried out prior to index test: none.</p> <p>Language of the study: English.</p>
Index tests	<p>gNIPT by TMPS (DANSR assay).</p> <p>Median fetal fraction DNA (interquartile range): euploids: 10.0% (7.8% to 13.0%), T21: 12.5% (9.2% to 21.3%), and T18: 9.3% (5.6% to 13.0%).</p> <p>Blood samples for gNIPT were collected before reference standard.</p> <p>Cutpoint: positive if FORTE algorithm risk score \geq 1%.</p> <p>Commercial test: Harmony™ Prenatal test by Ariosa Diagnostics, Inc.</p> <p>The traditional screening test (combined test at the first trimester) was also assessed.</p> <p>Cutpoint of combined test: 1 in 150.</p>
Target condition and reference standard(s)	<p>Target conditions: T21 and T18.</p> <p>Reference standards: fetal karyotype of chorionic villi or amniotic fluid or neonatal clinical examination.</p>
Flow and timing	<p>Blood samples were obtained prior to the invasive procedure (reference standard).</p> <p>gNIPT was a first-tier test.</p> <p>181/2230 samples were ineligible (no fetal karyotype or follow-up, miscarriage, stillbirth, termination of pregnancy or other abnormalities).</p> <p>100/2049 samples failed during sequencing process including 46 for low fetal DNA and 54 had assay failures (no gNIPT result).</p> <p>No repeated test reported.</p>
Comparative	
Aim to study	To assess performance of noninvasive prenatal testing for fetal trisomy in a routinely screened first-trimester pregnancy population.
Funding source or sponsor of the study	The study was supported by a grant from the Fetal Medicine Foundation (UK). The cost of collection and analysis of the samples was covered by Ariosa Diagnostics, Inc.
Informations about the authors contacted	No need for further contact.

Nicolaides 2012 (Continued)

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test TMPS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 2: Index Test Traditional screening tests			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		

Nicolaides 2012 (Continued)

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Nicolaides 2013
Study characteristics

Patient sampling Study design: blinded, prospective cohort study.
 Participants: pregnant women selected at high risk of fetal aneuploidy presenting for invasive testing.
 Inclusion criteria: singleton pregnancies at high risk of fetal aneuploidy between 11 to 13 weeks' gestation.
 Exclusion criteria: multifetal pregnancies.

Patient characteristics and setting Number enrolled: 242 pregnant women.
 Number available for 2 x 2 table: 229 pregnant women (subgroup of 95%).
 Setting: 1 centre. Fetal Medicine Centre, in UK.
 Recruitment period: not reported.
 Ethnicity: not reported.
 Median gestational age (range): 13.1 (11.3 to 13.9) weeks.
 Median maternal age (range): 35.7 (18.5 to 46.5) years.
 Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening.
 Language of the study: English.

Index tests gNIPT by TMPS (SNP-based method) on Illumina Genome Analyzer Ix or HiSeq sequencers with NATUS algorithm.
 Fetal fraction DNA: the lowest fetal fraction DNA on a case that returned a result was 3.95%.
 Blood samples for gNIPT were collected immediately before reference standard.
 Cutpoint: not reported.
 Commercial test: Natera's prenatal test.

Target condition and reference standard(s) Target conditions: T21, T18, 45,X, 47,XXY, 47,XYY and 47,XXX were also assessed but no case was found. T13 was also assessed but the only 1 case presented in this publication was published thereafter in [Hall 2014](#). T13 case was excluded to avoid double counting.
 Reference standard: fetal karyotype of chorionic villi.

Flow and timing Blood samples were obtained just before the invasive procedure (reference standard).

gNIPT was a second-tier test.
 13/242 samples failed sequencing process quality control (no gNIPT result).

No repeated test reported.

1 T13 cases was excluded to avoid double counting because it was published thereafter in [Hall 2014](#).

Comparative

Aim to study To assess the performance of ccfDNA testing in maternal blood for detection of fetal aneuploidy of chromosomes 13, 18, 21, X, and Y using TMPS of single-nucleotide polymorphisms.

Nicolaides 2013 (Continued)

Funding source or sponsor of the study	Study funded by a grant from the Fetal Medicine Foundation (UK Charity No: 1037116). Analysis of samples was performed at their own expense by Natera, Inc.
Informations about the authors contacted	No need for further contact.
Notes	T13 cases data are not shown in the present review. They were excluded to avoid double counting because they are also published in Hall 2014 .

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Unclear		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test TMPS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		

Nicolaides 2013 *(Continued)*

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Nicolaides 2014a
Study characteristics

Patient sampling	<p>Study design: case-control study.</p> <p>Participants: pregnant women selected from a high-risk population (archived maternal plasma samples).</p> <p>Inclusion criteria: singleton pregnancies.</p> <p>Exclusion criteria: cases of fetal mosaicism and multifetal pregnancies.</p>
Patient characteristics and setting	<p>Recruited participants: 177 archived maternal plasma.</p> <p>Number available for 2 x 2 table: 172 samples (subgroup of 97%).</p> <p>Setting: recruitment in London, UK.</p> <p>Ethnicity: Caucasian (90%), Afro-Caribbean (4%), Asian (5%) and other (1%).</p> <p>Gestational age range: 11.2 to 14.1 weeks.</p> <p>Maternal age range: 17.3 to 47.8 years.</p> <p>Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening.</p> <p>Language of the study: English.</p>
Index tests	<p>gNIPT by TMPS (DANSR assay) on Illumina HiSeq 2000 in 96-plex.</p> <p>Median fetal fraction DNA (range): euploids: 13.0% (4.8% to 32.0%), 45,X: 10.0% (6.3% to 18.0%), and 47,XXX, 47,XXY, and 47,XYY: 12.0% (6.4% to 16.0%).</p> <p>Blood samples for gNIPT were collected just before reference standard.</p> <p>Cutpoint: positive if FORTE algorithm risk score \geq 1%.</p> <p>Commercial test: Harmony™ Prenatal Test by Ariosa Diagnostics, Inc.</p>
Target condition and reference standard(s)	<p>Target conditions: 45,X, 47,XXX, 47,XXY, and 47,XYY.</p> <p>Reference standard: fetal karyotype of chorionic villi or amniotic fluid.</p>
Flow and timing	<p>Blood samples for gNIPT were collected just before invasive procedure (reference standard).</p> <p>gNIPT was a second-tier test.</p> <p>5/177 samples failed during sequencing process (no gNIPT result), including 1 sample failed laboratory quality control metrics and 4 samples failed for an insufficient fetal cfDNA fraction.</p>

Nicolaides 2014a (Continued)

No repeated test reported.

Comparative

Aim to study	To report the clinical performance of chromosome-selective sequencing of cfDNA in maternal blood and the FORTE algorithm for the assessment of fetal sex chromosome aneuploidies.
--------------	---

Funding source or sponsor of the study	No funding source was reported.
--	---------------------------------

Informations about the authors contacted	Author was contacted on: 10 February 2016. No reply received from the author.
--	--

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	No		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test TMPS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			

Nicolaides 2014a (Continued)

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Norton 2012
Study characteristics

Patient sampling Study design: blinded, prospective cohort study.
 Participants: pregnant women selected at high risk of fetal aneuploidy presenting for invasive testing.
 Inclusion criteria: pregnant women aged ≥ 18 years, at gestational age ≥ 10 weeks, with a singleton pregnancy, who were planning to undergo invasive prenatal diagnosis for any indication.
 Exclusion criteria: multifetal pregnancies, women with known aneuploidy, had active malignancy or a history of metastatic cancer, or had already undergone CVS or amniocentesis during the current pregnancy.

Patient characteristics and setting Number enrolled: 4002 pregnant women.
 Number available for 2 x 2 table: 3080 pregnant women (subgroup of 77%).
 Setting: 48 centres. Selected prenatal care Centres in USA, the Netherlands and Sweden.
 Recruitment period: not reported.
 Ethnicity: Caucasian (49.6%), Afro-American (6.4%), Asian (13.4%), Hispanic (22.7%) and other (7.9%).
 Mean gestational age (\pm SD; range): 16.9 (\pm 4.1; 10 to 38.7) weeks.
 Mean maternal age (\pm SD; range): 34.3 (\pm 6.4; 18 to 50) years.
 Relevant test carried out prior to index test: not reported.
 Language of the study: English.

Index tests gNIPT by TMPS (DANSR assay) on Illumina HiSeq 2000 in 96-plex.
 Mean fetal fraction DNA (\pm SD; range): euploids: 11% (\pm 4.5%; 4.2% to 51.3%), T21: 11.6% (\pm 4.2%; 5.1% to 23.3%), and T18: 10% (\pm 3.8%; 4.9% to 20.8%).
 Blood samples for gNIPT were collected before reference standard.
 Cutpoint: positive if FORTE algorithm risk score $\geq 1\%$.
 Commercial test: Ariosa Diagnostics, Inc's prenatal test.

Target condition and reference standard(s) Target conditions: T21 and T18.
 Reference standard: fetal karyotype of chorionic villi (74.7%) or amniotic fluid (25.3%).

Flow and timing Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard).
 gNIPT was a second-tier test.
 774/4002 samples excluded for ineligible criteria.
 148/3228 samples failed during sequencing process (no gNIPT result), including 57 samples failed for low fetal fraction DNA and 91 samples failed sequencing process.
 No repeated test reported.

Norton 2012 (Continued)

Comparative

Aim to study	To evaluate performance of a gNIPT of fetal T21 and T18.
--------------	--

Funding source or sponsor of the study	Study funded by Ariosa Diagnostics, Inc.
--	--

Informations about the authors contacted	No need for further contact.
--	------------------------------

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	No		
--	----	--	--

Was a case-control design avoided?	Yes		
------------------------------------	-----	--	--

Did the study avoid inappropriate exclusions?	No		
---	----	--	--

High
Low
DOMAIN 2: Index Test TMPS

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
---	-----	--	--

If a threshold was used, was it pre-specified?	Yes		
--	-----	--	--

Low
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes		
---	-----	--	--

Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
--	-----	--	--

Low
Low
DOMAIN 4: Flow and Timing

Norton 2012 *(Continued)*

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Norton 2015
Study characteristics

Patient sampling Study design: blinded, prospective cohort study.

Participants: unselected population of pregnant women undergoing aneuploidy screening (without prior risk of fetal aneuploidy).

Inclusion criteria: singleton pregnancies, at least 18 years of age, and between 10 to 14 weeks of gestation.

Exclusion criteria: women who had a miscarriage, chose to terminate the pregnancy or had a still-birth without confirmatory genetic testing.

Patient characteristics and setting Number enrolled: 18,955 pregnant women.

Number available for 2 x 2 table: 15,841 pregnant women (subgroup of 84%).

Setting: 35 centres in USA States, Canada, Sweden, the Netherlands, Belgium, and Italy.

Recruitment period: March 2012 to April 2013.

Ethnicity: Caucasian (70.9%), Afro-American (8.2%), Asian (10.5%), Native American (0.6%), multiracial (2.7%), other (6.7%) and missing data (0.5%).

Mean gestational age (range): 12.5 (10.0 to 14.3) weeks.

Mean maternal age (range): 31 (18 to 48) years whose 76% of pregnant women analysed had < 35 years old.

Relevant tests carried out prior to index test: none.

Language of the study: English.

Index tests gNIPT by TMPS (DANSR assay) on Illumina HiSeq 2000 sequencer in 96-plex.

Fetal fraction DNA: amount measured but not reported.

Blood samples for gNIPT were collected before reference standard.

Cutpoint: not reported. Usually, Harmony™ prenatal test uses FORTE algorithm; positive if FORTE risk score ≥ 1%.

Commercial test: Harmony™ Prenatal Test by Ariosa Diagnostics, Inc.

The traditional screening tests (combined test at the first trimester) were also assessed.

Norton 2015 (Continued)

Cutpoint of combined test: 1 in 270 for T21 or 1 in 150 for T18 and T13.

Target condition and reference standard(s)	<p>Target conditions: T21, T18 and T13.</p> <p>Reference standards: fetal karyotype of chorionic villi, amniotic fluid or products of conception or neonatal karyotype, neonatal clinical examination or medical record from birth.</p>
Flow and timing	<p>Blood samples were obtained prior to the invasive procedure (reference standard).</p> <p>gNIPT was a first-tier test.</p> <p>3114/18,955 samples excluded of analysis including 229 samples did not meet inclusion criteria or meet exclusion criteria, 31 had twins, 121 had unknown ovum-donor status, 64 withdrew or were withdrawn by investigator, 384 had sample-handling errors, 308 without standard screening test result, 488 failed sequencing and have no gNIPT result (192 for low fetal fraction DNA, 83 for non fetal fraction DNA and 213 for high assay variance or assay failures) and 1489 were lost to follow-up.</p>
Comparative	
Aim to study	<p>To test the hypothesis that gNIPT has better performance than standard first-trimester screening (with measurement of nuchal translucency and biochemical analytes) in risk assessment for trisomy 21 in a large, unselected population of women presenting for aneuploidy screening.</p> <p>To also evaluate the performance of gNIPT and standard screening in the assessment of risk for trisomies 18 and 13.</p>
Funding source or sponsor of the study	Study funded by Ariosa Diagnostics, Inc and Perinatal Quality Foundation.
Informations about the authors contacted	<p>Author was contacted on: 10 February 2016.</p> <p>Reply received on: 11 February 2016.</p>
Notes	This study is a clinical trial (Noninvasive Examination of Trisomy (NEXT) ClinicalTrials.gov number, NCT01511458).

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Unclear		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test TMPS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		

Norton 2015 (Continued)

If a threshold was used, was it pre-specified? Yes

Low

Low

DOMAIN 2: Index Test Traditional screening tests

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low

Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? No

Were all patients included in the analysis? No

High

Palomaki 2012
Study characteristics

Patient sampling	Study design: nested case-control (1:3) study. Participants: pregnant women at high risk of fetal aneuploidy presenting for invasive testing were selected. Inclusion criteria: ≥ 18 years old, between about 10 weeks and 21 weeks 6 days of gestation, at high risk of aneuploidies and who underwent a diagnostic procedure. Exclusion criteria: multifetal pregnancies or low risk of fetal aneuploidy.
------------------	---

Palomaki 2012 (Continued)

Patient characteristics and setting	<p>Number enrolled: 4664 pregnant women. 1776 pregnant women selected for this study and 212 re-analysed samples from Palomaki 2011.</p> <p>Number available for 2 x 2 table: 1971 pregnant women (1759 from this study + 212 from Palomaki 2011) (subgroup of 42%).</p> <p>Setting: 27 centres. Prenatal diagnostic centres (Canada, Italy, Spain, Czech Republic, Argentina, Ireland, Hungary, USA, Israel and Australia).</p> <p>Recruitment period: April 2009 to February 2011.</p> <p>Ethnicity (only for 293 pregnant women): Caucasian (84.9%), Afro-american (4.1%), Asian (5.5%) and unknown (5.5%).</p> <p>Mean gestational age (range): 14.7 (9 to 22) weeks.</p> <p>Mean maternal age (\pm SD): 37.2 (\pm 5) years.</p> <p>Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both.</p> <p>Language of the study: English.</p>
Index tests	<p>gNIPT by MPSS on Illumina HiSeq 2000 sequencer in 4-plex.</p> <p>Mean (geometric) fetal fraction DNA (range): 13.4% (4% to 50%).</p> <p>Blood samples for gNIPT were collected before reference standard.</p> <p>Cutpoint: positive if Z score > 3 (T21), > 3.88 (T18) or > 7.17 (T13).</p> <p>Commercial test: Sequenom's prenatal test.</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18 and T13.</p> <p>Reference standard: fetal karyotype of chorionic villi, amniotic fluid or products of conception.</p>
Flow and timing	<p>Blood samples were obtained immediately prior the invasive procedure (reference standard).</p> <p>gNIPT was a second-tier test.</p> <p>2888/4664 samples were not selected for this study.</p> <p>110/1776 samples failed the initial MPSS testing.</p> <p>105/110 samples required repeat testing using a second aliquot and 5/110 samples were re-sequenced with the same library. 93/110 samples obtained a gNIPT results.</p> <p>17/1776 samples failed during sequencing process, most for low fetal fraction DNA (no gNIPT result).</p>
Comparative	
Aim to study	To determine whether maternal plasma ccfDNA sequencing can identify T18 and T13 as well as T21.
Funding source or sponsor of the study	Study fully funded by Sequenom, Inc.
Informations about the authors contacted	No need for further contact.
Notes	This study is a clinical trial "A New Prenatal Blood Test for Down Syndrome" ClinicalTrials.gov number: NCT00877292.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		

Palomaki 2012 *(Continued)*

Was a case-control design avoided? No

Did the study avoid inappropriate exclusions? No

High
Low
DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High
Papageorghiou 2016a
Study characteristics

Patient sampling Study design: blinded, case-control study (1:9).
 Participants: pregnant women selected at high risk of fetal aneuploidy presenting for invasive testing.
 Inclusion criteria: at least 18 years of age, singleton or twin pregnancies of at least 10 weeks' gestation and a clinical indication for an invasive procedure.

Papageorghiou 2016a (Continued)

Exclusion criteria: higher-order multiple pregnancies (triplets or more), known mosaicism, partial trisomy or translocations, fetal demise, disappearing twin, malignancy or known aneuploidy in the pregnancy.

Patient characteristics and setting	<p>Number enrolled: 442 pregnant women. Number available for 2 x 2 table: 426 singleton pregnancies (subgroup of 96%). Setting: 6 hospital centres in England, UK. Recruitment period: April 2008 to November 2014. Ethnicity: not reported. Median gestational age (range): 15.4 (11 to 36.6) weeks. Median maternal age (range): 35 (18 to 55) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both. Language of the study: English.</p>
Index tests	<p>gNIPT by MPSS on Ion Proton™ sequencer in 8-plex.</p> <p>Fetal fraction DNA: amount measured but not reported. Blood samples for gNIPT were collected before reference standard. Cutpoint: positive if likelihood ratio > 1 and maternal age-adjusted probability risk score. Commercial test: IONA® test by Premaitha Health (public limited company).</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18 and T13. Reference standards: fetal karyotype of chorionic villi or amniotic fluid or medical record from birth.</p>
Flow and timing	<p>Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard). gNIPT was a second-tier test. 5/442 samples failed during sequencing process including 3 samples for low fetal fraction DNA and 2 samples did not have sufficient DNA fragment counts (no gNIPT result).</p> <p>11/437 twin pregnancies were not selected.</p> <p>No repeated test reported.</p>
Comparative	
Aim to study	<p>To investigate the accuracy of the IONA® test in the discrimination between euploid pregnancies and those affected by fetal trisomies 21, 18 and 13.</p>
Funding source or sponsor of the study	<p>Study funded by Premaitha Health (public limited company). Some authors are employees of Premaitha Health plc.</p>
Informations about the authors contacted	<p>Author was contacted on: 19 September 2016. Reply received on: 20 September 2016.</p>
Notes	<p>Data from singleton pregnancies only reported here. See Papageorghiou 2016b for data on twin pregnancies.</p>

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		

Papageorghiou 2016a *(Continued)*

Was a case-control design avoided? No

Did the study avoid inappropriate exclusions? No

High
Low
DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High
Papageorghiou 2016b
Study characteristics

Patient sampling Study design: blinded, case-control study (1:9).
 Participants: pregnant women selected at high risk of fetal aneuploidy presenting for invasive testing.
 Inclusion criteria: at least 18 years of age, a singleton or twin pregnancies of at least 10 weeks' gestation and a clinical indication for an invasive procedure.

Papageorghiou 2016b (Continued)

Exclusion criteria: higher-order multiple pregnancies (triplets or more), known mosaicism, partial trisomy or translocations, fetal demise, disappearing twin, malignancy or known aneuploidy in the pregnancy.

Patient characteristics and setting	<p>Number enrolled: 442 pregnant women. Number available for 2 x 2 table: 11 twin pregnancies (subgroup of 2%). Setting: 6 hospital centres in England, UK. Recruitment period: April 2008 to November 2014. Ethnicity: not reported. Median gestational age (range): 15.4 (11 to 36.6) weeks. Median maternal age (range): 35 (18 to 55) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both. Language of the study: English.</p>
Index tests	<p>gNIPT by MPSS on Ion Proton™ sequencer in 8-plex. Fetal fraction DNA: amount measured but not reported. Blood samples for gNIPT were collected before reference standard. Cutpoint: positive if likelihood ratio > 1 and maternal age-adjusted probability risk score. Commercial test: IONA® test by Premaitha Health (public limited company).</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18 and T13. Reference standards: fetal karyotype of chorionic villi or amniotic fluid or medical record from birth.</p>
Flow and timing	<p>Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard). gNIPT was a second-tier test. 5/442 samples failed during sequencing process including 3 samples for low fetal fraction DNA and 2 samples did not have sufficient DNA fragment counts (no gNIPT result). 426/437 singleton pregnancies were not selected. No repeated test reported.</p>
Comparative	
Aim to study	<p>To investigate the accuracy of the IONA® test in the discrimination between euploid pregnancies and those affected by fetal trisomies 21, 18 and 13.</p>
Funding source or sponsor of the study	<p>Study funded by Premaitha Health (public limited company). Some authors are employees of Premaitha Health plc.</p>
Informations about the authors contacted	<p>Author was contacted on: 19 September 2016. Reply received on: 20 September 2016.</p>
Notes	<p>Data from twin pregnancies only reported here. Data from singleton pregnancies reported in Papageorghiou 2016a.</p>

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		

Papageorghiou 2016b *(Continued)*

Was a case-control design avoided? No

Did the study avoid inappropriate exclusions? No

High
Low
DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High
Pergament 2014
Study characteristics

Patient sampling Study design: blinded, prospective cohort study.
 Participants: pregnant women from a population with mixed risk of fetal aneuploidy presenting for aneuploidy screening (51% high risk and 49% low risk).
 Inclusion criteria: women were 18 years of age or older with a singleton pregnancy of at least 7 weeks of gestation and signed an informed consent.

Pergament 2014 (Continued)

Exclusion criteria: women with confirmed sex chromosome abnormality (47,XXX, XXY, XYY), confirmed triploidy, confirmed fetal mosaicism or multifetal pregnancy or egg donor.

Patient characteristics and setting	<p>Number enrolled: 1064 pregnant women. Number available for 2 x 2 table: 963 pregnant women for T21, 964 for T18 and 45,X and 965 for T13 (subgroup of 91%). Setting: 35 centres. Prenatal care centres worldwide in Czech Republic, Japan, USA, Ireland and Spain. Recruitment period: not reported. Ethnicity: not reported. Mean gestational age (\pm SD; range): 17.0 (\pm 4.1; 7.6 to 40.6) weeks. Median gestational age: 14.3 weeks. Mean maternal age (\pm SD; range): 30.3 (\pm 7.4; 18 to 47) years. Median maternal age: 30.0 years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both. Language of the study: English.</p>
Index tests	<p>gNIPT by TMPS (SNP-based method) on Illumina Genome Analyzer IIx or HiSeq sequencers, 19,488-plex targeted PCR with NATUS algorithm.</p> <p>Range fetal fraction DNA: 2% to 50%. Blood samples for gNIPT were collected before (93%) or 4 days or later after (7%) reference standard. Cutpoint: not reported. Commercial test: Natera's prenatal test.</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18, T13 and 45,X. Reference standards: fetal karyotype with confirmatory fluorescence in situ hybridisation or cytogenetic karyotype analysis or by genetic testing of cord blood, buccal sample, saliva, or products of conception, post-natal or post-live birth follow-up.</p>
Flow and timing	<p>Blood samples for gNIPT were obtained prior (93%) or after (7%) to the invasive procedure (reference standard). gNIPT was a second-tier test.</p> <p>13/1064 samples excluded for other aneuploidies, including 6 cases with triploidy, 3 fetal mosaics, 2 cases with 47,XXY, 1 case with 47,XXX and 1 case with 47,XYY.</p> <p>85/1051 samples failed quality control (no gNIPT result) including 64 low fetal fraction DNA, 12 low DNA, 6 contaminations, 2 loss of heterozygosity and 1 poor model fit.</p> <p>Between 1 to 3 samples did not passed quality control for all 5 chromosomes.</p> <p>No repeated test reported.</p>
Comparative	
Aim to study	<p>To estimate performance of a single nucleotide polymorphism-based gNIPT (TMPS) for fetal aneuploidy in high-risk and low-risk populations on single venipuncture.</p>
Funding source or sponsor of the study	<p>Study funded by Natera, Inc. and a grant from the National Institute of Health, National Institute of Child Health and Human Development (4R44HD062114-02). The majority of the authors are employees of Natera, Inc. and hold stock or options to hold stock in the company.</p>
Informations about the authors contacted	<p>Author was contacted on: 22 June 2016. No reply received from the author.</p>
Notes	

Methodological quality

Pergament 2014 (Continued)

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Unclear		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	High
DOMAIN 2: Index Test TMPS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all analysed patients receive the reference standard?	Yes		
Were all patients included in the analysis?	No		
		High	

Persico 2016
Study characteristics

Patient sampling	Study design: blinded, prospective cohort study. Participants: pregnant women selected from a high-risk population. Inclusion criteria: singleton pregnancies. Exclusion criteria: multifetal pregnancies.
Patient characteristics and setting	Number enrolled: 259 pregnant women. Number available for 2 x 2 table: 249 pregnant women (subgroup of 96%). Setting: 4 fetal medicine centres in Italy. Recruitment period: March to December 2014. Ethnicity: not reported. Gestational age: not reported. Median maternal age (range): 36 (20 to 46) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both. Language of the study: English.
Index tests	gNIPT by TMPS (SNP-based method) on Illumina Genome Analyzer IIX or HiSeq sequencers, 19,488-plex targeted PCR with NATUS algorithm. Fetal fraction DNA: amount measured but not reported (usually NATERA used quality control criteria > 4%). Blood samples for gNIPT were collected just before reference standard. Cut-off value: positive if risk score > 1%. Commercial test: Natera's prenatal test.
Target condition and reference standard(s)	Target conditions: T21, T18, T13, 45,X, 47,XXY and 47,XXX. 47,YYY was also assessed but no case was found. Reference standard: fetal karyotype of chorionic villi or amniotic fluid.
Flow and timing	Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard). gNIPT was a second-tier test. 10/259 samples failed during sequencing process (no gNIPT result) including 2 samples failed internal quality control and 8 samples had low fetal fraction DNA. No repeated test reported.
Comparative	
Aim to study	To investigate a strategy for clinical implementation of ccfDNA testing in high-risk pregnancies after first-trimester combined screening.
Funding source or sponsor of the study	Study not funded by industry but the cost of ccfDNA testing were covered by Natera, Inc.
Informations about the authors contacted	No need for further contact.

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

DOMAIN 1: Patient Selection

Persico 2016 (Continued)

Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test TMPS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all analysed patients receive the reference standard?	Yes		
Were all patients included in the analysis?	No		
		High	

Poon 2016
Study characteristics

Patient sampling	Study design: retrospective cohort, blinded nested case-control study. Participants: archived maternal plasma from pregnant women selected at high risk of fetal aneuploidy presenting for invasive testing (CVS). Inclusion criteria: singleton pregnancies. Exclusion criteria: multifetal pregnancies.
------------------	--

Poon 2016 (Continued)

Patient characteristics and setting	Number enrolled: 242 pregnant women. Number available for 2 x 2 table: 241 pregnant women (subgroup of 99.6%). Setting: 1 centre at King's College Hospital, London, UK. Recruitment period: April 2007 to June 2012. Ethnicity: Caucasian (75%), Afro-Caribbean (17%), Asian (5%) and mixed (3%). Median gestational age (range): 12.7 (11.4 to 13.6) weeks. Median maternal age (range): 33.7 (29.2 to 40.5) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both. Language of the study: English.
Index tests	gNIPT by MPSS on Ion Proton™ sequencer. Fetal fraction DNA: amount measured but not reported. Blood samples for gNIPT were collected just before reference standard. Cutpoint: not reported but authors used the same prenatal test than Papageorghiou 2016a (cutpoint: positive if likelihood ratio > 1 and maternal age-adjusted probability risk score). Commercial test: IONA® test by Premaitha Health (public limited company).
Target condition and reference standard(s)	Target conditions: T21, T18 and T13. Reference standard: fetal karyotype of chorionic villi.
Flow and timing	Blood samples for gNIPT were obtained just before the invasive procedure (reference standard). gNIPT was a second-tier test. 1/242 samples failed for low fetal fraction DNA (no gNIPT result). No repeated test reported.
Comparative	
Aim to study	To assess the potential performance of screening for fetal T21, T18 and T13 by ccfDNA analysis of maternal blood using the IONA® test.
Funding source or sponsor of the study	Study not funded by industry but the IONA® test was provided by Premaitha Health plc, Manchester, UK. Study supported by a grant from The Fetal Medicine Foundation.
Informations about the authors contacted	Author was contacted on: 19 September 2016. No reply received from the author.

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	No		
Did the study avoid inappropriate exclusions?	No		
		High	Low

Poon 2016 (Continued)

DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High
Porreco 2014
Study characteristics

Patient sampling Study design: blinded, prospective cohort, observational study.
 Participants: pregnant women selected at high risk of fetal aneuploidy presenting for invasive testing when research personnel have been available.
 Inclusion criteria: singleton pregnancy in a patient 18 years of age or older who had provided written informed consent and who had made the decision to pursue invasive prenatal diagnosis by CVS or amniocentesis.
 Exclusion criteria: inability to give written informed consent, multifetal pregnancies, or fetal demise of an additional embryo during the current pregnancy at 8 weeks or more of gestation.

Patient characteristics and setting Number enrolled: 4170 pregnant women.
 Number available for 2 x 2 table: 3322 for autosomes (subgroup of 80%), 3278 for 45,X and 47,XXX (subgroup of 79%) and 3201 for 47,XXY and 47,XYY (subgroup of 77%).
 Setting: 31 centres in USA.
 Recruitment period: September 2009 to April 2011.
 Ethnicity: Caucasian (60,1%), Asian (18,7%), Afro-American (4,5%) and other (16.7%).

Porreco 2014 (Continued)

Mean gestational age (\pm SD; range): 16.3 (\pm 3.5; 9.0 to 37.0).
 Mean maternal age (\pm SD; range): 35.1 (\pm 5.6; 18.0 to 50.0).
 Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening.
 Language of the study: English.

Index tests	<p>gNIPT by MPSS on Illumina HiSeq 2000 in 12-plex.</p> <p>Range fetal fraction DNA: 4% to 50%. Blood samples for gNIPT were collected before reference standard. Cutpoint:</p> <ol style="list-style-type: none"> 1) for T21, positive if Z score \geq 3. 2) for T18 and T13, positive if Z score \geq 3.95. 3) positive for 45,X if Z score $<$ -3.5 (non-reportable regions between -2.5 and -3.5). 4) positive for 47,XXX if Z score $>$ 3.5 (non-reportable regions between 2.5 and 3.5). 5) positive for 47,XXY if Z score risk $<$ -3.5 with Chrom. Y representation. 6) positive for 47,XXY if Z score risk is between -3.5 and 3.5 with Chrom. Y representation. <p>Commercial test: Sequenom's prenatal test.</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18, T13, 45,X, 47,XXX, 47,XXY and 47,XXY. Reference standards: fetal karyotype of chorionic villi or amniotic fluid, or medical record from birth.</p>
Flow and timing	<p>Blood samples were obtained prior to the invasive procedure (reference standard).</p> <p>gNIPT was a second-tier test. 740/4170 samples excluded before sequencing process including 320 samples for insufficient sample volume, 120 samples processed outside of the 6 hours laboratory process window, 270 failed laboratory quality control set, 24 for incomplete case report form and 6 without invasive procedure performed).</p> <p>For autosomes: 54/3430 autosomes samples excluded for quality control deviation (low fetal DNA fraction, library concentration, total counts, and amplification bias).</p> <p>For autosomes: 54/3376 samples excluded for complex autosome karyotypes (mosaic, triploidies, unbalanced rearrangements with missing or duplicated genetic material).</p> <p>For 45,X and 47,XXX: 102/3430 samples excluded for low fetal fraction DNA or copy number variation of the Chrom. X is confounded by maternal component and cannot be determined.</p> <p>For 45,X and 47,XXX: 50/3328 samples excluded for complex SCA karyotype.</p> <p>For 47,XXY and 47,XXY: 182/3430 samples excluded for low fetal fraction DNA or copy number variation of the Chrom. X is confounded by maternal component and cannot be determined.</p> <p>For 47,XXY and 47,XXY: 47/3248 samples excluded for complex SCA karyotype.</p> <p>No repeated test reported.</p>
Comparative	
Aim to study	<p>To validate the clinical performance of MPSS of ccfDNA contained in specimens from pregnant women at high risk of fetal aneuploidy to test fetuses for T21, T18, T13, 45,X, 47,XXX, 47,XXY and 47,XXY.</p>
Funding source or sponsor of the study	<p>Study funded by Sequenom, Inc.</p>

Porreco 2014 (Continued)

Informations about the authors contacted
 Author was contacted on: 30 May 2016.
 Reply received on: 31 May 2016.

Notes
 This study is a clinical trial (Non-Invasive Screening for Fetal Aneuploidy) ClinicalTrials.gov number, NCT00847990.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	No		
--	----	--	--

Was a case-control design avoided?	Yes		
------------------------------------	-----	--	--

Did the study avoid inappropriate exclusions?	No		
---	----	--	--

High

Low

DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
---	-----	--	--

If a threshold was used, was it pre-specified?	Yes		
--	-----	--	--

Low

Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes		
---	-----	--	--

Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
--	-----	--	--

Low

Low

DOMAIN 4: Flow and Timing

Porreco 2014 (Continued)

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Quezada 2015
Study characteristics

Patient sampling Study design: prospective cohort study.
 Participants: self-selected pregnant women from the general population presenting for aneuploidy screening (without prior risk of fetal aneuploidy).
 Inclusion criteria: pregnant women between 10 to 11 weeks' gestation with singleton pregnancy who underwent the combined test.
 Exclusion criteria: multifetal pregnancies.

Patient characteristics and setting Number enrolled: 2905 pregnant women.
 Number available for 2 x 2 table: 2785 pregnant women (subgroup of 96%).
 Setting: 1 centre. Fetal Medicine Centre in London, UK.
 Recruitment period: October 2012 to January 2014.
 Ethnicity: Caucasian (88.5%), South Asian (6.0%), East Asian (3.3%), Afro-Caribbean (0.7%) and mixed (1.5%).
 Median gestational age (range): 10.6 (10 to 11.9) weeks.
 Median maternal age (range): 36.9 (20.4 to 51.9) years.
 Relevant tests carried out prior to index test: none.
 Language of the study: English.

Index tests gNIPT by TMPS (DANSR assay).
 Median fetal fraction DNA (range): 11% (4% to 40%).
 Blood samples for gNIPT were collected before reference standard.
 Cutpoint: not reported. Usually, Harmony™ prenatal test uses FORTE algorithm; positive if FORTE risk score ≥ 1%.
 Commercial test: Harmony™ Prenatal test by Ariosa Diagnostics, Inc.
 The traditional screening tests (combined test at the first trimester) was also assessed.
 Cutpoint of combined test: 1 in 100 for T21.

Target condition and reference standard(s) Target conditions: T21, T18 and T13.
 Reference standards: fetal karyotype of chorionic villi, amniotic fluid or products of conception, neonatal karyotype, neonatal clinical examination or medical record from birth.

Flow and timing Blood samples were obtained prior to invasive procedure (reference standard).
 gNIPT was a first-tier test.
 122/2905 failed the initial TMPS testing (122 = 123 - 1 sample lost in mail).

Genomics-based non-invasive prenatal testing for detection of fetal chromosomal aneuploidy in pregnant women (Review)

168

Quezada 2015 (Continued)

66/2851 samples without follow-up were excluded.

110/122 required repeat testing using a second blood sample and results were obtained in 69/110 samples.

53/2905 samples failed during sequencing process (41 samples failed second sequencing and 12 unrepeated tests) (no gNIPT result).

Comparative

Aim to study To examine, in a general population (pregnant women without prior risk of fetal aneuploidy), the performance of ccfDNA testing for T21, T18 and T13 at 10 to 11 weeks' gestation and compare it to that of the combined test at 11 to 13 weeks' gestation.

Funding source or sponsor of the study Study not funded by industry but Ariosa Diagnostics, Inc made sequencing and analyses.

Informations about the authors contacted Author was contacted on: 21 April 2016 and 30 May 2016.
No reply received from the author.

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	No		
--	----	--	--

Was a case-control design avoided?	Yes		
------------------------------------	-----	--	--

Did the study avoid inappropriate exclusions?	No		
---	----	--	--

High

Low

DOMAIN 2: Index Test TMPS

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
---	-----	--	--

If a threshold was used, was it pre-specified?	Yes		
--	-----	--	--

Low

Low

DOMAIN 2: Index Test Traditional screening tests

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
---	-----	--	--

If a threshold was used, was it pre-specified?	Yes		
--	-----	--	--

Quezada 2015 (Continued)

	Low	Low
DOMAIN 3: Reference Standard		
Is the reference standards likely to correctly classify the target condition?	Yes	
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear	
	Unclear	Low
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Yes	
Did all analysed patients receive the reference standard?	Yes	
Were all patients included in the analysis?	No	
	High	

Samango-Sprouse 2013

Study characteristics	
Patient sampling	Study design: blinded, prospective cohort study. Participants: pregnant women at high or low risk of fetal aneuploidy (known sex chromosome aneuploidy and euploid pregnancies). Inclusion criteria: women were at least 18 years of age, had singleton pregnancy, or with known sex chromosome aneuploidy. Exclusion criteria: pregnant women with known mosaicism, autosomal trisomy, or triploidy.
Patient characteristics and setting	Number enrolled: 201 pregnant women. Number available for 2 x 2 table: 186 pregnant women (subgroup of 93%). Setting: 8 prenatal care centres in UK, USA, Poland, and Czech Republic. Recruitment period: not reported. Ethnicity: not reported. Mean gestational age: euploid pregnancies 13.2 weeks, and aneuploid pregnancies 15.3 weeks. Gestational age range: overall 9.4 to 36.4 weeks. Maternal age: not reported. Relevant test carried out prior to index test: not reported. Language of the study: English.
Index tests	gNIPT by TMPS (SNP-based method) on Illumina HiSeq 2000 sequencer with NATUS algorithm. Mean fetal fraction DNA: euploids: 10.9% and aneuploids: 12.1%. Overall range: 2.9% to 37.7%. Blood samples for gNIPT were collected just before or at least 4 days after reference standard. Cutpoint: not reported.

Samango-Sprouse 2013 (Continued)

Commercial test: Natera's prenatal test.

Target condition and reference standard(s)	Target conditions: 45,X, 47,XXX, 47,XXY, and 47,XYY. Reference standards: fetal karyotype of chorionic villi or amniotic fluid or genetic testing of cord blood, buccal sample, saliva, or products of conception.
Flow and timing	Blood samples were collected just before or at least 4 days after invasive procedure (reference standard). gNIPT was a first- or second-tier test. 14/201 samples failed sequencing process quality control (no gNIPT result) including 12 for low fetal fraction or poor DNA quality and 2 samples did not return a result for SCA. 1/187 sample excluded for conflicting algorithm metrics (no meaningful gNIPT result). No repeated test reported.
Comparative	
Aim to study	To develop a SNP-based and informatics-based gNIPT that detects sex chromosome aneuploidies early in pregnancy.
Funding source or sponsor of the study	It is unclear if the study was funded by industry but all authors are employees of Natera, Inc. except the first author (Carole Samango-Sprouse). This study was supported in part by a grant from the National Institute of Health, National Institute of Child Health and Human Development.
Informations about the authors contacted	Author was contacted on: 22 April, 4 July and 29 September 2016. Replies received on: 29 and 30 September 2016.

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	High
DOMAIN 2: Index Test TMPS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		

Samango-Sprouse 2013 (Continued)

	Low	Low
DOMAIN 3: Reference Standard		
Is the reference standards likely to correctly classify the target condition?	Yes	
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes	
	Low	Low
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Yes	
Did all analysed patients receive the reference standard?	Yes	
Were all patients included in the analysis?	No	
	High	

Sehnert 2011

Study characteristics	
Patient sampling	Study design: blinded retrospective study (archived maternal plasma samples). Participants: pregnant women selected from a high risk of fetal aneuploidy population. Inclusion criteria: pregnant women age 18 years or older with singleton or multifetal pregnancy. Exclusion criteria: not reported.
Patient characteristics and setting	Number enrolled: overall: 1014 pregnant women including 71 women selected on 435 for the training set (not shown in the present review) and 48 women selected on 575 for the test set. Number available for 2 x 2 table: 47 (subgroup of 8%). Setting: 13 centres in USA. Recruitment period: January 2010 to June 2010. Ethnicity: Caucasian (62.7%), Hispanic (16.5%), Asian (6.2%), multiethnic (5.2%), Afro-American (4.0%), Native American (0.9 %) and other or not specified (1.8%). Mean gestational age (range): 15.4 (10.6 to 28.4) weeks. Mean maternal age (\pm SD; range): 34.2 (\pm 8.22; 18 to 46) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both. Language of the study: English.
Index tests	gNIPT by MPSS on Illumina Genome Analyzer Iix sequencer in uniplex. Fetal fraction DNA: not reported. Blood samples for gNIPT were collected before reference standard. Cutpoint prespecified with the training set:

Sehnert 2011 (Continued)

- 1) positive if NCV > 4 for autosomes. There is a "no call zone" between 2.5 and 4 considering as gNIPT positive result for the present review.
- 2) positive if NCV for Chrom. Y < -2.0 SDs from the mean of male samples and if NCV for Chrom. X < -3.0 SDs from the mean of female samples for 45,X.
 Commercial test: Verinata's prenatal test.

Target condition and reference standard(s)	Target conditions: T21, T18, T13 and 45,X. Reference standard: fetal karyotype of chorionic villi (58,3%) or amniotic fluid (41.7%).
Flow and timing	Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard). gNIPT was a second-tier test. 895/1014 samples were not selected for sequencing. 71/119 samples were selected for the training set (not shown in the present review). 1/48 sample from twin gestation in the test set was removed from the final analysis. No repeated test reported.
Comparative	
Aim to study	To develop and test an optimised algorithm from MPSS data and demonstrated the potential universality of the sequence tag mapping and chromosome quantification method for the detection of multiple chromosomal abnormalities.
Funding source or sponsor of the study	Study funded by Illumina (formerly Verinata Health). The funding organizations played a direct role in the design of the study, the choice of enrolled patients, the review and interpretation of data, and the preparation and final approval of the manuscript.
Informations about the authors contacted	No need for further contact.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		

Sehnert 2011 (Continued)

If a threshold was used, was it pre-specified? Yes

Low

Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? Yes

Low

Shaw 2014
Study characteristics

Patient sampling Study design: prospective cohort study.
 Participants: consecutive pregnant women were selected from a mixed-risk population. They were classified in extremely high-risk group for T21 with a screening T21 risk > 1:30 or nuchal translucency > 3.0 mm and low-risk group with a screening T21 risk < 1:1500.
 Inclusion criteria: pregnant women at > 12 weeks' gestation, singleton or multifetal pregnancies.
 Exclusion criteria: not reported.

Patient characteristics and setting Number enrolled: 201 pregnant women.
 Number available for 2 x 2 table: 200 pregnant women (subgroup of 99.5%).
 Setting: 11 medical centres in Taiwan.
 Recruitment period: June to December 2012.
 Ethnicity: Asian.
 Mean gestational age (\pm SD): high-risk pregnant women 17.3 (\pm 2.1) weeks, and low-risk pregnant women 16.1 (\pm 3.0) weeks.
 Gestational age range: overall 12 to 20 weeks.
 Mean maternal age (\pm SD): high-risk pregnant women 35.1 (\pm 3.2) years, and low-risk pregnant women 34.6 (\pm 2.6) years.
 Chorionicity: all dichorionic (4/4).

Shaw 2014 (Continued)

Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening.
 Language of the study: English.

Index tests	<p>gNIPT by MPSS on Illumina v2 HiSeq 2000 sequencer in 12-plex.</p> <p>Fetal fraction DNA: not reported. Blood samples for gNIPT were collected before reference standard. Cutpoint:</p> <ol style="list-style-type: none"> 1) positive if Z score > 3 (T21, T18, and T13). 2) positive if Z score Chrom. X < -3 and Z score Chrom. Y < 3 (45,X). 3) positive if Z score Chrom. X < -3 and Z score Chrom. Y > 3 (47,XXY). <p>Commercial test: Berry Genomics' prenatal test.</p> <p>The traditional screening test (combined test at the first trimester) was also assessed but complete data for 2 x 2 tables were unavailable.</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18, T13, 45,X, and 47,XXY. 47,XXX and 47,XXY were also screened but no case was found. Reference standards: fetal karyotype of amniotic fluid or medical record from birth.</p>
Flow and timing	<p>Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard). gNIPT was a second-tier test. 1 sample excluded for early gestational age (< 12 weeks).</p> <p>No repeated test reported.</p>
Comparative	
Aim to study	To evaluate the performance of gNIPT for all fetal chromosomal aneuploidies in an extremely high-risk group undergoing first-trimester combined T21 screening.
Funding source or sponsor of the study	Funding sources were not reported but 2 authors are affiliated to Berry Genomics Co. Ltd., Beijing, PR China.
Informations about the authors contacted	Author was contacted on: 10 February and 23 June 2016. No reply received from the author.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Unclear		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	High

Shaw 2014 (Continued)

DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low

Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? Yes

Low

Song 2013
Study characteristics

Patient sampling Study design: blinded, prospective cohort study.
 Participants: pregnant women without a priori risk of fetal aneuploidy who undergo routine prenatal screening.
 Inclusion criteria: singleton pregnancies and pregnant women younger than 35 years old.
 Exclusion criteria: not reported.

Patient characteristics and setting Number enrolled: 1916 pregnant women.
 Number available for 2 x 2 table: 1741 pregnant women (subgroup of 91%).
 Setting: 2 clinical centres in Beijing, China.
 Recruitment period: April 2011 to December 2011.
 Ethnicity: Asian.
 Mean gestational age (\pm SD; range): 16.57 (\pm 1.56; 11 to 21.9) weeks.
 Mean maternal age (\pm SD; range): 29.03 (\pm 2.70; 20 to 34) years.

Song 2013 (Continued)

Relevant tests carried out prior to index test: none.
 Language of the study: English.

Index tests	<p>gNIPT by MPSS on Illumina v2 HiSeq 2000 sequencer in 12-plex.</p> <p>Fetal fraction DNA: not reported. Blood samples for gNIPT were collected before reference standard. Cutpoint: positive if Z score ≥ 3. Commercial test: Berry Genomics' prenatal test.</p> <p>The traditional screening test (second-trimester triple test) was also assessed.</p> <p>Cutpoint of triple test: 1 in 270 for T21 and T18.</p>		
Target condition and reference standard(s)	<p>Target conditions: T21, T18 and T13. 45,X, 47,XXX, 47, XXY, 47,YYY were also screened but inappropriate reference standard for the present review was used. Reference standards: fetal karyotype of chorionic villi, amniotic fluid or cord blood or medical record from birth.</p>		
Flow and timing	<p>It is not reported if the blood samples were collected before or after invasive procedure (reference standard). It is not reported if the gNIPT was a first- or second-tier test. 64/1916 samples failed sequencing process (failed DNA quality control criteria or sequencing quality control) (no gNIPT result).</p> <p>102/1916 samples without follow-up were excluded.</p> <p>9/1916 samples were without follow-up and failed sequencing process (no gNIPT result).</p> <p>No repeated test reported.</p>		
Comparative			
Aim to study	<p>To evaluate the performance of gNIPT for detection of fetal aneuploidies in a Chinese cohort of women younger than 35 years old in a prospective clinical setting. Also, to compare the performance of gNIPT with the routine prenatal screening (second-trimester combined test).</p>		
Funding source or sponsor of the study	<p>Study not funded by industry. This study was supported by a grant (2006BAI05A10) from the National Key Technology Research and Development Program of China during the '11th Five-Year Plan'.</p>		
Informations about the authors contacted	<p>No need for further contact.</p>		
Notes	<p>SCA were also screened but inappropriate reference standard for the present review was used. gNIPT data from SCA were not shown in this review.</p>		
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Unclear		
Was a case-control design avoided?	Yes		

Song 2013 (Continued)

Did the study avoid inappropriate exclusions? No

High
Low
DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low
Low
DOMAIN 2: Index Test Traditional screening tests

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Unclear

Did all analysed patients receive the reference standard? No

Were all patients included in the analysis? No

High

Song 2015
Study characteristics

Patient sampling	<p>Study design: blinded, prospective cohort study.</p> <p>Participants: pregnant women selected arbitrarily at high risk of fetal aneuploid presenting for aneuploidy screening by gNIPT.</p> <p>Inclusion criteria: advanced maternal age (≥ 35 years) and singleton pregnancies.</p> <p>Exclusion criteria: multifetal pregnancies.</p>
Patient characteristics and setting	<p>Number enrolled: 213 pregnant women.</p> <p>Number available for 2 x 2 table: 204 pregnant women (subgroup of 96%).</p> <p>Setting: 1 centre. Peking Union Medical College Hospital (PUMCH), Beijing, China.</p> <p>Recruitment period: May 2012 to August 2013.</p> <p>Ethnicity: Asian.</p> <p>Median gestational age (range): 9.9 (8 to 12.9) weeks.</p> <p>Mean maternal age (range): 37.25 (35 to 45) years.</p> <p>Relevant tests carried out prior to index test: none.</p> <p>Language of the study: English.</p>
Index tests	<p>gNIPT by MPSS on Illumina v2 HiSeq 2000 sequencer in 12-plex.</p> <p>Median fetal fraction DNA (range): only male fetus: 8.54% (2.69% to 18.75%).</p> <p>Blood samples for gNIPT were collected before reference standard.</p> <p>Cutpoint: positive if Z score ≥ 3.</p> <p>Commercial test: Berry Genomics' prenatal test.</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18, T13, 45,X, 47,XXY and 47,XXX. 47,XYY were also assessed but no case was found.</p> <p>Reference standards: fetal karyotype of chorionic villi or amniotic fluid for all analysed women (178/178) and neonatal clinical examination (198/212).</p>
Flow and timing	<p>Blood samples for gNIPT were obtained prior the invasive procedure (reference standard). gNIPT was a first-tier test.</p> <p>1/213 sample failed quality control (haemolysis).</p> <p>8/212 samples without reference standard were excluded including 5 miscarriages, 2 in-traiterine fetal deaths and 1 termination of pregnancy.</p> <p>No repeated test reported.</p>
Comparative	
Aim to study	To evaluate the feasibility of gNIPT of maternal plasma samples collected from pregnant Chinese women in early gestation, between 8 and 12.9 weeks' gestation.
Funding source or sponsor of the study	<p>Study not funded by industry but gNIPTs were done and analysed at Berry Genomics Co. Ltd.</p> <p>Study funded by a grant from the National Natural Science Foundation of China.</p>
Informations about the authors contacted	No need for further contact.

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			

Song 2015 *(Continued)*

Was a consecutive or random sample of patients enrolled? Yes

Was a case-control design avoided? Yes

Did the study avoid inappropriate exclusions? No

High

Low

DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low

Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Sparks 2012a
Study characteristics

Patient sampling Study design: case-control study from a prospective cohort.
 Participants: pregnant women selected from a high risk of fetal aneuploidy population.
 Inclusion criteria: women at least 18 years of age, at least 10 weeks' gestation and have singleton pregnancy.

Sparks 2012a (Continued)

Exclusion criteria: multifetal pregnancies.

Patient characteristics and setting	<p>Number enrolled: not reported. A subset of 338 pregnant women including 171 women in the training set (data not shown in the present review) and 167 women in the validation set were selected for this study.</p> <p>Number available for 2 x 2 table: 167 pregnant women (subgroup of 49%).</p> <p>Setting: not reported.</p> <p>Recruitment period: not reported.</p> <p>Ethnicity: not reported.</p> <p>Mean gestational age (\pm SD; range): 18.6 (\pm 4.0; 11 to 36.1) weeks.</p> <p>Mean maternal age (\pm SD; range): 33.5 (\pm 7.1; 18 to 51) years.</p> <p>Relevant tests carried out prior to index test: not reported.</p> <p>Language of the study: English.</p>
Index tests	<p>gNIPT by TMPS (DANSR assay) on Illumina HiSeq 2000 sequencer in multiplex with FORTE algorithm.</p> <p>Range fetal fraction DNA: 3% to 33%.</p> <p>It is not reported if the blood samples for gNIPT were collected before or after reference standard.</p> <p>Cutpoint: not reported.</p> <p>Commercial test: Ariosa Diagnostics, Inc's prenatal test.</p>
Target condition and reference standard(s)	<p>Target conditions: T21 and T18.</p> <p>Reference standards: fetal karyotype or chromosome analysis by FISH or both.</p>
Flow and timing	<p>It is not reported if the blood samples were collected before or after invasive procedure (reference standard).</p> <p>gNIPT was a second-tier test.</p> <p>171/338 samples were excluded for the training set.</p> <p>No failed sample reported in the validation set.</p> <p>No repeated test reported.</p>
Comparative	
Aim to study	To develop a novel biochemical assay and algorithm for the prenatal evaluation of risk for fetal T21 and T18 using ccfDNA obtained from maternal blood.
Funding source or sponsor of the study	Study funded by Ariosa Diagnostics, Inc. All authors are employees of Aria Dx Inc. (now Ariosa Diagnostics). K Sparks is a member of the board of the company.
Informations about the authors contacted	Author was contacted on: 23 June 2016. No reply received from the author.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	No		

Sparks 2012a (Continued)

Did the study avoid inappropriate exclusions? No

High
Low
DOMAIN 2: Index Test TMPS

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Unclear

Unclear
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Yes

Low
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Unclear

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? Yes

Unclear
Stumm 2014
Study characteristics

Patient sampling Study design: prospective cohort study. Blinded for T21 and unblinded for T18 and T13. Participants: all consecutively enrolled pregnant women selected at high risk of fetal aneuploidy. Inclusion criteria: pregnant women at least 18 years old, at high risk for chromosomal aberrations, signed informed consent, planned a conventional karyotyping procedure (invasive diagnostic), had singleton pregnancy and blood drawn before the invasive procedure. Exclusion criteria: multifetal pregnancies.

Patient characteristics and setting Number enrolled: 522 pregnant women. Number available for 2 x 2 table: 472 pregnant women (subgroup of 90%).

Stumm 2014 (Continued)

Setting: 5 clinical centres in Germany and Switzerland.
 Recruitment period: not reported.
 Ethnicity: not reported.
 Mean gestational age (range): 15.6 (11.0 to 32.1) weeks.
 Mean maternal age (range): 36.0 (19 to 47) years.
 Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both.
 Language of the study: English.

Index tests	<p>gNIPT by MPSS on Illumina HiSeq 2000 sequencer in 12-plex with DAP.21 algorithm without CG correction.</p> <p>Mean fetal fraction DNA (range): male fetus only: 12.3% (3.7% to 36.8%). Blood samples for gNIPT were collected just before reference standard. Cutpoint:</p> <p>1) positive if MAD-based Z -score ≥ 3 for T21. 2) positive if MAD-based Z score ≥ 3.2 for T18. 3) positive if MAD-based Z score ≥ 3.9 for T13. Commercial test: LifeCodexx's prenatal test.</p>		
Target condition and reference standard(s)	<p>Target conditions: T21, T18 and T13. Reference standard: fetal karyotype of chorionic villi (30.3%), amniotic fluid (69.1%) or cord blood (0.6%).</p>		
Flow and timing	<p>Blood samples for gNIPT were obtained just prior the invasive procedure (reference standard). gNIPT was a second-tier test. 18/522 samples excluded, including 8 without reference standard result, 9 without consent and 1 was previously analysed.</p> <p>32/504 samples failed during sequencing process (no gNIPT result), including 14 samples failed sequencing quality criteria and 18 samples failed libraries.</p> <p>No repeated test reported.</p>		
Comparative			
Aim to study	To validate the diagnostic accuracy of a gNIPT for detecting T21, T18 and T13 for a population in Germany and Switzerland.		
Funding source or sponsor of the study	Study funded by LifeCodexx AG and GATC Biotech AG.		
Informations about the authors contacted	Author was contacted on: 22 February 2016, 24 February and 19 May 2016. Reply received on: 24 February 2016.		
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		

Stumm 2014 (Continued)

Was a case-control design avoided?	Yes
------------------------------------	-----

Did the study avoid inappropriate exclusions?	No
---	----

High

Low

DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard?	No
---	----

If a threshold was used, was it pre-specified?	No
--	----

High

Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes
---	-----

Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes
--	-----

Low

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes
--	-----

Did all analysed patients receive the reference standard?	Yes
---	-----

Were all patients included in the analysis?	No
---	----

High

Sukhikh 2015
Study characteristics

Patient sampling	Study design: prospective cohort study. Participants: pregnant women selected at high risk of fetal aneuploidy presenting for invasive testing. Inclusion criteria: not reported. Exclusion criteria: not reported.
------------------	--

Sukhikh 2015 (Continued)

Patient characteristics and setting	Number enrolled: 200 pregnant women. Number available for 2 x 2 table: 200 pregnant women (whole cohort included in analyses). Setting: not reported. Recruitment period: not reported. Ethnicity: not reported. Median gestational age (range): 14 (10 to 20) weeks. Maternal age: not reported. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both. Language of the study: Russian.
Index tests	gNIPT by MPSS on Ion Proton™ sequencer. Fetal fraction DNA: not reported. Blood samples for gNIPT were collected before reference standard. Cutpoint: 1) Positive for T21 and T18 if T score > 5. 2) Positive for T13 if T score > 4. 3) Positive for 45,X if T score for chrom. X > 0.04 and for chrom. Y < 0.04. In-house test.
Target condition and reference standard(s)	Target conditions: T21, T18, T13 and 45,X. Reference standard: fetal karyotype of chorionic villi, amniotic fluid or placenta.
Flow and timing	Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard). gNIPT was a second-tier test. No failed sample reported. No repeated test reported.
Comparative	
Aim to study	To estimate the feasibility of using a next-generation sequencing technique for the non-invasive prenatal diagnosis of fetal aneuploidies.
Funding source or sponsor of the study	Funding source not reported.
Informations about the authors contacted	Author was contacted on: 9 September and 4 October 2016. No reply received from the author.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		

Sukhikh 2015 (Continued)

Did the study avoid inappropriate exclusions? No

High
Low
DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard? Unclear

If a threshold was used, was it pre-specified? Yes

Unclear
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? Unclear

Unclear
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? Yes

Low
Sung-Hee 2015
Study characteristics

Patient sampling Study design: retrospective study from a prospective cohort.
 Participants: pregnant women selected from a high risk of fetal aneuploidy population.
 Inclusion criteria: singleton pregnancies.
 Exclusion criteria: multifetal pregnancies.

Patient characteristics and setting Number enrolled: 918 pregnant women.
 Number available for 2 x 2 table: 901 pregnant women (subgroup of 99%).
 Setting: various medical sites in Korea.
 Recruitment period: May 2012 to December 2013.
 Ethnicity: Asian.
 Mean gestational age (\pm SD; range): 16.6 (\pm 2.2; 11 to 25) weeks.
 Mean maternal age (\pm SD; range): 35.3 (\pm 4.1; 22 to 46) years.

Sung-Hee 2015 (Continued)

Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening (59%) or both.
 Language of the study: English.

Index tests	<p>gNIPT by MPSS on Illumina Genome Analyzer IIx or HiSeq 2000 sequencer in 12-plex.</p> <p>Fetal fraction DNA: amount measured but not reported. Blood samples for gNIPT were collected before reference standard. Cutpoint: positive if L score > 1 and t score > 2.5 (warning zone if t score risk > 2.5 or L score risk > 1). Commercial test: BGI-Shenzhen's test.</p> <p>Biochemical serum-screening results were reported in the study but 2 x 2 tables could not be derived.</p>
Target condition and reference standard(s)	<p>Target conditions: T21 and T18. T13 were also assessed but the only case found was without follow-up. SCA were also screened but inappropriate reference standard for the present review was used. Reference standards: fetal karyotype for gNIPT positive cases and medical record from birth for gNIPT negative cases.</p>
Flow and timing	<p>Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard). gNIPT was a first- or a second-tier test. 8/918 samples were ineligible.</p> <p>9/910 samples without follow-up were excluded (all samples had positive gNIPT result). 2/9 women had abortion and 7/9 women declined invasive testing.</p> <p>21/910 samples failed to give a risk score for gNIPT of the first blood samples including 1 haemolysed sample, 8 samples thawing due to transport delay, 3 due to cell-free DNA extraction failures and 9 samples had low fetal fraction.</p> <p>16/21 samples were repeated with new sampling. 14/16 samples obtained a gNIPT results and 2/16 samples failed to provide informative results and were classified as test failures.</p> <p>7/910 samples failed during sequencing process (no gNIPT result). 5/7 samples failed the initial MPSS testing and were not resequenced and 2/7 samples failed the second MPSS testing.</p>
Comparative	
Aim to study	<p>To report the initial clinical performance of gNIPT in detecting fetal chromosomal aneuploidies, especially T21, T18 and T13, in singleton pregnancies in Korea.</p>
Funding source or sponsor of the study	<p>Study not funded by industry but BGI performed sequencing and analysis. Study funded by Seoul Clinical Laboratories Research Grant (2015, President: Kyoung-Ryul Lee).</p>
Informations about the authors contacted	<p>Author was contacted on: 13, 19 and 26 September 2016. Replies received on: 19 and 25 September and 11 October 2016.</p>

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		

Sung-Hee 2015 *(Continued)*

Did the study avoid inappropriate exclusions? No

High

Low

DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard? Yes

If a threshold was used, was it pre-specified? Yes

Low

Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition? Yes

Were the reference standard results interpreted without knowledge of the results of the index tests? No

High

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all analysed patients receive the reference standard? Yes

Were all patients included in the analysis? No

High

Tynan 2016
Study characteristics

Patient sampling Study design: blinded, retrospective clinical evaluation study.
 Participants: pregnant women selected from 3 internal clinical studies (archived maternal plasma samples). 84.5% without prior risk and 15.8% had high risk of fetal aneuploidy.
 Inclusion criteria: singleton pregnancies.
 Exclusion criteria: multifetal pregnancies.

Tynan 2016 (Continued)

Patient characteristics and setting	<p>Number enrolled: 1100 pregnant women. Number available for 2 x 2 table: 1048 pregnant women (subgroup of 95%). Setting: multicentre. Recruitment period: beginning in November 2009. Ethnicity: not reported. Gestational age (range): 9 to 38.1 weeks. Maternal age (range): 18 to 45 years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both. Language of the study: English.</p>		
Index tests	<p>gNIPT by MPSS on Illumina HiSeq 2000 or HiSeq 2500 sequencer in multiplex.</p> <p>Mean fetal fraction DNA (\pm SD): low-risk group: 10% (\pm 3.1%), high-risk group (< 35 years): 11.9% (\pm 4.8%), and high-risk group (\geq 35 years): 11.1% (\pm 3.4%).</p> <p>Median fetal fraction DNA (range): low-risk group: 10.7% (3.1% to 22.9%), high-risk group (< 35 years): 10.7% (4.9% to 28.3%), and high-risk group (\geq 35 years): 11% (3.1% to 25.5%). Blood samples for gNIPT were collected before reference standard. Cutpoint: positive if risk score \geq 1%. Commercial test: VisibiliT™ test by Sequenom, Inc.</p>		
Target condition and reference standard(s)	<p>Target conditions: T21, T18 and T13. Reference standards: fetal karyotype of chorionic villi or amniotic fluid or medical record from birth.</p>		
Flow and timing	<p>Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard). gNIPT was a first- or second-tier test. 52/1100 samples failed during sequencing process (no gNIPT result) including 28 for technical failures (library preparation or low aligned reads counts) and 24 for discretionary non-reporting because of factors such as sequencing bias.</p> <p>No repeated test reported.</p>		
Comparative			
Aim to study	<p>To demonstrate the clinical performance of a simplified, low coverage, low cost MPSS assay (VisibiliT™) that combines a maternal age-based risk for T21, T18, and T13, the fractional concentration of fetal DNA, and the representation of chromosomes 21, 18, and 13 in the sample to provide a risk score for T21, T18 and T13, with classification of fetal sex result.</p>		
Funding source or sponsor of the study	<p>Study funded by Sequenom, Inc.</p>		
Informations about the authors contacted	<p>No need for further contact.</p>		
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		

Tynan 2016 *(Continued)*

Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	High
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all analysed patients receive the reference standard?	Yes		
Were all patients included in the analysis?	No		
		High	

Verweij 2013
Study characteristics

Patient sampling	Study design: blinded, prospective cohort study. Participants: consecutive pregnant women selected at high risk of fetal aneuploidy presenting for invasive testing. Inclusion criteria: women who sign informed consent, ≥ 18 years old and carrying a singleton pregnancy with a gestational age of at least 10 weeks.
------------------	--

Verweij 2013 (Continued)

Exclusion criteria: multifetal pregnancies, an invasive procedure performed prior to the blood sampling, history or active significant malignancy requiring major surgery or systemic chemotherapy, or language restriction with failure to understand the study information.

Patient characteristics and setting	<p>Number enrolled: 595 pregnant women. Number available for 2 x 2 table: 504 pregnant women (subgroup of 85%). Setting: multicentres in the Netherlands and Sweden. Recruitment period: May 2011 to March 2012. Ethnicity: Caucasian (84.8%), Mediterranean (6%), Asian (3.3%), Afro-European (1.3%), and other (4.6%). Mean gestational age (\pm SD; range): 14.0 (\pm 2.1; 10 to 28) weeks. Mean maternal age (\pm SD; range): 36.4 (\pm 4.6; 20 to 47) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both. Language of the study: English.</p>
Index tests	<p>gNIPT by TMPS (DANSR assay) on Illumina HiSeq 2000 in 96-plex with FORTE algorithm.</p> <p>Mean fetal fraction DNA (\pm SD; range): 11.1% (\pm 4.1%; 4% to 30%). Blood samples for gNIPT were collected just before reference standard. Cutpoint: positive if FORTE risk score \geq 1%. Commercial test: Ariosa Diagnostics, Inc's test.</p>
Target condition and reference standard(s)	<p>Target condition: T21. Reference standard: fetal karyotype of chorionic villi (54%) or amniotic fluid (46%).</p>
Flow and timing	<p>Blood samples for gNIPT were obtained just prior the invasive procedure (reference standard). gNIPT was a second-tier test. 75/595 samples were ineligible.</p> <p>51/520 samples failed the initial TMPS testing.</p> <p>51/51 samples were repeated with a second aliquot of the first sampling and 35/51 samples obtained a gNIPT results.</p> <p>16/520 samples failed during sequencing process (no gNIPT result), including 7 samples with low fetal DNA fraction and 9 samples failed laboratory processing or specimen issues.</p>
Comparative	
Aim to study	<p>To evaluate the performance of a directed gNIPT method of ccfDNA analysis for fetal T21 by shipping the whole blood samples from Europe to a laboratory in the USA.</p>
Funding source or sponsor of the study	<p>Study funded by Ariosa Diagnostics, Inc. 2 authors are paid employees of Ariosa Diagnostics. 1 author is a board member of Ariosa Diagnostics.</p>
Informations about the authors contacted	<p>Author was contacted on: 22 April 2016. Reply received on: 25 April 2016.</p>
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		

Verweij 2013 *(Continued)*

Was a case-control design avoided?	Yes
------------------------------------	-----

Did the study avoid inappropriate exclusions?	No
---	----

High

Low

DOMAIN 2: Index Test TMPS

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes
---	-----

If a threshold was used, was it pre-specified?	Yes
--	-----

Low

Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes
---	-----

Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes
--	-----

Low

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes
--	-----

Did all analysed patients receive the reference standard?	Yes
---	-----

Were all patients included in the analysis?	No
---	----

High

Wang 2014
Study characteristics

Patient sampling	Study design: prospective cohort study. Participants: pregnant women in the first trimester of pregnancy with advanced maternal ages or ultrasound abnormality (high risk of fetal aneuploidy). Inclusion criteria: singleton pregnancies between 11 to 14 weeks' gestation.
------------------	--

Wang 2014 (Continued)

Exclusion criteria: multifetal pregnancies.

Patient characteristics and setting	Number enrolled: 136 pregnant women. Number available for 2 x 2 table: 136 pregnant women (whole cohort included in analyses). Setting: 1 centre. General Hospital of PLA, Beijing, China. Recruitment period: March 2011 to August 2013. Ethnicity: Asian. Gestational age range: 11 to 13.9 weeks. Maternal age range: 35 to 44 years. Relevant test carried out prior to index test: ultrasonography for some women. Language of the study: Chinese.
Index tests	gNIPT by MPSS on Illumina HiSeq 2000 sequencer with NIFTY™ algorithm. Fetal fraction DNA: not reported. Blood samples for gNIPT were collected before reference standard. Cutpoint: not reported. Commercial test: BGI-Shenzhen's prenatal test.
Target condition and reference standard(s)	Target conditions: T21 and T18. T13 was also assessed but no case was found. 45,X was also screened but inappropriate reference standard for the present review was used for pregnant women with gNIPT negative result. gNIPT data from 45,X were not shown in this review. Reference standards: fetal karyotype of amniotic fluid or cord blood or neonatal clinical examination at 42 days after birth or both.
Flow and timing	Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard). gNIPT was a first- or second-tier test. No failed sample reported. No repeated test reported.
Comparative	
Aim to study	To investigate the value of maternal plasma ccfDNA examination in detection of fetal chromosomal aneuploidy in pregnant women at advanced maternal age during the first trimester of pregnancy.
Funding source or sponsor of the study	Study not funded by industry. Study funded by National Science & Technology Pillar Program during the Twelfth Five-year Plan Period (2012BA131B06).
Informations about the authors contacted	Author was contacted on: 19 May and 27 June 2016. No replies received from the author.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Unclear		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		

Wang 2014 (Continued)

		High	Low
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Unclear		
Did all analysed patients receive the reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Unclear	

Wang 2015a
Study characteristics

Patient sampling	Study design: prospective cohort study. Participants: pregnant women selected from a high risk of fetal aneuploidy population. Inclusion criteria: pregnant women at high risk of fetal aneuploidy between 14 and 26 weeks of gestation. Exclusion criteria: not reported.
Patient characteristics and setting	Number enrolled: 917 pregnant women. Number available for 2 x 2 table: 917 pregnant women (whole cohort included in analyses). Setting: 1 centre at prenatal clinic, Lianyungang Maternal and Child Health Hospital, Lianyungang, Jiangsu 222001, China. Recruitment period: January 2012 to December 2013. Ethnicity: Asian. Gestational age range: 14 to 26 weeks.

Wang 2015a (Continued)

Maternal age range: 18 to 46 years.
Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both.
Language of the study: English.

Index tests	gNIPT by MPSS on Illumina v2 HiSeq 2000 flow cell on a HiSeq sequencer. Fetal fraction DNA: not reported. Blood samples for gNIPT were collected before reference standard. Cutpoint: 1) for T21, T18 and T13, positive if Z score > 3. 2) for 47,XXY and 47,XYY, positive if Z score Chrom. X > -3 and Z score Chrom. Y < 3. 3) for 45,X and 47,XXX, positive if Z score Chrom. X between -3 and 3 without Chrom. Y representation. Commercial test: Berry Genomics' prenatal test.
Target condition and reference standard(s)	Target conditions: T21 and T18. T13 was also assessed but no case was found. SCA was also assessed but inappropriate reference standard for the present review was used. Reference standards: fetal karyotype of amniotic fluid or clinical follow-up (once per month) from birth to 6 months.
Flow and timing	Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard). gNIPT was a second-tier test. No failed sample reported. No repeated test reported.
Comparative	
Aim to study	To investigate the clinical efficiency of gNIPT identifying fetal chromosomal aneuploidies.
Funding source or sponsor of the study	Study not funded by industry but Berry Genomics Co. Ltd give technical support. Study funded by the Community Development Fund, granted by the Department of Family Planning and Healthcare, Jiangsu Province, China.
Informations about the authors contacted	No need for further contact.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		

Wang 2015a (Continued)

	High	Low
DOMAIN 2: Index Test MPSS		
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes	
If a threshold was used, was it pre-specified?	Yes	
	Low	Low
DOMAIN 3: Reference Standard		
Is the reference standards likely to correctly classify the target condition?	Yes	
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear	
	Unclear	Low
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Yes	
Did all analysed patients receive the reference standard?	Yes	
Were all patients included in the analysis?	Yes	
	Low	

Yao 2014

Study characteristics	
Patient sampling	Study design: retrospective study. Participants: pregnant women presenting with low-, high- or without prior risk factors of fetal aneuploidy (gNIPT was offered routinely as a prenatal screening test). Inclusion criteria: singleton pregnancies. Exclusion criteria: multifetal pregnancies.
Patient characteristics and setting	Number enrolled: 5950 pregnant women. Number available for 2 x 2 table: 5530 pregnant women (subgroup of 93%). Setting: 1 centre. The Prenatal Diagnosis Centre, Southwest Hospital, Chongqing, China. Recruitment period: June 2011 to December 2012. Ethnicity: Asian. Mean gestational age (range): 19.6 weeks (65% of the cohort were between 16 to 20.9 weeks).

Yao 2014 (Continued)

Mean maternal age (\pm SD): 30 (\pm 5) years.
Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both for some women.
Language of the study: English.

Index tests

gNIPT by MPSS on Illumina Genome Analyzer IIx or HiSeq 2000 sequencer in 12-plex with NIFTY™ algorithm.

Fetal fraction DNA: amount measured but not reported.
Blood samples for gNIPT were collected before reference standard.
Cutpoint:

- 1) positive if t score \geq 2.5 for autosomes.
- 2) positive if t score for Chrom. X $<$ -2.5 for female fetuses for 45,X.
- 3) positive if t score for Chrom. X $>$ 2.5 for female fetuses for 47,XXX.
- 4) positive if t score for Chrom. X $>$ 2.5 combined with estimation of fetal cfDNA concentration by Chrom. X (expected value of zero) for 47,XXY.
- 5) positive if t score for Chrom. X $>$ 2.5 and R-value (the ratio of the fetal DNA fraction estimated by chromosome Y to that estimated by chromosome X) between 1.8 and 2.2 for 47,XXY.

Commercial test: BGI-Shenzhen's prenatal test.

Target condition and reference standard(s)

Target conditions: T21, T18, T13. 45,X, 47,XXY, 47,XYY and 47,XXX were also screened but inappropriate reference standard for the present review was used. gNIPT data from SCA were not shown in this review.
Reference standards: fetal karyotype of chorionic villi or amniotic fluid or follow-up by telephone interview with the clinician after the expected delivery date.

Flow and timing

Blood samples for gNIPT were obtained prior to the invasive procedure (reference standard).
gNIPT was a first- or second-tier test.
420/5950 samples without follow-up were excluded.

No failed sample reported.
No repeated test reported.

Comparative

Aim to study

To evaluate the performance of a MPSS in detecting fetal sex chromosome aneuploidy (SCA) and to present a comprehensive clinical counselling protocol for SCA-positive patients. Author also assessed autosomes aneuploidies.

Funding source or sponsor of the study

Funding source not reported but many authors are employees of the Clinical Laboratory of BGI Health, BGI-Shenzhen or of the Shenzhen Birth Defect Screening Project Lab.

Informations about the authors contacted

No need for further contact.

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

DOMAIN 1: Patient Selection

Yao 2014 (Continued)

Was a consecutive or random sample of patients enrolled?	No
--	----

Was a case-control design avoided?	Yes
------------------------------------	-----

Did the study avoid inappropriate exclusions?	No
---	----

High
High
DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes
---	-----

If a threshold was used, was it pre-specified?	Yes
--	-----

Low
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes
---	-----

Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear
--	---------

Unclear
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes
--	-----

Did all analysed patients receive the reference standard?	Yes
---	-----

Were all patients included in the analysis?	No
---	----

High
Zhang 2016
Study characteristics

Patient sampling	Study design: blinded, prospective cohort study.
------------------	--

Zhang 2016 (Continued)

Participants: pregnant women selected from a high risk of fetal aneuploidy population.
 Inclusion criteria: women aged ≥ 35 years at the time of delivery, single birth, high risk of T21 or single abnormal multiple of the median, elevated fetal nuchal translucency in the early pregnancy, a soft marker in the genetic scan, or cardiac structural abnormalities in the *second-trimester* genetic sonography, not suitable for invasive prenatal diagnosis, such as those with human immunodeficiency virus infection, placenta previa, *low-set* placenta, oligohydramnios, Rh-negative blood type, a history of abortion, threatened abortion or precious pregnancy.
 Exclusion criteria: multifetal pregnancies, maternal with chromosomal diseases, or received allogeneic blood transfusion, organ transplantation, stem cell therapy, or with a gestational age of < 12 weeks.

Patient characteristics and setting	Number enrolled: 87 pregnant women. Number available for 2 x 2 table: 87 pregnant women (whole cohort included in analyses). Setting: 1 centre at the Obstetrics and Gynecology Hospital of Fudan University (Shanghai, China). Recruitment period: January 2012 to December 2013. Ethnicity: Asian. Median gestational age (range): 19 (12.4 to 32.5) weeks. Mean maternal age (\pm SD): 37.48 (\pm 2.17) years. Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) or biochemical screening or both. Language of the study: English.
Index tests	gNIPT by MPSS on Illumina Hiseq 2000 sequencer in 12-plex. Fetal fraction DNA: not reported. It is not reported if the blood samples for gNIPT were collected before or after reference standard. Cutpoint for T21: positive if Z score ≥ 3 . No other cutpoint reported. Commercial test: Berry Genomics' prenatal test.
Target condition and reference standard(s)	Target conditions: T21, T18, 45,X and 47,XXX were also screened but inappropriate reference standard for the present review was used. Reference standards: fetal karyotype of amniotic fluid or cord blood or neonatal clinical examination by neonatologists.
Flow and timing	It is not reported if the blood samples were collected before or after invasive procedure (reference standard). gNIPT was a second-tier test. No failed sample reported. No repeated test reported.
Comparative	
Aim to study	To evaluate the efficacy of using gNIPT technology in screening T21 among women of advanced maternal age and to provide evidence for prenatal screening of T21.
Funding source or sponsor of the study	Funding source not reported.
Informations about the authors contacted	Author was contacted on: 7 September 2016. No reply received from the author.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

Zhang 2016 (Continued)

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	No
--	----

Was a case-control design avoided?	Yes
------------------------------------	-----

Did the study avoid inappropriate exclusions?	No
---	----

High
Low
DOMAIN 2: Index Test MPSS

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes
---	-----

If a threshold was used, was it pre-specified?	Yes
--	-----

Low
Low
DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes
---	-----

Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes
--	-----

Low
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Unclear
--	---------

Did all analysed patients receive the reference standard?	Yes
---	-----

Were all patients included in the analysis?	Yes
---	-----

Unclear
Zhou 2014a
Study characteristics

Zhou 2014a (Continued)

Patient sampling	<p>Study design: blinded, prospective cohort study.</p> <p>Participants: pregnant women selected at high risk for T21 (51.3%), low risk for T21 (2.6%) or without a priori risk (46.1%). gNIPT was integrated in clinical workflow.</p> <p>Inclusion criteria: singleton pregnancies.</p> <p>Exclusion criteria: multifetal pregnancies.</p>
Patient characteristics and setting	<p>Number enrolled: 306 pregnant women.</p> <p>Number available for 2 x 2 tables: 301 pregnant women in the pilot validation set (subgroup of 98%). See Zhou 2014b for the integration set.</p> <p>Setting: 1 centre. Women's Hospital, Zhejiang University School of Medicine, Hangzhouin, China.</p> <p>Recruitment period: September 2011 to October 2011.</p> <p>Ethnicity: Asian.</p> <p>Gestational age range: 12 to 24 weeks.</p> <p>Maternal age: not reported.</p> <p>Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening for a part of this cohort.</p> <p>Language of the study: English.</p>
Index tests	<p>gNIPT by MPSS on Illumina Genome Analyzer Iix or HiSeq 2000 sequencer in 12-plex.</p> <p>Fetal fraction DNA: amount measured but not reported.</p> <p>Blood samples for gNIPT were collected before reference standard.</p> <p>Cutpoint: positive if T score > 2.5 and L score > 1 (warning zone if t score > 2.5 or L score > 1).</p> <p>Commercial test: NIFTY™ prenatal test by BGI-Shenzhen.</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18 and T13.</p> <p>Reference standards: fetal karyotype of amniotic fluid or neonatal karyotype or birth outcome.</p>
Flow and timing	<p>Blood samples were obtained prior to the invasive procedure (reference standard).</p> <p>gNIPT was a first- or second-tier test.</p> <p>For the pilot validation set: 5/306 samples without follow-up were excluded.</p> <p>No failed sample reported.</p> <p>No repeated test reported.</p>
Comparative	
Aim to study	To report the clinical application of gNIPT to detect chromosomal aneuploidies, especially T21, T18 and T13 in Chinese singleton pregnancies.
Funding source or sponsor of the study	Study not funded by industry but BGI-Shenzhen made sequencing and analysis. Some authors are employees of BGI-Shenzhen.
Informations about the authors contacted	Author was contacted on: 31 May 2016. No reply received from author.
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			

Zhou 2014a *(Continued)*

Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	High
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all analysed patients receive the reference standard?	Yes		
Were all patients included in the analysis?	No		
		High	

Zhou 2014b
Study characteristics

Patient sampling	Study design: blinded, prospective cohort study. Participants: pregnant women selected at high risk, low risk for T21 or without a priori risk. gNIPT was integrated in clinical workflow. Inclusion criteria: singleton pregnancies.
------------------	--

Zhou 2014b (Continued)

Exclusion criteria: multifetal pregnancies.

Patient characteristics and setting	<p>Number enrolled: 7705 pregnant women.</p> <p>Number available for 2 x 2 tables: 3950 pregnant women in the integration set (subgroup of 51%).</p> <p>Setting: 1 centre. Women's Hospital, Zhejiang University School of Medicine, Hangzhouin, China.</p> <p>Recruitment period: September 2011 to July 2013.</p> <p>Ethnicity: Asian.</p> <p>Gestational age range: 12 to 24 weeks.</p> <p>Maternal age: not reported.</p> <p>Relevant tests carried out prior to index test: ultrasonography (nuchal translucency measurement) and biochemical screening for a part of this cohort.</p> <p>Language of the study: English.</p>
Index tests	<p>gNIPT by MPSS on Illumina Genome Analyzer Ix or HiSeq 2000 sequencer in 12-plex.</p> <p>Fetal fraction DNA: amount measured but not reported.</p> <p>Blood samples for gNIPT were collected before reference standard.</p> <p>Cutpoint: positive if T score > 2.5 and L score > 1 (warning zone if t score > 2.5 or L score > 1).</p> <p>Commercial test: NIFTY™ prenatal test by BGI-Shenzhen.</p>
Target condition and reference standard(s)	<p>Target conditions: T21, T18 and T13.</p> <p>Reference standards: fetal karyotype of amniotic fluid or neonatal karyotype or birth outcome.</p>
Flow and timing	<p>Blood samples were obtained prior to the invasive procedure (reference standard).</p> <p>gNIPT was a first- or second-tier test.</p> <p>141/7705 samples failed the initial MPSS testing. 141/141 samples were repeated with a new sampling and 137/141 samples obtained a gNIPT results.</p> <p>4/7705 samples failed the second MPSS testing for low fetal fraction DNA (no gNIPT result).</p> <p>3751/7701 samples without birth outcome were excluded (no reference standard).</p>
Comparative	
Aim to study	<p>To report the clinical application of gNIPT to detect chromosomal aneuploidies, especially T21, T18 and T13 in Chinese singleton pregnancies.</p>
Funding source or sponsor of the study	<p>Study not funded by industry but BGI-Shenzhen made sequencing and analysis. Some authors are employees of BGI-Shenzhen.</p>
Informations about the authors contacted	<p>Author was contacted on: 31 May 2016.</p> <p>No reply received from the author.</p>
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		

Zhou 2014b (Continued)

Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	High
DOMAIN 2: Index Test MPSS			
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear		
If a threshold was used, was it pre-specified?	Yes		
		Unclear	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all analysed patients receive the reference standard?	Yes		
Were all patients included in the analysis?	No		
		High	

CVS: chorionic villi sampling
 DANSR: digital analysis of selected regions
 FISH: fluorescence in situ hybridisation
 gNIPT: genomics-based non-invasive prenatal testing
 MAD: Median absolute deviation
 MPSS: massively parallel shotgun sequencing
 NCV: normalised chromosome value
 SD: standard deviation
 SNP: single nucleotide polymorphism
 TMPS: targeted massively parallel sequencing

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Anderson 2015	Not a diagnostic test accuracy study. Poster abstract.
Anselem 2016	Decision making study. Observational study. Not a diagnostic test accuracy study.
Bayindir 2015	Samples overlap with Brady 2016 . Most gNIPT results unconfirmed by a reference standard test. Insufficient information to derive 2 x 2 tables.
Beamon 2013	Poster abstract of the 33 rd Annual Meeting of the Society for Maternal-Fetal Medicine: The Pregnancy Meeting. Observational study with incomplete follow-up. Samples overlap with Beamon 2014 .
Beamon 2014	Observational study with incomplete follow-up. Not a diagnostic test accuracy study. Some gNIPT results unconfirmed by a reference standard test.
Belloin 2016	Most women (95%) completed a questionnaire to report their birth outcome (inappropriate reference standard for this review).
Benachi 2015b	Tribune. Not a diagnostic test accuracy study.
Benachi 2016	All samples overlap with Benachi 2015 .
Benn 2015	Letter to the editor on Zhang 2015 without data. Not a diagnostic test accuracy study.
Bhatt 2014	Poster abstract of the 18 th International Conference on Prenatal Diagnosis and Therapy, ISPD 2014. Patients with gNIPT negative result were without follow-up (no reference standard). Incomplete 2 x 2 tables.
Bianchi 2012a	Samples overlap with Bianchi 2012 . Data excluded to avoid double counting.
Bianchi 2014b	Editorial. Not a diagnostic test accuracy study.
Bianchi 2014c	Data excluded to avoid double counting. Samples overlap with Bianchi 2015b .
Bianchi 2015a	Not a diagnostic test accuracy study. Author presented some false positive cases in women with malignancy.
Bianchi 2015b	Incomplete 2 x 2 table. In this observational study, most women (98.9%) had no follow-up (no reference standard).
Bianchi 2015c	Not a diagnostic test accuracy study. Author presented some false positive cases in women with malignancy. Samples overlap with Bianchi 2015a .
Bianchi 2015d	Poster abstract. Not a diagnostic test accuracy study. Author presented some false positive cases in women with malignancy. Samples overlap with Bianchi 2015a .
BlueCross BlueShield Association 2014	Technology Evaluation Center Assessment. Review.
Brady 2016	Review with new data but most gNIPT results unconfirmed by a reference standard test. Insufficient information to derive 2 x 2 tables.
Chen 2013	Poster abstract of the ISPD 17 th International Conference on Prenatal Diagnosis and Therapy. Samples overlap with Huang 2014 .

Study	Reason for exclusion
Chen 2014	Poster abstract. Samples overlap with Yeang 2014 .
Cherry 2014	Poster abstract. Samples overlap with Meck 2015 .
Cheung 2015	Incomplete 2 x 2 table. This letter presented women who had positive results after screening and were referred for invasive procedure to confirm the presence of fetal aneuploidy. Only, the true positive and false positive gNIPT results were reported.
Chiu 2008	Proof-of-concept. Not a diagnostic test accuracy study.
Chiu 2010	Proof-of-concept. Not a diagnostic test accuracy study.
Christina 2012	Proof-of-concept. Not a diagnostic test accuracy study.
Cinnioglu 2012	Poster abstract. Samples overlap with Rabinowitz 2012a .
Cirigliano 2013	Full poster from the 17 th International Conference on Prenatal Diagnosis and Therapy, ISPD 2013 received. Samples overlap with Cirigliano 2014 and Ordoñez 2015 . Insufficient information to derive 2 x 2 tables.
Cirigliano 2014	Full poster from the 18 th International Conference on Prenatal Diagnosis and Therapy, ISPD 2014 received. Samples overlap with Cirigliano 2013 and Ordoñez 2015 . Insufficient information to derive 2 x 2 tables.
Cuckle 2015	Review with simulation model for gNIPT.
Curnow 2014	Poster abstract of the 18 th International Conference on Prenatal Diagnosis and Therapy, ISPD 2014. Samples overlap with Dar 2014 .
Dan 2012	Incomplete 2 x 2 table. Women with gNIPT negative results completed a questionnaire to report their birth outcome (inappropriate reference standard for this review).
Dar 2014	Implementation study. Incomplete 2 x 2 table. Most patients with gNIPT negative result were without follow-up (no reference standard). Some women had follow-up by telephone (inappropriate reference standard for this review).
De Ligt 2013	Case report (deletion).
Denona 2016	Poster abstract. Retrospective observational study. Insufficient information to derive 2 x 2 tables.
Discenza 2015	Poster abstract. Some gNIPT results unconfirmed by a reference standard test. Insufficient information to derive 2 x 2 tables.
Dobson 2015	Poster abstract. Insufficient information to derive 2 x 2 tables (gNIPT positive results only). Decision making. Samples overlap with Dobson 2016 .
Dobson 2016	Insufficient information to derive 2 x 2 tables (gNIPT positive results only). Decision making.
Dong 2016	Sequencing not based on maternal plasma ccfDNA.
Duenwald 2016	Method development. Analytical accuracy. Not a diagnostic test accuracy study.
Ehrich 2011a	Editorial comment without new data.
Eiben 2014	Review. Not a diagnostic test accuracy study.

Study	Reason for exclusion
Ellison 2015	Poster abstract. All gNIPT results (TMPS) were confirmed with a previous gNIPT result (MPSS) (inappropriate reference standard for this review). Insufficient information to derive 2 x 2 tables.
Faas 2011	Poster abstract of the 8 th European Cytogenetics Conference. Samples overlap with Faas 2012 .
Faas 2012	Proof-of-concept. Not a diagnostic test accuracy study.
Fairbrother 2013a	Observational study. Incomplete 2 x 2 data. Most patients were without follow-up (no reference standard).
Fairbrother 2013b	Conference abstract of the 17 th International Conference on Prenatal Diagnosis and Therapy, ISPD 2013. Samples overlap with Fairbrother 2013a .
Fan 2008	Proof-of-concept. Not a diagnostic test accuracy study.
Fang 2015	Insufficient information to derive 2 x 2 tables.
Ferres 2013	Not a diagnostic test accuracy study (implementation study).
Fiorentino 2015	Poster abstract. All samples overlap with Fiorentino 2016 .
Fosler 2015	Poster abstract of the 35 th Annual Meeting of the Society for Maternal-Fetal Medicine: The Pregnancy Meeting. Observational study. Incomplete 2 x 2 table. Most patients with gNIPT negative result were without follow-up (no reference standard).
Futch 2013	Observational study with incomplete follow-up. Incomplete 2 x 2 table. Many gNIPT results unconfirmed by a reference standard test.
Gabriel 2014	Conference abstract. Proof-of-concept.
Galea 2014	Full poster from the 18 th International Conference on Prenatal Diagnosis and Therapy, ISPD 2014. Incomplete 2 x 2 table. Most patients with gNIPT negative result were without follow-up (no reference standard).
Gao 2014	News, comment on Liao 2014 and Yuan 2013 without new data.
Gao 2015	Poster abstract. Insufficient information to derive 2 x 2 tables.
Geifman-Holtzman 2013	Poster abstract of the 33 rd Annual Meeting of the Society for Maternal-Fetal Medicine: The Pregnancy Meeting. Samples overlap with Xiong 2015 .
Geifman-Holtzman 2014	Poster abstract of the 34 th Annual Meeting of the Society for Maternal-Fetal Medicine: The Pregnancy Meeting. Samples overlap with Xiong 2015 .
Gerundino 2017	Women were asked to complete a questionnaire to report their birth outcome (inappropriate reference standard for this review). Insufficient information to derive 2 x 2 tables.
Gil 2013	Most patients with gNITP negative result were without follow-up (no reference standard) because 962 women had not yet delivered at the time of writing the publication. Insufficient information to derive 2 x 2 tables. Some patients overlap with del Mar Gil 2014 .
Gil 2015	Decision making including gNIPT accuracy data. All samples overlap with Gil 2016 .
Gnetetskaya 2015	Poster abstract. Samples overlap with Kurtser 2015 .

Study	Reason for exclusion
Grati 2014	Not a diagnostic test accuracy study. No sequencing data.
Gray 2013	Observational study. Not a diagnostic test accuracy study. Full poster received from the authors. Poster of the 17 th International Conference on Prenatal Diagnosis and Therapy, ISPD 2013.
Gromminger 2014	Data excluded to avoid double counting. Blinded DNA sequencing libraries were provided by Sequenom from their clinical trial cohort (NCT00877292) and were resequenced by LifeCodexx.
Guex 2013	Research letter. Samples overlap with Pescia 2017 .
Halks-Miller 2015	In reply to Bianchi 2015a . Not a diagnostic test accuracy study.
Harasim 2016	Poster abstract. Insufficient information to derive 2 x 2 tables.
Hernandez-Gomez 2015	Implementation study. Not a diagnostic test accuracy study.
Hofmann 2013	Poster abstract. Samples overlap with Stumm 2014 .
Hofmann 2014	Conference abstract. Insufficient information to derive 2 x 2 tables.
Hofmann 2015	Method development. Data were reanalysed by a new algorithmic approach of PraenaTest®. Not a diagnostic test accuracy study.
Hu 2014	Not a next generation sequencing publication. NIPT was ultrasound and serum biomarkers.
Hu 2015	Incomplete 2 x 2 tables. Only gNIPT positive results presented.
Hui 2015a	Poster abstract. All samples overlap with Hui 2015b .
Hui 2015b	Implementation study. Incomplete 2 x 2 table. Most patients with gNIPT negative result were without follow-up (no reference standard).
Jackson 2013	Poster abstract. All samples overlap with Jackson 2014 .
Jensen 2013	Proof-of-concept. Not a diagnostic test accuracy study. Samples overlap with Palomaki 2012 .
Jensen 2015	Proof-of-concept study with unblinded samples. Not a diagnostic test accuracy study.
Jin 2014	Incomplete 2 x 2 table. Women with gNIPT negative results were followed-up by telephone (inappropriate reference standard for this review).
Johnson 2013	Not a next generation sequencing method.
Juneau 2014	Method development. Incomplete 2 x 2 table. Most patients were without follow-up (no reference standard).
Kagan 2015	Not a diagnostic test accuracy study. Simulation model.
Kalantar 2014	Not next generation sequencing method.
Karlsson 2015	Methodological publication. Not a diagnostic test accuracy study.
Kershberg 2015	Poster abstract. Some gNIPT results unconfirmed by a reference standard test. Insufficient information to derive 2 x 2 tables.

Study	Reason for exclusion
Kinde 2012	Methodological publication. Not a diagnostic test accuracy study.
Korabecna 2012	Bioinformatic simulation with Palomaki 2011 data.
Koumbaris 2016	Method development (proof-of-concept study). Development of an advanced fetal fraction estimation method and aneuploidy determination algorithm. Not a diagnostic test accuracy study.
Kurtser 2015	Most patients with gNIPT negative result were without follow-up (no reference standard). Incomplete 2 x 2 tables.
Lambert-Messerlian 2014	Samples overlap with Palomaki 2011 and Palomaki 2012 . Data excluded to avoid double counting.
Larion 2015	Poster abstract. Implementation study.
Lau 2012a	Incomplete 2 x 2 table. Women with gNIPT negative results were followed up by telephone or by email (inappropriate reference standard for this review). All samples overlap with Lau 2014 .
Lau 2013	Sample overlap with Lau 2014 .
Lau 2014	Incomplete 2 x 2 table. Women with gNIPT negative results were followed up by telephone or by email (inappropriate reference standard for this review).
Lebo 2015	Incomplete 2 x 2 table.
Leung 2013	Proof-of-concept. Not a diagnostic test accuracy study.
Levandoski 2015	Poster abstract. Observational study about discordant gNIPT results. Insufficient information to derive 2 x 2 tables.
Levy 2013	Poster abstract. Incomplete 2 x 2 table.
Levy 2013a	Poster abstract. Incomplete 2 x 2 table.
Levy 2013b	Proof-of-concept. Not a diagnostic test accuracy study. Samples overlap with Zimmermann 2013 .
Li 2012	Methodological publication about relation between fetal fraction and multiple clinical factors.
Li 2015	Observational study. Unavailable information about gNIPT approach used. It is unclear if patients with gNIPT negative result were followed up (no reference standard).
Liao 2011	Not a diagnostic test accuracy study. No aneuploid case.
Liao 2012	Proof-of-concept. Not a diagnostic test accuracy study.
Liao 2013	This is a poster abstract. The full publication was also excluded. See Liao 2014 for reasons of exclusion.
Liao 2014	Incomplete 2 x 2 table for the retrospective and the prospective cohort. In prospective cohort, most patients were without follow-up (no reference standard). For the retrospective cohort, number of gNIPT results was not reported. Sensitivity and specificity were presented for the retrospective cohort but 2 x 2 tables could not be derived.
Liao 2014a	Letter to the editor about Bianchi 2014b without new data.
Liu 2015	Incomplete 2 x 2 table. Women with gNIPT negative results were followed up by telephone (inappropriate reference standard for this review).

Study	Reason for exclusion
Lo 2014	Bioinformatic development. Comparison of sensitivity and specificity using 3 different count normalisation methods.
Lo 2014a	Poster abstract. Bioinformatic development. Comparison of sensitivity and specificity using 3 different count normalisation methods. Samples overlap with Lo 2014 .
Loucký 2013	Samples overlap with Palomaki 2012 . Data excluded to avoid double counting.
Louis-Jacques 2014	Full poster. Observational study. Not a diagnostic test accuracy study.
Ma 2015	Samples overlap with Ma 2016 .
Ma 2015a	Poster abstract. All samples overlap with Ma 2016 .
Manotaya 2016	Insufficient information to derive 2 x 2 tables. Women without invasive testing results were encouraged to report birth outcomes through the insurance policy reimbursed (inappropriate reference standard for this review).
Marchili 2015	Poster abstract. Implementation study. Not a diagnostic test accuracy study. Insufficient information to derive 2 x 2 tables.
Mayen 2015	Observational study. Not a diagnostic test accuracy study.
Mazloom 2013a	Poster abstract. Samples overlap with Mazloom 2013 .
McCullough 2014	Incomplete 2 x 2 table. The clinician of women who passed gNIPT was encouraged to send ad hoc feedback to the lab (inappropriate reference standard for this review).
McCullough 2014a	Poster abstract. Incomplete 2 x 2 table. Samples overlap with McCullough 2014 .
McCullough 2015	Full poster of the 19 th International Conference on Prenatal Diagnosis and Therapy, ISPD 2015. Most gNIPT results unconfirmed by a reference standard test. Insufficient information to derive 2 x 2 tables. Some patients overlap with McCullough 2014 .
McLennan 2016	Most patients with gNIPT negative result were without follow-up (no reference standard). Insufficient information to derive 2 x 2 tables.
Meck 2014	Poster abstract. Samples overlap with Meck 2015 .
Meck 2015	Not a diagnostic test accuracy study. Observational study.
Meck 2015a	Poster abstract. Samples overlap with Meck 2015 .
Mennuti 2015	Review without original data.
Minarik 2015	gNIPT negative results unconfirmed by a reference standard test. Not a diagnostic test accuracy study.
Miron 2011	Not a diagnostic test accuracy study. This these explore traditional screening tests.
Mundy 2008	Health Technology Assessment. Not a diagnostic test accuracy study.
Mundy 2009	Health Technology Assessment. Not a diagnostic test accuracy study.
Musci 2014	Poster abstract. Samples overlap with Norton 2015 .

Study	Reason for exclusion
Musci 2014a	Poster abstract. Samples overlap with Hooks 2014 and Nicolaides 2014a .
NCT00770458	Not a gNIPT method (other method).
NCT00877292	Not with ccfDNA (other sampling).
NCT00891852	Not a gNIPT method (other method).
NCT00971334	Completed clinical trial but no published data.
NCT01052688	Incomplete 2 x 2 data (ongoing study with cases only).
NCT01256606	Not a gNIPT method (other method).
NCT01451671	Incomplete 2 x 2 data (ongoing study with cases only).
NCT01451684	Observational study on gNIPT without fetal karyotype.
NCT01555346	Completed clinical trial but no published data.
NCT01574781	Completed clinical trial but no published data.
NCT01597063	Completed clinical trial but no published data.
NCT01661010	Not a diagnostic test accuracy study.
NCT01663675	Adult with T21. Not with pregnant women (other population).
NCT01668251	Not a diagnostic test accuracy study.
NCT01725438	Not with ccfDNA (other sampling).
NCT01837979	Incomplete 2 x 2 data.
NCT01966991	Completed clinical trial but no published data.
NCT02127515	Not a diagnostic test accuracy study. Pregnant women with gNIPT have not a reference standard.
NCT02226315	Inappropriate reference standard for this review (pregnancy outcome data obtained from the patient).
NCT02872948	Not a gNIPT method (other method).
Neufeld-Kaiser 2015	Observational study. Not a diagnostic test accuracy study. Incomplete 2 x 2 tables. Most gNIPT results unconfirmed by a reference standard test.
Neveling 2015	Method validation for the NextSeq 500 platform. Not a diagnostic test accuracy study.
Nickolich 2016	Not a diagnostic test accuracy study.
Nicolaides 2013a	Poster abstract. All samples overlap with Nicolaides 2012 .
Nicolaides 2014	Simulation model on gNIPT implantation in first- or second-tier test.
Nicolaides 2014b	Note on Nicolaides 2014a without new data.

Study	Reason for exclusion
Nicolaidis 2014c	Target condition presented in this publication is not the focus of this review. Publication of next generation sequencing with ccfDNA for fetal triploidy.
Norem 2015	Full poster received from authors. Most patients were without follow-up (no reference standard). Incomplete 2 x 2 tables.
Norton 2014	Bioinformatic simulation.
Norton 2014a	Poster abstract. Samples overlap with Norton 2015 .
Norton 2015a	Bioinformatic simulation.
Norton 2015b	Editorial on Norton 2015 without new data.
Norton 2015c	Author reply to comments from Sentilhes 2015 and Smith-Bindman 2015 about Norton 2015 without new data.
Norton 2016	Simulation model to compare sequential and ccfDNA screening with data published in the literature. Not a diagnostic test accuracy study.
O'Leary 2014	Bioinformatic simulation.
Oepkes 2015	Most patients were without follow-up (no reference standard). Insufficient information to derive 2 x 2 tables.
Oneda 2016	Poster abstract. Insufficient information to derive 2 x 2 tables.
Ordoñez 2015	Full poster received. Some gNIPT results unconfirmed by a reference standard test. Insufficient information to derive 2 x 2 tables.
Palomaki 2011	Samples overlap with Palomaki 2012 (samples in Palomaki 2011 have been reanalysed in Palomaki 2012). Study excluded to avoid double counting.
Palomaki 2012a	Samples overlap with Palomaki 2012 . Conference abstract about Palomaki 2012 data.
Palomaki 2012b	Editorial on Palomaki 2011 without new data.
Palomaki 2015	Not a diagnostic test accuracy study.
Palomaki 2015a	Note about Palomaki 2015 . Not a diagnostic test accuracy study.
Perez-Pedregosa 2015	Incomplete 2 x 2 tables. Some women with gNIPT negative results were followed up by telephone (inappropriate reference standard for this review).
Pescia 2017	Follow-up for gNIPT negative results was ensured by an inquiry of two sets of randomly selected samples (inappropriate reference standard for this review).
Petersen 2014	Not a next generation sequencing publication. NIPT was ultrasound measurement and serum biomarkers.
Pettit 2014	Most patients with gNIPT negative result were without follow-up (no reference standard). Insufficient information to derive 2 x 2 tables.
Porreco 2014a	Reply to Grati 2014 without sequencing data.
Rabinowitz 2012	Poster abstract. Proof-of-concept. Not a diagnostic test accuracy study.

Study	Reason for exclusion
Rabinowitz 2012a	Poster abstract with incomplete 2 x 2 tables.
Rabinowitz 2012b	Poster abstract. Samples overlap with Rabinowitz 2012a .
Rabinowitz 2013	Poster abstract. Samples overlap with Pergament 2014 .
Rabinowitz 2014	Poster abstract. Sample overlap with Pergament 2014 .
Rad 2014	Implementation study without sequencing data presented.
Radoi 2015	Incomplete 2 x 2 tables. Most patients were without follow-up (no reference standard).
Rava 2012	Poster abstract. Samples overlap with Bianchi 2012 .
Rava 2014	Methodological publication about fetal DNA fraction with MELISSA samples.
Reiff 2015	Insufficient information to derive 2 x 2 tables.
Reiff 2016	Insufficient information to derive 2 x 2 tables.
Reimers 2015	Conference abstract from the 19 th International Conference on Prenatal Diagnosis and Therapy, ISPD 2015. Simulation model. Not a diagnostic test accuracy study.
Revello 2016	Not a diagnostic test accuracy study. All samples overlap with Gil 2016 and Quezada 2015 .
Ryan 2016	Method development of version 2 to SNP-based gNIPT. Not a diagnostic test accuracy study.
Sachse 2015	Proof-of-concept of fetal fraction quantification by qPCR.
Samura 2015	Most patients were without follow-up (no reference standard). Insufficient information to derive 2 x 2 tables. Samples overlap with Sago 2015 .
Sarno 2016	Some women reported their birth outcome (inappropriate reference standard for this review). Information about false positive results were insufficient to derive all 2 x 2 tables.
Schöck 2015	Poster abstract. Bioinformatics development with unblinded samples.
Sehnert 2013	Poster abstract. Incomplete 2 x 2 table.
Sehnert 2014	Poster abstract. Samples overlap with Bianchi 2014b .
Sentilhes 2015	Comment about Norton 2015 without new data.
Seo 2015	Women with gNIPT result were without follow-up at birth.
Settler 2015	Full poster received. Insufficient information to derive 2 x 2 tables. Some gNIPT results unconfirmed by a reference standard test.
Shani 2016	Simulation model. Not a diagnostic test accuracy study.
Shaohua 2012	Poster abstract. Full poster not received. Incomplete 2 x 2 table.
Sharma 2015	Poster abstract about patient perceptions of gNIPT from the multi-centered Canadian PEGASUS trial. gNIPT results compared with first trimester combined test (inappropriate reference standard for this review).

Study	Reason for exclusion
Shaw 2013	Poster abstract. Samples overlap with Shaw 2014 .
Shen 2016	Method development. Not a diagnostic test accuracy study.
Shi 2015	Incomplete 2 x 2 table. gNIPT negative result unconfirmed by a reference standard test.
Shulman 2014	Poster abstract. Incomplete 2 x 2 table. Most patients with gNIPT negative result were without a reference standard test.
Sistermans 2015a	Letter to the editor on Bianchi 2015a without data.
Smith-Bindman 2015	Comment about Norton 2015 without new data.
Song 2012	Poster abstract. Some samples overlap with Sparks 2012a .
Sparks 2012	Method development (all unblinded samples). Incomplete 2 x 2 table. Most patients with gNIPT negative result were unconfirmed by a reference standard test.
Srinivasan 2013	Poster abstract. Samples from MELISSA study (potentially overlap).
Stokowski 2015	Not a next generation sequencing method.
Strah 2015	Women were followed up by telephone interview to find out their birth outcome (inappropriate reference standard for this review).
Straver 2014	Proof-of-concept.
Strom 2015	Incomplete 2 x 2 table. Only women with gNIPT positive result were reported.
Stumm 2011	Proof-of-concept. Not a diagnostic test accuracy study.
Stumm 2012	Proof-of-concept. Not a diagnostic test accuracy study.
Stumm 2012a	Poster abstract. Samples overlap with Stumm 2014 .
Stumm 2013	Poster abstract. Samples overlap with Stumm 2014 .
Stumm 2016	Not a diagnostic test accuracy study.
Swanson 2012	Publication about Bianchi 2012 without new data.
Syngelaki 2014	Not a diagnostic test accuracy study. Simulation model.
Tan 2016	Women with gNIPT negative results were followed up by telephone interview (inappropriate reference standard for this review). Insufficient information to derive 2 x 2 tables.
Taneja 2016	Incomplete follow-up. Incomplete 2 x 2 table. Many gNIPT results unconfirmed by a reference standard test.
Taneja 2017	Most patients with gNIPT negative result were without reference standard test. Providers were encouraged to report discordant clinical outcomes. Insufficient information to derive 2 x 2 tables.
TARRIER 2015	gNIPT results unconfirmed by a reference standard test. Their reference method is verifi [®] results (inappropriate reference standard for this review).

Study	Reason for exclusion
Taylor 2014	Not a diagnostic test accuracy study. Observational study and decision making about gNIPT uptake in their center.
Togneri 2016	Full poster received. Internal verification set and implantation in their centre. Not a diagnostic test accuracy study.
Tong 2016	Not a next-generation sequencing method with ccfDNA.
Valderramos 2016a	Poster abstract. Insufficient information to derive 2 x 2 tables.
Valderramos 2016b	Poster abstract. Samples overlap with Valderramos 2016c .
Valderramos 2016c	Insufficient information to derive 2 x 2 tables. Retrospective cohort of patients with gNIPT positive results.
van den Oever 2012a	Proof-of-concept. Not a diagnostic test accuracy study.
van den Oever 2012b	Proof-of-concept. Not a diagnostic test accuracy study.
van den Oever 2013	Proof-of-concept. Not a diagnostic test accuracy study.
Van Opstal 2016	Simulation model. Not a diagnostic test accuracy study.
Verweij 2013a	Poster abstract. All samples overlap with Verweij 2013 .
Wald 2015a	Not a diagnostic test accuracy study. Prenatal screening workflow proposed.
Wald 2015b	Not a diagnostic test accuracy study. Prenatal screening workflow proposed.
Wang 2012	Incomplete 2 x 2 table. Women with gNIPT negative results were followed up by telephone (inappropriate reference standard for this review).
Wang 2015b	Not a diagnostic test accuracy study.
Wang 2015c	Proof-of-concept. Not a diagnostic test accuracy study.
Wang 2015d	Editorial on Wang 2015b without new data.
Wang 2015e	Not a diagnostic test accuracy study.
Xiong 2015	Full poster received. Observational study and incomplete follow-up.
Yankova 2015	Simulation model for gNIPT implantation. Not a diagnostic test accuracy study.
Yaron 2015	Commentary about gNIPT for microdeletion syndromes and rare autosomal trisomies. Not a diagnostic test accuracy study.
Yeang 2014	Proof-of-concept. Not a diagnostic test accuracy study.
Yu 2014	Proof-of-concept. Not a diagnostic test accuracy study.
Yuan 2013	Proof-of-concept. Not a diagnostic test accuracy study.
Zhang 2015	Incomplete 2 x 2 table. Women with gNIPT negative results were followed-up by telephone (inappropriate reference standard for this review).

Study	Reason for exclusion
Zhou 2013	Poster abstract. Incomplete 2 x 2 table.
Zimmermann 2012	Proof-of-concept. Not a diagnostic test accuracy study.
Zimmermann 2013	Proof-of-concept. Not a diagnostic test accuracy study.
Zwiefelhofer 2013	Implementation assessment of 2 sequencing platforms for gNIPT in a routine clinical environment. Not a diagnostic test accuracy study.

ccfDNA: circulating cell-free DNA

gNIPT: genomics-based non-invasive prenatal testing

MPSS: massively parallel shotgun sequencing

TMPS: targeted massively parallel sequencing

Characteristics of ongoing studies [author-defined order]

Basaran 2015

Trial name or title	Publication's title: False positive and false negative results of cell free DNA testing.
Target condition and reference standard(s)	Target conditions: T21, T18, T13, 45,X, 47,XXY, 47,XYY and 47,XXX. Reference standard: fetal karyotype of chorionic villi or amniotic fluid.
Index and comparator tests	gNIPT by TMPS or MPSS by commercial company providing gNIPT in Turkey (Ariosa Diagnostics, Inc., BGI-Shenzhen, Illumina, Inc, Natera, Inc. and Sequenom, Inc). Blood samples for gNIPT were collected before reference standard.
Starting date	Not reported.
Contact information	Dr Seher Basaran Department of Medical Genetics Istanbul University, Istanbul Medical Faculty TURKEY 90 (212) 4142000 basarabs@istanbul.edu.tr
Aim to study	To demonstrate the importance of confirmation of fetus genotype by invasive testing after gNIPT.
Funding source or sponsor of the study	The genetic centre is not affiliated with any commercial company providing gNIPT.
Information about the authors contacted	Author was contacted on: 12, 14 and 18 January 2016. Last reply received on: 19 January 2016.
Notes	At the time of this writing, the authors plan to publish a full publication soon.

Buresch 2016

Trial name or title	Poster's title: Actual rates of recommended diagnostic testing after first-trimester screening vs same-day screening by cell free DNA.
Target condition and reference standard(s)	Target conditions: T21, T18, T13, 45,X, 47,XXY, 47,XYY and 47,XXX. Reference standard: not reported.
Index and comparator tests	MPS.
Starting date	January to June 2015.
Contact information	Susan Klugman Department of Obstetrics and Gynecology and Women's Health, Albert Einstein College of Medicine, Montefiore Medical Center 1695 Eastchester Road, Bronx, NY 10461, United States. sklugman@montefiore.org
Aim to study	To compare actual patient referrals for post-screen diagnostic tests following first-trimester screening vs same day ccfDNA.
Funding source or sponsor of the study	Not reported.
Information about the authors contacted	Author was contacted on: 1 and 23 September 2016. Reply received on: 23 September 2016.
Notes	Authors are working on data at the time of writing and they plan to submit for publication.

Chen 2011a

Trial name or title	Oral presentation's title: Noninvasive prenatal diagnosis of fetal aneuploidy by massively parallel sequencing of maternal plasma DNA.
Target condition and reference standard(s)	Target conditions: T21, T18, T13 and SCA. Reference standard: fetal karyotype.
Index and comparator tests	gNIPT by MPSS on Illumina GAIIx/HiSeq 2000 sequencer. Cutpoint: positive if t score < -4. Commercial test: BGI's test.
Starting date	Not reported.
Contact information	Fang Chen, Beijing Genomics Institute, Shenzhen, China
Aim to study	To assess gNIPT with ccfDNA performance on fetal aneuploidies.
Funding source or sponsor of the study	Not reported.
Information about the authors contacted	BGI was contacted on: 19 May 2016. No reply received from the author.

Chen 2011a (Continued)

Notes	Cohort of 5268 pregnant women. They successfully identified 62 cases of T21, 40 cases of T18, 3 cases of T13, 13 cases of SCA. In a cohort of karyotyping cases, the sensitivity and specificity of the aneuploidy fetus detection was 100% and 100%, respectively.
-------	---

Da Fonseca 2015

Trial name or title	Abstract's title: Non-Invasive prenatal testing for the most common aneuploidies (trisomies 21, 18, and 13) using a semiconductor-sequencing platform: a French multicenter pilot study.
Target condition and reference standard(s)	Target conditions: T21, T18 and T13. Reference standard: fetal karyotype.
Index and comparator tests	gNIPT on semiconductor sequencing platform (MPSS). Blood sample collection not reported. Cutpoint: not reported.
Starting date	Not reported.
Contact information	J.P. Da Fonseca, Inserm U1016 Plateforme Génomique, Paris, France.
Aim to study	To validate a common protocol and to evaluate the efficiency and reliability of gNIPT of the most common chromosomal aneuploidies using a semiconductor sequencing platform.
Funding source or sponsor of the study	Not reported.
Information about the authors contacted	Author was contacted on: 19 January 2016 and 23 March 2016. Reply received on: 16 February 2016.
Notes	Conference Abstract of the 10 th European Cytogenetics Conference of the European Cytogenetics Association, ECA 2015. Prospective study of 500 pregnant women at high risk of fetal aneuploidy who undergo fetal karyotyping. The NIPT results matched the fetal karyotyping results in all of the cases: all trisomies were detected.

ISRCTN11174071

Trial name or title	Comparison of false positive rates in prenatal combined screening and cell free DNA screening for trisomy 21 (ReFaPo study).
Target condition and reference standard(s)	Target condition: T21. Reference standard: prenatal or postnatal karyotype.
Index and comparator tests	gNIPT.
Starting date	July 2016.
Contact information	Karl Oliver Kagan University of Tuebingen Department of Obstetrics and Gynaecology Calwerstrasse 7 Tuebingen 72076

ISRCTN11174071 (Continued)

Germany

Aim to study	To compare the false positive rate of cell-free DNA and traditional screening methods in a randomised controlled trial in a cohort without prior risk of fetal aneuploidy.
Funding source or sponsor of the study	Study funded by CENATA GmbH who does the analysis.
Information about the authors contacted	No need for further contact.
Notes	<p>Target number of participants: 1400.</p> <p>Recruitment end date: March 2017.</p> <p>Intention to publish date: October 2018.</p> <p>DOI 10.1186/ISRCTN11174071</p>

Lin 2014

Trial name or title	Clinical implementation of noninvasive prenatal testing in twin pregnancies with assisted reproductive technique treatment.
Target condition and reference standard(s)	<p>Target conditions: T21, T18 and T13.</p> <p>Reference standards: fetal karyotype or clinical outcomes.</p>
Index and comparator tests	<p>gNIPT by MPSS.</p> <p>Commercial test: BGI Shenzhen's prenatal test.</p>
Starting date	Not reported.
Contact information	<p>BGI-Shenzhen</p> <p>Shenzhen, China</p>
Aim to study	To assess the clinical implementation of MPS-based NIPT in twin pregnancies with assisted reproductive technique treatment.
Funding source or sponsor of the study	Not reported but BGI-Shenzhen made sequencing and analyses.
Information about the authors contacted	No need for further contact.
Notes	Some women were still pregnant at the time of writing this poster abstract.

Mu 2014

Trial name or title	Maternal non-invasive fetal DNA test used in prenatal diagnosis.
Target condition and reference standard(s)	Target conditions: T21, T18, T13 and 45,X.

Mu 2014 (Continued)

Reference standard: fetal karyotype of amniotic fluid.

Index and comparator tests	gNIPT by MPSS. NIFTY™ prenatal test by BGI-Shenzhen.
Starting date	In 2012.
Contact information	Mu Y. Beijing United Family Hospital Beijing, China.
Aim to study	Not reported.
Funding source or sponsor of the study	Not reported.
Information about the authors contacted	Author was contacted on: 19 April and 19 May 2016. No reply received from the author.
Notes	Poster abstract. Some women were still pregnant at the time of writing this poster abstract.

NCT01429389

Trial name or title	Specimen collection from pregnant women at increased risk for fetal aneuploidy.
Target condition and reference standard(s)	Target condition: T21. Reference standard: fetal karyotype.
Index and comparator tests	gNIPT.
Starting date	May 2011.
Contact information	Sequenom, Inc.
Aim to study	To develop a prenatal aneuploidy test using ccfDNA from blood samples from pregnant women who have an increased risk indicator/s for fetal chromosomal aneuploidy detection (T21).
Funding source or sponsor of the study	Study funded by Sequenom, Inc.
Information about the authors contacted	No need for further contact.
Notes	

NCT01472523

Trial name or title	A safer pre-natal diagnosis using free DNA in maternal blood (IONA®).
Target condition and reference standard(s)	Target conditions: T21, T18, T13 and other chromosomal abnormalities yet to be determined.

NCT01472523 (Continued)

	Reference standards: prenatal karyotype and follow-up for 1 year.
Index and comparator tests	gNIPT by TMPS (selective amplification of fetal DNA) by Premaitha Health.
Starting date	April 2007.
Contact information	Brenda Kelly National Health Service, United Kingdom
Aim to study	To validate a novel gNIPT method that could increase the titre of fetal DNA within a given sample.
Funding source or sponsor of the study	Study funded by Premaitha Health.
Information about the authors contacted	No need for further contact.
Notes	

NCT01545674

Trial name or title	Prenatal Non-invasive Aneuploidy Test Utilizing SNPs trial (PreNATUS).
Target condition and reference standard(s)	Target conditions: aneuploidy in a fetus at chromosomes 13, 18, 21, X and Y. Reference standard: fetal karyotype.
Index and comparator tests	gNIPT by TMPS (SNP based technology by Natera, Inc.).
Starting date	January 2012.
Contact information	Ronald Wapner, MD, Columbia University
Aim to study	To assess the diagnostic capability of an informatics enhanced SNP based technology (Parental Support) to identify pregnant women who are carrying a fetus with an aneuploidy from free floating DNA in the maternal blood.
Funding source or sponsor of the study	Study funded by Natera, Inc.
Information about the authors contacted	No need for further contact.
Notes	

NCT01925742

Trial name or title	Study of the efficacy of new non-invasive prenatal tests for screening for fetal trisomies using maternal blood (PEGASUS).
Target condition and reference standard(s)	Target conditions: T21, T18 and T13.

NCT01925742 (Continued)

	Reference standards: prenatal or neonatal karyotype or medical record from birth.
Index and comparator tests	gNIPT by Semiconductor MPSS (Ion Torrent Proton™) or optical-based MPSS (Illumina) or by TMPS with Harmony™ prenatal test by Ariosa Diagnostics, Inc.
Starting date	August 2013.
Contact information	François Rousseau CHU de Québec Québec, Canada
Aim to study	To perform a pan-Canadian large-scale validation study comparing the relative effectiveness and clinical performances of 2 index gNIPT methods using fetal ccfDNA in maternal blood in Canadian clinical laboratories between themselves and with that of fetal karyotype for detecting fetal aneuploidy of chromosomes 13, 18 and 21 and to compare the accuracy of this new gNIPT method with traditional prenatal screening methods.
Funding source or sponsor of the study	Study funded by Centre Hospitalier Universitaire de Québec, Laval University, Genome Canada, Genome Quebec, Genome British Columbia and Canadian Institutes of Health Research (CIHR).
Information about the authors contacted	No need for further contact.
Notes	Recruitment of patients completed (near 5000 pregnant women enrolled). at the time of writing, they are sequencing 3600 pregnant women with the 2 gNIPT MPSS platforms. A subsample of about 2300 blood samples was analysed by Ariosa Diagnostics, Inc (TMPS). Estimated study completion date: June 2017.

NCT02201862

Trial name or title	Non-invasive Chromosomal Evaluation of Trisomy study (NICHE).
Target condition and reference standard(s)	Target conditions: T21, T18 and T13. Reference standard: fetal karyotype.
Index and comparator tests	gNIPT by TMPS by Ariosa Diagnostics, Inc.
Starting date	April 2014.
Contact information	Romielle Aquino 408-209-9098 raqino@ariosadx.com Or Thomas Musci 408-229-7500 tmusci@ariosadx.com

NCT02201862 (Continued)

Aim to study	To provide clinically annotated samples to support continued improvements in the Ariosa Diagnostics, Inc Test content, methodology, specimen processing and quality control.
Funding source or sponsor of the study	Study funded by Ariosa Diagnostics, Inc.
Information about the authors contacted	No need for further contact.
Notes	

NCT02278536

Trial name or title	Multiple gestation study.
Target condition and reference standard(s)	Target conditions: T21, T18, T13 and SCA. Reference standards: fetal karyotype (amniocentesis or CVS) or genetic testing from cheek swab or saliva from live-born children.
Index and comparator tests	gNIPT by TMPS by Natera, Inc.
Starting date	March 2013.
Contact information	Brian Kirshon Houston Perinatal Associates Or Zach Demko Natera, Inc.
Aim to study	To demonstrate the accuracy of our new NATUS diagnostic method to determine the genetic health of the developing fetuses in a multiple gestation pregnancy from a maternal blood sample.
Funding source or sponsor of the study	Study funded by Natera, Inc.
Information about the authors contacted	No need for further contact.
Notes	

NCT02278874

Trial name or title	High risk multiple gestation study.
Target condition and reference standard(s)	Target conditions: T21, T18, T13 and SCA. Reference standards: fetal karyotype (amniocentesis or CVS) or genetic testing from cheek swab or saliva from live-born children.

NCT02278874 (Continued)

Index and comparator tests	gNIPT by TMPS by Natera, Inc.
Starting date	August 2014.
Contact information	Joanne Stone Mt. Sinai Hospital, New York
Aim to study	To demonstrate the accuracy of our proprietary algorithm method to determine the genetic health of the developing fetuses in a multiple gestation pregnancy from a maternal blood sample.
Funding source or sponsor of the study	Study funded by Natera, Inc., Mount Sinai Hospital New York, Montefiore Medical Center, Long Island Jewish Medical Center and Tufts Medical Center.
Information about the authors contacted	No need for further contact.
Notes	

NCT02317965

Trial name or title	Non-invasive screening for fetal aneuploidy.
Target condition and reference standard(s)	Target conditions: T21 and T18. Reference standard: fetal karyotype.
Index and comparator tests	gNIPT by MPSS by Progenity, Inc.
Starting date	March 2015.
Contact information	Richard Porreco Obstetrix Medical Group of Colorado
Aim to study	To detect whole chromosome abnormalities on all chromosomes 13, 16, 18, 21, X and Y, in the fetus through analysis of ccfDNA and compound sample DNA in maternal blood.
Funding source or sponsor of the study	Study funded by Progenity, Inc.
Information about the authors contacted	No need for further contact.
Notes	

NCT02424474

Trial name or title	T21,18 and 13 screening by cell free fetal DNA in low risk patients (DEPOSA).
Target condition and reference standard(s)	Target conditions: T21, T18 and T13. Reference standard: fetal karyotype.
Index and comparator tests	gNIPT by MPSS.

NCT02424474 (Continued)

Starting date	June 2015.
Contact information	Alexandra Benachi Antoine Bécèle Hospital
Aim to study	To evaluate the performance of gNIPT in a population of pregnant women with and without in vitro fertilisation (IVF) concomitantly to regular first-trimester trisomy 21 (T21) screening using maternal age, nuchal fold measurement and serum screening.
Funding source or sponsor of the study	Study funded by Assistance Publique - Hôpitaux de Paris.
Information about the authors contacted	No need for further contact.
Notes	Recruitment of patients completed (933 pregnant women enrolled).

NCT02787486

Trial name or title	Expanded Noninvasive Genomic Medical Assessment: the Enigma study.
Target condition and reference standard(s)	Target conditions: T21, T18, T13, microdeletion syndromes, sex chromosome abnormalities, infectious and other diseases, and blood group typing. Reference standard: fetal karyotype or medical records.
Index and comparator tests	gNIPT by MPSS provided by Progenity, Inc.
Starting date	October 2015.
Contact information	Paul Bien 760-494-1743 paul.bien@progenity.com
Aim to study	To evaluate the relative clinical sensitivity, specificity, and performance of the laboratory-developed test as a screening test for fetal chromosomal aneuploidy, infectious and other diseases, and RhD genotyping in the general population of pregnant women.
Funding source or sponsor of the study	Study funded by Progenity, Inc.
Information about the authors contacted	No need for further contact.
Notes	

Sago 2015

Trial name or title	Nationwide demonstration project of next-generation sequencing of cell-free DNA in maternal plasma in Japan: 1-year experience
---------------------	--

Sago 2015 (Continued)

Target condition and reference standard(s)	Target conditions: T21, T18 and T13. Reference standards: fetal karyotype of chorionic villi or amniotic fluid or tissues of the miscarriage or medical record from birth.
Index and comparator tests	gNIPT by MPSS. Blood samples were collected before invasive procedure. Commercial test: MaterniT21™ prenatal test from Sequenom, Inc.
Starting date	15 November 2012. Recruitment period between April 2013 to March 2014.
Contact information	Haruhiko Sago National center for Child-health and development Perinatal Center 2-10-1Ookura, Setagaya-ku, Tokyo 03-3416-0181 sagou-h@ncchd.go.jp
Aim to study	To evaluate the quality of the genetic counselling in Japan. Sago 2015 reported the 1-year experience of a nationwide demonstration project to introduce gNIPT of fetal aneuploidy from maternal plasma and discuss how to implement this program in Japan.
Funding source or sponsor of the study	Study supported by the Grant of the National Center for Child Health and Development 24-3, Japan. Sequenom, Inc made gNIPT.
Information about the authors contacted	Author was contacted on: 6 April and 14 June 2016. No reply received from the author.
Notes	Authors continue collecting follow-up data in the study population.

Sanchez-Usabiaga 2015

Trial name or title	Clinical implementation of non-invasive prenatal study for detecting aneuploidies by fetal DNA based on single nucleotide polymorphisms: 2 years in Mexico.
Target condition and reference standard(s)	Target conditions: T21, T18, T13, 45,X, 47,XXY, 47,XYY and 47,XXX. Reference standards: fetal karyotype of chorionic villi or amniotic fluid or medical record from birth.
Index and comparator tests	gNIPT by TPMS. Commercial test: Natera's prenatal test.
Starting date	Recruitment period: March 2013 to February 2015.
Contact information	Dr. Rafael Sánchez Usabiaga rsanchez@medicafertil.com.mx
Aim to study	To describe our experience of 2 years integrating gNIPT by ccfDNA in its variant of single nucleotide polymorphism (SNPs) as a screening method for the detection of common aneuploidies, since 9 weeks of gestation.

Sanchez-Usabiaga 2015 *(Continued)*

Funding source or sponsor of the study	Not reported but Natera, Inc. made gNIPT sequencing and analyses.
Information about the authors contacted	No need for further contact.
Notes	There are 270 pregnant women included in this study.

Sisternans 2015

Trial name or title	TRIDENT: or monitored NIPT implementation in the Netherlands.
Target condition and reference standard(s)	Target conditions: T21, T18 and T13. Reference standard: fetal karyotype of chorionic villi or amniotic fluid is recommended in case of abnormal gNIPT test results. Neonatal clinical examination not mentioned.
Index and comparator tests	gNIPT by MPSS.
Starting date	01 April 2014.
Contact information	Dr. Erik Sisternans. VU University Medical Center Dept. of Clinical and Human Genetics Van der Boechorststraat 7 1081 BT Amsterdam NETHERLANDS +31-20-020-4448346 Email: e.sisternans@vumc.nl
Aim to study	To investigate and evaluate all relevant aspects of the introduction of NIPT in the Dutch prenatal screening program.
Funding source or sponsor of the study	The TRIDENT study was designed and proposed by the national multidisciplinary NIPT consortium.
Information about the authors contacted	Author have been contacted on: 9 December 2015 and 15 March 2016. Reply received on: 16 March 2016.
Notes	Conference abstract presented at the Annual conference of the European Society of Human Genetics at Glasgow, Scotland, UK. http://www.emgo.nl/research/quality-of-care/research-projects/1451/trident-study-trial-by-dutch-laboratories-for-evaluation-of-non-invasive-prenatal-testing-nipt/background/ The authors plan to publish a full publication soon.

Torres 2015

Trial name or title	Genetic non invasive prenatal testing: A clinical and technical experience of 3000 cases with follow-up.
Target condition and reference standard(s)	Target conditions: T21, T18, T13, 45,X, 47,XXY, 47,XYY and 47,XXX. Reference standard: fetal karyotype of amniotic fluid.

Torres 2015 (Continued)

Index and comparator tests	gNIPT by MPSS. Commercial test: TrisoNIM® prenatal test by NIMGenetics Genomics.
Starting date	Not reported.
Contact information	Juan C Cigudosa NIMGenetics Genomics Madrid, Spain.
Aim to study	To show a NIPT protocol, called TrisoNIM®, which has been partially performed in our laboratory, based in massive parallel sequencing.
Funding source or sponsor of the study	Not reported.
Information about the authors contacted	Author were contacted on: 29 February, 22 March, 15 and 27 June 2016. Reply received on: 20 June 2016.
Notes	Full poster received from the authors.

Van Wymersch 2015

Trial name or title	Introduction of noninvasive prenatal testing for fetal trisomies: preliminary results and consequences on invasive samplings.
Target condition and reference standard(s)	Target conditions: T21, T18, T13, 45,X, 47,XXY, 47,XYY and 47,XXX. Microdeletion syndromes can also be detected. Reference standards: fetal karyotype of chorionic villi or amniotic fluid or medical record from birth.
Index and comparator tests	gNIPT by MPSS. Blood samples were collected before invasive procedure. Cutpoint: not reported. Commercial test: NIFTY™ test (Bejing Genomics Institute, Hong-Kong, China).
Starting date	December 2013.
Contact information	Dr Van Wymersch Didier, Service de Gynécologie Obstétrique, Centre Hospitalier de Luxembourg, 4 Rue Barblé, L-1210 Luxembourg E-mail: vanwymersch.didier@chl.lu
Aim to study	To analyse a year of gNIPT implantation in our institute and to analyse gNIPT implication in chromosomal abnormalities screening politic.
Funding source or sponsor of the study	No reported. Samples analysed at BGI.
Information about the authors contacted	Author was contacted on: 12 September 2016. Reply received on: 29 September 2016.

Van Wymersch 2015 *(Continued)*

Notes	This publication showed the first 683 samples. At the time of writing, authors have a much larger population of 2132 pregnant women. No false negative results have been observed to date for all the pregnancies that have already come to term.
-------	---

Willems 2014

Trial name or title	The first 3000 Non-Invasive Prenatal Tests (NIPT) with the Harmony test in Belgium and the Netherlands.
Target condition and reference standard(s)	Target conditions: T21, T18 and T13. Reference standard: fetal karyotype of chorionic villi or amniotic fluid.
Index and comparator tests	gNIPT by TMPS. Commercial test: Harmony™ prenatal test by Ariosa Diagnostics, Inc.
Starting date	Recruitment period: March 2013 to December 2013.
Contact information	Patrick Willems patrick.willems@genetic-diagnostic.net
Aim to study	To report the results of the first 3000 consecutive gNIPT tests performed in pregnant women from Belgium and the Netherlands.
Funding source or sponsor of the study	Not reported. Ariosa Diagnostics, Inc made sequencing and analysis.
Information about the authors contacted	No need for further contact.
Notes	

Yu 2014a

Trial name or title	Maternal non-invasive fetal DNA test used in prenatal diagnosis.
Target condition and reference standard(s)	Target conditions: T21, T18, T13 and 45,X. Reference standards: cytogenetic tests and postnatal follow-up.
Index and comparator tests	gNIPT by MPSS by BGI-Shenzhen.
Starting date	Patients recruited in 2012.
Contact information	Yu M or Fei S. Beijing United Family Hospital.
Aim to study	To determine gNIPT accuracy in Chinese population.
Funding source or sponsor of the study	Not reported.

Yu 2014a (Continued)

Information about the authors contacted	Author was contacted on: 15 May 2016. BGI was contacted on: 19 May 2016. No reply received from the author or BGI.
Notes	Conference abstract. Some women were still pregnant at the time of writing their conference abstract.

Zwiefelhofer 2014

Trial name or title	Prenatal detection of fetal aneuploidy on the Ion Torrent Proton™ platform.
Target condition and reference standard(s)	Target condition: T21. Reference standard: fetal karyotype.
Index and comparator tests	gNIPT by MPSS on the Proton™ platform.
Starting date	Not reported.
Contact information	Sequenom, Inc.
Aim to study	To examine the performance of a gNIPT for fetal aneuploidy on the Ion Torrent Proton™ platform.
Funding source or sponsor of the study	Study funded by Sequenom, Inc.
Information about the authors contacted	Author was contacted on: 19 April and 15 June 2016. Reply received on: 22 June 2016.
Notes	Full poster received from authors. This study includes 156 samples including 16 women carrying a T21 fetus. All patient samples were correctly identified according to their karyotype results.

CVS: chorionic villi sampling
 gNIPT: genomics-based non-invasive prenatal testing
 MPSS: massively parallel shotgun sequencing
 TMPS: targeted massively parallel sequencing

DATA

Presented below are all the data for all of the tests entered into the review.

Table Tests. Data tables by test

Test	No. of studies	No. of participants
1 MPSS T21	41	50133
2 MPSS T18	38	49003
3 MPSS T13	29	46090

Test	No. of studies	No. of participants
4 MPSS 45,X	14	7867
5 MPSS 47,XXX	5	5449
6 MPSS 47,XXY	8	6588
7 MPSS 47,XXY	8	6629
8 MPSS all 7 aneuploidies	44	50864
9 MPSS, autosomes	43	50453
10 MPSS, SCA	14	7911
11 TMPS T21	16	32487
12 TMPS T18	12	30319
13 TMPS T13	10	22868
14 TMPS 45,X	6	2214
15 TMPS 47,XXX	2	586
16 TMPS 47,XXY	4	1021
17 TMPS 47,XXY	2	358
18 TMPS all 7 aneuploidies	21	35275
19 TMPS, autosomes	18	34473
20 TMPS, SCA	6	2214
21 Traditional screening tests, autosomes	5	24279
22 Traditional screening tests T21	2	17753
23 Traditional screening tests T18	2	17747
24 Traditional screening tests T13	1	11185

Test 1. MPSS T21.

Test 2. MPSS T18.

Test 3. MPSS T13.

Test 4. MPSS 45,X.

Test 5. MPSS 47,XXX.

Test 6. MPSS 47,XXY.

Test 7. MPSS 47,XXY.

Test 8. MPSS all 7 aneuploidies.

Test 9. MPSS, autosomes.

Test 10. MPSS, SCA.

Test 11. TMPS T21.

Test 12. TMPS T18.

Test 13. TMPS T13.

Test 14. TMPS 45,X.

Test 15. TMPS 47,XXX.

Test 16. TMPS 47,XXY.

Test 17. TMPS 47,XY.

Test 18. TMPS all 7 aneuploidies.

Test 19. TMPS, autosomes.

Test 20. TMPS, SCA.

Test 21. Traditional screening tests, autosomes.

Test 22. Traditional screening tests T21.

Test 23. Traditional screening tests T18.

Test 24. Traditional screening tests T13.

ADDITIONAL TABLES

Table 1. Characteristics of target conditions

Target condition	Affected births ^a /100,000	Clinical features	Prognosis
T21	140 to 230 ^{b,c}	Intellectual disability (mild to moderate), neurodevelopmental problems, characteristic dysmorphic features, congenital defects (cardiac (44% to 58%) and gastrointestinal system (4% to 10%)), vision or hearing impairment (38% to 80%) and obstructive sleep apnoea syndrome (57%) ^{d,e}	Mean and median life expectancies are estimated to be 51 and 58 years old ^f
T18	59 ^c	Severe intellectual disability and a wide range of significant malformations (cardiac defects, gastrointestinal system defects, renal anomalies, central nervous system defects (apnoea and seizures)) ^{d,g}	Most affected fetuses die in utero. Median survival has been estimated at 14 days (95% confidence interval (CI) 10 to 20) and 8% (95% CI 4 to 14) reach 1 year of age ^h

Table 1. Characteristics of target conditions (Continued)

T13	23 ^c	Severe intellectual disability, seizures and several dysmorphic features, malformations of the extremities, cardiac defects, renal anomalies, and abdominal wall defects ^{d,i}	Most affected fetuses die in utero. Median survival time has been estimated at 10 days (95% CI 7 to 19) and 8% (95% CI 4 to 14) reach 1 year of age ^h
45,X	30 to 50 ^{c,j}	Learning disabilities (70%), short stature, congenital heart diseases (30%) and gonadal dysgenesis (90% with amenorrhoea and infertility due to early ovarian failure) ^{k,l}	Mortality in 45,X women is 3-fold higher than in the general population with an average life span of 69 years ^m
47,XXY	12 ^c	Learning disabilities (> 75%), small testes (> 95%), azoospermia (> 95%), male infertility (91% to 99%), decreased testosterone level (63% to 85%) and gynaecomastia (38% to 75%) ^{l,n}	Life expectancy is slightly shorter (approximately 2 years) than euploid men ^o
47,XXX	6 ^c	Developmental delays (motor and speech), learning or intellectual disability, attention deficits (25% to 35%), mood disorders (anxiety and depression), tall stature (80% to 89%), clinodactyly (42% to 65%), hypotonia in infancy (55% to 71%), genitourinary malformations and congenital heart defects ^o	Mortality significantly increased with a median survival age of 70.9 years compared to 81.7 years for euploid females ^p
47,XYY	3 ^c	Developmental delays (speech, language and motor), attention deficit disorder (52%), tall stature (78%), central adiposity, macrocephaly (33%), hypotonia (63%), clinodactyly (52%), hypertelorism (59%) and testicular enlargement for age (50%) but no increase in genital anomalies ^q	Mortality increased with a reduction of life span of 10.3 years compared to euploid men ^r

45,X: Turner syndrome, 47,XXX: triple X syndrome, 47,XXY: Klinefelter syndrome, T21: trisomy 21, T18: trisomy 18, T13: trisomy 13.

^aIncluding live births, fetal deaths and terminations of pregnancy.

^b(Christianson 2006; Parker 2010)

^c(Wellesley 2012)

^d(Driscoll 2009)

^e(Irving 2012; Weijerman 2010)

^f(Wu 2013b)

^g(Cereda 2012)

^h(Wu 2013a)

ⁱ(Chen 2009)

^j(Stochholm 2006)

^k(Karnis 2012; Mazzanti 1998; Sybert 2004)

^l(Tyler 2004)

^m(Saenger 1996; Schoemaker 2008)

ⁿ(Groth 2013)

^o(Tartaglia 2010)

^p(Stochholm 2010b)

^q(Bardsley 2013; Leggett 2010)

^r(Stochholm 2010a).

Table 2. Reported accuracy of commercially available genomics-based non-invasive prenatal tests^a

Test name (Company,	Method	Aneuploidy	Reported sensitivity	Reported specificity	Reported false positive
------------------------	--------	------------	-------------------------	-------------------------	----------------------------

Genomics-based non-invasive prenatal testing for detection of fetal chromosomal aneuploidy in pregnant women (Review)

234

Table 2. Reported accuracy of commercially available genomics-based non-invasive prenatal tests^a (Continued)

country)			% (95% CI)	% (95% CI)	rate %
Bambni™	MPSS	T21	100.0 (ND)	> 99.9 (ND)	< 0.1
Test		T18	100.0 (ND)	> 99.9 (ND)	< 0.1
(Berry Genomics Co. Ltd, China)		T13	100.0 (ND)	> 99.9 (ND)	< 0.1
		45,X	100.0 (ND)	99.8 (ND)	0.0
		47,XXX	100.0 (ND)	100.0 (ND)	0.1
		47,XXY	100.0 (ND)	100.0 (ND)	0.0
		47,XYY	100.0 (ND)	100.0 (ND)	0.0
GENOMOM	MPSS	T21, T18 and T13	99.0 (ND)	ND	ND
(Genome Care, Korea)		SCA	95.0 (ND)	ND	ND
Harmony™	Oligo TMPS	T21	> 99.0 (ND)	> 99.9 (ND)	< 0.1
prenatal test		T18	97.4 (ND)	> 99.9 (ND)	< 0.1
(Ariosa Diagnostics, Inc., USA)		T13	93.8 (ND)	> 99.9 (ND)	< 0.1
		45,X ^b	96.3 (81.7 to 99.8)	99.5 (98.1 to 99.9)	0.5
		47,XXX ^b	100.0 (ND)	99.5 (98.1 to 99.9)	0.5
		47,XXY ^b	100.0 (61.0 to 100.0)	100.0 (99.0 to 100.0)	0.0
IONA® test	MPSS	T21	> 99.0 (ND)	> 99.0 (ND)	< 1.0
(Premaita Health plc, UK)		T18	> 99.0 (ND)	> 99.0 (ND)	< 1.0
		T13	> 99.0 (ND)	> 99.0 (ND)	< 1.0
(Laboratoire CERBA, France)	MPSS	T21, T18 and T13	> 99.8 (ND)	> 99.8 (ND)	< 0.2
MaterniT21™	MPSS	T21	99.1 (96.6 to 99.9)	99.9 (99.7 to 99.9)	0.1
Plus test		T18	> 99.9 (93.9 to 100.0)	99.6 (99.3 to 99.7)	0.4
(Sequenom Inc., USA)		T13	91.7 (61.0 to 99.0)	99.7 (98.5 to 99.5)	0.3
		combined sex aneuploidies	96.2 (ND)	99.7 (ND)	0.3
MomGuard™	MPSS	T21, T18, T13, 45,X, 47,XXX,	> 99.0 (ND)	ND	ND

Table 2. Reported accuracy of commercially available genomics-based non-invasive prenatal tests^a (Continued)

Korea)		47,XXY, 47,XYY			
NIFTY™ test (Beijing Genomics Institute (BGI), China)	MPSS	T21	99.2 (ND)	100 (ND)	0
		T18	98.2 (ND)	100 (ND)	0
		T13	100 (ND)	100 (ND)	0
		45,X	> 99.9 (ND)	> 99.9 (ND)	< 0.1
Panorama™ prenatal test ^c (Natera, Inc., USA)	SNP TMPS	T21	> 99.9 (ND)	100 (ND)	0
		T18	> 96.4 (ND)	> 99.9 (ND)	< 0.1
		T13	> 99.9 (ND)	100 (ND)	0
		45,X	> 92.9 (ND)	> 99.9 (ND)	< 0.1
PrenaTest® (LifeCodexx AG, Germany)	MPSS	T21	98.7 (ND)	99.9 (ND)	0.1
		T18	100 (ND)		
		T13	100 (ND)		
		45,X	90.9 (ND)	98.8 (ND)	1.2
		47,XYY	100 (ND)		
Prendia (Genesupport, Switzerland)	MPSS	T21	100.0 (88.8 to 100.0)	100.0 (98.0 to 100.0)	0.0
		T18	95.8 (76.8 to 99.7)	100.0 (97.0 to 100.0)	0.0
		T13	100.0 (74.6 to 100.0)	100.0 (98.1 to 100.0)	0.0
		45,X	100.0 (74.6 to 100.0)	100.0 (98.1 to 100.0)	0.0
		47,XXX	100.0 (46.2 to 100.0)	100.0 (98.2 to 100.0)	0.0
Tranquility (Genoma, Switzerland)	MPSS	T21	99.9 (ND)	99.8 (ND)	0.2
		T18	99.9 (ND)	99.9 (ND)	0.1
		T13	99.9 (ND)	99.7 (ND)	0.3
verifi® prenatal test (Illumina, Inc., USA)	MPSS	T21	99.5 (98.7 to 99.5)	99.8 (98.9 to 99.9)	0.2
		T18	97.3 (94.2 to 98.2)	99.7 (99.5 to 99.9)	0.3
		T13	98.0 (95.6 to 98.9)	99.8 (99.8 to 99.9)	0.2
		45,X	95.0 (75.1 to 99.9)	99.0 (97.6 to 99.7)	1.0
VisibiliT™ (Sequenom Inc., USA)	MPSS	T21	> 99.0 (80.8 to 100)	> 99.9 (99.5 to 100)	< 0.1
		T18	> 99.0 (65.5 to 100)	> 99.9 (99.5 to 100)	< 0.1

45,X: Turner syndrome, 47,XXX: triple X syndrome, 47,XXY: Klinefelter syndrome, T21: trisomy 21, T18: trisomy 18, T13: trisomy 13 CI: confidence interval, MPSS: massively parallel shotgun sequencing, ND: no data available, TMPS: targeted massively parallel sequencing and SNP: single nucleotide polymorphism.

^a(Ariosa Diagnostics 2016; BGI 2014; BGI 2016; Berry Genomics 2016; Genoma 2016; Genome Care 2016; Illumina 2014; Illumina 2016; LabGenomics 2016; LifeCodexx 2016; Natera 2016; Genesupport 2016; Premaita Health plc 2016; Sequenom 2016).

^b(Hooks 2014).

^cDNA of maternal and paternal origin are needed.

Table 3. Traditional screening tests (mostly for T21)^a

Screening tests	First trimester (before 14 weeks' gestation)	Second trimester (14 to 20 weeks' gestation)
Ultrasonography	<ul style="list-style-type: none"> • NT measurement 	<ul style="list-style-type: none"> • Various morphologic measurements that modify the prior risk established
Combined test	<ul style="list-style-type: none"> • hCG (free β or total) • PAPP-A • NT measurement 	NA
Triple test	NA	<ul style="list-style-type: none"> • hCG (free β or total) • uE3 • AFP
Quadruple test	NA	<ul style="list-style-type: none"> • hCG (free β or total) • uE3 • AFP • inhibin A
Sequential test ^b	<ul style="list-style-type: none"> • free β hCG • PAPP-A • NT measurement 	<ul style="list-style-type: none"> • Invasive test is offered if 1st trimester result is positive • Quadruple test is offered if 1st trimester result is negative
Contingent test ^b	<ul style="list-style-type: none"> • free β hCG • PAPP-A • NT measurement 	<ul style="list-style-type: none"> • Invasive test is offered if 1st trimester result is positive • Quadruple test is offered after an intermediate 1st trimester result • No test is offered after a low-risk result
Serum integrated test ^c	<ul style="list-style-type: none"> • PAPP-A 	<ul style="list-style-type: none"> • Triple or Quadruple test
Integrated test ^c	<ul style="list-style-type: none"> • PAPP-A • NT measurement 	<ul style="list-style-type: none"> • Quadruple test

Maternal age is often included in the algorithm for prenatal screening tests. AFP: alpha-fetoprotein, hCG: human chorionic gonadotropin, NA: not applicable, NT: nuchal translucency, PAPP-A: pregnancy associated plasma protein A and uE3: unconjugated estriol.

^a(Gekas 2009; Okun 2008; Wald 2005).

^bA test result was available after first-trimester screening test.

^cSingle test result available after second-trimester screening test.

Table 4. Characteristics of included studies by type of gNIPT

Study ID	Target condition(s)	Study design and participants	Prior risk	Index test details	Cutpoint	Reference standard	Comparator
MPSS							
Alberti 2015	T21	<ul style="list-style-type: none"> Case-control study (1:2) from a prospective cohort 976 singleton pregnancies enrolled, 183 were analysed 	High risk	<ul style="list-style-type: none"> Illumina HiSeq 2000 sequencer without multiplexing In-house test FF measured 	Z score of 3	Fetal karyotype ^a	
Benachi 2015	T21, T18, T13	<ul style="list-style-type: none"> Blinded retrospective study 900 singleton or twin pregnancies enrolled, 886 were analysed 	High risk	<ul style="list-style-type: none"> Illumina v3 flow-cell on a HiSeq 1500 sequencer in 12-plex Commercial - Laboratoire CERBA FF measured 	Z score of 3 for T21; 3.95 for T18 and T13	Fetal karyotype or neonatal clinical examination	
Bianchi 2012	T21, T18, T13, 45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> Nested case-control study (1:4) from a prospective cohort (MELISSA) 2882 singleton pregnancies enrolled, 503 for T21, 502 for T18, 501 for T13 and 489 for 45,X were analysed 	High risk	<ul style="list-style-type: none"> Illumina HiSeq 2000 sequencer in 6-plex Commercial test - Verinata FF measured 	Different cut-points used for autosomes and SCA ^b	Fetal karyotype	
Bianchi 2013	T21, T18, T13, 45,X	<ul style="list-style-type: none"> Retrospective study from stored plasma 2882 singleton pregnancies enrolled, 113 were analysed 	High risk	<ul style="list-style-type: none"> Illumina TrueSeq 3.0 sequencing chemistry Commercial test - Verinata 	Different cut-points used for autosomes and SCA ^b	Fetal karyotype	
Bianchi 2014a	T21, T18, T13	<ul style="list-style-type: none"> Blinded prospective cohort study 2052 singleton pregnancies enrolled, 1952 for T21 and T18, and 1914 for T13 were analysed 	High, low and without prior risk	<ul style="list-style-type: none"> Illumina HiSeq 2000 in 8-plex Commercial - verifi[®] prenatal test FF measured 	NCV of 4; resequenced if NCV is between 3 and 4	Fetal or post-natal karyotype, neonatal clinical examination or medical record from birth	Standard screening (T21 only with mixed cut-points) which include first-trimester combined test or a second-trimester result

(quadruple, serum integrated, fully integrated, or sequential).

Table 4. Characteristics of included studies by type of gNIPT *(Continued)*

Bijok 2014	T21, T18, T13	<ul style="list-style-type: none"> Prospective cohort study 10 singleton pregnancies enrolled, 9 were analysed 	High risk	<ul style="list-style-type: none"> Illumina Genome Analyzer Iix or HiSeq 2000 sequencer in multiplex Commercial - NIFTY™ test, BGI-Shenzhen FF measured 	NR	Fetal karyotype
Canick 2012	T21, T18, T13	<ul style="list-style-type: none"> Case-control study 4664 pregnant women enrolled, 27 multifetal pregnancies were analysed 	High risk	<ul style="list-style-type: none"> Illumina HiSeq 2000 sequencer in 4-plex Commercial test - Sequenom, Inc. FF measured 	Z score of 3	Fetal karyotype
Chen 2011	T18, T13	<ul style="list-style-type: none"> Nested case-control study from prospective and retrospective cohorts 392 singleton pregnancies enrolled, 289 were analysed 	High risk	<ul style="list-style-type: none"> Illumina Genome Analyzer Iix in 2-plex Commercial test - Sequenom, Inc. 	Z score of 3	Fetal karyotype
Chiu 2011	T21	<ul style="list-style-type: none"> Blinded case-control study (1:5) from prospective and retrospective cohorts 824 singleton pregnancies enrolled, 753 were analysed by 8-plex method and 314 by 2-plex method 	Mostly high (> 1/300) and some intermediate risk (between 1/300 and 1/1000)	<ul style="list-style-type: none"> Illumina Genome Analyzer II in 8-plex and 2-plex Commercial test - Sequenom, Inc. FF measured 	Z score of 3	Fetal karyotype
Ehrich 2011	T21	<ul style="list-style-type: none"> Blinded case-control study (1:11) from prospective cohort 480 pregnant women enrolled, 449 were analysed 	High risk	<ul style="list-style-type: none"> Illumina Genome Analyzer Iix sequencer in 4-plex Commercial test - Sequenom, Inc. FF measured 	Z score of 2.5	Fetal karyotype

Table 4. Characteristics of included studies by type of gNIPT (Continued)

Fiorentino 2016	T21, T18, T13	<ul style="list-style-type: none"> Blinded prospective cohort study 7103 singleton pregnancies enrolled, 7082 were analysed 	Mostly high risk and without prior risk	<ul style="list-style-type: none"> Illumina HiSeq 2500 sequencer in 15-plex, SAFeR™ algorithm. Commercial - Genoma's prenatal test FF measured 	NCV of 4; aneuploidy suspected if NCV is between 3 and 4	Fetal karyotype or neonatal clinical examination
Hou 2012	T21, T18, T13, 45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> Prospective cohort study 308 singleton pregnancies enrolled, 205 were analysed 	High risk	<ul style="list-style-type: none"> Illumina HiSeq 2000 sequencer Commercial test - BGI-Shenzhen 	NR	Fetal karyotype
Huang 2014	T21, T18	<ul style="list-style-type: none"> Blinded prospective cohort study 189 twin pregnancies enrolled, 189 were analysed 	High risk	<ul style="list-style-type: none"> Illumina Genome Analyzer Ix or HiSeq 2000 sequencer Commercial test - BGI-Shenzhen 	L score of 1 and t score of 2.5 including warning zone	Fetal karyotype
Jeon 2014	T21, T18	<ul style="list-style-type: none"> Prospective cohort study 155 singleton pregnancies enrolled, 155 were analysed 	High risk	<ul style="list-style-type: none"> Ion Torrent PGM or HiSeq 2000 sequencers, 10 samples per Chip Commercial test - Genome Care 	Z score of 2.566 for T21; 2.459 for T18.	Fetal karyotype
Jiang 2012	T21, T18, T13, 45,X, 47,XXY, 47, XYY	<ul style="list-style-type: none"> Prospective cohort study 903 pregnant women enrolled, 903 were analysed 	High risk	<ul style="list-style-type: none"> Illumina Genome Analyzer Ix or HiSeq 2000 sequencer in multiplex Commercial - NIFTY™ test, BGI-Shenzhen FF measured 	Different cut-points used for autosomes and SCA ^b	Fetal karyotype
Johansen 2016	T21, T18, T13	<ul style="list-style-type: none"> Prospective cohort study 375 singleton pregnancies enrolled, 173 were analysed 	High risk	<ul style="list-style-type: none"> Ion Proton™ sequencer in 5-plex In-house test FF measured 	Z score of 4 (unclassified if Z score is between 3 and 4) and WISECONDOR of 1%	Fetal karyotype
Ke 2015	T21, T18, T13	<ul style="list-style-type: none"> Prospective cohort study 2340 singleton pregnancies enrolled, 2340 were analysed 	High risk	<ul style="list-style-type: none"> High throughput sequencing platform Commercial test - BGI-Shenzhen 	T score of 3	Fetal karyotype or newborn outcome

Table 4. Characteristics of included studies by type of gNIPT (Continued)

Kim 2016	T21	<ul style="list-style-type: none"> Blinded prospective cohort study 101 pregnant women enrolled, 101 were analysed 	High risk	<ul style="list-style-type: none"> Ion Proton™ sequencer in multiplex Commercial test - Genome Care 	Z score of 2.10 for Ion Proton™	Fetal karyotype
Lau 2012	T21, T18, T13, 45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> Blinded prospective cohort study 108 singleton pregnancies enrolled, 108 were analysed 	Mostly high risk	<ul style="list-style-type: none"> Illumina HiSeq 2000 sequencers in 12-plex Commercial - NIFTY™ test, BGI-Shenzhen 	Different cut-points used for autosomes and SCA ^b	Fetal karyotype
Lee 2015	T21, T18, T13 and SCA (no case found)	<ul style="list-style-type: none"> Blinded prospective cohort study 93 singleton and multifetal pregnancies enrolled, 92 were analysed 	High risk	<ul style="list-style-type: none"> Illumina MiSeq sequencer in 12-plex or NextSeq sequencer in 96-plex Commercial test - Mom-Guard™, LabGenomics FF measured 	Z score of 4 (intermediate risk if Z score is between 2.5 and 4) for T21 and T18; 2.8 for T13 (intermediate risk if Z score is between 1.9 and 2.8)	Fetal or neonatal karyotype
Lefkowitz 2016	T21, T18, T13, 45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> Retrospective cohort, blinded case-control study 5321 pregnant women enrolled but 1222 were selected and 1166 were analysed 	High risk	<ul style="list-style-type: none"> Illumina HiSeq 2000 sequencer in 6-plex or uniplex Commercial test - Sequenom, Inc. FF measured 	Different cut-points used for autosomes and SCA ^b	Fetal karyotype
Liang 2013	T21, T18, T13, 45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> Blinded prospective cohort study 435 singleton and twin pregnancies enrolled, 412 were analysed 	High risk	<ul style="list-style-type: none"> Illumina HiSeq 2000 sequencer in 8-plex or 12-plex Commercial test - Berry Genomics Co. Ltd. FF measured 	Different cut-points used for autosomes and SCA ^b	Fetal karyotype
Liu 2012	T21, T18, T13, 45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> Prospective cohort study 153 pregnant women enrolled, 153 were analysed 	High risk	<ul style="list-style-type: none"> Illumina HiSeq sequencer in multiplex. 	Z score of 3	Fetal karyotype
Ma 2016	T21, T18, T13	<ul style="list-style-type: none"> Blinded retrospective (archived samples) and prospective cohorts study 	High and low risk	<ul style="list-style-type: none"> Sequencing on BGISEQ-1000 in 16 or 24-plex Commercial test - BGI-Shenzhen 	Z score of 3	Fetal karyotype or post-natal follow-up

Table 4. Characteristics of included studies by type of gNIPT (Continued)

		<ul style="list-style-type: none"> 10,598 singleton pregnancies enrolled, 10,579 were analysed 				
Mazloom 2013	45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> Blinded prospective cohort study 1975 singleton pregnancies enrolled, 411 samples from the validation set were analysed 	High risk	<ul style="list-style-type: none"> Illumina v3 flow-cell on a HiSeq 2000 sequencer in 12-plex Laboratory test development by Sequenom, Inc. FF measured 	Different cut-points used for the four SCA ^b	Fetal karyotype
Palomaki 2012	T21, T18, T13	<ul style="list-style-type: none"> Nested case-control study (1:3) 4664 pregnant women enrolled but 1988 singleton pregnancies were selected and 1971 were analysed 	High risk	<ul style="list-style-type: none"> Illumina HiSeq 2000 sequencer in 4-plex Commercial test - Sequenom, Inc. FF measured 	Z score of 3 for T21; 3.88 for T18; 7.17 for T13	Fetal karyotype
Papa-georghiou 2016a	T21, T18, T13	<ul style="list-style-type: none"> Retrospective cohort, case-control study (1:9) 442 singleton and twin pregnancies enrolled, 426 singleton pregnancies were analysed 	High risk	<ul style="list-style-type: none"> Ion Proton™ sequencer in 8-plex Commercial - IONA® test, Premaitha Health (public limited company in UK) FF measured 	Likelihood ratio of 1 and maternal age-adjusted probability risk score	Fetal karyotype or medical record from birth
Papa-georghiou 2016b	T21, T18, T13	<ul style="list-style-type: none"> Retrospective cohort, case-control study (1:9) 442 singleton and twin pregnancies enrolled, 11 twin pregnancies were analysed 	High risk	<ul style="list-style-type: none"> Ion Proton™ sequencer in 8-plex Commercial - IONA® test, Premaitha Health (public limited company in UK) FF measured 	Likelihood ratio of 1 and maternal age-adjusted probability risk score	Fetal karyotype or medical record from birth
Poon 2016	T21, T18, T13	<ul style="list-style-type: none"> Retrospective cohort, blinded nested case-control study 242 singleton pregnancies enrolled, 241 were analysed 	High risk	<ul style="list-style-type: none"> Ion Proton™ sequencer, IONA® software algorithm Commercial - IONA® test, Premaitha Health (public limited company in UK) FF measured 	NR (authors used the same gNIPT than Papa-georghiou 2016a)	Fetal karyotype
Porreco 2014	T21, T18, T13, 45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> Blinded prospective cohort study 4170 singleton pregnancies enrolled, 3322 for auto- 	High risk	<ul style="list-style-type: none"> Illumina HiSeq 2000 sequencer in 12-plex Commercial test - Sequenom, Inc. 	Different cut-points used for autosomes and SCA ^b	Fetal karyotype or medical record from birth

Table 4. Characteristics of included studies by type of gNIPT *(Continued)*

		<ul style="list-style-type: none"> • some, 3278 for 45,X and 47,XXX and 3201 for 47,XXY and 47,XYY were analysed 		<ul style="list-style-type: none"> • FF measured 			
Sehnert 2011	T21, T18, T13, 45,X	<ul style="list-style-type: none"> • Retrospective (archived samples) cohort study • 1014 singleton and multifetal pregnancies enrolled but only 47 singleton pregnancies in the test set were analysed in this review. 	High risk	<ul style="list-style-type: none"> • Illumina Genome Analyzer Ix sequencer in uniplex • Commercial test - Verinata 	Different cut-points used for autosomes and SCA ^b	Fetal karyotype	
Shaw 2014	T21, T18, T13, 45,X, 47, XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> • Prospective cohort study • 201 singleton and multifetal pregnancies enrolled, 200 were analysed 	High and low risk	<ul style="list-style-type: none"> • Illumina v2 HiSeq 2000 sequencer in 12-plex • Commercial test - Berry Genomics Co. Ltd. 	Different cut-points used for autosomes and SCA ^b	Fetal karyotype or medical record from birth	
Song 2013	T21, T18, T13, 45,X, 47,XXX, 47, XXY, 47,XYY (SCA data not shown in this review)	<ul style="list-style-type: none"> • Blinded prospective cohort study • 1916 singleton pregnancies enrolled, 1741 were analysed 	Without prior risk	<ul style="list-style-type: none"> • Illumina v2 HiSeq2000 in 12-plex • Commercial test- Berry Genomics Co. Ltd. 	Z score of 3	Fetal or post-natal karyotype or medical record from birth	Triple test for T21 and T18 (cutpoint of 1 in 270).
Song 2015	T21, T18, T13, 45,X, 47,XXX, 47,XYY	<ul style="list-style-type: none"> • Blinded prospective cohort study • 213 singleton pregnancies enrolled, 204 were analysed 	High risk	<ul style="list-style-type: none"> • Illumina v2 HiSeq 2000 sequencer in 12-plex • Commercial test - Berry Genomics Co. Ltd. • FF measured 	Z score of 3	Fetal karyotype or neonatal clinical examination or both	
Stumm 2014	T21, T18, T13	<ul style="list-style-type: none"> • Prospective cohort, blinded study for T21 and unblinded for T18 and T13 • 522 singleton pregnancies enrolled, 472 were analysed 	High risk	<ul style="list-style-type: none"> • Illumina HiSeq 2000 sequencer in 12-plex (DAP.21 algorithm without CG correction) • Commercial test - Life-Codexx AG • FF measured 	MAD-based Z score of 3 for T21; 3.2 for T18; 3.9 for T13	Fetal karyotype	
Sukhikh 2015	T21, T18, T13, 45,X	<ul style="list-style-type: none"> • Prospective cohort study • 200 pregnant women enrolled, 200 were analysed 	High risk	<ul style="list-style-type: none"> • Ion Proton™ sequencer • In-house test 	T score of 5 for T21 and T18; 4 for T13; 0.04 Chrom. X and	Fetal karyotype	

Table 4. Characteristics of included studies by type of gNIPT (Continued)

					0.04 Chrom. Y for 45,X	
Sung-Hee 2015	T21, T18, T13, 45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> Retrospective study 918 singleton pregnancies enrolled, 901 were analysed 	High risk	<ul style="list-style-type: none"> Illumina Genome Analyzer Ix or HiSeq 2000 sequencer in 12-plex Commercial - NIFTY™ test, BGI-Shenzhen FF measured 	L score of 1 and t score of 2.5	Fetal karyotype or medical record from birth
Tynan 2016	T21, T18, T13	<ul style="list-style-type: none"> Blinded retrospective cohort study 1100 singleton pregnancies enrolled, 1048 were analysed 	High and without prior risk	<ul style="list-style-type: none"> Illumina HiSeq 2000 or HiSeq 2500 sequencers in multiplex Commercial - VisibiliT™ test, Sequenom, Inc. FF measured 	risk score of 1%	Fetal karyotype or medical record from birth
Wang 2014	T21, T18, T13, 45,X	<ul style="list-style-type: none"> Prospective cohort study 136 singleton pregnancies enrolled, 136 were analysed 	High risk	<ul style="list-style-type: none"> Illumina HiSeq 2000 sequencer Commercial - NIFTY™ test, BGI-Shenzhen 	NR	Fetal or neonatal karyotype or clinical examination at 42 days after birth or both
Wang 2015a	T21, T18, T13, 45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> Prospective cohort study 917 pregnant women enrolled, 917 were analysed 	High risk	<ul style="list-style-type: none"> Illumina v2 HiSeq 2000 flow cell on a HiSeq sequencer Commercial test - Berry Genomics Co. Ltd 	Z score of 3 for T21, T18 and T13; -3 for Chrom. X and 3 for Chrom. Y for sex Chrom. classification.	Fetal karyotype or clinical follow-up to 6 months from birth
Yao 2014	T21, T18, T13 and SCA (SCA data not shown in this review)	<ul style="list-style-type: none"> Retrospective study 5950 singleton pregnancies enrolled, 5530 were analysed 	High, low and without prior risk	<ul style="list-style-type: none"> Illumina Genome Analyzer Ix or HiSeq 2000 sequencer in 12-plex Commercial - NIFTY™ test, BGI-Shenzhen FF measured 	Different cut-points used for autosomes and SCA ^b	Fetal karyotype or clinical follow-up
Zhang 2016	T21, T18, 45,X, 47,XXX (SCA data not shown in this review)	<ul style="list-style-type: none"> Blinded prospective cohort study 87 singleton pregnancies enrolled, 87 were analysed 	High risk	<ul style="list-style-type: none"> Illumina HiSeq 2000 sequencer in 12-plex Commercial test - Berry Genomics Co. Ltd. 	Z score of 3 for T21 (no other cutpoint reported)	Fetal or neonatal karyotype or neonatal clinical

Table 4. Characteristics of included studies by type of gNIPT (Continued)

	shown in this review)					ical examination
Zhou 2014a	T21, T18, T13	<ul style="list-style-type: none"> Blinded prospective cohort study 306 singleton pregnancies enrolled, 301 were analysed 	High, low and without prior risk	<ul style="list-style-type: none"> Illumina Genome Analyzer Ix or HiSeq 2000 sequencer in 12-plex Commercial - NIFTY™ test, BGI-Shenzhen FF measured 	L score of 1 and t score of 2.5	Fetal or neonatal karyotype or birth outcome
Zhou 2014b	T21, T18, T13	<ul style="list-style-type: none"> Blinded prospective cohort study 7705 singleton pregnancies enrolled, 3950 were analysed 	High, low and without prior risk	<ul style="list-style-type: none"> Illumina Genome Analyzer Ix or HiSeq 2000 sequencer in 12-plex Commercial - NIFTY™ test, BGI-Shenzhen FF measured 	L score of 1 and t score of 2.5	Fetal or neonatal karyotype or birth outcome
TMPS						
Ashoor 2012	T21, T18	<ul style="list-style-type: none"> Nested case-control study (1:3) from a prospective cohort 400 singleton pregnancies enrolled, 397 were analysed 	High risk	<ul style="list-style-type: none"> DANSR assay (FORTE algorithm), Illumina HiSeq 2000 in 96-plex Commercial - Harmony™ prenatal test, Ariosa Diagnostics, Inc. 	NR (usually Harmony™ prenatal test uses FORTE risk score of 1%)	Fetal karyotype
Ashoor 2013	T13	<ul style="list-style-type: none"> Blinded prospective cohort study 2167 singleton pregnancies enrolled, 1949 were analysed 	High and low risk	<ul style="list-style-type: none"> DANSR assay (FORTE algorithm), Illumina HiSeq 2000 in 96-plex Commercial - Harmony™ prenatal test, Ariosa Diagnostics, Inc. FF measured 	FORTE risk score of 1%	Fetal karyotype or neonatal clinical examination
Bevilacqua 2015	T21, T18, T13	<ul style="list-style-type: none"> Prospective cohort study 515 multifetal pregnancies enrolled, 340 were analysed Women with singleton pregnancies were excluded (incomplete 2 x 2 table). 	High and without prior risk	<ul style="list-style-type: none"> DANSR assay (FORTE algorithm), Illumina HiSeq 2000 in 96-plex Commercial - Harmony™ prenatal test, Ariosa Diagnostics, Inc. FF measured 	NR (usually Harmony™ prenatal test uses FORTE risk score of 1%)	Fetal or neonatal karyotype

Table 4. Characteristics of included studies by type of gNIPT (Continued)

Comas 2015	T21, T18, T13, 45,X, 47,XXX, 47,XXY, 47,XYY (SCA data not shown in this review)	<ul style="list-style-type: none"> Blinded prospective cohort study 333 singleton pregnancies enrolled, 312 were analysed 	High and without prior risk	<ul style="list-style-type: none"> DANSR assay (FORTE algorithm) or SNP-based method Commercial - Panorama™ test, Natera, Inc. or Harmony™ prenatal test, Ariosa Diagnostics, Inc. FF measured 	Harmony™ prenatal test: NR (usually Harmony™ prenatal test uses FORTE risk score of 1%) Panorama™ test: NR	Fetal karyotype or neonatal clinical examination
del Mar Gil 2014	T21, T18, T13	<ul style="list-style-type: none"> Retrospective cohort study 207 multifetal pregnancies enrolled, 192 twin pregnancies were analysed 	Without prior risk	<ul style="list-style-type: none"> DANSR assay (FORTE algorithm), Illumina HiSeq 2000 in 96-plex Commercial - Harmony™ prenatal test, Ariosa Diagnostics, Inc. FF measured 	NR (usually Harmony™ prenatal test uses FORTE risk score of 1%)	Fetal karyotype
Gil 2016	T21, T18, T13	<ul style="list-style-type: none"> Prospective cohort study 11,692 singleton pregnancies enrolled, 3633 were analysed 	High and intermediate risk ^c	<ul style="list-style-type: none"> DANSR assay (usually with FORTE algorithm) Commercial - Harmony™ prenatal test, Ariosa Diagnostics, Inc. 	NR (usually Harmony™ prenatal test uses FORTE risk score of 1%)	Fetal or postnatal karyotype or neonatal clinical examination
Hall 2014	T13	<ul style="list-style-type: none"> Case-control study (1:3)/1000 singleton pregnancies enrolled, 64 were analysed. 	High risk	<ul style="list-style-type: none"> SNP-based method (NATUS algorithm), Illumina Genome Analyzer Iix or HiSeq sequencer, 11,000 or 19,488-plex targeted PCR Commercial - Natera's prenatal test FF measured 	NR	Fetal karyotype or genetic testing of cord blood, buccal, saliva or products of conception
Hooks 2014	45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> Case-control study from archived samples 432 singleton pregnancies enrolled, 414 were analysed 	High risk	<ul style="list-style-type: none"> DANSR assay (FORTE algorithm), Illumina HiSeq 2000 in 96-plex Commercial - Harmony™ prenatal test, Ariosa Diagnostics, Inc. FF measured 	NR (usually Harmony™ prenatal test uses FORTE risk score of 1%)	Fetal karyotype

Table 4. Characteristics of included studies by type of gNIPT (Continued)

Jackson 2014	T21, T18, T13	<ul style="list-style-type: none"> Prospective cohort study 1228 pregnant women enrolled, 1161 were analysed 	High and low risk	<ul style="list-style-type: none"> DANSR assay (FORTE algorithm) Commercial - Harmony™ prenatal test, Ariosa Diagnostics, Inc. 	NR (usually Harmony™ prenatal test uses FORTE risk score of 1%)	Fetal karyotype or medical record from birth	
Korostelev 2014	T21, T18, T13, 45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> Prospective cohort study 1968 singleton pregnancies enrolled, 685 were analysed 	High and without prior risk	<ul style="list-style-type: none"> SNP-based method (NATUS algorithm), Illumina Genome Analyzer IIx or HiSeq sequencer, > 19,000-plex targeted PCR Commercial - Natera's prenatal test FF measured 	NR	Fetal karyotype or medical record from birth	
Nicolaides 2012	T21, T18	<ul style="list-style-type: none"> Retrospective study from archived plasma 2230 singleton pregnancies enrolled, 1949 were analysed 	Without prior risk	<ul style="list-style-type: none"> DANSR assay (usually with FORTE algorithm) Commercial - Harmony™ prenatal test, Ariosa Diagnostics, Inc. FF measured 	Risk score of 1%	Fetal karyotype or neonatal clinical examination	First-trimester combined test (cutpoint of 1 in 150).
Nicolaides 2013	T21, T18, T13, 45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> Blinded prospective cohort study 242 singleton pregnancies enrolled, 229 were analysed 	High risk	<ul style="list-style-type: none"> SNP-based method (NATUS algorithm), Illumina Genome Analyzer IIx or HiSeq sequencer, 19,488-plex targeted PCR Commercial - Natera's prenatal test FF measured 	NR	Fetal karyotype	
Nicolaides 2014a	45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> Case-control study (archived samples) 177 singleton pregnancies enrolled, 172 were analysed 	High risk	<ul style="list-style-type: none"> DANSR assay (FORTE algorithm), Illumina HiSeq 2000 in 96-plex Commercial - Harmony™ prenatal test FF measured 	FORTE risk score of 1%	Fetal karyotype	
Norton 2012	T21, T18	<ul style="list-style-type: none"> Blinded prospective cohort study 4002 singleton pregnancies enrolled, 3080 were analysed 	High risk	<ul style="list-style-type: none"> DANSR assay (FORTE algorithm), Illumina HiSeq 2000 in 96-plex Commercial test- Ariosa Diagnostics, Inc. 	FORTE risk score of 1%	Fetal karyotype	

Table 4. Characteristics of included studies by type of gNIPT (Continued)

				<ul style="list-style-type: none"> • FF measured 			
Norton 2015	T21, T18, T13	<ul style="list-style-type: none"> • Blinded prospective cohort study • 18,955 singleton pregnancies enrolled, 15,841 were analysed 	Without prior risk	<ul style="list-style-type: none"> • DANSR assay (FORTE algorithm) • Commercial - Harmony™ prenatal test, Ariosa Diagnostics, Inc. • FF measured 	NR (usually Harmony™ prenatal test uses FORTE risk score of 1%)	Fetal or post-natal karyotype, neonatal clinical examination or medical record from birth	First-trimester combined test (cutpoint of 1 in 270 for T21 and 1 in 150 for T18 and T13).
Pergament 2014	T21, T18, T13, 45,X	<ul style="list-style-type: none"> • Blinded prospective cohort study • 1064 singleton pregnancies enrolled, 963 were analysed 	High and low risk	<ul style="list-style-type: none"> • SNP-based method (NATUS algorithm), Illumina Genome Analyzer Iix or HiSeq sequencer, 19,488-plex targeted PCR • Commercial - Natera's prenatal test • FF measured 	NR	Fetal karyotype or genetic testing of cord blood, buccal, saliva or products of conception or birth outcome	
Persico 2016	T21, T18, 45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> • Blinded prospective cohort study • 259 singleton pregnancies enrolled, 249 were analysed 	High risk	<ul style="list-style-type: none"> • SNP-based method (NATUS algorithm), Illumina Genome Analyzer Iix or HiSeq sequencer, 19,488-plex targeted PCR • Commercial - Natera's prenatal test • FF measured 	Risk score of 1%	Fetal karyotype	
Quezada 2015	T21, T18, T13	<ul style="list-style-type: none"> • Prospective cohort study • 2905 singleton pregnancies enrolled, 2785 were analysed 	Without prior risk	<ul style="list-style-type: none"> • DANSR assay (FORTE algorithm) • Commercial - Harmony™ prenatal test • FF measured 	NR (usually Harmony™ prenatal test uses FORTE risk score of 1%)	Fetal or post-natal karyotype, neonatal clinical examination or medical record from birth	First-trimester combined test (cutpoint of 1 in 100 for T21).
Saman-go-Sprouse 2013	45,X, 47,XXX, 47,XXY, 47,XYY	<ul style="list-style-type: none"> • Blinded prospective cohort study • 201 singleton pregnancies (with known SCA and euploid pregnancies) enrolled, 186 were analysed 	High and low risk	<ul style="list-style-type: none"> • SNP-based method (NATUS algorithm), Illumina HiSeq sequencer, 19,488-plex targeted PCR • Commercial - Natera's prenatal test 	NR	Fetal karyotype or genetic testing of cord blood, buccal, saliva	

Table 4. Characteristics of included studies by type of gNIPT (Continued)

Study	Trisomy	Study Design	Risk Level	Methods	FF measured	Products of Conception
Sparks 2012a	T21, T18	Case-control study from a prospective cohort	High risk	DANSR assay (FORTE algorithm), Illumina HiSeq 2000 in 96-plex Commercial test- Ariosa Diagnostics, Inc.	NR	Fetal karyotype
Verweij 2013	T21	Blinded prospective cohort study	High risk	DANSR assay (FORTE algorithm), Illumina HiSeq 2000 in 96-plex Commercial test- Ariosa Diagnostics, Inc.	FORTE risk score of 1%	Fetal karyotype

45,X: Turner syndrome, 47,XXX: triple X syndrome, 47,XXY: Klinefelter syndrome, DANSR: digital analysis of selected regions, FF: fetal fraction DNA, FORTE: fetal-fraction optimised risk of trisomy evaluation, MAD: Median absolute deviation, MPSS: massively parallel shotgun sequencing, NATUS: Next-generation Aneuploidy Test Using SNPs, NCV: normalised chromosome value, SCA: sex chromosome aneuploidy, SNP: single-nucleotide polymorphism, TMPS: targeted massively parallel sequencing, T21: trisomy 21, T18: trisomy 18 and T13: trisomy 13.

^aFetal karyotype include traditional banding techniques, spectral karyotype, fluorescence in situ hybridisation, array comparative genomic hybridisation or quantitative fluorescence polymerase chain reaction.

^bDifferent cutpoints used for autosomes or SCA as follows:

Bianchi 2012: NCV of 4 (aneuploidy suspected if NCV is between 2.5 and 4) for T21, T18, and T13; NCV for Chrom. X of -4 and NCV for Chrom. Y of 2.5 for 45,X; NCV for Chrom. X of 4 and NCV for Chrom. Y of 2.5 for 47,XXX; NCV for Chrom. X between -2.5 and 2.5 and NCV for Chrom. Y > 33 for 47,XXY; NCV for Chrom. X of -4 and NCV for Chrom. Y of 4 for 47,XXY with NCV for Chrom. Y is two times greater than expected NCV Chrom. X.

Bianchi 2013: NCV of 4 (aneuploidy suspected if NCV is between 3 and 4) for T21, T18, and T13; NCV for Chrom. X of -3 and NCV for Chrom. Y of 3 for 45,X.

Jiang 2012: t score of 3 and logarithmic LR of 1 for T21, T18 and T13; if female fetus, t score of -2.5 for 45,X and 47,XXX; t score of 2.5 combined with estimation of fetal ccfDNA concentration by Chrom. X and Y independently for 47,XXY and 47,XXY.

Lau 2012: Z score of 3 for T21, T18 and T13; if female fetus, Z score for Chrom. X of -3 for 45,X; if female fetus, Z score for Chrom. X of 3 for 47,XXX; if male fetus, Z score for Chrom. Y of 3 for 47,XXY.

Lefkowitz 2016: Z score of 3 for T21; Z score of 3.95 for T18 and T13; Z scores for SCA see Mazloom 2013.

Liang 2013: Z score of 3 for T21; 5.91 for T18; 5.72 for T13; ± 2.91 for Chrom. X and ± 3 for Chrom. Y for sex chromosome classification.

Mazloom 2013: Z score of 3.5 for 47,XXX (non-reportable regions between 2.5 and 3.5); Z score of -3.5 for 45,X (non-reportable regions between -2.5 and -3.5); Z score of -3.5 for 47,XXY with Chrom. Y representation; between -3.5 and 3.5 for 47,XXY with Chrom. Y representation.

Porreco 2014: Z score of 3 for T21; Z score of 3,95 for T18 and T13; Z score of 3.5 for 47,XXX (non-reportable regions between 2.5 and 3.5); Z score of -3.5 for 45,X (non-reportable regions between -2.5 and -3.5); Z score of -3.5 for 47,XXY with Chrom. Y representation; Z score between -3.5 and 3.5 for 47,XXY with Chrom. Y representation.

Sehnert 2011: NCV of 4 (unclassified if NCV is between 2.5 and 4) for T21, T18, and T13; NCV for Chrom. Y of -2.0 SDs from the mean of male samples and NCV for Chrom. X of -3.0 SDs from the mean of female samples for sex chromosome classification.

Shaw 2014: Z score of 3 for T21, T18, and T13; Z score of -3 for Chrom. X and 3 for Chrom. Y for sex chromosome classification.

Yao 2014: T score of 2.5 for T21, T18 and T13; if female fetus, T score for Chrom. X of -2.5 for 45,X and 2.5 for 47,XXX; if male fetus, T score for Chrom. X of 2.5 combined with estimation of fetal ccfDNA concentration by Chrom. X (expected value of zero) for 47,XXY; if male fetus, T score for Chrom. X of 2.5 and R-value (the ratio of the fetal DNA fraction estimated by chromosome Y to that estimated by chromosome X) between 1.8 and 2.2 for 47,XXY.

cPregnant women with a first-trimester combined test selected for their risk of fetal aneuploidy (cutpoint of 1 in 100 for high risk and 1 in 101 to 1 in 2500 for intermediate risk).

Table 5. Manufacturers of gNIPT used in the included studies by prior risk of fetal aneuploidy

Company	Number of studies	Number of affected/unaffected pregnancies ^a	Number of studies with pregnant women without prior risk of fetal aneuploidy	Number of studies with high-risk pregnant women	Number of studies with mixed risk ^b cohort
Ariosa Diagnostics, Inc.	15	594/32,302	4	6	5
Beijing Genomics Institute (BGI)	12	427/24,724	0	7	5
Sequenom, Inc.	9	904/8486	0	7	2
Berry Genomics Co. Ltd	6	147/3414	1	4	1
Natera, Inc.	6	276/2103	0	3	3
Illumina, Inc.	4	273/2342	0	3	1
In-house	3	114/442	0	3	0
Premaitha Health plc	3	99/579	0	3	0
Genome Care	2	21/235	0	2	0
CERBA	1	113/745	0	1	0
Genoma	1	105/6977	0	0	1
LabGenomics	1	8/84	0	1	0
LifeCodexx AG	1	55/417	0	1	0
Not reported	1	5/148	0	1	0
Total	65	3141/82,998	5	42	18

^aWe included pregnancies with any other aneuploidy than the one under analysis with all euploid cases as "unaffected" pregnancies.

^bMixed-risk cohort included a mix of pregnant women without prior risk, low risk or high risk of fetal aneuploidy.

Table 6. Reasons for patient exclusion

Study ID	Number of pregnant women enrolled	Reasons for exclusion	Number of women with results for 2 x 2 table analysis
Alberti 2015	976	<ul style="list-style-type: none"> 701 not selected for the case-control study 23 selected for reference set 8 selected for pretesting phase 47 low amount of DNA 11 low fetal fraction DNA or assay failure 3 haemolysed samples Total: 793	183
Ashoor 2012	400	<ul style="list-style-type: none"> 3 samples failed amplification and sequencing 	397
Ashoor 2013	2167	<ul style="list-style-type: none"> 165 selected for first phase (case-control study not included in this review) 53 failed amplification or sequencing Total: 218	1949
Benachi 2015	900	<ul style="list-style-type: none"> 8 without reference standard result 6 low fetal fraction DNA or result appeared atypical Total: 14	886
Bevilacqua 2015	2362	<ul style="list-style-type: none"> 1847 not selected 159 without follow-up 11 failed samples 5 failed samples and were without follow-up Total: 2022	340
Bianchi 2012	2882	<ul style="list-style-type: none"> 127 ineligible 45 without karyotype 85 multifetal pregnancies 2091 not selected for this case-control study 2 for tracking issue 16 without fetal DNA detected Total: 2366 In addition, other samples excluded from 2 x 2 tables for censored complex karyotype: <ul style="list-style-type: none"> 13 for T21 14 for T18 15 for T13 27 for 45,X 	503 (T21) 502 (T18) 501 (T13) 489 (45,X)
Bianchi 2013	2882	<ul style="list-style-type: none"> 2769 not selected for the study 	113
Bianchi 2014a	2052	<ul style="list-style-type: none"> 10 failed blood quality control 72 without clinical outcome 17 without gNIPT result 28 without standard screening result 	1952 (T21 and T18) 1914 (T13)

Table 6. Reasons for patient exclusion (Continued)

		<ul style="list-style-type: none"> • 1 without gNIPT result and without standard screening result 	
		Total for T21 and T18: 100	
		Total for T13: 128	
Bijok 2014	10	<ul style="list-style-type: none"> • 1 low fetal fraction DNA 	9
Canick 2012	4664	<ul style="list-style-type: none"> • 4637 not selected for the case-control study 	27
Chen 2011	392	<ul style="list-style-type: none"> • 103 selected for reference control 	289
Chiu 2011	824	<ul style="list-style-type: none"> • 46 failed quality control for blood sampling • 12 without karyotype • 2 twin pregnancies • 11 failed quality control for sequencing 	753 (8-plex)
		Total: 71 (8-plex)	
Comas 2015	333	<ul style="list-style-type: none"> • 17 without follow-up • 3 unrepeated tests • 1 failed test second time^a and without follow-up 	312
		Total: 21	
del Mar Gil 2014	207	<ul style="list-style-type: none"> • 11 low fetal fraction DNA • 4 laboratory processing failures 	192
		Total: 15	
Ehrich 2011	480	<ul style="list-style-type: none"> • 13 preanalytic failure (including 9 for low plasma volume and 4 processing errors) • 18 failed quality control at second time (including 7 for low fetal fraction DNA) 	449
		Total: 31	
Fiorentino 2016	7103	<ul style="list-style-type: none"> • 21 failed quality control (unrepeated tests) 	7082
Gil 2016	11,692	<ul style="list-style-type: none"> • 7994 patients did not undergo a gNIPT • 45 failed tests first time^b • 20 failed tests second time 	3633
		Total: 8059	
Hall 2014	> 1000	<ul style="list-style-type: none"> • About 932 samples not selected for the case-control study • 4 failed quality control 	64
		Total: 936	
Hooks 2014	432	<ul style="list-style-type: none"> • 18 low fetal fraction DNA, unusually high variation in ccfDNA counts or failed QC 	414
Hou 2012	308	<ul style="list-style-type: none"> • 103 patients did not undergo a gNIPT 	205
Huang 2014	189	NR	189
Jackson 2014	1228	<ul style="list-style-type: none"> • 7 with other abnormal ultrasound • 14 opted for CVS without gNIPT 	1161

Table 6. Reasons for patient exclusion (Continued)

		<ul style="list-style-type: none"> • 32 declined all testing • 14 failed tests twice 	
		Total: 67	
Jeon 2014	155	NR	155
Jiang 2012	903	NR	903
Johansen 2016	375	<ul style="list-style-type: none"> • 191 not selected for validation set • 11 low fetal fraction DNA 	173
		Total: 202	
Ke 2015	2340	NR	2340
Kim 2016	101	NR	101
Korostelev 2014	1968	<ul style="list-style-type: none"> • 1043 without follow-up • 240 samples did not undergo a gNIPT 	685
		Total: 1283	
Lau 2012	108	NR	108
Lee 2015	93	<ul style="list-style-type: none"> • 1 low fetal fraction DNA 	92
Lefkowitz 2016	5321	<ul style="list-style-type: none"> • 4099 not selected for the study • 11 for incomplete follow-up • 3 with confirmed mosaicism • 11 low fetal fraction DNA • 29 for technical reasons • 2 for maternal event 	1166 (autosomes) 1144 (SCA)
		Total: 4155 (autosomes)	
		In addition:	
		<ul style="list-style-type: none"> • 22 sequencing failures for SCA 	
		Total: 4177 (SCA)	
Liang 2013	435	<ul style="list-style-type: none"> • 11 without karyotype • 12 failed quality control 	412
		Total: 23	
Liu 2012	153	NR	153
Ma 2016	10,598	<ul style="list-style-type: none"> • 14 with incomplete follow-up • 5 failed quality control 	10,579
		Total: 19	
Mazloom 2013	1975	<ul style="list-style-type: none"> • 1564 selected for the training set 	411
Nicolaides 2012	2230	<ul style="list-style-type: none"> • 181 ineligible • 46 low fetal fraction DNA • 54 assay failures 	1949

Table 6. Reasons for patient exclusion (Continued)

		Total: 281	
Nicolaides 2013	242	<ul style="list-style-type: none"> 13 failed quality control 	229
Nicolaides 2014a	177	<ul style="list-style-type: none"> 1 failed quality control 4 low fetal fraction DNA 	172
		Total: 5	
Norton 2012	4002	<ul style="list-style-type: none"> 774 ineligible 57 low fetal fraction DNA 91 assay failures 	3080
		Total: 922	
Norton 2015	18,955	<ul style="list-style-type: none"> 381 ineligible 64 withdrawn 384 handling errors 308 without standard screening test result 1489 without follow-up 192 low fetal fraction DNA 83 no fetal fraction DNA 213 high assay variance or assay failures 	15,841
		Total: 3114	
Palomaki 2012	4876	<ul style="list-style-type: none"> 2888 not selected for this study 17 failed tests second time (mostly for low fetal fraction DNA) 	1971
		Total: 2905	
Papageorghiou 2016a	442	<ul style="list-style-type: none"> 11 twin not selected 3 low fetal fraction DNA 2 failed quality control 	426
		Total: 16	
Papageorghiou 2016b	442	<ul style="list-style-type: none"> 426 singleton not selected 3 low fetal fraction 2 failed quality control 	11
		Total: 431	
Pergament 2014	1064	<ul style="list-style-type: none"> 13 not selected (other aneuploidies) 85 samples failed quality control for all five chromosomes (including 65 for low fetal fraction DNA) 	963 (T21) 964 (T18 and 45,X) 965 (T13)
		Total: 98	
		In addition,	
		<ul style="list-style-type: none"> 3 samples failed only for T21 (total for T21: 101) 2 samples failed only for T18 and 45,X (total for T18 and 45,X: 100) 1 sample failed only for T13 (total for T13: 99) 	
Persico 2016	259	<ul style="list-style-type: none"> 8 low fetal fraction DNA 2 failed internal quality control 	249

Table 6. Reasons for patient exclusion (Continued)

			Total: 10
Poon 2016	242	<ul style="list-style-type: none"> 1 low fetal fraction DNA 	241
Porreco 2014	4170	<ul style="list-style-type: none"> 320 for insufficient sample volume 390 failed quality control 24 with incomplete follow-up 6 without invasive procedure <p>In addition,</p> <ul style="list-style-type: none"> 54 failed quality control and 54 for complex autosome karyotypes^c (total: 108 for autosomes) 102 failed quality control or other^d and 50 for complex SCA karyotype (total: 152 for 45,X and 47,XXX) 182 low fetal fraction DNA or other^d and 47 for complex SCA karyotype (total: 229 for 47,XXY and 47,XYY) 	3322 (T21, T18, T13) 3278 (45,X, 47,XXX) 3201 (47,XXY, 47,XYY)
Quezada 2015	2905	<ul style="list-style-type: none"> 66 without follow-up 1 lost in mail 38 low fetal fraction DNA 15 assay failures <p>Total: 120</p>	2785
Samango-Sprouse 2013	201	<ul style="list-style-type: none"> 12 low fetal fraction DNA or poor DNA quality 2 without gNIPT result 1 with conflicting algorithm metrics <p>Total: 15</p>	186
Sehnert 2011	1014	<ul style="list-style-type: none"> 895 not selected for sequencing 71 selected for training set 1 twin pregnancy <p>Total: 967</p>	47
Shaw 2014	201	<ul style="list-style-type: none"> 1 for early GA 	200
Song 2013	1916	<ul style="list-style-type: none"> 102 without follow-up 64 failed quality control 9 failed quality control and without follow-up <p>Total: 175</p>	1741
Song 2015	213	<ul style="list-style-type: none"> 8 without follow-up 1 failed quality control <p>Total: 9</p>	204
Sparks 2012a	338	<ul style="list-style-type: none"> 171 selected for training set 	167
Stumm 2014	522	<ul style="list-style-type: none"> 8 without reference standard 9 without consent 1 previously analysed 14 failed sequencing quality control 18 failed libraries 	472

Table 6. Reasons for patient exclusion (Continued)

Total: 50			
Sukhikh 2015	200	NR	200
Sung-Hee 2015	918	<ul style="list-style-type: none"> • 8 ineligible • 9 without follow-up 	901
Total: 17			
Tynan 2016	1100	<ul style="list-style-type: none"> • 28 library preparation failures or failed quality control • 24 for discretionary non reporting 	1048
Total: 52			
Verweij 2013	595	<ul style="list-style-type: none"> • 75 ineligible • 7 low fetal fraction DNA • 9 laboratory processing failures or specimen issues 	504
Total: 91			
Wang 2014	136	NR	136
Wang 2015a	917	NR	917
Yao 2014	5950	<ul style="list-style-type: none"> • 420 without follow-up 	5530
Zhang 2016	87	NR	87
Zhou 2014a	306	<ul style="list-style-type: none"> • 5 without follow-up 	301
Zhou 2014b	7705	<ul style="list-style-type: none"> • 4 low fetal fraction DNA • 3751 without follow-up 	3950
Total: 3755			

ccfDNA: circulating cell-free DNA, CVS: chorionic villi sampling, GA: gestational age, gNIPT: genomics-based non-invasive prenatal testing, NR: not reported by authors.

^aSecond time: sample failed the second gNIPT assay.

^bFirst time: sample failed the initial gNIPT assay.

^cComplex autosome karyotypes are mosaic, triploidies, unbalanced rearrangements with missing or duplicated genetic material.

^dOther are copy number variation of the X chromosome is confounded by maternal component and cannot be determined.

Table 7. Proportion of pregnant women with a reference standard and assay failure during gNIPT process

Study ID	Failure rate at first attempt (%)	Repeated tests ^a (%)	Failure rate of repeated tests (%)	Final failure rate total (%)	Aneuploid ^b samples (%)	Euploid ^b samples (%)
MPSS						
Alberti 2015	61/244 (25%)	0	NA	61/244 (25%)	NR	NR

Table 7. Proportion of pregnant women with a reference standard and assay failure during gNIPT process (Continued)

Benachi 2015	42/892 (4.7%)	42 (100%) with second aliquot	6/42 (14.3%)	6/892 (0.7%)	2.7%	0.4%
Bianchi 2012	16/519 (3.1%)	0	NA	16/356 (3.1%)	NR	NR
Bianchi 2014a	18/1970 (0.9%)	0 ^c	NA	T21 and T18: 18/1970 (0.9%) T13: 18/1932 (0.9%)	NR	NR
Bijok 2014	1/10 (10.0%)	0	NA	1/10 (10.0%)	50%	0%
Chiu 2011	11/764 (1.4%)	0	NA	11/764 (1.4%)	NR	NR
Ehrich 2011	20/467 (4.3%)	20 (100%) re-sequenced	18/20 (90%)	18/467 (3.9%)	NR	NR
Fiorentino 2016	100/7103 (1.4%)	79 (79%) with new sampling	0 (0%)	21/7103 (0.3%)	0%	0.3%
Johansen 2016	NR	2 with second aliquot or resequenced were in the grey zone (between affected and unaffected)	NR	11/184 (6%) ^d	5.8%	6.1%
Lee 2015	1/93 (1.1%)	0	NA	1/93 (1.1%)	NR	NR
Lefkowitz 2016	Autosomes: 42/1208 (3.5%) SCA: 64/1208 (5.3%)	0	NA	Autosomes: 42/1208 (3.5%) SCA: 64/1208 (5.3%)	Autosomes: 3.8% SCA: 29.7%	Autosomes: 3.4% SCA: 4.5%
Liang 2013	12/424 (2.8%)	0	NA	12/424 (2.8%)	NR	NR
Ma 2016	5/10,584 (0.05%)	0	NA	5/10,584 (0.05%)	NR	NR
Mazloom 2013	21/432 (4.9%)	0	NA	21/432 (4.9%)	11.8%	4.3%
Palomaki 2012	110/1988 (5.5%)	105 (95.5%) with second aliquot and 5 (4.5%) resequenced	17/110 (15.5%)	17/1988 (0.9%)	1.0%	0.8%
Papageorghiou 2016a	5/431 (1.2%)	0	NA	5/431 (1.2%)	NR	NR

Table 7. Proportion of pregnant women with a reference standard and assay failure during gNIPT process (Continued)

 Papageorgiou
 2016b

Poon 2016	1/242 (0.4%)	0	NA	1/242 (0.4%)	0%	0.5%
Porreco 2014	Autosomes: 108/3430 (3.1%) 45,X and 47,XXX: 152/3430 (4.4%) 47,XXY and 47,XYY: 229/3430 (6.7%)	0	NA	Autosomes: 108/3430 (3.1%) 45,X and 47,XXX: 152/3430 (4.4%) 47,XXY and 47,XYY: 229/3430 (6.7%)	NR	NR
Song 2013	73/1814 (4.0%)	0	NA	73/1814 (4.0%)	0%	4.0%
Song 2015	1/205 (0.5%)	0	NA	1/205 (0.5%)	NR	NR
Stumm 2014	32/504 (6.3%)	0	NA	32/504 (6.3%)	3.5%	6.7%
Sung-Hee 2015	21/908 (2.3%)	16 (76.2%) with new sampling	2/16 (12.5%)	7/908 (0.8%)	NR	NR
Tynan 2016	52/1100 (4.7%)	0	NA	52/1100 (4.7%)	0%	4.9%
Yao 2014	0	0	NA	0	NA	NA
Zhou 2014a	0	0	NA	0	NA	NA
Zhou 2014b	141/3954 (3.6%)	141 (100%) with new sampling	4/141 (2.8%)	4/3954 (0.1%)	NR	NR
Overall range of final assay failure for MPSS				0% to 25%	0% to 50%	0% to 6.7%
TMPS						
Ashoor 2012	3/400 (0.8%)	0	NA	3/400 (0.8%)	0%	1%
Ashoor 2013	53/2002 (2.6%)	0	NA	53/2002 (2.6%)	0%	2.7%
Bevilacqua 2015	29/356 (8.1%)	26 (90%) with 2nd aliquot	13/26 (50%)	16/356 (4.5%)	NR	NR
Comas 2015	9/316 (2.8%)	6 (67%) with new sampling	1/6 (16.7%)	4/316 (1.3%)	NR	NR
del Mar Gil 2014	15/207 (7.2%)	0	NA	15/207 (7.2%)	23%	6%

Table 7. Proportion of pregnant women with a reference standard and assay failure during gNIPT process (Continued)

Gil 2016	99/3698 (2.8%)	54 (54.5%) with new sampling	20/54 (37%)	65/3698 (1.8%)	NR	NR
Hall 2014	4/68 (5.9%)	0	NA	4/68 (5.9%)	11.8%	3.9%
Hooks 2014	18/432 (4.2%)	0	NA	18/432 (4.2%)	NR	NR
Jackson 2014	NR	NR	14 (NR)	14/1175 (1.2%)	NR	NR
Nicolaides 2012	100/2049 (4.9%)	0	NA	100/2049 (4.9%)	9.1%	4.9%
Nicolaides 2013	13/242 (5.4%)	0	NA	13/242 (5.4%)	6.3%	5.2%
Nicolaides 2014a	5/177 (2.8%)	0	NA	5/177 (2.8%)	5.1%	1.7%
Norton 2012	148/3228 (4.6%)	0	NA	148/3228 (4.6%)	NR	NR
Norton 2015	488/16,329 (3.0%)	0	NA	488/16,329 (3.0%)	20.6%	2.9%
Pergament 2014	T21: 88/1051 (8.4%) T18, 45,X: 87/1052 (8.3%) T13: 86/1053 (8.2%)	0	NA	T21: 88/1051 (8.4%) T18, 45,X: 87/1052 (8.3%) T13: 86/1053 (8.2%)	All five chromo- somes (n = 85): 15.2%	All five chromo- somes (n = 85): 7.1%
Persico 2016	10/259 (3.9%)	0	NA	10/259 (3.9%)	8.4%	2.1%
Quezada 2015	122 ^e /2838 (4.2%)	110 (90.1%) with new sampling	41/110 (37.3%)	53/2838 (1.9%)	4.1%	1.8%
Saman-go-Sprouse 2013	15/201 (7.5%)	0	NA	15/201 (7.5%)	6.3%	7.6%
Verweij 2013	51/520 (9.8%)	51 (100%) with 2 nd aliquot	16/51 (31.4%)	16/520 (3.1%) NR	NR	NR
Overall range of final assay failure for TMPS				0.8% to 7.5%	0% to 23%	1% to 7.63%

CVS: chorionic villi sampling, FF: fetal fraction DNA, GA: gestational age, NA: not applicable, NR: not reported by authors, QC: quality control.

^aRepeated tests included second aliquot (aliquot from first sampling), resequenced (same library) or new sampling.

^baneuploid: proportion of failed samples of aneuploid cases out of all aneuploid tested with reference standard and gNIPT result. euploid: proportion of failed samples of euploid cases out of all euploid tested with reference standard and gNIPT result.

^cAuthors decided to resequence 12 samples with gNIPT results. They were in the grey zone (between affected and unaffected) and were resequenced in uniplex. All repeated tests were in affected or unaffected zone.

^dOnly the final failure rate was reported. The failure rate at first attempt was not reported nor the failure rate of repeated tests.

^eAuthor reported 123 failed tests but this number included one sample lost in the mail and so did not undergo the sequencing process.

Table 8. Data for 47,XXX, 47,XXY and 47,XYY according to the prior risk of fetal aneuploidy and gNIPT approach

Test		Number of studies	Number of affected pregnancies	Number of unaffected pregnancies ^a
47,XXX				
Selected high risk pregnant women	MPSS	5	8	5441
	TMPS	2	6	580
47,XXY				
Selected high risk pregnant women	MPSS	7	14	6466
	TMPS	3	8	827
47,XYY				
Selected high risk pregnant women	MPSS	7	11	6418
	TMPS	1	3	169

^aUnaffected pregnancies: we included pregnancies with any other aneuploidy than the one under analysis with all euploid cases as "unaffected".

Table 9. Subgroup analyses of MPSS and TMPS (type of pregnancy and gestational age)

Test subgroups		Number of studies	Number of affected pregnancies	Number of unaffected pregnancies ^a	Sensitivity ^b % (95% CI)	Specificity ^b % (95% CI)
Pregnancy type						
Autosomes (T21, T18 and T13 combined), unselected population						
MPSS	singleton	1	11	1730	100 (74.1 to 100)	99.9 (99.7 to 100)
TMPS	singleton	3	107	20,468	95.5 (87.4 to 98.4)	99.9 (99.8 to 100)
	multifetal	1	11	181	90.9 (62.3 to 98.4)	100 (97.9 to 100)
Autosomes (T21, T18 and T13 combined), selected high-risk population						
MPSS	singleton	19	1087	11,180	98.3 (97.3 to 98.9)	99.6 (99.5 to 99.7)
	multifetal	3	21	206	95.2 (72.9 to 99.3)	100 (98.2 to 100) ^c
TMPS	singleton	7	378	4282	98.9 (97.2 to 99.6)	99.9 (99.8 to 100)
SCA (45,X, 47,XXX, 47,XXY and 47,XYY combined), selected high-risk population						
MPSS	singleton	7	101	4690	88.3 (52.9 to 98.1)	99.3 (97.5 to 99.8)

Table 9. Subgroup analyses of MPSS and TMPS (type of pregnancy and gestational age) (Continued)

TMPS		4	96	968	93.8 (86.8 to 97.2)	99.6 (98.1 to 99.9)
Gestational age						
Autosomes (T21, T18 and T13 combined), unselected population						
MPSS	≤29 weeks	1	11	1730	100 (74.1 to 100)	99.9 (99.7 to 100)
TMPS	≤15 weeks	4	118	20,649	94.9 (89.1 to 97.7)	99.9 (99.8 to 99.9)
Autosomes (T21, T18 and T13 combined), selected high-risk population						
MPSS	≤15 weeks	3	49	532	100 (92.7 to 100) ^c	100 (99.3 to 100) ^c
	≤29 weeks	12	594	4605	98.3 (96.9 to 99.1)	99.3 (99.0 to 99.5)
	≤42 weeks	13	729	7831	98.9 (95.0 to 99.8)	99.9 (99.8 to 99.9)
TMPS	≤15 weeks	2	128	498	99.2 (95.7 to 99.9) ^c	100 (99.2 to 100) ^c
	≤29 weeks	2	33	535	97.0 (84.7 to 99.5) ^c	100 (99.3 to 100) ^c
	≤42 weeks	2	163	3084	99.4 (95.8 to 99.9)	99.9 (99.7 to 100)
SCA (45,X, 47,XXX, 47,XXY and 47,XYY combined), selected high-risk population						
MPSS	≤15 weeks	1	2	202	0.00 (0.00 to 65.8)	99.5 (97.2 to 99.9)
	≤29 weeks	5	58	996	86.5 (63.1 to 96.0)	95.1 (93.5 to 96.3)
	≤42 weeks	5	89	6103	95.8 (80.3 to 99.2)	99.6 (99.4 to 99.7)
TMPS	≤15 weeks	2	58	343	93.1 (83.0 to 97.4)	99.7 (98.0 to 100)
	≤42 weeks	1	34	380	97.1 (85.1 to 99.5)	98.9 (97.3 to 99.6)

45,X: Turner syndrome, 47,XXX: triple X syndrome, 47,XXY: Klinefelter syndrome, T21: trisomy 21, T18: trisomy 18, T13: trisomy 13 CI: confidence interval, MPSS: massively parallel shotgun sequencing, SCA: sex chromosome aneuploidies, TMPS: targeted massively parallel sequencing.

^aWe included pregnancies with any other aneuploidy than the one under analysis with all euploid cases as "unaffected" pregnancies.

^bFor two or more studies, the sensitivities and specificities are the summary estimates obtained from meta-analysis.

^cSimple pooling used to obtain summary estimates of sensitivity, specificity or both.

Table 10. Direct comparisons of gNIPT and traditional screening tests for autosomes (T21, T18 and T13 combined) in unselected population of pregnant women undergoing aneuploidy screening

Study	Sensitivity (true positives/cases)		Difference		Specificity (true negatives/unaffected ^a)		Difference	
	%		% (95% CI)		%		% (95% CI)	
	MPSS	Traditional screening tests			MPSS	Traditional screening tests		

Table 10. Direct comparisons of gNIPT and traditional screening tests for autosomes (T21, T18 and T13 combined) in unselected population of pregnant women undergoing aneuploidy screening (Continued)

	TMPS	Traditional screening tests		TMPS	Traditional screening tests	
Song 2013	100 (11/11)	54.6 (6/11)	45.5 (10.0 to 72.0)	99.9 (1729/1730)	86.0 (1487/1730)	14.0 (12.4 to 15.7)
Nicolaides 2012	100 (10/10)	100 (10/10)	0.00 (-27.8 to 27.8)	99.9 (1937/1939)	95.5 (1852/1939)	4.38 (3.51 to 5.40)
Norton 2015	98.0 (49/50)	78.0 (39/50)	20.0 (7.44 to 33.3)	99.9 (15,779/15,791)	94.1 (14,860/15,791)	5.82 (5.46 to 6.20)
Quezada 2015	91.5 (43/47)	100 (49/49)	-8.51 (-19.9 to 0.40)	99.7 (2730/2738)	95.6 (2663/2787)	4.16 (3.40 to 5.00)

CI: confidence interval, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing.

^aWe included pregnancies with any other aneuploidy than the one under analysis with all euploid cases as "unaffected" pregnancies.

Table 11. Sensitivity analyses

Test	Number of studies	Number of affected pregnancies	Number of unaffected pregnancies ^a	Summary sensitivity % (95% CI)	Summary specificity % (95% CI)	P value ^b
Case-control studies excluded						
Autosomes (T21, T18 and T13 combined), selected high-risk population						
MPSS	22	696	11,293	98.3 (95.1 to 99.4)	99.9 (99.8 to 100)	0.72
TMPS	4	219	3,813	98.6 (95.8 to 99.6)	99.9 (99.8 to 100)	
SCA (45,X, 47,XXX, 47,XXY and 47,XYY combined), selected high-risk population						
MPSS	10	98	5,872	91.9 (73.8 to 97.9)	99.5 (98.8 to 99.8)	0.41
TMPS	2	6	472	93.8 (86.8 to 97.2)	99.6 (98.1 to 99.9)	
Exclusion of studies with less than 10 pregnancies with aneuploidy						
Autosomes (T21, T18 and T13 combined), selected high-risk population						
MPSS	21	1458	13,921	98.7 (96.8 to 99.4)	99.8 (99.5 to 100)	0.07
TMPS	7	378	4,282	98.9 (97.2 to 99.6)	99.9 (99.8 to 100)	
SCA (45,X, 47,XXX, 47,XXY and 47,XYY combined), selected high-risk population						
MPSS	6	130	5,761	94.5 (80.6 to 98.6)	99.4 (97.6 to 99.8)	0.28
TMPS	2	90	496	94.4 (87.3 to 97.7)	99.0 (97.6 to 99.6)	

45,X: Turner syndrome, 47,XXX: triple X syndrome, 47,XXY: Klinefelter syndrome, T21: trisomy 21, T18: trisomy 18, T13: trisomy 13 CI: confidence interval, MPSS: massively parallel shotgun sequencing, SCA: sex chromosome aneuploidies, TMPS: targeted massively parallel sequencing.

^aWe included pregnancies with any other aneuploidy than the one under analysis with all euploid cases as "unaffected" pregnancies.

^bThe P value indicates the statistical significance of the difference in model fit and was obtained from likelihood ratio tests comparing models with and without a covariate for test type.

APPENDICES

Appendix 1. Screening tests and medical terms glossary^a

Terms	Definitions
Amniocentesis	Invasive procedure under continuous ultrasound guidance (performed between 15 to 19 weeks of gestational age). A sterile needle is passed through the mother's abdomen, uterus and amniotic sac. A sample of fetal cells present in the amniotic fluid surrounding the fetus is aspirated with a syringe and sent for analysis to test for a range of chromosomal and inherited disorders.
Aneuploidy	The state of having a different (additional or missing) number of chromosomes than the 23 pairs normally present in humans.
Attention deficit disorder (ADD)	ADD is a neurodevelopmental disorder defined by impairing levels of inattention and disorganisation. Inattention manifests behaviourally in ADD as wandering off task, lacking persistence, having difficulty sustaining focus and being disorganised.
Case-control study	In the context of diagnostic accuracy, existing records are used to identify a group of people known to have the target condition (cases) and another group (controls) without the target condition. The control group may consist of healthy individuals or those with other conditions similar to the target condition. Cases and controls are then compared with respect to certain variables hypothesised to increase the risk of having the disease.
Chorionic villus sampling (CVS)	An abdominal or cervical procedure performed under continuous ultrasound guidance to obtain a sample of placental tissue for chromosomal or genetic analysis (between 12 to 19 weeks of gestational age). The range of chromosomal and genetic conditions that can be detected is similar to those for amniocentesis.
Clinodactyly	Permanent deflection of one or more fingers.
Cut-off	Synonyms: cutpoint or threshold.
Cutpoint	A value for a test result measured on an ordinal or continuous scale which divides the group of people tested into a group at lower risk of the condition being screened for and a group at higher risk (for whom further investigations may be offered). Synonyms: cut-off or threshold.
Detection rate	The proportion of affected individuals with a positive screening result. The detection rate is the same as the sensitivity of a test.
Developmental delay	An individual with this neurodevelopmental disorder fails to meet expected developmental milestones in several areas of intellectual functioning.
Diagnostic accuracy	The ratio of true positive and true negative results to the total number of test results (true positives, true negatives, false positives and false negatives). Represents the level of agreement between the information from the index test and the reference standard.

(Continued)

Diagnostic test	A test recognised as having best performances to provide sufficient information allowing a definite diagnosis (as opposed to screening test results that need to be confirmed before a final diagnosis can be reached).
Dysgenesis	Defective or abnormal formation of an organ or part, primarily during embryonic development. Gonadal dysgenesis is a defective development of the gonads, which may be accompanied by abnormalities of the sex chromosomes.
False negative	A negative test result in someone with the target condition.
False positive	A positive test result in someone without the target condition.
Fluorescence in situ Hybridisation	Describes a type of DNA analysis by the hybridisation of fluorescently-labelled probes complementary to certain genomic regions. In the context of fetal aneuploidy detection, describes a diagnostic test in which chromosome-specific fluorescently-labelled DNA probes are used on uncultured cells from chorionic villi or amniotic fluid to assess the number of homologous chromosome copies present.
High risk of fetal aneuploidy	A pregnancy is considered at high risk of fetal aneuploidy if the result of the prenatal screening test puts the fetus at increased risk for aneuploidy.
Hypertelorism	Abnormal distance between two paired organs.
Intellectual disability or intellectual developmental disorder	This neurodevelopmental disorder is characterised by deficits in general mental abilities, such as reasoning, problem solving, planning, abstract thinking, judgment, academic learning and learning from experience. It is a condition diagnosed before age 18. In the past, the term mental retardation was used to describe this condition but this term is no longer used.
Invasive procedure	A method used to obtain a biological sample that involves significant disruption of the physical integrity of a patient. Examples include amniocentesis and chorionic villi sampling.
Karyotype	A photomicrograph of an individual's chromosomes arranged in a standard format and visualised by various staining methods, showing the number, size, and shape of each chromosome; used to correlate chromosomal anomalies with specific diseases (karyotyping). In humans, there are a total of 23 pairs of homologous chromosomes (total of 46 chromosomes).
Learning disability	Learning disability refers to inadequate development of specific academic, language and speech skills such as reading disability, mathematics disability and writing disability.
Low risk of fetal aneuploidy	A pregnancy is considered at low risk of fetal aneuploidy if the result of the prenatal screening test puts the fetus at decreased risk for aneuploidy.
Meta-analysis	The use of statistical techniques in a systematic review to integrate the results of included studies. Sometimes misused as a synonym for systematic reviews which may or may not include a meta-analysis.
Mosaic	An individual who has some cells with an unusual genetic or chromosomal make-up while the rest of the cells in the body have the typical genetic or chromosomal constitution.
Mixed risk population	Mixed risk population included a mixture of selected pregnant women with low, high or no prior risk of fetal aneuploidy.
Negative predictive value	A measure of test performance. Defined as the proportion of people with a negative test result who do not have the target condition.

(Continued)

Nuchal translucency scan	The thickness of fluid in the tissue space within the nape of the fetal neck typically measured by ultrasonography. An increased amount of fluid is associated with Down syndrome and other structural or genetic anomalies.
Positive predictive value	A measure of test performance. Defined as the proportion of people with a positive test result who do have the target condition.
Probability	The chance or risk of an event happening.
Prospective study	A study in which a group of individuals is followed through time in order to detect the occurrence of a disease or another outcome of interest.
Reference standard	The best available test to detect the presence or absence of the target condition.
Retrospective study	A study in which all or part of the data collection occurred before initiation of the study.
Screening	Testing asymptomatic people for the likelihood of the presence of a disease, either with the aim of reducing risk of an adverse outcome, or with the aim of giving information about risk.
Seizure	A sudden attack, spasm, or convulsion caused by abnormal electrical conduction in the brain.
Sensitivity	A measure of test performance. Defined as the proportion of individuals with the target condition who have a positive test result. Higher sensitivity values means that a higher proportion of affected individuals will be detected by the test (few false negatives). Sensitivity is the same as the detection rate.
Single nucleotide polymorphism	Single nucleotide polymorphisms are the most common type of genetic variation among people. A difference in a single DNA nucleotide (A, T, C or G) in a DNA sequence.
Specificity	A measure of test performance. Defined as the proportion of individuals without the target condition who have a negative test result. Higher specificity values means that a smaller proportion of unaffected individuals will be wrongly classified as having the target condition (few false positives).
Threshold	Synonyms: cutpoint or cut-off.
True negative	An individual with a negative test result who does not have the target condition.
True positive	An individual with a positive test result who has the target condition.
Trisomy	Three copies of a particular chromosome rather than the usual pair.
Unselected pregnant women	A pregnant women who did not undergo any prenatal screening test at the time of enrolment.

^aAdapted in part from the United Kingdom National Screening Committee Glossary, MedlinePlus Medical Encyclopedia, American Psychiatric Association and The Cochrane Collaboration's Glossary of terms (APA 2013; Cochrane Glossary 2014; MedlinePlus 2014; UK Screening Glossary 2012).

Appendix 2. List of acronyms and abbreviations

Acronyms or abbreviations	Terms
45,X	monosomy X or Turner syndrome

(Continued)

47,XXX	trisomy X or triple X syndrome
47,XXY	Klinefelter syndrome
aCGH	array comparative genomic hybridisation
AFP	alpha-fetoprotein
Bioch/US	biochemical or ultrasound or both screening test
ccfDNA	circulating cell-free DNA
Chrom. 21	chromosome 21
CVS	chorionic villi sampling
FISH	fluorescence in situ hybridisation
gNIPT	genomics-based non-invasive prenatal testing
hCG	human chorionic gonadotropin
HSROC	hierarchical summary receiver operating characteristic
MPSS	massively parallel shotgun sequencing
NA	not applicable
ND	no data available
NGS	next generation sequencing
NR	not reported
NT	nuchal translucency
PAPP-A	pregnancy associated plasma protein A
QF-PCR	quantitative fluorescent polymerase chain reaction
QUADAS-2	QUality Assessment of Diagnostic Accuracy Studies
Ref. Chrom	reference chromosome
RS	reference standard
SCA	sex chromosome aneuploidy
SNP	single nucleotide polymorphism
T13	trisomy 13 or Patau syndrome
T18	trisomy 18 or Edward syndrome
T21	trisomy 21 or Down syndrome

(Continued)

TMPS	targeted massively parallel sequencing
uE3	unconjugated estriol

Appendix 3. Index test technical details

Typically, blood samples from pregnant women are obtained by venous puncture in the first or second trimester. After two centrifugation steps, plasma is separated from maternal whole blood and ccfDNA is extracted from plasma with commercial kits. DNA is converted into a genomic library where each of the DNA fragments are ligated with platform specific adapters. For TMPS only, libraries are clonally amplified before being sequenced. Then the libraries of several pregnant women are loaded on a next generation sequencer. The produced sequencing reads are aligned on a reference human genome to their respective chromosomal location and the number of sequence reads from each chromosome is computed (Rothberg 2011). MPSS randomly sequences DNA fragments from across the whole genome while TMPS sequences DNA fragments from selected regions (Figure 1). Ultimately, all gNIPT for aneuploidies rely on assigning sequence reads of DNA fragments to their chromosome of origin and comparing total number or proportions of reads or single nucleotide polymorphisms (SNP) genotype between each chromosome of interest (e.g. 13, 18, 21, X and Y) and a reference set of chromosomes. A Z score (or other statistics) are computed and a patient-specific risk can be assessed based on a risk threshold determined from read counts from a series of known euploid and aneuploid pregnancies. For MPSS, the counts from chromosomes of interest are normalised using the counts from all other chromosome sequences, while, for TMPS, the counts are normalised against a subset of selected sequences. Bioinformatic approaches vary according to the testing approach (MPSS or TMPS) and research team. Besides the use of normalised chromosome read counts, TMPS also allows for the use of additional allelic information when polymorphic loci such as SNP are targeted, such as an estimate of fetal DNA concentration (fetal DNA proportion) (Liao 2012). Thus, while MPSS produces a larger number of total sequence reads, TMPS will generate a larger number of reads from each targeted chromosomes.

Appendix 4. Search strategy

MEDLINE (Ovid)

Steps	Text words and subject headings	Sets of search
1	'cell-free dna'.mp	Index test
2	'cell free dna'.mp	
3	cfdna.mp	
4	ffdna.mp	
5	cffdna.mp	
6	'free foetal dna'.mp	
7	'free fetal dna'.mp	
8	nipd.mp	
9	nipt.mp	
10	(non invasive or noninvasive or non-invasive).mp	
11	(genetic adj2 (diagnos* or detect* or test* or screen*)).mp	
12	exp Genetic Testing/	
13	exp Sequence Analysis, DNA/	

(Continued)

14	((antenatal or ante natal) adj2 (diagnos* or detect* or test* or screen*)).mp	
15	((prenatal or pre natal) adj2 (diagnos* or detect* or test* or screen*)).mp	
16	exp Prenatal Diagnosis/	
17	or/1-16	
18	maternal.mp	Patient description
19	exp Pregnancy/	
20	exp Pregnancy Complications/	
21	pregnant.mp	
22	pregnanc*.mp	
23	exp Fetus/	
24	fetus.mp	
25	foetus.mp	
26	fetal.mp	
27	foetal.mp	
28	or/18-27	
29	trisom*.mp	Target condition
30	aneuploid*.mp	
31	(down* adj syndrome*).mp	
32	exp Aneuploidy/	
33	exp Trisomy/	
34	exp Down Syndrome/	
35	chromosome disorders.mp	
36	or/29-35	
37	or/1-5,11-13	Combined sets
38	6 or 7 or 8 or 9 or 14 or 15 or 16	
39	36 and 37 and 28 and 10	
40	38 and 37 and 36	
41	39 or 40	Final combined set

Embase (Embase.com)

Steps	Text words and subject headings	Sets of search
1	'cell-free dna'	Index test
2	'cell free dna'	
3	cfdna	
4	ffdna	
5	cffdna	
6	'free foetal dna'	
7	'free fetal dna'	
8	nipd	
9	nipt	
10	'non invasive'	
11	noninvasive	
12	'non-invasive'	
13	genetic NEXT/1 (diagnos* or screen* or test* or detect*)	
14	'genetic screening'/exp	
15	'genetic testing'/exp	
16	'sequence analysis dna'/exp	
17	antenatal NEXT/1 (diagnos* or screen* or test* or detect*)	
18	prenatal NEXT/1 (diagnos* or screen* or test* or detect*)	
19	pre?natal NEXT/1 (diagnos* or screen* or test* or detect*)	
20	'prenatal diagnosis'/exp	
21	1 OR 2 OR 3 OR 4 OR 5 OR 6 OR 7 OR 8 OR 9 OR 10 OR 11 OR 12 OR 13 OR 14 OR 15 OR 16 OR 17 OR 18 OR 18 OR 19 OR 20	
22	antenatal	Patient description
23	prenatal	
24	pre?natal	

(Continued)

25	maternal	
26	foetus	
27	fetus	
28	foetal	
29	fetal	
30	pregnanc*	
31	pregnant	
32	'pregnancy'/exp	
33	'pregnancy complications'/exp	
34	'pregnant woman'/exp	
35	'pregnant women'/exp	
36	22 OR 23 OR 24 OR 25 OR 26 OR 27 OR 28 OR 29 OR 30 OR 31 OR 32 OR 33 OR 34 OR 35	
37	trisom*	Target condition
38	aneuploid*	
39	down* NEXT/1 syndrome	
40	'aneuploid'/exp	
41	'aneuploidy'/exp	
42	'trisomy'/exp	
43	'downs syndrome'/exp	
44	'down syndrome'/exp	
45	'chromosome disorders'	
46	37 OR 38 OR 39 OR 40 OR 41 OR 42 OR 43 OR 44 OR 45	
47	21 and 36 and 46	Final combined set

Web of Science (ISI)

Steps	Text words and subject headings	Sets of search
-------	---------------------------------	----------------

(Continued)

1	TOPIC: (down* syndrome) <i>OR</i> TOPIC: (trisom*) <i>OR</i> TOPIC: (aneuploid*)	Target condition
2	TOPIC: (pregnan*)	Patient description
3	TOPIC: (dna) <i>OR</i> TOPIC: (blood)	Index test
4	TOPIC: (pre?natal screen*) <i>OR</i> TOPIC: (prenatal screen*) <i>OR</i> TOPIC: (pre?natal test*) <i>OR</i> TOPIC: (prenatal test*) <i>OR</i> TOPIC: (genetic test*) <i>OR</i> TOPIC: (genetic screen*) <i>OR</i> TOPIC: (prenatal diagnos*) <i>OR</i> TOPIC: (pre?natal diagnos*) <i>OR</i> TOPIC: (detection) <i>OR</i> TOPIC: (genetic diagnos*) <i>OR</i> TOPIC: (non invasive) <i>OR</i> TOPIC: (non-invasive) <i>OR</i> TOPIC: (noninvasive)	Index test
5	#1 AND #2 AND #3 AND #4	Final combined set

Cochrane Register of Diagnostic Test Accuracy Studies

Hand search: in diagnostic test accuracy database, there are 18 publications from the Cochrane Pregnancy and Childbirth group.

Clinicaltrials.gov

Steps	Text words and subject headings	Sets of search
1	(down syndrome OR trisomy OR aneuploidy)	Target condition
2	(testing OR screening OR diagnosis OR detection)	Index test
3	#1 AND #2	Final combined set

European Clinical Trials Register

Steps	Text words and subject headings	Sets of search
1	pregnan*	Population
2	trisom* OR aneuploid*	Target condition
3	#1 OR #2	Final combined set

Who ICTRP

Steps	Text words and subject headings	Sets of search
1	screen* OR detect* OR diagnos* OR test* OR pregnan*	Index test and population
2	«down syndrome » OR trisom* OR aneuploid*	Target condition
3	#1 AND #2	Final combined set

NTIS.gov

Steps	Text words and subject headings	Sets of search
1	(down syndrome OR trisomy OR aneuploidy)	Target condition
2	(testing OR screening OR diagnosis OR detection)	Index test
3	#1 AND #2	Final combined set

OpenGrey

Steps	Text words and subject headings	Sets of search
1	"down syndrome" OR trisom* OR aneuploid*	Target condition
2	screen* OR detect* OR diagnos* OR test* OR pregnan*	Index test and population
3	#1 AND #2	Final combined set

National Guideline Clearing House (NGCH)

Steps	Text words and subject headings	Sets of search
1	diagnosis (guideline category)	Index test
2	screening (guideline category)	Index test
3	aneuploid OR trisomy OR « down syndrome »	Target condition
4	(#1 OR #2) AND #3	Final combined set

TheseNet

Steps	Text words and subject headings	Sets of search
1	trisomy	Target condition
2	screening	Index test
3	#1 AND #2	Final combined set

These Canada Portal

Steps	Text words and subject headings	Sets of search
1	trisomy OR (down AND syndrome)	Target condition
2	screening	Index test
3	#1 AND #2	Final combined set

Appendix 5. Data collection form for study classification during full-text assessment

Heading	Detailed instructions	Data
Study ID	Last name of the first author and year of publication	Name: Year:
Reference details	Details allowing identification of the publication	Journal: Volume: Issue: Pages: Accession number (e.g. PMID ^a):
Multiple reports of this study	For example, duplicate publications or follow-up studies. Provide the study ID linked to this classified study	Study ID:
Type of report	Check the appropriate box	Journal article # Conference/abstract # Ongoing trial # Others # Specify:
Language	In which language was the study written	Language:

(Continued)

		Translation needed? Yes # No #
Eligibility	Provide reason for exclusion or awaiting classification (e.g. why authors should be contacted and what issues should be clarified)	Study excluded? Yes # No # Reason: Awaiting classification? Yes # No # Reason:
Report author contact details for further information	Date when the authors were contacted (dd/mm/yyyy)	No need for further contact # Authors have been contacted on: Reply received on:
Review author ID	Who completed the form	Name:
Date of classification	(dd/mm/yyyy)	Date:
Notes, questions or reminders		

^aPMID: PubMed identifier.

Appendix 6. QUADAS-2 tool for assessing methodological quality of included studies

	Signalling question	Signalling question	Signalling question	Risk of bias	Concerns about applicability
Domain 1: Patient selection					
Patient selection	Was a consecutive or random sample of patients enrolled?	Was a case-control design avoided?	Did the study avoid inappropriate exclusions?	Could the selection of patients have introduced bias?	Are there concerns that the included patients and setting do not match the review question?
	<p>Yes: if all consecutive or random samples or convenient samples or all eligible pregnant women were enrolled.</p> <p>No: if selected pregnant women were enrolled.</p> <p>Unclear: if this was not clear from the report.</p>	<p>Yes: if a case-control design was avoided.</p> <p>No: If a case-control design was not avoided.</p> <p>Unclear: if this was not clear from the report.</p>	<p>Yes: if the study avoided inappropriate exclusions.</p> <p>No: if pregnant woman was excluded based on family's situation, maternal age, ethnicity, maternal cancer history, type of pregnancy, gestational age, assisted reproductive technology or</p>	<p>Low risk: if 'yes' for all signalling questions.</p> <p>High or unclear risk: if 'no' or 'unclear' was reported for at least one signalling question.</p>	<p>Low concern: if the selected pregnant women represent the women indicated by the review question^a.</p> <p>High concern: if selected pregnant women differ from those targeted by the review question^a.</p> <p>Unclear concern: if insufficient information was available.</p>

(Continued)

any other aneuploidies.

Unclear: if this was not clear from the report.

Domain 2: Index test

Index test ^b	Were the index test results interpreted without knowledge of the results of the reference standard ^c ?	If a threshold was used, was it prespecified?	Could the conduct or interpretation of the index test have introduced bias?	Are there concerns that the index test, its conduct, or interpretation differ from the review question?
	<p>Yes: if the gNIPT results were interpreted without knowledge of the results of the reference standard^c.</p> <p>No: if the gNIPT results were interpreted with knowledge of the results of the reference standard^c.</p> <p>Unclear: if this was not clear from the report.</p>	<p>Yes: if criteria for a positive test were prespecified.</p> <p>No: if the criteria for a positive test were not prespecified.</p> <p>Unclear: if this was not clear from the report.</p>	<p>Low risk: if 'yes' for all signalling questions.</p> <p>High or unclear risk: if 'no' or 'unclear' was reported for at least one signalling question.</p>	<p>Low concern: if the gNIPT was performed such as described in the review question^a.</p> <p>High concern: if gNIPT vary from those specified in the review question.</p> <p>Unclear concern: if insufficient information was available.</p>

Domain 3: Reference standard

Reference Standard ^c	Is the reference standard ^c likely to correctly classify the target condition ^d ?	Were the reference standard results interpreted without knowledge of the results of the index test ^b ?	Could the reference standard, its conduct, or its interpretation have introduced bias?	Are there concerns that the target condition as defined by the reference standard does not match the review question?
	<p>Yes: if one appropriate reference standard^c was used.</p> <p>No: if pregnant women did not undergo appropriate reference standard^c.</p> <p>Unclear: if this was not clear from the report.</p>	<p>Yes: if karyotype results were interpreted without knowledge of results of the index test^b.</p> <p>No: if karyotype results were interpreted with the knowledge of results of the index test^b.</p> <p>Unclear: if this was not clear from the report.</p>	<p>Low risk: if 'yes' for all signalling questions.</p> <p>High or unclear risk: if 'no' or 'unclear' was reported for at least one signalling question.</p>	<p>Low concern: if the reference standard^c were used as described in the review question^a.</p> <p>High concern: if the reference standard^c vary from those specified in the review question^a.</p> <p>Unclear concern: if insufficient information was available.</p>

(Continued)

Domain 4: Flow and timing

Flow and timing	Was there an appropriate interval between gNIPT and reference standard?	Did all analysed patients receive the reference standard?	Were all patients included in the analysis?	Could the patient flow have introduced bias?
	<p>Yes: if the interval between blood collection for gNIPT and fluid collection for reference standard^c was more than one day (only if blood collection occurred after fluid collection). If blood collection occurred before fluid collection, there is no time limit^e.</p> <p>No: if the interval between blood collection for gNIPT and the fluid collection for reference standard^c was less than one day if the blood collection occurred after the fluid collection.</p> <p>Unclear: if this was not clear from the report.</p>	<p>Yes: if all pregnant women analysed have appropriate reference standard^c.</p> <p>No: if some pregnant women analysed do not have a karyotype result.</p> <p>Unclear: if this was not clear from the report.</p>	<p>Yes: if all pregnant women recruited into the study were included in the analysis or if failed samples occurred before NGS process.</p> <p>No: if all pregnant women recruited into the study were not included in the analysis or if failed samples occurred during NGS process.</p> <p>Unclear: if this was not clear from the report.</p>	<p>Low risk: if 'yes' for all signalling questions.</p> <p>High or unclear risk: if 'no' or 'unclear' was reported for at least one signalling question.</p>

^aReview question: what is the diagnostic accuracy of massively parallel shotgun sequencing (MPSS) and targeted massively parallel sequencing (TMPS) using circulating cell-free DNA (ccfDNA) in maternal blood for the detection of common fetal aneuploidies (T21, T18, T13, 45,X, 47,XXY, 47,XXX and 47,XYY) in pregnant women according to their prior risk of fetal aneuploidy?

^bIndex test refers to genomics-based non-invasive prenatal testing (gNIPT) methods such as MPSS or TMPS.

^cThe appropriate reference standard is karyotyping (traditional banding techniques or spectral karyotyping from invasive methods like chorionic villi sampling or amniocentesis), chromosome analysis (e.g. FISH, aCGH and QF-PCR), clinical examination or medical record from birth (for T21, T18 or T13). For sex chromosome aneuploidies, only fetal karyotype was appropriate reference standard because they usually have a normal phenotype.

^dTarget conditions (aneuploidies) are T21, T18, T13, 45,X, 47,XXY, 47,XXX and 47,XYY.

^eTarget conditions (aneuploidies) do not vary over time.

Appendix 7. gNIPT accuracy in mixed prior risk of fetal aneuploidy

Test	Number of studies	Number of affected	Number of unaffected pregnancies ^a	Sensitivity ^b % (95% CI)	Specificity ^b % (95% CI)	P value ^c
------	-------------------	--------------------	---	--	--	----------------------

(Continued)

**pregnan-
cies**

T21, mixed risk						
MPSS	10	445	30,962	96.0 (93.7 to 97.4)	99.9 (99.9 to 100)	
TMPS	6	169	6925	98.2 (94.6 to 99.4)	100 (99.9 to 100)	
Difference between MPSS and TMPS				-2.27 (-4.97 to 0.43)	-0.04 (-0.08 to -0.002)	0.10
Tradition- al screening tests ^d	1	3	1909	100 (43.9 to 100)	96.4 (95.5 to 97.1)	
T18, mixed risk						
MPSS	9	113	30,637	100 (98.3 to 100) ^e	99.9 (99.8 to 100)	
TMPS	4	53	5569	98.1 (87.8 to 99.7)	99.9 (99.8 to 100)	
Tradition- al screening tests	1	1	1905	100 (20.7 to 100)	99.4 (99.0 to 99.7)	
T13, mixed risk^f						
MPSS	8	27	30,384	100 (87.5 to 100) ^e	100 (> 99.9 to 100) ^e	
TMPS	5	31	8362	78.6 (48.3 to 93.5)	99.9 (99.8 to 100)	
45X, mixed risk						
MPSS	2	12	296	91.7 (58.7 to 98.8)	100 (98.7 to 100) ^e	
TMPS	2	22	1128	90.9 (70.0 to 97.7)	99.9 (99.4 to 100)	
47,XXY, mixed risk						
MPSS	1	1	107	ND		
TMPS	1	2	184	ND		
47,XYY, mixed risk						
MPSS	1	1	199	ND		
TMPS	1	1	185	ND		
Autosomes (T21, T18 and T13 combined), mixed risk						
MPSS	10	585	30,822	96.9 (95.2 to 98.1)	99.9 (99.9 to 99.9)	
TMPS	7	253	8793	96.0 (92.8 to 97.9)	99.8 (99.7 to 99.9)	
Difference between MPSS and TMPS				0.88 (-1.90 to 3.65)	0.07 (-0.02 to 0.16)	0.25

(Continued)

Traditional screening tests	1	4	1908	100 (51.0 to 100)	95.8 (94.8 to 96.6)
-----------------------------	---	---	------	-------------------	---------------------

SCA (45,X, 47,XXX, 47,XXY and 47,XYY combined), mixed risk

MPSS	2	14	294	92.9 (63.0 to 99.0)	100 (98.7 to 100) ^e
TMPS	2	25	1125	92.0 (73.1 to 98.0)	99.9 (99.4 to 100)

45,X: Turner syndrome, 47,XXX: triple X syndrome, 47,XXY: Klinefelter syndrome, ND: no data available, T21: trisomy 21, T18: trisomy 18, T13: trisomy 13, CI: confidence interval, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, SCA: sex chromosome aneuploidies.

^aWe included pregnancies with any other aneuploidy than the one under analysis with all euploid cases as "unaffected" pregnancies.

^bFor two or more studies, the sensitivities and specificities are the summary estimates obtained from meta-analysis. Sensitivity and specificity, and their 95% CIs are reported as percentages.

^cThe P value indicates the statistical significance of the difference in model fit and was obtained from likelihood ratio tests comparing models with and without a covariate for test type.

^dTraditional screening test are first trimester combined test, second trimester quadruple test, second trimester fully integrated test, second trimester sequential test or second trimester triple test.

^eSimple pooling used to obtain summary estimates of sensitivity and/or specificity.

^fTest comparison analysis did not converge.

Appendix 8. Investigation of heterogeneity

Test subgroups		Number of studies	Number of affected pregnancies	Number of unaffected pregnancies ^a
Reference standard				
Autosomes, unselected population				
MPSS	mixed reference standard ^b	1	11	1730
TMPS	karyotyping ^c	1	11	181
	mixed reference standard	3	107	20,468
Autosomes, selected high-risk population				
MPSS	karyotyping	22	1075	7028
	mixed reference standard	10	433	8769
TMPS	karyotyping	7	378	4282

(Continued)

SCA, selected high-risk population

MPSS	karyotyping	10	134	3943
	mixed reference standard	2	17	3509
TMPS	karyotyping	4	96	968

Ethnicity
Autosomes, unselected population

MPSS	more than 50% Asian ^d	1	11	1730
TMPS	more than 50% Caucasian ^e	3	107	20,468
	not reported	1	11	181

Autosomes, selected high-risk population

MPSS	more than 50% Asian	14	206	6589
	more than 50% Caucasian	7	843	6262
	not reported	11	459	2946
TMPS	more than 50% Caucasian	3	237	3744
	not reported	4	141	538

SCA, selected high-risk population

MPSS	more than 50% Asian	5	25	1852
	more than 50% Caucasian	5	96	4286
	not reported	2	30	1314
TMPS	more than 50% Caucasian	1	56	116
	not reported	3	40	852

45,X: Turner syndrome, 47,XXX: triple X syndrome, 47,XXY: Klinefelter syndrome, T21: trisomy 21, T18: trisomy 18, T13: trisomy 13, MPSS: massively parallel shotgun sequencing, TMPS: targeted massively parallel sequencing, SCA: sex chromosome aneuploidies

^aWe included pregnancies with any other aneuploidy than the one under analysis with all euploid cases as "unaffected" pregnancies.

^bMixed RS: include karyotyping and neonatal clinical examination or medical records from birth.

^cKaryotyping: include fetal karyotyping performed on cells obtained from chorionic villi sampling (CVS), amniotic fluid, placental tissue, a fetus lost by miscarriage or other equivalent and recognised methods on the same materials.

^dMore than 50% Asian: in the cohort, more than 50% of all pregnant women were Asian ethnicity.

^eMore than 50% Caucasian: in the cohort, more than 50% of all pregnant women were Caucasian ethnicity.

HISTORY

Protocol first published: Issue 7, 2015

Review first published: Issue 11, 2017

Date	Event	Description
2 July 2015	Amended	Minor changes to author affiliations.

CONTRIBUTIONS OF AUTHORS

FR initiated this review project and recruited all authors. MB co-ordinated the production of the review.

MB, CL, JB, YT and FR drafted the protocol and all authors critically commented on the protocol. The final protocol was read and approved by all authors.

MB, CL and WW developed and applied the literature search. MB, CL, JB and LN applied eligibility criteria, collected, filled the data form and assessed methodological quality of the included studies. MB contacted authors of the included studies. YT checked data collection, undertook statistical analysis and data synthesis. MB, CL and FR wrote the draft of the review. MB, CL, JB, LN, YT, SL, YG and FR contributed with redrafting. SL, FL, YG and FR provide senior clinical input. FR oversaw the review process. The final review was read and approved by all authors.

DECLARATIONS OF INTEREST

Mylene Badeau: none known.

Jonatan Blais: none known.

Yves Giguère (and some other members of the review team - FR, SL, FL) are investigators in a Research Project funded under the auspices of Genome Canada and the Canadian Institutes for Health Research (both non-for-profit organisations funded by the Canadian government) but that call for some mandatory in-kind contributions from other partners. This Research Project thus receives in-kind funding from private corporations which either offer commercial NIPT tests (Ariosa Diagnostics Inc, San Jose, CA) or offer reagents and/or equipment that can be used to perform NIPT assays (Life Technologies Inc, NY, USA; Illumina, San Diego, CA, USA; QIAGEN, Hilden, GER; Perkin Elmer, Waltham, MASS, USA). This funding is at arms length from the scientific components of the research project. The present review is not funded by this Project. Some of the authors may eventually publish results from clinical trials that could be considered in this review. None of the four authors involved in gNIPT studies will take part in the selection of studies, nor in any decisions/analyses related to their own studies. All authors declared no other conflict of interest

Sylvie Langlois (and some other members of the review team - FR, YG, FL) are investigators in a Research Project funded under the auspices of Genome Canada and the Canadian Institutes for Health Research (both non-for-profit organisations funded by the Canadian government) but that call for some mandatory in-kind contributions from other partners. This Research Project thus receives in-kind funding from private corporations which either offer commercial NIPT tests (Ariosa Diagnostics Inc, San Jose, CA) or offer reagents and/or equipment that can be used to perform NIPT assays (Life Technologies Inc, NY, USA; Illumina, San Diego, CA, USA; QIAGEN, Hilden, GER; Perkin Elmer, Waltham, MASS, USA). This funding is at arms length from the scientific components of the research project. The present review is not funded by this Project. Some of the authors (FR, SL, YG and FL) may eventually publish results from clinical trials that could be considered in this review. None of these four authors involved in gNIPT studies will take part in the selection of studies, nor in any decisions/analyses related to their own studies. All authors declared no other conflict of interest. This review was supported by a Canadian Institutes of Health Research (CIHR) Knowledge Synthesis Grant: Fall 2014 Competition (2014-11-17) awarded to AFT, FR, FL, and SL. CIHR in no way influenced the results or conclusions of this review.

France Légaré (and some other members of the review team - FR, YG, SL) are investigators in a Research Project funded under the auspices of Genome Canada and the Canadian Institutes for Health Research (both non-for-profit organisations funded by the Canadian government) but that call for some mandatory in-kind contributions from other partners. This Research Project thus receives in-kind funding from private corporations which either offer commercial NIPT tests (Ariosa Diagnostics Inc, San Jose, CA) or offer reagents and/or equipment that can be used to perform NIPT assays (Life Technologies Inc, NY, USA; Illumina, San Diego, CA, USA; QIAGEN, Hilden, GER; Perkin Elmer, Waltham, MASS, USA). This funding is at arms length from the scientific components of the research project. The present review is not funded by this Project. Some of the authors (FR, SL, YG and FL) may eventually publish results from clinical trials that could be considered in this review. None of these four authors involved in gNIPT studies will take part in the selection of studies, nor in any decisions/analyses related to their own studies. All authors declared no other conflict of interest. This review was supported by a Canadian Institutes of Health

Research (CIHR) Knowledge Synthesis Grant: Fall 2014 Competition (2014-11-17) awarded to AFT, FR, FL, and SL. CIHR in no way influenced the results or conclusions of this review.

Carmen Lindsay: none known.

Leon Nshimyumukiza: none known.

François Rousseau (and some other members of the review team - SL, YG, FL) are investigators in a Research Project funded under the auspices of Genome Canada and the Canadian Institutes for Health Research (both non-for-profit organisations funded by the Canadian government) but that call for some mandatory in-kind contributions from other partners. This Research Project thus receives in-kind funding from private corporations which either offer commercial NIPT tests (Ariosa Diagnostics Inc, San Jose, CA) or offer reagents and/or equipment that can be used to perform NIPT assays (Life Technologies Inc, NY, USA; Illumina, San Diego, CA, USA; QIAGEN, Hilden, GER; Perkin Elmer, Waltham, MASS, USA). This funding is at arms length from the scientific components of the research project. The present review is not funded by this Project. Some of the authors (FR, SL, YG and FL) may eventually publish results from clinical trials that could be considered in this review. None of these four authors involved in gNIPT studies will take part in the selection of studies, nor in any decisions/analyses related to their own studies. All authors declared no other conflict of interest. This review was supported by a Canadian Institutes of Health Research (CIHR) Knowledge Synthesis Grant: Fall 2014 Competition (2014-11-17) awarded to AFT, FR, FL, and SL. CIHR in no way influenced the results or conclusions of this review.

Yemisi Takwoingi: The University of Birmingham received consultancy fees from Universite Laval for statistical analysis.

Alexis F Turgeon: This review was supported by a Canadian Institutes of Health Research (CIHR) Knowledge Synthesis Grant: Fall 2014 Competition (2014-11-17) awarded to AFT, FR, FL, and SL. CIHR in no way influenced the results or conclusions of this review.

William Witteman: none known.

SOURCES OF SUPPORT

Internal sources

- No sources of support supplied

External sources

- Research Chair in Health Technology Assessment in laboratory medicine to FR, Canada.

From Ministère de la Santé et des Services Sociaux, Fonds de la Recherche en Santé du Québec, Centre Hospitalier Universitaire de Québec.

- Canadian Institutes of Health Research (CIHR), Canada.

Knowledge Synthesis Grant: Fall 2014 Competition (2014-11-17) (AFT, FR, FL, SL)

- Canadian Institutes of Health Research (CIHR), Canada, Canada.

Canada Research Chair in Critical Care Neurology and Trauma (to AFT).

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

In the “reference standards” section in the methods, the protocol did not specify that neonatal clinical examination or medical records from birth are not appropriate reference standards for SCA diagnosis. These aneuploidies (SCA) usually have a normal phenotype and can not be detected with neonatal clinical examination or medical records from birth.

In the “sensitivity analyses” section, of the protocol, the authors stated they intended to investigate “studies where not all pregnant women received neither reference standard (no karyotyping confirmation nor birth follow-up); authors who have taken for granted that the baby is normal”. We decided to remove this analysis because this goes against one of our criteria for considering studies for this review (pregnant women with MPSS or TMPS and a reference standard). These type of studies were excluded.

We decide to remove Google scholar from our electronic searches databases list. Google scholar found more than 100,000 publications about our topic but only the first 1000 are retrievable. This database is not reproducible and search fields did not allow us to specify the search strategy. All first most relevant publications were already found with other databases.

We changed OpenSIGLE to Opengrey because the first has been replaced by the second.

Although studies used different cutpoints, there was little or no variation in threshold and no requirement to estimate the correlation between sensitivity and specificity across studies in a meta-analysis. Therefore, we did not estimate summary ROC curves using the HSROC model. As the cutpoints were regarded as qualitative, we estimated summary sensitivities and specificities using random-effects and

fixed-effect logistic regression models, and simple pooling as appropriate. Further details are available in the statistical analysis and data synthesis section. We used the Stata software package for the analyses instead of SAS.

We made changes in the QUADAS-2 tool. In domain 1, at third signalling question, we added "maternal cancer history, type of pregnancy, gestational age, assisted reproductive technology" in "No" answer. In concerns of applicability in domain 3, concerns about applicability, we modified conditions for low and high concern according to the review question. In domain 4, at second signalling question, we added "analysed" for clarify and we changed "yes" answer by removing "karyotype result" for "appropriate reference standard".

INDEX TERMS

Medical Subject Headings (MeSH)

*Aneuploidy; Cell-Free Nucleic Acids [*blood]; Chromosome Disorders [*diagnosis] [genetics]; Disorders of Sex Development [diagnosis] [genetics]; Fetal Diseases [*diagnosis] [genetics]; High-Throughput Nucleotide Sequencing [*methods]; Pregnancy, High-Risk; Prenatal Diagnosis [*methods]

MeSH check words

Female; Humans; Pregnancy