

Do Stroke Patients have an Increased Risk of Developing Suicidal Ideation or Dying by Suicide? An Overview of the Current Literature

Maurizio Pompili,^{1,2} Paola Venturini,³ Sandra Campi,³ Maria E. Seretti,¹ Franco Montebovi,³ Dorian A. Lamis,⁴ Gianluca Serafini,¹ Mario Amore⁵ & Paolo Girardi¹

1 Department of Neurosciences, Mental Health and Sensory Organs, Suicide Prevention Center, Sant'Andrea Hospital, Sapienza University of Rome, Rome, Italy

2 McLean Hospital, Harvard Medical School, Belmont, MA, USA

3 Psychiatry Residency Program, Faculty of Medicine and Psychology, Sapienza University of Rome, Rome, Italy

4 Department of Psychiatry and Behavioral Sciences, Emory University School of Medicine, Atlanta, GA, USA

5 Department of Neurosciences, Section of Psychiatry, University of Parma, Parma, Italy

Keywords

Intervention; Poststroke depression; Prevention; Stroke; Suicide.

Correspondence

M. Pompili, M.D., Ph.D., Department of Neurosciences, Mental Health and Sensory Organs, Suicide Prevention Center, Sant'Andrea Hospital, "Sapienza" University of Rome, 1035-1039, Via di Grottarossa, 00189 Rome, Italy.

Tel.: +39-06-33775675;

Fax: +39-06-33775342;

E-mail: maurizio.pompili@uniroma1.it

Received 7 April 2012; revision 8 May 2012;

accepted 10 May 2012.

doi: 10.1111/j.1755-5949.2012.00364.x

The first two authors contributed equally to this work.

Introduction

Suicidal behavior and stroke are both dramatic events that contribute to an increased risk of death and functional impairments. The World Health Organization has established that 15 million people suffer from stroke worldwide each year with 5 million of them dying and another 5 million becoming permanently disabled [1]. Stroke is considered one of the most devastating of all neurological disorders, often causing death or gross physical impairment and disability [2]. Although strokes can and do occur at any age, in the United States in 2001, 74.5% of all strokes occurred among people age 75 or older, while 88% occurred among individuals aged 65 or older: the risk of having a stroke more than doubles each decade after 55 years of age [3].

In the United States, approximately 32,000 individuals die by suicide each year [4] and 5000 (14%) of them are older than

SUMMARY

Stroke is a leading cause of death that affects 15 million people worldwide each year. Increasing evidence suggests that stroke confers substantial risk for suicide and following a stroke, patients frequently develop poststroke depression, which is a well-established risk factor for suicide. In this overview of the current literature, we examined the association between suffering a stroke and subsequent risk for suicide and suicidal ideation. We performed a careful MedLine, Excerpta Medica, PsycLit, PsycInfo, and Index Medicus search to identify all articles and book chapters in English. We initially selected 31 articles published between 1990 and 2011; however, only 16 studies were included in this review. All articles identified stroke as a significant risk factor for suicide, especially among depressed patients, providing further support for poststroke depression and suicidality. The results also indicated that there were differences between patients who developed acute-onset suicidal plans and those who reported delayed-onset plans, which occurred more frequently. Many of the stroke patients who died by suicide suffered from depression prior to their death, suggesting that being diagnosed with a mood disorder contributes to an increased risk of suicide in this population. Suffering from a stroke increases the risk of dying by suicide and developing suicidal ideation, particularly in young adults and women. The factors found to confer the most risk for suicidality were depression, previous mood disorder, prior history of stroke, and cognitive impairment.

65 years of age. In 2007, the rate of suicide in the elderly was 14.3 per 100,000 compared to 10.9 suicides per 100,000 in the general population [5]. Previous studies have shown that physical illnesses, especially neurological disorders, increase suicide risk [6–10]. Moreover, it has been well established that people who survive a stroke often develop poststroke depression which is consistently implicated as an important risk factor for suicide [11–13]. Given that stroke and suicide are both leading causes of death and contribute to significant societal costs, the purpose of this review is to investigate whether or not individuals who suffer a stroke are at an increased risk of having suicidal thoughts or dying by suicide.

Material and Methods

To provide a new and timely review examining the association between stroke and suicide, we performed a careful MedLine, Ex-

cerpta Medica, PsycLit, PsycInfo, and Index Medicus search to identify all relevant refereed journal articles and book chapters in English. The following search terms were used: suicide* AND stroke, suicide* after stroke, suicide* as cause of death in stroke patients. Only articles published in English were considered in this review, including studies that added an original contribution to the literature. If a title or abstract appeared to describe a study eligible for inclusion, the full article was obtained and assessed for relevance based on our predetermined inclusion criteria. Any discrepancies in literature eligibility between the two independent reviewers (MP, PV) were resolved by consultations with senior authors (GS, PG). In addition, we also examined reference lists and contacted experts in the field. This search strategy identified 31 articles; of these 16 were eligible for the purpose of this review. All of the studies included are reported in Table 1.

Suicidal Ideation and Suicidal Plans in Stroke Patients

The original studies investigating the relation between stroke and suicidal plans were conducted by Kishi and colleagues in 1996 [14,15]. The researchers recruited 301 patients and evaluated their suicidal plans after suffering an acute stroke (i.e., baseline) and at 3, 6, 12, and 24 months follow-up. They identified stroke as a risk factor for suicide in both studies, demonstrating that 6.6% of patients reported acute-onset suicidal plans, whereas 11.3% of stroke victims reported delayed-suicidal plans within 2 years of their stroke. The development of both acute- and delayed-onset suicidal plans in stroke patients has been found to be related to the presence of depressive disorders, particularly major depression, as well as to a prior history of stroke. The authors also noted some differences between the two kinds of onset, suggesting that the etiology of these two types of suicidal plans may be different. Specifically, delayed-onset suicidal plans have been found to be related to greater physical impairment from the stroke, and these patients had more frequent posterior lesions; whereas, acute-onset plans were associated with alcohol abuse and younger age. Furthermore, they found that suicidal ideation during the acute medical treatment period was associated with the presence of a diagnosable depressive disorder and that the severity of the illness was not an independent risk factor for developing a suicidal plan. This finding suggests that suicidal plans were not simply psychological reactions to physical impairment after the acute event.

Five years later, Pohjasvaara et al. [16] conducted a case-control study that substantially confirmed Kishi's [14,15] previous results providing further support for stroke being a significant risk factor for developing suicidal ideation. The authors of this study examined the frequency and clinical determinants of suicidal ideas at 3 and 15 months after stroke. They noted that the patients with suicidal ideas were more depressed at 3 and 15 months after stroke as compared to patients without suicidal ideas. Moreover, patients with suicidal ideas more often had a history of stroke, right-sided stroke, were more disabled, and more dependent in living 15 months after stroke. The researchers also found that motivated aspects of depression (e.g., anxiety, hyperemotionalism) were more common in depressed stroke patients, whereas the unmotivated or biologically determined aspects (e.g., suicidal

ideas, depressed mood, feelings of guilt) were more frequently reported in patients with an endogenous form of major depression. Unmotivated aspects, if present in stroke patients, were often attributed to the actual condition (stroke), but not to endogenous depression. This may explain why recurrent strokes have been found to be an independent correlate of suicidal ideas 15 months after stroke.

Rao et al. [17] found that individuals with cerebrovascular disease were at an increased risk of suicide. Specifically, the researchers evaluated patients with stroke, transient ischemic attack (TIA), and with peripheral vascular disease (PVD) and concluded that the impairment of the frontal lobe increased both impulsivity and suicide risk. In accordance with these results, Santos et al. [18] examined 117 stroke patients and reported that 15% of acute stroke patients presented with suicidal thoughts. Additionally, the researchers demonstrated that stroke patients with suicidal ideation had a lower educational level and were more likely to have had a previous mood disorder or acute depression than those without suicidal thoughts. Similarly, Fuller-Thompson et al. [19] concluded that stroke survivors had twice the odds of depression and suicidal ideation as compared to those who had never suffered a stroke. Further, Caucasians and younger individuals had higher odds of poststroke depression and suicidal ideation.

Tang et al. [20] investigated the role of fatigue in stroke survivors and revealed that poststroke fatigue was a significant predictor of suicidal ideation independent of depression.

Finally, Chan et al. [21] found that high cerebrovascular risk factor (CVFR) scores (derived by the weighted sum of severities of specific factors such as systolic blood pressure, antihypertensive treatment, coronary vessel disease, diabetes mellitus, hypertension, atrial fibrillation, left-ventricular hypertrophy, and cigarette smoking) were associated with an increased risk for suicide in middle age and older adults after accounting for age, sex, depression diagnosis, the presence of stroke, and functional status.

Suicide and Suicide Attempt in Stroke Patients

To estimate the percentage of suicides in patients who suffered a stroke, Stenager et al. [22] conducted an epidemiological study in Denmark over a period of 25 years from 1973 to 1998. In a sample of almost 38,000 patients with stroke, the researchers reported that 140 (7.2%) patients had died by suicide and suicide risk was significantly higher after stroke, especially in women and in the age group up to 60 years. Similar findings were reported in another study conducted in Denmark during the same time period [23]. Specifically, Teasdale and Engberg [23] found that there was an increased risk of suicide after stroke with the greatest risk being among younger patients, in relatively less severe cases, and within the first 5 years after stroke. Interestingly, the authors also concluded that the risk for suicide decreased with the duration of hospitalization and suggested that patients who had suffered more severe strokes requiring longer hospitalization could have more functional impairments reducing their capacity to carry out a suicidal act. In a recent population-based study examining the prevalence of stroke in suicide victims in Northern Finland [24], results indicated that 70% of stroke patients who died by suicide suffered from prestroke depression, with the highest risk for suicide being within 2 years after stroke. As previously noted, Teasdale and En-

Table 1 Main relevant studies about the association between stroke and suicide

Study	Study design	Eventual follow-up	Participants	Main findings	Limitations
Kishi et al. [14]	Case-control study	No	301 patients (cases n = 20 controls n = 281) evaluated for suicidal plans after acute stroke	<p>19 patients had suicidal plans and 1 attempted suicide (6.6%). Of those patients, 15 (75%) had major depressive disorder (MDD), 3 had minor depression (15%), and 1 (10%) was not depressed compared to 281 (93.4%) who did not have suicidal plans</p> <p>Those who had suicidal plans were more likely to have greater frequency of alcohol abuse, psychiatric disorders especially major depression, a previous personal history of cerebrovascular accidents, and sensory deficits compared to those who had not</p> <p>Suicidal patients were younger, had poorer social functioning, and greater cognitive impairment than nonsuicidal patients</p> <p>57 patients were diagnosed with MDD, of which 15 (26.3%) reported suicidal plans and 42 did not have a plan</p> <p>Depressed patients who reported suicidal plans had greater social withdrawal and brooding than depressed patients without suicidal plans</p>	<p>Authors evaluated only suicidal thoughts without investigating suicidal behavior so they were unable to know whether these patients attempted or died by suicide</p> <p>Patients with deficits in comprehension were excluded; therefore, findings may be not applicable to all patients with stroke</p> <p>Participants were predominantly of low socioeconomic status; so, the findings may not be generalizable to other socioeconomic classes</p>
Kishi et al. [15]	Longitudinal case-control study	24 months	301 patients (cases with acute-onset n = 20, delayed-onset n = 16, controls n = 117) evaluated for suicidal plans after acute stroke	<p>16 (11.3%) who did not have suicidal plans at the initial evaluation developed suicidal plans later, including 2 who attempted suicide, 7 after 6 months, 3 after 12 months, and 6 after 24 months. 10 of them had an initial diagnosis of MDD, 2 minor depression and 4 are not depressed vs. 117 who never developed suicidal plans over 24 months</p> <p>Greater frequency of alcohol abuse in acute-onset suicidal patients than in delayed-onset or nonsuicidal groups. Higher frequency of previous cerebrovascular accidents in both acute- and delayed-onset suicidal groups than in nonsuicidal group. Acute-onset</p>	<p>Authors evaluated suicidal thoughts only without investigating suicidal behavior so they are unable to ascertain whether or not these patients attempted or died by suicide</p> <p>Patients with deficits in comprehension were excluded; so findings may not be applicable to all patients who have suffered from stroke</p> <p>Participants were predominantly of low socioeconomic status. so the findings may not generalize to other classes</p> <p>Authors did not consider "passive suicide" (such as failure to follow medical advice that might constitute a form of passive suicide)</p>

(continued)

Table 1 (Continued)

Study	Study design	Eventual follow-up	Participants	Main findings	Limitations
Katayama et al. [25]	Case reports of suicides 6 months after stroke between 1999 and 2006	No	6 (n = 5 men, n = 1 woman) patients who died unexpectedly by suicide within 6 months after stroke	<p>patients were younger than nonsuicidal patients.</p> <p>Higher frequency of posterior lesions in delayed group compared to acute-onset group</p> <p>Higher frequency of social, physical, and cognitive impairment in both acute and delayed-onset suicidal groups</p> <p>Higher frequency of early morning awakening in acute-onset compared to delayed-onset groups</p> <p>4 patients attempted suicide, 2 survived</p> <p>None were diagnosed with MDD prior to stroke</p>	<p>Small sample size that did not allow findings to be generalized</p> <p>No data from any statistical analysis</p> <p>The study was retrospective in nature; therefore, it may be subject to recall bias</p>
Scott et al. [27]	Retrospective study, data from 14 countries of the World Mental Health	No	37,915	Those who had a heart attack and stroke were more likely (OR 1.9; 1.1–3.5) to have lifetime attempts in the total sample as well as among ideators with a plan (OR 2.7; 0.9–8.0) and to have lifetime ideation in the total sample (OR 1.9; 1.3–2.8)	<p>The study did not directly investigate the association between stroke and suicide but rather the association between suicide and several physical conditions</p> <p>Retrospective self-reports of the occurrence and timing of suicidal behavior, mental disorders, and covariates such as childhood adversity</p> <p>Possible bias in terms of sample selection: mixed results about stroke and heart attack were reported</p> <p>Physical conditions that were assessed via self-report</p>
Tang et al. [20]	Cross-sectional survey	No	595 patients with acute ischemic stroke admitted in Hong-Kong from 2005 to 2008	<p>76 patients (12.8%) reported suicidality</p> <p>Subjects who reported suicidality scored higher on the Fatigue Severity Scale (FSS) than those without suicidality</p> <p>Association between fatigue and suicidality is independent of depression</p> <p>55 (10%) patients with stroke died by suicide compared to 2865 who did not</p>	<p>Having recruited an hospital-based sample may preclude generalizability of findings to patients treated in other settings</p>
Voaklander et al. [28]	Case-control study	No	3601 (cases n = 602, controls n = 2999) evaluated between 1993 and 2002	<p>The study did not directly investigate the association between stroke and suicide</p>	<p>The study did not directly investigate the association between stroke and suicide</p>

(continued)

Table 1 (Continued)

Study	Study design	Eventual follow-up	Participants	Main findings	Limitations
Placido et al. [29]	Time series of 27 years between 1979 and 2005	No	3,364,908 deaths from suicide, ischemic heart disease or stroke	Depression, psychosis, neurosis, stroke, cancer, incontinence, liver disease, and previous history of self-harm were risk factors for suicide Incidence of death from suicide was positively correlated with ischemic heart disease and stroke ($r = 0.92$; $P < 0.0001$)	Potential underestimation of comorbidities The study was retrospective in nature Authors did not include several variables (e.g., social isolation, loss of spouse, family history of suicide) known to be risk factors for suicide in the elderly The study did not directly investigate the association between stroke and suicide Time-series analysis may suffer from several limitations, especially the association between life events in different subjects Cardiovascular risk related to highly lethal events was underestimated Cardiovascular risk in suicide is underestimated because of early death
Forsstrom et al. [24]	Retrospective population-based study of suicide victims in northern Finland from 1988 to 2007	No	2,283 suicide victims	75 (3.4%) patients suffered from stroke 80% were not under the influence of alcohol Compared to other victims, suicide victims were about 20 years older and more often women Approximately 70% of suicide victims with a previous stroke suffered from prestroke depression compared to 30% who never suffered from depression 359 suicides occurred in the study Standardized mortality ratio (SMR) was greater in men than in women SMR greatest in patients under 50 and lowest for patients 80 or older and relatively less severe cases Suicides were negatively related to duration of hospitalization, suicidal risk was lowest for those hospitalized for more than 3 months and highest for those hospitalized for less than 2 weeks	Data derived from only patients who needed hospitalization The study did not include a control group The severity of stroke in suicide victims was not evaluated
Teasdale et al. [23]	Retrospective study	No	114,098 patients discharged from hospital during 1979 and 1993		Over 50% of the diagnoses of stroke were of the undefined subtype Recording of suicide as cause of death may be unreliable and under-reported There are no data on laterality of stroke

(continued)

Table 1 (Continued)

Study	Study design	Eventual follow-up	Participants	Main findings	Limitations
Stenager et al. [22]	Retrospective study	No	37,869	Risk factors for suicide is greatest up to about 5 years after stroke 140 (7.2%) died by suicide Stroke increases suicidal risk especially in the age group up to age 60 and in women	The study did not include a control group The study was retrospective in nature
Rao [17]	Case-control study	No	100 subjects of which 75 were cases (25 patients with TIA, 25 patients with PVD, 25 patients with stroke) 25 controls aged 65 years or older	Peripheral vascular disease (PVD) patients did not perform significantly worse than control subjects on any of the neuropsychological tests The PVD group differed significantly from the stroke group on all neuropsychological tests except on aspects of frontal lobe function (verbal fluency F and BDCS) and memory; the TIA group only differed from the stroke group on the Trail-Making Test and verbal fluency 5	Small sample size Patients included in this study may have cognitive impairment potentially influencing neuropsychological performance.
Pohjasvaara [16]	Longitudinal follow-up study	15 months	286 patients	The general pattern of neuropsychological impairment appears similar in patients with TIA and in those with PVD, suggesting some similarity in etiological mechanisms TIA may result in more prolonged cognitive impairment, particularly in frontal lobe function. This group may be at increased risk of vascular dementia as well as impulsivity and suicide The frequency of suicidal ideas was found to be high at both 3 and 15 months of follow-up after stroke Patients with suicidal ideas were not more cognitively disabled in comparison with patients without suicidal ideas Patients with stroke have been shown to have an increased risk of suicide. The frequency of suicidal ideas increased with the time elapsed from stroke	Authors assessed only suicidal ideas instead of suicidal behavior, so they did not know whether someone attempted suicide or died by suicide

(continued)

Table 1 (Continued)

Study	Study design	Eventual follow-up	Participants	Main findings	Limitations
Chan et al. [21]	Case-control study	No	172 patients (cases were 84, controls were 74)	<p>Depression was the main correlate of suicidal ideas at 3 months after stroke, but at 15 months after stroke, suicidal ideas were also correlated with more disabling and right-sided stroke, especially in patients with a history of previous strokes.</p> <p>A higher CVRF score (derived by the weighted sum of severities of specific factors) was reported in cases (8.01 ± 6.46) than in controls (4.97 ± 3.81)</p> <p>Risk factors for cerebrovascular diseases confer higher risk for suicide in middle aged and older adults</p>	<p>Prevalence of affective disorders in the community controls was higher than the medium value previously reported, which was probably due to sampling bias</p> <p>Systematic misclassification of suicide because of different criteria adopted by judicial systems across different socio-cultural settings</p> <p>The cross-sectional nature of the sample precluded the causal interpretation of study variables. The authors did not know the information regarding treatment received or stroke's severity or when it occurred</p> <p>Sample was biased toward healthier individuals with stroke because of its exclusion of institutionalized individuals</p>
Fuller-Thomson et al. [19]	Case-control study	No	132,221 patients	<p>7.4% of patients with stroke were depressed, whereas only 5.2% were depressed without stroke</p> <p>Suicidal ideation: 15.2% in cases; 9.4% in controls</p> <p>After adjusting for socio-demographic factors, stroke survivors had twice the odds of depression (OR: 2.21) and suicidal ideation (OR: 2.07)</p> <p>After a reduction in functional limitation and activities in daily living limitations, both ORs (OR depression: 1.29; OR suicidal ideation: 1.37) decreased</p> <p>Functional limitations confer a higher risk of developing suicidal ideation</p> <p>Higher frequency of both depression and suicidal ideation in Caucasian and younger individuals</p>	<p>The cross-sectional nature of the sample precluded the causal interpretation of study variables. The authors did not know the information regarding treatment received or stroke's severity or when it occurred</p> <p>Sample was biased toward healthier individuals with stroke because of its exclusion of institutionalized individuals</p>
Santos et al. [18]	Cross-sectional study	No	177 (104 men, 73 women) acute stroke patients	<p>27 patients (15%) developed suicidal thoughts and 6 (22%) had a plan to die by suicide, but none completed the act. Among these, 23 (85%) were depressed, 13 (48%) had a previous mood disorder, and 5 (19%) had a recurrent stroke. 63</p>	<p>Lack of long-term follow-up data</p> <p>Small part of sample was probably not reliable because of cognitive impairment</p> <p>Young age of the sample limits the external validity of the results</p>

(continued)

Table 1 (Continued)

Study	Study design	Eventual follow-up	Participants	Main findings	Limitations
Garden et al. [26]	Case report study	No	2	<p>patients were depressed without suicidal ideation.</p> <p>After conducting a logistic regression analysis, suicidal thoughts were more frequent in patients with a lower educational level (OR: 5.13), previous mood disorder (OR: 4.22), diabetes and acute depression (OR: 7.82)</p> <p>Acute depression, previous mood disorder, low educational level are independent risk factors for suicidal thoughts in stroke patients</p> <p>Neither patient expressed suicidal thoughts, but both developed poststroke depression</p>	<p>Lack of imaging data that document the localization of lesions associated with suicidal thoughts</p> <p>Small sample size that did not allow findings to be generalized</p> <p>No data from any statistical analysis</p> <p>The study was retrospective in nature</p>

berg [23] concluded that the high risk period for suicide was 5 years after stroke; however, the authors did not consider the accelerated effect that depression has on suicidal behaviors. Conversely, Forsstrom *et al.* [24] recommend that suicide risk should be assessed and monitored not only in the long-term follow-up, but also during the 2 years immediately following stroke.

Moreover, two case reports [25,26] have investigated the association between stroke and suicide providing further support for this relation. Katayama *et al.* [25] evaluated 6 patients who attempted suicide within 6 months after stroke and hypothesized that these cases might have similar underlying organic processes contributing to both stroke and suicide attempts. The results revealed that four of the six patients suffered from depression and five of the six patients evidenced a moderate disability following stroke. In a second case study, retrospective analysis of medical records was examined from two patients who developed mood disorders in the acute poststroke period and eventually went on to die by suicide, be suicide. The researchers found that although neither patient openly expressed suicidal thoughts, both developed post-stroke depression, highlighting the importance of assessing for depressive symptoms and suicidality during the rehabilitation period following stroke [26]. In Table 1, we reported the findings from two additional studies [27–29] that are of particular interest, although they included mixed samples. Those studies did not directly investigate the role of stroke in suicidality but substantially confirmed the role of stroke as a risk factor for both attempted and completed suicide.

Discussion

The findings from the current review suggest that patients who have suffered a stroke are at an increased risk of suicide. Stroke patients commonly develop depression following their stroke, which significantly contributes to suicidal ideation, attempts, and completed suicide. Further, the existing literature suggests that patients should be closely monitored for depression and suicide during the 5 years following stroke. Regarding age and gender, suicide rates among stroke victims seemed to have increased in the younger population and decreased among the elderly as well as occurred more frequently in women [30,31]. However, given that stroke usually affects people age 65 and older, suicide victims who had suffered from a stroke were about 20 years older than suicide victims who had never suffered a stroke [3].

Although there were only two studies that indicated having a diagnosis of a previous mood disorder in general and prestroke depression in particular could be a clinical determinant for suicide, several studies considered in this review demonstrated that most patients who developed suicidal ideation and plans or died by suicide after a stroke were depressed. This finding suggests that assessing for a diagnosis of depression and/or depressive symptomatology is critical for preventing suicide in patients who have suffered a stroke as a result of disease. Also, these findings are in line with the study conducted by Arciniegas and Anderson [10] who reported that a suicide attempt and completed suicide were significantly associated with depression, feelings of hopelessness or helplessness, and social isolation in patients with neurological disorders.

The role of depression in suicide after stroke is an important issue given that the depression is the most common psychiatric complication after stroke and is highly associated with suicidality.

Kouwenhoven *et al.* [32], in fact, argued that poststroke depression (PSD) is one of the most prevalent affective disorders after stroke and occurs in at least one quarter of all stroke survivors during the first year. In this longitudinal study, the researchers described the lived experience as it develops over time among stroke survivors suffering from early depressive symptoms. Depression following stroke is a complex phenomenon and is not fully understood. The onset of depressive symptoms following stroke may be the result of a brain injury or a psychological reaction, which may complicate and delay stroke rehabilitation, contributing to poorer outcomes and increased functional dependence as compared to nondepressed stroke survivors. Furthermore, PSD may also become chronic in nature or even lead to death [28,30].

Generally, having an affective disorder is considered a major health problem and is often a burden impeding effective medical care [33]. Major depression has been identified as an independent stroke risk factor [34]. Also, subjects with major depression and bipolar disorder have a greater prevalence of established risk factors for vascular disease, including diabetes mellitus [35–38], obesity, [39–45], and less consistently hypertension [46–51].

Another major issue concerns adherence to treatment. Untreated depression (and particularly untreated bipolar disorder) is a significant predictor of negative outcome in patients having affective disorders. Medication nonadherence is a relevant obstacle to translating treatment efficacy from research settings into effectiveness in clinical practice [52]. Estimates of medication nonadherence for unipolar and bipolar disorders range from 10 to 60%; reduced adherence is associated with increased suicide risk. Extended courses of medication may be critical in improving the long-term outcome of patients with affective disorders [52]. Although effective treatment strategies exist, they are frequently misused in clinical practice; therefore, more efforts should be directed to improve treatment adherence among patients with major affective disorders to decrease patient symptoms burden, stroke risk, mortality and suicidal risk as well as to enhance poststroke healthcare utilization [52,53]. Early identification and treatment for depressive symptoms may significantly reduce suicidal risk in stroke patients [20].

Overall, the nature of the relationship between depression and stroke mortality is not well understood and many questions remain regarding prevention and treatment. Although depression after stroke has been associated with greater mortality, it is not entirely clear whether depression is a contributor to poststroke mortality as an individual risk factor or as an independent factor negatively influencing other stroke risk factors [54].

Other additional factors such as affective temperaments and personality dimensions may play a relevant role in determining the link between stroke and depression and contributing to suicide risk [55–58]. Personality disorders may have a marked impact on patients' interpersonal relationships, and social and occupational functioning leading to problematic interactions in the medical setting. The treatment for medical and psychiatric disorders is

more complicated in patients with comorbid personality disorders [55]. Moreover, affective temperaments may be useful tools in screening and identifying those affective disorder patients, particularly bipolar patients who are at higher risk of suicidal behavior [56,57]. An increase in behavioral and emotional disorders has also been observed after stroke in childhood. Daseking et al. [58] investigated 111 children aged 3–18 years who suffered a stroke and found that children may be at higher risk of behavioral (social and attention) and school-based problems related to stroke. Subjects with perinatal stroke and stroke in childhood had more behavioral and temperamental problems when compared to children with stroke in infancy. Further longitudinal research in this area is warranted to elucidate the relationship between depression and suicide in stroke victims.

This review has a number of limitations. First, no meta-analytic technique has been used to evaluate the results. Secondly, the authors chose to report those studies available in the literature that could support a broad analysis of the topic so as to offer a

tutorial study. Despite a careful and systematic search, we extrapolated those studies that presented original data; however, a number of additional articles could have been added as useful sources of information.

Conclusion

Stroke confers substantial risk for suicide, particularly in young adults and in women. The factors most frequently implicated in suicide among stroke victims are poststroke depression, previous mood disorder, prior history of stroke, and cognitive impairment. Given the frequency of stroke and poststroke depression, clinicians should assess and monitor depression in stroke patients, particularly during the rehabilitation period, to reduce suicide risk.

Conflict of Interest

The authors declare no conflict of interest.

References

- World Health Organization. *Atlas of heart disease and stroke*. Geneva, Switzerland: World Health Organization, 2004. Available from: <http://www.who.int/mediacentre/news/releases/2004/pr68/en/index.html> [Assessed 27 January, 2012].
- Mukherjee D, Patil CG. Epidemiology and the global burden of stroke. *World Neurology* 2011;**76**: S85–S90.
- Centers for Disease Control and Prevention. National Center for Chronic Disease Prevention and Health Promotion. The burden of chronic diseases and their risk factors. Section II: The burden of heart disease, stroke, cancer, and diabetes. United States. Online. 11 February 2002. Available from: http://www.cdc.gov/nccdphp/burdenbook2002/02_stroke.htm [Assessed 9 February, 2012].
- Conwell Y, Thompson C. Suicidal behavior in elders. *Psychiatr Clin North Am* 2008;**31**: 333–356.
- Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. Web-based Injury Statistics Query and Reporting System (WISQARS): Available from: <http://www.cdc.gov/ncipc/wisqars> [Assessed 13 January, 2012].
- Harris EC, Barraclough B. Suicide as an outcome for mental disorders. A meta-analysis. *Br J Psychiatry* 1997;**170**: 205–228.
- Juurink DN, Herrmann N, Szalai JP, Kopp A, Redelmeier DA. Medical illness and the risk of suicide in the elderly. *Arch Intern Med* 2004;**164**: 1179–1184.
- Quan H, Arboleda-Flórez J, Fick GH, Stuart HL, Love EJ. Association between physical illness and suicide among the elderly. *Soc Psychiatry Psychiatr Epidemiol* 2002;**37**: 190–197.
- Mackenzie TB, Popkin MK. Suicide in medical patient. *Int J Psychiatry Med* 1987;**17**: 3–22.
- Arciniegas DB, Anderson CA. Suicide in neurologic illness. *Curr Treat Options Neurol* 2002;**4**: 457–468.
- Hadidi N, Treat-Jacobson DJ, Lindquist R. Poststroke depression and functional outcome: a critical review of literature. *Heart Lung* 2009;**38**: 151–162.
- Fleminger S, Oliver DL, Williams WH, Evans J. The neuropsychiatry of depression after brain injury. *Neuropsychol Rehabil* 2003;**13**: 65–87.
- Beekman AT, Penninx BW, Deeg DJ, Ormel J, Smit JH, Braam AW, van Tilburg W. Depression in survivor of stroke: a community-based study of prevalence, risk factors and consequences. *Soc Psychiatry Psychiatr Epidemiol* 1998;**33**: 463–470.
- Kishi Y, Kosier JT, Robinson RG. Suicidal plans in patients with acute stroke. *J Nerv Ment Dis* 1996a;**184**: 274–280.
- Kishi Y, Robinson RG, Kosier JT. Suicidal plans in patients with stroke: comparison between acute-onset and delayed-onset suicidal plans. *Int Psychogeriatr* 1996b;**8**: 623–634.
- Pohjasvaara T, Vataja R, Leppävuori A, Kaste M, Erkinjuntti T. Suicidal ideas in stroke patients 3 and 15 months after stroke. *Cerebrovasc Dis* 2001;**12**: 21–26.
- Rao R, Jackson S, Howard R. Neuropsychological impairment in stroke, carotid stenosis, and peripheral vascular disease, a comparison with healthy community residents. *Stroke* 1999;**30**: 2167–2173.
- Santos CO, Caeiro L, Ferro JM, Figueira ML. A study of suicidal thoughts in acute stroke patients. *J Stroke Cerebrovasc Dis*. 2011. DOI: 10.1016/j.jstrokecerebrovasdis.2011.04.001 (in press).
- Fuller-Thomson E, Tulipano MJ, Song M. The association between depression, suicidal ideation, and stroke in a population-based sample. *Int J Stroke* 2012;**7**: 188–194.
- Tang WK, Lu JY, Mok V, Ungvari GS, Wong KS. Is fatigue associated with suicidality in stroke? *Arch Phys Med Rehabil* 2011;**92**: 1336–1338.
- Chan SS, Lyness JM, Conwell Y. Do cerebrovascular factors confer risk for suicide in later life? A case-control study. *Am J Geriatr Psychiatry* 2007;**15**: 541–544.
- Stenager EN, Madsen C, Stenager E, Boldsen J. Suicide in patients with stroke: epidemiological study. *BMJ* 1998;**316**: 1206.
- Teasdale TW, Engberg AW. Suicide after a stroke: a population study. *J Epidemiol Community Health* 2001;**55**: 863–866.
- Forsström E, Hakko H, Nordström T, Räsänen P, Mainio A. Suicide in patients with stroke: a population-based study of suicide victims during the years 1988–2007 in northern Finland. *J Neuropsychiatry Clin Neurosci* 2010;**22**: 182–187.
- Katayama M, Naritomi H, Oomura M, et al. Case reports of unexpected suicides in patients within six months after stroke. *Kobe J Med Sci* 2010;**56**: 184–194.
- Garden FH, Garrison SJ, Jain A. Assessing suicide risk in stroke patients: review of two cases. *Arch Phys Med Rehabil* 1990;**71**: 1003–1005.
- Scott KM, Hwang I, Chiu WT, et al. Chronic physical conditions and their association with first onset of suicidal behavior in the world mental health surveys. *Psychosom Med* 2010;**72**: 712–719.
- Voaklander DC, Rowe BH, Dryden DM, Pahal J, Saar P, Kelly KD. Medical illness, medication use and suicide in seniors: a population-based case-control study. *J Epidemiol Community Health* 2008;**62**: 138–146.
- Placido A, Sposito AC. Association between suicide and cardiovascular disease: time series of 27 years. *Int J Cardiol* 2009;**135**: 261–262.
- Hawton K, van Heeringen K. Suicide. *Lancet* 2009;**373**: 1372–1381.
- Ellis C, Zhao Y, Egede L. Depression and increased risk of death in adults with stroke. *J Psychosom Res* 2010;**68**: 545–551.
- Kouwenhoven SE, Kirkevold M, Engedal K, Biong S, Kim HS. The lived experience of stroke survivors with early depressive symptoms: a longitudinal perspective. *Int J Qual Stud Health Well-being* 2011;**6**: 8491–8504.
- Saravane D, Feve B, Frances Y, et al. [Drawing up guidelines for the attendance of physical health of patients with severe mental illness]. *Encephale* 2009;**35**: 330–339.
- Williams LS. Depression and stroke: cause or consequence? *Semin Neurol* 2005a;**25**: 396–409.
- Cassidy F, Ahearn E, Carroll BJ. Elevated frequency of diabetes mellitus in hospitalized manic depressive patients. *Am J Psychiatry* 1999;**156**: 1417–1420.
- Kilbourne AM, Cornelius JR, Han X, et al. Burden of general medical conditions among individuals with bipolar disorder. *Bipolar Disord* 2004;**6**: 368–373.
- Birkenaes AB, Opjordsmoen S, Brunborg C, et al. The level of cardiovascular risk factors in bipolar disorder equals that of schizophrenia: a comparative study. *J Clin Psychiatry* 2007a;**68**: 917–923.
- Golden SH, Lazo M, Carnethon M, et al. Examining a bidirectional association between depressive symptoms and diabetes. *JAMA* 2008;**299**: 2751–2759.
- Eaton WW, Armenian H, Gallo J, Pratt L, Ford DE. Depression and risk for onset of type II diabetes. A prospective population-based study. *Diabetes Care* 1996;**19**: 1097–1102.
- Mezuk B, Eaton WW, Golden SH, Ding Y. The influence of educational attainment on depression and risk of type 2 diabetes. *Am J Public Health* 2008;**98**: 1480–1485.
- McElroy SL, Frye MA, Suppes T. Correlates of overweight and obesity in 644 patients with bipolar disorder. *J Clin Psychiatry* 2002;**63**: 207–213.

42. Fagiolini A, Frank E, Scott JA, Turkin S, Kupfer DJ. Metabolic syndrome in bipolar disorder: findings from the Bipolar Disorder Center for Pennsylvanians. *Bipolar Disord* 2005;**7**: 424–430.
43. Shah A, Shen N, El-Mallakh RS. Weight gain occurs after onset of bipolar illness in overweight bipolar patients. *Ann Clin Psychiatry* 2006;**18**: 239–241.
44. Fiedorowicz JG, Palagummi NM, Forman-Hoffman VL, Miller del D, Haynes WG. Elevated prevalence of obesity, metabolic syndrome, and cardiovascular risk factors in bipolar disorder. *Ann Clin Psychiatry* 2008;**20**: 131–137.
45. Birkenaes AB, Opjordsmoen S, Brunborg C, et al. The level of cardiovascular risk factors in bipolar disorder equals that of schizophrenia: a comparative study. *J Clin Psychiatry* 2007b;**68**: 917–923.
46. Birkenaes AB, Opjordsmoen S, Brunborg C, et al. The level of cardiovascular risk factors in bipolar disorder equals that of schizophrenia: a comparative study. *J Clin Psychiatry* 2007c;**68**: 917–923.
47. Yates WR, Wallace R. Cardiovascular risk factors in affective disorder. *J Affect Disord* 1987;**12**: 129–134.
48. Johannessen L, Strudsholm U, Foldager L, Munk-Jorgensen P. Increased risk of hypertension in patients with bipolar disorder and patients with anxiety compared to background population and patients with schizophrenia. *J Affect Disord* 2006;**95**: 13–17.
49. Patten SB, Williams JV, Lavorato DH, Modgill G, Jette N, Eliasziw M. Major depression as a risk factor for chronic disease incidence: longitudinal analyses in a general population cohort. *Gen Hosp Psychiatry* 2008;**30**: 407–413.
50. Davis J, Fujimoto RY, Juarez DT, Hodges KA, Asam JK. Major depression associated with rates of cardiovascular disease state transitions. *Am J Manag Care* 2008;**14**: 125–128.
51. Meyer CM, Armenian HK, Eaton WW, Ford DE. Incident hypertension associated with depression in the Baltimore Epidemiologic Catchment area follow-up study. *J Affect Disord* 2004;**83**: 127–133.
52. Pompili M, Serafini G, Del Casale A, et al. Improving adherence in mood disorders: the struggle against relapse, recurrence and suicide risk. *Expert Rev Neurother* 2009;**9**: 985–1004.
53. Williams LS. Depression and stroke: cause or consequence? *Semin Neurol* 2005b;**25**: 396–409.
54. Prithard C, Hansen L. Comparison of suicide in people aged 65–74 and 75 + by gender in England and Wales and the major Western countries 1979–1999. *Int J Geriatr Psychiatry* 2005;**20**: 17–25.
55. Pompili M, Ruberto A, Girardi P, Tatarelli R. Suicidality in DSM IV cluster B personality disorders. An overview. *Ann Ist Super Sanita* 2004;**40**: 475–483.
56. Pompili M, Innamorati M, Rihmer Z, et al. Cyclothymic-depressive-anxious temperament pattern is related to suicide risk in 346 patients with major mood disorders. *J Affect Disord* 2012a;**136**: 405–411.
57. Pompili M, Rihmer Z, Akiskal H, et al. Temperaments mediate suicide risk and psychopathology among patients with bipolar disorders. *Compr Psychiatry* 2012b;**53**: 280–285.
58. Daseking M, Petermann F, Simonis A. Behavioral disorders and psycho-social outcome after stroke in children]. *Fortschr Neurol Psychiatr* 2008;**76**: 662–671.