



HHS Public Access

Author manuscript

J Community Health. Author manuscript; available in PMC 2020 June 01.

Published in final edited form as:

J Community Health. 2019 June ; 44(3): 451–462. doi:10.1007/s10900-018-00612-7.

The association between perceived stress and hypertension among Asian Americans: does social support and social network make a difference?

Xiaoxiao Lu, PhD, MD, MPH¹, Hee-Soon Juon, PhD², Xin He, PhD¹, Cher M Dallal, PhD¹, Ming Qi Wang, PhD³, and Sunmin Lee, ScD¹

¹Department of Epidemiology and Biostatistics, University of Maryland School of Public Health, College Park, Maryland ²Department of Medical Oncology, Thomas Jefferson University, Philadelphia, Pennsylvania ³Department of Behavioral and Community Health, University of Maryland School of Public Health, College Park, Maryland

Abstract

Background: Prior research suggests that stress plays role in the etiology and progression of hypertension. To lend a more accurate depiction of the underlying mechanisms between stress and hypertension, this study aims to assess the associations between perceived stress and hypertension across varying levels of social support and social network among Asian Americans.

Methods: We conducted a cross-sectional study using data on 530 Chinese, Korean and Vietnamese Americans recruited from a liver cancer prevention program in the Washington D.C. – Baltimore metropolitan area.

Results: Hypertension prevalence was 29.1%. Individuals with high perceived stress were 61% more likely to have hypertension compared to those with low levels of perceived stress (Odds Ratio (OR): 1.61, 95% Confidence interval (CI): 1.15, 2.46). There was no evidence that social support and social network acted as effect modifiers. Social support had a direct beneficial effect on hypertension, irrespective of whether individuals were under stress. The relationship between perceived stress and hypertension was modified by gender and ethnicity whereby a significant positive association was only observed among male or Chinese participants.

Conclusion: Our study highlights the importance of understanding the associations between stress, social support, and hypertension among Asian American subgroups. Findings from the study can be used to develop future stress management interventions, and incorporate culturally and linguistically appropriate strategies into community outreach and education to decrease hypertension risk within the Asian population.

Corresponding author: Xiaoxiao Lu, PhD, MD, MPH, Department of Epidemiology and Biostatistics, University of Maryland School of Public Health, 2242 Valley Drive, College Park, MD 20742, Tel: 202-412-7885, xxluumd@gmail.com.

Conflict of interest

Xiaoxiao Lu, Hee-Soon Juon, Xin He, Cher M Dallal, Ming Qi Wang, and Sunmin Lee declare that they have no conflict of interest.

Keywords

Hypertension; Asian American; Stress; Social support; Social network

Introduction

Hypertension is a major risk factor for the development of cardiovascular disease (CVD), and a main cause of morbidity and mortality worldwide (1). Asian Americans, a rapidly growing minority population in the U.S., have been found to have a high prevalence of hypertension. According to data from the National Health and Nutrition Examination Survey (NHANES), about one-quarter (25.6%) of Asian Americans 20 years of age and older had hypertension during 2011-2012 (2). While there is a lack of national studies that observe the prevalence of hypertension in Asian American subgroups, local data reveals that significant differences in hypertension prevalence across various Asian American subgroups (3-5).

Emerging evidence indicates that chronic exposure to psychological stress/stressors plays a central role in the pathophysiology of hypertension and CVD (6, 7). Stress is a complex process and can influence the pathogenesis of physical disease by exerting direct effects on biological processes or indirect effects on behavioral patterns (8). When stress exceeds adaptive capacity, it results in maladaptive processes that negatively impact cardiovascular health (9). Prior evidence suggests that perceived stress contributes to the elevated blood pressure and to the development of hypertension and CVD (10, 11).

Regarding social relationships, a growing body of literature indicates that social support and social isolation can have a direct effect on hypertension (12-16). Prior research suggests that a low level of social support is a risk factor for the development of CVD in healthy individuals (17-19). In terms of social network, several studies have shown that a smaller social network is associated with higher blood pressure levels (20-22). Social isolation limits the amount of interactions and support that this population needs in order to maintain a healthy lifestyle. Failure to comply with certain medical recommendations because of social isolation could increase blood pressure in stressful situations. To the contrary, a large social network and strong social support may attenuate the negative cardiovascular response in situations of stress, thereby avoiding the increase in blood pressure (23).

Prior research has examined the mechanisms of social support and social network that underlie the association between stress and hypertension. In 1985, Cohen and Wills introduced two mechanisms in which social support may contribute to health: the Main Effect Hypothesis and the Stress-Buffering Hypothesis (24). The Main Effect Hypothesis proposes that social resources have a beneficial health effect by providing positive experience and stability in life situation, irrespective of whether persons are under stress or not (15, 24). The expression of negative feelings, for example, could lead to positive physiological, and concurrent, immunological benefits. The Stress-Buffering Hypothesis predicts that social support will have stronger positive effects on adjustment and physical well-being when a stressor becomes more intense or persistent (24). Unfortunately, there is a lack of studies of the inter-relationship between perceived stress, social support, social network, and hypertension, particularly among Asian Americans.

To bridge the missing gaps in previous literature, this study aims to: (1) assess the relationship between perceived stress and hypertension among Chinese, Korean and Vietnamese Americans in the Washington, DC metropolitan area; (2) test the main effect hypothesis by examining the direct effects of social support and social network on hypertension; and (3) test the stress-buffering hypothesis by assessing the relationship between perceived stress and hypertension across varying levels of social support and social network. Given the high prevalence of hypertension among Asian Americans, utilizing multiple psychosocial measures allows us to capture the differences in the associations between various psychosocial aspects and hypertension, and to better understand the contributing psychosocial factors to hypertension.

Methods

Study Population and Recruitment Procedure

We used the baseline survey data collected for the Asian American Liver Cancer Prevention Program (a randomized controlled trial to increase hepatitis B vaccination) for this ancillary study (25). To examine the relationship between perceived stress and hypertension, foreign-born Asian American adults, aged 18 years and older, were recruited from the community in the Baltimore-Washington Metropolitan Area. After obtaining informed consent, all the participants were asked to complete a self-administered questionnaire in English, Chinese, Korean, or Vietnamese, with the assistance of a bilingual interviewer when necessary. While completing the survey, blood pressure was measured three times by the research team at the study site. This study was approved by the Institutional Review Board at Johns Hopkins Bloomberg School of Public Health.

Of the 645 eligible volunteers, 30 did not participate in the program and 15 of those who attended the program did not complete either the survey or the hepatitis B screening. Thus, a total of 600 participants (201 Chinese, 198 Korean, and 201 Vietnamese) completed the screening and survey. We excluded 70 participants who had missing values on the following variables: social support (19), stress (32), social network index (11), education (6), employment (3), body mass index (BMI) (3), smoke (7), binge drinking (7), and self-rated identity (3). The final sample included 530 subjects for descriptive and regression analyses. Compared to the excluded sample ($N = 70$), women remaining ($N = 530$) in the analysis sample tended to be more educated, employed, self-identified very Asian, and more likely to smoke. We addressed this potential selection bias in the discussion section (see Table 6 in Appendix).

Measures

Dependent variable—While participants were seated, three readings of blood pressure were taken at 5-minute intervals in the right arm using the OMRON HEM 907 blood pressure monitor. Based on the American Heart Association (AHA)'s recommendations (26), the mean of the latter two readings were used in the analysis. The following criteria were used to define hypertension in this study: 1. systolic blood pressure (SBP) of 140 mmHg or higher or diastolic blood pressure (DBP) of 90 mmHg or higher; or 2. use of antihypertensive drugs. Participants were considered to be aware of their hypertension status

if they answered yes to the question “Have you been told by a doctor or other healthcare professional that you have hypertension or high blood pressure?”.

Independent variable

Perceived Stress.: Perceived stress was measured using the 10-item version of Perceived Stress Scale (PSS) which captures how unpredictable, uncontrollable, and overloaded respondents find their lives (8). Response categories are based off a 5-point Likert scale ranging from never (0) to very often (4) (27). PSS scores can range from 0 to 40, with a higher score indicates more perceived stress. Cronbach’s alpha for the scale with our data is 0.77. We categorized the PSS score based on the median into low (0-15) and high (16 and above) groups.

Social Support.: Social support was assessed using the 8-item version of the Duke-UNC Functional Social Support Questionnaire (DUFSS) which measures the amount and type of perceived functional social support. The 8-item version of the DUFSS includes two subscales: 6-item confidant support (including having someone to talk to, trust with, socialize with, receive advice from, and receive help from) and 2-item-affective support (being shown love and affection). The responses are on 5 point Likert scale, ranging from 5 (as much as I would like) to 1 (much less than I would like). The scale yields a single total support score, ranging between 8 and 40. The higher total score reflects a higher perceived social support. Cronbach’s alpha for this scale from our data is 0.94. We categorized the DUFSS score based on the median into low (0-30) and high (31 and above) groups.

Social Network Index Score.: Social network index was measured using Berkeman and Syme’s Social Network Index (SNI). This measure is a composite measure of four types of social connections: marital status (married vs. not); sociability (number and frequency with close relatives and close friends); church membership (yes vs. no); and membership in other community organizations (yes vs. no) (28). The SNI allows researchers to categorize individuals into four levels of social connection: socially isolated, moderately isolated, moderately integrated, and socially integrated (28). The SNI considers both the number and the relative importance of social ties across these four network categories and combines this information into a single summary measure (ranging from 0 to 4). Four network categories were developed to reflect differences in type and extent of social contact: married (no = 0; yes = 1); close friends and relatives (0-2 friends and 0-2 relatives (including children) = 0; all other scores = 1); group participation (no = 0; yes = 1); and participation in religious meetings or services (never or almost never = 0; greater than or equal to once or twice a month = 1). Scores were summed: 0 or 1 being the most isolated category (socially isolated); 2 (moderately isolated), 3 (moderately integrated), or 4 (socially integrated) formed the other three categories of increasing social connectedness. The procedures by which this index was developed and the precise description of methods used to score it are available elsewhere (28-30).

Covariates.: Based on existing literature, the following confounders were included in the analysis: age, gender, ethnicity, education, employment status, marital status, body mass index (BMI), smoking, binge drinking, self-rated identity and English speaking. We

obtained information on these variables from the self-administered questionnaire mentioned previously. Since age was not linearly related to the outcome based on LOWESS analysis (31, 32), age was categorized into 3 groups: less than 40 years, 40-59 years, and 60 years and over. With regards to education, we categorized participants into three groups: less than high school, high school or some college, college graduate or higher. We collapsed the marital status variable into two categories: married/living with a partner, or others (separated/divorced/widowed/never married). Employment status was dichotomous: employed vs unemployed. BMI was assessed as a continuous variable using anthropometric measure of weight and height, and was defined as body weight (in kilograms) divided by the square of height (in meters). Smoking was grouped as smoke (had smoked at least 100 cigarettes in entire life) vs non-smoke. Binge drinking status was categorized as yes (women had 4 or more drinks on an occasion during the past 30 days; men had 4 or more drinks on an occasion during the past 30 days) vs no (reference level). For the acculturation variables, English proficiency was measured as fluent, well, so-so, poorly, or not at all. Due to the low cell counts, we collapsed this variable into three groups: native/very well, so-so, and poorly/not at all. Self-rated identity was assessed using a single item measure on a 5-point scale: very Asian, mostly Asian, bicultural, mostly westernized, and very westernized. The last three self-rated identity categories were combined into one group: bicultural/westernized.

Statistical Analysis

Descriptive statistics calculated included estimating means of continuous variables and proportions of categorical variables. Hypertension status was compared by acculturation and covariates using the t-test for continuous variables and the Pearson's chi-squared test for categorical variables. The statistical significance was based on a p value less than 0.05.

Age-adjusted logistic regression and multiple logistic regression were performed to assess the association between each psychological measure and hypertension. Potential moderation by social support, social network index, age categories, gender, education, employment status, self-identity, English proficiency, obesity, smoking and binge drinking were examined by including the corresponding interaction term. The potential moderators were selected based on prior literature. The stratified analysis was conducted if the interaction term was significant. The statistical analysis of the study was performed using SAS 9.4 (SAS Institute, Cary, NC).

Results

Participant characteristics are presented in Table 1. Of the 530 participants, the majority (63%) were between age of 40 and 59 years. About 58% were female and most of the participants were married (78%). Chinese, Korean, and Vietnamese ethnic composition was about the same, each making up about one third of the sample. About half of the study population received a college education or higher while 18% reported a history of smoking and 17% were binge drinkers. Based on the objective measure of BMI, 31% of the participants were overweight ($BMI > 25 \text{ kg/m}^2$).

The mean of the perceived stress score in our sample was 15.5 (SD = 5.8). The level of perceived stress varied with ethnicity groups, English proficiency, and social support. A larger proportion of Korean Americans reported high perceived stress compared to Chinese and Vietnamese Americans (64% vs 48% and 45%, $p < 0.01$). In addition, those with high level of perceived stress tended to receive less social support and rate their English fluency as “poorly/not at all” or “so-so”. In regards to social support, 53% of the participants reported low level of social support. The level of social support also differed by gender and English proficiency. Females and individuals who spoke English well tended to receive a high level of social support. For the social network index score, 13% were socially isolated, 22% were moderately isolated, 30% were moderately integrated, and 35% were socially integrated. Socially isolated individuals tended to be less than 40 years or older than 60 years, not married, more Korean than Chinese or Vietnamese Americans, lower educated, not employed, and rated their English fluency as “poorly/not at all”.

The overall prevalence of hypertension in this population was 29%. As shown in Table 1, those with hypertension tended to be older, male, less educated, higher BMI, rate their English fluency as “poorly/not at all” or “so-so”, and have a history of smoking compared to those without hypertension. Perceived stress and social support were significantly associated with hypertension in the bivariate analyses.

Table 2 shows age-adjusted models for the association between psychological factors and hypertension. Individuals who had high level of perceived stress were 71% more likely to have hypertension as compared to those with low level of perceived stress (OR: 1.71, 95% CI: 1.15, 2.54) after adjusting for age. Participants with high social support were 51% less likely to have hypertension comparing with those with low social support (OR: 0.49, 95% CI: 0.33, 0.74). However, social network index score was not significantly associated with hypertension after controlling for age.

Table 3 reports multivariable adjusted models for the association between psychological factors and hypertension. Age and BMI had the largest confounding effect among all covariates. After adjusting for other variables, individuals who had a high level of perceived stress were 61% more likely to have hypertension as compared to those with low level of perceived stress (OR: 1.61, 95% CI: 1.15, 2.46). Social support also had a direct impact to hypertension: participants with high social support were 48% less likely to have hypertension comparing with those with low social support (OR: 0.52, 95% CI: 0.33, 0.81). However, social network index score was not significantly associated with hypertension in general. Moderately integrated individuals were 58% less likely to have hypertension than socially isolated individuals (OR: 0.42, 95% CI: 0.19, 0.92). Interactions between perceived stress and social support or social network index were not statistically significant.

There was a significant interaction between perceived stress and gender (Table 4), suggesting that perceived stress had a stronger association with hypertension in men than in women. Male participants with high level of perceived stress were 95% more likely to have hypertension compared with those with low level of perceived stress (OR: 1.95, 95% CI: 1.05, 3.60). This association was not significant in women.

The relationships between perceived stress and hypertension also varied by three ethnic groups (Table 5). The adjusted model suggests that Chinese participants who had a high level of perceived stress were 278% more likely to have hypertension compared to those with low level of perceived stress (OR: 3.78, 95% CI: 1.13, 6.80). Similar to perceived stress, Chinese participants with high social support were 64% less likely to have hypertension as compared to those who had low social support (OR: 0.36, 95% CI: 0.15, 0.87). Among Korean and Vietnamese groups, no significant difference in hypertension status was found for various psychological measures.

Discussion

Our study is one of the first to examine multiple measures of psychosocial factors in relation to hypertension among Asian Americans overall and by ethnicity. The findings in this study clearly indicate that perceived stress is a strong predictor of hypertension. Our results support the main effect hypothesis that social support had a direct beneficial effect on hypertension, irrespective of whether persons were under stress. However, a stress-buffering effect of social support or social network was not demonstrated in this study. Our findings also suggest the relationship between perceived stress and hypertension was modified by gender. Among Asian American men, those with high level of perceived stress were significantly more likely to develop hypertension compared to those with low level of perceived stress. For women, no association between perceived stress and hypertension was found. In addition, the impact of perceived stress on hypertension may differ by Asian American subgroups as the significant association between perceived stress and hypertension was only observed among Chinese Americans.

The prevalence of hypertension was found to be slightly higher among Asian Americans in our sample (29.1%) compared to the data of NHANES 2011-2012 (25.6%) (2). This is probably because our sample was mostly comprised of a foreign-born population. According to the NHANES data, the prevalence of hypertension was higher among foreign-born (26.1%) individuals compared to those U.S.- born (21.2%) (2). Regarding the disaggregated data, consistent with a study that observed differences in hypertension prevalence among Asian subgroups (5), we found some variations in hypertension prevalence: hypertension prevalence was highest among Vietnamese Americans (33%), followed by Korean (31%) and Chinese Americans (23%).

Another notable finding is the level of perceived stress observed among Asian Americans in this study. The mean of the perceived stress score in our sample was 15.5, which is slightly higher than the mean score reported among Whites in the study by Cohen et al. (15.2) (33). Prior evidence has suggested that immigrant population experienced more psychosocial stress compared to dominant western cultural groups (34). Studies have shown that most common difficulties which immigrants experience include language barriers, adjusting to an unfamiliar environment, differences between Asian and American cultures, and problems relating to children who are often more acculturated than their parents (35). Our study also found that perceived stress was strongly related to acculturation whereby those with perceived low stress tended to rate English as “native fluency/well” or “so-so” (37% native fluency/well vs 35% so-so vs 29% poorly/not at all, $p < 0.01$). In addition, our data suggests

that a larger proportion of Korean Americans reported high perceived stress relative to Chinese and Vietnamese Americans (64% vs 48% and 45%, $p < 0.01$). This finding is consistent with previous studies that reveal high levels of psychosocial stress, anxiety, and depression in Korean Americans (36-38). Potential explanatory factors may include higher rates of under-employment (a better education but lower-prestige jobs); limited ability in English; and a higher concentration in small business located in high-risk minority districts (39).

A strong relationship between perceived stress and hypertension among Asian Americans was concluded in our study. Prior research connecting stress with hypertension has produced mixed findings, with some studies identifying a positive association (40, 41), some showing no correlation (37) while others even demonstrating a negative correlation (42). However, few of these studies were conducted among Asian Americans (37). Logan et al. explored the relationship between perceived stresses and blood pressure among 102 Korean Americans in North Carolina (37). They found that there was no significant association between perceived stress and SBP/DBP.

The mechanisms by which stress may be linked to hypertension are complex and involve a variety of interrelated physiological and behavioral pathways (43). Early studies have shown that the primary biological pathway linking emotions to disease is hormone (44). Excessive discharge of certain hormones have been implicated in the pathogenesis of cardiovascular disease and diseases involving the immune system including cancer, infectious diseases, and autoimmune diseases (44). Behavioral changes may also occur as adaptations to stressors. People exposed to stressors or with perceived stress tend to engage in poor health behaviors, such as smoking, alcohol consumption, inactivity, unhealthy diets, and poor sleep patterns (27, 45). Additionally, it is believed that immigrants change their lifestyle to cope with the increased stress (46). These lifestyle changes, such as unhealthy diet, have been found to increase vulnerability to developing hypertension (46, 47). Therefore, well-established behavioral risk factors may represent an intermediate stage through which stress increases hypertension risk (48). However, the established risk factors do not account completely for the stress and CVD association. Consistent with earlier studies (49), our study found adjusting for intermediate behavioral health indicators (smoking, binge drinking and BMI) resulted in only a small change in the strength of associations of perceived stress with hypertension.

We found mixed evidence regarding the effects of social relationship on hypertension. In this analysis, social support was significantly associated with hypertension. Similar findings have been reported from Tomaka et al (50). They examined relations between social support and health outcomes in a senior sample from New Mexico. The results show that the odds of hypertension were 24% lower for a one unit increase in the social support score (OR: 0.76, $p < 0.05$). Inconsistent with prior studies (20, 51, 52), we found there was no significant association between social network and hypertension. In addition, we observed no difference in the associations between perceived stress and hypertension across various levels of social support or social network. This suggests that the impact of the subjective experience of stress on hypertension was not moderated by social network or social support. But the stress-buffering effect of social support was demonstrated in other studies (53, 54). The findings

from these studies highlight the importance of examining the joint contribution of stress and coping resources to hypertension.

The gender discrepancies in the association between stress and hypertension found in our study is consistent with prior research (30, 55). Our results revealed that Asian American men with a high level of perceived stress were significantly more likely to develop hypertension compared to those with a low level of perceived stress. Interestingly enough, there was no association between perceived stress and hypertension among women. The mechanism for these gender differences of stress-related hypertension remains unclear. It has been reported that Asian men experience a more difficult time in adjusting to new cultures compared with Asian women (55). Asian culture had instilled in these men more authority that the impact of stress would be more profound (55). In addition, prior research has observed gender differences in the use of stress coping strategies (56). Women preferred the emotion-focused coping strategy to mobilize their social networks, especially peers, to talk about what was troubling them as a way of releasing stress; while men usually repressed their emotions to either fight or escape (57). Although this stress-buffering effect was not observed in our data, social support has been demonstrated as an effect modifier in other research (53, 54).

Another noteworthy finding is the heterogeneity among Asian subgroups in the associations between psychosocial factors and hypertension. Stratified analyses concluded that the significant associations between perceived stress, social support and hypertension only presented among Chinese participants. This suggests that the negative effect of perceived stress and positive effect of social support on hypertension may be particularly potent among Chinese Americans. Chinese culture emphasizes interdependence and values group consensus more than individual attributes (35). In the current study, we also found the mean of social support score was higher among Chinese (30.2) than Korean (28.1) and Vietnamese Americans (29.0). Future study may elucidate the underlying mechanisms in the heterogeneous associations among Asian subgroups by examining the culturally relevant stress and social support measures.

We acknowledge a number of limitations in this study. First, our study was based on cross-sectional data and does not lend itself to causal inference. Future studies using longitudinal data are needed to corroborate our findings. Second, the study used non-probability sampling methods because the target population is a hard-to-reach population; however, our sample composition closely follows the composition of the United States 2010 Census data (58). Therefore, generalizability of the study may not be largely compromised. Third, the majority of our study population was first generation immigrants because the goal of the parent study was to study hepatitis B. Thus, variability for perceived stress and social support might have been smaller than studies that included more US-born Asian Americans. This might have reduced our ability in detect potential associations between psychosocial factors and hypertension. Fourth, we did not have data on a number of established risk factors for hypertension that are also related to stress. Physical activity and diet are known to be associated with the risk of hypertension (59, 60), but these information were not available in our data.

Nevertheless, this study advanced our understanding of psychosocial factors and hypertension among Asian Americans. To our best knowledge, this is one of the first studies examining the synergistic effect of perceived stress, functional social support, and structural social support on hypertension among Asian Americans. By using a large sample size, this study focused on three of the largest Asian American populations: Chinese, Koreans, and Vietnamese. This highlights the importance of identifying differences in health effects of perceived stress among disaggregated Asian Americans subgroups. Moreover, as opposed to using self-reported hypertension information, our study used objective measure of hypertension as our outcome. Three blood pressure readings were taken at the time of the survey for increased accuracy. There is ample evidence in previous literature of large disparities between the objective and subjective health measures in even very high quality surveys (61).

The implication of this work is directed towards expanding our understanding of the interrelationship between stress, social support, and hypertension among Asian Americans. It is critical to this discussion that researchers and clinicians understand the fundamental cultural differences that Asian Americans have relative to the general population. At the same time, it is also essential to recognize the diversity that exists within the Asian American culture. Because of this, our study underlines the importance of identifying differences in health effects of stress among disaggregated Asian American subgroups in order to help health professionals prioritize which subgroups need the most urgent intervention in terms of stress management. Successful interventions to reduce the modifiable risk factors among individuals under stress may prevent the risk of long-term health hazards. In general, findings from the study can be used to develop future stress management interventions, and incorporate culturally and linguistically appropriate strategies into community outreach and education to decrease hypertension risk within the Asian Americans.

Acknowledgement

This study was funded by the National Cancer Institute (R01CA163805; PI: Juon), AND Center for Strategic Scientific Initiatives, National Cancer Institute (5P30CA056036-17). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Appendix

Table 6.

Compare characteristics between analysis sample and excluded sample with missing values

	Analysis sample (n = 530)	Excluded sample (n = 70)	P-Value
Age			0.10
< 40 years	126 (23.8)	10 (14.3)	
40–59 years	335 (63.2)	46 (65.7)	
60 years and over	69 (13.0)	14 (20.0)	
Gender			0.72
Male	224 (42.3)	28 (40.0)	

	Analysis sample (n = 530)	Excluded sample (n = 70)	P-Value
Female	306 (57.7)	42 (60.0)	
Ethnicity			0.49
Chinese	177 (33.4)	24 (34.3)	
Korean	179 (33.8)	19 (27.1)	
Vietnamese	174 (33.8)	27 (38.6)	
BMI (mean[SD])	23.4 (3.4)		
Marital status			
Married	411 (77.6)		
Not married	119 (22.4)		
Education			<0.01
Less than high school	68 (12.8)	18 (25.7)	
High school or some college+	202 (38.1)	21 (30.0)	
College graduate +	260 (49.1)	25 (35.7)	
Missing	0	6 (8.6)	
Employment status			<0.01
Employed	356 (67.2)	42 (60.0)	
Unemployed	174 (32.8)	25 (35.7)	
Missing	0	3 (4.3)	
Self-rated Identity			<0.01
Very Asian	304 (57.2)	37 (52.9)	
Mostly Asian	88 (16.5)	11 (15.7)	
Bicultural/Westernized	140 (26.3)	19 (27.1)	
Missing		3 (4.3)	
English speaking			0.31
Poorly/not at all	194 (36.6)	31 (44.3)	
So-so	207 (39.1)	27 (38.6)	
Native/very well	129 (24.3)	12 (17.1)	
Missing	0		
Smoke			<0.01
Smoker	94 (17.7)	10 (14.3)	
Non-smoker	438 (82.3)	54 (77.1)	
Missing	0	6 (8.6)	
Binge drink			0.53
Yes	67 (12.6)	63 (90.0)	
No	463 (82.4)	7 (10.0)	

Reference

1. Nwankwo T, Yoon SS, Burt V, Gu Q. Hypertension among adults in the United States: National Health and Nutrition Examination Survey, 2011-2012. *NCHS data brief.* 2013(133):1–8.
2. Aoki Y, Yoon SS, Chong Y, Carroll MD. Hypertension, abnormal cholesterol, and high body mass index among non-Hispanic Asian adults: United States, 2011-2012. *NCHS data brief.* 2014 1;(140): 1–8.

3. Jose PO, Frank AT, Kapphahn KI, Goldstein BA, Eggleston K, Hastings KG, Cullen MR, Palaniappan LP. Cardiovascular disease mortality in Asian Americans. *Journal of the American College of Cardiology*. 2014;64(23):2486–94. [PubMed: 25500233]
4. Kim MT, Kim KB, Juon HS, Hill MN. Prevalence and factors associated with high blood pressure in Korean Americans. *Ethn Dis*. 2000;10(3):364–74. Epub 2000/12/08. PubMed PMTD: 11110353. [PubMed: 11110353]
5. Wu TY, Hsieh HF, Wang J, Yao L, Oakley D. Ethnicity and cardiovascular risk factors among Asian Americans residing in Michigan. *J Community Health*. 2011;36(5):811–8. doi: 10.1007/s10900-011-9379-1. PubMed PMTD: 21380579. [PubMed: 21380579]
6. Cohen S, Janicki-Deverts D, Miller GE. Psychological stress and disease. *Jama*. 2007;298(14):1685–7. [PubMed: 17925521]
7. Everson-Rose SA, Lewis TT. Psychosocial factors and cardiovascular diseases. *Annu Rev Public Health*. 2005;26:469–500. [PubMed: 15760298]
8. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *Journal of health and social behavior*. 1983:385–96. [PubMed: 6668417]
9. McEwen BS, Gianaros PJ. Central role of the brain in stress and adaptation: links to socioeconomic status, health, and disease. *Annals of the New York Academy of Sciences*. 2010;1186(1):190–222. [PubMed: 20201874]
10. Richardson S, Shaffer JA, Falzon L, Krupka D, Davidson KW, Edmondson D. Meta-analysis of perceived stress and its association with incident coronary heart disease. *The American journal of cardiology*. 2012;110(12):1711–6. Epub 2012/09/15. doi: 10.1016/j.amjcard.2012.08.004. PubMed PMID: 22975465; PMCID: Pmc3511594. [PubMed: 22975465]
11. Palagini L, Bruno RM, Cheng P, Mauri M, Taddei S, Ghiadoni L, Drake CL, Morin CM. Relationship between insomnia symptoms, perceived stress and coping strategies in subjects with arterial hypertension: psychological factors may play a modulating role. *Sleep medicine*. 2016;19:108–15. [PubMed: 27198955]
12. Holt-Lunstad J, Smith TB, Layton JB. Social relationships and mortality risk: a meta-analytic review. *PLoS medicine*. 2010;7(7):859.
13. Orth-Gomer K, Rosengren A, Wilhelmsen L. Lack of social support and incidence of coronary heart disease in middle-aged Swedish men. *Psychosomatic Medicine*. 1993;55(1):37–43. [PubMed: 8446739]
14. Cacioppo JT, Hawkley LC. Social isolation and health, with an emphasis on underlying mechanisms. *Perspectives in biology and medicine*. 2003;46(3):S39–S52. [PubMed: 14563073]
15. House JS, Umberson D, Landis KR. Structures and processes of social support. *Annual review of sociology*. 1988:293–318.
16. Strawbridge WJ, Cohen RD, Shema SJ, Kaplan GA. Frequent attendance at religious services and mortality over 28 years. *American Journal of Public Health*. 1997;87(6):957–61. [PubMed: 9224176]
17. Barth J, Schneider S, von Kanel R. Lack of social support in the etiology and the prognosis of coronary heart disease: a systematic review and meta-analysis. *Psychosomatic medicine*. 2010;72(3):229–38. [PubMed: 20223926]
18. Hemingway H, Marmot M. Evidence based cardiology-Psychosocial factors in the aetiology and prognosis of coronary heart disease: systematic review of prospective cohort studies. *Bmj*. 1999;318(7196):1460–7. [PubMed: 10346775]
19. Lett HS, Blumenthal JA, Babyak MA, Strauman TJ, Robins C, Sherwood A. Social support and coronary heart disease: epidemiologic evidence and implications for treatment. *Psychosomatic medicine*. 2005;67(6):869–78. [PubMed: 16314591]
20. Redondo-Sendino Á, Guallar-Castillón P, Banegas JR, Rodríguez-Artalejo F. Relationship between social network and hypertension in older people in Spain. *Revista Espanola de Cardiologia (English Edition)*. 2005;58(11):1294–301.
21. Gump BB, Polk DE, Kamarck TW, Shiffman SM. Partner interactions are associated with reduced blood pressure in the natural environment: Ambulatory monitoring evidence from a healthy, multiethnic adult sample. *Psychosomatic medicine*. 2001;63(3):423–33. [PubMed: 11382269]

22. Bland SH, Krogh V, Winkelstein W, Trevisan M. Social network and blood pressure: a population study. *Psychosomatic medicine*. 1991;53(6):598–607. [PubMed: 1758945]
23. Glynn LM, Christenfeld N, Gerin W. Gender, social support, and cardiovascular responses to stress. *Psychosomatic medicine*. 1999;61(2):234–42. [PubMed: 10204977]
24. Cohen S, Wills TA. Stress, social support, and the buffering hypothesis. *Psychological bulletin*. 1985;98(2):310. [PubMed: 3901065]
25. Juon H-S, Strong C, Kim F, Park E, Lee S. Lay Health Worker Intervention Improved Compliance with Hepatitis B Vaccination in Asian Americans: Randomized Controlled Trial. *PLoS ONE*. 2016;11(9):e0162683. doi: 10.1371/journal.pone.0162683. [PubMed: 27617742]
26. Pickering TG, Hall JE, Appel LJ, Falkner BE, Graves J, Hill MN, Jones DW, Kurtz T, Sheps SG, Roccella EJ. Recommendations for blood pressure measurement in humans and experimental animals: part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Circulation*. 2005;111(5):697–716. Epub 2005/02/09. doi: 10.1161/01.cir.0000154900.76284.f6. PubMed PMID: 15699287. [PubMed: 15699287]
27. Cohen S Perceived stress in a probability sample of the United States 1988.
28. Berkman LF, Syme SL. Social networks, host resistance, and mortality: a nine-year follow-up study of Alameda County residents. *American journal of Epidemiology*. 1979; 109(2): 186–204. [PubMed: 425958]
29. Kawachi I, Berkman LF. Social ties and mental health. *Journal of urban health : bulletin of the New York Academy of Medicine*. 2001;78(3):458–67. Epub 2001/09/21. doi: 10.1093/jurban/78.3.458. PubMed PMID: 11564849; PMCID: Pmc3455910. [PubMed: 11564849]
30. Loucks EB, Sullivan LM, D'AGOSTINO RB Sr, Larson MG, Berkman LF, Benjamin EJ. Social networks and inflammatory markers in the Framingham Heart Study. *Journal of biosocial science*. 2006;38(06):835–42. [PubMed: 16441967]
31. Cleveland W Robust locally weighted regression and smoothing scatterplots. *Journal of the American Statistical Association*; 1979 p. 829–36.
32. Cleveland W, Devlin S. Locally-weighted regression: An approach to regression analysis by local fitting. *Journal of the American Statistical Association*; 1988 p. 596–610.
33. Cohen S, JANICKI-DEVERTS D Who's stressed? Distributions of psychological stress in the United States in probability samples from 1983, 2006, and 2009. *Journal of applied social psychology*. 2012;42(6): 1320–34.
34. Fazel M, Wheeler J, Danesh J. Prevalence of serious mental disorder in 7000 refugees resettled in western countries: a systematic review. *The Lancet*. 2005;365(9467): 1309–14.
35. Lee LC, Zane NW. *Handbook of Asian American psychology*: Sage Publications, Inc; 1998.
36. Chun M, Knight B, Youn G. Differences in stress and coping models of emotional distress among Korean, Korean-American and White-American caregivers. *Aging and Mental Health*. 2007; 11(1):20–9. [PubMed: 17164154]
37. Logan JG, Barksdale DJ, Carlson J, Carlson BW, Rowsey PJ. Psychological stress and arterial stiffness in Korean Americans. *Journal of psychosomatic research*. 2012;73(1):53–8. [PubMed: 22691560]
38. Kuo WH. Prevalence of depression among Asian-Americans. *The Journal of nervous and mental disease*. 1984; 172(8):449–57. [PubMed: 6747614]
39. Kim KC, Hurh WM. Ethnic resources utilization of Korean immigrant entrepreneurs in the Chicago minority area. *International Migration Review*. 1985:82–111.
40. Wiernik E, Pannier B, Czernichow S, Nabi H, Hanon O, Simon T, Simon J-M, Thomas F, Bean K, Consoli SM. Occupational Status Moderates the Association Between Current Perceived Stress and High Blood Pressure Evidence From the IPC Cohort Study. *Hypertension*. 2013;61(3):571–7. [PubMed: 23319539]
41. Gebreab SY, Diez-Roux AV, Hickson DA, Boykin S, Sims M, Sarpong DF, Taylor HA, Wyatt SB. The contribution of stress to the social patterning of clinical and subclinical CVD risk factors in African Americans: The Jackson Heart Study. *Social Science & Medicine*. 2012;75(9):1697–707. [PubMed: 22841454]

42. WINKLEBY MA, Ragland DR, Syme SL. Self-reported stressors and hypertension: evidence of an inverse association. *American Journal of Epidemiology*. 1988;127(1):124–34. [PubMed: 3276154]
43. Schneiderman N, Ironson G, Siegel SD. Stress and health: psychological, behavioral, and biological determinants. *Annual review of clinical psychology*. 2005;1:607.
44. Glaser R, Kiecolt-Glaser JK. Stress-induced immune dysfunction: implications for health. *Nature Reviews Immunology*. 2005;5(3):243–51.
45. Conway TL, Vickers RR Jr, Ward HW, Rahe RH. Occupational stress and variation in cigarette, coffee, and alcohol consumption. *Journal of Health and Social Behavior*. 1981:155–65. [PubMed: 7240714]
46. Anand SS, Yusuf S, Yusuf S, Cairns J, Camm J, Fallen E, Gersh B. Ethnicity and cardiovascular disease. *Evidence based cardiology*. 1998:329–52.
47. Sheth T, Nair C, Nargundkar M, Anand S, Yusuf S. Cardiovascular and cancer mortality among Canadians of European, south Asian and Chinese origin from 1979 to 1993 an analysis of 1.2 million deaths. *Canadian Medical Association Journal*. 1999;161(2):132–8. [PubMed: 10439820]
48. Steptoe A, Kivimaki M. Stress and cardiovascular disease: an update on current knowledge. *Annual review of public health*. 2013;34:337–54.
49. Gallo LC, Roesch SC, Fortmann AL, Carnethon MR, Penedo FJ, Perreira K, Birnbaum-Weitzman O, Wassertheil-Smoller S, Castaneda SF, Talavera GA. Associations of chronic stress burden, perceived stress, and traumatic stress with cardiovascular disease prevalence and risk factors in the HCHS/SOL Sociocultural Ancillary Study. *Psychosomatic medicine*. 2014;76(6):468. [PubMed: 24979579]
50. Tomaka J, Thompson S, Palacios R. The relation of social isolation, loneliness, and social support to disease outcomes among the elderly. *Journal of aging and health*. 2006;18(3):359–84. [PubMed: 16648391]
51. Grant N, Hamer M, Steptoe A. Social isolation and stress-related cardiovascular, lipid, and cortisol responses. *Annals of Behavioral Medicine*. 2009;37(1):29–37. [PubMed: 19194770]
52. Shankar A, McMunn A, Banks J, Steptoe A. Loneliness, social isolation, and behavioral and biological health indicators in older adults. *Health Psychology*. 2011;30(4):377. [PubMed: 21534675]
53. Lee J-S, Koeske GF, Sales E. Social support buffering of acculturative stress: A study of mental health symptoms among Korean international students. *International Journal of Intercultural Relations*. 2004;28(5):399–414.
54. Clark R Self-reported racism and social support predict blood pressure reactivity in Blacks. *Annals of Behavioral Medicine*. 2003;25(2):127–36. [PubMed: 12704015]
55. Koh KB. Perceived stress, psychopathology, and family support in Korean immigrants and nonimmigrants. *Yonsei medical journal*. 1998;39(3):214–21. [PubMed: 9664825]
56. Matud MP. Gender differences in stress and coping styles. *Personality and individual differences*. 2004;37(7):1401–15.
57. Martínez-Hernández A, Carceller-Maicas N, DiGiacomo SM, Ariste S. Social support and gender differences in coping with depression among emerging adults: a mixed-methods study. *Child and Adolescent Psychiatry and Mental Health*. 2016;10(1):2. doi: 10.1186/s13034-015-0088-x. [PubMed: 26744601]
58. U.S. Census Bureau. 2013 American Community Survey: Maryland 2013 [cited 2015 3 11]. Available from: <https://www.census.gov/quickfacts/table/PST045215/00>.
59. Paffenbarger RS, Jung DL, Leung RW, Hyde RT. Physical activity and hypertension: an epidemiological view. *Annals of medicine*. 1991;23(3):319–27. [PubMed: 1930924]
60. Beilin L Diet and hypertension: critical concepts and controversies. *Journal of hypertension Supplement: official journal of the International Society of Hypertension*. 1987;5(5):S447–57. [PubMed: 3327929]
61. Gupta S, McDade T, Adam E. Objective versus subjective measures of health: Systematic differences and determinants. Preliminary version. 2010.

Table 1.

Baseline Characteristics of the Participants (N = 530)

	Total N = 530	Non-Hypertensive N = 376	Hypertensive N= 154	P-value
Age				<0.01
< 40 years	126 (23.8)	111 (29.5)	15 (9.8)	
40-59 years	335 (63.2)	227 (60.4)	108 (70.1)	
60 years and over	69 (13.0)	38 (10.1)	31 (20.1)	
Gender				<0.01
Male	224 (42.3)	131 (34.8)	93 (60.4)	
Female	306 (57.7)	245 (65.2)	61 (39.6)	
Ethnicity				0.07
Chinese	177 (33.4)	137 (36.4)	40 (26.0)	
Korean	179 (33.8)	121 (32.2)	58 (37.7)	
Vietnamese	174 (33.8)	118 (31.4)	56 (36.3)	
BMI (mean[SD])	23.4 (3.4)	22.6 (3.1)	25.4 (3.2)	<0.01
Marital status				0.29
Married	411 (77.6)	287 (76.3)	124 (80.5)	
Not married	119 (22.4)	89 (23.7)	30 (19.5)	
Education				<0.01
Less than high school	68 (12.8)	40 (10.6)	28 (18.2)	
High school or some college+	202 (38.1)	135 (35.9)	67 (43.5)	
College graduate +	260 (49.1)	201 (53.5)	59 (38.3)	
Employment status				0.77
Employed	356 (67.2)	122 (32.4)	52 (33.8)	
Unemployed	174 (32.8)	254 (67.6)	102 (66.2)	
Self-rated Identity				0.07
Very Asian	304 (57.2)	205 (54.5)	99 (64.3)	
Mostly Asian	88 (16.5)	63 (16.8)	25 (16.2)	
Bicultural/Westernized	140 (26.3)	108 (28.7)	30 (19.5)	
English speaking				0.0017
Poorly/not at all	194 (36.6)	126 (33.5)	68 (44.2)	
So-so	207 (39.1)	143 (38.0)	64 (41.5)	
Native/very well	129 (24.3)	107 (28.5)	22 (14.3)	
Smoke				<0.01
Smoker	94 (17.7)	51 (13.6)	43 (27.9)	
Non-smoker	438 (82.3)	325 (86.4)	111 (72.1)	
Binge drink				0.31
Yes	67 (12.6)	44 (11.7)	23 (14.9)	
No	463 (82.4)	332 (88.3)	131 (85.1)	
Perceived Stress				0.01
Low	251 (47.2)	192 (51.1)	59 (38.3)	
High	281 (52.8)	184 (48.9)	95 (61.7)	

	Total N = 530	Non-Hypertensive N = 376	Hypertensive N= 154	P-value
Social Network Index Score				0.84
socially isolated	67 (12.6)	46 (12.2)	21 (13.6)	
moderately isolated	115 (21.6)	83 (22.1)	31 (20.1)	
moderately integrated	162 (30.5)	117 (31.1)	44 (28.6)	
socially integrated	188 (35.3)	130 (34.6)	58 (37.7)	
Social support				<0.01
Low	282 (53.0)	179 (47.6)	101 (65.6)	
High	250 (47.0)	197 (52.4)	53 (34.4)	

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2.

Age adjusted logistic regression models of psychological factors and hypertension (N = 530)

	Model 1	Model 2	Model 3
Perceived Stress			
Low		ref	
High	1.71 (1.15, 2.54)*		
Social Network Index Score			
socially isolated		ref	
moderately isolated		0.66 (0.33, 1.34)	
moderately integrated		0.60 (0.31, 1.17)	
socially integrated		0.65 (0.34, 1.24)	
Social support			
Low			ref
High			0.49 (0.33, 0.74)*
Age			
< 40 years		ref	ref
40-59 years	3.52 (1.95, 6.33)*	3.88 (2.11, 7.12)*	3.50 (1.94, 6.32)*
60 years and over	6.18 (3.00, 12.74)*	6.45 (3.10, 13.39)*	5.69 (2.76, 11.75)*

*
p<0.05

Table 3.

Multivariate logistic regression models of psychological factors and hypertension (N = 530)

	Model 1	Model 2	Model 3
Perceived Stress			
Low	ref		
High	1.61 (1.03, 2.51)*		
Social Network Index Score			
socially isolated		ref	
moderately isolated		0.51 (0.23, 1.12)	
moderately integrated		0.42 (0.19, 0.92)*	
socially integrated		0.52 (0.25, 1.12)	
Social support			
Low			ref
High			0.52 (0.33, 0.81)*
Age			
< 40 years	ref	ref	ref
40-59 years	3.04 (1.54, 5.98)*	3.52 (1.72, 7.22)*	3.13 (1.59, 6.20)*
60 years and over	4.32 (1.86, 10.06)*	4.82 (2.01, 11.52)*	4.18 (1.79, 9.76)*
Gender			
Male	ref	ref	ref
Female	0.43 (0.26, 0.73)*	0.44 (0.26, 0.74)*	0.47 (0.28, 0.80)*
BMI (mean[SD])	1.28 (1.19, 1.38)*	1.29 (1.20, 1.39)*	1.30 (1.21, 1.39)*
Education			
Less than high school	ref	ref	ref
High school or some college+	0.85 (0.44, 1.65)	0.86 (0.43, 1.69)	0.89 (0.46, 1.74)
College graduate +	0.54 (0.26, 1.09)	0.57 (0.28, 1.17)	0.56 (0.28, 1.15)
Employment status			
Unemployed	ref	ref	ref
Employed	0.89 (0.54, 1.47)	0.94 (0.56, 1.56)	0.85 (0.51, 1.41)
Self-rated Identity			
Very Asian	ref	ref	ref
Mostly Asian	1.05 (0.57, 1.94)	1.06 (0.57, 1.98)	1.08 (0.58, 2.02)
Bicultural/Westernized	0.67 (0.38, 1.19)	0.64 (0.36, 1.15)	0.70 (0.39, 1.23)
English speaking			
Poorly/not at all	ref	ref	ref
So-so	1.07 (0.63, 1.81)	1.06 (0.63, 1.81)	1.09 (0.64, 1.85)
Native/very well	0.57 (0.27, 1.23)	0.49 (0.23, 1.05)	0.60 (0.28, 1.30)
Smoke			
Non-smoker	ref	ref	ref
Smoker	1.24 (0.67, 2.29)	1.30 (0.71, 2.39)	1.28 (0.70, 2.36)
Binge drink			

	Model 1	Model 2	Model 3
No	ref	ref	ref
Yes	0.90 (0.46, 1.77)	0.95 (0.49, 1.87)	0.92 (0.47, 1.81)

*
p<0.05

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 4.

Logistic regression models of psychological factors and hypertension by gender (N = 530)

	Male N = 224	Female N = 306
Perceived Stress		
Low	ref	ref
High	1.95 (1.05, 3.60)*	1.53 (0.78, 3.00)
Social Network Index Score		
socially isolated	ref	ref
moderately isolated	0.22 (0.07, 0.71)	1.22 (0.35, 4.29)
moderately integrated	0.23 (0.07, 0.76)	0.79 (0.24, 2.66)
socially integrated	0.25 (0.08, 0.82)	1.03 (0.31, 3.36)
Social support		
Low	ref	ref
High	0.50 (0.26, 0.93)*	0.48 (0.24, 0.94)*

*
p<0.05

Adjusted for age, gender, BMI, education, employment status, self-identity, English speaking, smoke and binge drinking

Table 5.

Logistic regression models of psychological factors and hypertension by ethnicity (N = 530)

	Chinese N = 177	Korean N = 179	Vietnamese N = 174
Perceived Stress			
Low	ref	ref	ref
High	3.78 (1.13, 6.80)*	1.03 (0.46, 2.31)	1.73 (0.84, 3.57)
Social Network Index Score			
socially isolated	ref	ref	ref
moderately isolated	0.28 (0.07, 1.06)	2.03 (0.26, 16.08)	0.64 (0.16, 2.53)
moderately integrated	0.29 (0.07, 1.24)	0.89 (0.11, 6.95)	0.57 (0.16, 2.03)
socially integrated	0.40 (0.10, 1.65)	0.94 (0.13, 7.03)	1.02 (0.28, 3.73)
Social support			
Low	ref	ref	ref
High	0.36 (0.15, 0.87)*	0.47 (0.20, 1.07)	0.58 (0.26, 1.29)

*
p<0.05

Adjusted for Age, gender, BMI, education, employment status, self-identity, English speaking, smoke and binge drinking