



Cochrane
Library

Cochrane Database of Systematic Reviews

Progestogen for treating threatened miscarriage (Review)

Wahabi HA, Fayed AA, Esmail SA, Bahkali KH

Wahabi HA, Fayed AA, Esmail SA, Bahkali KH.
Progestogen for treating threatened miscarriage.
Cochrane Database of Systematic Reviews 2018, Issue 8. Art. No.: CD005943.
DOI: [10.1002/14651858.CD005943.pub5](https://doi.org/10.1002/14651858.CD005943.pub5).

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	3
BACKGROUND	5
OBJECTIVES	6
METHODS	6
RESULTS	9
Figure 1.	10
Figure 2.	13
Figure 3.	14
DISCUSSION	16
AUTHORS' CONCLUSIONS	17
ACKNOWLEDGEMENTS	17
REFERENCES	18
CHARACTERISTICS OF STUDIES	23
DATA AND ANALYSES	42
Analysis 1.1. Comparison 1 Progestogens versus placebo or no treatment, Outcome 1 Miscarriage.	43
Analysis 1.2. Comparison 1 Progestogens versus placebo or no treatment, Outcome 2 Preterm birth.	43
Analysis 1.3. Comparison 1 Progestogens versus placebo or no treatment, Outcome 3 Pregnancy-induced hypertension.	44
Analysis 1.4. Comparison 1 Progestogens versus placebo or no treatment, Outcome 4 Antepartum haemorrhage.	44
Analysis 1.5. Comparison 1 Progestogens versus placebo or no treatment, Outcome 5 Stillbirth.	44
Analysis 1.6. Comparison 1 Progestogens versus placebo or no treatment, Outcome 6 Congenital abnormalities.	45
APPENDICES	45
WHAT'S NEW	45
HISTORY	45
CONTRIBUTIONS OF AUTHORS	46
DECLARATIONS OF INTEREST	46
SOURCES OF SUPPORT	46
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	47
INDEX TERMS	47

[Intervention Review]

Progestogen for treating threatened miscarriage

Hayfaa A Wahabi¹, Amel A Fayed², Samia A Esmail³, Khawater Hassan Bahkali³

¹Chair of Evidence-Based Healthcare and Knowledge Translation, King Saud University, Riyadh, Saudi Arabia. ²College of Medicine, Clinical Department, Princess Nourah Bint Abdulrahman University, Riyadh, Saudi Arabia. ³Department of Family and Community Medicine, King Saud University, Riyadh, Saudi Arabia

Contact address: Hayfaa A Wahabi, Chair of Evidence-Based Healthcare and Knowledge Translation, King Saud University, Riyadh, 11451, Saudi Arabia. umlana@yahoo.com.

Editorial group: Cochrane Pregnancy and Childbirth Group.

Publication status and date: Edited (no change to conclusions), published in Issue 9, 2018.

Citation: Wahabi HA, Fayed AA, Esmail SA, Bahkali KH. Progestogen for treating threatened miscarriage. *Cochrane Database of Systematic Reviews* 2018, Issue 8. Art. No.: CD005943. DOI: [10.1002/14651858.CD005943.pub5](https://doi.org/10.1002/14651858.CD005943.pub5).

Copyright © 2018 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Miscarriage is a common complication encountered during pregnancy. It is defined as spontaneous pregnancy loss before 20 weeks' gestation. Progesterone's physiological role is to prepare the uterus for the implantation of the embryo, enhance uterine quiescence and suppress uterine contractions, hence, it may play a role in preventing rejection of the embryo. Inadequate secretion of progesterone in early pregnancy has been linked to the aetiology of miscarriage and progesterone supplementation has been used as a treatment for threatened miscarriage to prevent spontaneous pregnancy loss. This update of the Cochrane Review first published in 2007, and previously updated in 2011, investigates the evidence base for this practice.

Objectives

To determine the efficacy and the safety of progestogens in the treatment of threatened miscarriage.

Search methods

We searched Cochrane Pregnancy and Childbirth's Trials Register, [ClinicalTrials.gov](https://www.clinicaltrials.gov) and the WHO International Clinical Trials Registry Platform (ICTRP) (8 August 2017) and reference lists of retrieved trials.

Selection criteria

Randomised, quasi-randomised or cluster-randomised controlled trials, that compared progestogen with placebo, no treatment or any other treatment for the treatment of threatened miscarriage in women carrying singleton pregnancy.

Data collection and analysis

At least two review authors assessed the trials for inclusion in the review, assessed trial quality and extracted the data and graded the body of evidence.

Main results

We included seven trials (involving 696 participants) in this update of the review. The included trials were conducted in different countries, covering the full spectrum of the World Bank's economic classification, which enhances the applicability of evidence drawn from this review. Two trials were conducted in Germany and Italy which are high-income countries, while four trials were conducted in upper-middle income countries; two in Iran, one in Malaysia and the fourth in Turkey, and the seventh trial was conducted in Jordan, which is a lower-middle income country. In six trials all the participants met the inclusion criteria and in the seventh study, we included in the meta-analysis only the subgroup of participants who met the inclusion criteria. We assessed the body of evidence for the main outcomes using the GRADE

tool and the quality of the evidence ranged from very low to moderate. Downgrading of evidence was based on the high risk of bias in six of the seven included trials and a small number of events and wide confidence intervals for some outcomes.

Treatment of miscarriage with progestogens compared to placebo or no treatment probably reduces the risk of miscarriage; (risk ratio (RR) 0.64, 95% confidence interval (CI) 0.47 to 0.87; 7 trials; 696 women; moderate-quality evidence). Treatment with oral progestogen compared to no treatment also probably reduces the miscarriage rate (RR 0.57, 95% CI 0.38 to 0.85; 3 trials; 408 women; moderate-quality evidence). However treatment with vaginal progesterone compared to placebo, probably has little or no effect in reducing the miscarriage rate (RR 0.75, 95% CI 0.47 to 1.21; 4 trials; 288 women; moderate-quality evidence). The subgroup interaction test indicated no difference according to route of administration between the oral and vaginal subgroups of progesterone.

Treatment of miscarriage with the use of progestogens compared to placebo or no treatment may have little or no effect in reducing the rate of preterm birth (RR 0.86, 95% CI 0.52 to 1.44; 5 trials; 588 women; low-quality evidence).

We are uncertain if treatment of threatened miscarriage with progestogens compared to placebo or no treatment has any effect on the rate of congenital abnormalities because the quality of the evidence is very low (RR 0.70, 95% CI 0.10 to 4.82; 2 trials; 337 infants; very-low quality evidence).

Authors' conclusions

The results of this Cochrane Review suggest that progestogens are probably effective in the treatment of threatened miscarriage but may have little or no effect in the rate of preterm birth. The evidence on congenital abnormalities is uncertain, because the quality of the evidence for this outcome was based on only two small trials with very few events and was found to be of very low quality.

PLAIN LANGUAGE SUMMARY

Progestogen for treating threatened miscarriage

What is the issue?

Spontaneous miscarriage occurs in about 15% to 20% of pregnancies. Threatened miscarriage occurs when a mother might be losing her baby at less than 20 weeks' gestation. The symptoms of threatened miscarriage are vaginal bleeding, with or without abdominal pain, while the cervix of the womb is closed and the baby inside the womb is alive. Progesterone is a hormone that is known to prepare the uterus for implantation of the fertilized egg and suppress uterine contractions until term. Medications that mimic the action of progesterone are known as progestogens. Treatment with progestogens may be effective in reducing the rate of miscarriage in women who have threatened miscarriage. This Cochrane Review examines whether progestogens could reduce miscarriage for women with threatened miscarriage, and also addresses the safety of these medications for mother and baby.

Why is this important?

We were interested to investigate if progestogens are effective and safe in the treatment of threatened miscarriage, which may increase the women's chances of having a successful pregnancy and a live birth.

What evidence did we find?

In this review of the literature, up to August 2017, we identified seven randomised trials involving 696 women that compared the use of progestogens in the treatment of threatened miscarriage with either placebo or no treatment. We found that the use of a progestogen probably reduces the rate of spontaneous miscarriage and this was supported by moderate-quality evidence. Five trials, involving 588 women, reported on the effectiveness of progestogens given for threatened miscarriage in reducing the rate of preterm delivery and showed little or no effect, with low-quality evidence. Two trials, involving 337 women, reported on the effect of treatment with progestogens given for threatened miscarriage on the rate of occurrence of congenital abnormalities in the newborns. The evidence on congenital abnormalities is uncertain, because the quality of the evidence for this outcome was based on only two small trials with very few events and was found to be of very low quality.

What does this mean?

The evidence suggests that progesterone probably reduces the rate of spontaneous miscarriage but may make little or no difference to the number of preterm deliveries. The evidence for congenital abnormalities is uncertain because the quality of the evidence for this outcome was based on only two small trials with very few events and was found to be of very low quality.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. Progesterone compared to placebo or no treatment for treating threatened miscarriage

Progesterone compared to placebo or no treatment for treating threatened miscarriage

Patient or population: women with threatened miscarriage

Setting: two in high-income countries (Germany and Italy), four in upper-middle income countries (two in Iran, one in Malaysia and one in Turkey) and one in lower-income country (Jordan). All in hospitals (university or women's) or medical centres

Intervention: progesterone

Comparison: placebo or no treatment

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	N° of participants (trials)	Certainty of the evidence (GRADE)	Comments
	Risk with placebo or no treatment	Risk with progesterone treatment				
Miscarriage	Study population		RR 0.64 (0.47 to 0.87)	696 (7 RCTs)	⊕⊕⊕⊖ Moderate ¹	
	242 per 1000	138 per 1000 (102 to 189)				
Preterm birth	Study population		RR 0.86 (0.52 to 1.44)	588 (5 RCTs)	⊕⊕⊖⊖ Low ^{1, 2}	
	91 per 1000	84 per 1000 (49 to 142)				
Congenital abnormalities	Study population		RR 0.70 (0.10 to 4.82)	337 (2 RCTs)	⊕⊖⊖⊖ Very low ^{1, 2, 3}	
	13 per 1000	9 per 1000 (1 to 62)				

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **RCT:** randomised controlled trial; **RR:** risk ratio

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

- 1Random sequence generation and/or allocation concealment had a high or unclear risk of bias in the majority of the trials (methodological quality of trials) (-1).
- 2Wide confidence interval crossing the no effect line (imprecision) (-1).
- 3A small number of events (-1).

BACKGROUND

Description of the condition

"Miscarriage is pregnancy loss before 20 weeks' gestation based on the first day of the last menstrual period or, if gestation age is unknown, it is the loss of an embryo or a fetus of less than 400 g" (Zegers-Hochschild 2009). Threatened miscarriage is manifested by vaginal bleeding, with or without abdominal pain, while the cervix is closed and the fetus is viable and inside the uterine cavity (Cunningham 2001a). Once the cervix begins to dilate, miscarriage and pregnancy loss are inevitable. When the fetus is non-viable and the cervix is closed, this is known as missed miscarriage or missed abortion (Cunningham 2001a). The presence of a short cervix or funnelling of the internal cervical os in the gestation period between 16 and 24 weeks has been found to indicate increased risk or threat to miscarriage (Owen 2004; Rust 2005).

Miscarriage is a common complication of pregnancy occurring in 15% to 20% of all clinically recognised pregnancies (Blohm 2008; Cohain 2017). As many as 50% of the miscarried fetuses and embryos have normal chromosomes (Suzumori 2010; Vorsanova 2005). In many cases, the cause of miscarriage cannot be identified, however, among the recognised risk factors of miscarriage are maternal age over 34 years and paternal age over 40 years (De La Rochebrochard 2002), previous history of two or more miscarriages (Sugiura-Ogasawara 2009), and maternal autoimmune factors such as phospholipids antibodies (Check 2011). In addition, many modifiable risk factors were recognised including maternal obesity (Hahn 2014), maternal infection such as genital herpes simplex, HIV-1 and vaginal colonisation of group B streptococci (Nigro 2011; Rocchetti 2011; Temmerman 1992). Maternal endocrine abnormalities such as uncontrolled diabetes mellitus (Galindo 2006), polycystic ovary syndrome (Arredondo 2006) and insufficient production of progesterone are other known risk factors. Progesterone insufficiency is due to corpus luteum (CL) deficiency, which is a group of cells in the ovary that are formed after the release of the ovum (Cunningham 2001b). The main function of the CL is to produce sufficient progesterone to prepare and support the lining of the uterus and facilitate implantation of the ovum (Cunningham 2001b). It is suggested that progesterone deficiency in early pregnancy may be an aetiological factor for miscarriage, hence, treatment with progesterone may prevent miscarriage. However the evidence support for the effectiveness and safety of such treatment is uncertain based on the results of recently published trials (Coomarasamy 2015; Saccone 2017).

Description of the intervention

Progestogens are a group of hormones that bind to the progesterone receptors; they include both the natural female sex hormone progesterone and the synthetic forms. Medications that mimic the action of progesterone are known as progestational agents. Progesterone and progestational agents are known as progestogens. Progesterone is secreted during early pregnancy from the ovary by CL (Cunningham 2001b). It is an essential hormone for the establishment and maintenance of pregnancy. Progestogens can be given to women with threatened miscarriage orally, as intramuscular injection or in the form of vaginal suppositories. Due to the wide range of doses, preparations and types of progestogens used for the treatment of threatened miscarriage (Carp 2012; Mirza 2016), it is uncertain what is the

best type, dose and route of administration for the treatment of threatened miscarriage.

How the intervention might work

Progesterone induces secretory changes in the lining of the uterus, which are important for implantation of the fertilised ovum (Jin 2006). In addition, it modulates the immune response of the mother to prevent rejection of the embryo through a protein called progesterone induced blocking factor (PIBF), which is produced by the lymphocytes (white blood cells) of the pregnant woman (Szekeres-Bartho 2010), and it enhances uterine quiescence and suppresses uterine contractions (Szekeres-Bartho 2008; Szekeres-Bartho 2009). Low progesterone levels have been linked to increased risk of first trimester miscarriage (Osmanağaoğlu 2010).

Owing to the documented physiological role of progesterone in maintaining pregnancy, it has been used to treat women with threatened miscarriage and presumed progesterone deficiency to improve expectations for continuity of pregnancy (Palagianò 2004). The therapeutic value of progesterone in preventing or treating threatened miscarriage has not been established (Kalinka 2005), although more recent evidence suggests that it may be effective in preventing miscarriage in women with a history of recurrent miscarriage of unclear etiology (Haas 2018). This might be due to the poor designs of the trials done to evaluate its effectiveness (Kalinka 2005), and the inclusion of women in these trials with different aetiologies for threatened miscarriage.

Why it is important to do this review

Miscarriage is associated with considerable physical and psychological morbidity. Women who had threatened miscarriage were found to have increased rate of antepartum haemorrhage, pre-labour rupture of the membranes, preterm delivery, and intrauterine growth restriction when compared with women who did not have threatened miscarriage (Saraswat 2010). The emotional response to miscarriage can be profound; it includes depression, sleep disturbance, anger, and marital disturbances (Marcinko 2011). The introduction of ultrasound scans in the management of bleeding in early pregnancy has improved the diagnosis by rapid confirmation of viability and has improved the management by introducing prognostic factors such as fetal bradycardia and discrepancy between gestational age and crown-to-rump length (Dede 2010; Makrydimas 2003). This has rationalised the management of threatened miscarriage, as attempts to maintain a pregnancy are likely to be effective only if the fetus is viable and has no chromosomal abnormalities (Lede 2005).

The importance of progesterone on the maintenance of pregnancy was demonstrated by the successful use of progesterone antagonists, such as mifepristone (RU 486) in the elective induction of abortion (Dabash 2015). In a recently published systematic review vaginally administered progesterone was more effective in the prevention of preterm birth compared to the injectable progesterone (Oler 2017). This raised the question about the importance of the route of administration and the type of progestogen used to prevent preterm labour (Di Renzo 2005). These same questions might apply to the use of progestogens in the treatment of threatened miscarriage.

In earlier reports progestogen therapy has been linked to the development of hypospadias (deformity of the penis) in the male fetus (Silver 1999); however, recent reports did not show an increase in rate of hypospadias in infants of mothers who received progestogens in early pregnancy (Källén 2010).

The aim of this review is to study all the available data on the effectiveness of administration of progestogens for the treatment of threatened miscarriage.

OBJECTIVES

To determine the efficacy and the safety of progestogens in the treatment of threatened miscarriage.

METHODS

Criteria for considering studies for this review

Types of studies

We included randomised, quasi-randomised or cluster-randomised controlled trials, that assessed the effectiveness and safety of progestogens in the treatment of threatened miscarriage compared to placebo, no treatment or other intervention, if viability of the embryo or the fetus was confirmed before the commencement of treatment.

Types of participants

We included pregnant women, with threatened miscarriage at or less than 23 weeks, with singleton embryo or fetus, and who had a confirmed viable pregnancy. Fetal viability was to ensure that we excluded from this review trials that included women with bleeding in early pregnancy due to missed miscarriage. We have placed no restriction on the age of the woman or past obstetric history.

Types of interventions

We included any type of progestogens, natural or synthetic, used in the treatment of threatened miscarriage regardless of the dose, duration or route of administration compared with placebo, no treatment or other intervention.

Types of outcome measures

Primary outcomes

1. Miscarriage

Secondary outcomes

Mother

1. Pain relief (as defined by the study authors)
2. Thromboembolism
3. Preterm birth
4. Depression (as defined by the study authors)
5. Any other adverse outcomes that were reported (pregnancy-induced hypertension; antepartum haemorrhage)

Child

1. Preterm birth
2. Stillbirth
3. Neonatal death
4. Congenital abnormalities, including genital malformations

5. Any other adverse neonatal outcomes that were reported (intrauterine growth restriction; respiratory distress syndrome)
6. Low birthweight (not prespecified)
7. Birthweight (not prespecified)

Search methods for identification of studies

The following methods section of this review is based on a standard template used by Cochrane Pregnancy and Childbirth.

Electronic searches

We searched Cochrane Pregnancy and Childbirth's Trials Register by contacting their Information Specialist (8 August 2017)

The Register is a database containing over 23,000 reports of controlled trials in the field of pregnancy and childbirth. For full search methods used to populate Pregnancy and Childbirth's Trials Register including the detailed search strategies for CENTRAL, MEDLINE, Embase and CINAHL; the list of handsearched journals and conference proceedings, and the list of journals reviewed via the current awareness service, please follow this link to the editorial information about the [Cochrane Pregnancy and Childbirth](#) in the Cochrane Library and select the '**Specialized Register**' section from the options on the left side of the screen.

Briefly, Cochrane Pregnancy and Childbirth's Trials Register is maintained by their Information Specialist and contains trials identified from:

1. monthly searches of the Cochrane Central Register of Controlled Trials (CENTRAL);
2. weekly searches of MEDLINE (Ovid);
3. weekly searches of Embase (Ovid);
4. monthly searches of CINAHL (EBSCO);
5. handsearches of 30 journals and the proceedings of major conferences;
6. weekly current awareness alerts for a further 44 journals plus monthly BioMed Central email alerts.

Two people screen the search results and review the full text of all relevant trial reports identified through the searching activities described above. Based on the intervention described, each trial report is assigned a number that corresponds to a specific Pregnancy and Childbirth review topic (or topics), and is then added to the Register. The Information Specialist searches the Register for each review using this topic number rather than keywords. This results in a more specific search set that has been fully accounted for in the relevant review sections ([Included studies](#); [Excluded studies](#); [Studies awaiting classification](#); [Ongoing studies](#)).

In addition, we searched [ClinicalTrials.gov](#) and the WHO International Clinical Trials Registry Platform ([ICTRP](#)) for unpublished, planned and ongoing trial reports (8 August 2017) using the search terms given in [Appendix 1](#).

Searching other resources

We scanned bibliographies of identified papers.

We did not apply any language restrictions.

Data collection and analysis

We have outlined the methods of data collection and analysis that we used to assess [Gerhard 1987](#) and [Palagiano 2004](#) in the previous version of this review, [Wahabi 2011](#).

The following methods section of this review is based on a standard template used by Cochrane Pregnancy and Childbirth.

Selection of studies

Three review authors independently assessed for inclusion all the potential trials we identified as a result of the search strategy. We resolved any disagreement through discussion.

Data extraction and management

For methods used in the previous version of this review, see [Wahabi 2011](#).

For this update, we used the following methods for assessing the 29 reports that the Information Specialist identified as a result of the updated search.

Assessment of risk of bias in included studies

The four review authors independently assessed risk of bias for each study using the criteria outlined in the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011b](#)). We resolved any disagreement by discussion.

(1) Random sequence generation (checking for possible selection bias)

We described the method used to generate the allocation sequence in sufficient detail to allow an assessment of whether it should produce comparable groups.

For each included study we assessed the method as being at:

- low risk of bias (any truly random process, e.g. random number table; computer random number generator);
- high risk of bias (any non-random process, e.g. odd or even date of birth; hospital or clinic record number);
- unclear risk of bias.

(2) Allocation concealment (checking for possible selection bias)

For each included study we described the method used to conceal allocation to interventions prior to assignment and assessed whether intervention allocation could have been foreseen in advance of, or during recruitment, or changed after assignment.

We assessed the methods as being at:

- low risk of bias (e.g. telephone or central randomisation; consecutively numbered sealed opaque envelopes);
- high risk of bias (open random allocation; unsealed or non-opaque envelopes, alternation; date of birth);
- unclear risk of bias.

(3.1) Blinding of participants and personnel (checking for possible performance bias)

For each included study we described the methods used, if any, to blind study participants and personnel from knowledge of which intervention a participant received. We considered that trials were

at low risk of bias if they were blinded, or if we judged that the lack of blinding was unlikely to affect results. We assessed blinding separately for different outcomes or classes of outcomes.

We assessed the methods as being at:

- low, high or unclear risk of bias for participants;
- low, high or unclear risk of bias for personnel.

(3.2) Blinding of outcome assessment (checking for possible detection bias)

For each included study we described the methods used, if any, to blind outcome assessors from knowledge of which intervention a participant received. We assessed blinding separately for different outcomes or classes of outcomes.

We assessed methods used to blind outcome assessment as being at:

- low, high or unclear risk of bias.

(4) Incomplete outcome data (checking for possible attrition bias due to the amount, nature and handling of incomplete outcome data)

For each included study, and for each outcome or class of outcomes, we described the completeness of data including attrition and exclusions from the analysis. We stated whether attrition and exclusions were reported and the numbers included in the analysis at each stage (compared with the total randomised participants), reasons for attrition or exclusion where reported, and whether missing data were balanced across groups or were related to outcomes. Where sufficient information was reported, or could be supplied by the trial authors, we planned to re-include missing data in the analyses that we undertook.

We assessed methods as being at:

- low risk of bias (e.g. no missing outcome data; missing outcome data balanced across groups);
- high risk of bias (e.g. numbers or reasons for missing data imbalanced across groups; 'as treated' analysis done with substantial departure of intervention received from that assigned at randomisation);
- unclear risk of bias.

We excluded trials where more than 20% of participants were lost to follow-up.

(5) Selective reporting (checking for reporting bias)

For each included study we described how we investigated the possibility of selective outcome reporting bias and what we found.

We assessed the methods as being at:

- low risk of bias (where it was clear that all of the study's prespecified outcomes and all expected outcomes of interest to the review have been reported);
- high risk of bias (where not all the study's prespecified outcomes were reported; one or more reported primary outcomes were not prespecified; outcomes of interest were reported incompletely and so could not be used; study failed to include

results of a key outcome that would have been expected to have been reported);

- unclear risk of bias.

(6) Other bias (checking for bias due to problems not covered by (1) to (5) above)

For each included study we described any important concerns we had about other possible sources of bias.

(7) Overall risk of bias

We made explicit judgements about whether trials were at high risk of bias, according to the criteria given in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011b). With reference to (1) to (6) above, we planned to assess the likely magnitude and direction of the bias and whether we considered it was likely to have an impact on the findings. In future updates, we will explore the impact of the level of bias through undertaking sensitivity analyses (Sensitivity analysis).

Assessment of the quality of the evidence using the GRADE approach

For this update of the review, we used the GRADE approach, as outlined in the GRADE handbook (GRADE 2013), to assess the quality of the body of evidence relating to the following outcomes for the main comparisons, progesterone versus placebo or no treatment.

1. Miscarriage
2. Preterm birth
3. Congenital abnormalities

We used the GRADEpro Guideline Development tool (GRADEpro GDT 2015) to import data from Review Manager 5 (RevMan 5) (RevMan 2014) in order to create 'Summary of findings' tables. We produced a summary of the intervention effect and using the GRADE approach, a measure of quality for each of the above outcomes. The GRADE approach uses five considerations (study limitations, consistency of effect, imprecision, indirectness and publication bias) to assess the quality of the body of evidence for each outcome. The evidence can be downgraded from 'high quality' by one level for serious (or by two levels for very serious) limitations, depending on assessments for risk of bias, indirectness of evidence, serious inconsistency, imprecision of effect estimates or potential publication bias.

Measures of treatment effect

Dichotomous data

For dichotomous data, we present results as summary risk ratio (RR) with 95% confidence intervals (CIs).

Continuous data

None of the outcomes were reported as continuous data in this update. In future updates, if trials measure outcomes in the same way, we will use the mean difference. We will use the standardised mean difference to combine trials that measure the same outcome, but use different methods.

Unit of analysis issues

Cluster-randomised trials

We did not identify any cluster-randomised trials for inclusion. If we identify any cluster-randomised trials in future updates of this review, we will include them in the analyses along with individually randomised trials. We will adjust their sample sizes using the methods described in the *Cochrane Handbook for Systematic Reviews of Interventions*, using an estimate of the intra-cluster correlation co-efficient (ICC) derived from the trial (if possible), from a similar trial or from a study of a similar population (Higgins 2011c). If we use ICCs from other sources, we will report this and conduct sensitivity analyses to investigate the effect of variation in the ICC. If we identify both cluster-randomised trials and individually-randomised trials, we plan to synthesise the relevant information. We will consider it reasonable to combine the results from both if there is little heterogeneity between the study designs and the interaction between the effect of intervention and the choice of randomisation unit is considered to be unlikely.

We will also acknowledge heterogeneity in the randomisation unit and perform a subgroup analysis to investigate the effects of the randomisation unit.

Dealing with missing data

For included trials, we noted levels of attrition. For all outcomes, we carried out analyses, as far as possible, on an intention-to-treat basis, that is, we attempted to include all participants randomised to each group in the analyses, and analysed all participants in the group to which they were allocated, regardless of whether or not they received the allocated intervention. The denominator for each outcome in each trial was the number randomised minus any participants whose outcomes were known to be missing.

Assessment of heterogeneity

We assessed statistical heterogeneity in each meta-analysis using the Tau², I² (Higgins 2003) and Chi² statistics (Deeks 2011). We regarded heterogeneity as substantial if I² was greater than 30% and either Tau² was greater than zero, or there was a low P value (less than 0.10) in the Chi² test for heterogeneity.

Assessment of reporting biases

In future updates of this review, if there are 10 or more trials in the meta-analysis we will investigate reporting biases (such as publication bias) using funnel plots. We will assess funnel plot asymmetry visually. If asymmetry is suggested by a visual assessment, we will perform exploratory analyses to investigate it (Sterne 2011).

Data synthesis

We carried out statistical analysis using the RevMan 5 software (RevMan 2014). We used fixed-effect meta-analysis for combining data where it was reasonable to assume that trials were estimating the same underlying treatment effect: that is, where trials were examining the same intervention, and the trials' populations and methods were judged sufficiently similar. In future updates, if there is clinical heterogeneity sufficient to expect that the underlying treatment effects would differ between trials, or if we detect substantial statistical heterogeneity, we will use random-effects meta-analysis to produce an overall summary, if an average treatment effect across trials is considered clinically meaningful.

We will treat the random-effects summary as the average range of possible treatment effects and we will discuss the clinical implications of treatment effects differing between trials. If the average treatment effect is not clinically meaningful we will not combine trials (Deeks 2011).

If we use random-effects analyses, we will present the results as the average treatment effect with 95% CIs, and the estimates of Tau² and I².

Subgroup analysis and investigation of heterogeneity

We did not identify substantial heterogeneity. However, we carried out the following subgroup analyses for the primary outcome.

1. Route of administration

If we identify substantial heterogeneity in future updates of this review, we will investigate it using subgroup analyses and sensitivity analyses. We will consider whether an overall summary is meaningful, and if it is, use random-effects analysis to produce it. We will include the following subgroups, in addition to route of administration.

1. Type of progestogen
2. Progestogen dose

3. Effect of progestogens in early (no more than 12 weeks) and late (more than 12 weeks) threatened miscarriage

We will restrict subgroup analysis to the primary outcome. We will assess subgroup differences by interaction tests available within RevMan 5 (RevMan 2014). We will report the results of subgroup analyses quoting the Chi² statistic and P value, and the interaction test I² value (Deeks 2011).

Sensitivity analysis

We included seven trials in this review. All but one were similar in terms of risk of bias. For future updates of this review, we will carry out sensitivity analysis to explore the effect of risk of bias. This will involve analysis based on our 'Risk of bias' judgements for allocation concealment, performance bias and attrition bias. We will compare the results of those trials assessed as being rated 'low' risk of bias for these domains with those rated as 'high' or 'unclear' risk of bias.


RESULTS

Description of studies

Results of the search

We assessed 29 new trial reports of potential relevance identified by the updated search (see: Figure 1).

Figure 1. Trial flow diagram



Included studies

Trial location

We included seven trials (involving 696 participants) in this update of the review. The included trials were conducted in different countries, covering the full spectrum of the World Bank's economic classification ([World Bank 2017](#)), which may enhance the applicability and generalisation of evidence drawn from this review. Two trials were conducted in Germany and Italy ([Gerhard 1987](#); [Palagiano 2004](#)), which are high-income countries, while four trials were conducted in upper-middle income countries; two in Iran ([Alimohamadi 2013](#); [Yassae 2014](#)), one in Malaysia ([Pandian 2009](#)) and the fourth in Turkey ([Turgal 2017](#)). The seventh trial was conducted in Jordan ([El-Zibdeh 2009](#)), which is a lower-middle income country. All trials were conducted in hospital settings.

Trial design

We included six randomised controlled trials ([Alimohamadi 2013](#); [Gerhard 1987](#); [Palagiano 2004](#); [Pandian 2009](#); [Turgal 2017](#); [Yassae 2014](#)) and one quasi-randomised controlled trial ([El-Zibdeh 2009](#)).

Sample size

A total of 696 women were included in this review. The largest trial involved 191 women ([Pandian 2009](#)) and the smallest involved 35 women ([Gerhard 1987](#)).

Participants

We included pregnant women, with threatened miscarriage at or less than 23 weeks and who had a confirmed viable pregnancy. Fetal viability was to ensure that we excluded from this review trials that included women with bleeding in early pregnancy due to missed miscarriage. We placed no restriction on the age of the woman or past obstetric history.

Type of intervention

The seven included trials compared progestogens to either placebo ([Alimohamadi 2013](#); [Gerhard 1987](#); [Palagiano 2004](#); [Turgal 2017](#)) or no treatment ([El-Zibdeh 2009](#); [Pandian 2009](#); [Yassae 2014](#)). Three trials investigated oral progestogen ([El-Zibdeh 2009](#); [Pandian 2009](#); [Turgal 2017](#)) and four investigated vaginal progesterone ([Alimohamadi 2013](#); [Gerhard 1987](#); [Palagiano 2004](#); [Yassae 2014](#)).

Outcome measure

Primary outcome

Miscarriage

All the included trials investigated this primary outcome. Three trials investigated oral progestogen ([El-Zibdeh 2009](#); [Pandian 2009](#); [Turgal 2017](#)), and four trials investigated vaginal progesterone ([Alimohamadi 2013](#); [Gerhard 1987](#); [Palagiano 2004](#); [Yassae 2014](#)).

Secondary outcomes

Pain relief

Two trials ([Palagiano 2004](#); [Yassae 2014](#)) reported pain relief as an outcome of treatment with progesterone.

Preterm birth

Five trials ([Alimohamadi 2013](#); [El-Zibdeh 2009](#); [Gerhard 1987](#); [Pandian 2009](#); [Turgal 2017](#)); with 588 women, reported the effect of oral or vaginal progestogens on preterm birth.

Pregnancy-induced hypertension

Two trials ([El-Zibdeh 2009](#); [Pandian 2009](#)) reported pregnancy-induced hypertension as a maternal adverse outcome.

Antepartum haemorrhage

Two trials ([El-Zibdeh 2009](#); [Pandian 2009](#)) reported antepartum haemorrhage as a maternal adverse outcome.

Perinatal death

One trial investigated the effect of progestogens on neonatal death ([Alimohamadi 2013](#)) and two trials investigated the effect of progestogens on stillbirth ([Pandian 2009](#); [Turgal 2017](#)).

Congenital abnormalities

Two trials ([El-Zibdeh 2009](#); [Pandian 2009](#)) reported congenital abnormalities as an outcome.

Intrauterine growth restriction and low birthweight

Two trials ([Turgal 2017](#), [El-Zibdeh 2009](#)) reported on the frequency of intrauterine growth restriction in the intervention and control groups. Another trial ([Alimohamadi 2013](#)) reported on the frequency of low birthweight in the progesterone and the placebo group.

Respiratory distress syndrome

One trial ([Alimohamadi 2013](#)) investigated the rate of respiratory distress syndrome in the progesterone and the placebo group.

Birthweight

Only one trial ([Turgal 2017](#)) reported the difference in birthweight as an outcome.

Date of trials

Six trials were published between 2004 and 2017 ([Alimohamadi 2013](#); [El-Zibdeh 2009](#); [Palagiano 2004](#); [Pandian 2009](#); [Turgal 2017](#); [Yassae 2014](#)). One trial was published in 1987 ([Gerhard 1987](#)).

Funding source

Solvay Pharmaceuticals funded two trials ([El-Zibdeh 2009](#); [Pandian 2009](#)). The rest of the trials did not mention the source of funding.

Declaration of interest

None of the study authors reported any conflict of interest.

For further details, see [Characteristics of included studies](#).

Excluded studies

In this update of the review, we excluded 10 (plus two duplicate) full-text trials, in addition to the 32 excluded from the previous version. The reasons for exclusion of the 42 trials was as follows: 18 trials had a different population to this review' criteria ([Brenner 1962](#); [Corrado 2002](#); [El Zibdeh 2000](#); [El Zibdeh 2002](#); [El Zibdeh 2005](#); [Fuchs 1966](#); [Goldzieher 1964](#); [Johnson 1975](#); [Klopper 1965](#); [Le Vine 1964](#); [Nyboe 2002](#); [Porcaro 2015](#); [Prietl 1992](#); [Reijnders 1988](#);

Shearman 1963; Smitz 1992; Swyer 1953; Turner 1966); 14 trials, either compared progesterone to another type of progesterone or used it in combination with another therapy (Beigi 2016; Berle 1977; Check 1995; Chye 2014 (and duplicate); Czajkowski 2007; Famina 2015; Hui 2015; Luz 1988 [pers comm]; Pustotina 2018; Siew 2014; Siew 2015; Song 2007; Vincze 2006; Zhang 2000); seven trials did not use a reliable method to confirm the viability of the fetus (Berle 1980; Crowder 1950; Govaerts-Videtzky 1965; Moller 1965; Sondergaard 1985; Souka 1980; Tognoni 1980); one study

(and duplicate) had a mixed group of participants and we could not obtain separate results for the threatened-miscarriage group (Costantino 2016); one was not an RCT (Akhtar 2012); and one study had a high rate of attrition, more than 20% (Omar 2005).

For further details, see [Characteristics of excluded studies](#).


Risk of bias in included studies

([Figure 2](#), [Figure 3](#))

Figure 2. Risk of bias summary: review authors' judgements about each risk of bias item for each included trial

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Alimohamadi 2013	+	+	+	+	+	+	+
El-Zibdeh 2009	-	-	-	+	+	+	-
Gerhard 1987	?	?	+	+	+	+	?
Palagiano 2004	?	+	+	+	+	-	+
Pandian 2009	+	+	-	+	+	+	+
Turgal 2017	+	?	+	+	+	+	+
Yassaei 2014	?	?	-	+	+	+	+

Figure 3. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included trials



We judged only three of the seven included trials (42%) to be at low risk of bias for random sequence generation and allocation concealment, and four trials (57%) to be at low risk for performance bias. We judged all trials (100%) at low risk of detection bias and attrition bias. We judged six trials at low risk of selective reporting bias (85%), and five trials (71%) at low risk of other biases.

Allocation

In three trials (Alimohamadi 2013; Pandian 2009; Turgal 2017), the investigators described a random component in the sequence-generation process by using computer random-number generators, so we judged these trials to have low risk of selection bias. In three trials (Gerhard 1987; Palagiano 2004; Yassaee 2014), there was insufficient information about the sequence-generation process to permit judgement of low or high risk, so we judged them as unclear risk of selection bias. We judged one study (El-Zibdeh 2009) as high risk for selection bias, as the investigators described a non-random component in the sequence-generation process, which was based on the day of the week that women presented to the clinic.

Participants and investigators enrolling participants could not foresee assignment to intervention or control group in three trials because Alimohamadi 2013 and Palagiano 2004 used sequentially numbered drug containers of identical appearance and Pandian 2009 used sequentially numbered concealed envelopes; we judged these trials at low risk of selection bias. Three trials (Gerhard 1987; Turgal 2017; Yassaee 2014), did not describe the method of concealment sufficiently to allow a definite judgement, so we judged them at unclear risk of selection bias. In El-Zibdeh 2009, participants and investigators enrolling participants could foresee assignments, as the allocation was performed by the attending physician based on the day of the week the women visited the clinic, and thus the risk of selection bias introduction was high.

Blinding

Four trials (Alimohamadi 2013; Gerhard 1987; Palagiano 2004; Turgal 2017) ensured blinding of participants and key study personnel, and it is unlikely that the blinding could have been

broken, so we judged the risk of performance bias to be low. Three trials did not use placebo for the control group. Two of these (El-Zibdeh 2009; Pandian 2009) did not blind either personnel or women to the treatment received and one (Yassaee 2014) was a single-blind study, in which the researchers were unaware which woman had received progesterone. Although outcomes such as miscarriage, preterm birth and fetal structural malformations are not likely to be influenced by lack of blinding, lack of blinding of participants or personnel could have introduced performance bias if participants inadvertently took a different type of progestogen than the one in the trial or exaggerated their reports of subjective outcomes, such as pain.

Five of the seven trials blinded outcome assessors to the treatment the women had received (Alimohamadi 2013; El-Zibdeh 2009; Gerhard 1987; Palagiano 2004; Yassaee 2014). Pandian 2009 did not blind the assessors, and Turgal 2017 did not give any information about blinding. Outcomes such as miscarriage, preterm birth and fetal structural malformations are not likely to be influenced by lack of assessor's blinding, so we judged all seven trials as having low risk of detection bias.

Incomplete outcome data

There was no missing outcome data in five of the seven included trials (Alimohamadi 2013; El-Zibdeh 2009; Palagiano 2004; Pandian 2009; Yassaee 2014); Gerhard 1987 excluded two women (5.7%) from the study after randomisation, and Turgal 2017 reported incomplete outcome data for 12 women (14.5%), equal in both groups. We judged all seven trials to be at low risk of attrition bias.

Selective reporting

All the included trials, except one (Palagiano 2004), reported all of the expected prespecified (primary and secondary) outcomes, so we assessed these at low risk of reporting bias. In Palagiano 2004, the study authors mentioned in the methods that they would report all the adverse effects, but in the results there was no explicit mention of the absence or presence of any adverse effects, so we judged it to be at high risk of reporting bias.

Other potential sources of bias

We assessed one study at high risk of other biases due to the difference in the number of participants recruited to the experimental and the control groups (86 versus 60; [El-Zibdeh 2009](#)). In another study ([Gerhard 1987](#)), the risk of other bias was not clear because we included only participants with evidence of fetal viability, which amounts to only half of the participants.

Effects of interventions

See: [Summary of findings for the main comparison Progesterone compared to placebo or no treatment for treating threatened miscarriage](#)

Seven trials met the inclusion criteria, involving 696 participants. We performed subgroup analysis for the effect of progestogens by route of administration; however, due to paucity of data we could not carry out subgroup analysis for type and dose of progestogen. We undertook a total of six meta-analyses.

Primary outcome

Miscarriage

The seven included trials, with 696 participants, compared progestogens to either placebo ([Alimohamadi 2013](#); [Gerhard 1987](#); [Palagiano 2004](#); [Turgal 2017](#)) or no treatment ([El-Zibdeh 2009](#); [Pandian 2009](#); [Yassaee 2014](#)). Treatment of miscarriage with progestogens compared to placebo or no treatment probably reduces the risk of miscarriage; (risk ratio (RR) 0.64, 95% confidence interval (CI) 0.47 to 0.87; 7 trials; 696 women; moderate-quality evidence).

Subgroup analysis by route of administration

Three trials investigated treatment with oral progestogen ([El-Zibdeh 2009](#); [Pandian 2009](#); [Turgal 2017](#)), and showed that treatment with oral progestogen compared to no treatment probably reduces the miscarriage rate (RR 0.57, 95% CI 0.38 to 0.85; 3 trials; 408 women; moderate-quality evidence; [Analysis 1.1.1](#)). Four trials investigated treatment with vaginal progesterone compared to placebo ([Alimohamadi 2013](#); [Gerhard 1987](#); [Palagiano 2004](#); [Yassaee 2014](#)), which probably has little or no effect in reducing the miscarriage rate (RR 0.75, 95% CI 0.47 to 1.21; 4 trials; 288 women; moderate-quality evidence; [Analysis 1.1.2](#)). However, it should be noted that there were no differences between these two subgroups according to the subgroup interaction test (test for subgroup differences: $\text{Chi}^2 = 0.76$, $\text{df} = 1$ ($P = 0.38$), $I^2 = 0\%$).

Secondary outcomes

Maternal outcomes

Pain relief

Two trials ([Palagiano 2004](#); [Yassaee 2014](#)) reported pain relief as an outcome of treatment with progesterone. [Palagiano 2004](#) reported reduction on the mean pain score with the use of progesterone from 2.6 ± 0.9 before treatment to 0.4 ± 0.7 (mean \pm standard deviation) after treatment ($P < 0.01$), while no change in pain score was observed in women who received placebo from 2.5 ± 1.0 before treatment to 2.4 ± 0.8 (mean \pm standard deviation) after treatment ($P > 0.05$). They reported pain using a progressive score from 0 to 4, where a score of 0 indicated 'no pain' and a score of 4 indicated 'extreme pain'. In [Yassaee 2014](#), nine (30%) of the women in the progesterone and seven (23.3%) of the control group, continued to

have pain after intervention ($P = 0.55$). We could not pool the results of these two trials due to the different types of data (dichotomous and continuous).

Preterm birth

Five trials ([Alimohamadi 2013](#); [El-Zibdeh 2009](#); [Gerhard 1987](#); [Pandian 2009](#); [Turgal 2017](#)); reported the effect of progestogens on preterm birth. Treatment of preterm birth with the use of progestogens compared to placebo or no treatment may have little or no effect in reducing the rate of preterm birth (RR 0.86, 95% CI 0.52 to 1.44; 5 trials; 588 women; low-quality evidence; [Analysis 1.2](#)).

Other maternal adverse outcomes

Pregnancy-induced hypertension

Two trials ([El-Zibdeh 2009](#); [Pandian 2009](#)) reported pregnancy-induced hypertension as a maternal adverse outcome. There was no difference in the occurrence of pregnancy-induced hypertension between the progestogen and the control group (RR 1.00, 95% CI 0.54 to 1.88; 2 trials; 337 women; [Analysis 1.3](#)).

Antepartum haemorrhage

Two trials ([El-Zibdeh 2009](#); [Pandian 2009](#)) reported antepartum haemorrhage as maternal adverse outcome. Progestogens have little or no difference in the occurrence of antepartum haemorrhage (RR 0.76, 95% CI 0.30 to 1.94; 2 trials; 337 women; [Analysis 1.4](#)).

Child outcomes

Stillbirth

Two trials investigated the effect of progestogens on the stillbirth rate ([Pandian 2009](#); [Turgal 2017](#)). Progestogens have little or no difference effects on the rate of stillbirth (RR 1.94, 95% CI 0.18 to 20.49; 2 trials; 262 women; [Analysis 1.5](#)).

Congenital abnormalities

Two trials ([El-Zibdeh 2009](#); [Pandian 2009](#)) reported congenital abnormalities as an outcome. We are uncertain if treatment of threatened miscarriage with progestogens compared to placebo or no treatment has any effect on the rate of congenital abnormalities, (RR 0.70, 95% CI 0.10 to 4.82; 2 trials; 337 infants; very low-quality evidence; [Analysis 1.6](#)).

Neonatal death

Only one trial reported the effect of progestogens on neonatal death ([Alimohamadi 2013](#)). Progestogens have little or no effect in the neonatal death rate (RR 1.35, 95% CI 0.31 to 5.83; 1 trial; 145 women).

Other neonatal adverse outcomes

Intrauterine growth restriction and low birthweight

One trial ([Turgal 2017](#)) reported on the frequency of intrauterine growth restriction in the intervention and control group and showed no difference between the two groups; 0 (0%) in the progestogen group, one (3.5%) in the control group. Another trial ([Alimohamadi 2013](#)) reported on the frequency of low birthweight in the progesterone and the placebo group and showed no difference between the two groups; five (7%) for the progesterone group and seven (9.8%) for the placebo group.

Birthweight

Only one trial (Turgal 2017) reported the difference in birthweight as an outcome. It showed no difference in the mean birthweight between the progestogen and the control group; 3.1156 ± 0.643 kg for the intervention group and 3.2076 ± 0.500 kg for the control group.

Respiratory distress syndrome

One trial (Alimohamadi 2013) investigated the rate of respiratory distress syndrome in the progesterone and the placebo groups, and showed no difference between the two groups; there were two neonates with respiratory distress syndrome (2.8%) in the progesterone group and one (1.4%) in the placebo group.

Outcomes not reported in the primary trials

Due to paucity of data, we could not evaluate the following secondary outcomes.

1. Thromboembolism
2. Depression

In addition we could not perform the following subgroup analyses.

1. Route of progestogen administration
2. Type of progestogen
3. Progestogen dose
4. Effect of progestogens in early and late miscarriage

DISCUSSION

Summary of main results

We included seven trials (involving 696 participants) in this update of the review. The included trials were conducted in different countries, covering the full spectrum of the World Bank's economic classification (World Bank 2017), which enhances the applicability of evidence drawn from this review. Two trials were conducted in Germany and Italy which are high-income countries, while four trials were conducted in upper-middle income countries; two in Iran, one in Malaysia and the fourth in Turkey, and the seventh trial was conducted in Jordan, which is a lower-middle income country. In six trials all the participants met the inclusion criteria and in the seventh study, we included only the subgroup of participants who met the inclusion criteria in the meta-analysis. Using the GRADE tool, we assessed the body of evidence for the main outcomes as ranging from very low-quality to moderate-quality evidence. Downgrading of evidence was based on the high risk of bias in six of the seven included trials and imprecision for some of the outcomes.

Treatment of miscarriage with progestogens compared to placebo or no treatment probably reduces the risk of miscarriage. Subgroup analysis by route of administration showed that treatment with oral progestogen compared to no treatment probably reduces the miscarriage rate, although treatment with vaginal progesterone compared to placebo, probably has little or no effect in reducing the miscarriage rate. However, the subgroup interaction test indicated no difference according to route of administration between the oral and vaginal subgroups of progesterone. Treatment of preterm birth with the use of progestogens compared to placebo or no treatment may have little or no effect in reducing the rate of preterm birth. We are uncertain if treatment of threatened miscarriage with

progestogens compared to placebo or no treatment has any effect on the rate of congenital abnormalities.

Overall completeness and applicability of evidence

Threatened miscarriage is a common health problem. In a longitudinal study by Blohm 2008, one in four multiparous women had experienced a miscarriage and more than 7% of the studied population had experienced three or more miscarriages. In a large population-based study by Cohain 2017, more than 40% of parous women reported having experienced one or more first trimester spontaneous miscarriages.

This update of the review includes seven trials, with a relatively small number of women (696), from six countries. All trials were conducted in hospital settings. Two trials were conducted in high-income countries (Gerhard 1987; Palagiano 2004), four in upper-middle income countries (Alimohamadi 2013; Pandian 2009; Turgal 2017; Yassaee 2014) and one in lower-middle income country (El-Zibdeh 2009). Although no trial was conducted in a low-income country, the participants in the trials of the review have diverse ethnic and economic backgrounds (World Bank 2017). This may make the evidence in this review applicable to a large sector of women with threatened miscarriage.

Quality of the evidence

In this updated review we assessed the body of evidence for miscarriage as moderate quality, mainly due to limitations in study design of the included trials. From the seven included trials we judged only three (42%) to be at low risk of bias for random sequence generation and allocation concealment. Four trials (57%) were at low risk for performance bias, and all were at low risk of detection bias. We assessed all trials at low risk for attrition bias and six were at low risk of selective reporting bias. Only five trials (71%) trials were at low risk for other biases (Summary of findings for the main comparison).

For preterm birth we assessed the body of evidence as low quality due to imprecision (wide confidence intervals crossing the no-effect line) and limitations with study design of the included trials. From the five included trials, three (60%) were at low risk of bias for random sequence generation, while three (60%) were at low risk of performance bias and all trials were at low risk of detection bias. In addition, one study randomised different numbers of participants to the two arms of the study, which we considered as additional bias and in one study we included the results of only 50% of the participants who met the inclusion criteria.

For the outcome congenital abnormality, we assessed the body of evidence as very low due to imprecision (wide confidence intervals crossing the no-effect line), small number of trials (two trials) and participants (337 women) and limitations with study design. We assessed one of the two included trials to be at high risk of selection bias and performance bias and the other study at high risk of performance bias.

Potential biases in the review process

In this review, we took steps to minimise bias, although we are aware that bias may be present in our review: two review authors independently assessed trials for eligibility and extracted the data as necessary. We resolved discrepancies through discussion or, if required, we consulted a third review author; two review

authors also performed GRADE assessments independently and resolved discrepancies through discussion; and we conducted a comprehensive search in order to identify all relevant published and unpublished literature.

Agreements and disagreements with other studies or reviews

Our assessment of the body of evidence in this update of the review did not differ much from the previous version of the review (Wahabi 2011). For the main comparison, we have included three additional trials, however, neither the magnitude nor the direction of evidence changed markedly from the earlier version of the review. We have used the recent methodology introduced for Cochrane Reviews in 2008, which assesses risk of bias in the individual trials more carefully than in the past (Higgins 2011b). In addition, we assessed the quality of the evidence using the GRADE approach as outlined in the GRADE Handbook (GRADE 2013).

The results of this review agree with a recently published systematic review of randomised and non-randomised controlled trials of dydrogesterone (oral progestogen) for the treatment of threatened miscarriage (Carp 2012) and with non-randomised controlled trials (Akhtar 2012), which investigated the effectiveness of oral progestogens in the treatment of threatened miscarriage. The results of this trial showed that a larger proportion of women who received progestogen continued to be pregnant compared to the control group, however, the difference was not significant.

Recently published trials suggested that the route of administration and the type of progestogen influence its effectiveness in the prevention of miscarriage. In a multicenter, randomised, placebo-controlled trial on women with unexplained recurrent miscarriages, vaginal progesterone was found not to be significantly different from placebo in the prevention of miscarriage (Coomarasamy 2015). However, the results of a review of randomised trials on the use of progestogen in the prevention of recurrent miscarriage, concluded that synthetic progestogens, but not natural progesterone, were associated with a lower risk of recurrent miscarriage, nevertheless, the conclusion was limited by the low quality of evidence (Saccone 2017).

Our results suggested that progestogen may not be as effective in the prevention of preterm birth (Analysis 1.2). There are conflicting

published reports about the effectiveness of progestational agents in the prevention of preterm birth with apparent influence of type of progestogen and route of administration, as well as different subgroups of women, on the effectiveness of treatment (Facchinetti 2017; Heyborne 2016; O'Brien 2016).

Our results relating to the safety of progestogen use for the mother and infant are limited by the small number of trials and participants in addition to the low quality of the evidence. However, our systematic review is in agreement with other systematic reviews, which also suggest that there are no increased risks of adverse outcomes to the mother or the infant from the use of progestogen compared to the control group (Haas 2018).

AUTHORS' CONCLUSIONS

Implications for practice

The results of this review suggest that progestogens are probably effective in the treatment of threatened miscarriage but may have little or no effect in the rate of preterm birth. The evidence on congenital abnormalities is uncertain, because the quality of the evidence for this outcome was based on only two small trials with very few events and was found to be of very low quality.

Implications for research

Further research could investigate the effectiveness and safety of progestogens in the treatment of threatened miscarriage by conducting methodologically sound, randomised trials, with special focus on the type of progestogen and the route of administration.

ACKNOWLEDGEMENTS

The authors extend their gratitude to King Saud University, Deanship of Scientific Research, Research Chairs, for funding this update of the review.

This project was supported by the National Institute for Health Research, via Cochrane Infrastructure funding to Cochrane Pregnancy and Childbirth. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, NHS or the Department of Health.

REFERENCES

References to studies included in this review

Alimohamadi 2013 {published data only}

Alimohamadi S, Javadian P, Gharedaghi MH, Javadian N, Alinia H, Khazardoust S, et al. Progesterone and threatened abortion: a randomized clinical trial on endocervical cytokine concentrations. *Journal of Reproductive Immunology* 2013;**98**(1-2):52-60.

El-Zibdeh 2009 {published data only}

El-Zibdeh MY, Yousef LT. Dydrogesterone support in threatened miscarriage. *Maturitas* 2009;**65 Suppl 1**:S43-S46.

Gerhard 1987 {published data only}

Gerhard I, Gwinner B, Eggert-Kruse W, Runnebaum B. Double-blind controlled trial of progesterone substitution in threatened abortion. *Biological Research in Pregnancy and Perinatology* 1987;**8**:26-34.

Palagiano 2004 {published data only}

Palagiano A, Bulletti C, Pace MC, De Ziegler D, Cicinelli E, Izzo A. Effects of vaginal progesterone on pain and uterine contractility in patients with threatened abortion before twelve weeks of pregnancy. *Annals of the New York Academy of Sciences* 2004;**1034**:200-10.

Pandian 2009 {published data only}

Pandian RU. Dydrogesterone in threatened miscarriage: a Malaysian experience. *Maturitas* 2009;**65 Suppl 1**:S47-S50.

Turgal 2017 {published data only}

Turgal M, Aydin E, Ozyuncu O. Effect of micronized progesterone on fetal-placental volume in first-trimester threatened abortion. *Journal of Clinical Ultrasound* 2017;**45**(1):14-9.

Yassaee 2014 {published data only}

Yassaee F, Shekarriz-Foumani R, Afsari S, Fallahian M. The effect of progesterone suppositories on threatened abortion: a randomized clinical trial. *Journal of Reproduction & Infertility* 2014;**15**(3):147-51.

References to studies excluded from this review

Akhtar 2012 {published data only}

Akhtar M, Fatima N, Jabeen S, Akram M. Role of progesterone in the management of threatened miscarriage. *Pakistan Journal of Medical and Health Sciences* 2012;**6**(1):261-4.

Beigi 2016 {published data only}

Beigi A, Esmailzadeh A, Pirjani R. Comparison of risk of preterm labor between vaginal progesterone and 17-alpha-hydroxyprogesterone caproate in women with threatened abortion: a randomized clinical trial. *International Journal of Fertility & Sterility* 2016;**10**(2):162-8. [IRCT2014123120504N1]

Berle 1977 {published data only}

Berle P, Behnke K. The treatment of threatened abortion. *Geburtshilfe und Frauenheilkunde* 1977;**37**:139-42.

Berle 1980 {published data only}

Berle P, Budenz M, Michaelis J. Is hormonal therapy still justified in imminent abortion. *Zeitschrift für Geburtshilfe und Perinatologie* 1980;**184**:353-8.

Brenner 1962 {published data only}

Brenner WE, Hendricks CH. Effect of medroxyprogesterone acetate upon the duration and characteristics of human gestation and labor. *American Journal of Obstetrics and Gynecology* 1962;**83**:1094-8.

Check 1995 {published data only}

Check JH, Tarquini P, Gandy P, Lauer C. A randomized study comparing the efficacy of reducing the spontaneous abortion rate following lymphocyte immunotherapy and progesterone treatment vs progesterone alone in primary habitual aborters. *Gynecologic and Obstetric Investigation* 1995;**39**:257-61.

Chye 2014 {published data only}

Chye TT, ChiCTR-IOR-17011593. The randomised controlled trial of micronised progesterone and dydrogesterone (TRoMaD) for threatened miscarriage. chictr.org.cn/showproj.aspx?proj=19668 (first received 6 July 2017).

* Chye TT, ChiCTR-IPR-14005251. Micronised progesterone versus dydrogesterone: a randomised controlled trial. chictr.org.cn/showproj.aspx?proj=9539http://www.chictr.org.cn/showproj.aspx?proj=9539 (first received 21 September 2014).

Corrado 2002 {published data only}

Corrado F, Dugo C, Cannata ML, Di Bartolo M, Scilipoti A, Stella NC. A randomised trial of progesterone prophylaxis after midtrimester amniocentesis. *European Journal of Obstetrics & Gynecology and Reproductive Biology* 2002;**100**:196-8.

Costantino 2016 {published data only}

Costantino D, NCT02601898. Vaginal administration of alpha lipoic acid vs progesterone for the subchorionic hematoma treatment. clinicaltrials.gov/show/NCT02601898 (first received 6 November 2015).

* Costantino M, Guaraldi C, Costantino D. Resolution of subchorionic hematoma and symptoms of threatened miscarriage using vaginal alpha lipoic acid or progesterone: clinical evidences. *European Review for Medical and Pharmacological Sciences* 2016;**20**(8):1656-63.

Crowder 1950 {published data only}

Crowder RE, Bills ES, Broadbent JS. The management of threatened abortion: a study of 100 cases. *American Journal of Obstetrics and Gynecology* 1950;**60**:896-9.

Czajkowski 2007 {published data only}

Czajkowski K, Sienko J, Mogilinski M, Bros M, Szczecina R, Czajkowska A. Uteroplacental circulation in early pregnancy complicated by threatened abortion supplemented with vaginal micronized progesterone or oral dydrogesterone. *Fertility and Sterility* 2007;**87**(3):613-8.

El Zibdeh 2000 {published data only}

El Zibdeh MY. Randomized study comparing the efficacy of reducing spontaneous abortion following treatment with progesterone and human chorionic gonadotropin (hCG). *Fertility and Sterility* 2000;**70**(3 Suppl 1):S77-S78.

El Zibdeh 2002 {published data only}

El Zibdeh M. Randomized clinical trial comparing the efficacy of dydrogesterone and human chorionic gonadotropin [abstract]. 10th World Congress on Menopause; 2002 July 10-14; Berlin, Germany. 2002:136.

El Zibdeh 2005 {published data only}

El-Zibdeh MY. Dydrogesterone in the reduction of recurrent spontaneous abortion. *Journal of Steroid Biochemistry & Molecular Biology* 2005;**97**(5):431-4.

Famina 2015 {published data only}

Famina M. The effect of dydrogesterone on placental angiogenesis and pregnancy outcomes in women with threatened miscarriages. *Giornale Italiano di Ostetricia e Ginecologia* 2015;**37**(4):155-8.

Fuchs 1966 {published data only}

Fuchs F, Olsen P. An attempted double-blind controlled trial of progesterone therapy in habitual abortion. *Ugeskrift for Laeger* 1966;**128**:1461-2.

Goldzieher 1964 {published data only}

Goldzieher JW. Double-blind trial of a progestin in habitual abortion. *JAMA* 1964;**188**(7):651-4.

Govaerts-Videtzky 1965 {published data only}

Govaerts-Videtzky M, Martin L, Hubinont PO. A double-blind study of progestogen treatment in spontaneous abortion (preliminary report). *Journal of Obstetrics and Gynaecology of the British Commonwealth* 1965;**72**:1034.

Hui 2015 {published data only}

Hui CYY, Siew SJY, Tan TC. Biochemical and clinical outcomes following the use of micronised progesterone and dydrogesterone for threatened miscarriage - a randomised controlled trial. *BJOG: an international journal of obstetrics and gynaecology* 2015;**122**(Suppl S1):276.

Johnson 1975 {published data only}

Johnson JWC, Austin KL, Jones GS, Davis GH, King TM. Efficacy of 17alpha-hydroxyprogesterone caproate in the prevention of premature labor. *New England Journal of Medicine* 1975;**293**:675-80.

Klopper 1965 {published data only}

Klopper AI, MacNaughton MC. Hormones in recurrent abortion. *Journal of Obstetrics and Gynaecology of the British Commonwealth* 1965;**72**:1022-8.

Le Vine 1964 {published data only}

Le Vine L. Habitual abortion. A controlled clinical study of progestational therapy. *Western Journal of Surgery* 1964;**72**:30-6.

Luz 1988 [pers comm] {published data only}

Luz NP. Hormones in the treatment of patients with threatened early pregnancy loss. Letter to: Oxford Database of Perinatal Trials, John Radcliffe Infirmary, Oxford 22 October 1988.

Moller 1965 {published data only}

Moller KJA, Fuchs F. Double blind controlled trial of 6-methyl-17-acetoxyprogesterone in threatened abortion. *Journal of Obstetrics and Gynaecology of the British Commonwealth* 1965;**72**:1042-4.

Nyboe 2002 {published data only}

Nyboe Andersen A, Popovic-Todorovic B, Schmidt KT, Loft A, Lindhard A, Hojgaard A, et al. Progesterone supplementation during early gestations after IVF or ICSI has no effect on the delivery rates: a randomized controlled trial. *Human Reproduction* 2002;**17**(2):357-61.

Omar 2005 {published data only}

Omar MH, Mashita MK, Lim PS, Jamil MA. Dydrogesterone in threatened abortion: pregnancy outcome. *Journal of Steroid Biochemistry & Molecular Biology* 2005;**97**(5):421-5.

Porcaro 2015 {published data only}

Porcaro G, Brillo E, Giardina I, Di Iorio R. Alpha lipoic acid (ALA) effects on subchorionic hematoma: preliminary clinical results. *European Review for Medical and Pharmacological Sciences* 2015;**19**(18):3426-32.

Priehl 1992 {published data only}

Priehl G, Diedrich K, Van der Ven HH, Luckhaus J, Krebs D. The effect of 17alpha-hydroxyprogesterone caproate/oestradiol valerate on the development and outcome of early pregnancies following in vitro fertilization and embryo transfer: a prospective and randomized controlled trial. *Human Reproduction* 1992;**7**(1):1-5.

Pustotina 2018 {published data only}

Pustotina OA. Effectiveness of dydrogesterone, 17-OH progesterone and micronized progesterone in prevention of preterm birth in women with a short cervix. *Journal of Maternal-Fetal & Neonatal Medicine* 2018;**31**(14):1830-8.

Reijnders 1988 {published data only}

Reijnders FJL, Thomas CMG, Doesburg WH, Rolland R, Eskes TKAB. Endocrine effects of 17 alpha-hydroxyprogesterone caproate during early pregnancy: a double-blind clinical trial. *British Journal of Obstetrics and Gynaecology* 1988;**95**:462-8.

Shearman 1963 {published data only}

Shearman RP, Garrett WJ. Double-blind study of effect of 17-hydroxyprogesterone caproate on abortion rate. *BMJ* 1963;**1**:292-5.

Siew 2014 {published data only}

Siew JYS, Allen JC, Malhotra R, Ostbye T, He S, Tan TC. Progesterone for threatened miscarriage: a randomised controlled trial comparing micronised progesterone and dydrogesterone. *Journal of Maternal-Fetal & Neonatal Medicine* 2014;**27**(Suppl 1):243.

Siew 2015 {published data only}

Siew S, Hui CY, Tan TC, Allen JC, Malhotra R. Micronized progesterone compared with dydrogesterone for threatened miscarriage: a randomized controlled trial. *Obstetrics & Gynecology* 2015;**125**(5 Suppl):104S.

Smits 1992 {published data only}

Smits J, Devroey P, Faguer B, Bourgain C, Camus M, Van Steirteghem AC. Randomized prospective trial comparing supplementation of the luteal phase and of early pregnancy by natural progesterone given by intramuscular or vaginal administration. *Revue Francaise de Gynecologie et d Obstetrique* 1992;**87**:507-16.

Sondergaard 1985 {published data only}

Sondergaard F, Ottesen B, Detlefsen GU, Schierup L, Pederson SC, Lebech PE. Progesterone treatment of cases of threatened pre-term delivery in women with a low level of plasma progesterone. *Contraception Fertilite Sexualite* 1985;**13**:1227-31.

Song 2007 {published data only}

Song YL, Zhu LP. The fetus protection effects of Zhixue Baotai Decoction on women of early threatened abortion with dark area surrounding pregnancy sac. *Chinese Journal of Integrated Traditional and Western Medicine* 2007;**27**(11):1025-8.

Souka 1980 {published data only}

Souka AR, Osman M, Sibaie F, Einen MA. Therapeutic value of indomethacin in threatened abortion. *Prostaglandins* 1980;**19**:457-60.

Swyer 1953 {published data only}

Swyer GIM, Daley D. Progesterone implantation in habitual abortion. *British Medical Journal* 1953;**1**:1073-86.

Tognoni 1980 {published data only}

Tognoni G, Ferrario L, Inzalaco M, Crosignani PG. Progestagens in threatened abortion. *Lancet* 1980;**2**:1242-3.

Turner 1966 {published data only}

Turner SJ, Mizock GB, Feldman GL. Prolonged gynecologic and endocrine manifestations subsequent to administration of medroxyprogesterone acetate during pregnancy. *American Journal of Obstetrics and Gynecology* 1966;**95**:222-7.

Vincze 2006 {published data only}

Vincze E, Molnar GB, Foldesi I, Pal A. Treatment possibilities using progestagens and progesteron-like preparations in threatened abortion [A fenyegeto veteles kezelesenek lehetosegei progeszteron es progeszteron-szeru keszitmenyekkel]. *Magyar Noorvosok Lapja* 2006;**69**(4):281-4.

Zhang 2000 {published data only}

Zhang J, Zhang Y, Liu G. Clinical and experimental study on yun'an granule in treating threatened abortion. *Chung-Kuo Chung Hsi i Chieh Ho Tsa Chih* 2000;**20**(4):251-4.

References to studies awaiting assessment
Yadav 2015 {published data only}

Yadav R, Yadav S, Jain A. To study the impact of progesterone (dydrogesterone) on proinflammatory (IL-6 and TNF-alpha) and anti-inflammatory (IL-10) cytokines in threatened abortion. *International Journal of Gynecology and Obstetrics* 2015;**131**(Suppl 5):E196.

References to ongoing studies
ACTRN12611000405910 {published data only}

ACTRN12611000405910. Does using progesterone in threatened miscarriage increase the live birth rate? Supporting Threatened Outcomes with Progesterone (STOP trial). anzctr.org.au/Trial/Registration/TrialReview.aspx?ACTRN=12611000405910 (first received 19 April 2011).

McLindon L, Beckmann M. Supporting threatened outcomes with progesterone - the stop trial; a protocol. *BJOG: an international journal of obstetrics and gynaecology* 2015;**122**(Suppl S1):23.

IRCT201012035294N1 {published data only}

IRCT201012035294N1. To investigate the possible effect of Progesterone Supp on cytokines endocervical concentration of women at risk of abortion. en.search.irct.ir/view/4741 (first received 17 March 2012).

IRCT2015020217035N2 {published data only}

IRCT2015020217035N2. Effect of oral progesterone (dydrogestron) on incidence of glucose intolerance in pregnant females with threatened abortion. en.search.irct.ir/view/22059 (first received 19 September 2016).

IRCT2016040627248N1 {published data only}

IRCT2016040627248N1. The efficacy of a special food (fried egg with grape molasses) on threatened abortion: a randomized controlled trial. en.search.irct.ir/view/29438 (first received 9 July 2016).

NCT015018902011 {published data only}

NCT01501890. The impact of progesterone treatment on obstetrical outcome among women with first trimester vaginal bleeding. clinicaltrials.gov/show/NCT01501890 (first received 9 September 2011).

NCT02128685 {published data only}

Chan DM, Cheung KW, Yung SS, Lee VC, Li RH, Ng EH. A randomized double-blind controlled trial of the use of dydrogesterone in women with threatened miscarriage in the first trimester: study protocol for a randomized controlled trial. *Trials* 2016;**17**(1):408.

NCT02128685. A randomized double-blind controlled trial of use of dydrogesterone in women with threatened miscarriage in the first trimester. clinicaltrials.gov/show/NCT02128685 (first received 29 April 2014).

NCT02145767 {published data only}

NCT02145767. Progesterone for the prevention of miscarriage and preterm birth in women with first trimester bleeding: PREEMPT trial. clinicaltrials.gov/show/NCT02145767 (first received 14 May 2014).

NCT02633878 {published data only}

ChiCTR-IOR-15007526. Chinese herbal medicine and micronized progesterone for threatened miscarriage: an international cooperation multicenter randomized controlled trial. chictr.org.cn/showproj.aspx?proj=12677 (first received 12 March 2015).

NCT02633878. Chinese herbal medicine and micronized progesterone for threatened miscarriage RCT. clinicaltrials.gov/ct2/show/record/NCT02633878 (first received 7 December 2015).

NCT02690129 {published data only}

NCT02690129. Vaginal progesterone for treatment of threatened miscarriage; randomized clinical trial. clinicaltrials.gov/ct2/show/NCT02690129 (first received 19 February 2016).

NCT02950935 {published data only}

NCT02950935. Prospective, double-blind, randomised, placebo controlled, phase iii clinical study assessing the efficacy of natural progesterone 25 mg/bid administered subcutaneously in the maintenance of early pregnancy in women with symptoms of threatened abortion. clinicaltrials.gov/show/NCT02950935 (first received 27 October 2016).

Additional references
Arredondo 2006

Arredondo F, Noble LS. Endocrinology of recurrent pregnancy loss. *Seminars in Reproductive Medicine* 2006;**24**:33-9.

Blohm 2008

Blohm F, Friden B, Milsom I. A prospective longitudinal population-based study of clinical miscarriage in an urban Swedish population. *BJOG: an international journal of obstetrics and gynaecology* 2008;**115**:176-82.

Carp 2012

Carp H. A systematic review of dydrogesterone for the treatment of threatened miscarriage. *Gynecological Endocrinology* 2012;**28**(12):983-90.

Check 2011

Check JH. A practical approach to the prevention of miscarriage: part 5--antiphospholipid syndrome as a cause of spontaneous abortion. *Clinical and Experimental Obstetrics and Gynecology* 2011;**38**:5-9.

Cohain 2017

Cohain JS, Buxbaum RE, Mankuta D. Spontaneous first trimester miscarriage rates per woman among parous women with 1 or more pregnancies of 24 weeks or more. *BMC Pregnancy and Childbirth* 2017;**17**(1):437.

Coomarasamy 2015

Coomarasamy A, Williams H, Truchanowicz E, Seed PT, Small R, Quenby S, et al. A randomized trial of progesterone in women with recurrent miscarriages. *New England Journal of Medicine* 2015;**373**(22):2141-8.

Cunningham 2001a

Cunningham FG. Section IX. Reproductive success and failure. Williams Obstetrics. 21st Edition. New York: McGraw-Hill, 2001.

Cunningham 2001b

Cunningham FG, editor(s). Physiology of pregnancy. Williams Obstetrics. 21st Edition. New York: McGraw-Hill, 2001.

Dabash 2015

Dabash R, Chelli H, Hajri S, Shochet T, Raghavan S, Winikoff B. A double-blind randomized controlled trial of mifepristone or placebo before buccal misoprostol for abortion at 14-21 weeks of pregnancy. *International Journal of Gynecology & Obstetrics* 2015;**130**(1):40-4.

De La Rochebrochard 2002

De La Rochebrochard E, Thonneau P. Paternal age and maternal age are risk factors for miscarriage; results of a multicentre European study. *Human Reproduction* 2002;**17**:1649-56.

Dede 2010

Dede FS, Ulucay U, Kose MF, Dede H, Dilbaz S. Fetal loss in threatened abortion after demonstration of fetal cardiac activity in a low socioeconomic population. *Journal of Obstetrics and Gynaecology* 2010;**30**:622-5.

Deeks 2011

Deeks JJ, Higgins JPT, Altman DG (editors). Chapter 9: Analysing data and undertaking meta-analyses. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Di Renzo 2005

Di Renzo GC, Rosati A, Mattei A, Gojinic M, Gerli S. The changing role of progesterone in preterm labour. *BJOG: an international journal of obstetrics and gynaecology* 2005;**112** Suppl 1:57-60.

Facchinetti 2017

Facchinetti F, Vergani P, Di Tommaso M, Marozio L, Acaia B, Vicini R, et al. Progestogens for maintenance tocolysis in women with a short cervix: a randomized controlled trial. *Obstetrics and Gynecology* 2017;**130**(1):64-70.

Galindo 2006

Galindo A, Burguillo AG, Azriel S, Fuente P de L. Outcome of fetuses in women with pregestational diabetes mellitus. *Journal of Perinatal Medicine* 2006;**34**(4):323-31.

GRADE 2013

Editors: Schünemann H, Brożek J, Guyatt G, Oxman A. Handbook for grading the quality of evidence and the strength of recommendations using the GRADE approach. Updated October 2013. Available from gdt.guidelinedevelopment.org/app/handbook/handbook.html.

GRADEpro GDT 2015 [Computer program]

McMaster University (developed by Evidence Prime). GRADE pro GDT. Version 'date accessed' 12/09/17. Hamilton (ON): McMaster University (developed by Evidence Prime), 2015.

Haas 2018

Haas DM, Hathaway TJ, Ramsey PS. Progestogen for preventing miscarriage in women with recurrent miscarriage of unclear etiology. *Cochrane Database of Systematic Reviews* 2018, Issue [in press].

Hahn 2014

Hahn KA, Hatch EE, Rothman KJ, Mikkelsen EM, Brogly SB, Sørensen HT, et al. Body size and risk of spontaneous abortion among Danish pregnancy planners. *Paediatric and Perinatal Epidemiology* 2014;**28**(5):412-23.

Heyborne 2016

Heyborne KD. 17- α hydroxyprogesterone caproate for the prevention of recurrent preterm birth: one size may not fit all. *Obstetrics & Gynecology* 2016;**128**(4):899-90.

Higgins 2003

Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**:557-60.

Higgins 2011a

Higgins JPT, Green S, editors. *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from www.handbook.cochrane.org.

Higgins 2011b

Higgins JPT, Altman DG, Sterne JAC (editors). Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Higgins 2011c

Higgins JPT, Deeks JJ, Altman DG (editors). Chapter 16: Special topics in statistics. In: Higgins JPT, Green S (editors), *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Jin 2006

Jin S, Li SW, Long J, Li L, Tan ZJ. The role of progesterone in human early pregnancy is mediated by insulin-like growth factors binding protein1-3. *Sichuan Da Xue Xue Bao Yi Xue Ban* 2006;**37**:399-403.

Kalinka 2005

Kalinka J, Szekeres-Bartho J. The impact of dydrogesterone supplementation on hormonal profile and progesterone-induced blocking factor concentrations in women with threatened abortion. *American Journal of Reproductive Immunology* 2005;**53**:166-71.

Källén 2010

Källén B, Finnström O, Lindam A, Nilsson E, Nygren KG, Otterblad PO. Congenital malformations in infants born after in vitro fertilization in Sweden. *Birth Defects Research. Part A Clinical and Molecular Teratology* 2010;**88**:137-43.

Lede 2005

Lede R, Duley L. Uterine muscle relaxant drugs for threatened miscarriage. *Cochrane Database of Systematic Reviews* 2005, Issue 3. [DOI: [10.1002/14651858.CD002857.pub2](https://doi.org/10.1002/14651858.CD002857.pub2)]

Makrydimas 2003

Makrydimas G, Sebire NJ, Lolis D, Vlassis N, Nicolaidis KH. Fetal loss following ultrasound diagnosis of a live fetus at 6-10 weeks of gestation. *Ultrasound in Obstetrics and Gynecology* 2003;**22**:368-72.

Marcinko 2011

Marcinko VM, Marcinko D, Dordevic V, Oreskovic S. Anxiety and depression in pregnant women with previous history of spontaneous abortion. *Collegium Antropologicum* 2011;**35 Suppl 1**:225-8.

Mirza 2016

Mirza FG, Patki A, Pexman-Fieth C. Dydrogesterone use in early pregnancy. *Gynecological Endocrinology* 2016;**32**(2):97-106.

Nigro 2011

Nigro G, Mazzocco M, Mattia E, Di Renzo GC, Carta G, Anceschi MM. Role of the infections in recurrent spontaneous abortion. *Journal of Maternal Fetal Neonatal Medicine* 2011;**24**:983-9.

O'Brien 2016

O'Brien JM, Lewis DF. Prevention of preterm birth with vaginal progesterone or 17-alpha-hydroxyprogesterone caproate: a critical examination of efficacy and safety. *American Journal of Obstetrics and Gynecology* 2016;**214**(1):45-56.

Oler 2017

Oler E, Eke AC, Hesson A. Meta-analysis of randomized controlled trials comparing 17 α -hydroxyprogesterone caproate and vaginal progesterone for the prevention of recurrent spontaneous preterm delivery. *International Journal of Gynecology & Obstetrics* 2017;**138**(1):12-6.

Osmanağaoğlu 2010

Osmanağaoğlu MA, Erdoğan I, Eminağaoğlu S, Karahan SC, Özgün S, Can G, et al. The diagnostic value of beta-human chorionic gonadotropin, progesterone, CA125 in the prediction of abortions. *Journal of Obstetrics and Gynaecology* 2010;**30**:288-93.

Owen 2004

Owen J, Yost N, Berghella V, MacPherson C, Swain M, Dildy GA 3rd, et al. Can shortened midtrimester cervical length predict very early spontaneous preterm birth?. *American Journal of Obstetrics and Gynecology* 2004;**191**(1):298-303.

RevMan 2014 [Computer program]

Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager 5 (RevMan 5). Version 5.3. Copenhagen: Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Rocchetti 2011

Rocchetti TT, Marconi C, Rall VL, Borges VT, Corrente JE, Da Silva MG. Group B streptococci colonization in pregnant women: risk factors and evaluation of the vaginal flora. *Archives of Gynecology and Obstetrics* 2011;**283**:717-21.

Rust 2005

Rust OA, Atlas RO, Kimmel S, Roberts WE, Hess LW. Does the presence of a funnel increase the risk of adverse perinatal outcome in a patient with a short cervix?. *American Journal of Obstetrics and Gynecology* 2005;**192**(4):1060-6.

Saccone 2017

Saccone G, Schoen C, Franasiak JM, Scott RT Jr, Berghella V. Supplementation with progestogens in the first trimester of pregnancy to prevent miscarriage in women with unexplained recurrent miscarriage: a systematic review and meta-analysis of randomized, controlled trials. *Fertility and Sterility* 2017;**107**(2):430-8.

Saraswat 2010

Saraswat L, Bhattacharya S, Maheshwari A, Bhattacharya S. Maternal and perinatal outcome in women with threatened miscarriage in the first trimester: a systematic review. *BJOG: an international journal of obstetrics and gynaecology* 2010;**117**:245-57.

Silver 1999

Silver RI, Rodriguez R, Chang TS, Gearhart JP. In vitro fertilization is associated with an increased risk of hypospadias. *Journal of Urology* 1999;**161**(6):1954-7.

Sterne 2011

Sterne JAC, Egger M, Moher D (editors). Chapter 10: Addressing reporting biases. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Intervention*. Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Sugiura-Ogasawara 2009

Sugiura-Ogasawara M, Ozaki Y, Kitaori T, Suzumori N, Obayashi S, Suzuki S. Live birth rate according to maternal age and previous number of recurrent miscarriages. *American Journal of Reproductive Immunology* 2009;**62**:314-9.

Suzumori 2010

Suzumori N, Sugiura-Ogasawara M. Genetic factors as a cause of miscarriage. *Current Medicinal Chemistry* 2010;**17**:3431-7.

Szekeres-Bartho 2008

Szekeres-Bartho J, Wilczynski JR, Basta P, Kalinka J. Role of progesterone and progestin therapy in threatened abortion and preterm labour. *Frontiers in Bioscience* 2008;**13**:1981-90.

Szekeres-Bartho 2009

Szekeres-Bartho J. Progesterone-mediated immunomodulation in pregnancy: its relevance to leukocyte immunotherapy of recurrent miscarriage. *Immunotherapy* 2009;**1**:873-82.

Szekeres-Bartho 2010

Szekeres-Bartho J, Polgar B. PIBF: the double edged sword. Pregnancy and tumor. *American Journal of Reproductive Immunology* 2010;**64**(2):77-86.

Temmerman 1992

Temmerman M, Lopita MI, Sanghvi HC, Sinei SK, Plummer FA, Piot P. The role of maternal syphilis, gonorrhoea and HIV-1 infection in spontaneous abortion. *International Journal of Sexually Transmitted Diseases and AIDS* 1992;**3**:418-22.

Vorsanova 2005

Vorsanova SG, Kolotii AD, Iourov IY, Monakhov VV, Kirillova EA, Soloviev IV, et al. Evidence for high frequency of chromosomal mosaicism in spontaneous abortions revealed by interphase FISH analysis. *Journal of Histochemistry and Cytochemistry* 2005;**53**:375-80.

World Bank 2017

World Bank. World Bank list of economies. iccmoot.com/wp-content/uploads/2017/07/World-Bank-List-of-Economies.pdf.

Zegers-Hochschild 2009

Zegers-Hochschild F, Adamson GD, De MJ, Ishihara O, Mansour R, Nygren K, et al. International Committee for Monitoring Assisted Reproductive Technology (ICMART) and the World Health Organization (WHO) revised glossary of ART terminology. *Fertility and Sterility* 2009;**92**:1520-4.

References to other published versions of this review
Wahabi 2011

Wahabi HA, Fayed AA, Esmaeil SA, Al Zeidan RA. Progestogen for treating threatened miscarriage. *Cochrane Database of Systematic Reviews* 2011, Issue 12. [DOI: [10.1002/14651858.CD005943.pub4](https://doi.org/10.1002/14651858.CD005943.pub4)]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES
Characteristics of included studies [ordered by study ID]

Alimohamadi 2013

Methods	<p>Design: RCT, double-blind, placebo-controlled, parallel-group</p> <p>Recruitment method: unclear</p> <p>Method of randomisation: computer-generated block randomisation</p> <p>Setting: Obstetrics and Gynaecology ward of Vali-e-Asr teaching hospital in Tehran</p>
Participants	<p>Inclusion: pregnant women with clinical symptoms of threatened abortion before 20th week of pregnancy.</p> <p>Exclusion: women with systemic diseases, including maternal hypertension before or during pregnancy, cardiac disease, renal disease, genital tract anomalies of the mother or/and diabetes. Women with uterine tenderness, congenital defects of the fetus, and those who had used a progestational drug during pregnancy, prior to being recruited into the study</p> <p>Participants randomised: 160</p>
Interventions	<p>Intervention group: (80 women) vaginal progesterone, 200 mg twice daily for 1 week</p> <p>Control group: (80 women) placebo with the same description and duration of treatment as experimental group</p>
Outcomes	<p>Endocervical concentrations of different cytokines</p> <p>Miscarriage</p> <p>Preterm delivery</p> <p>Neonatal death</p> <p>Low birth weight</p> <p>Respiratory distress syndrome</p>
Notes	<p>Journal: Journal of Reproductive Immunology</p> <p>Year of publication: 2013</p> <p>Country: Iran</p> <p>Income status of the country: upper-middle-income</p> <p>Source of funding: not mentioned</p> <p>Conflict of interest: no conflict of interest</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated block randomisation
Allocation concealment (selection bias)	Low risk	Pharmacy prepared
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Participants and clinicians were unaware of allocation

Alimohamadi 2013 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Blinded assessment
Incomplete outcome data (attrition bias) All outcomes	Low risk	Incomplete outcome data of 15 participants (9.4%); 8 in intervention and 7 in control groups (balanced attrition)
Selective reporting (reporting bias)	Low risk	All outcomes were reported
Other bias	Low risk	No other risk of biases

El-Zibdeh 2009

Methods	<p>Design: quasi-experimental</p> <p>Recruitment method: pregnant women consecutively presented to the clinic during study period</p> <p>Method of randomisation: according to the day of the week the women attended the clinic; women attending on Saturday, Monday, and Wednesday were assigned to intervention group and those attending on Sunday, Tuesday, and Thursday were assigned to the control group</p> <p>Setting: Amman Islamic Hospital clinic</p>
Participants	<p>Inclusion: women with threatened miscarriage (mild or moderate vaginal bleeding during the first trimester of pregnancy)</p> <p>Exclusion: presence of a systemic illness or fever, the suspected passage of any fetal or pregnancy materials, and the absence of a normal gestational sac at 5 weeks gestational age, a yolk sac at 5.5–6 weeks gestational age, an embryo at 6–6.5 weeks gestational age or cardiac activity at 7 weeks gestational age</p> <p>Participants randomised: 146</p>
Interventions	<p>Intervention group: (86 women) synthetic progesterone, dydrogesterone, oral 10 mg twice daily. Until 1 week after the bleeding stopped. Standard supportive care, including iron, folic acid and multivitamin supplements and bed rest</p> <p>Control group: (60 women) no treatment. Standard supportive care, including iron, folic acid and multivitamin supplements and bed rest</p>
Outcomes	<p>Miscarriage</p> <p>Preterm delivery</p> <p>Fetal structural malformations, including genital malformations</p> <p>Maternal hypertension</p> <p>Intra-uterine growth restriction</p> <p>APH</p>
Notes	<p>Journal: Maturitas</p> <p>Year of publication: 2009</p> <p>Country: Jordan</p>

El-Zibdeh 2009 (Continued)

Income status of the country: lower-middle-income

Source of funding: Solvay Pharmaceuticals

Conflict of interest: no actual or potential conflict

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	According to the day of the week the women presented to the clinic. Women attending the clinic on Saturday, Monday or Wednesday were allocated to the intervention group and those attending on Sunday, Tuesday or Thursday were allocated to the control group.
Allocation concealment (selection bias)	High risk	The randomisation was performed by the attending physician, who also gave the treatment to the women.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No blinding of the participants and study personnel
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Blinding was ensured for the outcome assessors.
Incomplete outcome data (attrition bias) All outcomes	Low risk	No loss to follow-up
Selective reporting (reporting bias)	Low risk	All the prespecified outcomes were addressed.
Other bias	High risk	The difference in the number of participants recruited to experimental and the control groups (86 versus 60) might have introduced bias.

Gerhard 1987

Methods	<p>Design: RCT, double-blind</p> <p>Recruitment method: women with vaginal bleeding were referred to the hospital</p> <p>Method of randomisation: unclear</p> <p>Setting: Women's Hospital, University of Heidelberg</p>
Participants	<p>Inclusion: vaginal bleeding and the internal cervical os being closed</p> <p>Exclusion: none</p> <p>Participants randomised: 35</p>
Interventions	<p>Intervention group: (17 women) 1 vaginal suppository twice daily, containing 25 mg progesterone, and bed rest</p> <p>Control group: (18 women) 1 vaginal suppository twice daily, containing polyethylene glycol, and bed rest</p>

Gerhard 1987 (Continued)

Outcomes

Miscarriage

14 days of being symptom free

Preterm delivery

Notes

Journal: Biological Research in Pregnancy

Year of publication: 1987

Country: Germany

Income status of the country: high-income

Source of funding: not mentioned

Conflict of interest: not mentioned

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	The method of randomisation was not mentioned
Allocation concealment (selection bias)	Unclear risk	The method of allocation concealment was not mentioned
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Blinding of participants and study personnel
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Blinding of the outcome assessors
Incomplete outcome data (attrition bias) All outcomes	Low risk	2 women (5.7%) were omitted from the study after randomisation
Selective reporting (reporting bias)	Low risk	All the prespecified outcomes were addressed
Other bias	Unclear risk	As we included only participants with evidence of fetal viability, which amounted to only half the participants, we are unclear how much bias was introduced.

Palagiano 2004

Methods

Design: RCT

Recruitment method: unclear

Method of randomisation: unclear

Setting: a medical centre

Palagianò 2004 (Continued)

Participants	<p>Inclusion: women with threatened abortion; viable baby; amenorrhoea 6-12 weeks; closed uterine cervix</p> <p>Exclusion: women with previous adequate luteal phase; women who were using hormonal treatment or other drugs affecting uterine contractility; women with vaginal infection; absence of embryo's heart-beat; open cervix (> 2 cm measured by U/S); embryo's size 1 week more than the corresponding amenorrhoea.</p> <p>Participants randomised: 50</p>
Interventions	<p>Intervention group: (25 women) 90 mg vaginal progesterone once daily for 5 days</p> <p>Control group: (25 women) placebo</p>
Outcomes	<p>Pain score</p> <p>Uterine contractility</p> <p>Abortion rate</p> <p>(Followed for 60 days for the occurrence of miscarriage and for 5 days for the other outcomes)</p>
Notes	<p>Journal: Annals New York Academy of Sciences</p> <p>Year of publication: 2004</p> <p>Country: Italy</p> <p>Income status of the country: high-income</p> <p>Source of funding: not mentioned</p> <p>Conflict of interest: not mentioned</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	This is a randomised trial but the method of randomisation was not mentioned
Allocation concealment (selection bias)	Low risk	Code numbers were attached to each container of the gel. The researcher recorded the progressive number of the container given to the patients enrolled, who received a correspondent code. Only 3 months after the end of the database recording, the codes were open and showed to the Data Monitoring Board (DMB) to be evaluated by a statistician.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Participants and study personnel were blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The effect on pain symptom amelioration was evaluated by a 5-score intensity grading, whereas uterine contractions were evaluated by standard transvaginal ultrasound, the code was broken only after the analysis by the statistician.
Incomplete outcome data (attrition bias) All outcomes	Low risk	No loss to follow-up

Palagiano 2004 (Continued)

Selective reporting (reporting bias)	High risk	In the methods study authors mentioned they will report all adverse effects, but there was no explicit mention of the absence or presence of any adverse effects in the results section.
Other bias	Low risk	No risk of other biases

Pandian 2009

Methods	<p>Design: RCT</p> <p>Recruitment method: all women presenting with vaginal bleeding up to 16 weeks of pregnancy were assessed for inclusion</p> <p>Method of randomisation: sealed envelopes picked by the participants</p> <p>Setting: Department of Obstetrics and Gynaecology, Seberang Jaya Hospital</p>
Participants	<p>Inclusion: women with threatened abortion. Viability of fetus confirmed by U/S. no systematic illness or fever and no loss of conception tissue</p> <p>Exclusion: women with recurrent miscarriage (> 3), heavy bleeding, cervical polyp, multiple gestation, empty sac > 26 mm</p> <p>Participants randomised: 191 women</p>
Interventions	<p>Intervention group: (96 women) 40 mg followed by 10 mg twice daily</p> <p>Control group: (95 women) conservative treatment with bed rest only</p>
Outcomes	<p>Miscarriage rate</p> <p>Placenta previa</p> <p>Preterm delivery</p> <p>Congenital anomalies</p> <p>APH</p> <p>Caesarean section</p> <p>Low birth weight</p> <p>Preganancy induced hypertension</p> <p>Perinatal death</p>
Notes	<p>Journal: Maturitas</p> <p>Year of publication: 2009</p> <p>Country: Malaysia</p> <p>Income status of the country: upper middle-income</p> <p>Source of funding: Solvay Pharmaceutical Company</p> <p>Conflict of interest: no conflicts</p>

Risk of bias
Progestogen for treating threatened miscarriage (Review)

Pandian 2009 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random allocation of each women using sealed envelope
Allocation concealment (selection bias)	Low risk	Concealed envelope
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants and the study personnel were not blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Only the statistician was blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	There was no loss of follow-up
Selective reporting (reporting bias)	Low risk	All the prespecified outcomes were addressed
Other bias	Low risk	No apparent other sources of bias

Turgal 2017

Methods	<p>Design: RCT</p> <p>Recruitment method: through assessment of women presented to the hospital</p> <p>Method of randomisation: computerised random number generator program</p> <p>Setting: a university hospital</p>
Participants	<p>Inclusion: women with threatened abortion and with presence singleton pregnancy and live embryo, before 9 weeks of gestation</p> <p>Exclusion: non viable fetus, twin pregnancy, presence of subchorionic haematoma and history of hypertension, diabetes mellitus, severe hepatic disorders, uterine leiomyoma, congenital uterine anomaly and recurrent pregnancy loss</p> <p>Participants randomised: 83</p>
Interventions	<p>Intervention group: (42 women) oral micronised progesterone, 400 mg/d for 4 weeks</p> <p>Control group: (41 women) placebo</p>
Outcomes	<p>Primary outcome:</p> <ul style="list-style-type: none"> fetal-placental volume change <p>Secondary outcomes:</p> <ul style="list-style-type: none"> miscarriage rate live birth rate preterm birth

Turgal 2017 (Continued)

- birthweight
- LBW
- intrauterine growth restriction
- neonatal respiratory distress syndrome

Notes

Journal: Journal of clinical Ultrasound

Year of publication: 2017

Country: Turkey

Income status of the country: upper middle-income

Source of funding: not mentioned

Conflict of interest: not mentioned

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computerised random number by using generator program
Allocation concealment (selection bias)	Unclear risk	Unclear
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Placebo was used
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Outcome is not affected by blinding of outcomes assessors
Incomplete outcome data (attrition bias) All outcomes	Low risk	Incomplete outcome data in 12 women (14.5%), equal in both groups
Selective reporting (reporting bias)	Low risk	All the prespecified outcomes were addressed
Other bias	Low risk	No other biases

Yassaee 2014

Methods

Design: RCT

Recruitment method: through assessment of women presented to the hospital

Method of randomisation: unclear

Setting: Taleghani Hospital affiliated to Shahid Beheshti University of Medical Sciences

Participants

Inclusion: pregnant women with threatened abortion. The presence of singleton pregnancy and detection of fetal heart activity, besides gestational age of < 20 weeks was verified by U/S

Yassaee 2014 (Continued)

Exclusion: women were excluded if they had reaction to Cyclogest, repeated abortions, multiple gestation, absence of fetus or fetal heart tone, uterine anomaly or fetal anomaly

Participants randomised: 60 women

Interventions	<p>Intervention group: 400 mg of vaginal progesterone suppository (Cyclogest) each day until their bleeding stopped in < 1 week</p> <p>Control group: no treatment</p>
Outcomes	<p>Successful delivery</p> <p>Miscarriage rates</p> <p>Pain relief</p>
Notes	<p>Journal: Journal of Reproduction and Infertility</p> <p>Year of publication: 2014</p> <p>Country: Iran</p> <p>Income status of the country: upper middle-income</p> <p>Source of funding: not mentioned</p> <p>Conflict of interest: no conflicts</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details on random allocation
Allocation concealment (selection bias)	Unclear risk	Unclear
Blinding of participants and personnel (performance bias) All outcomes	High risk	There was no placebo, so participants were not blind
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Assessors were blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	No loss of participants to follow-up
Selective reporting (reporting bias)	Low risk	All the prespecified outcomes were addressed
Other bias	Low risk	No apparent other risk of bias

APH: antepartum haemorrhage; **LBW:** low birthweight; **RCT:** randomised controlled trial; **U/S:** ultrasound

Characteristics of excluded studies [ordered by study ID]

Progestogen for treating threatened miscarriage (Review)

Study	Reason for exclusion
Akhtar 2012	Not RCT
Beigi 2016	Comparing 2 types of progestogens
Berle 1977	Combination therapy of progesterone and oestrogen was used in this study
Berle 1980	Viability of the fetus was not confirmed before commencement of progesterone treatment
Brenner 1962	Different population
Check 1995	Combination therapy of progestogen and immunotherapy
Chye 2014	Compares 2 types of progestogens
Corrado 2002	Different population
Costantino 2016	Participants were mixed group of women with threatened abortion and subchorionic haematoma, results of intervention cannot be obtained separately
Crowder 1950	Viability was not confirmed before the commencement of the study
Czajkowski 2007	The study compared 2 types of progestogen
El Zibdeh 2000	Different population. Population with recurrent miscarriage rather than threatened miscarriage
El Zibdeh 2002	Different population. Population with recurrent miscarriage rather than threatened miscarriage
El Zibdeh 2005	Different population. Population with recurrent miscarriage rather than threatened miscarriage
Famina 2015	Comparing 2 types of progestogens
Fuchs 1966	Different population. Patients with recurrent miscarriage not threatened miscarriage
Goldzieher 1964	Different population. Population with recurrent miscarriage rather than threatened miscarriage
Govaerts-Videtzky 1965	Viability was not confirmed before the commencement of the study
Hui 2015	Comparing 2 types of progestogens
Johnson 1975	Different population. Progestogen was used for the prevention of miscarriage in women with recurrent miscarriage or preterm delivery not threatened miscarriage
Klopper 1965	Different population. Progestogen was used for the prevention of miscarriage in women with recurrent miscarriage not threatened miscarriage
Le Vine 1964	Different population. Progestogen was used for the prevention of miscarriage in women with recurrent miscarriage not threatened miscarriage
Luz 1988 [pers comm]	Combination therapy of progestogen and oestrogen was used in this study
Moller 1965	Viability was not confirmed before the commencement of the study
Nyboe 2002	Different population. Progestogen was used for the prevention of miscarriage rather than the treatment of threatened miscarriage

Study	Reason for exclusion
Omar 2005	Loss to follow-up was more than 20%
Porcaro 2015	Different population
Prietl 1992	Different population. Progesterone was used for the prevention of miscarriage rather than the treatment of threatened miscarriage in IVF patients
Pustotina 2018	Comparing 2 types of progestogens
Reijnders 1988	Different population. Progestogen was used for the prevention not the treatment of miscarriage
Shearman 1963	Different population. Progestogen was used for population with recurrent miscarriage not threatened miscarriage
Siew 2014	Comparing 2 types of progestogen.
Siew 2015	Comparing 2 types of progestogens
Smitz 1992	Different population. Progestogen was used for the prevention of miscarriage rather than the treatment of threatened miscarriage in IVF patients
Sondergaard 1985	Viability was not confirmed before the commencement of the study.
Song 2007	Combination progestogen with vitamin E
Souka 1980	Viability was not confirmed before the commencement of the study
Swyer 1953	Different population. Progestogen was used for population with recurrent miscarriage not threatened miscarriage
Tognoni 1980	Viability was not confirmed before the commencement of the study
Turner 1966	Different population. The study participants do not have threatened miscarriage
Vincze 2006	Comparing 2 types of progesterone
Zhang 2000	Combination progesterone with vitamin E

IVF: in vitro fertilisation; **RCT:** randomised controlled trial

Characteristics of studies awaiting assessment [ordered by study ID]

Yadav 2015

Methods	RCT
Participants	100
Interventions	Dydrogesterone, 30 mg/d
Outcomes	Serum levels of IL-6, TNF-alpha and IL-10 were measured at the time of admission, 10 days, 14 weeks
	Abortion rate

Yadav 2015 (Continued)

Notes Only abstract is available, waiting response from trial authors

RCT: randomised controlled trial

Characteristics of ongoing studies [ordered by study ID]

ACTRN12611000405910

Trial name or title	Does using progesterone in threatened miscarriage increase the live birth rate? Supporting threatened outcomes with progesterone
Methods	RCT intervention model: parallel assignment. Blinding (masking use)
Participants	386 women age: ≥ 18 Inclusion criteria: threatened miscarriage, live intrauterine pregnancy; gestation < 10 weeks + 0 days
Interventions	Intervention group: progesterone pessary 400 mg nightly, until 12 weeks + 0 days Control group: placebo
Outcomes	Primary outcome: <ul style="list-style-type: none">live birth rate
Starting date	Jan 2012
Contact information	Dr Luke McLindon address Mater Mothers' Hospital, Raymond Terrace, South Brisbane, Queensland, 4101
Notes	

IRCT201012035294N

Trial name or title	Clinical trial comparison of progesterone suppository and placebo on the concentration of cervical inflammatory cytokines in patients at risk of miscarriage
Methods	Randomised, parallel, double-blinded study
Participants	All women who are 20 weeks pregnant and who have been referred for pain or increased cervical secretion and spotting are included in the study.
Interventions	Intervention group: progesterone suppository 400 mg twice/d for 1 week Control group: placebo
Outcomes	Primary <ul style="list-style-type: none">miscarriage rate Secondary <ul style="list-style-type: none">measurement of cervical inflammatory cytokines
Starting date	June 2006

Progestogen for treating threatened miscarriage (Review)

IRCT201012035294N (Continued)

Contact information Dr Sedigheh Hanustah Zadeh
Valiasr Hospital, Tehran University of Medical Sciences
Iran, Tehran
00982166581617
hantoushzadeh@tums.ac.ir

Notes

IRCT2015020217035N2

Trial name or title Effect of oral progesterone (dydrogesterone) on incidence of glucose intolerance in pregnant females with threatened miscarriage

Methods **Allocation:** randomised double blind placebo controlled trial.
Intervention model: parallel assignment.
Masking: triple (participant, care provider, investigator).
Primary purpose: treatment.

Participants A total of 100 women who are 14 weeks pregnant and presented with threatened miscarriage will be recruited. Participants will be tested to exclude abnormal blood glucose level. Women will be randomised to have either oral Dydrogesterone 10 mg twice daily or placebo for the control group.
Inclusion criteria: singleton pregnancy, the absence of any systemic disease, age < 30 years, BMI < 25, no history of stillbirth, no family history of diabetes, non-smoking,
Exclusion criteria: history of diabetes, gestational diabetes and other systemic diseases, any fetal abnormalities on ultrasound examination, multiple pregnancy, and previous macrosomia.
Interventions: oral administration of dydrogesterone 10 mg twice daily or placebo.

Interventions **Intervention group:** oral dydrogesterone 10 mg twice daily
Control group: placebo

Outcomes Treatment of threatened miscarriage and preterm delivery
To study the effect of oral progesterone (dydrogesterone) on incidence of glucose intolerance in pregnant women with threatened miscarriage

Starting date March 2015

Contact information Hafez Hospital
Shiraz
Fars
Iran, Islamic Republic Of
009871361222220
00989173138847

IRCT2015020217035N2 (Continued)

maryamkasraeian@gmail.com

Notes

IRCT2016040627248N1

Trial name or title	The efficacy of a special food (fried egg with grape molasses) on threatened miscarriage: a randomised controlled trial
Methods	RCT, single-blind, no placebo
Participants	90 women Inclusion criteria: pregnant women, (up to 12 weeks), presenting with mild-moderate vaginal bleeding with normal U/S and closed cervix in vaginal examination; singleton pregnancy; and who consent to participate in the study and are ready to consume the special food.
Interventions	Intervention group: in addition to intravaginal Cyclogest, the food recipe of fried eggs with butter and grape molasses will be given to women. The other conditions of intervention group, including the duration of eating this food, are similar to the control group.
Outcomes	Primary outcome: <ul style="list-style-type: none"> determine the miscarriage rate (loss of an intrauterine pregnancy at = 20 weeks' gestation) in patients with threatened miscarriage. Timepoint: weekly follow-up until vaginal bleeding has stopped and follow-up at 20 weeks of pregnancy Secondary outcome: <ul style="list-style-type: none"> the average duration of vaginal bleeding
Starting date	July 2016
Contact information	Dr Fatemeh Moradi School of Traditional medicine, Tehran University of Medical Sciences, SarParast St., Taleghani St., ValiAsr Blvd. 1449614353 Tehran Iran, Islamic Republic Of Tel:00982166976527 e-mail:dfmoradi@gmail.com
Notes	

NCT015018902011

Trial name or title	The impact of progesterone treatment on obstetrical outcome among women with first trimester vaginal bleeding
Methods	Allocation: randomised Intervention model: parallel assignment Masking: triple (participant, care provider, investigator) Primary purpose: prevention
Participants	Women attending the Gynecological Emergency Unit due to first trimester vaginal bleeding, with a viable singleton pregnancy at a gestational age of 6-13 completed weeks of gestation
Interventions	Intervention group: progesterone

Progestogen for treating threatened miscarriage (Review)

NCT015018902011 (Continued)

Control group: placebo

Outcomes	<p>Primary outcome measures:</p> <ul style="list-style-type: none"> • adverse pregnancy outcomes • miscarriage (also gestational age of miscarriage) • sonographic intrauterine hematoma • IUGR • placenta previa • pregnancy-induced hypertension and pre-eclampsia/eclampsia, • gestational age at delivery, preterm delivery (before 37 weeks); Early preterm delivery (before 34 weeks); and very early (before 28 weeks) • mode of delivery • placental abruption • PPROM • induction of labour • PPH • Apgar score • umbilical cord blood PH at birth • birthweight • fetal malformations • perinatal mortality • admission to the neonatal unit <p>Secondary outcome measures:</p> <ul style="list-style-type: none"> • uterine artery blood flow velocimetry, systolic to diastolic ratio, pulsatility index, resistance index and peak systolic velocity • placental pathological examination; placental weight and presence of infarcts, calcifications, fibrin deposits or signs of inflammation
Starting date	January 2012
Contact information	Ralika Hershkovitch, MD Soroka University Medical Center Beer Sheva, Israel
Notes	

NCT02128685

Trial name or title	A randomised double-blind controlled trial of the use of dydrogesterone in women with threatened miscarriage in the first trimester: study protocol for a randomised controlled trial
Methods	Double-blind, RCT
Participants	A total of 400 women presenting with first-trimester threatened miscarriage will be enrolled
Interventions	<p>Intervention group: Dydrogesterone 40 mg orally, followed by 10 mg orally 3 times/d</p> <p>Control group: placebo</p>
Outcomes	<ul style="list-style-type: none"> • Primary outcome: percentage of miscarriage before 20 weeks of gestation

NCT02128685 (Continued)

Starting date	Registered on 29 April 2014
Contact information	Department of Obstetrics and Gynaecology, Queen Mary Hospital, 102 Pokfulam Road, Hong Kong Special Administrative Region, China. dcmanka@gmail.com
Notes	

NCT02145767

Trial name or title	Progesterone for the prevention of miscarriage and preterm delivery in women with first trimester bleeding: PREEMPT trial
Methods	<p>Allocation: randomised</p> <p>Intervention model: parallel assignment</p> <p>Masking: quadruple (participant, care provider, investigator, outcomes assessor)</p> <p>Primary purpose: prevention</p>
Participants	<p>Ages eligible for study: 18 years-45 years</p> <p>Accepts healthy volunteers: yes</p> <p>Live intrauterine singleton pregnancy of < 14 weeks by crown-rump length on U/S with documented fetal cardiac activity presence of progesterational (subchorionic) hematoma on U/S</p>
Interventions	<p>Intervention group: progesterone 200 mg suppository administered vaginally at bedtime until 34 completed weeks of pregnancy</p> <p>Control group: placebo similar appearing suppository containing vehicle alone administered vaginally at bedtime until 34 completed weeks of pregnancy</p>
Outcomes	<p>Primary outcomes</p> <ul style="list-style-type: none"> • miscarriage • preterm delivery (< 37 weeks) <p>Secondary outcomes:</p> <ul style="list-style-type: none"> • maternal outcomes <ul style="list-style-type: none"> * antenatal admissions * treatment of preterm delivery, etc • Neonatal outcomes: <ul style="list-style-type: none"> * malformations * growth restriction * prematurity * associated morbidity, etc • Healthcare outcomes: <ul style="list-style-type: none"> * hospital costs
Starting date	December 2014
Contact information	Contact: Andrea Spence 514-418-0875

NCT02145767 (Continued)

preempttrial@gmail.com

Notes

NCT02633878

Trial name or title	Chinese herbal medicine and micronized progesterone for threatened miscarriage
Methods	RCT intervention model: parallel assignment Masking: quadruple (participant, care provider, investigator, outcomes assessor)
Participants	1656 women age:18-37 years. Inclusion criteria: pregnant (as confirmed by positive urinary pregnancy tests), vaginal bleeding with or without abdominal pain, while the cervix is closed by visual exam and the fetus is viable inside the uterine cavity during early pregnancy (5-10 weeks' gestation), No previous treatment for miscarriage, ability and willingness to give informed consent and willingness to be randomised and to take daily study medications for up to several months
Interventions	Chinese herbal medicine versus progesterone Chinese herbal medicine versus progesterone placebo Chinese herbal medicine placebo versus progesterone Chinese herbal medicine placebo + progesterone placebo
Outcomes	Primary outcome: <ul style="list-style-type: none"> live birth rate Secondary outcome measures: <ul style="list-style-type: none"> ongoing pregnancy rate (beyond gestation 12 weeks, ongoing pregnancy rate (beyond gestation 20 weeks), ongoing pregnancy rate (beyond gestation 32 weeks), live births, premature live births anti-ovarian antibody, angiosperm antibody, anticardiolipin antibody, anti-uterus endometrial antibody, antinuclear antibody resistive index of uterus pulsatility index of ovary; pulsatility Index of uterus biochemical pregnancy loss rate, pregnancy loss rate serum progesterone level pregnancy-induced hypertension diabetes APH preterm delivery, postdate delivery rate pre-eclampsia IUGR rate small-for-gestational-age infant stillbirth rate, neonatal death congenital anomaly, birth defect rate Zung Self-Rating Anxiety Scale, Zung Self-Rating Depression Scale
Starting date	Dec 2015

NCT02633878 (Continued)

Contact information No contacts or locations provided

Notes

NCT02690129

Trial name or title Vaginal progesterone for treatment of threatened miscarriage; randomised clinical trial

Methods RCT intervention model: parallel assignment
Masking: single (participant)

Participants 290 women age: 20 years-35 years

Inclusion criteria: pregnant < 24 weeks pregnant presented with bleeding with or without pain, with single viable fetus (confirmed by U/S examination) and accepting to participate in the trial.
Exclusion criteria: currently under medication for any chronic diseases, hypersensitivity to progesterone, congenital fetal anomaly, on hormonal treatment in the current pregnancy and women conceived via assisted reproduction technique

Interventions **Intervention group:** (progesterone group): complete bed rest for first 48-72 hours. Single daily dose of natural micronised progesterone (Prontogest® 200 mg) at bedtime for 15 days. If needed, a pain killer as Indomethacin 50 mg/rectally twice daily up to control of uterine pain. Complete abstaining from sexual activity or strenuous effort. Rh-ve women will be given a shot of anti-D immunoglobulin 300 uG/IM if they continue to bleed; after 12 weeks' gestation or if undergo surgical evacuation.
Control group: (placebo): will follow the same plan of management without progesterone support.

Outcomes Primary outcome:
 • miscarriage rate up to 28 weeks of gestation
 Secondary outcome:
 • gestational age at delivery or termination of pregnancy

Starting date Feb 2016

Contact information Omar M Shaaban, MD Address: Faculty of Medicine Assiut, Egypt Tel:+201223971457 Email:omshaaban2000@yahoo.com

Notes

NCT02950935

Trial name or title Prospective, double-blind, randomised, placebo controlled, phase iii clinical study assessing the efficacy of natural progesterone 25 mg/bid administered subcutaneously in the maintenance of early pregnancy in women with symptoms of threatened miscarriage

Methods RCT: parallel assignment
Masking: quadruple (participant, care provider, investigator, outcomes assessor)

Participants 268 women age: 18-37 years, BMI: 18-28 kg/m2

NCT02950935 (Continued)

Inclusion criteria: pregnant women attending the study sites with the following characteristics: accept to sign the informed consent form and adhere to the study visit schedule; symptoms of threatened miscarriage including vaginal bleeding and pelvic pain); U/S proof of viable singleton intrauterine pregnancy (positive fetal heart beat); gestation week \geq 6 weeks (5 week + 1 day) and < 12 weeks (11 week + 1 day) according to U/S dating (CRL); closed uterine cervix; subchorionic hematoma, if detected, with < 50% placental detachment

Interventions	<p>Intervention group: subcutaneous injection of progesterone solution will be performed twice a day from onset of threatened miscarriage symptoms until week 12 of pregnancy</p> <p>Control group: placebo; subcutaneous injection of placebo solution will be performed twice a day from onset of threatened miscarriage symptoms until week 12 of pregnancy</p>
Outcomes	<p>Primary outcome:</p> <ul style="list-style-type: none"> ongoing pregnancy rate at 12 weeks of gestation <p>Secondary outcome:</p> <ul style="list-style-type: none"> reduction of the frequency of uterine contractions pain reduction (using a numerical rating scale) reduction of subchorionic hematoma (size of subchorionic hematoma will be measured (in mm) at screening and after treatment) number of women with onset of new threatened miscarriage
Starting date	April 2017
Contact information	Barbara PS Cometti, PHD. Tel:41583601000. Email:barbara.cometti@ibsa.ch
Notes	

APH: antepartum haemorrhage; **IUGR:** intrauterine growth restriction; **PPH:** postpartum haemorrhage; **PPROM:** preterm pre-labour rupture of membranes; **U/S:** ultrasound

DATA AND ANALYSES


Comparison 1. Progestogens versus placebo or no treatment

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Miscarriage	7	696	Risk Ratio (M-H, Fixed, 95% CI)	0.64 [0.47, 0.87]
1.1 Oral progestogen versus no treatment	3	408	Risk Ratio (M-H, Fixed, 95% CI)	0.57 [0.38, 0.85]
1.2 Vaginal progesterone versus placebo	4	288	Risk Ratio (M-H, Fixed, 95% CI)	0.75 [0.47, 1.21]
2 Preterm birth	5	588	Risk Ratio (M-H, Fixed, 95% CI)	0.86 [0.52, 1.44]
3 Pregnancy-induced hypertension	2	337	Risk Ratio (M-H, Fixed, 95% CI)	1.00 [0.54, 1.88]
4 Antepartum haemorrhage	2	337	Risk Ratio (M-H, Fixed, 95% CI)	0.76 [0.30, 1.94]


Progestogen for treating threatened miscarriage (Review)


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
5 Stillbirth	2	262	Risk Ratio (M-H, Fixed, 95% CI)	1.94 [0.18, 20.49]
6 Congenital abnormalities	2	337	Risk Ratio (M-H, Fixed, 95% CI)	0.70 [0.10, 4.82]

Analysis 1.1. Comparison 1 Progestogens versus placebo or no treatment, Outcome 1 Miscarriage.




Analysis 1.2. Comparison 1 Progestogens versus placebo or no treatment, Outcome 2 Preterm birth.







Analysis 1.3. Comparison 1 Progestogens versus placebo or no treatment, Outcome 3 Pregnancy-induced hypertension.




Analysis 1.4. Comparison 1 Progestogens versus placebo or no treatment, Outcome 4 Antepartum haemorrhage.



Analysis 1.5. Comparison 1 Progestogens versus placebo or no treatment, Outcome 5 Stillbirth.



Analysis 1.6. Comparison 1 Progestogens versus placebo or no treatment, Outcome 6 Congenital abnormalities.



APPENDICES

Appendix 1. Search terms for ICTRP and ClinicalTrials.gov

progest* AND miscarriage

progest* AND abortion

dydrogesterone AND miscarriage

dydrogesterone AND abortion

WHAT'S NEW

Date	Event	Description
27 September 2018	Amended	Corrected typo in result in Abstract for the outcome 'preterm birth'.

HISTORY

Protocol first published: Issue 2, 2006

Review first published: Issue 3, 2007

Date	Event	Description
8 August 2018	Amended	Corrected affiliation for Dr Amel Fayed.
8 August 2017	New citation required but conclusions have not changed	With the addition of the data from the new included trials, analysis suggested that progestogens are still probably effective in the treatment of threatened miscarriage but may have little or no effect in the rate of preterm birth. The evidence on congenital abnormalities is uncertain, because the quality of the evidence for this outcome was based on only two small trials with very few events and was found to be of very low quality.
8 August 2017	New search has been performed	Search updated, 29 new trial reports identified, from which three trials were included in this update of the review (Alimohamadi 2013; Turgal 2017; Yassaee 2014), one is awaiting classification

Date	Event	Description
		(Yadav 2015), 10 (in addition to three duplicate), are ongoing and 10 (in addition to two duplicate) were excluded. We updated the risk of bias for all included trials. We have assessed the quality of evidence and included a 'Summary of findings' table in this update.
30 September 2011	New citation required and conclusions have changed	With the addition of the data from the new included studies, meta-analysis suggested that oral progestogen is effective in treating threatened miscarriage. Data analysis also suggested that treatment of women with threatened miscarriage by progestogens did not increase the risk of congenital abnormalities, pregnancy induced hypertension nor antepartum haemorrhage. However these results should be approached with caution due to the small sample size and the poor methodological quality of the included studies.
30 September 2011	New search has been performed	Search updated. We have identified two new studies (El-Zibdeh 2009; Pandian 2009) and included both. Two new authors (Amel A Fayed and Samia A Esmaeil) helped prepare this update. We have updated the methods to reflect the latest Cochrane Handbook (Higgins 2011a).
14 February 2011	New citation required but conclusions have not changed	New author helped prepare the latest update.
1 December 2009	New search has been performed	Search updated. Three new reports identified and excluded (Czajkowski 2007; Song 2007; Vincze 2006). One trial previously awaiting classification has been excluded (Zhang 2000).
20 September 2008	Amended	Converted to new review format.

CONTRIBUTIONS OF AUTHORS

For the 2017 update, Dr Hayfaa Wahabi participated in the selection of eligible trials, assessment of the trials for inclusion, data extraction and analysis and grading of evidence. She participated in writing both the initial and final version of the review.

Dr Amel Fayed participated in the selection of eligible trials, assessment of the trials for inclusion, data extraction and analysis.

Dr. Khawater Bahkali participated in the selection of eligible trials, assessment of the trials for inclusion, data extraction and analysis and grading of evidence. She participated in writing the draft of the review.

Dr Samia Ahmed participated in the selection of eligible trials, assessment of the trials for inclusion, and data extraction. She participated in writing the draft of the review.

DECLARATIONS OF INTEREST

Hayfaa A Wahabi: none known

Amel A Fayed: none known

Samia A Esmaeil: none known

Khawater Hassan Bahkali: none known

SOURCES OF SUPPORT

Internal sources

- Cochrane Review Initiative Project, Saudi Arabia.

External sources

- No sources of support supplied

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

1. The primary outcome 'Miscarriage', was previously listed as 'Early miscarriage up to 12 weeks' and 'Miscarriage later than 12 weeks and less than 23 weeks'. We grouped both outcomes together because the protocol stated that a subgroup analysis for early and late miscarriage would be carried out when data were available
2. The following outcomes are included in this update.
 - a. For the mother:
 - i. pregnancy-induced hypertension
 - ii. antepartum haemorrhage
 - b. For the child:
 - i. intrauterine growth restriction
 - ii. low birthweight
 - iii. birthweight
 - iv. respiratory distress syndrome
3. The methods have been updated to reflect the latest *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011a).

For this update, we assessed trial quality for seven selected outcomes using the GRADE approach (see [Summary of findings for the main comparison](#)).

INDEX TERMS

Medical Subject Headings (MeSH)

Abortion, Spontaneous [epidemiology]; Abortion, Threatened [*drug therapy]; Administration, Intravaginal; Congenital Abnormalities [epidemiology]; Premature Birth [drug therapy] [epidemiology] [prevention & control]; Progestins [adverse effects] [*therapeutic use]; Randomized Controlled Trials as Topic

MeSH check words

Female; Humans; Pregnancy