

CORRIGENDUM

Persistent upregulation of the β -tubulin tubb6, linked to muscle regeneration, is a source of microtubule disorganization in dystrophic muscle

Davide Randazzo¹, Umara Khalique¹, Joseph J. Belanto², Aster Kenea¹, Dana M. Talsness², John T. Olthoff², Michelle D. Tran¹, Kristien J. Zaal¹, Katherine Pak³, Iago Pinal-Fernandez^{3,4}, Andrew L. Mammen^{3,4}, Dan Sackett⁵, James M. Ervasti² and Evelyn Ralston^{1,*}

¹Light Imaging Section, Office of Science and Technology, National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS), National Institutes of Health (NIH), Bethesda, MD, USA, ²Department of Biochemistry, Molecular Biology, and Biophysics, and Program in Molecular, Cellular, Developmental Biology, and Genetics, University of Minnesota-Twin Cities, Minneapolis, MN, USA, ³Laboratory of Muscle Stem Cells and Gene Regulation, Muscle Disease Unit, NIAMS, NIH, Bethesda, MD, USA, ⁴Johns Hopkins University School of Medicine, Baltimore, MD, USA and ⁵Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD), NIH, Bethesda, MD, USA

*To whom correspondence should be addressed at: Light Imaging Section, Office of Science and Technology, National Institute of Arthritis and Musculoskeletal and Skin Diseases, National Institutes of Health 9000 Rockville Pike, Bldg. 50, Rm. 1535, Bethesda, MD 20892-8023, USA. Tel: +1-301-496-6164; Fax: +1-301-402-3417; Email: evelyn.ralston@nih.gov

Human Molecular Genetics, 2019, 28(7), 1117–1135
doi: 10.1093/hmg/ddy418.

The authors would like to apologize for errors within the legend of Figure 10. The legend has been corrected online and in print as below:

- “2.300.30-fold” changed to “2.30±0.30-fold”
- “4.100.41-fold” changed to “4.10±0.41-fold”
- “1.460.05-fold” changed to “1.46±0.05-fold”